Chlamydomonas reinhardtii and Microbacterium fakhimi sp. nov., a mutualistic

association that favor sustainable hydrogen production.

Neda Fakhimi [a, b], Alexandra Dubini[a,*], María Jesus Torres[a], Emilio Fernandez[a], Aurora

Galván[a] and David González-Ballester[a]

a. Universidad de Córdoba, Departamento de Bioquímica y Biología Molecular, Campus Universitario de Rabanales, Ed. C6, Planta Baja, 14071, Córdoba, Spain.
b. Carnegie Institution for Science. Department of Biosphere Sciences and Engineering, 260 Panama Street, Stanford, CA 94305, USA

Emails: Neda Fakhimi (nfakhimi@carnegiescience.edu); A. Dubini (alexandra.dubini@uco.es); M. J. Torres (mjtorres_@msn.com); E. Fernández (bb1feree@uco.es); A. Galván (bb1gacea@uco.es); D. González-Ballester (dgballester@uco.es)

*Corresponding author: Alexandra Dubini (alexandra.dubini@uco.es)

Related research articles:
1) Draft genome sequence of *Microbacterium fakhimi* sp. nov.*,* a novel bacterium associated with the alga *Chlamydomonas reinhardtii*
2) Draft genome sequence of *Stenotrophomonas goyi* sp. nov., a novel bacterium associated with the alga *Chlamydomonas reinhardtii*

Abstract

A multispecies bacterial community including *Microbacterium fakhimi* sp. nov.,

Stenotrophomonas goyi sp. nov., and *Bacillus cereus* greatly promoted sustained hydrogen

production by the microalga *Chlamydomonas reinhardtii* when cocultivated in mannitol- and

yeast extract-containing medium (up to 313 mL·L^{-1}). Alga viability was also largely prolonged

in the cocultures (>45 days) without any nutrient supplementation. Among the bacterial

community, *Microbacterium fakhimi* sp. nov. was the main responsible for the hydrogen

production improvement. Nonetheless, the use of the entire bacterial community allowed a

better growth of the alga during hydrogen production. *Chlamydomonas reinhardtii* and

Microbacterium fakhimi sp. nov. established a mutualistic association, based on the release

of ammonium and acetic acid from the bacterium, while the alga provided sulfur-containing

metabolites and complemented the bacterial auxotrophy for biotin and thiamine. This study uncovers the potential of the Chlamydomonas-bacteria consortia for durable and stable H_2 production while allowing the simultaneous production of biomass.

Keywords: Chlamydomonas, Microbacterium, hydrogen, biomass, algae-bacteria consortia

Introduction

Hydrogen (H_2) is a very interesting clean fuel that do not release CO_2 after combustion and that can be easily interconvertible with electricity, allowing the transport and storage of renewable energy (Abdin et al., 2020). The clean and sustainable H_2 production (green H_2) is currently part of the main energy policies of many governments (Kakoulaki et al., 2021). Among the different technologies to produce green H_2, the biological production of H_2 (bioH_2) offers the possibility of culturing living organisms (bacteria, cyanobacteria, and algae) that liberate H_2. This kind of technology, which is currently in TRL 3-4 stage, allows for H_2 production at low temperatures (23-30ºC) and atmospheric pressure, which is key to produce clean and sustainable energy at low costs (Lepage et al., 2021). BioH_2 is particularly interesting when combined with other biological industrial applications such as wastewater treatment (Aydin et al., 2021) or any valorization of the biomass (e.g., biofertilizers, cosmetics, bioplastics, biofuels, etc.) (Fabris et al., 2020; Leong et al., 2021). Importantly, when using algae, the possibility of using light-driven energy and CO_2 capture is also an advantage of this technology. Still, bioH_2 is currently impeded by some technological (high cost, low TRL stage) and physiological barriers (low yield) that prevent its industrial application (Dębowski et al., 2020).

Heterotrophic bacteria link H_2 production to their fermentative metabolism (dark H_2) (Dahiya et al., 2021), while microalgae and cyanobacteria can produce H_2 linked to their

photosynthetic activity (photobioH$_2$) (Ghirardi et al., 2009). In both cases, H$_2$ production requires anoxic/hypoxic conditions since the hydrogenases/nitrogenases are oxygen sensitive. H$_2$ production in microalgae have been widely studied in *Chlamydomonas reinhardtii* (Chlamydomonas throughout), a microalga able to grow photoautotrophycally or heterotrophically/mixotrophically in the presence of acetic acid (Chaiboonchoe et al., 2014). In this alga, there are two main pathways for H$_2$ photoproduction, namely the PSII-dependent and PSII-independent pathways. The PSII-dependent uses electrons derived from water photolysis to produce H$_2$, while the PSII-independent relies on the non-photochemical reduction of the plastoquinone pool (Fouchard et al., 2005; Hemschemeier et al., 2008). S deprivation have been widely employed to elicit sustainable PSII-dependent H$_2$ production in Chlamydomonas, and many genetic engineered Chlamydomonas strains have been designed to improved H$_2$ production during S deprivation (Dubini and Ghirardi, 2015). However, S deprivation is a stress condition that causes loss of cell viability, lowers biomass yield, and increases cultivation costs, which reduce the potential for industrial applications. On the other hand, H$_2$ production in nutrient-replete media is also possible, mostly by controlling illumination and/or using acetate in the media (Degrenne et al., 2011; Scoma et al., 2014; Jurado-Oller et al., 2015). Assimilation of acetate under low light can elicit H$_2$ production via the PSII-independent pathway (Jurado-Oller et al., 2015; González-Ballester et al., 2017). However, H$_2$ production rates in nutrient-replete media are lower than during S-deprivation. In the last years, algae-bacteria cocultures have emerged as effective strategy to elicit H$_2$ production in both S-depleted and nutrient-replete media (reviewed in (Fakhimi et al., 2020)).

In natural ecosystems, alga-bacteria associations can be complex and versatile, with many bidirectional metabolic collaborations consisting of the exchange of a variety of

molecules (Hom et al., 2015; Ramanan et al., 2016; Yao et al., 2019). Bacteria can provide macronutrients like algal assimilable carbon (Cho et al., 2015) and nitrogen sources. The later can be provided by either fixing nitrogen (Kim et al., 2014) or mineralizing amino acids and peptides (Calatrava et al., 2018). Other nutrient exchanges from the bacteria to the algae include phytohormones (Teplitski and Rajamani, 2011), vitamins (Helliwell et al., 2011) and micronutrients (Amin et al., 2009). In return, the alga can supply O_2 and fixed organic carbon sources (Cho et al., 2015b; Calatrava et al., 2018; Yao et al., 2019)

An evident advantage of coculturing heterotrophic bacteria with algae for H_2 production, is that bacteria can efficiently remove O_2 from the media, which is the most critical bottleneck associated with H_2 photoproduction. However, other than O_2 depletion, little information has been provided explaining the increased H_2 production in algae-bacteria cocultures. In different Chlamydomonas-bacteria cocultures incubated with sugars, H_2 photo-production was linked to the ability of the bacteria to produce acetic acid (Fakhimi et al., 2019a). Moreover, when H_2 producing bacteria and Chlamydomonas are cocultured in sugar-containing media, a synergetic H_2 production can be obtained by integrating the photobiological and fermentative production. This synergetic H_2 production is based on acetic acid exchange (Fakhimi et al., 2019a).

This study describes a multispecies bacterial community composed of *Bacillus cereus* and two novel bacteria (*Microbacterium fakhimi* sp. nov. and *Stenotrophomonas goyi* sp. nov.) capable of associating with Chlamydomonas enabling large sustained algal H_2 production and long-term cell viability of the alga. The genome sequences and growth features of *M. fakhimi* and *S. goyi* are discussed in the related publications (Fakhimi et al., 2023a; Fakhimi et al., 2023b). These two novel bacteria are unable to use sulfate as S source but require S-reduced forms such as cysteine and methionine to grow, while *M. fakhimi* sp.

nov. is additionally auxotroph for the vitamins biotin and thiamine (Fakhimi et al., 2023a; Fakhimi et al., 2023b). *Microbacterium fakhimi* sp. nov. was the solely bacteria responsible of the increase in algal H_2 production. The metabolic interactions established between *Microbacterium fakhimi* sp. nov. and Chlamydomonas are analyzed and discussed.

Materials and methods

Algal and bacterial strains

Chlamydomonas reinhardtii strain 704 (cw15, arg7[+], Nia1:Ars, mt[+]) (Loppes et al., 1999) was used in all the experiments. Chlamydomonas precultures was grown photomixotrophically on Tris Acetate Phosphate (TAP) medium (Harris, 2008) at 24°C under continuous Photosynthetic Photon Flux Density (PPFD) of 60-90 µmol photon·m^{-2}·s^{-1}.

Three bacteria were identified in this work from a bacterial community. Individual members of this bacterial community were isolated by sequential rounds of plate streaking in Yeast Extract Mannitol (YEM) medium, until 3 different types of bacterial colonies were visually identified (**Supplemental Fig. 1**). Colonies were grown separately, and the subsequent isolated DNA was used for PCR-amplification of their partial RNA 16S sequences. After sequencing, the three isolated bacteria were identified as members of the genus *Microbacterium*, *Stenotrophomonas*, *and Bacillus*. Further PacBio whole genome sequencing and genome assembling by SNPsaurus LLC identified these three bacteria as *Microbacterium fakhimi* sp. nov (Fakhimi et al., 2023a), *Stenotrophomonas goyi* sp. nov. (Fakhimi et al., 2023b), and *Bacillus cereus*. *Microbacterium fakhimi* sp. nov. and *Stenotrophomonas goyi* sp. nov. have been deposited in the Spanish Type Culture Collection (CECT) with accession number CECT30765 and CECT30764, respectively. Genome sequences for *Microbacterium fakhimi* sp. nov. and *Stenotrophomonas goyi* sp. nov. have been

deposited in GenBank (NCBI) as CP116871 and SUB12685123. *M. fakhimi* sp. nov. has been submitted for patenting (OEPM Submission P202330306)

All the bacterial precultures were grown on Yeast Extract Mannitol (YEM) medium or LB medium. The bacteria cultures were incubated at 25-28±1°C and under agitation (130 rpm).

Coculturing alga and bacteria

Chlamydomonas precultures were grown to mid-log phase, harvested by centrifugation (3.000×g for 3 min) and resuspended in the corresponding media. Bacterial precultures grown until the Optical Density at 600 nm (OD_{600}) reached 0.6-0.8 (logarithmic phase), then harvested by centrifugation (8.000×g for 3 min), washed twice and resuspended in the corresponding media to a final OD_{600} of 1.0. Alga and bacteria were mixed in the corresponding media with final chlorophyll concentration of 10 $\mu g \cdot mL^{-1}$ and bacterial OD of 0.05 or 0.1. The cocultures and their respective control monocultures were incubated in a growth chamber equipped with LED panels (AlgaeTron AG 230, Photon System Instruments) at 25°C and continuously agitated (180-200 rpm).

Media used in this work were: Minimal Medium (MM) (Harris, 2008); Tris Acetate Phosphate (TAP) (Harris, 2008); TAP supplemented with 56 mM of mannitol (TM); TAP supplemented with 0.8 $g \cdot L^{-1}$ of yeast extract (TY); TAP supplemented with 56 mM mannitol and 0.8 $g \cdot L^{-1}$ of yeast extract (TYM). In some experiments the mannitol and yeast extract concentrations of these media were lowered to 28 mM and 0.4 $g \cdot L^{-1}$, respectively, and the corresponding media denoted as TM_L, TY_L and TYM_L (L standing for Low). Synthetic dairy wastewater was composed of TAP medium supplemented with 2.5 $g \cdot L^{-1}$ of tryptone and 2.5 $g \cdot L^{-1}$ of lactose.

Cultures used for growth test were performed in Erlenmeyer flasks (100 mL) containing 70 ml of the corresponding medium (termed as aerobic condition throughout).

Cultures for H_2 production were performed in hermetically sealed bioreactors (155 mL) containing 100 mL of culture media (termed as hypoxic condition throughout). Bioreactors for H_2 production were opened under sterile conditions every 24 h (aeration) to release the H_2 partial pressure and replace the gas composition in the headspace with atmospheric air (Jurado-Oller et al., 2015). The daily H_2 production and O_2 levels in the headspace were measured before aeration of the bioreactors. All cultures were continuously agitated under 50 µmol photons·m^{-1}·s^{-1}, unless otherwise is indicated.

Measurement of alga and bacteria growth

Alga growth was assessed in terms of chlorophyll concentration (Wintermans and De Mots, 1965) and by counting cells number using a hemocytometer (Sysmex Microcellcounter F-500).

Bacteria growth in monocultures was estimated spectrophotometrically in term of OD_{600} evolution (DU 800, Beckman Coulter). In cocultures, bacterium cells were separated from alga cells by a customized Selective Centrifugal Sedimentation (SCS) approach (Torres et al., 2022). The goal of the technique consists in finding the centrifugation parameters that led to maximize algal cell sedimentation while minimize bacterial cell sedimentation. Thus, measuring the OD of the supernatant after centrifugation can provide an estimation of the bacterial growth in the cocultures. To do this, the % of precipitated cells of each monoculture were calculated at different forces (from 300 x g to 1000 x g) and times (30 s to 2 min) using the measured absorbances of individual monocultures before centrifugation (A_{BC}) and after centrifugating the mixture of those cultures (A_{AC}) (**Supplemental Table 1**). By using 500 x g for 1 min percentage of precipitated *M. fakhimi* cells in cocultures was minimal (meaning that 95.6% of the *M. fakhimi* cells remained in the supernatant). This condition

was chosen as a good compromise for SCS and used to evaluate the contribution of the bacteria to the OD in cocultures ($^{SCS}OD_{600}$).

Colony Forming Unit (CFU)

Colony Forming Unit (CFU) was quantified by making serial 10-fold dilutions of each culture, and 100 µl of each dilution was plated on LB plates. Plates were incubated for 3-4 days at 25°C. Later, the plates containing 50-100 colonies were selected, counted and further calculations were applied using corresponding dilution factors.

Analyses of metabolites in gas and liquid phases

Gas chromatography (GC) (Agilent 7820A, Agilent Technologies) was used to detect H_2 and O_2 evolution in the headspace of the bioreactors. Gas samples from the headspaces (250 µl) were collected using a 1 mL Hamilton's SampleLock™ syringe and manually injected into a Gas Chromatograph (GC) (Agilent 7820A, AgilentTechnologies). H_2 and O_2 were separated using a packed column (60/80 Molecular Sieve 5A, Ref. 13133-U, Supelco) at 75 °C and detected using a Thermal Conductivity Detector (TCD). Argon was used as the carrier gas.

Acetic acid, ethanol, lactic acid, formic acid, and mannitol analyses were performed by HPLC (Agilent series 1200, Agilent Technologies) using an ion-exchange column (Agilent Hi-Plex H, 300 × 7.7 mm, 6 µm I.D.) with isocratic elution using 5 mM H_2SO_4 at 50 °C. Samples (500 µL) were centrifuged, filtered (0.2 µm) and injected (20 µL) onto the HPLC system at a flow rate of 0.6 mL · min^{-1}. Retention peaks were observed using UV (205 nm) and Refractive Index detectors.

RESULTS

Contaminated Chlamydomonas cultures enhanced H$_2$ production in acetate-containing medium (TAP medium).

Preliminary experiments showed that the accidental bacterial contamination of Chlamydomonas TAP cultures enhanced H$_2$ production (bacterial contamination derived from YEM medium). Supernatant from the contaminated Chlamydomonas cultures were preserved and used as inoculum for upcoming experiments. This Initial Bacterial Inoculum will be referred as IBI throughout.

Chlamydomonas TAP cultures were inoculated with IBI in sealed bioreactors and used for H$_2$ production experiments at different light intensities (12, 50 and 100 PPDF) (**Table 1**). As previously reported, Chlamydomonas control cultures produced more H$_2$ at low light intensities (Jurado-Oller et al., 2015). In the IBI-Chlamydomonas cocultures, the H$_2$ production was 1.4, 3.6 and 13.5 times higher than the respective monocultures incubated at 12, 50 and 100 PPDF, respectively. Similar results were obtained previously using cocultures of Chlamydomonas with different bacterial strains including *Pseudomonas putida, Pseudomonas stutzeri, E. coli* and *Rhizobium etli* where the highest improvements in H$_2$ production, relative to Chlamydomonas monocultures, were obtained at high light intensities (Fakhimi et al., 2019b; Fakhimi et al., 2019a). No H$_2$ production was observed in the solely bacterial IBI cultures, indicating that the bacterial community did not produce H$_2$ under these conditions.

The Chlamydomonas-IBI consortium can sustain H$_2$ production and prolong the algal cell viability when incubated in the medium containing mannitol and yeast extract (TYM medium).

The H$_2$ production capability of the Chlamydomonas-IBI cocultures was studied under different nutrient sources that are more suitable for the bacteria growth. TAP medium was either supplemented with mannitol (TM), yeast extract (TY) or both (TYM). H$_2$ production, O$_2$ evolution, growth, and acetic acid and mannitol uptake were monitored for 30 days (**Table 2** and **Fig. 1**).

All Chlamydomonas monocultures except TM monocultures produced H$_2$ (6.03 to 12.34 mL·L^{-1}) during the first two days, afterward no more H$_2$ production was observed (**Table 2**). None of the IBI cultures produced H$_2$, confirming that Chlamydomonas was the only H$_2$ producer (**Table 2**). However, a very remarkable boost in H$_2$ production was observed in cocultures incubated in TYM (164 mL·L^{-1}; 21-fold total H$_2$ production) (**Table 2** and **Fig. 1a**). A moderate improvement in H$_2$ production was also observed in the TY cocultures (18.77 mL·L^{-1}; 1.56-fold total H$_2$ production). No H$_2$ production was observed in the TM cocultures (**Table 2**). Moreover, H$_2$ production in the TYM medium extended for 25 days, while in the rest of the cultures, the production was only sustained for 2-3 days. H$_2$ production in the TYM cocultures experienced 3 phases. The first phase consisted in 2 days of relatively high H$_2$ production rate (8 mL H$_2$·L^{-1}·d^{-1}), afterwards 2 more days with very low H$_2$ rate (< 1 mL·L^{-1}·d^{-1}), and then a third phase started from day 4 with increasing rates of H$_2$ production to a maximum of 16.12 mL·L^{-1}·d^{-1} on the 23rd day (**Fig. 1a**). The TYM cocultures were the only Chlamydomonas-containing cultures able to sustain hypoxia for a long period (23 days) (**Fig. 1b**).

The growth of Chlamydomonas (by the mean of chlorophyll concentration) under H$_2$ production condition was determined (**Fig. 1c**). In the TM medium, Chlamydomonas monocultures and cocultures turned yellow and died after 24h, indicating a likely toxic effect of the high mannitol concentration (56 mM) for the alga. Surprisingly,

Chlamydomonas cells incubated in the TYM media did not suffer such a toxic effect. On the contrary, the chlorophyll content of the TYM monocultures and cocultures peaked at 76 and 59 $\mu g \cdot mL^{-1}$ on day 15, respectively, which was higher than in the standard TAP medium (48 $\mu g \cdot ml^{-1}$ for both monocultures and cocultures). These data suggest that the simultaneous supplementation with yeast extract and mannitol not only protected Chlamydomonas cells from the toxic effect of mannitol, but also provided a better growth condition relative to the standard TAP medium. Previously, it was shown that mannitol cannot be used as carbon source by Chlamydomonas (Chaiboonchoe et al., 2014; Fakhimi et al., 2019a). It may be possible that yeast extract is playing two roles simultaneously: a) compensating the potential osmotic stress caused by mannitol, and b) providing a N source for the alga through the activity of the extracellular L-amino acid oxidase (LAO1), which liberates ammonium and different α-keto-acids from small peptides and amino acids (Calatrava et al., 2019). This second role could also explain the higher growth observed in Chlamydomonas monocultures incubated in TY relative to TAP (**Fig. 1c**).

There was no difference in term of chlorophyll concentrations between the alga monocultures and cocultures incubated in TAP and TY, which reflect the neutral effect of the bacteria on Chlamydomonas growth in these media. However, in the TYM medium, less chlorophyll concentration was observed in the cocultures than in the monocultures. Still, a decline in chlorophyll concentration was observed in all the cultures in the 3rd week, except in the TYM cocultures where the chlorophyll remained stable for 30 days (**Fig. 1c**), indicating a longer viability of the alga when cocultured with IBI in the TYM medium under hypoxia.

Solely bacterial cultures (IBI cultures) grew very efficiently in the presence of mannitol (TM and TYM media), showing a two-phase growth kinetic (**Fig. 1d**). During the first 5 days, the TYM and TY bacterial cultures grew similarly and faster than in TAP and TM.

Afterwards, only TYM and TM showed a pronounced second growth phase reaching similar final ODs. The final growth of the cultures incubated in TAP and TY were also paired. Therefore, it appears that the presence of yeast extract promoted a rapid initial growth of the IBI cultures, although only the presence of mannitol allowed for sustained growth.

Despite mannitol not being assimilated by Chlamydomonas, continuous consumption of mannitol was observed in the TYM cocultures (**Fig. 1e**), indicating the ability of the bacteria community to uptake this nutrient. Acetic acid was quickly consumed in cocultures and algal monocultures (**Fig. 1f**). As previously reported, the slower the acetic acid is consumed the more sustained is hypoxia and H_2 production in Chlamydomonas (Fakhimi et al., 2019b). Accordingly, the slightly slower consumption of acetic acid observed in the TY cocultures led to a bit more sustained H_2 production relative to the TAP cocultures (3 vs 2 days) (**Table 2**). The TYM cocultures also showed a slower acetic acid uptake but a much more sustained H_2 production phase (25 days). In this case, it is possible that the excretion of acetic acid from the bacteria through fermentation of mannitol supported sustained H_2 production by the alga. Accumulation of acetic acid in the TYM cocultures was detectable only from day 18, although it is possible that experimentally undetectable levels of accumulated acetic acid were present from the 4$^{\text{th}}$ day (**Fig. 1f**).

In another experimental set up, Chlamydomonas monocultures incubated in sealed bioreactors with TAP, TY and TYM were left for 16 days. During this period, a short (2 days) H_2 production phase (< 12 mL·L^{-1}) was obtained in all the cultures (**Fig. 2a**). Initial acetic acid was totally consumed after 16 days (**Fig. 2d**). On day 16, a set of cultures were inoculated with the IBI, and the rest remained as Chlamydomonas monocultures. A great and sustained H_2 production was observed only in the IBI-inoculated cultures incubated in TYM (total of 313 mL H_2·L^{-1} along 17 days) (**Fig. 2a**). O_2 depletion was also observed only in the TYM-

inoculated cultures (**Fig. 2b**). The TY and TAP cocultures did not produce extra H_2 or deplete the O_2. Acetic acid in the TYM medium was detectable 4 days after the IBI inoculation, which probably explains the algal H_2 production observed in this medium (**Fig. 2d**). In all the cultures, but in the TYM cocultures, the chlorophyll content present on the 16^{th} day continuously dropped along the experiment (**Fig. 2c**). However, in the TYM cocultures the chlorophyll content remained constant (around 80 $\mu g \cdot mL^{-1}$) until the end of the experiment (day 30) and conserved a green color, confirming the previous observation that the IBI community can extend the viability of the Chlamydomonas cultures cultivated in the TYM medium under hypoxia.

Overall, the Chlamydomonas-IBI cocultures incubated in TYM under hypoxia promoted sustained algal H_2 production (up to 313 $mL \cdot L^{-1}$ for >16 days) (**Fig. 3a**) that was compatible with algal growth (up to 59 $\mu g \cdot mL^{-1}$) (**Fig. 1c**) and further extended Chlamydomonas cells viability for more than 30 days without any nutrient supplementation (**Fig. 1c and 2c**). This long viability of the Chlamydomonas cells was not observed in any other conditions. The extended viability could be linked to the acetic acid availability derived from the bacterial fermentation of mannitol (**Fig. 2d**), although the possibility that other nutrient exchanges (e.g., N or S sources), supporting Chlamydomonas survival cannot be discarded.

To our knowledge, the H_2 production detected in this consortium is the highest reported using non-genetically modified Chlamydomonas strains. Previously, the highest Chlamydomonas H_2 production yields were obtained with the *pgr5* (850 $mL \cdot L^{-1}$; 9 days) (Steinbeck et al., 2015), *pgrl1* (437 $mL \cdot L^{-1}$; 5 days) (Tolleter et al., 2011), *stm6* (540 $mL \cdot L^{-1}$; 14 days) (Kruse et al., 2005), D1-*L159I-N230Y* (504 $mL \cdot L^{-1}$; 12 days) (Torzillo et al., 2009) and

Stm6Glc4L01 (361 mL·L^{-1}; 8 days) (Oey et al., 2013) mutant lines in acetate-containing S-depleted medium (TAP-S).

The production of H$_2$ observed in the Chlamydomonas-IBI cocultures is also the highest amount reported in any Chlamydomonas-bacteria cocultures (Fakhimi et al., 2020). Previously, the highest Chlamydomonas-bacteria H$_2$ production was reported with *Bradyrhizobium japonicum* (170-141 mL·L^{-1}; 14-16 days) (Wu et al., 2012; Xu et al., 2016) and *Pseudomonas* spp. (130 mL·L^{-1}; 12 days) (Ban et al., 2018); all of them incubated under S-deficiency (TAP-S).

However, the use of mutant strains and/or S deficiency may have some biotechnological drawbacks. First, implementation of S deficiency (like any other nutrient limitation) implies a two-stage bioreactor process and often designing of synthetic media which are not the cheapest options, and greatly limit the utilization of inexpensive nutrient sources such as wastewaters. Moreover, H$_2$ production in S-depleted media not only does not support growth but also declines the initial chlorophyll content and cell viability of the alga. Accordingly, no growth was reported during H$_2$ production in the mutant strains described above. Moreover, all the mentioned above Chlamydomonas mutant strains have growth or/and photosynthetic deficiencies, which may further hinder the long-term sustainability of the cultures and the biomass production. Finally, genetically modified organisms could also raise some environmental concerns and face legal drawbacks that may further limit their use.

Importantly, the H$_2$ production in the Chlamydomonas-IBI cocultures incubated in the TYM medium was concomitant with the growth of the alga and supported a long alga viability (more than in any other conditions) without any further nutrient supplementation.

Identification of the members of the bacterial IBI community

To clarify the specific roles of each bacterium within the Chlamydomonas-IBI cocultures, 3 different bacteria present in this community were isolated and identified (**Supplemental Fig. 1**). Whole genome sequences were obtained and used to genetically identify and classified these 3 bacteria through the Type (Strain) Genome Server (TYGS) (Meier-Kolthoff and Göker, 2019). As a result, *Bacillus cereus* was identified together with two other novel species belonging to the genus of *Microbacterium* and *Stenotrophomonas,* which were named as *Microbacterium fakhimi* sp. nov. (*M. fakhimi*) (Fakhimi et al., 2023a) and *Stenotrophomonas goyi* sp. nov. (*S. goyi*) (Fakhimi et al., 2023b), respectively.

Interestingly, *Microbacterium sp.* and *Stenotrophomonas sp.* have been previously isolated from a contaminated Chlamydomonas culture with moderate enhanced H_2 production in TAP-S (25-34 mL $H_2 \cdot L^{-1}$) (Li et al., 2013).

Microbacterium sp., *Bacillus* sp. and *Stenotrophomonas sp.* are among the most common inhabitants of the rhizosphere (Rosenblueth and Martínez-Romero, 2006; Marquez-Santacruz et al., 2010; Abadi et al., 2020). They are also recurrent bacteria frequently found within algal cultures (Burmølle et al., 2006; Li et al., 2013; Jones et al., 2016; Ji et al., 2019; Martelli et al., 2021; Mitra et al., 2021), which is probably reflecting frequent coexistence of these organisms in natural ecosystems.

Deconvolution of the individual roles of each member of the IBI community in H_2 production and extension of the alga viability

H_2 production was evaluated in pairwise cocultures of Chlamydomonas with *M. fakhimi* sp. nov., *S. goyi* sp. nov. or *B. cereus* in the TYM medium. Also, to test any potential multiple collaboration between the three bacteria, Chlamydomonas was cocultured with

two-by-two bacteria combinations, with the three bacteria, as well as with the original IBI community (**Fig. 3**). Chlamydomonas monocultures were cultivated in the TYM medium for three days, afterwards, once the initial acetic acid content was totally consumed and the initial H_2 production stopped, the different bacteria combinations were inoculated. Although all the cultures produced some H_2, only the *M. fakhimi*-containing cultures showed noticeable and sustained H_2 production (**Fig. 3a**). H_2 production among all the *M. fakhimi*-containing cocultures was similar (35.8-47.1 mL·L^{-1}) and sustained for at least 7 days. Moreover, although all the cocultures eventually reached hypoxia after bacteria inoculation, only the *M. fakhimi*-containing cocultures were able to prolong hypoxia for several days (**Fig. 3b**). None of the bacteria monocultures produced H_2, confirming that H_2 was only produced by the alga. From these results, it was evident that only *M. fakhimi* sp. nov. promoted sustained H_2 production in Chlamydomonas, while *B. cereus* or *S. goyi* sp. nov. did not show a relevant role in H_2 production.

The capability of the individual bacteria to uptake mannitol and acetic acid was also analyzed in the bacterial monocultures (**Fig. 3c** and **d**). Only *M. fakhimi* and *B. cereus* were able to uptake some mannitol (**Fig. 3c**). *S. goyi* and *B. cereus* consumed acetic acid, especially the former which showed a very efficient acetic acid uptake (**Fig. 3d**). Importantly, the *M. fakhimi* monocultures did not consume any acetic acid, but its excretion was observed, which probably derived from mannitol fermentation (**Fig. 3d**). This results also pointed out to *M. fakhimi* as the only bacterium responsible for the acetic acid excretion observed in the Chlamydomonas-IBI cocultures (**Fig. 1f** and **2d**), which is also likely linked to the enhanced algal H_2 production.

Once *M. fakhimi* sp. nov. was identified as the only IBI member able to promote sustained H_2 production in Chlamydomonas, the next question addressed was if this

bacterium was also responsible for the prolonged viability of the Chlamydomonas cultures in the TYM medium. To answer this, Chlamydomonas and *M. fakhimi* sp. nov. were cocultured in the TYM medium in not-sealed bioreactors (aerobic condition) for 70 days (**Fig. 4**). Due to the mannitol toxic effect on the alga previously encountered in the TM medium (**Fig. 1c**), the TYM mannitol concentration was reduced from 56 mM to 28 mM and referred as TYM_L (yeast extract was also lowered to 0.4 g·L^{-1} in the TYM_L medium). Initially, the Chlamydomonas monocultures and cocultures incubated in TYM_L grew similarly for 4-5 days (**Fig. 4a** and **4b**). After this initial growth, both the chlorophyll content and the alga cell number drastically decreased in the monocultures but not in the cocultures. The chlorophyll and cell number of the cocultures remained stable for more than 30 days (**Fig. 4a** and **4b**, and **Supplemental Fig. 2**), proving that *M. fakhimi* sp. nov. can prolong the alga cell viability in the TYM_L medium under aerobic conditions.

As commented before, this extended algal cell viability is probably related with the capacity of *M. fakhimi* sp. nov. to excrete acetic acid (Fig. 3d), which is used by Chlamydomonas as C source. But also, very important, the bacterial proteolytic activity led to the release of ammonium from amino acids/peptides, which provided the alga with a N source for a long period (Fig. 4d).

Although the above results revealed a positive effect of *M. fakhimi* sp. nov. on algal H_2 production and alga viability, it remained unclear whether the bacterium also benefited from the alga (mutualism versus commensalism relationship). To clarify this situation, bacterial Colony Forming Units (CFU) were measured at the beginning of the experiment and on day 45 in both cocultures and bacteria control cultures. The CFU increased from 0.18×10^9 to 3×10^9 in cocultures, while no colonies appeared in the bacterial monocultures after 45 days (**Fig. 4c**). These results show that the alga and the bacterium established a

mutualistic relationship where the growth of both microorganisms was benefited when incubated in TYM_L. Similar results were obtained with *S. goyi* sp. nov. when cocultured with Chlamydomonas in TYM_L medium (Fakhimi et al., 2023b).

In sum, under aerobic conditions, Chlamydomonas and *M. fakhimi* sp. nov. established a mutualistic relationship where the growth of both microorganisms was benefited. Similarly, in the Chlamydomonas-*Microbacterium paraoxydans* consortium used for phenol bioremediation, a significant increase in the growth rate of both microorganisms was found (Mora-Salguero et al., 2019). Yet, in another study, no alga growth promotion was observed in Chlamydomonas-*Microbacterium* sp. cocultures incubated in nutrient-repleted media (TAP) (Li et al., 2013).

Various strains of *Microbacterium sp.* are known to promote plant growth through indole-3-acetic acid and ammonia production, ACC deaminase activity (Singh and Singh, 2019) and by affecting S and N metabolism (Cordovez et al., 2018). However, the potential role and mechanisms of *Microbacterium* sp. as algal growth-promoting bacteria are unknown. In the Chlamydomonas-*M. fakhimi* sp. nov. consortium incubated in TYM_L, the extended algal viability observed under aerobiosis was likely supported by the acetic acid and ammonium secreted by the bacteria after metabolization of the mannitol and amino acids, respectively (**Fig. 3d, 5h and 5i**).

On the other hand, in a related publication has been shown that *M. fakhimi* sp. nov. is auxotroph for biotin and thiamine (Fakhimi et al., 2023a). Moreover, the bacterium can grow on different C sources although is unable to use sulfate as S source; its growth depends on S-reduced forms such as methionine or cysteine (Fakhimi et al., 2023a). Thereby, Chlamydomonas promoted the bacterium growth probably by alleviating the

bacterial auxotrophy for biotin and thiamine and by providing S-reduced forms. Methionine and cystathionine are known S-containing metabolites secreted by Chlamydomonas (Vogel et al., 1978; Lorincz et al., 2010). In addition to these nutrients exchange, photosynthetic O_2 probably also contributed to promote the bacterium growth in cocultures.

Relevance of media composition during H_2 production in the Chlamydomonas-*M. fakhimi* sp. nov. consortium

H_2 production in Chlamydomonas-*M. fakhimi* sp. nov. cocultures and respective monocultures was assessed in sealed bioreactors (hypoxic condition) with different media: TAP, TM_L, TY_L and TYM_L (**Fig. 5a**). As expected, H_2 production was promoted and sustained in the TYM_L medium. Similarly, probably due to the decreased content of mannitol employed in the TM_L medium, and in turn the lower toxicity for Chlamydomonas, H_2 was also produced in this medium, although less sustained and in less quantity compared with the production in the TYM_L medium. As previously observed, sustainability of the H_2 production phase correlated very well with the presence of acetic acid in the medium (**Fig. 5h**) and the duration of the hypoxia phase (**Fig. 5b**).

Mannitol consumption in TM_L and TYM_L cocultures was similar (**Fig. 5c**). Mannitol uptake was also observed TYM_L bacterial monocultures, but not in TM_L bacterial monocultures (**Fig. 5c**). Accordingly, in the absence of yeast extract, *M. fakhimi* sp. nov. monocultures did not grow efficiently on mannitol (TM_L medium) (**Fig 5e**). The growth of the bacterium in cocultures was estimated using a Selective Centrifugal Sedimentation (SCS) approach (Torres et al., 2022) (**Supplementary Table 1**). *M. fakhimi* sp. nov. grew very efficiently on mannitol when co-cultivated with the alga, and independent of the addition of yeast extract (TYM_L and TM_L media) (**Fig. 5f**). From these results, it can be concluded that *M.*

fakhimi sp. nov. can grow efficiently on mannitol either when the medium is co-supplemented with yeast extract or when it is cocultured with Chlamydomonas (**Fig. 1c, 5e and 5f**). Thereby, the alga must be providing some nutrients to the bacterium allowing the bacterial utilization of mannitol and growth in the absence of yeast extract. Indeed, mannitol uptake and bacterial growth was higher in the TM cocultures than in the TYM monocultures, implying a better performance of the bacterium when cocultured with the alga without yeast extract than when incubated as a monoculture in TYM. As commented before, Chlamydomonas is likely providing biotin, thiamine, cystathionine, and methionine that can support the bacterium growth on the medium without yeast extract (TM medium).

Moreover, *M. fakhimi* sp. nov. monocultures were able to uptake acetic acid only in the TY_L medium (**Fig 5i**), which suggests that the bacterium only uptakes this nutrient if there is no other carbon source and yeast extract is available. Furthermore, the acetic acid was consumed faster in the TM_L than in the TYM_L cocultures (**Fig. 5h**). As commented before, slow acetic acid uptake rates benefit H_2 production (Fakhimi et al., 2019b), which likely explains why the H_2 production observed in TYM_L was higher than in TM_L (**Fig. 5a**). Thereby, the presence of yeast extract reduced the acetic acid uptake in the cocultures, which in turn benefited H_2 production. Finally, the cocultures incubated in TYM_L showed higher accumulation of lactic acid and ethanol while less formic acid than the TM_L cocultures (**Supplemental Fig. 3**). These metabolic differences could also impact H_2 production.

In sum, the presence of yeast extract in the media allowed the bacterial monocultures to uptake mannitol and grow. Nevertheless, in the absence of yeast extract, *M. fakhimi* sp. nov. can also grow over mannitol if cocultivated with Chlamydomonas, which likely provides biotin, thiamine, cystathionine, and methionine. Still, the presence of yeast

extract slowed down the uptake of acetate by the Chlamydomonas-*M. fakhimi* sp. nov. cocultures, which is key to maximize H_2 production.

Remarkably, no significant alga growth was observed in the TYM_L cocultures during H_2 production condition (**Fig. 5d).** This fact greatly contrasts with the data obtained when Chlamydomonas was cocultivated with the IBI community under the same condition (hypoxia) (**Fig. 1c**) or with *M. fakhimi* sp. nov. incubated under aerobic condition (**Fig. 4a**), where substantial alga growth was obtained. It is possible that other members of the IBI community (*Bacillus sp.* and/or *S. goyi*) counteract this negative impact of *M. fakhimi* sp. nov. over the alga growth when cultivated under hypoxia.

Potential of Chlamydomonas-*M. fakhimi* sp. nov. consortium to produce H_2 in a synthetic dairy wastewater.

Due to the ability of *M. fakhimi* sp. nov. to grow on sugars and peptides/amino acids, the potential of the consortium to produce H_2 in synthetic dairy wastewaters was investigated. The consortium was cultivated in a synthetic dairy wastewater containing lactose and casein-derived tryptone. Sustained H_2 production was obtained for at least 11 days, with a total production of 70 mL·L^{-1} (**Fig. 6**).

This result highlights the potential biotechnological interest of Chlamydomonas-*M. fakhimi* sp. nov. consortium.

Suitability of the IBI- and *M. fakhimi*-Chlamydomonas consortia for biotechnological purposes.

Apart from the large sustained H_2 production obtained with the IBI-Chlamydomonas consortium, one of the most interesting features of these consortium is the possibility of simultaneously obtaining H_2 and algae biomass. Algae biomass is of biotechnological interest for the designing of new downstream process based on algae biomass valorization

(Fabris et al., 2020; Leong et al., 2021), which can reduce the cost and thereby increase the economic viability of the industrial $bioH_2$ production (Dębowski et al., 2020; Lepage et al., 2021). In this regard, $BioH_2$ production and biomass valorization are particularly interesting when combined with wastewater treatment (Aydin et al., 2021). Some studies have shown the feasibility of producing $bioH_2$ in wastewater environment, although the H_2 production rate quickly and considerably declined (Hwang et al., 2018; Shetty et al., 2019). In the Chlamydomonas-IBI and Chlamydomonas-*M. fakhimi* cocultures sustainable H_2 production is possible if the media is enriched with sugar and proteins/peptides. Dairy wastewater could be an ideal broth to assay concomitant $bioH_2$ production, bioremediation, and biomass valorization with these consortia, although further research with real dairy wastewater is needed to confirm this potential.

When using different Chlamydomonas-bacteria consortia, various parameters have been associated with the interruption of algal H_2 production: a) rapid death of bacteria (Lakatos et al., 2014); b) acetic acid depletion (Lakatos et al., 2014; Fakhimi et al., 2019b); and c) loss of alga cells viability due to low pH and/or excessive accumulation of bacterial fermentative products, especially of acetic acid (Fakhimi et al., 2019a). For example, in Chlamydomonas-*E. coli* consortia incubated in glucose, there is a rapid acetic acid accumulation and pH decrease that provoke the alga death. On the other hand, in Chlamydomonas cocultures with *P. putida* or *R. etli* incubated with glucose and mannitol, respectively, acetic acid accumulation in the media was low and in turn the alga produced low H_2 rates (Fakhimi et al., 2019a).

All these drawbacks were successfully bypassed in the Chlamydomonas-IBI and -*M. fakhimi* sp. nov. consortia incubated in TYM: a) the bacterial growth was promoted in cocultures relative to bacterial monocultures, at least for *M. fakhimi and S. goyi* (**Fig. 4c** and

5e,f) (Fakhimi et al., 2023b)*; b*) out of the three members of the IBI community only *S. goyi* showed intense acetic acid uptake when incubated in TYM, whereas *M. fakhimi* excreted acetic acid (**Fig. 3d**), which allowed sustained algal H_2 production (**Fig. 1a** and **5a**); and c) the mannitol fermentation, especially by *M. fakhimi* sp. nov., was fast enough to provide sufficient acetic acid allowing sustainable H_2 production (~5.5 $mL \cdot L^{-1} \cdot d^{-1}$), but slow enough to avoid acidification of the medium and impair the viability of the alga. In fact, algal viability was largely promoted in the Chlamydomonas-IBI and Chlamydomonas-*M. fakhimi* sp. nov. cocultures under hypoxic and aerobic conditions, respectively.

On the other hand, it is intriguing that the Chlamydomonas-*M. fakhimi* cocultures, unlike the Chlamydomonas-IBI cocultures, did not promoted the alga growth when incubated under hypoxia (**Fig. 5d**). Hence, the presence of *S. goyi* sp. nov. and *B. cereus* could be crucial to obtain a growth equilibrium that benefited the alga growth and the H_2 production in TYM_L medium under hypoxia.

The potential metabolic complementation that these four microorganisms could established in TYM_L medium could be complex and interdependent. Interestingly, *S. goyi* sp. nov. and *M. fakhimi* sp. nov. are unable to use sulfate as S source and they show a dependency on S-reduced sources such as cysteine and methionine (Fakhimi et al., 2023a; Fakhimi et al., 2023b). Hence, the survival of *M. fakhimi* sp. nov. and *S. goyi* sp. nov. in natural ecosystems could be highly dependent on the surrounding microbial communities. In this multispecies association, the alga could provide S-reduced sources in the form of cystathionine and methionine to both *S. goyi* sp. nov. and *M. fakhimi*. Moreover, Chlamydomonas, *S. goyi* and *B. cereus* could be alleviating the auxotrophy for biotin and thiamine of *M. fakhimi* sp. nov. (the genomes of *S. goyi* sp. nov. and *B. cereus* have entire biosynthetic pathways for biotin and thiamine, whereas the biosynthesis of these vitamins is

well documented in Chlamydomonas (Aaronson et al., 1977; Croft et al., 2007)). On the other hand, the bacteria could provide CO_2 and acetic acid to the alga through their fermentative activity. Acetic acid is mostly provided by *M. fakhimi* sp. nov. Moreover, S. *goyi* sp. nov. and *M. fakhimi* have great proteolytic capacity and can grow over amino acids/peptides as only C and N sources (Fakhimi et al., 2023b; Fakhimi et al., 2023a). Thereby, they both could also provide the alga with ammonium derived from the mineralization of the amino acids/peptides.

In any case, this multispecies association was mutually beneficial and guaranties the viability of the consortium by preventing an excessive bacterial growth, which could be one of the main drawbacks when algae-bacteria cocultures are used for biotechnological applications. Further research is needed to better resolve the relationships of these four microorganisms when incubated together.

Conclusion

The consortium composed of the alga Chlamydomonas, *M. fakhimi* sp. nov., S. *goyi* sp. nov. and *B. cereus* can concomitantly sustain H_2 and biomass production when incubated in mannitol and yeast extract. *M. fakhimi* sp. nov. is the sole bacterium responsible to promote H_2 production in Chlamydomonas, although S. *goyi* sp. nov. and *B. cereus* could be important to generate alga biomass and extend the alga viability under hypoxia. *M. fakhimi* sp. nov. and S. *goyi* sp. nov. can be mutualistically associated with the alga Chlamydomonas. This consortium could be of biotechnological interest for H_2 and biomass production using wastewaters. The potential biotechnological application and the precise metabolic complementation of this consortium need to be further investigated under different growth conditions.

Figure 1. Comparison of Chlamydomonas-IBI cocultures in different media. H_2 production (**a**), O_2 evolution (**b**), chlorophyll concentration (**c**), bacterial OD_{600} (**d**), and mannitol (**e**) and acetic acid uptake (**f**) of Chlamydomonas-IBI (cocultures), and respective control cultures (algal monocultures and IBI cultures) incubated in TAP, TYM, TY or TM under 50 PPFD. For the sake of clarity, in panel (a) only cocultures are plotted. Mannitol and acetic acid uptake were not assayed in sole bacterial cultures. OD_{600} measurements were performed only over IBI cultures. Initial chlorophyll and bacterial OD_{600} were of 10 µg/mL and 0.05, respectively. Cultures were incubated in sealed bioreactors and aerated daily. H_2 production in the headspaces was measured daily before aeration, and accumulative H_2 evolution is plotted. Represented data are the average of three independent experiments.

Figure 2. Effect of IBI inoculation on Chlamydomonas cultures. Chlamydomonas monocultures were incubated in TAP, TY and TYM media for 16 days (initial chlorophyll 10 µg·mL^{-1}). On day 16th, a set of cultures were inoculated with the IBI (final OD_{600} of 0.05) (indicated with an arrow) and the rest were left as algal monocultures. H_2 production (**a**), O_2 evolution (**b**) during the 33 days and, chlorophyll concentration (**c**), and acetic acid levels (**d**) from day 15th are plotted. Cultures were incubated in sealed bioreactors and aerated daily. H_2 production in the headspaces was measured daily before aeration, and accumulative H_2 evolution is plotted. Data represent only one biological sample.

Figure 3. H_2 production in cocultures of Chlamydomonas with *M. fakhimi*., *B.* cereus., *S. goyi* or the original IBI community. H_2 production (**a**), O_2 evolution (**b**) of the different Chlamydomonas cocultures and respective monocultures, and mannitol (**c**) and acetic acid uptake (**d**) in bacterial monocultures. Chlamydomonas monocultures were incubated in the TYM medium, and on day 3, these cultures were inoculated with either, one, two or three different bacteria simultaneously (indicated by an arrow). Initial chlorophyll and bacterial OD_{600} were 10 µg/mL and 0.1, respectively. Cultures were incubated in sealed bioreactors and aerated daily. H_2 production in the headspaces was measured daily before aeration, and accumulative H_2 evolution is plotted. All cocultures are color-coded, while grey symbols

refer to bacterial monocultures. Represented data are the average of three independent experiments.

Figure 4. Long-term growth performance of Chlamydomonas-*M. fakhimi* sp. nov. consortium under aerobic conditions. Chlorophyll content (**a**), Chlamydomonas cell number (**b**) and (**c**) bacterial Colony Forming Units (CFU), were determined in the Chlamydomonas-*M. fakhimi* sp. nov. consortium and respective monocultures incubated in the TYM_L medium for 70 days. Ammonium released into the medium was also measured in cocultures under the same growth conditions for 28 days (**d**). CFU was measured on day 0 and 45. Initial chlorophyll and bacterial OD_{600} were 10 μg/mL and 0.1, respectively. Represented data are the average of three independent experiments. N.D., not detected.

Figure 5. Comparison of Chlamydomonas-*M. fakhimi* sp. nov. in different media. H_2 production (**a**), O_2 evolution (**b**), mannitol uptake (**c**) chlorophyll concentration (**d**), bacterial growth in monocultures (OD_{600}) (**e**) and cocultures ($^{SCS}OD_{600}$) (**f**), and acetic acid uptake (**f, h, i**) of Chlamydomonas-*Microbacterium* cocultures and respective monocultures. For the sake of clarity, in panel (a) only cocultures are plotted. Mannitol was not uptaken by Chlamydomonas monocultures (data not shown). Initial chlorophyll and bacterial OD_{600} were 10 μg/mL and 0.05, respectively. Cultures were aerated daily. H_2 production in the headspaces was measured daily before aeration, and accumulative H_2 evolution is plotted. Represented data are the average of three independent experiments.

Figure 6. H_2 production by Chlamydomonas-*M. fakhimi* sp. nov. cocultures incubated in synthetic dairy wastewater. Synthetic dairy wastewater consisted of TAP medium supplemented with 2.5 $g·L^{-1}$ of tryptone and 2.5 $g·L^{-1}$ of lactose. Initial chlorophyll and bacterial OD_{600} were 10 μg/mL and 0.1, respectively. Alga and bacterium monocultures were used as controls. Cultures were aerated daily. H_2 production in the headspaces was measured daily before aeration, and accumulative H_2 evolution is plotted. Represented data are the average of three independent experiments.

Table 1: **H_2 production ($mL·L^{-1}$) in the TAP medium under different light intensities**

Table 2. H$_2$ production (mL·L^{-1}) in different media. TAP was supplemented with 56 mM of mannitol (TM), or 0.8 g·L^{-1} of yeast extract (TY) or both (TYM). Numbers between parentheses indicated the length of the H$_2$ production phase (days). Cultures were incubated under 50 PPFD.

Supplemental Figure 1. Visual characterization of the IBI community under the microscope (**a**), and the isolated colonies of *Stenotrophomonas goyi* sp. nov. (**b**), *Bacillus cereus* (**c**), and *Microbacterium fakhimi* sp. nov. (**d**).

Supplemental Figure 2. Chlamydomonas monocultures and Chlamydomonas-*M. fakhimi* cocultures incubated in TYM along 60 days.

Supplementary Figure 3. Lactic acid (**a**, **b**, **c**, **d**), formic acid (**e**, **f**, **g**, **h**) and ethanol (**i**, **j**, **k**, **l**) concentrations found in the media of Chlamydomonas-*M. fakhimi* sp. nov. cocultures and respective monocultures incubated in TAP, TM$_L$, TY$_L$, TYM$_L$. Growth conditions are as in Fig 5. Represented data are the average of three independent experiments.

Supplemental Table 1. Centrifugation forces and times used to determine the Selective Centrifugal Sedimentation (SCS) approach for Chlamydomonas and *M. fakhimi*.
Absorbance for Chlamydomonas (750nm) and *M. fakhimi* monocultures (600nm) were measured as monocultures before centrifugation (A$_{BC}$). Both monocultures were mixed (1:1) and measured again after centrifugation (A$_{AC}$) at a specific centrifugation force and time. Dilutions factors were applied. Error % = 100 − [(A$_{AC}$ · 100)/A$_{BC}$]. This value indicates the percentage of the error obtained in estimating the absorbance of Chlamydomonas or *M. fakhimi* cultures in the mixed cultures compared to their initial monocultures (before mixing). The optimized centrifugation force (x *g*) and time (min) used for the Chlamydomonas and *M. fakhimi* consortium is highlighted in bold. Data represent the means from at least two different biological samples and assayed with technical duplicates.

Author Contributions: Writing—original draft preparation, D.G.-B and N.F..; writing—review & editing, N.F and A.D.; Bacteria isolation, N.F.; design and execution of experiments, N.F.,

M.J.T.; in silico analysis, D.G.-B.; results analysis and interpretation, N.F., D.G.-B.; supervision, A.D. and D.G.-B.; project administration, A.G., E.F., A.D. and D.G.-B.; funding acquisition, A.G., E.F., A.D. and D.G.-B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the European ERANETMED program [ERANETMED2-72-300], the Spanish Ministerio de Ciencia e Innovación [PID2019-105936RB-C22 and TED2021-130438B-I00]; the UCO-FEDER [UCO-1381175], and the Plan Propio of University of Córdoba [MOD.4.1 P.P.2016 A. DUBINI]

Acknowledgments: The authors acknowledge Dr. Gregorio Galvéz Valdivieso for his unvaluable contribution to this research: perfection sometimes kills new discoveries.

Conflicts of Interest: The authors declare no conflict of interest. The sponsors had no role in the design, execution, interpretation, or writing of the study.

References

Aaronson S, Dhawale SW, Patni NJ, Deangelis B, Frank O, Baker H (1977) The cell content and secretion of water-soluble vitamins by several freshwater algae. Arch Microbiol **112**: 57–59

Abadi VAJM, Sepehri M, Rahmani HA, Zarei M, Ronaghi A, Taghavi SM, Shamshiripour M (2020) Role of Dominant Phyllosphere Bacteria with Plant Growth–Promoting Characteristics on Growth and Nutrition of Maize (Zea mays L.). J Soil Sci Plant Nutr. doi: 10.1007/s42729-020-00302-1

Abdin Z, Zafaranloo A, Rafiee A, Mérida W, Lipiński W, Khalilpour KR (2020) Hydrogen as an energy vector. Renewable and Sustainable Energy Reviews. doi: 10.1016/j.rser.2019.109620

Amin SA, Green DH, Hart MC, Küpper FC, Sunda WG, Carrano CJ (2009) Photolysis of iron-siderophore chelates promotes bacterial-algal mutualism. Proc Natl Acad Sci U S A **106**: 17071–17076

Aydin MI, Karaca AE, Qureshy AMMI, Dincer I (2021) A comparative review on clean hydrogen production from wastewaters. J Environ Manage **279**: 111793

Ban S, Lin W, Wu F, Luo J (2018) Algal-bacterial cooperation improves algal photolysis-mediated hydrogen production. Bioresour Technol **251**: 350–357

Burmølle M, Webb JS, Rao D, Hansen LH, Sørensen SJ, Kjelleberg S (2006) Enhanced biofilm formation and increased resistance to antimicrobial agents and bacterial invasion are caused by synergistic interactions in multispecies biofilms. Appl Environ Microbiol. doi: 10.1128/AEM.03022-05

Calatrava V, Hom EFY, Llamas Á, Fernández E, Galván A (2018) OK, thanks! A new mutualism between Chlamydomonas and methylobacteria facilitates growth on amino acids and peptides. FEMS Microbiol Lett **365**: 1–9

Calatrava V, Hom EFY, Llamas Á, Fernández E, Galván A (2019) Nitrogen scavenging from amino acids and peptides in the model alga Chlamydomonas reinhardtii. The role of extracellular L-amino oxidase. Algal Res **38**: 101395

Chaiboonchoe A, Dohai BS, Cai H, Nelson DR, Jijakli K, Salehi-Ashtiani K (2014) Microalgal metabolic network model refinement through high-throughput functional metabolic profiling. Front Bioeng Biotechnol **2**: 1–12

Cho DH, Ramanan R, Heo J, Lee J, Kim BH, Oh HM, Kim HS (2015) Enhancing microalgal biomass productivity by engineering a microalgal-bacterial community. Bioresour Technol **175**: 578–585

Cordovez V, Schop S, Hordijk K, de Boulois HD, Coppens F, Hanssen I, Raaijmakers JM, Carrióna VJ (2018) Priming of plant growth promotion by volatiles of rootassociated Microbacterium spp. Appl Environ Microbiol **84**: 1865–1883

Croft MT, Moulin M, Webb ME, Smith AG (2007) Thiamine biosynthesis in algae is regulated by riboswitches. Proc Natl Acad Sci U S A **104**: 20770–20775

Dahiya S, Chatterjee S, Sarkar O, Mohan SV (2021) Renewable hydrogen production by dark-fermentation: Current status, challenges and perspectives. Bioresour Technol **321**: 124354

Dębowski M, Zieliński M, Kazimierowicz J, Kujawska N, Talbierz S (2020) Microalgae cultivation technologies as an opportunity for bioenergetic system development—advantages and limitations. Sustainability (Switzerland) **12**: 1–37

Degrenne B, Pruvost J, Legrand J (2011) Effect of prolonged hypoxia in autotrophic conditions in the hydrogen production by the green microalga Chlamydomonas reinhardtii in photobioreactor. Bioresour Technol **102**: 1035–1043

Dubini A, Ghirardi ML (2015) Engineering photosynthetic organisms for the production of biohydrogen. Photosynth Res **123**: 241–253

Fabris M, Abbriano RM, Pernice M, Sutherland DL, Commault AS, Hall CC, Labeeuw L, McCauley JI, Kuzhiuparambil U, Ray P, et al (2020) Emerging Technologies in Algal Biotechnology: Toward the Establishment of a Sustainable, Algae-Based Bioeconomy. Front Plant Sci. doi: 10.3389/fpls.2020.00279

Fakhimi N, Dubini A, Tavakoli O, González-Ballester D (2019a) Acetic acid is key for synergetic hydrogen production in Chlamydomonas-bacteria co-cultures. Bioresour Technol **289**: 121648

Fakhimi N, Gonzalez-Ballester D, Fernández E, Galván A, Dubini A (2020) Algae-Bacteria Consortia as a Strategy to Enhance H2 Production. Cells. doi: 10.3390/cells9061353

Fakhimi N, Tavakoli O, Marashi S-AA, Moghimi H, Mehrnia MR, Dubini A, González-Ballester D (2019b) Acetic acid uptake rate controls H2 production in Chlamydomonas-bacteria co-cultures. Algal Res **42**: 101605

Fakhimi N, Torres MJ, Delgado-Luque J, Fernandez E, Galván A, Dubini A, Gonzalez-Ballester D (2023a) Draft genome sequence of Microbacterium fakhimi sp. nov., a novel bacterium associated with the alga Chlamydomonas reinhardtii. Genomics

Fakhimi N, Torres MJ, Dubini A, Gonzalez-Ballester D (2023b) Draft genome sequence of Stenotrophomonas goyi sp. nov., a novel bacterium associated with the alga Chlamydomonas reinhardtii. F1000Res

Fouchard S, Hemschemeier A, Caruana A, Pruvost J, Legrand J, Happe T, Peltier G, Cournac L (2005) Autotrophic and mixotrophic hydrogen photoproduction in sulfur-deprived Chlamydomonas cells. Appl Environ Microbiol **71**: 6199–6205

Ghirardi ML, Dubini A, Yu J, Maness PC (2009) Photobiological hydrogen-producing systems. Chem Soc Rev **38**: 52–61

Harris E (2008) The Chlamydomonas Sourcebook: Introduction into Chlamydomonas and its laboratory use.

Helliwell KE, Wheeler GL, Leptos KC, Goldstein RE, Smith AG (2011) Insights into the evolution of vitamin B 12 auxotrophy from sequenced algal genomes. Mol Biol Evol **28**: 2921–2933

Hemschemeier A, Fouchard S, Cournac L, Peltier G, Happe T (2008) Hydrogen production by Chlamydomonas reinhardtii: an elaborate interplay of electron sources and sinks. Planta **227**: 397–407

Hom EFY, Aiyar P, Schaeme D, Mittag M, Sasso S (2015) A Chemical Perspective on Microalgal-Microbial Interactions. Trends Plant Sci **20**: 689–693

Hwang JH, Church J, Lim J, Lee WH (2018) Photosynthetic biohydrogen production in a wastewater environment and its potential as renewable energy. Energy **149**: 222–229

Ji X, Li H, Zhang J, Saiyin H, Zheng Z (2019) The collaborative effect of Chlorella vulgaris - Bacillus licheniformis consortia on the treatment of municipal water. J Hazard Mater **365**: 483–493

Jones KJ, Moore K, Sambles C, Love J, Studholme DJ, Aves SJ (2016) Draft genome sequences of Achromobacter piechaudii GCS2, Agrobacterium sp. strain SUL3, Microbacterium sp. strain GCS4, Shinella sp. strain GWS1, and Shinella sp. strain SUS2 isolated from consortium with the hydrocarbon-producing alga Botryococcus brauni. Genome Announc. doi: 10.1128/genomeA.01527-15

Jurado-Oller JL, Dubini A, Galván A, Fernández E, González-Ballester D (2015) Low oxygen levels contribute to improve photohydrogen production in mixotrophic non-stressed Chlamydomonas cultures. Biotechnol Biofuels **8**: 1–14

Kakoulaki G, Kougias I, Taylor N, Dolci F, Moya J, Jäger-Waldau A (2021) Green hydrogen in Europe – A regional assessment: Substituting existing production with electrolysis powered by renewables. Energy Convers Manag. doi: 10.1016/j.enconman.2020.113649

Kim BH, Ramanan R, Cho D-HH, Oh HM, Kim HS (2014) Role of Rhizobium, a plant growth promoting bacterium, in enhancing algal biomass through mutualistic interaction. Biomass Bioenergy **69**: 95–105

Kruse O, Rupprecht J, Bader K, Thomas-hall S, Schenk PM, Finazzi G, Hankamer B (2005) Improved Photobiological H 2 Production in Engineered Green Algal Cells *. **280**: 34170–34177

Lakatos G, Deák Z, Vass I, Rétfalvi T, Rozgonyi S, Rákhely G, Ördög V, Kondorosi É, Maróti G (2014) Bacterial symbionts enhance photo-fermentative hydrogen evolution of Chlamydomonas algae. Green Chemistry **16**: 4716–4727

Leong HY, Chang CK, Khoo KS, Chew KW, Chia SR, Lim JW, Chang JS, Show PL (2021) Waste biorefinery towards a sustainable circular bioeconomy: a solution to global issues. Biotechnol Biofuels. doi: 10.1186/s13068-021-01939-5

Lepage T, Kammoun M, Schmetz Q, Richel A (2021) Biomass-to-hydrogen: A review of main routes production, processes evaluation and techno-economical assessment. Biomass Bioenergy. doi: 10.1016/j.biombioe.2020.105920

Li X, Huang S, Yu J, Wang Q, Wu S (2013) Improvement of hydrogen production of *Chlamydomonas reinhardtii by co-cultivation with isolated bacteria. Int J Hydrogen Energy 38: 10779–10787*

*Loppes R, Radoux M, Ohresser MCP, Matagne RF (1999) Transcriptional regulation of the Nia1 gene encoding nitrate reductase in Chlamydomonas reinhardtii: Effects of various environmental factors on the expression of a reporter gene under the control of the Nia1 promoter. Plant Mol Biol **41**: 701–711*

Lorincz Z, Preininger É, Kósa A, Pónyi T, Nyitrai P, Sarkadi L, Kovács GM, Böddi B, Gyurján I (2010) Artificial tripartite symbiosis involving a green alga (Chlamydomonas), a bacterium (Azotobacter) and a fungus (Alternaria): Morphological and physiological characterization. Folia Microbiol (Praha). doi: 10.1007/s12223-010-0067-9

Marquez-Santacruz HA, Hernandez-Leon R, Orozco-Mosqueda MC, Velazquez-Sepulveda I, Santoyo G (2010) Diversity of bacterial endophytes in roots of Mexican husk tomato plants (Physalis ixocarpa) and their detection in the rhizosphere. Genet Mol Res **9**: 2372–2380

Martelli F, Marrella M, Lazzi C, Neviani E, Bernini V (2021) Microbiological contamination of ready-to-eat algae and evaluation of bacillus cereus behavior by microbiological challenge test. J Food Prot. doi: 10.4315/JFP-20-407

Meier-Kolthoff JP, Göker M (2019) TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. Nat Commun. doi: 10.1038/s41467-019-10210-3

Mitra M, Nguyen KMAK, Box TW, Berry TL, Fujita M (2021) Isolation and characterization of a heavy metal- and antibiotic-tolerant novel bacterial strain from a contaminated culture plate of Chlamydomonas reinhardtii, a green micro-alga. F1000Res **10**: 1–34

Mora-Salguero D, Vives Florez MJ, Husserl Orjuela J, Fernández-Niño M, González Barrios AF (2019) Evaluation of the phenol degradation capacity of microalgae-bacteria consortia from the bay of Cartagena, Colombia. TecnoLógicas **22**: 149–158

Oey M, Ross IL, Stephens E, Steinbeck J, Wolf J, Radzun KA, Kügler J, Ringsmuth AK, Kruse O, Hankamer B (2013) RNAi Knock-Down of LHCBM1, 2 and 3 Increases Photosynthetic H2Production Efficiency of the Green Alga Chlamydomonas reinhardtii. PLoS One **8**: e61375

Ramanan R, Kim BH, Cho DH, Oh HM, Kim HS (2016) Algae-bacteria interactions: Evolution, ecology and emerging applications. Biotechnol Adv **34**: 14–29

Rosenblueth M, Martínez-Romero E (2006) Bacterial endophytes and their interactions with hosts. Molecular Plant-Microbe Interactions **19**: 827–837

Scoma A, Durante L, Bertin L, Fava F (2014) Acclimation to hypoxia in Chlamydomonas reinhardtii: can biophotolysis be the major trigger for long-term H2 production? New Phytologist **204**: 890–900

Shetty P, Boboescu IZ, Pap B, Wirth R, Kovács KL, Bíró T, Futó Z, White RA, Maróti G (2019) Exploitation of Algal-Bacterial Consortia in Combined Biohydrogen Generation and Wastewater Treatment. Front Energy Res **7**: 52

Singh T, Singh DK (2019) Rhizospheric Microbacterium sp. P27 Showing Potential of Lindane Degradation and Plant Growth Promoting Traits. Curr Microbiol **76**: 888–895

Steinbeck J, Nikolova D, Weingarten R, Johnson X, Richaud P, Peltier G, Hermann M, Magneschi L, Hippler M (2015) Deletion of Proton Gradient Regulation 5 (PGR5) and PGR5-Like 1 (PGRL1) proteins promote sustainable light-driven hydrogen production in Chlamydomonas reinhardtii due to increased PSII activity under sulfur deprivation. Front Plant Sci **6**: 892

Teplitski M, Rajamani S (2011) Signal and Nutrient Exchange in the Interactions Between Soil Algae and Bacteria. Springer, Berlin, Heidelberg, pp 413–426

Tolleter D, Ghysels B, Alric J, Petroutsos D, Tolstygina I, Krawietz D, Happe T, Auroy P, Adriano JM, Beyly A, et al (2011) Control of hydrogen photoproduction by the proton gradient generated by cyclic electron flow in Chlamydomonas reinhardtii. Plant Cell **23**: 2619–2630

Torres MJ, González-Ballester D, Gómez-Osuna A, Galván A, Fernández E, Dubini A (2022) *Chlamydomonas-Methylobacterium oryzae cooperation leads to increased biomass, nitrogen removal and hydrogen production. Bioresour Technol. doi: 10.1016/j.biortech.2022.127088*

Torzillo G, Scoma A, Faraloni C, Ena A, Johanningmeier U (2009) *Increased hydrogen photoproduction by means of a sulfur-deprived Chlamydomonas reinhardtii D1 protein mutant. Int J Hydrogen Energy* **34**: 4529–4536

Vogel SL, Frisch HL, Gotham IJ (1978) *QUALITATIVE ASSAY OF DISSOLVED AMINO ACIDS AND SUGARS EXCRETED BY CHLAMYDOMONAS REINHARDTII (CHLOROPHYCEAE) AND EUGLENA GRACILIS (EUGLENOPHYCEAE). J Phycol. doi: 10.1111/j.1529-8817.1978.tb02459.x*

Wintermans JFGM, De Mots A (1965) *Spectrophotometric characteristics of chlorophylls a and b and their phenophytins in ethanol. Biochimica et Biophysica Acta (BBA) - Biophysics including Photosynthesis* **109**: 448–453

Wu S, Li X, Yu J, Wang Q (2012) *Increased hydrogen production in co-culture of Chlamydomonas reinhardtii<i/> and Bradyrhizobium japonicum<i/>. Bioresour Technol* **123**: 184–188

Xu L, Li D, Wang Q, Wu S (2016) *Improved hydrogen production and biomass through the co-cultivation of Chlamydomonas reinhardtii and Bradyrhizobium japonicum. Int J Hydrogen Energy* **41**: 9276–9283

Yao S, Lyu S, An Y, Lu J, Gjermansen C, Schramm A (2019) *Microalgae–bacteria symbiosis in microalgal growth and biofuel production: a review. J Appl Microbiol* **126**: 359–368

△ Alga TAP	▲ Cocultures TAP	▲ IBI TAP	▽ Alga TM
□ Alga TY	■ Cocultures TY	■ IBI TY	▼ Cocultures TM
			○ Alga TYM
			● Cocultures TYM
			▼ IBI TM
			● IBI TYM