Cities as parasitic amplifiers? Malaria prevalence and diversity in

great tits along an urbanization gradient

Authors

Aude E. Caizergues[1,2]†*, Benjamin Robira[3]†, Charles Perrier[4], Mélanie Jeanneau[1], Arnaud

Berthomieu[1], Samuel Perret[1], Sylvain Gandon[1] & Anne Charmantier[1]

[1]CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

[2] Department of Biology, University of Toronto Mississauga; Mississauga, ON, Canada

[3]Animal Ecology Unit, Research and Innovation Centre, Fondazione Edmund Mach,

San Michele all' Adige, TN, Italy

[4] UMR CBGP, INRAE, CIRAD, IRD, Institut Agro, Université Montpellier, Montpellier, France

† These authors contributed equally to this work.

* Corresponding author

Corresponding author

Aude E. Caizergues : audeemiliecaizergues@gmail.com

ORCID

Aude E. Caizergues : 0000-0003-4467-3912

Benjamin Robira : 0000-0002-3168-6573

Charles Perrier : 0000-0001-5820-9374

Sylvain Gandon : 0000-0003-2624-7856

Anne Charmantier : 0000-0002-0691-2647

22
23

ABSTRACT

Urbanization is a worldwide phenomenon that modifies the environment. By affecting the reservoirs of pathogens and the body and immune conditions of hosts, urbanization alters the epidemiological dynamics and diversity of diseases. Cities could act as areas of pathogen dilution or amplification, depending on whether urban features have positive or negative effects on vectors and hosts. In this study, we focused on a host species and investigated the prevalence and diversity of avian malaria parasites (*Plasmodium/Haemoproteus* sp. and *Leucocytozoon* sp.) in great tits (*Parus major*) living across an urbanization gradient. In general, we observed high prevalence in adult birds (from 95% to 100%), yet lower prevalence in fledglings (from 0% to 38%). We found a slight tendency for increased Plasmodium sp. prevalence with increasing urbanization in adults. Urban nestlings had higher *Plasmodium* sp. infection rates than non-urban nestlings. We found evidence of higher diversity of parasites in the most natural urban park; however, parasite diversity was similar across other urbanization levels (e.g. from a little artificialized park to a highly anthropized industrial area). Parasite lineages were not habitat specific. Only one *Plasmodium* sp. lineage (YWT4) was associated with urban areas and some rare lineages (e.g., AFR065) were present only in a zoo area, perhaps because of the presence of African birds. This study suggests that urbanization can lead to a parasite amplification effect and can favor different avian malaria lineages.

KEYWORDS: avian malaria, epidemiology, diversity, parasite, *Parus major*, prevalence, urbanization, south of France

INTRODUCTION

Urbanization is a worldwide phenomenon driving environmental change and leading to the emergence of artificial habitats (Marzluff 2001; Gaston et al. 2015). Urban areas are a combination of remnant natural habitats and a complex assemblage of anthropogenic perturbations. They are characterized by new environmental conditions such as high levels of chemical, light, and sound pollution, increased impervious surfaces, and altered vegetation communities dominated by exotic plants (Forman and Godron 1986). Such extensive habitat modifications affect biodiversity at multiple ecological levels, from individual phenotypes to community assemblages. Notably, some species thrive in cities while others are not able to cope with urban conditions. Hence, urban communities are altered and mainly composed of fewer, often generalist and/or exotic, species with higher population densities compared to natural habitats (Shochat et al. 2006; Faeth et al. 2011).

Urbanization not only impacts individual species but also species interactions (Faeth et al. 2011), which can affect species evolution (Ots and Hõrak 1998; Marzal et al. 2005; Dyrcz et al. 2005). In particular, host-parasite interactions can be altered in urban habitats (Martin and Boruta 2013; Becker et al. 2015) because of variation in both the occurrence and abundance of species enabling the spread and transmission of the parasite (i.e., the vector species) (Reyes et al. 2013; Giraudeau et al. 2014; Neiderud 2015), changes in vectors' feeding preferences in urban areas (Santiago-Alarcon et al. 2012; Abella-Medrano et al. 2018), and shifts in body condition and immune system efficiency of host species (Bailly et al. 2016; Capilla-Lasheras et al. 2017; Partecke et al. 2020). Depending on the positive and/or negative impact on the vector and host species, the effect of urbanization on disease prevalence can be twofold. First, in cases where urbanization predominantly negatively impacts vector species and/or predominantly favors the host species (e.g., if environmental requirements for parasite development are not met, Calegaro-Marques and Amato 2014), urban areas may act as a parasite dilution factor and urban animal populations should face lower risks of infections compared to their non-urban counterparts (Geue and Partecke 2008; Evans et al. 2009). Second, if the host species is predominantly negatively impacted by the urban conditions (e.g., immune depression in the host species, Bailly et al. 2016) urban individuals may suffer higher parasite burdens due to an amplification effect (e.g., Bichet et al. 2013).

Empirical evidence support both of these two scenarios, revealing case- and host-species

dependence (Evans et al. 2009; Belo et al. 2011; Bichet et al. 2013b; Santiago-Alarcon et al. 2018).

This might be because of the binary view of comparing urban *versus* non-urban habitats, with the

postulate that the urban and non-urban environments stand as homogeneous and dichotomic

environments. Yet, at a finer resolution, the urban matrix consists of a heterogeneous mosaic of local

environments, some of which might be covered by impervious surfaces that contrast with green

spaces. For example, parks offer great potential for multiple species to be supported (Nielsen et al.

2014; Lepczyk et al. 2017), sometimes leading to more diverse and species-rich areas than in nearby

wild habitats (McKinney 2008). It therefore seems necessary to move from a binary perspective (i.e.

the comparison between urban and non-urban habitats) to the study of a continuous urbanization

gradient (e.g., French et al. 2008). Despite the growing body of literature on host-parasite interactions

in urban habitats, their variations along an urbanization gradient are still poorly understood (Bradley

and Altizer 2007; Delgado-V. and French 2012; Ferraguti et al. 2020).

Avian malaria parasites belong to *Haemoproteus*, *Plasmodium*, or *Leucocytozoon* genera and

are widely studied in the context of host-parasite interactions (Rivero and Gandon 2018). Avian

haemosporidians are ubiquitous parasites and encompass a vast diversity of species and strains that

can be generalists, and infect a broad number of bird species (e.g. SGSI *Plasmodium relictum* strain),

or specialist, and infect only one or few species (Valkiunas, 2004). They are vector-borne parasites

infecting blood cells and mainly transmitted by five families of Diptera insects: *Culicidae*,

Hippoboscidae, *Simuliidae*, *Ceratopogonidae*, and *Psychodidae* (Valkiunas and Iezhova 2018). These

vectors are frequently encountered both in non-urban and urban areas, although their diversity and

richness varies with habitat (Coene 1993). Indeed, the presence of water sources (river or pond) in

urban areas is important for vector reproduction and population survival (Asghar et al. 2011). Among

these vectors, some are known to be generalists and to feed on several vertebrate groups, especially in

urban habitats (Jansen et al. 2009).

Great tits are common birds in Eurasia and are abundant in a wide range of habitats, from

natural forests to heavily urbanized city centers (Fink et al. 2022). They are a good model species for

ecologists and evolutionary biologists because they nest in human-provided nest boxes and are easy to

capture and manipulate. Infection by avian malaria in Passeriformes is known to often induce an

increase in immune response, lower survival, and reduced reproductive success (Ots and Hõrak 1998;

Hõrak et al. 2001; Asghar et al. 2011; Lachish et al. 2011; Christe et al. 2012; Pigeault et al. 2018) ;

therefore, if host-parasite interactions are affected by urbanization levels, the outcome for bird

populations could depend on their habitat preference along the urban gradient.

In this study, we investigated the prevalence and diversity of avian malaria parasites in great

tits (*Parus major*) in and around the city of Montpellier, south of France. Specifically, we (1)

compared the prevalence in nestlings and adult individuals across different urbanization levels

measured at the different scales, (2) characterized parasite molecular lineage richness and diversity

along the gradient of urbanization, and (3) assessed the role of urbanization levels on parasite

diversity.

METHODS

Study sites along an urbanization gradient

We studied nest boxes at two anthropogenically contrasted areas that had different levels of urban

impacts. First the city of Montpellier, in southern France (43°36′N 3°53′E), which is a metropolitan

area hosting 480,000 inhabitants. Second, the Rouvière oak forest, which is located 20 km northwest

of Montpellier (Figure 1). In these city and forest contexts (hereafter urban and non-urban,

respectively), long-term monitoring programs of the breeding populations of great tits have been

conducted since 2011 and 1991, respectively (Charmantier et al. 2017). Monitoring consists of weekly

visits mid-March to mid-July to document great tit reproduction in artificial nest boxes scattered in

eight sites across the city (222 nest boxes) (Figure 1, see Figure S1 for a satellite image of the Zoo

site) and across the forest of La Rouvière (94 nest boxes). The climate is typically Mediterranean, with

mild winters and dry summers. Spring is marked by a sudden rise in temperature, coinciding with the

great tit breeding season. This region of France hosts high densities of avian malaria *Plasmodium*

vectors such as *Culex pipiens*, for which massive insecticide-based control treatments have been

deployed for more than 60 years, with a focus on coastal areas (i.e., ca 15 km from Montpellier

historical center, EID, 2020).

We characterized the level of urbanization and anthropogenic disturbance around each nest

box, considering the area defined by a 50 m circular buffer around each nest-box where parents and

nestlings were captured and sampled. This area is typically considered representative of a breeding

great tit foraging area (Perrins 1979). We quantified four environmental features relevant for great tits

breeding performances and fitness: (1) the extent of the vegetation cover (reflecting abundance of

resources), (2) the motorized traffic disturbance (reflecting background noise pollution and chemical

pollution), (3) the pedestrian disturbance (reflecting direct human disturbance), and (4) the amount of

light pollution (affecting birds' circadian rhythm, immunity and behavior). We measured the surface

of vegetation cover (canopy and grass) around each nest box based on satellite images from Google

maps. We quantified the motorized traffic perturbation by counting the number of motorized engines

passing in the area during a 5 min count performed for each box in the early morning (between 7am

and 11am). This count showed a 0.85 Pearson correlation with traffic data provided by the city of

Montpellier (opendata.montpelliernumerique.fr/) in a given area (Demeyrier et al. 2016). We similarly

estimated pedestrian disturbance with counts of pedestrians, bikes, and scooters. Finally, we defined

local light pollution as the area covered by artificial light from lamp posts, assuming that a lamp post

would illuminate a circular area of 50 m from its location. We summarized those four metrics using a

principal component analysis as in Caizergues et al. (2021) to describe urbanization and disturbance at

the nest level along two composite measures (Figure S2). In brief, we retained the two main axes

explaining 67.8% of the variation in urban features, from which we used only the first axis in the

present study. This first axis explained 42.4% of variance and was defined as the "naturalness"

gradient, with positive values associated with larger vegetation cover, lower traffic disturbance, and

lower light pollution. The second axis, defined as the "pedestrian frequency" gradient (25.4% of

variance explained), was not used in the current study since it was not correlated with the habitat

artificialization of an area but rather logically to the number of pedestrians passing by each nest box.

We obtained site-level measures of "naturalness" for the eight urban sites and La Rouvière (sites

hereafter referred to by acronyms made of their first three letters, see Table S1) by averaging these

composite measures considering all nest boxes within a given site. This ranged from the most

"natural" site, La Rouvière (ROU), to the most urbanized one, Mas Nouguier (MAS) in Montpellier

city. Among urban sites, the zoo (ZOO; Figure 1C) was the least urbanized site, but was well settled in

the urban matrix (Figure S1), with short distances from CEF and FAC allowing great tits from the zoo

to interact with birds in these neighboring sites (see low genetic differentiation described in Perrier et

al. 2017).

Serologic sampling and molecular analyses

Blood sample collection

Between 2014 and 2019, we collected serologic samples between mid-March and mid-July. Samples

were collected from 15 days old nestling and adult great tits across the urban and non-urban sites. We

captured the parents when nestlings were 10-15 days old using traps inside nest boxes. All nestlings

and adults were uniquely identified with rings provided by the Centre de Recherches sur la Biologie

des Populations d'Oiseaux (CRBPO, Paris, France). We had a total of 296 adults (154 females 142

males) and 90 nestlings (not sexed and all sampled in 2014), see sampling detail Table S2.

170   We collected 10 µL of blood by performing a venipuncture in either the ulnar (i.e., wing) vein

or a small subepidermal neck vein. We transferred blood samples using a capillary into an Eppendorf

filled with 1 mL of Queen's lysis buffer, then stored in 4°C refrigerators at the end of the field day

until DNA extraction.

DNA extraction

We extracted total genomic DNA from blood samples using the DNeasy Blood and Tissue kit

(Qiagen). We adapted the standard protocol by mixing 500 µL of solution of blood and Queen's buffer

(~1/100 of blood) with 40 µL of proteinase K and 250 µL AL buffer. We then incubated the mixture at

56°C for 1.5 h. Afterwards, we added 8 µL of RNase A (100 mg/ml). We then performed DNA

precipitation by adding 400 µL of ethanol.

Figure 1: Maps of the sampling locations A) at European scale, B) at regional scale and C) at the city
scale, where each polygon represents the limits of an urban sampling site and the colour represents the
naturalness score of the site.

Infection detection

We detected and identified parasites adapting Hellgren et al. (2004) protocol. We first amplified

possible large fragments of mtDNA from *Plasmodium* spp., *Haemoproteus* spp. and *Leucocytozoon*

spp. using polymerase chain reaction (PCR) with the HaemNF, HaemNR2 and HaemNR3 primers.

PCR conditions included 15 min at 94°C, followed by 25 cycles of 30 s at 94°C, 40 s at 50°C, 1 min at

72°C, and a last cycle of 10 min at 60°C. Using 1 µL from the first amplified reaction, we then

performed a secondary and more specific PCR to separately identify *Leucocytozoons* spp. and

Plasmodium-Haemoproteus spp. presence with two different sets of primers: (i) we used the HaemF/

HaemR2 primers to amplify *Plasmodium* spp. And *Haemoproteus* spp. (test PH); (ii) and

HaemFL/HaemRL primers to amplify *Leucocytozoon* spp. (test L). We performed this second PCR

using Multiplex PCR kit Qiagen in a final volume of 10 µL following one cycle of 15 min at 94°C, 35

cycles of 30 s at 94°C, 40 s at 51°C/52°C (for *Leucocytozoon* spp.*/Plasmodium* spp. or *Haemoproteus*

spp., respectively), 1 min at 72°C and one last cycle of 10 min at 60°C. We assessed amplification in

2% agarose gels leading to four possible infection outcomes: (1) uninfected (negative test PH and L),

(2) infected by *Plasmodium* spp. and/or *Haemoproteus* spp. (positive test PH, negative test L)*,* (3)

infected by *Leucocytozoon* spp. (positive test L, negative test PH), and (4) coinfected by *Plasmodium*

spp. and/or *Haemoproteus* spp. and *Leucocytozoon* spp. (positive test PH and L).

Lineage identification

We sent positive samples to Eurofins Genomics Company for Sanger sequencing. We then blasted

sequences against the MalAvi database for molecular lineage identification (Bensch et al. 2009). We

identified single and multiple infections of *Plasmodium* sp. and *Haemoproteus* sp. In contrast, the

Leucocytozoon sp. sequencing quality was poor (i.e., there was an uncertain multiple-base identity in

the sequence), and we were unable to identify a unique lineage (i.e., 100% blast score with a sequence

from the database) for each sample. Therefore, we only identified a set of 5 likely lineages (blast

>96%) for each sample. As no infection by any parasite from *Haemoproteus* genus was detected in our

samples, we hereafter refer to *Plasmodium* sp. only.

Statistical analyses

We performed all analyses with *R* software (version 4.2.1, R Core Team 2022). A complete list of the packages, associate versions, and reference used for data processing, analyses, and plotting, are provided in Supplementary Material Table S32.

Quantifying parasitic prevalence at the different sites

We estimated the site-level prevalence in nestlings and adults of *Plasmodium* sp. and *Leucocytozoon* sp. as the proportion of infected individuals as well as their 95% confidence intervals (Wilson score interval, "propCI" function of the *prevalence* package).

To further assess the role of urbanization in shaping prevalence patterns, we ran linear models separately on nestlings and adult individuals, and for the different parasite genera (*Plasmodium* sp. and *Leucocytozoon* sp., respectively*)*, to link infection probability to the urban context across different spatial scales: the site level (i.e., average urbanization level of around all nests from a given site) and the local level (i.e., the urbanization level around the nest). To ease comparability with previous studies, we also carried out the analyses considering the habitat along the urban vs non-urban dichotomy (in this case the site and nest level always matched in their classification). To do so, we fitted three logistic regressions ("glm" and "glmer" function with a log-link function *stat* R package, Bates et al. 2015) with a binary response of infection (0 as not infected, 1 as infected) as a function of either habitat type (binary variable, 0 as non-urban, 1 as urban), the site-level naturalness (first axis of the PCA averaged on all the nest boxes of a sample site, see above), or the local nest-level naturalness (per nest-box first axis of the PCA value, and site as a random effect when the analysis was conducted considering nest-scale urbanization). For models ran on data from adult individuals, we further controlled for sex, age (in years), as well as the year of sampling. Because of the low number of samples in years 2017 (N = 6) and 2018 (N = 18), we removed these data from analysis to limit model convergence issues. We assessed the significance of each predictor using likelihood ratio tests ("drop1" function of the *stats* package) while dropping one predictor at a time.

We verified that linear models' assumptions were not violated using various visual controls of

residual distributions and associated statistical tests (histogram of residuals, Q-Q plot of expected

residuals vs observed residuals, scattered plot of residuals vs estimates) using the *DHARMa* package

as well as the *performance* package (see Supplementary Material: Supplementary text 1, Tables S2 to

S31 and Figures S2 to S24). This raised no problem of collinearity, singular fit, convergence, or

influential points.

Characterising lineage diversity and habitat specificity at the different sites

For subsequent analyses, we focused on adult individuals, as the quality of nestling malaria sequences

was low and prevented us from correctly identifying lineages. Given the uncertainty in the

Leucocytozoon sp. lineage identification (i.e., only a subset of likely lineages could be identified), we

repeated the analyses (below) 1000 times for this parasite genus, each iteration randomly sampling a

unique lineage (out of the 5 identified lineages) per individual. Thus, for *Leucocytozoon* sp. we

provide the median estimates and associated 95% confidence intervals.

Lineage diversity

Haemosporidian lineages richness and abundance were analyzed with the *vegan* and *BiodiversityR*

packages. To analyze patterns of lineage diversity per site, we estimated lineage richness and the

combination between richness and evenness using the Shannon (roughly giving more weight to rare

species) and inverse-Simpson (roughly giving more weight to common species) diversity indices. We

also plotted the rank abundance curves for each study site, which highlight the richness (absolute value

in the curve) and the evenness (slope of the curve) of parasite assemblages (Nagendra 2002).

We estimated dissimilarities in lineage composition between sites using the Bray-Curtis

dissimilarity index ("vegdist" function of the *vegan* package). We computed this index on the binary

sequence (i.e., indicating whether a given lineage was present or absent), and on the sequence of

individual prevalence for each lineage (i.e., percent of infected individuals having the lineage). The

former would provide insight into parasite composition resemblance (hereafter Bray-Curtis

composition) and the latter into prevalence resemblance (hereafter Bray-Curtis prevalence).

Habitat specificity

To investigate whether some lineages occurred more frequently than randomly expected in urban

versus non-urban environment, we compared the proportion of urban nest boxes at which each lineage

was present to the overall number of nest boxes sampled using a binomial test ("binom.test" function).

To avoid false negative habitat association in rare lineages, we computed the test only for lineages for

which the type II error was below 0.20 (Cohen, 2013), and that occurred at least 10 times overall.

In addition, we investigated if parasitic community similarity was linked to urbanization at

two scales: the sampling site and the nest box. We analyzed the correlation between naturalness

distance (absolute difference in "naturalness" level) and parasite dissimilarity (Bray-Curtis

composition) matrices using a mantel test with 999 permutations ("mantel.test" function of the *ape*

package). We also controlled for spatial autocorrelation by testing whether parasitic community

similarity was related to geographic proximity, repeating those analyses comparing the Euclidean

distance between pairs of sites or nest boxes ("st_distance" function of the *sf* package) to parasite

dissimilarity.

RESULTS

***Plasmodium*, *Haemoproteus* and *Leucocytozoon* prevalences**

Parasitic prevalence in nestlings

In 15-day-old nestlings, avian haemosporidian prevalence was < 40% in both habitats, with some

heterogeneity among sites (Figure 2A). No nestling was simultaneously infected by *Plasmodium* sp.

and *Leucocytozoon* sp. parasites.

The prevalence in *Plasmodium* parasites ranged from 0% to 38%, with an average of 16.33%

(Figure 2A). Prevalence was significantly higher in the urban nestlings compared to non-urban

nestlings (16.67% averaged on all urban sites vs. 0% in the non-urban site; χ^2_1 = 9.854, P = 0.002).

However, overall small sample sizes and the presence of only one replicate of "non-urban" site

potentially inflated these differences.). Plasmodium prevalence was not related to the nest- and site-

level naturalness gradient ($\chi^2_1 = 0.012$, P = 0.908; $\chi^2_1 = 1.186$, P = 0.276, respectively).

The prevalence in *Leucocytozoon* sp. ranged from 0% to 40%, with an average of 9.90% and

did not substantially differ consistently between urban and non-urban nestlings (11.11% averaged on

all urban sites vs. 2.78% in the non-urban site; $\chi^2_1 = 2.383$, P = 0.123). *Leucocytozoon* sp. prevalence

was unrelated to the nest- or site-level naturalness gradient ($\chi^2_1 = 1.837$, P = 0.175; $\chi^2_1 = 1.291$, P =

0.256, respectively).

Parasitic prevalence in breeding individuals

Avian haemosporidian prevalence ranged from 95% to 100% for *Plasmodium* sp. (mean = 97.04%),

and 80% to 100% for *Leucocytozoon* sp. in breeding great tits, (mean = 92.93%) (Figure 2). Double

infection was frequent (91.9% of individuals). In particular, all individuals infected with

Leucocytozoon sp. were systematically infected with *Plasmodium* sp.

Prevalence of *Plasmodium* sp. and *Leucocytozoon* sp. did not vary significantly between urban and

non-urban sites ($\chi^2_1 = 0.003$, P = 0.955; $\chi^2_1 = 1.71$, P = 0.191, respectively) nor with the site-level

naturalness ($\chi^2_1 = 0.360$, P = 0.548; $\chi^2_1 = 0.012$, P = 0.911, respectively). However, nest-level

naturalness gradient was potentially related to *Plasmodium* sp. prevalence (glmer: est. ± S.E. = -0.616

± 0.397, $\chi^2_1 = 2.937$, P = 0.087), with a weak tendency for lower prevalence in less urbanized areas. In

contrast, *Leucocytozoon* prevalence was not related to the nest-level naturalness gradient ($\chi^2_1 = 0.567$,

P = 0.452). In addition, prevalence of both parasites genera did not vary between males and females

(all P >> 0.05) or with age (all P >> 0.05). *Leucocytozoon* prevalence models showed a significant

year effect when urbanization was considered dichotomous (glm: est. ± S.E. = 1.051 ± 0.484, $\chi^2_1 =$

5.075, P = 0.024), with greater prevalence in 2019 compared to 2014. In contrast, *Plasmodium* sp.

prevalence did not vary by year (all P >> 0.05).

Figure 2: Mean prevalence of avian *Plasmodium* sp. (light grey) and *Leucocytozoon* sp. (dark grey) per site in great tit (A) nestlings and (B) adults. Error bars represent 95% confidence intervals. Sites are ordered by increasing urbanization level and sample size is detailed below each site.

Prevalence in nestlings versus breeding individuals

Plasmodium sp. and *Leucocytozoon* sp. prevalence at each site were not correlated between nestling and adult stages (Spearman correlation test, ⍴ = 0.133, P = 0.803 and ⍴ = -0.577, P = 0.231, respectively).

Parasite molecular lineage diversity

A combination of 47 lineages of *Plasmodium* spp. and *Leucocytozoon* spp. species were recorded across all study sites (Figure 3), including 5 *Plasmodium* spp. and 42 *Leucocytozoon* spp. (total number of lineages identified by BLAST, not accounting for uncertainty in lineage identification). The *Plasmodium* sp. lineage SGS1 was the most represented of all lineages, with 272 infected birds out of 296 individuals sampled. Comparisons of lineage diversity depended on how diversity was quantified.

The least urbanized urban site (ZOO) had the highest richness and Shannon's evenness (richness = 22,

evenness = 2.20, Table 1). The non-urban site (ROU) had intermediate richness (richness = 16) and

was among the lowest in terms of Shannon's evenness (evenness = 1.79). In contrast, FAC had the

highest inverse Simpson's diversity (Simpson's index = 4.98). The non-urban site ROU had the lowest

inverse Simpson's diversity (Simpson's index = 3.08, Table 1). Overall, diversity patterns among site

were similar, , with all sites having low evenness in parasite types which systematically consisted of a

small subset of the 47 lineages (Figure 4).

Habitat specificity of lineages in breeders

Regarding lineage habitat specificity, we found one lineage, YWT4 (*Plasmodium* sp.), that occurred

more in urban habitats than expected by chance (Figure 5). None of the other *Plasmodium* sp. or

Leucocytozoon sp. lineages were statistically more associated with one habitat type than the other.

Resemblances between sites were globally homogenous between pairs of sites (Figure 6), both in

composition (i.e., in terms of lineage diversity) and prevalence (i.e., in terms of infection rate for a

given lineage). Anecdotically, BOT and CEF, the smallest and least sampled sites, were the most

dissimilar to other sites (Figure 6).

Table 1: Haemosporidian lineages richness and diversity indices (Shannon and Inverse Simpson)
across the eight urban sites and the non-urban site (ROU).

Site	N	Naturalness index	Richness	Shannon	Inverse Simpson
MAS	41	-2.383	18 (15 - 20)	1.98 (1.88 - 2.06)	3.65 (3.56 - 3.72)
MOS	37	-0.865	16 (14 - 19)	2.09 (1.99- 2.17)	4.53 (4.41 - 4.61)
FON	40	-0.854	18 (16 - 21)	2.09 (2.00 - 2.17)	4.11 (4.02 - 4.17)
FAC	21	-0.750	15 (13 - 17)	2.14 (2.01 - 2.23)	4.98 (4.77 - 5.13)
BOT	12	-0.406	9 (7 - 11)	1.71 (1.54 - 1.84)	3.51 (3.33-3.64)
GRA	45	0.254	16 (13 - 18)	1.86 (1.76 - 1.94)	3.33 (3.25 - 3.38)
CEF	8	0.458	8 (6 - 9)	1.71 (1.49 - 1.80)	3.81 (3.46 - 3.95)
ZOO	41	0.687	22 (19 - 25)	2.20 (2.10 - 2.28)	4.11 (4 - 4.17)
ROU	52	1.221	16 (14 - 19)	1.79 (1.69 - 1.88)	3.08 (3.02 - 3.13)

Figure 3: Proportions of (A) *Plasmodium* sp. and (B) *Leucocytozoon* sp. lineages found in each study site. For *Leucocytozoon* sp., only the most abundant lineages are shown in detail and lineages with less than 15 total occurrences were grouped as "other".

Figure 4: Rank-abundance curves for avian Haemosporidian lineages in each urban site and a non-urban site. Abundance is defined as the prevalence of a lineage at a given site. The *x*-axis represents the rank-abundance. The shape of the curve highlights the evenness: the steeper the curve, the less even distribution of lineage abundance. A flat curve indicates an evenly distributed community).

Figure 5: Occurrence probability of avian Haemosporidian lineages in the urban habitat for A)
Plasmodium sp. and B) *Leucocytozoon sp*. Error bars represent 95% confidence intervals. The dashed
line represents the expected probability of occurrence of a lineage in the urban habitat under random
distribution. Grey dots and error bars represent lineages that are found statistically more in the urban
habitat, and black, lineages that are not habitat-specific.

Figure 6: Heatmap of Bray-Curtis dissimilarity between each site considering binary sequences of
lineage composition (bottom) or the prevalence of each lineage among infected individuals (top).
Darker colors represent higher values of Bray Curtis index and stronger differences in lineage
composition/prevalence between a pair of sites. Values in the cells indicate the upper border of the
95% confidence interval.

We found no statistical link between parasitic community similarity and naturalness gradient

or geographical proximity at both the site or the nest box levels (Mantel test: P >> 0.05 for all the 1000

subsampled datasets; p-values were adjusted to maintain the false discovery rate to 5%).

DISCUSSION

In this study, we investigated the link between urbanization and avian malaria prevalence and lineage

diversity at different scales across wild populations of great tits in and around Montpellier, a

metropolis of almost half a million inhabitants in southern France. We found marked differences in

parasite prevalence between life stages, with 15-day-old nestlings showing substantially lower parasite

prevalence than adult birds. Malaria parasite prevalence also varied depending on the environment,

with urban nestlings significantly more infected than non-urban nestlings. There was also a tendency

for a higher *Plasmodium* prevalence in adults from more urbanized nests. Altogether, this suggests the

possible parasitic amplification effect in the city. However, the overall parasite diversity was unvariant

across urbanization contexts. Only some haemosporidian lineages occurred solely or more often in

urban areas, suggesting the possibility for habitat specificity of parasite strains.

<u>Life stage and habitat-dependent prevalence</u>

Overall, infection by *Plasmodium* was greater than infection by *Leucocytozoon*, which is a common

pattern observed across bird species (Pigeault et al. 2018, but see Merino et al. 2008). Haemosporidian

prevalence was overall low in nestlings (from 0% to 38%) but high in adults (from 95% to 100%), and

this pattern was consistent in both urban and non-urban areas. Such prevalence levels are comparable

to previous studies for adult great tits (Glaizot et al. 2012; Rooyen et al. 2013). To our knowledge, this

is the first time it is tested in 15 day-old urban great tit nestlings. In fact, only lower prevalence in

young juvenile (one year-old) birds compared to adults was previously described in great tits and other

passerine species (Wood et al. 2007; Santiago-Alarcon et al. 2016). The higher infection detection in

adults than in nestlings frames coherently with the vector (e.g., *Culex pipiens*) life cycle, with a

progressive increase in adult mosquitos and associated infection risk from spring to summer (Zélé et

al. 2014). As a consequence, the risk for 15 day-old nestlings of being infected is expected to be low

as they were sampled during spring. Similarly, Valkiunas and Iezhova (2018) found that young adults

presented lower prevalence, which is in line with the fact that Haemosporidian infections yield an

acute infection followed by a life-long chronic infection. Hence, the longer the exposure to the

parasites, the higher the probability of eventually being infected. Possibly, the lower infection in 15

day-old nestlings could also be due to the delay of detection that is not immediate after infection

(Cosgrove et al. 2006).

Our results tend to support the hypothesis of the existence of a parasitic burden in more

urbanized areas. Specifically, the observed statistical effect of nest-level urbanization on *Plasmodium*

was marginal. It would thus deserve further attention to clearly determine the role of the urbanization

at such scale. This potential effect would contrast frequent reports of lower parasitic prevalence in

urban areas including in our focal species (Bailly et al. 2016). Whether avian malaria is more or less

prevalent in cities thus appears strongly case-specific (Evans et al. 2009).. Differences in parasite

prevalence between habitats may be directly induced by variations in the presence and/or density of

vectors (e.g., Martínez-de la Puente et al. 2013). These variations should be the consequence of

presence or absence of their suitable ecological niches. For instance, among the 11 paired populations

of blackbirds *Turdus merula* studied by Evans et al. (2009), in 3 cases, avian malaria prevalence was

found to be higher in urban areas as a consequence of underwater area presence. While fine scale

densities of vectors are not yet known for the city of Montpellier and its surrounding area, a tendency

towards higher malaria prevalence in more urbanized areas could indicate higher population size or

densities of vectors in such areas, perhaps given the marshes nearby. This, however, remains to be

empirically demonstrated.

In addition, urban nestlings showed higher prevalence than non-urban ones, although the non-

urban site had only one replicate, limiting robust generalization. Replication of the work done here at

other sites is therefore crucial to (in)validate this finding. Provided that this is generalizable, one can

hypothesize that early malaria infection in urban nestlings might be an indirect result of the heat island

effect. Indeed, Paz and Albersheim (2008) showed that higher temperatures in urban areas proved

beneficial to *Culex pipiens* mosquitoes growth and that some diseases (i.e., the human West Nile

Fever) transmitted by this vector appeared earlier in the season in the city compared with surrounding

countryside areas. Hence, environmental shifts observed in urban areas can be directly linked to spatial

and temporal parasite infections. In addition, malaria infections are known to vary in time (Zélé et al.

2014). Given the role of the urban area in buffering on climatic variations, urbanization could be

responsible for major changes in seasonality of parasitic infection. As shown here, this could cascade

onto the emergence of earlier disease outbreak and earlier nestling contamination. The link between

urban specific climatic features and seasonality of vectors and disease outbreaks in urban areas

remains overlooked and should be the focus of further research avenues.

<u>Spatial heterogeneity in lineage diversity</u>

When exploring diversity of Haemosporidian lineages across sites, we mostly found similar levels of

diversity along the urbanization gradient and no strong 'cluster' of similar lineages in similarly

urbanized or closer sites. Anecdotally, the non-urban sites had the lowest Haemosporidian lineage

diversity, whereas the large zoo urban park had the highest. Interestingly, previous studies reported

that urban parks with higher diversity of plant and bird species were also the most diverse in terms of

Haemosporidian lineages (in multiple species: Carbó-Ramírez et al. 2017 ; in the House Sparrow:

Jiménez-Peñuela et al. 2021). In our case, the Zoo du Lunaret consists of an 80-ha natural area where a

large diversity of both native and exotic plant and bird species coexist. Interestingly, the only

occurrence of *Plasmodium* sp. AFR065 lineage was in this zoo. According to the *MalAvi* database

(Bensch et al. 2009), this lineage was found previously only on the African continent, in two   bird

genus in Malawi (*Cercotrichas* and *Andropadus,* Lutz et al. 2015). Hence, the presence of such

lineages in this particular area of the city is most probably linked to the presence of captive African

birds in the zoo (see next section for details on these birds).

The diversity of Haemosporidian lineages at the non-urban site of La Rouvière, 20 km away

from the city of Montpellier, ranked among the lowest in richness and evenness (Table 1 and Figures 3

and 4), which contrasts with previous results found and showed opposite trends when comparing

urban and non-urban sites (e.g., in the House Sparrow : Jiménez-Peñuela et al. 2021). The difference

in diversity highlighted by these indices may however be biologically small, as the dissimilarity

between ROU and the other sites was in the range of any other pairs of sites. In our study site, the

overall urban habitat presents numerous ornamental plant species, whereas the non-urban habitat,

which is a Mediterranean forest, is mainly dominated by oak trees. Hence, even with lower density of

vegetation, the urban areas might be prone to a maintain high diversity of pathogens (Carbó-Ramírez

et al. 2017). However, such hypothesis remains to be further tested. In particular, since vector

distribution, abundance, and diversity are likely to play a major role in malaria infection patterns

observed in bird hosts, disentangling the processes underpinning parasite prevalence and diversity

patterns in different urban conditions will require combining parallel investigation of vectors and hosts

along gradients of urbanization. In this study, we however only focused on the host.

<u>Habitat specific lineages</u>

While none of the sampled sites revealed a striking divergent composition in Haemosporidian

lineages, hence similar diversity trends, we still observed some heterogeneity in lineage type

occurrence. Overall, the *Plasmodium* sp. infections were mainly dominated by SGS1 lineage

(*Plasmodium relictum*). SGS1 is known to be a generalist lineage, present in multiple avian species

and environments (Rooyen et al. 2013) and transmitted by *Culex pipiens* (Ventim et al. 2012; Inci et

al. 2012), which is widely present in the south of France in both habitats. Aside from SGS1, some

lineages were found in low occurrence exclusively in the urban habitat: AFR065 occurred once in

ZOO and DELURB4 occurred in urban sites only. Habitat specificity analyses controlling for unequal

sampling across the sites revealed that only one *Plasmodium* sp. lineage (YWT4) occurred more in

urban habitats. No lineage was found to be associated with the forest habitat. Yet, this is possible that

it is due to a lack of power in our forest dataset as it included only NN individuals. When investigating

the previous occurrences of the 3 specific urban associated lineages (i.e., AFR065, DELURB4 and

YWT4) in the MalAvi database, we found that they were relatively rarely encountered, at least in great

tits.

AFR065 was reported only twice, once in the Miombo scrub robin (*Cercotrichas barbata*,

Muscicapidae) and once in the western greenbul (*Andropadus tephrolaemus,* Pycnonotidae) in Malawi

and never on the European continent nor in the great tits (Lutz et al. 2015). As mentioned before, the

individual infected by AFR065 was captured in the Zoo du Lunaret (most natural urban site). At the

time of the sampling for this study, the zoo hosted 65 African birds from 14 different species. While

malaria infection status of these captive birds held in the zoo are low (<5%, unpublished data), we can

hypothesize that they were the initial carriers of AFR065 that was then transferred to a great tit via the

contaminated vectors. This result raises concern regarding local wildlife epidemiology when

introducing or keeping exotic wildlife captive in contact with native species.

We found no previous occurrence of the DELURB4 lineage in great tits in the *MalAvi*

database, even if this lineage was previously shown to be the second most common lineage present in

the vector *C. pipiens* in the area (Zélé et al. 2014), and numerously recorded in the close sister species

the Blue tit *Cyanistes caeruleus* (Ferrer et al. 2012) and in other bird families (e.g., *Passeridae*,

Turdidae and *Muscicapidae*) in several European countries (Spain, Italy, Bulgaria, Russia according to

the *MalAvi* database). Similarly, YWT4 is a rare lineage with only 7 occurrences in the whole *MalAvi*

database, mainly in the Western yellow wagtail (*Motacilla flava*), but was found 25 times in the

studied urban great tits, and once in a non-urban bird. Reasons why these lineages were more common

in urban areas than in non-urban habitats remain to be explored. A possible explanation could be the

difference in bird community composition between habitats, leading to contact with different bird

species, each with their own body of specific Haemosporidian parasite lineages as suggested by the

occurrence of a rare lineage in the zoo that is hosting African species. Testing this hypothesis would

require a thorough scan of Haemosporidian infections in multiple host species from both urban and

non-urban habitats in replicated cities combined with a thorough characterization of bird community

assemblages and abundances along urbanization gradients.

Conclusion

While we found no striking difference in malaria prevalence between urban and non-urban great tits,

urbanization was associated with earlier infections in nestlings. In addition, we found a weak tendency

for *Plasmodium* sp. prevalence to increase with urbanization. While our results will need to be

replicated with higher number of sampled sites and individuals, they could suggest that urbanization

does not decrease parasitic load but may, on the contrary, lead to a parasitic burden for urban great tits. Interestingly, although sites displayed no major differences in Haemosporidian lineage community composition, urban sites hosted preferentially lineages that rarely occurred in malaria databases. This suggests that urbanization could play a role in the emergence and spread of previously rare disease strains, especially when zoos are present.

STATEMENTS AND DECLARATIONS

Funding

This work was funded by the Agence Nationale de la Recherche (grant "EVOMALWILD", ANR-17-CE35-0012) and long-term support from the OSU-OREME (Observatoire des Sciences de l'Univers – Observatoire de REcherche Montpellierain de l'Environnement).

Conflict of interest

The authors declare no conflict of interest.

Ethical statement

Captures were performed under personal ringing permits delivered by the CRBPO (Centre de Recherches par le Baguage des Populations d'Oiseaux, e.g., ringing permit for Anne Charmantier number 1907) for the Research Ringing Programme number 369. All experimental protocols were approved by the ethics committee for animal experimentation of Languedoc Roussillon (CEEA-LR, most recent approval in 2018 for APAFIS#8608-2017012011062214) as well as by Regional Institutions (most recent bylaw issued on 07/04/2022 by the Prefecture n° 2B-2022-04-07-00002).

Data and code sharing

Data and code used for this study are freely available on Zenodo via Github (DOI : 10.5281/zenodo.8329693 & https://github.com/AudeCaizergues/Malaria_Great_Tits).

Authors contribution

A.E.C., S.P & A.C. collected the samples along with field collaborators. M.J. & A.B. performed the molecular analyses. A.E.C. & B.R. conducted the statistical analyses and wrote the manuscript. C.P., S.G. & A.C. conceptualised the research. S.G. & A.C. financed the project. All authors contributed to writing the manuscript.

Acknowledgements

We are grateful to the managers and the employees of the Zoo de Lunaret, Montpellier, especially

Baptiste Genet, David Gomis, Marc Romans as well as the PLT platform of the CEFE for their help in

data collection and their feedback on our research. We also thank the city Council of Montpellier for

permitting us to carry out this long-term research project.

BIBLIOGRAPHY

Abella-Medrano CA, Ibáñez-Bernal S, Carbó-Ramírez P, Santiago-Alarcon D (2018) Blood-meal

preferences and avian malaria detection in mosquitoes (Diptera: Culicidae) captured at different

land use types within a neotropical montane cloud forest matrix. Parasitol Int 67:313–320.

https://doi.org/10.1016/J.PARINT.2018.01.006

Asghar M, Hasselquist D, Bensch S, et al (2011) Are chronic avian haemosporidian infections costly

in wild birds? J Avian Biol 42:530–537. https://doi.org/10.1111/J.1600-048X.2011.05281.X

Bailly J, Scheifler R, Belvalette M, et al (2016) Negative impact of urban habitat on immunity in the

great tit Parus major. Oecologia 182:1053–1062. https://doi.org/10.1007/S00442-016-3730-

2/TABLES/2

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using **lme4**. J

Stat Softw 67:. https://doi.org/10.18637/jss.v067.i01

Becker DJ, Streicker DG, Altizer S (2015) Linking anthropogenic resources to wildlife–pathogen

dynamics: a review and meta-analysis. Ecol Lett 18:483–495. https://doi.org/10.1111/ELE.12428

Belo NO, Pinheiro RT, Reis ES, et al (2011) Prevalence and Lineage Diversity of Avian

Haemosporidians from Three Distinct Cerrado Habitats in Brazil. PLoS One 6:e17654.

https://doi.org/10.1371/JOURNAL.PONE.0017654

Bensch S, Hellgren O, PÉrez-Tris J (2009) MalAvi: a public database of malaria parasites and related

haemosporidians in avian hosts based on mitochondrial cytochrome b lineages. Mol Ecol Resour

9:1353–1358. https://doi.org/10.1111/J.1755-0998.2009.02692.X

Bichet C, Scheifler R, Cœurdassier M, et al (2013a) Urbanization, Trace Metal Pollution, and Malaria

Prevalence in the House Sparrow. PLoS One 8:e53866.

https://doi.org/10.1371/JOURNAL.PONE.0053866

Bichet C, Scheifler R, Cœurdassier M, et al (2013b) Urbanization, Trace Metal Pollution, and Malaria

Prevalence in the House Sparrow. PLoS One 8:e53866.

https://doi.org/10.1371/JOURNAL.PONE.0053866

Bradley CA, Altizer S (2007) Urbanization and the ecology of wildlife diseases. Trends Ecol Evol

22:95–102. https://doi.org/10.1016/J.TREE.2006.11.001

Caizergues AE, Charmantier A, Lambrechts MM, et al (2021) An avian urban morphotype: how the

city environment shapes great tit morphology at different life stages. Urban Ecosyst 1–13.

https://doi.org/10.1007/s11252-020-01077-0

Calegaro-Marques C, Amato SB (2014) Urbanization Breaks Up Host-Parasite Interactions: A Case

Study on Parasite Community Ecology of Rufous-Bellied Thrushes (Turdus rufiventris) along a

Rural-Urban Gradient. PLoS One 9:e103144. https://doi.org/10.1371/JOURNAL.PONE.0103144

Capilla-Lasheras P, Dominoni DM, Babayan SA, et al (2017) Elevated Immune Gene Expression Is

Associated with Poor Reproductive Success of Urban Blue Tits. Front Ecol Evol 5:64.

https://doi.org/10.3389/FEVO.2017.00064

Carbó-Ramírez P, Zuria I, Schaefer HM, Santiago-Alarcon D (2017) Avian haemosporidians at three

environmentally contrasting urban greenspaces. J Urban Ecol 3:juw011.

https://doi.org/10.1093/JUE/JUW011

Charmantier A, Demeyrier V, Lambrechts M, et al (2017) Urbanization Is Associated with Divergence

in Pace-of-Life in Great Tits. Front Ecol Evol 5:53. https://doi.org/10.3389/fevo.2017.00053

Christe P, Glaizot O, Strepparava N, et al (2012) Twofold cost of reproduction: an increase in parental

effort leads to higher malarial parasitaemia and to a decrease in resistance to oxidative stress.

Proc R Soc B Biol Sci 279:1142. https://doi.org/10.1098/RSPB.2011.1546

Coene J (1993) Malaria in urban and rural Kinshasa: the entomological input. Med Vet Entomol

7:127–137. https://doi.org/10.1111/j.1365-2915.1993.tb00665.x

Cohen J (2013) Statistical power analysis for the behavioral sciences. Academic press.
Cosgrove CL, Knowles SC, Day KP, Sheldon BC (2006). No evidence for avian malaria infection

during the nestling phase in a passerine bird. Journof Parasito, 92:1302-1304.

Delgado-V. CA, French K (2012) Parasite–bird interactions in urban areas: Current evidence and

emerging questions. Landsc Urban Plan 105:5–14.

https://doi.org/10.1016/J.LANDURBPLAN.2011.12.019

Demeyrier V, Lambrechts MM, Perret P, Grégoire A (2016) Experimental demonstration of an

ecological trap for a wild bird in a human-transformed environment. Anim Behav 118:181–190.

https://doi.org/10.1016/J.ANBEHAV.2016.06.007

Dyrcz A, Wink M, Kruszewicz A, Leisler B (2005) Male Reproductive Success is Correlated With

Blood Parasite Levels and Body Condition in the Promiscuous Aquatic Warbler (Acrocephalus

Paludicola). Auk 122:558–565. https://doi.org/10.1093/AUK/122.2.558

Evans KL, Gaston KJ, Sharp SP, et al (2009) Effects of urbanisation on disease prevalence and age structure in blackbird Turdus merula populations. Oikos 118:774–782. https://doi.org/10.1111/J.1600-0706.2008.17226.X

Faeth SH, Bang C, Saari S (2011) Urban biodiversity: Patterns and mechanisms. Ann N Y Acad Sci 1223:69–81. https://doi.org/10.1111/j.1749-6632.2010.05925.x

Ferraguti M, Hernández-Lara C, Sehgal RNM, Santiago-Alarcon D (2020) Anthropogenic effects on avian haemosporidians and their vectors. Avian Malar Relat Parasites Trop Ecol Evol Syst 451–485. https://doi.org/10.1007/978-3-030-51633-8_14/FIGURES/4

Ferrer ES, García-Navas V, Sanz JJ, Ortego J (2012) Molecular characterization of avian malaria parasites in three Mediterranean blue tit (Cyanistes caeruleus) populations. Parasitol Res 111:2137–2142. https://doi.org/10.1007/S00436-012-3062-Z/TABLES/1

Fink, D., T. Auer, A. Johnston, M. Strimas-Mackey, S. Ligocki, O. Robinson, W. Hochachka, L. Jaromczyk, A. Rodewald, C. Wood, I. Davies AS (2022) Statuts et tendances eBird, Version des donnees : 2021 ; Diffuse : 2022. Ithaca, New York.

Forman RTT, Godron M (1986) Landscape Ecology. John Wiley and Sons, New York

French SS, Fokidis HB, Moore MC (2008) Variation in stress and innate immunity in the tree lizard (Urosaurus ornatus) across an urban-rural gradient. J Comp Physiol B 178:997–1005. https://doi.org/10.1007/S00360-008-0290-8

Gaston KJ, Visser ME, Hölker F (2015) The biological impacts of artificial light at night: the research challenge. Philos Trans R Soc B Biol Sci 370:20140133. https://doi.org/10.1098/rstb.2014.0133

Geue D, Partecke J (2008) Reduced parasite infestation in urban Eurasian blackbirds (Turdus merula): A factor favoring urbanization? Can J Zool 86:1419–1425. https://doi.org/10.1139/Z08-129/ASSET/IMAGES/LARGE/Z08-129F3.JPEG

Giraudeau M, Mousel M, Earl S, McGraw K (2014) Parasites in the City: Degree of Urbanization Predicts Poxvirus and Coccidian Infections in House Finches (Haemorhous mexicanus). PLoS One 9:e86747. https://doi.org/10.1371/JOURNAL.PONE.0086747

Glaizot O, Fumagalli L, Iritano K, et al (2012) High Prevalence and Lineage Diversity of Avian Malaria in Wild Populations of Great Tits (Parus major) and Mosquitoes (Culex pipiens). PLoS One 7:e34964. https://doi.org/10.1371/JOURNAL.PONE.0034964

Hellgren O, Waldenström J, Bensch S (2004) A new PCR assay for simultaneous studies of Leucocytozoon, Plasmodium, and Haemoproteus from avian blood. J Parasitol 90:797–802. https://doi.org/10.1645/GE-184R1

Hõrak P, Ots I, Vellau H, et al (2001) Carotenoid-based plumage coloration reflects hemoparasite
infection and local survival in breeding great tits. Oecologia 126:166–173.
https://doi.org/10.1007/S004420000513/METRICS

Inci A, Yildirim A, Njabo KY, et al (2012) Detection and molecular characterization of avian
Plasmodium from mosquitoes in central Turkey. Vet Parasitol 188:179–184.
https://doi.org/10.1016/J.VETPAR.2012.02.012

Jansen CC, Webb CE, Graham GC, et al (2009) Blood sources of mosquitoes collected from urban
and peri-urban environments in eastern Australia with species-specific molecular analysis of
avian blood meals. Am J Trop Med Hyg 81:849–857. https://doi.org/10.4269/AJTMH.2009.09-
0008

Jiménez-Peñuela J, Ferraguti M, Martínez-de la Puente J, et al (2021) Urbanization effects on temporal
variations of avian haemosporidian infections. Environ Res 199:111234.
https://doi.org/10.1016/J.ENVRES.2021.111234

Lachish S, Knowles SCL, Alves R, et al (2011) Fitness effects of endemic malaria infections in a wild
bird population: the importance of ecological structure. J Anim Ecol 80:1196–1206.
https://doi.org/10.1111/J.1365-2656.2011.01836.X

Lepczyk CA, Aronson MFJ, Evans KL, et al (2017) Biodiversity in the City: Fundamental Questions
for Understanding the Ecology of Urban Green Spaces for Biodiversity Conservation. Bioscience
67:799–807. https://doi.org/10.1093/BIOSCI/BIX079

Lutz HL, Hochachka WM, Engel JI, et al (2015) Parasite Prevalence Corresponds to Host Life History
in a Diverse Assemblage of Afrotropical Birds and Haemosporidian Parasites. PLoS One
10:e0121254. https://doi.org/10.1371/JOURNAL.PONE.0121254

Martin LB, Boruta M (2013) The impacts of urbanization on avian disease transmission and
emergence. Avian Urban Ecol.
https://doi.org/10.1093/ACPROF:OSOBL/9780199661572.003.0009

Martínez-De La Puente J, Martínez J, Rivero-De-Aguilar J, et al (2013) Vector abundance determines
Trypanosoma prevalence in nestling blue tits. Parasitology 140:1009–1015.
https://doi.org/10.1017/S0031182013000371

Marzal A, De Lope F, Navarro C, Møller AP (2005) Malarial parasites decrease reproductive success:
An experimental study in a passerine bird. Oecologia 142:541–545.
https://doi.org/10.1007/S00442-004-1757-2/TABLES/1

Marzluff JM (2001) Worldwide urbanization and its effects on birds. In: Avian Ecology and
Conservation in an Urbanizing World. Springer US, Boston, MA, pp 19–47

657 McKinney ML (2008) Effects of urbanization on species richness: A review of plants and animals.
658   Urban Ecosyst 11:161–176. https://doi.org/10.1007/S11252-007-0045-4/TABLES/9

659 Merino S, Moreno J, Vasquez RA, Martinez J, Sanchez-Monsalvez I, Estades CF, Ippi S, Sabat P,
660   Rozzi R and McGehee (2008) Haematozoa in forest birds from southern Chile: Latitudinal
661   gradients in prevalence and parasite lineage richness. Austral Ecology 33:329-340.
662   https://doi.org/10.1111/j.1442-9993.2008.01820.x

663 Nagendra H (2002) Opposite trends in response for the Shannon and Simpson indices of landscape
664   diversity. Appl Geogr 22:175–186. https://doi.org/10.1016/S0143-6228(02)00002-4

665 Neiderud C-J (2015) How urbanization affects the epidemiology of emerging infectious diseases.
666   Infect Ecol Epidemiol 5:27060. https://doi.org/10.3402/IEE.V5.27060

667 Nielsen AB, van den Bosch M, Maruthaveeran S, van den Bosch CK (2014) Species richness in urban
668   parks and its drivers: A review of empirical evidence. Urban Ecosyst 17:305–327.
669   https://doi.org/10.1007/S11252-013-0316-1/TABLES/2

670 Ots I, Hõrak P (1998) Health impact of blood parasites in breeding great tits. Oecologia 1998 1164
671   116:441–448. https://doi.org/10.1007/S004420050608

672 Partecke J, Hegyi G, Fitze PS, et al (2020) Maternal effects and urbanization: Variation of yolk
673   androgens and immunoglobulin in city and forest blackbirds. Ecol Evol 10:2213–2224.
674   https://doi.org/10.1002/ECE3.6058

675 Paz S, Albersheim I (2008) Influence of warming tendency on Culex pipiens population abundance
676   and on the probability of West Nile fever outbreaks (Israeli case study: 2001-2005). Ecohealth
677   5:40–48. https://doi.org/10.1007/S10393-007-0150-0/TABLES/1

678 Perrier C, Lozano del Campo A, Szulkin M, Demeyrier V, Gregoire A, & Charmantier A. (2018).
679   Great tits and the city: Distribution of genomic diversity and gene–environment associations
680   along an urbanization gradient. *Evolutionary Applications*, *11*(5), 593-613.
681   https://doi.org/10.1111/eva.12580

682 Perrins CM (1979) British tits. London, UK

683 Pigeault R, Cozzarolo CS, Choquet R, et al (2018) Haemosporidian infection and co-infection affect
684   host survival and reproduction in wild populations of great tits. Int J Parasitol 48:1079–1087.
685   https://doi.org/10.1016/J.IJPARA.2018.06.007

686 Reyes R, Ahn R, Thurber K, Burke TF (2013) Urbanization and Infectious Diseases: General
687   Principles, Historical Perspectives, and Contemporary Challenges. Challenges Infect Dis 123.
688   https://doi.org/10.1007/978-1-4614-4496-1_4

Rivero A, Gandon S (2018) Evolutionary Ecology of Avian Malaria: Past to Present. Trends Parasitol
34:712–726. https://doi.org/10.1016/J.PT.2018.06.002

Rooyen J van, Lalubin F, Glaizot O, Christe P (2013) Altitudinal variation in haemosporidian parasite
distribution in great tit populations. Parasites and Vectors 6:1–10. https://doi.org/10.1186/1756-
3305-6-139

Santiago-Alarcon D, Carbó-Ramírez P, Macgregor-Fors I, et al (2018) The prevalence of avian
haemosporidian parasites in an invasive bird is lower in urban than in non-urban environments.
Ibis (Lond 1859) 162:201–214. https://doi.org/10.1111/ibi.12699

Santiago-Alarcon D, Havelka P, Schaefer HM, Segelbacher G (2012) Bloodmeal Analysis Reveals
Avian Plasmodium Infections and Broad Host Preferences of Culicoides (Diptera:
Ceratopogonidae) Vectors. PLoS One 7:e31098.
https://doi.org/10.1371/JOURNAL.PONE.0031098

Santiago-Alarcon D, MacGregor-Fors I, Kühnert K, et al (2016) Avian haemosporidian parasites in an
urban forest and their relationship to bird size and abundance. Urban Ecosyst 19:331–346.
https://doi.org/10.1007/S11252-015-0494-0/TABLES/2

Shochat E, Warren PS, Faeth SH, et al (2006) From patterns to emerging processes in mechanistic
urban ecology. Trends Ecol Evol 21:186–191. https://doi.org/10.1016/J.TREE.2005.11.019

Valkiunas G, Iezhova TA (2018) Keys to the avian malaria parasites. Malar. J. 17:1–24

Ventim R, Ramos JA, Osório H, et al (2012) Avian malaria infections in western European
mosquitoes. Parasitol Res 111:637–645. https://doi.org/10.1007/S00436-012-2880-3

Wood MJ, Cosgrove CL, Wilkin TA, et al (2007) Within-population variation in prevalence and
lineage distribution of avian malaria in blue tits, Cyanistes caeruleus. Mol Ecol 16:3263–3273.
https://doi.org/10.1111/J.1365-294X.2007.03362.X

Zélé F, Vézilier J, L'Ambert G, et al (2014) Dynamics of prevalence and diversity of avian malaria
infections in wild Culex pipiens mosquitoes: The effects of Wolbachia, filarial nematodes and
insecticide resistance. Parasites and Vectors 7:. https://doi.org/10.1186/1756-3305-7-437