Conditional independence as an assessment of evidence integration processes

Emilio Salinas[1][*] **and Terrence R Stanford**[1]

[1] Department of Neurobiology & Anatomy, Wake Forest University School of Medicine, Winston-Salem, North Carolina, United States of America

[*] Corresponding author

e-mail: esalinas@wakehealth.edu (ES)

draft May 3, 2023, 19:51

Abstract

Intuitively, combining multiple sources of evidence should lead to more accurate decisions than considering single sources of evidence individually. In practice, however, the proper computation may be difficult, or may require additional data that are inaccessible. Here, based on the concept of conditional independence, we consider model-free expressions that can serve either as simplified recipes for integrating evidence or as benchmarks for characterizing evidence integration processes. Consider three events, A, B, and C. If A and B are conditionally independent with respect to C, then the probability that C occurs given that both A and B are known, $P(C|A, B)$, can be easily calculated without the need to measure the full three-way dependency between A, B, and C. This simplified approach can be used in two general ways: to generate predictions by combining multiple (independent) sources of evidence, or to test whether separate sources of evidence are functionally independent of each other. These applications are demonstrated with four computer-simulated examples, which include detecting a disease based on repeated diagnostic testing, inferring biological age based on multiple biomarkers of aging, discriminating two spatial locations based on multiple cue stimuli (multisensory integration), and examining how behavioral performance in a visual search task depends on selection histories. The results indicate that this methodology could be useful for analyzing data in a wide range of experimental systems and conditions.

Introduction

Decisions are often based on multiple sources of evidence. For instance, to determine whether a patient has an infection, a doctor may take their temperature and order a lab test, and combine both results to form a diagnosis. The question is, how exactly should those two pieces of evidence be combined? Such inference process requires computational recipes and knowledge of past outcomes under equivalent circumstances, and the equivalence is key; for accurate estimation, the prior data must conform to the specific evidence samples being evaluated. In the case of the doctor's diagnosis, the patient's results should be compared (via appropriate probabilistic calculus) to prior measurements of temperature, lab test data, and infection status *taken simultaneously*. That is, in general, knowing the relationship between temperature and infection status on one hand, and lab test and infection status on the other, separately, is not sufficient for making an optimal informed decision — unless a particular relationship is satisfied between the three variables. In that case, optimal decisions and inferences can indeed be made based on more limited data.

This particular relationship is conditional independence [1]. When two sources of evidence, A and B, are conditionally independent with respect to a third quantity, C, then characterizing separately the relationships between A and C and between B and C suffices for completely characterizing how C will vary when A and B are both known; there is no need to make prior, simultaneous, three-way measurements to establish the necessary reference. This is important because, in practice, making pairwise measurements generally requires many fewer resources than making simultaneous measurements of more than two variables, everything else being equal, and the difference becomes dramatically larger as the number of variables increases beyond three.

Here we derive expressions for the performance that is expected when two or more sources of evidence are conditionally independent with respect to a variable of interest, and demonstrate their applicability to the life sciences with four examples. These expressions are potentially useful in two broad types of situation. One is when there are good reasons to *assume* that the conditional independence requirement is satisfied. In that case, the results can be used to make predictions easily; that is, to establish expected outcomes based on multiple, current evidence samples together with prior knowledge of pairwise measurements only. The other scenario in which the de-

rived expressions are potentially useful is when simultaneous measurements of all the variables of interest are available. In that case, the results serve as an intuitive benchmark to *test* whether separate pieces of evidence are being combined independently or not, which can provide insight into the underlying mechanistic interactions giving rise to the data. The concrete examples are chosen for clarity and to be representative of real-life contexts in which these situations may arise, but the range of potential applications is wide.

Methods

All numerical calculations were performed using the MATLAB programming environment (version R2013b and later; The Mathworks, Natick, MA). The code for generating all the shown examples and plots is available from the Zenodo repository at https://doi.org/10.5281/zenodo.7894053. Functions for computing posterior probabilities according to Equations 7–13 are included as part of the software package.

Generating correlated outcomes

Simulating the outcome of a diagnostic test, a coin flip, or any other binary outcome is simple; it amounts to drawing a random sample from a uniform distribution and comparing it to the event's probability [2]. To generate positively correlated data for N events (i.e., N diagnostic tests or biomarker readings), we used two methods. Both methods start by drawing a set of N independent random samples, and both use a parameter ρ, which varies between 0 and 1, to determine the level of correlation between them. In method 1, an additional random sample η is drawn, and each of the N samples is reset to equal η with a probability ρ. Thus, as ρ increases, it becomes more likely that a subset of the random samples are identical to each other, and thus that the outcomes of the simulated events are the same. Method 2 also considers N independent random samples and an additional one, η, but now the N random samples move toward η by an amount that is proportional to ρ. In this case, ρ determines how close the samples are to each other; when $\rho = 1$, all of them converge on the same point. Data shown in the figures are based on method 1, but results were qualitatively the same with method 2. The Matlab code that implements these methods is included in the deposited software package.

Simulating behavior in the oddball task

The model for the oddball task is described in the main text. The Matlab code that simulates it is as follows (note that the percent sign, %, indicates a comment).

```
%
% Simulate performance in a sequence of Nt trials of the oddball task
%
Nt = 100000;
tLoc = randi(4, Nt, 1);   % sequence of target locations
tClr = randi(2, Nt, 1);   % sequence of target colors
hit = NaN(Nt,1);          % initialize sequence of outcomes

pmax = 0.95;     % maximum accuracy for informed choices
alpha1 = 0.5;    % decay constant for motor bias (1/trial)
alpha2 = 0.40;   % decay constant for color sensitivity (1/trial)

mval = 0.01 + zeros(1,4);   % initialize motor biases
cval = 0.01 + zeros(1,2);   % initialize color sensitivities

```

```
92   for j=1:Nt
93       itar = tLoc(j);
94       iclr = tClr(j);
95       % probability of making an informed choice
96       pinf = 1./(1 + exp(-( cval(iclr) - 0.9)));
97       % make a choice
98       if rand < pinf
99           % informed choice based on color discrimination
100          hit(j) = (rand < pmax);
101      else
102          % guess based on motor bias
103          mguess = mval/sum(mval);
104          iguess = find((rand < cumsum(mguess)), 1);
105          hit(j) = (iguess == itar);
106      end
107      % update values for the 4 motor biases
108      mval = alpha1*mval + 0.1;
109      mval(itar) = mval(itar) + 1;
110      % update values for two color sensitivities
111      cval = alpha2*cval;
112      if hit(j) > 0
113          % color values are reinforced after a reward
114          if iclr == 1
115              cval = cval + [1 -1]*2.0;
116          else
117              cval = cval + [-1 1]*2.0;
118          end
119      end
120  end
```

For analysis of the simulated task data, the target color and target location in a series of con­secutive trials can be encoded literally, as in {Red, Green, Red, Red}, or as same/different relative to the oddball color in the current trial, as in {Same, Different, Same, Same}. We use the latter encoding because it automatically pools the data for red and green targets, which were statisti­cally identical. Likewise for target location; we use the same/different encoding because all four locations were statistically identical.

Results and Discussion

Expected accuracy based on conditionally independent sources of evidence

The starting concept is that of conditional independence. Two events A and B, are said to be conditionally independent relative to a third event C if their joint probability given C is

$$P(A, B|C) = P(A|C)\, P(B|C) \tag{1}$$

This means that, if C is known, A and B occur independently of each other. Said differently, if condition C is known, the probability that B occurs is fixed, regardless of whether A occurs or not. Indeed, the equality

$$P(B|A, C) = P(B|C) \tag{2}$$

which directly states this idea, is equivalent to the first expression. Conditional independence is entirely separate from general independence; that is, Equation 1 does not tell us whether

$$P(A, B) = P(A)\, P(B) \tag{3}$$

Fig 1. Frequency tree for three events. C corresponds to 'Carol is home', A corresponds to 'time is after 6 pm', and B corresponds to 'the dog is barking', with 1 and 0 indicating true and false. Numbers of observations are in red. Probabilities for all event combinations can be readily calculated from the tree. For example, the joint probability that all three events occur simultaneoulsy, $P(A\!=\!1, B\!=\!1, C\!=\!1)$, is equal to $3/100$, as given by the rightmost branch. The probability that the time is not past 6 pm given that Carol is not home, $P(A\!=\!0|C\!=\!0)$, is equal to $40/60$, as given by the leftmost branch, and so on. This example is constructed so that A and B occur independently of each other given C.

is true or not, and this relationship has no bearing on the results that follow.

Here is a concrete example presented as a frequency tree (Fig 1), which facilitates probabilistic reasoning and calculation [3, 4]. Suppose that the probability that Carol is home ($C = 1$) is 0.6 when it is after 6 pm ($A\!=\!1$), but only 0.2 when it is before 6 pm ($A\!=\!0$), and futhermore, her dog only barks ($B\!=\!1$) with a probability of 0.1 when Carol is home, but barks with a probability of 0.6 when she is away ($C\!=\!0$). Note that the barking depends only on whether Carol is home or not, so knowing the time of day makes no difference in assessing what the dog will do, if we already know where Carol is. Translated into numbers, it can be verified from Fig 1 that

$$P(B\!=\!1|A\!=\!1, C\!=\!1) = P(B\!=\!1|A\!=\!0, C\!=\!1) = P(B\!=\!1|C\!=\!1) = 0.1 \tag{4}$$

and that similar equalities are also satisfied when $C = 0$ and for the case $B = 0$, in agreement with conditional independence as defined in Equation 2. However, *in general* the barking is not independent of time; for instance

$$P(B\!=\!1, A\!=\!1) = 0.15 \neq 0.2 = 0.4 \times 0.5 = P(B\!=\!1)\,P(A\!=\!1) \tag{5}$$

What is interesting about this example is that, because of their conditional independence, the dog barking and the time of day can be used as independent pieces of evidence regarding whether Carol is home or not. Now the quantity that we seek is the conditional probability of C given A and B, or

$$P(C|A, B) = P(A, B|C)\,\frac{P(C)}{P(A, B)} \tag{6}$$

where the right side is the standard identity based on Bayes theorem. The calculation has no general solution because there is no universal way to obtain the first term on the right; the three-way relationship between A, B, and C simply has to be measured — unless A and B are conditionally independent given C.

If that is the case, then Equation 1 can be used to factorize all the terms where A and B appear jointly in Equation 6, and everything can be put in terms of $P(C|A)$ and $P(C|B)$, which are the separate, individual pieces of information about the event of interest, whether Carol is home or

not. When C is a binary variable representing two possible states or outcomes (0 and 1), the result of combining Equations 1 and 6 is, after rearranging terms,

$$P(C=1|A, B) = \frac{P(C=1|A)\,P(C=1|B)}{P(C=1|A)\,P(C=1|B) + P(C=0|A)\,P(C=0|B)\,\frac{P(C=1)}{P(C=0)}} \tag{7}$$

The result is stated for $C = 1$, but it is entirely analogous for $C = 0$ (the $C = 1$ and $C = 0$ terms simply exchange places). This expression provides a recipe for determining the probability of event C given two sources of evidence, A and B, and knowledge of how each of them predicts C separately, assuming that A and B act independently of each other with respect to C. One can also think of this equation as an alternative, operationally distinct way to define conditional independence: if the relationship is satisfied for all the values of the variables, it means that A and B are conditionally independent with respect to C. The applications considered further below illustrate both uses of this result.

For the example above (Fig 1), the prior probability that Carol is home is $P(C=1) = 0.4$. This is the estimate of her status without any further knowledge or measurement. The probability that Carol is home given that it is past 6 pm is $P(C = 1|A = 1) = 0.6$, and the probability that Carol is home given that the dog is not barking is $P(C = 1|B = 0) = 0.6$. Therefore, because the time and the barking are conditionally independent, it is straightforward to apply these numbers to Equation 7 to infer the probability that Carol is home given that, both, it is past 6 pm and the dog is not barking: $P(C=1|A=1, B=0) = 27/(27 + 8) \approx 0.77$. Certainty about her status grows when the two sources of evidence are in agreement, as one would expect, and Equation 7 tells exactly by how much.

Another way to view the implications of conditional independence is to consider to the odds of the two outcomes for event C. In general, the odds depend on two three-way measurements,

$$\frac{P(C=1|A, B)}{P(C=0|A, B)} = \frac{P(A, B, C=1)}{P(A, B, C=0)} \tag{8}$$

but under conditional independence (i.e., using Equation 2) this turns into

$$\frac{P(C=1|A, B)}{P(C=0|A, B)} = \frac{P(C=1|A)}{P(C=0|A)}\,\frac{P(C=1|B)}{P(C=0|B)}\,\frac{P(C=0)}{P(C=1)} \tag{9}$$

which is simply a product of individual odds. Seen this way, it is clear that the events A and B are uninformative when they predict C with the same odds as the prior. When one of them is informative and the other is not (e.g., if $P(C=1|B) = P(C=1)$), the odds for C simply adhere to those given by the informative event.

Expected accuracy in higher dimensions

Equation 7 can be generalized in several ways. We develop two special cases first, and then present a more general expression that encompasses both.

The first extension of Equation 7 is to the case where more than just two sources of evidence may be predictive of event C; for example, in addition to the time of day and the dog barking, the presence of a car in the garage may also be informative of whether Carol is home or not. If instead of A and B there are N events $E_1, E_2, \ldots, E_N$, all of which are conditionally independent given C, then the probability that C is in state 1 given knowledge of all those events is

$$P(C=1|E_1, \ldots E_N) = \frac{\prod_{j=1}^{N} P(C=1|E_j)}{\prod_{j=1}^{N} P(C=1|E_j) + \prod_{j=1}^{N} P(C=0|E_j) \left[\frac{P(C=1)}{P(C=0)}\right]^{N-1}} \tag{10}$$

Equation 7 is the same as above for the case $N=2$, and again, it is written for one particular outcome, $C=1$, but the probability of the alternative outcome is entirely analogous.

The second straightforward extension of Equation 7 is to the case where the C variable can take more than two states; for example, we may be interested in whether Carol is home, at work, or at the coffee shop, which would require three values. If C can take any of M possible values $c_1, c_2, \ldots, c_M$ and A and B are still conditionally independent across all of them, then the probability that C is equal to c_j (i.e., that it is in state j), is

$$P(C\!=\!c_j|A,B) = \frac{P(C\!=\!c_j|A)\,P(C\!=\!c_j|B)}{P(C\!=\!c_j|A)\,P(C\!=\!c_j|B) + \sum_{i\neq j}^{M} P(C\!=\!c_i|A)\,P(C\!=\!c_i|B)\frac{P(C=c_j)}{P(C=c_i)}} \tag{11}$$

This expression is also the same as Equation 7 when $M\!=\!2$.

Finally, a more general expression that allows for both multiple sources of evidence and multiple states of the variable of interest is the following

$$f(C\!=\!c_j; E_1, \ldots E_N) \;\equiv\; \frac{\prod_{k=1}^{N} P(C\!=\!c_j|E_k)}{[P(C\!=\!c_j)]^{N-1}} \tag{12}$$

$$P(C\!=\!c_j|E_1, \ldots E_N) \;=\; \frac{f(C\!=\!c_j; E_1, \ldots E_N)}{\sum_i^M f(C\!=\!c_i; E_1, \ldots E_N)} \tag{13}$$

where the denominator in the last equation is simply a normalization factor. This is the probability that the variable of interest is in state j (one of M possible states) given that N conditionally independent sources of evidence are known. This is the main analytic result. Equations 7, 10, and 11 are all lower-dimension versions of this expression (with terms arranged slightly differently). In this form, it is clear that what matters is how each conditional probability $P(C\!=\!c_j|E_k)$ deviates from the prior, $P(C\!=\!c_j)$, in terms of their ratio.

The critical point is that these equations provide a statistical expectation for what *should* happen if multiple sources of evidence act independently given the quantity of interest, C. This provides a model-free benchmark for analyzing the efficacy of evidence-integration processes in a variety of situations, as discussed in the rest of the paper.

Repeated diagnostic tests for a disease

A paradigmatic situation in which probabilities are "inverted," as in Equation 6, is in clinical settings, when a diagnostic or screening test is applied to detect a disease. Let the variable D represent the presence ($D=1$) or absence ($D=0$) of the disease, and let the variable T represent the outcome of the test, which can be positive ($T=+$) or negative ($T=-$). In this context, two probabilities characterize the effectiveness of the test,

$$\begin{aligned} P(T\!=\!+|D\!=\!1) &= \text{ sensitivity } = s^+ \\ P(T\!=\!-|D\!=\!0) &= \text{ specificity } = s^- \end{aligned} \tag{14}$$

where s^+ and s^- are just abbreviations for these quantities, which are known as sensitivity and specificity [5, 6]. Knowing these two probabilities, together with the prevalence of the disease, which is its prior probability, $P(D\!=\!1)$, allows one to use the test as a diagnostic; that is, it makes it possible to test a new patient and determine, for example, the probability that the disease is

indeed present when the test is positive,

$$
\begin{aligned}
P(D{=}1|T{=}+) &= \frac{P(T{=}+|D{=}1)\,P(D{=}1)}{P(T{=}+)} \\
&= \frac{s^+\,P(D{=}1)}{s^+\,P(D{=}1) + (1-s^-)\,P(D{=}0)} \\
&= r^+
\end{aligned}
\tag{15}
$$

or the probability that the disease is indeed absent when the test is negative,

$$
\begin{aligned}
P(D{=}0|T{=}-) &= \frac{P(T{=}-|D{=}0)\,P(D{=}0)}{P(T{=}-)} \\
&= \frac{s^-\,P(D{=}0)}{s^-\,P(D{=}0) + (1-s^+)\,P(D{=}1)} \\
&= r^-
\end{aligned}
\tag{16}
$$

where $P(D{=}0)$ is simply $1 - P(D{=}1)$ and, again for convenience, r^+ and r^- are simply abbreviations for the probabilities defined on the left-hand sides of these equations. These probabilities are often referred to as the positive predictive value (r^+) and the negative predictive value (r^-) of the diagnostic test [7].

Now suppose that the test can be repeated multiple times, for instance by drawing multiple samples from the same patient [8]. In that case, if the outcomes of these repeated tests are conditionally independent given the disease state, D, then Equation 10 can be used to determine the probability that the disease is present given that n out of N samples turn out to be positive. Because all the test repetitions have the same sensitivity and specificity, and thus the same posterior probabilities r^+ and r^-, Equation 10 becomes

$$
P(D{=}1|n + \text{tests}) = \frac{(r^+)^n\,(1-r^-)^{N-n}}{(r^+)^n\,(1-r^-)^{N-n} + (1-r^+)^n\,(r^-)^{N-n}\left[\frac{P(D=1)}{P(D=0)}\right]^{N-1}}
\tag{17}
$$

The behavior of this expression for different values of n and N demonstrates the power of combining independent evidence samples (Fig 2). Suppose that both r^+ and r^- are relatively close to 0.5, so the test is rather unreliable and a single application provides little information (Fig 2a, top). If the test is applied 4 times (Fig 2a, second row), its accuracy increases substantially when all 4 tests are positive or all are negative, but the procedure is still not very useful because for the most common outcomes (1 or 2 positive tests; see dashed lines) the resulting probabilities still do not deviate from 0.5 very much. However, as the test is repeated more times, the results eventually become both accurate (probabilities close to 1 or 0) and unambiguous, as the range in which the number of positive tests provides an uncertain diagnosis becomes progressively less likely (Fig 2a, bottom plot; note little overlap between dashed lines).

Such convergence to high accuracy is also observed when one test result (positive or negative) is much more informative than the other. In an example that approximates the case of a rare disease, i.e., one with relatively low incidence, a single negative test indicates that the disease is very likely absent, but a single positive test provides negligible evidence that the disease is present (Fig 2b, top). However, in spite of the strong asymmetry, multiple repetitions of the test eventually yield accurate and unambigous diagnoses for both disease presence and absence (Fig 2b, bottom) — as long as the tests remain conditionally independent of each other given the disease state.

Fig 2. Multiple conditionally-independent repetitions of a diagnostic test increase diagnostic accuracy. The plots show results from Equation 17 for different repetitions of a test (N, varying across rows) as a function of the number of positive tests obtained (n, along the x-axes). Red bars and data points correspond to $P(D = 1|n + \text{tests})$; blue bars and data points correspond to $P(D = 0|n + \text{tests})$. Dashed lines are proportional to the likelihood of obtaining a particular number of positive tests when the disease is present (red) or absent (blue). **a**, Example where a single test has low sensitivity and specificity. In this case, $s^+ = 0.5$, $s^- = 0.75$, and $P(D = 1) = 0.5$, so one test is relatively uninformative. **b**, Example where a single test has low sensitivity and high specificity. In this case, $s^+ = 0.67$, $s^- = 0.91$, and $P(D = 1) = 0.0148$, so a single negative test is clearly informative but a positive one not so much. With increasing repetitions, the diagnosis becomes more accurate and less ambiguous.

251   This example illustrates how to generate a prediction for the performance of a detector based
252 on multiple repetitions of a test when the conditional independence between replicates of this test
253 is either assumed or known to be applicable. However, the same relationship (Equation 17) can
254 be used to determine whether multiple repetitions of a test are, in fact, conditionally independent
255 given the state of the tested individual, or to quantify the degree to which the data deviate from
256 conditional independence.

257   Suppose that the repeated-test protocol is applied many times and that the ground truth about
the presence or absence of the disease can be determined by a separate, gold-standard test. In
that case, the predictive values (i.e., posterior probabilities) can be determined empirically from
the observed results. For instance, for a four-test protocol one would compute the probability that
the disease was actually present given that 0, 1, 2, 3, or 4 tests were positive. These measured
probabilities would then be compared against those predicted based on the single-test data and
the assumption of conditional independence, via Equation 17 (i.e., the data in Fig 2, row 2). For
repeated tests that are indeed conditionally independent given the disease state, a plot of the
empirical probabilities against the predicted ones would produce data points lying along the main
diagonal. In contrast, for repeated tests that violate the independence condition, the data points
would deviate from the diagonal, and the magnitude of the deviations would be indicative of the
degree of correlation between the repeated tests (Fig 3).

269   To illustrate the possible results, a subset of the conditions depicted in Fig 2 were simulated
with diagnostic tests that, upon repetition, were correlated to varying degrees (Fig 3; Methods).
What is notable is that the discrepancy between the predicted and "empirical" results can vary
widely depending on the specific conditions of the simulation. Of course, a higher degree of
correlation produced differences of larger magnitude (Fig 3, compare left versus right columns
in panels a and b), as one would expect. However, the impact of the correlation was milder
for fewer repetitions than for more (Fig 3a, b, compare top versus bottom rows), and depended

Fig 3. Determining conditional independence across repetitions of a diagnostic test. The plots compare the results obtained from Equation 17 (x axes), which are based on test outcomes that are conditionally independent given the disease state, with the results of simulations (y axes) in which the test outcomes were correlated. The number of repeated tests is indicated in each case, along with their correlation coefficient (ρ). Points correspond to $P(D = 1|n + \text{tests})$ (orange) and $P(D = 0|n + \text{tests})$ (blue). **a**, Example where a single test has low sensitivity and specificity (same as in Fig 2a). **b**, Example where a single test has low sensitivity and high specificity (same as in Fig 2b). The discrepancy between the independent and non-independent conditions increases with the number of test repetitions and their degree of correlation (ρ). CI, conditional independence.

on the symmetry of the sensitivity and specificity values as well (Fig 3, compare a versus b). The examples suggest that, under certain conditions, the assumption of conditional independence may be a reasonable approximation even when it is not strictly true.

Calculating biological age

There is much interest in understanding the fundamental processes that lead to performance decline during aging — for instance, how environmental versus genetic factors contribute to it. An important concept in this context is that of biological age, which represents the functional state of a biological system along its lifelong trajectory irrespective of its chronological age [9]. Practical determination of biological age is thought to be potentially useful as a more accurate predictor of death or disease than standard chronological age [10].

Numerous physiological processes have been identified that evolve in stereotypical ways over the normal course of aging. Some of these aging biomarkers are widespread molecular processes [11, 12]. For instance, telomere length is indicative of the number of cell division cycles that the organism has gone through; DNA methylation is indicative of gene transcription activity and serves as an "epigenetic clock"; the levels of certain proteins are indicative of cellular senescence and correlate strongly with chronological age; etc. Many other physiological markers of aging are organ-specific [9]. For example, lung function as measured by peak aerobic capacity is one of the most reliable correlates of aging, but cardiovascular, immune, neurocognitive, and other systems show characteristic trajectories with aging that have both macroscopic and cellular manifestations. Knowing the actual capabilities of these systems at a particular moment, i.e., their effective wear and tear, so to speak, can serve to define the biological age of a subject. For our purposes, the key question is how. How should multiple aging biomarkers be combined into one quantity that

Fig 4. Estimation of biological age. a, Probability curves for three hypothetical biomarkers of age. Each biomarker is assumed to have two possible states, and each plotted function corresponds to $P(B=1|\text{age})$. **b**, Each curve shows the probability that a subject has a particular age given the measured states of the three biomarkers above, assuming that they are conditionally independent of each other with respect to age. The 8 curves plotted correspond to the 8 possible states of the 3 (binary) biomarkers. The left-most curve (dark red) corresponds to $P(\text{age}|000)$, which is when all biomarkers are in the 0 state; the next curve (bright red) corresponds to $P(\text{age}|100)$, which is when the first biomarker is in state 1 and the other two are in state 0, and so on. Colored curves are for congruent states, in which the biomarkers are activated in an order consistent with the midpoints of their probability curves (states 000, 100, 110, and 111); light gray curves are for the remaining, incongruent states (states 010, 001, 011, 101). **c, d**, As in **a**, **b**, but with 8 biomarkers. Now 256 curves are plotted in panel **d**, 8 congruent and 248 incongruent. **e, f**, As in **c**, **d**, but with biomarkers that generally switch state more abruptly as functions of age.

corresponds to biological age?

There are several possible approaches [13, 14, 15, 16], but Equations 12, 13 provide a path that
is potentially data efficient and leads to a simple, easily interpretable result. Furthermore, because
many of the identified biomarkers of aging relate to widely different biological processes, at least
a subset of them are likely to be conditionally independent with respect to chronological age. We
illustrate this with a hypothetical example (Fig 4).

Consider three aging biomarkers, each of which can be in two states, 0 or 1, and suppose
that the probability of being in state 1 increases with age (Fig 4a, top). One such probability
function could describe, for instance, the fraction within a given subject population for which
telomere length has dipped below a certain threshold, or the fraction for which the concentration
of a particular hormone is below a given cutoff, etc. Whether the functions increase or decrease
does not matter; they do not even need to be monotonic. In the examples, all the biomarker
functions are shown as monotonically increasing simply for ease of visualization, but what is
important is that they represent probabilities. With this in mind, the expression

$$P(B=1|a) = \frac{1}{2}\left(1 + \text{erf}\left(\frac{a-\alpha}{\sqrt{2}\sigma}\right)\right) \qquad (18)$$

312 where erf is the error function, corresponds to the probability of biomarker B being in state 1
313 at each value of a, which represents chronological age, and this probability is characterized by
314 two biomarker-specific parameters: α, which determines the midpoint of the curve, and σ, which

determines the steepness of its rise. In the example with three biomarkers, each one follows an equation like this but with different parameter values (Fig 4a, top).

Another necessary ingredient for the computation of biological age is knowledge of the overall prior age distribution, $P(a)$. For illustration purposes, we consider the age distribution to be flat between 0 and 100 years, with zero probability elsewhere. This simplifies the discussion, but is by no means a requirement and does not affect the conclusions.

Having defined the prior age distribution, $P(a)$, the prior for each biomarker state, i.e., $P(B=1)$ and $P(B=0)$, can be obtained by combining $P(a)$ with Equation 18. Then, Equation 18 can be inverted according to Bayes theorem to generate the probability function $P(a|B)$ for each biomarker. Finally, using Equations 12 and 13, we can compute the probability that a subject has a particular age given knowledge of the three biomarkers. With 3 binary biomarkers there are a total of $2^3 = 8$ possible state vectors, so 8 possible curves describing the likely values of age given the measured physiological state (Fig 4b). For instance, when the state vector is 000, none of the biomarkers have transitioned to state 1, so the subject is likely to be very young (Fig 4b, dark red, leftmost curve). When the state vector is 110, the first two biomarkers have transitioned but the third has not, so the subject is most likely to be about 65 years old (Fig 4b, gold curve, third colored curve from left). Some state vectors are more likely (Fig 4b, colored curves) than others (Fig 4b, gray curves), but they all lead to a distribution of ages that reflects the measured condition of the subject. By this procedure, biological age is defined as the age that would be inferred given the physiological condition of the subject in reference to comparable historical data.

The situation is entirely analogous when, say, 8 biomarkers instead of 3 are used (Fig 4c, e), except that now $2^8 = 256$ state vectors are potentially observable (Fig 4d, f, note the number of curves). With more biomarkers, the distributions of likely ages given the states are typically narrower than with fewer biomarkers (Fig 4, compare b versus d), but the spread of these resulting distributions also depends on the steepness of the biomarker state transitions (Fig 4, compare c versus e, and d versus f).

The point of this example is to show that, having identified a set of physiological quantities that vary reliably with age, and having formulated their dependencies in terms of state-transition probabilities, there is a natural way to define biological age — namely, as the age that one would infer based on historical data, a subject's current physiological condition, and the rules of probability. And this inference is particularly easy to compute when the physiological quantities are conditionally independent of each other with respect to age.

As with the example in the previous section, the same equations can be used to infer a quantity of interest (age in this case) or, alternatively, to test the degree to which the measured variables (the biomarkers) deviate from conditional independence. To illustrate the latter process, we simulated the measurement of the 8 binary biomarkers with probability curves shown in Fig 4c. For a given age and a given iteration, the 8 responses were generated by comparing 8 randomly drawn samples from a uniform distribution to the corresponding probability-curve values, as is customary for simulating probabilistic outcomes. However, these random samples were correlated (Methods). As a result, the biomarker readouts for each iteration — that is, for each simulated subject — were also correlated. This way, it was possible to produce non-independent biomarker readings, as shown by the resulting matrix of correlation coefficients (Fig 5a); when one biomarker was in state 1, other biomarkers from the same simulated subject were more likely to be in state 1 too. That the resulting biomarkers were not conditionally independent with respect to age can be verified by comparing the probability of any given biomarker state vector $\mathbf{B}$ at each age in two cases: as expected from the probability curves and conditional independence (Fig 5b, x axis), and as observed from the actual simulated biomarker states (Fig 5b, y axis). Although the data points do not fall too far off the diagonal, the underlying correlations generated state-vector probabilities

Fig 5. Determining conditional independence between biomarkers of aging. Results are for the same set of 8 binary biomarkers depicted in Fig 4c, but either satisfying or violating the conditional independence assumption. **a**, Matrix of correlation coefficients obtained after simulating the correlated responses of the 8 biomarkers. **b**, Data points indicate the probability of a biomarker state **B** given the age of the subject, either based on the conditional independence assumption (x axis) or on the results of simulations (y axes) in which the biomarker outcomes were correlated. Results are for the nine congruent state vectors, that is, vectors 00000000, 10000000, 110000000,... and 11111111. Same color scheme as in Fig 4d. Note deviations from the diagonal demonstrating the impact of the correlations between outcomes. **c**, Data points indicate the probability of a particular age given the biomarker vector state. Values on the x axis are as expected from conditional independence (Equations 12, 13; same values as on the y axis of Fig 4d). Values on the y axis are as determined from the simulated biomarker states with correlations. With binary biomarkers, conditional independence provides a good approximation to the true posterior probability $P(\text{age}|\mathbf{B})$ even when the assumption is not strictly true.

that clearly deviate from those obtained with zero correlation. More importantly, however, the correlations only caused a relatively modest distortion of the posterior probability for age given the state vector. That is, the age distributions inferred given the actual, correlated biomarker state vectors (Fig 5c, y axis) were fairly similar to the distributions obtained by ignoring those correlations and assuming conditional independence (Fig 5c, x axis). The results were qualitatively the same for different numbers of biomarkers (from 3 to 12) and regardless of the specific parameter values used (α and σ in Equation 18).

This example illustrates both applications of the expressions derived at the beginning in the context of an interesting biomedical problem, but still within a highly simplified scenario. Addressing the problem in reality would require much more detail, but this is a proof of concept. The point is that conditional independence is a potentially useful tool either for selecting informative biomarkers that are non-redundant, or as an alternative method or benchmark for estimating biological age.

Integration of sensory cues from multiple modalities

Everyday behaviors are often guided by diverse types of sensory input. For instance, hearing the muffled sound of broken leaves while gleaning a grayish object suddenly moving in the periphery may lead one to conclude that a squirrel just hid behind a tree. There is much interest in understanding the mechanisms whereby neuronal circuits combine multiple sources of information, as in this example, to generate a signal that is more reliable and informative than its individual components. When the individual cues consist of sensory stimuli of different qualities or modalities, this problem often falls under the rubric of multisensory integration [17, 18, 19].

In practice, cue integration studies often rely on behavioral tasks in which a choice must be made based on two cues — we will call them A and V in this case — and success in the task is measured for various combinations of cue values. A common design is to interleave trials in

Fig 6. Conditional independence as a benchmark for performance in a hypothetical multisensory integration task. **a**–**c**, Sequence of events in the multisensory spatial discrimination task. After the subject fixates their gaze on a central point (Fixation), a cue stimulus is briefly presented (Cue) either to the left or the right of fixation. The subject's task is to respond by making an eye movement (Saccade) to the choice target (open circle) that is on the same side as the stimulus. Three trial types are interleaved: unisensory auditory trials (**a**), unisensory visual trials (**b**), and multisensory trials (**c**). Black arrows indicate eye movements. **d**, Simulated fraction correct as a function of visual cue luminance in five experimental conditions. Blue horizontal lines mark performance in unisensory auditory trials with two sound intensities (A_1 and A_2). Black symbols indicate performance in unisensory visual trials (V). Light and dark red symbols indicate the multisensory performance expected from conditional independence when combining the visual cues with the first ($V + A_1$) or the second ($V + A_2$) auditory stimulus. Note that chance performance is at 50% correct (dotted line).

which A is presented alone, V is presented alone, and A and V are presented together (Fig 6a-c). A fundamental question in this context is whether the performance of the subject (or the response of a recorded neuron) in the multi-cue condition is higher than would be expected based on the performance levels seen in the single-cue conditions. Choice accuracy above such expectation may be indicative of active mechanisms dedicated to generating combined signals that are most effective or advantageous; and vice versa, performance below expectation may be diagnostic of internal correlations or other constraints that may limit the processing power of the underlying circuitry [17, 18, 19].

Interestingly, though, it is often unclear what the default expectation should be. Prior approaches have been at least in part tailored according to the measurement of interest, which may be choice accuracy (percent correct), reaction time, or the firing activity of a neuron, for instance. In behavioral studies, a popular benchmark is a Bayesian model that makes broad statistical assumptions about the underlying signal detection processes and relevant noise distributions; i.e., about the variability in the internal representations of the cues [20, 21, 22]. Alternatively, there have been various proposals for estimating the increase in performance that should be expected based simply on the fact that in the multicue condition the system has two sensory signals to 'choose' from, rather than one [23, 24, 25, 26, 27]. The idea in this case is that each sensory channel can independently drive the decision process, but at any point in time only the stronger signal contributes to the subject's response. We submit that the performance expected when the two cues are conditionally independent relative to the choice provides a principled, model-free benchmark that in many cases is ideal for evaluating the effectiveness of multisensory integration.

To illustrate this, we consider a simple hypothetical task that captures the essence of the problem. In the multisensory spatial discrimination task, a cue is given in each trial and the subject must indicate which side it originated from, left or right. Auditory (Fig 6a), visual (Fig 6b), and multisensory trials (Fig 6c) are randomly interleaved, and all stimuli are brief (say, 50 ms). To

demonstrate the full range of possible results, we assume that the luminance of the visual stimulus varies between 0 and a maximum value, such that performance in the unisensory visual trials covers the full range between chance (50% correct) and 100% correct (Fig 6b, black curve). The auditory stimulus is instead delivered at two intensities, both of them yielding performance levels in between the two extremes (Fig 6b, blue horizontal lines). The key question is: given the performance in the two unisensory conditions, what performance should be expected in the multisensory case?

Conditional independence provides a model-free answer that is consistent with the absence of a functional interaction between the two stimuli. When the visual and auditory signals are conditionally independent with respect to the subject's choice, the expected performance curves (Fig 6d, light and dark red curves) demonstrate improved accuracy that stays within the limits imposed by the representation of probabilities. At the lowest luminance, multisensory choices are just as accurate as with the auditory stimulus alone; this makes sense because a visual stimulus that is entirely uninformative (thus yielding chance accuracy) cannot possibly improve performance when combined with an informative auditory cue. At the other end, when the visual stimulus is of the highest luminance, choice accuracy in visual-only trials is already near 100% correct, so an additional auditory cue cannot increase performance much further. Overall, in the multisensory conditions, the fraction of correct choices shows a graded increase with luminance that parallels that in the visual-only condition, but bracketed between the two limits just discussed.

The predicted multisensory curves in this example were generated from a formula that is essentially identical to Equation 7, but there is a subtle difference that is important to explain. Consider an auditory stimulus of fixed intensity and a visual stimulus of fixed luminance, and take $V=1$ to mean that the visual cue was shown and $V=0$ to mean that it was not. The quantity of interest is then $P(C=1|A=1,V=1)$, which is the probability of a correct outcome given that both cues were presented. Application of Equation 7 would seem straightforward, except that the probabilities on the right hand side of the equation do not match the probabilities obtained from the unisensory trials. For instance, the term $P(C=1|A=1)$ corresponds to the probability of a correct choice when we know that the auditory cue was shown and we *do not know* whether the visual cue was shown or not; in contrast, the unisensory auditory experiment yields $P(C=1|A=1,V=0)$, which is the probability of a correct choice when we know that the auditory cue was shown and that the visual cue was not. The same goes for the visual unisensory experiment. So, Equation 7 is not applicable to the unisensory data at hand.

However, the conditional probability for the multicue trials can be cast in terms of the probabilities obtained from the unisensory trials. To do this, first write down the four versions of Equation 9 (with V taking the place of B) that result when the four possible combinations of the A and V binary events are considered. Then combine those expressions to eliminate the terms conditioned on A alone or V alone. After some algebra, the result is

$$P(1|1,1) = \frac{P(1|1,0)\,P(1|0,1)}{P(1|1,0)\,P(1|0,1) + P(0|1,0)\,P(0|0,1)\,\frac{P(1|0,0)}{P(0|0,0)}} \tag{19}$$

where we have used the abbreviations

$$\begin{aligned} P(1|1,1) &= P(C=1|A=1,V=1) \\ P(0|1,0) &= P(C=0|A=1,V=0) \end{aligned}$$

and so on, always assuming the same order, $(C|A,V)$. Equation 19 has the exact same form as Equation 7 but with the terms on the right hand side correctly matching the knowledge obtained

from the two unisensory experiments. Another notable aspect of this expression is that the terms that previously represented the priors for correct and incorrect choices, $P(C = 1)$ and $P(C = 0)$, have now turned into $P(C = 1|A = 0, V = 0)$ and $P(C = 0|A = 0, V = 0)$, which are the probabilities for correct and incorrect choices when neither cue is shown, which simply correspond to the chance levels of the experiment. The multisensory curves in this example (Fig 6d, light and dark red curves) were obtained by plugging the performance values in the simulated unisensory conditions into the above expression, along with 50% chance probabilities for correct and incorrect choices.

Equation 19 is an alternative formulation of the conditional independence relationships derived earlier (Equation 7). Although the data that these two expressions use is slightly different, they give rise to the same bounded arithmetic between the combined probabilities. This latest version matches the typical structure of a multisensory experiment, wherein the performance observed with a multisensory cue is compared to that obtained with separate unisensory cues. However, consistent with the earlier expressions, deviations from the predicted accuracy should be indicative of functional interactions between the two cues. This benchmark for multisensory interacton should be applicable under a wide range of conditions; namely, whenever (1) the primary measurements can be framed in terms of probabilities, and (2) chance performance is well defined within the experimental design.

How past events interact to influence ongoing performance

Much is known about the mechanisms that guide visuomotor behaviors. For instance, it is well established that the decision of where to look next depends in lawful ways on two factors, the salience of the objects in view, and the subject's internal goals, e.g., what a person might be looking for, such as a pen on a desk, a face in a crowd, and so on [28, 29, 30]. However, more recently it has been recognized that there is a third, independent factor that also plays a key role in guiding our attention and eye movements, which is generally referred to as selection history [31, 32, 33]. This includes effects attributable to recent experiences, such as past stimuli seen or past rewards gained or lost, that do not directly affect goals and intentions. The signature manifestation of a relevant history variable is a difference in performance when the variable is repeated across trials (e.g., successive red targets) versus when it switches value (e.g., a red target following successive green targets). The final example illustrates the application of conditional independence to the problem of disentangling how different types of selection history interact to strongly influence a subject's behavior in a visuomotor task.

Consider a simple search task in which the objective is to locate the oddball [34, 35], the object that has a different color (Fig 7a). After the subject fixates a central spot, a cue stimulus is briefly displayed and then masked, and the subject is required to make an eye movement to the location where the oddball was presented. From trial to trial, the oddball position varies randomly between four possible locations (up, down, left, right) and its color varies randomly between red (with green distracters) and green (with red distracters). The cue presentation period is short enough, and the chromatic contrast between target and distracters small enough, that average performance is above chance (25% correct) but well below 100% correct.

Although this particular task is hypothetical, it captures key elements of real tasks used to characterize visuomotor performance and its underlying neural mechanisms. That is, behavior in this task was simulated with a simple model (Methods) that generated many choices, one trial after another, such that selection history effects would be observed based on target features, target location, and outcome (correct/incorrect) — dependencies found in many prior studies [35, 36, 37, 38, 39, 40]. The underlying mechanisms whereby the model produces such sequential effects are

Fig 7. Selection history effects in a hypothetical oddball search task. a, Sequence of events in the oddball task. After the subject fixates their gaze on a central point (Fixation), the cue stimulus is briefly presented (Cue) and followed by a mask (Mask). The subject then indicates the location of the target with an eye movement (Saccade) and receives feedback indicating what the true location was (not shown). **b**, Fraction correct (x axis) for choices preceded by different color histories (y axis) going 1–4 trials back, with prior target stimuli having been of the same (S) or different (D) color as the target in the current trial. **c**, As in **b**, but for performance conditioned on the history of preceding target locations. In this case, the prior targets could have been at the same (S) or at a different (D) location as the target in the current trial. **d**, As in **b**, but for performance conditioned on the history of prior outcomes. In this case, the 1–4 trials preceding the current one could have been correct choices (C) or errors (E). Results in b–d are from the same simulation, which consisted of 100,000 trials. Black lines indicate 95% confidence intervals.

most important, but to appreciate them we must first describe how those history effects manifest in the data.

To reveal the influence of selection history, we parsed the simulated choices (100,000 trials) according to their preceding histories as done in real psychophysical experiments (Fig 7b–d). In the case of target color, for instance, we computed the fraction of choices that were correct given that the color of the oddball in the previous trial was the same as in the current trial (Fig 7b, case S), or the fraction of choices that were correct given that the color of the oddball in the previous 3 trials was different from that in the current trial (Fig 7b, case DDD), and so forth. In the simulated data, performance got progressively better as the target color was repeated (Fig 7b, red bars going from 0.62 for S to 0.79 for SSSS), but got progressively worse as more color repetitions were followed by a switch in color (Fig 7b, blue bars going from 0.39 for D to 0.31 for DDDD). Analogous sorting of the simulated trials but now according to target location demonstrated a similar trend: the fraction correct increased noticeably when the target appeared repeatedly at the same location (Fig 7c, red bars going from 0.68 for S to 0.86 for SSSS); in contrast, performance dropped considerably when the prior target location was different from that in the current trial (Fig 7c, case D), and a location switch that followed up to 4 location repeats produced somewhat worse performance (Fig 7c, blue bars going from 0.45 for D to 0.38 for DDDD). Finally, the performance of the model also varied according to outcome history: choice accuracy was higher in trials that followed a correct choice than in trials that followed an error (Fig 7d, compare cases C and E), but the difference was relatively modest (0.54 vs. 0.47) and further repetitions had a barely visible impact ($\lesssim 0.01$) in either direction.

Based on these descriptive results, it is clear that the hypothetical subject's performance on the current trial depends on the color of the prior targets, the location of the prior targets, and to a lesser degree, whether those prior trials were successful or not. But what drives these dependencies? And, are the underlying mechanisms responsible for these three effects independent or interrelated? These are questions that would be asked if the data were from a real experiment, and

conditional independence can be used to address them.

The above analyses produced three sets of probabilities associated with the three individual selection histories. To understand the dependencies between the three variables (color, location, outcome), we can compute the probability of a correct choice given any given pairwise combination of such history terms and compare it to the probability predicted by assuming conditional independence. For example, consider the probability of success given that the oddball color in the prior trial was the same as in the current trial

$$P(C = 1|\text{S color}) = 0.62 \tag{20}$$

and the probability of success given that the prior choice was correct

$$P(C = 1|\text{C outcome}) = 0.54 \tag{21}$$

which were already obtained (Fig 7b, d). Then compute the probability of success given the joint condition

$$P(C = 1|\text{S color}, \text{C outcome}) \tag{22}$$

in two ways, (1) as measured from the data, by considering all the trials in which the preceding trial was both correct and of the same target color, and (2) as predicted by inserting the values in Equations 20 and 21 into Equation 7. The resulting numbers are 0.80 and 0.65, respectively (Fig 8a, top row, point marked SC). The difference between them is indicative of an interdependence between color and outcome. However, a more complete picture is obtained by repeating the same process for all the possible combinations of color and outcome going back one trial (Fig 8a, top row, points marked DE, DC, SE, SC). The resulting four points are clearly off the diagonal, and the actual, measured probabilities differ from the predicted ones in a particular pattern: the former deviate more from the overall probability of success, $P(C = 1) = 0.51$, which is simply the prior for a correct choice. That is, knowledge of the true joint history is more informative — it produces a more certain prediction — than expected based on conditional independence.

This pattern becomes more evident when the same comparison is made for joint histories going back 2, 3, and 4 trials (Fig 8, a, rows 2–4), which give rise to 4^2, 4^3, and 4^4 different conditional probability terms, respectively. For example, the term

$$P(C = 1|\text{DDS color}, \text{ECC outcome}) \tag{23}$$

gives the probability of success given that the target colors of the three preceding trials were different, different, and the same as that in the current trial, and that the outcomes of those same preceding trials were error, correct, and correct. In all cases, the regression lines (Fig 8a, blue lines) for the resulting clouds of points cross the diagonal near the value of the prior (0.51) and have similar slopes < 1, confirming that the color and outcome histories are far from being conditionally independent. This suggests that there is some form of interaction between target color and outcome in the underlying mechanism whereby they influence the current choice.

A very different result is obtained when the same analysis is performed by combining target location and outcome histories (Fig 8b). Now the data points line up tightly along the diagonal, indicating high consistency between the empirical and predicted success probabilities. For histories going back 1–3 trials, most deviations from the diagonal are within the 95% confidence intervals expected simply from the finite numbers of trials sampled in each condition; it is only when histories go back four trials that large deviations can be observed, typically for points that have large uncertainties (Fig 8b, black dots in bottom row indicate 95% confidence interval span

Fig 8. Interactions between selection history variables manifest as deviations from the predictions based on conditional independence. Each point indicates the probability of a correct choice given a joint set of histories, either measured from appropriately sorted, simulated trials (x axes) or as predicted by assuming conditional independence between the individual histories (y axes). **a**, Probabilities based on joint consideration of target color and choice outcome in prior trials. Rows correspond to histories going back 1–4 trials ($H = 1$, $H = 2$, $H = 3$, $H = 4$, respectively). Lines behind circles indicate 95% confidence intervals based on binomial statistics. Black dots are data points that had large uncertainties (95% confidence interval span > 0.15). Dashed diagonal lines mark equality between x- and y-axis values. Blue lines indicate linear regressions based on red circles. Regression slopes are indicated (β_2, with 95% confidence intervals). **b**, Probabilities based on joint consideration of target location and choice outcome in prior trials. **c**, Probabilities based on joint consideration of target color and location in prior trials.

> 0.15). Overall, the data suggest that target location and outcome do not interact; they influence
each choice independently.

Thus, based on their individual effects, it would appear that color and target histories exert
qualitatively similar influences on task performance (Fig 7a, b). However, this superficial simi-
larity is deceiving. Analyses of their effects jointly with outcome — when benchmarked against
the performace expected based on conditional independence — reveals a stark difference between
them (Fig 8a vs. b). And indeed, within the dynamics of the model, color and location histories
act in entirely different ways mechanistically.

The model is very simple (Methods). It produces two kinds of response, guesses and informed
choices. (1) During guesses, the model picks one of the four possible response locations based
on where the oddball has previously appeared. Each of the four locations has a bias, and the
probability of guessing toward location j is directly proportional to its bias. The biases are updated
after every trial; whenever the oddball is presented at location j, the bias in favor of location j is
increased and others are decreased. (2) During informed choices, there is a 0.95 probability of
making a correct response toward the oddball and a 0.05 probability of making an erroneous
response elsewhere (a lapse). The key to the model is that the probability of making an informed

choice versus a guess depends on the match between the current color of the oddball and the prior colors experienced recently. Each of the two oddball colors (red and green) has a sensitivity. In each trial, the probability of making an informed choice depends on the sensitivity associated with the color of the current target: the probability is high if the oddball is red and the sensitivity for red is high, which is typically the case when prior targets have been red; and conversely, the probability is low if the oddball is red and the sensitivity for red is low, which is typically the case when prior targets have been green. Similarly when the oddball is green. Crucially, the sensitivities are reinforced after correct responses only. After a correct choice with a red oddball, the sensitivity to red is increased and that for green is decreased. Similarly for green. After an incorrect choice, both sensitivities decay slightly toward their neutral state.

In summary, the model simulates a subject that implements a simple strategy: if target and distracters can be discriminated (when sensitivity is high), the choice is informed and has a 95% chance of being correct, but if target and distracters cannot be discriminated (when sensitivity is low), the response is a guess that is biased toward recent target locations. Choice outcome enters into the dynamics, albeit indirectly; its only role is to gate the update of the color sensitivities. This is why the apparent effect of outcome history alone is weak and short-lived (Fig 7d).

So, color and location histories derive from fundamentally different mechanisms for driving performance, and this is highly consistent with the qualitative differences observed when success probability is conditioned on their respective joint histories combined with outcome (Fig 8a vs. b). Interestingly, further indication of their distinct mechanistic roles is obtained by a similar analysis in which success probability is conditioned on the joint history of color and location (Fig 8c). In this case, the data points also line up along the diagonal, indicating that, overall, color and location histories exert independent influences on performance.

In conclusion, this example illustrates the use of conditional independence as a benchmark for gaining some insight about how history variables modulate visuomotor performance. Specifically, analysis of the synthetic data suggested that the effect of color history was mediated by outcome (or reward, in a typical experiment), whereas that of location history was not, and furthermore, it indicated that the color and location influences were largely independent. All this was indeed in agreement with the model implementation. In additional computational experiments, alterations to the model produced corresponding alterations in the pattern of deviations from conditional independence, but the interpretation of the latter was always consistent with the former. The broader lesson is that, applied in this way, conditional independence may be a useful tool for assessing the presence of functional interactions between stochastic variables that modulate visuomotor performance.

Conclusion

In this report, we discussed the concept of conditional independence and its application as a tool for analyzing a variety of evidence integration problems in biomedical sciences. The underlying statistical question is both intuitive and general: what can be inferred from two pieces of evidence known simultaneously relative to what can be inferred from them when known individually? Specifically, we showed that when two variables, A and B, are conditionally independent with respect to a third one, C, predicting the state of the latter based on observation of the former is greatly simplified in terms of both the computational recipe (Equation 7) and the data ($P(C|A)$ and $P(C|B)$) that are necessary for generating the prediction. In two examples, diagnosing a disease and inferring biological age, we demonstrated application of the analytical results to situations in which the primary goal would be precisely to generate such predictions. In two additional examples, such predictions served as benchmarks for comparison to empirical data. In these cases,

the goal was to determine whether two relevant input variables exerted independent effects on an output variable (accuracy in visuomotor performance) or, alternatively, whether they interacted in some way. The chosen examples are neither exahaustive nor highly detailed; they are meant as proofs of concept, and as guidelines for future applications to real data.

The broader theme is that conditional independence is a useful statistical tool for understanding how multiple sources of evidence are, or can be, integrated. This is certainly not a new mathematical insight [1, 41]. However, to our knowledge, the use of conditional independence for analyzing experimental data in psychology or biomedicine is at the very least uncommon, and perhaps altogether novel. Measuring a correlation coefficient is a first step toward understanding the functional or causal relationship between two variables. Likewise, for some problems such as characterizing the effectiveness of multisensory integration or determining how selection histories modulate task performance, determining conditional independence can be a first step toward understanding the functional or causal relationships between three or more variables. In this context, conditional independence provides an intuitive reference for determining whether distinct pieces of evidence interfere with each other, amplify their individual effects, or act independently when they are considered simultaneously.

Acknowledgments

We thank Denise Anderson for logistical and administrative support.

References

1. Dawid AP. Conditional independence in statistical theory. J R Statist Soc B. 1979; 41:1–31.

2. Press WH, Flannery BP, Teukolsky SA, Vetterling WT. Numerical Recipes in C. 2nd ed. New York: Cambridge University Press; 1992.

3. Sedlmeier P, Gigerenzer G. Teaching Bayesian reasoning in less than two hours. J Exp Psychol Gen. 2001; 130:380–400.

4. Zhu L, Gigerenzer G. Children can solve Bayesian problems: the role of representation in mental computation. Cognition. 2006; 98:287–308.

5. Altman DG, Bland JM. Diagnostic tests. 1: Sensitivity and specificity. BMJ. 1994; 308:1552. doi: 10.1136/bmj.308.6943.1552.

6. Fletcher RH, Fletcher SW. Clinical epidemiology: the essentials. 4th ed. Baltimore: Lippincott Williams & Wilkins; 2005.

7. Altman DG, Bland JM. Diagnostic tests 2: Predictive values. BMJ. 1994; 309:102. doi: 10.1136/bmj.309.6947.102.

8. Balayla J. Derivation of generalized equations for the predictive value of sequential screening tests. arXiv:2007.13046v1 [Preprint]. 2007. Available from: https://doi.org/10.48550/arXiv.2007.13046.

9. Khan SS, Singer BD, Vaughan DE. Molecular and physiological manifestations and measurement of aging in humans. Aging Cell. 2017; 16:624–633.

10. Ferrucci L, Gonzalez-Freire M, Fabbri E, Simonsick E, Tanaka T, Moore Z et al. Measuring biological aging in humans: A quest. Aging Cell. 2020; 19:e13080.

11. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. Cell. 2013; 153:1194–1217.

12. Jylhävä J, Pedersen NL, Hägg S.. Biological age predictors. EBioMedicine. 2017; 21:29–36.

13. Klemera P, Doubal S. A new approach to the concept and computation of biological age.
Mech Ageing Dev. 2006; 127:240–248.

14. Belsky DW, Caspi A, Houts R, Cohen HJ, Corcoran DL, Danese A, et al. Quantification of
biological aging in young adults. Proc Natl Acad Sci USA. 2015; 112:E4104-10.

15. Levine ME. Modeling the rate of senescence: Can estimated biological age predict mortality
more accurately than chronological age? J Gerontol A Biol Sci Med Sci. 2013; 68:667–674.

16. Levine ME, Lu AT, Quach A, Chen BH, Assimes TL, Bandinelli S, et al. An epigenetic
biomarker of aging for lifespan and healthspan. Aging (Albany NY). 2018; 10:573–591.

17. Stein BE, editor. The New Handbook of Multisensory Processing. 1st ed. Cambridge: The
MIT Press; 2012.

18. Fetsch CR, DeAngelis GC, Angelaki DE. Bridging the gap between theories of sensory cue
integration and the physiology of multisensory neurons. Nat Rev Neurosci. 2013; 14:429–
442.

19. Wallace MT, Woynaroski TG, Stevenson RA. Multisensory integration as a window into
orderly and disrupted cognition and communication. Annu Rev Psychol. 2020; 71:193–219.

20. Ernst MO, Banks MS. Humans integrate visual and haptic information in a statistically opti-
mal fashion. Nature. 2012; 415:429–433.

21. Alais D, Burr D. The ventriloquist effect results from near-optimal bimodal integration. Curr
Biol. 2004; 14:257–262.

22. Ernst MO. Optimal multisensory integration: assumptions and limits. In: Stein BE, editor.
The New Handbook of Multisensory Processes. Cambridge: The MIT Press; 2012. pp. 527–
543.

23. Miller JO. Divided attention: evidence for coactivation with redundant signals. Cognitive
Psychology 1982; 14:247–279

24. Beauchamp MS. Statistical criteria in FMRI studies of multisensory integration. Neuroinfor-
matics. 2005; 3:93–113.

25. Stein BE, Stanford TR, Ramachandran R, Perrault TJ Jr, Rowland BA. Challenges in quanti-
fying multisensory integration: alternative criteria, models, and inverse effectiveness. Exp
Brain Res. 2009; 198:113–126.

26. Otto TU, Dassy B, Mamassian P. Principles of multisensory behavior. J Neurosci. 2013;
33:7463–7474.

27. Smyre SA, Wang Z, Stein BE, Rowland BA. Multisensory enhancement of overt behavior
requires multisensory experience. Eur J Neurosci. 2021; 54:4514–4527.

28. Itti L, Koch C. Computational modelling of visual attention Nat Rev Neurosci. 2001; 2:194–
203.

29. Theeuwes J. Top-down and bottom-up control of visual selection. Acta Psychol (Amst). 2010;
135:77–99.

30. Wolfe JM, Horowitz TS. Five factors that guide attention in visual search. Nature Hum
Behav. 2017; 1:0058.

31. Awh E, Belopolsky AV, Theeuwes J. Top-down versus bottom-up attentional control: a failed
theoretical dichotomy. Trends Cogn Sci. 2012; 16:437–443.

32. Theeuwes J. Feature-based attention: it is all bottom-up priming. Philos Trans R Soc Lond B

Biol Sci. 2013; 368:20130055.

33. Anderson BA, Kim H, Kim AJ, Liao MR, Mrkonja L, Clement A, Grégoire L. The past, present, and future of selection history. Neurosci Biobehav Rev. 2021; 130:326–350.

34. Scerra VE, Costello MG, Salinas E, Stanford TR. All-or-none context dependence delineates limits of FEF visual target selection. Curr Biol. 2019; 29:294–305.e3.

35. Oor EE. The roles of distinct attentional mechanisms and their neural correlates in the Frontal Eye Field to perceptual decision making and oculomotor choice. Ph.D. Thesis, Wake Forest University. 2022. Available from: https://www.proquest.com/dissertations-theses/roles-distinct-attentional-mechanisms-their/docview/2669597234/se-2?accountid=14868

36. Maljkovic V, Nakayama K. Priming of pop-out: II. The role of position. Percept Psychophys. 1996; 58:977–991.

37. McPeek RM, Maljkovic V, Nakayama K. Saccades require focal attention and are facilitated by a short-term memory system. Vision Res. 1999; 39:1555–1566.

38. Busse L, Ayaz A, Dhruv NT, Katzner S, Saleem AB, Schölvinck ML, Zaharia AD, Carandini M (2011) The detection of visual contrast in the behaving mouse. J Neurosci 31:11351–11361.

39. Hauser CK, Zhu D, Stanford TR, Salinas E. Motor selection dynamics in FEF explain the reaction time variance of saccades to single targets. eLife. 2018; pii:e33456.

40. Hermoso-Mendizabal A, Hyafil A, Rueda-Orozco PE, Jaramillo S, Robbe D, de la Rocha J. Response outcomes gate the impact of expectations on perceptual decisions. Nat Commun. 2020; 11:1057.

41. Pearl J. Causality: Models, Reasoning, and Inference. 1st ed. Cambridge: Cambridge University Press; 2000.