**Sex-specific relationship between maternal and neonate cortisol in**

**a free-ranging large mammal**

Bawan Amin[1, 5*], Ruth Fishman[2], Matthew Quinn[1, 3], Devorah Matas[2], Rupert Palme[4], Lee Koren[2],

Simone Ciuti[1]

1) Laboratory of Wildlife Ecology and Behaviour, School of Biology and Environmental Science,

University College Dublin, Dublin, Ireland

2) Faculty of Life Sciences, Bar-Ilan University, Ramat-Gan, Israel

3) Department of Biosciences, Durham University, Durham, United Kingdom

4) Unit of Physiology, Pathophysiology and Experimental Endocrinology, Department of Biomedical

Sciences, University of Veterinary Medicine, Vienna, Austria

5) Faculty of Social and Behavioural Sciences, Utrecht University, Utrecht, Netherlands

* Correspondence: b.k.r.amin@uu.nl

**Acknowledgements**

We thank the Office of Public Works (OPW), Ireland, for funding (grant no. R18625) and support. We

thank the SBES in University College Dublin (UCD), and the British Deer Society (R21845) for funding

this project. We also would like to thank the Higher Education Authority of Ireland for providing a

Covid-19 related Research Cost Extensions to BA. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript. We declare that none

of the authors have a conflict of interest.

**Open science and data availability statement**

Following Kane & Amin (2023), all versions, updates and additional material are stored transparently

on OSF (DOI 10.17605/OSF.IO/4YMC8). In addition, we have published the data and the script used

for the analysis under CC-BY 4.0 license, also on OSF (Amin et al., 2023). These can be found via the

following link: https://doi.org/10.17605/OSF.IO/WYNKE.

**Author contributions**

BA: Conceptualization, data curation, formal analysis, investigation, visualization, writing -original

draft preparation; RF: Conceptualization, writing – original draft preparation, writing – review &

editing; MQ: Data curation, investigation, writing – review & editing; DM: Investigation,

methodology, writing – review & editing; RP: Investigation, methodology, writing – review & editing;

LK: Investigation, methodology, writing – review & editing; SC: Conceptualization, funding

acquisition, resources, project administration, supervision, writing – review & editing

**Abstract**

Maternal phenotypes can have long-term effects on the offspring phenotype starting at gestation.

Proximate mechanisms include foetal hormone levels affected by maternal levels. Recent studies

have highlighted that these maternal effects may be different between the sexes. How maternal

glucocorticoid (GC) levels relate to foetal levels is, however, still not completely understood. Here

we related, for the first time in a free-ranging large mammal, the fallow deer (*Dama dama*),

maternal GC levels with foetal *in utero* GC levels by using novel state-of-the-art methods. We found

that foetal GC levels were positively associated with maternal GC levels, but only in females. These

findings highlight sex differences, which may have evolved to optimize male growth at the cost of

survival.

**Keywords**

Glucocorticoids, free-ranging, fallow deer, hair-testing, faecal metabolites

**Introduction**

Parental phenotypes can be drivers of offspring variation, and in many mammalian species, these parental effects are often labelled as maternal effects since it is the mother that mostly takes care of the offspring. Maternal effects are seen as an adaptive way in which mothers can either fine tune their offspring for the current environmental conditions (Sheriff & Love, 2013), or maximise their own reproductive output (Marshall & Uller, 2007). These effects on the offspring are present from the earliest stages of development and can have long lasting influence on offspring phenotype (Seckl & Meaney, 2004; Weinstock, 2008).

Acute stress typically leads to an increase in circulating glucocorticoid (GC) levels due to activation of the hypothalamic-pituitary-adrenal (HPA) axis (Sapolsky et al., 2000; Weinstock, 2008). Maternal GC levels during gestation can affect offspring phenotype in many ways, including affecting offspring birth weight and HPA-axis reactivity (Dantzer et al., 2013; Seckl & Holmes, 2007; Seckl & Meaney, 2004) and these effects can linger until offspring are well into adulthood (e.g. 8), which is also referred to as foetal programming (Seckl & Holmes, 2007). There is increasing evidence, however, that there are sex differences in foetal programming (Braithwaite et al., 2018; Fishman et al., 2022; Liu et al., 2001).

Most studies investigating hormonal maternal effects investigate offspring GC levels after birth and most evidence comes from either clinical studies on humans (Gitau et al., 1998), rodents (Weinstock, 2008) or birds (Groothuis et al., 2019; Jenkins et al., 2014), with a paucity of data on free-ranging mammals. This is usually due to the challenges of measuring maternal and foetal hormone levels during gestation in non-captive mammals. Recent developments, however, have enabled measuring foetal hormonal levels post-parturition by quantifying steroids in neonate hair (Amin et al., 2021; Fishman et al., 2019; Kapoor et al., 2016). Hair steroid levels represent long-term levels, accumulated over the last weeks to months (Gormally & Romero, 2020), and therefore reflect *in utero* integrated levels when quantified in neonates (Amin et al., 2021). Whether these levels are associated with circulating maternal GC levels, however, is currently not well understood.

Here we explored, for the first time in free-ranging large mammals, the relationship

between circulating maternal GC levels and offspring *in utero* accumulated GC levels. For that

purpose, we collected faeces of mothers during late gestation, from which we quantified GC

metabolites (Palme, 2019), and related that to their offspring hair GC levels (Amin et al., 2021),

which were collected during the first days post-parturition at neonate captures. We made sure to

sample females within their last five weeks of gestation, since that is when the foetus is fully covered

in fur (Chapman & Chapman, 1997). To allow for sex differences in the association between

maternal and foetal GC levels, we ran an additional analysis to explore differences between male

and female offspring.

**Methods**

*Study population*

We conducted this study in Phoenix Park, a 7.07 km$^2$ urban park located in Dublin, Ireland. There is a

resident population of free-ranging fallow deer that has been introduced in the 17$^{th}$ century, with a

population size of about 600 individuals in late summer, after fawn births (Griffin et al., 2022). Most

births occur between early and late June of each year. Fallow deer are a hider species and fawns

remain hidden, usually in tall grass or understory vegetation, away from the main doe herd during

the first weeks of life after which they are brought into the doe herd by their mothers (Chapman &

Chapman, 1997; Ciuti et al., 2006). Fawns are occasionally predated upon by red foxes (*Vulpes*

*vulpes*), the only natural predator in the park, and domestic dogs who are brought into the park by

public visitors.

*Faeces collection and analysis of maternal faecal cortisol metabolites (FCMs)*

Faecal samples were collected between 8 AM and 3 PM, from May 19th until May 29th 2020, when

the does were in late gestation. Groups of deer were observed from a distance of 50 meters using a

spotting scope to identify individuals (>80% of the population is identifiable via unique colour coded

ear-tags). Fresh faecal samples were collected within a minute of defecation and immediately stored

in zip locked bags. These were kept in a cooler bag until they could be stored in a freezer at -20˚C,

which was always within a few hours. Samples (see *Sample sizes*-section below) were kept frozen on

dry ice during transportation to the University of Veterinary Medicine (Vienna, Austria), where

cortisol metabolites in the samples were quantified (Palme, 2019). We added 5 mL of methanol

(80%) to weighed aliquots (0.5 g) of homogenized faecal samples, after which the samples were

shaken and centrifuged as previously described (Palme et al., 2013). FCMs were analysed with an 11-

oxoaetiocholanolone enzyme immunoassay (for details see Möstl et al., 2002), which has already

been validated for fallow deer (Konjević et al., 2011).

*Neonate hair collection and hair GC level quantification*

Neonate hair samples were collected during the fawn captures in June 2020. Fawns were routinely

captured with fishing nets, as part of the long-term management of the population. During handling,

we collected physiological and behavioural data (see Amin et al., 2021 (Amin et al., 2021) for full

details on the capture protocol), as well as hair samples (>100 mg) from the belly of the fawn using

an electric trimmer (Wahl model 9639; Wahl Clipper Corporation). We extracted cortisol levels from

fawn hair using a standardized protocol for hair-testing (e.g. 15,25), by using commercial enzyme

immunoassay (EIA; Salimetrics; Ann Arbor, MI, USA, item no. 1-3002) kits. Full details on these

extractions and validations are described in Amin et al., 2021 and in the supplementary material

(S1).

*Mother-fawn pairing*

Mother-fawn pairs were based on field observations with data collection starting in July, when

young fawns were making their first entrances into the female herds. For details regarding the

pairing, see Griffin et al., 2022 and Supplementary S1.

*Sample sizes*

We collected and analysed a total of 164 faecal samples of 99 different pregnant does. For

individuals with multiple values, we took the mean of these values to create a dataset that had one

estimate for each individual.  Out of these, we were able to pair 41 does with their fawns through

the mother-fawn observations during summer. We removed one extreme outlier from the male

neonate cortisol dataset (value: 29.5 pg/mg hair; range of other values: 7.7-18.3 pg/mg hair).

Running our analysis with and without this outlier revealed that removing this outlier did not affect

the conclusions of our analysis. This led to our final sample size of 40 fawns (18 females and 22

males) paired with their mothers.

*Statistical analysis*

All analyses were performed in RStudio (Version 1.3.1093) using R version 4.0.2 (R Core Team, 2020).

We first checked whether maternal or neonatal GC levels differed between the offspring sexes by

performing a t-test, considering unequal variances.

To investigate the relationship between neonate hair GC and maternal FCMs, we ran a linear

model for each fawn sex separately. In both models, we had the neonate hair GC as the response

variable and the maternal FCMs as the explanatory variable. Maternal FCM levels were log-

transformed to improve model fit, since these suffered from a slight positive skew. Model

assumptions were checked with the *DHARMa*-package (Version 0.4.3; Hartig, 2021) and were

successfully met. Statistical inferences were made based on the estimate and the associated 95%

Confidence Interval from the models. During the preliminary analysis, we included the number of

days between the collection of mothers' faeces and the day of fawn birth in our models. This was to

account for potential variation in FCM levels as a function of gestational day of collection (i.e. mid

May vs late May when closer to parturition). This additional predictor had no clear effect and made

our models worse (higher AIC); we therefore decided not to include it. We plotted our results using

the *ggplot2*-package (Wickham, 2016).

Finally, we decided to run an additional model *post-hoc*, including both sexes in the same

model, to further investigate whether the slopes of our models were different between male and

female offspring. We ran a linear model with neonate GC as the response variable. As explanatory

variables, we included maternal FCMs (log transformed), fawn sex and the interaction between both

these explanatory variables.

**Results**

Considerable variation was found in both the individual maternal FCM levels (range: 91-608 ng/g

faeces) as well as the neonate hair GC levels (range: 6.50-18.30 pg/mg hair). We found no clear sex

differences, neither for mean maternal FCM levels (t-test; t = 0.56, 95% CI [-51.99, 91.50], p = 0.58)

nor mean neonate hair GC levels (t-test; t = 0.26, 95% CI [-1.74, 2.25], p = 0.80).

We found that female hair cortisol was positively associated with maternal FCMs (LM: $\beta$ =

5.88, 95% CI [0.20, 11.57], p = 0.04, N = 18, $R^2_{adjusted}$ = 0.18; Fig. 1A). For males, however, there was

no clear association between both measures (LM: $\beta$ = 0.59, 95% CI [-1.32, 2.49], p = 0.53, N = 22,

$R^2_{adjusted}$ = -0.03; Fig. 1B). Our post-hoc model confirmed that there was indeed an interaction

between fawn sex and maternal levels, where the slope between maternal and foetal levels was

lower in male offspring compared to female offspring (LM: $\beta$ = -5.30, 95% CI [-10.39, -0.20], p = 0.04,

N = 40, $R^2_{adjusted}$ = 0.10).

**Figure 1:** The relationship between maternal faecal cortisol metabolites (FCMs) and neonate hair

GCs, for female (A) and male (B) offspring. Trendlines and their 95% confidence intervals are fitted in

each plot. Solid trendline indicates a statistically meaningful relationship.

**Discussion**

In this study we found that maternal FCM levels were positively related to foetal GC levels, but only in females. In male offspring, we found no clear relationship between maternal and foetal GC levels. These findings suggest that maternal GC levels may affect male offspring less then female offspring. We discuss here how, from an evolutionary point of view, sex-specific mechanisms may have evolved to optimize life-history trade-offs.

Like many other species, fallow deer have a skewed mating distribution (McElligott et al., 2001; Moore et al., 1995). While females tend to produce one fawn per year (Chapman & Chapman, 1997), male reproduction skew is steep. Only a small proportion of the males reproduces, with the majority having no or very little offspring (Ciuti et al., 2011; McElligott et al., 2001; Moore et al., 1995). Therefore, there may be a strong selection for males to prioritize growth over survival, since low quality sons will end up costing resources, with no fitness benefits in the long run. A recent study has shown that male birthweights in our populations are indeed higher than female birthweights (Griffin et al., 2023).

GC levels play a key role during gestation and have major effects on offspring phenotype early in life. In addition to being crucial for organ development during the late stages of pregnancy (Kitterman et al., 1981; Liggins, 1994), GCs play a role in fighting inflammation (Auphan et al., 1995) and thereby help to keep the foetus vital. At the same time, high GC levels may restrict growth (Edwards et al., 1996; Meakin et al., 2021; Seckl & Holmes, 2007). As males tend to prioritize growth (Meakin et al., 2021), it may be that there would be sex differences in the mechanisms that evolved. For example, maternal GC contribution may be limited in male foetuses to prevent growth retardation, at the cost of prenatal survival. This may contribute in some degree to why males generally face higher prenatal mortality rates (Desportes et al., 1994; Eriksson et al., 2010; Kruuk et al., 1999).

The mammalian placenta likely plays a crucial role in mediating these sex differences. Previous studies, in rodents and humans, have shown sex differences in the structure and activity of

the placenta (Meakin et al., 2021; Murphy et al., 2003; O'Connell et al., 2013; Rosenfeld, 2015;

Wieczorek et al., 2019). There is also evidence that maternal GCs may transport more easily through

female placentas than males' (Wieczorek et al., 2019). This may be, for instance, through different

levels of expression and activity of enzymes or transporters that mitigate different maternal GC level

spill over between males and females.

One of the key enzymes, 11β-hydroxysteroid dehydrogenase type 2 (11β-HSD2), inactivates

cortisol to cortisone (Edwards et al., 1996; Tomlinson & Stewart, 2001). Placental 11β-HSD has been

frequently shown to be positively related to birthweight (Edwards et al., 1996; Meakin et al., 2021;

Seckl et al., 2000; Seckl & Meaney, 2004), thus likely playing a role in preventing growth restriction.

Previous studies have shown that male foetuses have higher expression or activity of 11β-HSD2 in

response to maternal GC surges (Murphy et al., 2003; Wieczorek et al., 2019). Similarly in ovine,

male foetuses increased 11β-HSD2 in response to maternal dexamethasone administration, whereas

female foetuses did not (Braun et al., 2009). This may explain why a different ovine study also found

that, following an antenatal GC treatment, growth restriction was lower in males than in females

(Miller et al., 2012). In addition, male mice were shown to also have increased ATP-binding cassette

transporters, which mediate GC efflux toward maternal circulation (Wieczorek et al., 2019),

indicating that there are different mechanisms through which males may be able to limit maternal

GC levels (Montano et al., 1993). This altogether may explain why females may have higher foetal

survival (Meakin et al., 2021), whereas males may be larger and faster growing.

We acknowledge that there are some shortcomings to this study. We report patterns taken

over only one year, whereas ecological patterns may vary between years. Furthermore, we have

modest sample sizes (due to the difficulty of the design), which restrict us in doing more elaborate

analyses. Nevertheless, as one of the first studies quantifying the relationship between maternal and

foetal GC levels non-invasively, we do provide novel insights into how these fundamental

relationships may function in a free-ranging large mammal population. It is important to note that

the nature of the study described here is exploratory, meaning that it generates hypotheses that can

be tested in the future.

To conclude, we explored in this study the relationship between maternal and foetal GC

levels in a free-ranging population of fallow deer. Our findings suggest that there are sex differences

in the underlying evolutionary processes, which may have optimized growth at the cost of survival

for male offspring, whereas the opposite is true for female offspring. Most of the existing literature

uses the rodent or human model, with very little existing studies on other taxa. Although some

features of the placenta may be conserved, there are also evolutionary differences between species

(Fowden, 2003; Rosenfeld, 2015). Non-model species may provide different results (Fishman et al.,

2019), through which we can gain a better fundamental understanding of evolutionary mechanisms.

By studying free-ranging fallow deer, this study therefore also provides novel insights into possible

sex-specific maternal effects in a non-model species, and provides a method that can be viable for

other populations as well, enabling the study of patterns rarely studied before in the wild.

**References**

Amin, B., Fishman, R., Quinn, M., Matas, D., Palme, R., Koren, L., & Ciuti, S. (2023). Data from: Sex-

specific relationship between maternal and neonate cortisol in a free-ranging large mammal.

*Open Science Framework*. https://doi.org/10.17605/OSF.IO/WYNKE

Amin, B., Jennings, D. J., Smith, A. F., Quinn, M., Chari, S., Haigh, A., Matas, D., Koren, L., & Ciuti, S.

(2021). In utero accumulated steroids predict neonate anti-predator response in a wild

mammal. *Functional Ecology*, *35*(6), 1255–1267. https://doi.org/10.1111/1365-2435.13790

Auphan, N., Didonato, J. A., Rosette, C., Helmberg, A., & Karin, M. (1995). *Immunosuppression by*

*glucocorticoids: Inhibition of NF-κB activity through induction of IκB synthesis*. *270*(5234),

286–290. https://doi.org/10.1126/science.270.5234.286

Braithwaite, E. C., Hill, J., Pickles, A., Glover, V., O'Donnell, K., & Sharp, H. (2018). Associations

between maternal prenatal cortisol and fetal growth are specific to infant sex: Findings from

the Wirral Child Health and Development Study. *Journal of Developmental Origins of Health*

*and Disease*, *9*(4), 425–431. https://doi.org/10.1017/S2040174418000181

Braun, T., Li, S., Sloboda, D. M., Li, W., Audette, M. C., Moss, T. J. M., Matthews, S. G., Polglase, G.,

Nitsos, I., Newnham, J. P., & Challis, J. R. G. (2009). Effects of maternal dexamethasone

treatment in early pregnancy on pituitary-adrenal axis in fetal sheep. *Endocrinology*,

*150*(12), 5466–5477. https://doi.org/10.1210/en.2009-0086

Chapman, D., & Chapman, N. (1997). Fallow deer: Their history, distribution and biology (2nd ed.).

*Coch-y-bonddu Books*.

Ciuti, S., Bongi, P., Vassale, S., & Apollonio, M. (2006). Influence of fawning on the spatial behaviour

and habitat selection of female fallow deer (Dama dama) during late pregnancy and early

lactation. *Journal of Zoology*, *268*(1), 97–107. https://doi.org/10.1111/j.1469-

7998.2005.00003.x

Ciuti, S., De Cena, F., Bongi, P., & Apollonio, M. (2011). Benefits of a risky life for fallow deer bucks

(Dama dama) aspiring to patrol a lek territory. *Behaviour*, *148*(4), 435–460.

https://doi.org/10.1163/000579511X563981

Dantzer, B., Newman, A. E. M., Boonstra, R., Palme, R., Boutin, S., Humphries, M. M., & McAdam, A.

G. (2013). Density triggers maternal hormones that increase adaptive offspring growth in a

wild mammal. *Science*, *340*(6137), 1215–1217. https://doi.org/10.1126/science.1235765

Desportes, G., Andersen, L. W., & Bloch, D. (1994). Variation in foetal and postnatal sex ratios in

long-finned pilot whales. *Ophelia*, *39*(3), 183–196.

https://doi.org/10.1080/00785326.1994.10429543

Edwards, C. R. W., Benediktsson, R., Lindsay, R. S., & Seckl, J. R. (1996). 11β-Hydroxysteroid

dehydrogenases: Key enzymes in determining tissue-specific glucocorticoid effects. *Steroids*,

*61*(4), 263–269. https://doi.org/10.1016/0039-128X(96)00033-5

Eriksson, J. G., Kajantie, E., Osmond, C., Thornburg, K., & Barker, D. J. P. (2010). Boys live dangerously

in the womb. *American Journal of Human Biology*, *22*(3), 330–335.

https://doi.org/10.1002/ajhb.20995

Fishman, R., Kralj-Fišer, S., Marglit, S., Koren, L., & Vortman, Y. (2022). Quantitative genetic study

suggests sex-specific genetic architecture for fetal testosterone in a wild mammal. *BioRxiv*

*(Preprint)*. https://doi.org/10.1101/2022.02.18.480975

Fishman, R., Vortman, Y., Shanas, U., & Koren, L. (2019). Non-model species deliver a non-model

result: Nutria female fetuses neighboring males in utero have lower testosterone. *Hormones*

*and Behavior*, *111*(February), 105–109. https://doi.org/10.1016/j.yhbeh.2019.02.011

Fowden, A. (2003). The insulin-like growth factors and feto-placental growth. *Placenta*, *24*(8–9),

803–812. https://doi.org/10.1016/S0143-4004(03)00080-8

Gitau, R., Cameron, A., Fisk, N. M., & Glover, V. (1998). Fetal exposure to maternal cortisol. *Lancet*,

*352*(9129), 707–708. https://doi.org/10.1016/S0140-6736(05)60824-0

Gormally, B. M. G., & Romero, L. M. (2020). What are you actually measuring? A review of

techniques that integrate the stress response on distinct time-scales. *Functional Ecology*,

*34*(10), 2030–2044. https://doi.org/10.1111/1365-2435.13648

Griffin, L. L., Haigh, A., Amin, B., Faull, J., Corcoran, F., Baker-Horne, C., & Ciuti, S. (2023). Does

artificial feeding impact neonate growth rates in a large free-ranging mammal? *Royal Society*

*Open Science*, *10*(3), 221386. https://doi.org/10.1098/rsos.221386

Griffin, L. L., Haigh, A., Amin, B., Faull, J., Norman, A., & Ciuti, S. (2022). Artificial selection in human-

wildlife feeding interactions. *Journal of Animal Ecology*, *91*(9), 1892–1905.

https://doi.org/10.1111/1365-2656.13771

Groothuis, T. G. G., Hsu, B. Y., Kumar, N., & Tschirren, B. (2019). Revisiting mechanisms and functions

of prenatal hormone-mediated maternal effects using avian species as a model.

*Philosophical Transactions of the Royal Society B: Biological Sciences*, *374*(1770), 20180115.

https://doi.org/10.1098/rstb.2018.0115

Hartig, F. (2021). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed) Regression*

*Models* (0.4.3). https://CRAN.R-project.org/package=DHARMa

Jenkins, B. R., Vitousek, M. N., Hubbard, J. K., & Safran, R. J. (2014). An experimental analysis of the

heritability of variation in glucocorticoid concentrations in a wild avian population.

*Proceedings of the Royal Society B: Biological Sciences*, *281*(1790), 20141302.

https://doi.org/10.1098/rspb.2014.1302

Kane, A., & Amin, B. (2023). Amending the literature through version control. *Biology Letters*, *19*(1),

20220463. https://doi.org/10.1098/rsbl.2022.0463

Kapoor, A., Lubach, G. R., Ziegler, T. E., & Coe, C. L. (2016). Hormone levels in neonatal hair reflect

prior maternal stress exposure during pregnancy. *Psychoneuroendocrinology*, *66*, 111–117.

https://doi.org/10.1016/j.psyneuen.2016.01.010

Kitterman, J. A., Liggins, G. C., Campos, G. A., Clements, J. A., Forster, G. C., Lee, C. H., & Creasy, R. K.

(1981). Prepartum maturation in fetal sheep: Relation of the lung to cortisol. *Journal of*

*Applied Physiology*, *51*(2), 384–390. https://doi.org/10.1152/jappl.1981.51.2.384

Konjević, D., Janicki, Z., Slavica, A., Severin, K., Krapinec, K., Božić, F., & Palme, R. (2011). Non-

invasive monitoring of adrenocortical activity in free-ranging fallow deer (Dama dama L.).

*European Journal of Wildlife Research*, *57*(1), 77–81. https://doi.org/10.1007/s10344-010-

0401-1

Koren, L., & Geffen, E. (2009). Androgens and social status in female rock hyraxes. *Animal Behaviour*,

*77*(1), 233–238. https://doi.org/10.1016/j.anbehav.2008.09.031

Kruuk, L. E., Clutton-Brock, T. H., Albon, S. D., Pemberton, J. M., & Guinness, F. E. (1999). Population

density affects sex ratio variation in red deer. *Nature*, *399*(6735), 459–461.

https://doi.org/10.1038/20917

Liggins, G. C. (1994). The role of cortisol in preparing the fetus for birth. *Reproduction, Fertility and*

*Development*, *6*(2), 141–150. https://doi.org/10.1071/RD9940141

Liu, L., Li, A., & Matthews, S. G. (2001). Maternal glucocorticoid treatment programs HPA regulation

in adult offspring: Sex-specific effects. *American Journal of Physiology-Endocrinology and*

*Metabolism*, *280*(5), E729–E739. https://doi.org/10.1152/ajpendo.2001.280.5.E729

Marshall, D. J., & Uller, T. (2007). When is a maternal effect adaptive? *Oikos*, *116*, 1957–1963.

https://doi.org/10.1111/j.2007.0030-1299.16203.x

McElligott, A. G., Gammell, M. P., Harty, H. C., Paini, D. R., Murphy, D. T., Walsh, J. T., & Hayden, T. J.

(2001). Sexual size dimorphism in fallow deer (Dama dama): Do larger, heavier males gain

greater mating success? *Behavioral Ecology and Sociobiology*, *49*(4), 266–272.

https://doi.org/10.1007/s002650000293

Meakin, A. S., Cuffe, J. S. M., Darby, J. R. T., Morrison, J. L., & Clifton, V. L. (2021). Let's talk about

placental sex, baby: Understanding mechanisms that drive female-and male-specific fetal

growth and developmental outcomes. *International Journal of Molecular Sciences*, *22*(12).

https://doi.org/10.3390/ijms22126386

Miller, S. L., Sutherland, A. E., Supramaniam, V. G., Walker, D. W., Jenkin, G., & Wallace, E. M. (2012).

Antenatal glucocorticoids reduce growth in appropriately grown and growth-restricted ovine

fetuses in a sex-specific manner. *Reproduction, Fertility and Development*, *24*(5), 753.

https://doi.org/10.1071/RD11143

Montano, M. M., Wang, M., & vom Saal, F. S. (1993). Sex differences in plasma corticosterone in

mouse fetuses are mediated by differential placental transport from the mother and

eliminated by maternal adrenalectomy or stress. *Journal of Reproduction and Fertility*, *99*,

283–290. https://doi.org/10.1530/jrf.0.0990283

Moore, N. P., Kelly, P. F., Cahill, J. P., & Hayden, T. J. (1995). Mating strategies and mating success of

fallow (Dama dama) bucks in a non-lekking population. *Behavioral Ecology and Sociobiology*,

*36*(2), 91–100. https://doi.org/10.1007/BF00170713

Möstl, E., Maggs, J. L., Schrötter, G., Besenfelder, U., & Palme, R. (2002). Measurement of cortisol

metabolites in faeces of ruminants. *Vetinary Research Communications*, *26*, 127–139.

https://doi.org/10.1023/A:1014095618125

Murphy, V. E., Gibson, P. G., Giles, W. B., Zakar, T., Smith, R., Bisits, A. M., Kessell, C. G., & Clifton, V.

L. (2003). Maternal asthma is associated with reduced female fetal growth. *American Journal*

*of Respiratory and Critical Care Medicine*, *168*(11), 1317–1323.

https://doi.org/10.1164/rccm.200303-374OC

O'Connell, B. A., Moritz, K. M., Walker, D. W., & Dickinson, H. (2013). Sexually dimorphic placental

development throughout gestation in the spiny mouse (Acomys cahirinus). *Placenta*, *34*(2),

199–126. https://doi.org/10.1016/j.placenta.2012.11.009

Palme, R. (2019). Non-invasive measurement of glucocorticoids: Advances and problems. *Physiology*

*& Behavior*, *199*, 229–243. https://doi.org/10.1016/j.physbeh.2018.11.021

Palme, R., Touma, C., Arias, N., Dominchin, M. F., & Lepschy, M. (2013). Steroid extraction: Get the

best out of faecal samples. *Wiener Tierärztliche Monatsschrift*, *100*(9–10), 238–248.

R Core Team. (2020). *R: A language and environment for statistical computing*. https://www.R-

project.org/

Rosenfeld, C. S. (2015). Sex-specific placental responses in fetal development. *Endocrinology*,

*156*(10), 3422–3434. https://doi.org/10.1210/en.2015-1227

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). *How do glucocorticoids influence stress

responses? Integrating permissive, suppressive, stimulatory, and preparative Actions*. *21*(1),

55–89. https://doi.org/10.1210/edrv.21.1.0389

Seckl, J. R., Cleasby, M., & Nyirenda, M. J. (2000). Glucocorticoids, 11β-hydroxysteroid

dehydrogenase, and fetal programming. *Kidney International*, *57*(4), 1412–1417.

https://doi.org/10.1046/j.1523-1755.2000.00984.x

Seckl, J. R., & Holmes, M. C. (2007). Mechanisms of disease: Glucocorticoids, their placental

metabolism and fetal'programming'of adult pathophysiology. *Nature Clinical Practice

Endocrinology & Metabolism*, *3*, 479–488. https://doi.org/10.1038/ncpendmet0515

Seckl, J. R., & Meaney, M. J. (2004). Glucocorticoid programming. *Annals of the New York Academy

of Sciences*, *1032*(44), 63–84. https://doi.org/10.1196/annals.1314.006

Sheriff, M. J., & Love, O. P. (2013). Determining the adaptive potential of maternal stress. *Ecology

Letters*, *16*(2), 271–280. https://doi.org/10.1111/ele.12042

Tomlinson, J. W., & Stewart, P. M. (2001). Cortisol metabolism and the role of 11β-hydroxysteroid

dehydrogenase. *Best Practice & Research Endocrinology & Metabolism*, *15*(1), 61–78.

https://doi.org/10.1053/beem.2000.0119

Weinstock, M. (2008). The long-term behavioural consequences of prenatal stress. *Neuroscience and

Biobehavioral Reviews*, *32*(6), 1073–1086. https://doi.org/10.1016/j.neubiorev.2008.03.002

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer.

Wieczorek, A., Perani, C. V., Nixon, M., Constancia, M., Sandovici, I., Zazara, D. E., Leone, G., Zhang,

M. Z., Arck, P. C., & Solano, M. E. (2019). Sex-specific regulation of stress-induced fetal

glucocorticoid surge by the mouse placenta. *American Journal of Physiology - Endocrinology*

*and Metabolism*, *317*(1), E109–E120. https://doi.org/10.1152/ajpendo.00551.2018

400