Sex-specific relationship between maternal and neonate cortisol in a free-ranging large mammal

Bawan Amin[1, 5*], Ruth Fishman[2], Matthew Quinn[1, 3], Devorah Matas[2], Rupert Palme[4], Lee Koren[2], Simone Ciuti[1]

1) Laboratory of Wildlife Ecology and Behaviour, School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

2) Faculty of Life Sciences, Bar-Ilan University, Ramat-Gan, Israel

3) Department of Biosciences, Durham University, Durham, United Kingdom

4) Unit of Physiology, Pathophysiology and Experimental Endocrinology, Department of Biomedical Sciences, University of Veterinary Medicine, Vienna, Austria

5) Faculty of Social and Behavioural Sciences, Utrecht University, Utrecht, Netherlands

* Correspondence: b.k.r.amin@uu.nl

Acknowledgements

We thank the Office of Public Works (OPW), Ireland, for funding (grant no. R18625) and support. We thank the SBES in University College Dublin (UCD), and the British Deer Society (R21845) for funding this project. We also would like to thank the Higher Education Authority of Ireland for providing a Covid-19 related Research Cost Extensions to BA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We declare that none of the authors have a conflict of interest.

Data availability statement

All data used in this manuscript will be made available on OSF or Zenodo at acceptance, and is

already available for the review process as supplementary material. We commit to providing the DOI

of the dataset in this section in the final version of the manuscript.

Author contributions

BA: Conceptualization, data curation, formal analysis, investigation, visualization, writing -original

draft preparation; RF: Conceptualization, writing – original draft preparation, writing – review &

editing; MQ: Data curation, investigation, writing – review & editing; DM: Investigation,

methodology, writing – review & editing; RP: Investigation, methodology, writing – review & editing;

LK: Investigation, methodology, writing – review & editing; SC: Conceptualization, funding

acquisition, resources, project administration, supervision, writing – review & editing

Abstract

Maternal phenotypes can have long-term effects on the offspring phenotype starting at gestation.

Proximate mechanisms include foetal hormone levels affected by maternal levels. Recent studies

have highlighted that these maternal effects may be different between the sexes. How maternal

glucocorticoid (GC) levels relate to foetal levels is, however, still not completely understood. Here

we related, for the first time in a free-ranging large mammal, the fallow deer (*Dama dama*),

maternal GC levels with foetal *in utero* GC levels by using novel state-of-the-art methods. We found

that foetal GC levels were positively associated with maternal GC levels, but only in females. These

findings highlight sex differences, which may have evolved to optimize male growth at the cost of

survival.

Keywords

Glucocorticoids, free-ranging, fallow deer, hair-testing, faecal metabolites

Introduction

Parental phenotypes can be drivers of offspring variation, and in many mammalian species, these parental effects are often labelled as maternal effects since it is the mother that mostly takes care of the offspring. Maternal effects are seen as an adaptive way in which mothers can either fine tune their offspring for the current environmental conditions [1], or maximise their own reproductive output [2]. These effects on the offspring are present from the earliest stages of development and can have long lasting influence on offspring phenotype [3,4].

Acute stress typically leads to an increase in circulating glucocorticoid (GC) levels due to activation of the hypothalamic-pituitary-adrenal (HPA) axis [4,5]. Maternal GC levels during gestation can affect offspring phenotype in many ways, including affecting offspring birth weight and HPA-axis reactivity [3,6,7] and these effects can linger until offspring are well into adulthood (e.g. 8), which is also referred to as foetal programming [6]. There is increasing evidence, however, that there are sex differences in foetal programming [8–10].

Most studies investigating hormonal maternal effects investigate offspring GC levels after birth and most evidence comes from either clinical studies on humans [11], rodents [4] or birds [12,13], with a paucity of data on free-ranging mammals. This is usually due to the challenges of measuring maternal and foetal hormone levels during gestation in non-captive mammals. Recent developments, however, have enabled measuring foetal hormonal levels post-parturition by quantifying steroids in neonate hair [14–16]. Hair steroid levels represent long-term levels, accumulated over the last weeks to months [17], and therefore reflect *in utero* integrated levels when quantified in neonates [16]. Whether these levels are associated with circulating maternal GC levels, however, is currently not well understood.

Here we explored, for the first time in free-ranging large mammals, the relationship between circulating maternal GC levels and offspring *in utero* accumulated GC levels. For that purpose, we collected faeces of mothers during late gestation, from which we quantified GC metabolites [18], and related that to their offspring hair GC levels [16], which were collected during

the first days post-parturition at neonate captures. We made sure to sample females within their

last five weeks of gestation, since that is when the foetus is fully covered in fur [19]. To allow for sex

differences in the association between maternal and foetal GC levels, we ran an additional analysis

to explore differences between male and female offspring.

Methods

Study population

We conducted this study in Phoenix Park, a 7.07 km^2 urban park located in Dublin, Ireland. There is a

resident population of free-ranging fallow deer that has been introduced in the 17th century, with a

population size of about 600 individuals in late summer, after fawn births [20]. Most births occur

between early and late June of each year. Fallow deer are a hider species and fawns remain hidden,

usually in tall grass or understory vegetation, away from the main doe herd during the first weeks of

life after which they are brought into the doe herd by their mothers [19,21]. Fawns are occasionally

predated upon by red foxes (*Vulpes vulpes*), the only natural predator in the park, and domestic dogs

who are brought into the park by public visitors.

Faeces collection and analysis of maternal faecal cortisol metabolites (FCMs)

Faecal samples were collected between 8 AM and 3 PM, from May 19th until May 29th 2020, when

the does were in late gestation. Groups of deer were observed from a distance of 50 meters using a

spotting scope to identify individuals (>80% of the population is identifiable via unique colour coded

ear-tags). Fresh faecal samples were collected within a minute of defecation and immediately stored

in zip locked bags. These were kept in a cooler bag until they could be stored in a freezer at -20˚C,

which was always within a few hours. Samples (see *Sample sizes*-section below) were kept frozen on

dry ice during transportation to the University of Veterinary Medicine (Vienna, Austria), where

cortisol metabolites in the samples were quantified [18]. We added 5 mL of methanol (80%) to

weighed aliquots (0.5 g) of homogenized faecal samples, after which the samples were shaken and

centrifuged as previously described [22]. FCMs were analysed with an 11-oxoaetiocholanolone

enzyme immunoassay (for details see Möstl et al., 2002 [23]), which has already been validated for

fallow deer [24].

Neonate hair collection and hair GC level quantification

Neonate hair samples were collected during the fawn captures in June 2020. Fawns were routinely

captured with fishing nets, as part of the long-term management of the population. During handling,

we collected physiological and behavioural data (see Amin et al., 2021 [16] for full details on the

capture protocol), as well as hair samples (>100 mg) from the belly of the fawn using an electric

trimmer (Wahl model 9639; Wahl Clipper Corporation). We extracted cortisol levels from fawn hair

using a standardized protocol for hair-testing (e.g. 15,25), by using commercial enzyme

immunoassay (EIA; Salimetrics; Ann Arbor, MI, USA, item no. 1-3002) kits. Full details on these

extractions and validations are described in Amin et al., 2021 [16] and in the supplementary material

(S1).

Mother-fawn pairing

Mother-fawn pairs were based on field observations with data collection starting in July, when

young fawns were making their first entrances into the female herds. For details regarding the

pairing, see Griffin et al., 2022 [20] and Supplementary S1.

Sample sizes

We collected and analysed a total of 164 faecal samples of 99 different pregnant does. For

individuals with multiple values, we took the mean of these values to create a dataset that had one

estimate for each individual. Out of these, we were able to pair 41 does with their fawns through

the mother-fawn observations during summer. We removed one extreme outlier from the male

neonate cortisol dataset (value: 29.5 pg/mg hair; range of other values: 7.7-18.3 pg/mg hair).

Running our analysis with and without this outlier revealed that removing this outlier did not affect

the conclusions of our analysis. This led to our final sample size of 40 fawns (18 females and 22

males) paired with their mothers.

Statistical analysis

All analyses were performed in RStudio (Version 1.3.1093) using R version 4.0.2 [26].

We first checked whether maternal or neonatal GC levels differed between the offspring sexes by

performing a t-test, considering unequal variances.

To investigate the relationship between neonate hair GC and maternal FCMs, we ran a linear

model for each fawn sex separately. In both models, we had the neonate hair GC as the response

variable and the maternal FCMs as the explanatory variable. Maternal FCM levels were log-

transformed to improve model fit, since these suffered from a slight positive skew. Model

assumptions were checked with the *DHARMa*-package (Version 0.4.3[27]) and were successfully

met. Statistical inferences were made based on the estimate and the associated 95% Confidence

Interval from the models. During the preliminary analysis, we included the number of days between

the collection of mothers' faeces and the day of fawn birth in our models. This was to account for

potential variation in FCM levels as a function of gestational day of collection (i.e. mid May vs late

May when closer to parturition). This additional predictor had no clear effect and made our models

worse (higher AIC); we therefore decided not to include it. We plotted our results using the *ggplot2*-

package [28].

Finally, we decided to run an additional model *post-hoc*, including both sexes in the same

model, to further investigate whether the slopes of our models were different between male and

female offspring. We ran a linear model with neonate GC as the response variable. As explanatory

variables, we included maternal FCMs (log transformed), fawn sex and the interaction between both

these explanatory variables.

Results

Considerable variation was found in both the individual maternal FCM levels (range: 91-608 ng/g

faeces) as well as the neonate hair GC levels (range: 6.50-18.30 pg/mg hair). We found no clear sex

differences, neither for mean maternal FCM levels (t-test; t = 0.56, 95% CI [-51.99, 91.50], p = 0.58)

nor mean neonate hair GC levels (t-test; t = 0.26, 95% CI [-1.74, 2.25], p = 0.80).

165   We found that female hair cortisol was positively associated with maternal FCMs (LM: β =

5.88, 95% CI [0.20, 11.57], p = 0.04, N = 18, $R^2_{adjusted}$ = 0.18; Fig. 1A). For males, however, there was

no clear association between both measures (LM: β = 0.59, 95% CI [-1.32, 2.49], p = 0.53, N = 22,

$R^2_{adjusted}$ = -0.03; Fig. 1B). Our post-hoc model confirmed that there was indeed an interaction

between fawn sex and maternal levels, where the slope between maternal and foetal levels was

lower in male offspring compared to female offspring (LM: β = -5.30, 95% CI [-10.39, -0.20], p = 0.04,

N = 40, $R^2_{adjusted}$ = 0.10).

Figure 1: The relationship between maternal faecal cortisol metabolites (FCMs) and neonate hair

175   GCs, for female (A) and male (B) offspring. Trendlines and their 95% confidence intervals are fitted in

176   each plot. Solid trendline indicates a statistically meaningful relationship.

Discussion

In this study we found that maternal FCM levels were positively related to foetal GC levels, but only in females. In male offspring, we found no clear relationship between maternal and foetal GC levels. These findings suggest that maternal GC levels may affect male offspring less then female offspring. We discuss here how, from an evolutionary point of view, sex-specific mechanisms may have evolved to optimize life-history trade-offs.

Like many other species, fallow deer have a skewed mating distribution [29,30]. While females tend to produce one fawn per year [19], male reproduction skew is steep. Only a small proportion of the males reproduces, with the majority having no or very little offspring [29–31]. Therefore, there may be a strong selection for males to prioritize growth over survival, since low quality sons will end up costing resources, with no fitness benefits in the long run. A recent study has shown that male birthweights in our populations are indeed higher than female birthweights [32].

GC levels play a key role during gestation and have major effects on offspring phenotype early in life. In addition to being crucial for organ development during the late stages of pregnancy [33,34], GCs play a role in fighting inflammation [35] and thereby help to keep the foetus vital. At the same time, high GC levels may restrict growth [6,36,37]. As males tend to prioritize growth [36], it may be that there would be sex differences in the mechanisms that evolved. For example, maternal GC contribution may be limited in male foetuses to prevent growth retardation, at the cost of prenatal survival. This may contribute in some degree to why males generally face higher prenatal mortality rates [38–40].

The mammalian placenta likely plays a crucial role in mediating these sex differences. Previous studies, in rodents and humans, have shown sex differences in the structure and activity of the placenta [36,41–44]. There is also evidence that maternal GCs may transport more easily through female placentas than males' [43]. This may be, for instance, through different levels of expression and activity of enzymes or transporters that mitigate different maternal GC level spill over between males and females.

One of the key enzymes, 11β-hydroxysteroid dehydrogenase type 2 (11β-HSD2), inactivates

cortisol to cortisone [37,45]. Placental 11β-HSD has been frequently shown to be positively related

to birthweight [3,36,37,46], thus likely playing a role in preventing growth restriction. Previous

studies have shown that male foetuses have higher expression or activity of 11β-HSD2 in response to

maternal GC surges [41,43]. Similarly in ovine, male foetuses increased 11β-HSD2 in response to

maternal dexamethasone administration, whereas female foetuses did not [47]. This may explain

why a different ovine study also found that, following an antenatal GC treatment, growth restriction

was lower in males than in females [48]. In addition, male mice were shown to also have increased

ATP-binding cassette transporters, which mediate GC efflux toward maternal circulation [43],

indicating that there are different mechanisms through which males may be able to limit maternal

GC levels [49]. This altogether may explain why females may have higher foetal survival [36],

whereas males may be larger and faster growing.

We acknowledge that there are some shortcomings to this study. We report patterns taken

over only one year, whereas ecological patterns may vary between years. Furthermore, we have

modest sample sizes (due to the difficulty of the design), which restrict us in doing more elaborate

analyses. Nevertheless, as one of the first studies quantifying the relationship between maternal and

foetal GC levels non-invasively, we do provide novel insights into how these fundamental

relationships may function in a free-ranging large mammal population. It is important to note that

the nature of the study described here is exploratory, meaning that it generates hypotheses that can

be tested in the future.

To conclude, we explored in this study the relationship between maternal and foetal GC

levels in a free-ranging population of fallow deer. Our findings suggest that there are sex differences

in the underlying evolutionary processes, which may have optimized growth at the cost of survival

for male offspring, whereas the opposite is true for female offspring. Most of the existing literature

uses the rodent or human model, with very little existing studies on other taxa. Although some

features of the placenta may be conserved, there are also evolutionary differences between species

[44,50]. Non-model species may provide different results [15], through which we can gain a better

fundamental understanding of evolutionary mechanisms. By studying free-ranging fallow deer, this

study therefore also provides novel insights into possible sex-specific maternal effects in a non-

model species, and provides a method that can be viable for other populations as well, enabling the

study of patterns rarely studied before in the wild.

References

1. Sheriff MJ, Love OP. 2013 Determining the adaptive potential of maternal stress. *Ecol. Lett.* **16**,
271–280. (doi:10.1111/ele.12042)

2. Marshall DJ, Uller T. 2007 When is a maternal effect adaptive? *Oikos* **116**, 1957–1963.
(doi:10.1111/j.2007.0030-1299.16203.x)

3. Seckl JR, Meaney MJ. 2004 Glucocorticoid programming. *Ann. N. Y. Acad. Sci.* **1032**, 63–84.
(doi:10.1196/annals.1314.006)

4. Weinstock M. 2008 The long-term behavioural consequences of prenatal stress. *Neurosci.*
Biobehav. Rev. **32**, 1073–1086. (doi:10.1016/j.neubiorev.2008.03.002)

5. Sapolsky RM, Romero LM, Munck AU. 2000 How do glucocorticoids influence stress responses?
Integrating permissive, suppressive, stimulatory, and preparative Actions. **21**, 55–89.
(doi:10.1210/edrv.21.1.0389)

6. Seckl JR, Holmes MC. 2007 Mechanisms of disease: glucocorticoids, their placental metabolism
and fetal'programming'of adult pathophysiology. *Nat. Clin. Pract. Endocrinol. Metab.* **3**, 479–
488. (doi:10.1038/ncpendmet0515)

7. Dantzer B, Newman AEM, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam AG. 2013
Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal.
Science **340**, 1215–1217. (doi:10.1126/science.1235765)

8. Liu L, Li A, Matthews SG. 2001 Maternal glucocorticoid treatment programs HPA regulation in
adult offspring: sex-specific effects. *Am. J. Physiol.-Endocrinol. Metab.* **280**, E729–E739.
(doi:10.1152/ajpendo.2001.280.5.E729)

9. Braithwaite EC, Hill J, Pickles A, Glover V, O'Donnell K, Sharp H. 2018 Associations between
maternal prenatal cortisol and fetal growth are specific to infant sex: findings from the Wirral
Child Health and Development Study. *J. Dev. Orig. Health Dis.* **9**, 425–431.
(doi:10.1017/S2040174418000181)

10. Fishman R, Kralj-Fišer S, Marglit S, Koren L, Vortman Y. 2022 Quantitative genetic study suggests
sex-specific genetic architecture for fetal testosterone in a wild mammal. *BioRxiv Prepr.*
(doi:10.1101/2022.02.18.480975)

11. Gitau R, Cameron A, Fisk NM, Glover V. 1998 Fetal exposure to maternal cortisol. *Lancet* **352**,
707–708. (doi:10.1016/S0140-6736(05)60824-0)

12. Jenkins BR, Vitousek MN, Hubbard JK, Safran RJ. 2014 An experimental analysis of the
heritability of variation in glucocorticoid concentrations in a wild avian population. *Proc. R. Soc.*
B Biol. Sci. **281**, 20141302. (doi:10.1098/rspb.2014.1302)

13. Groothuis TGG, Hsu BY, Kumar N, Tschirren B. 2019 Revisiting mechanisms and functions of
prenatal hormone-mediated maternal effects using avian species as a model. *Philos. Trans. R.*
Soc. B Biol. Sci. **374**, 20180115. (doi:10.1098/rstb.2018.0115)

14. Kapoor A, Lubach GR, Ziegler TE, Coe CL. 2016 Hormone levels in neonatal hair reflect prior
maternal stress exposure during pregnancy. *Psychoneuroendocrinology* **66**, 111–117.
(doi:10.1016/j.psyneuen.2016.01.010)

15. Fishman R, Vortman Y, Shanas U, Koren L. 2019 Non-model species deliver a non-model result:
Nutria female fetuses neighboring males in utero have lower testosterone. *Horm. Behav.* **111**,
105–109. (doi:10.1016/j.yhbeh.2019.02.011)

16. Amin B, Jennings DJ, Smith AF, Quinn M, Chari S, Haigh A, Matas D, Koren L, Ciuti S. 2021 In
utero accumulated steroids predict neonate anti-predator response in a wild mammal. *Funct.*
Ecol. **35**, 1255–1267. (doi:10.1111/1365-2435.13790)

17. Gormally BMG, Romero LM. 2020 What are you actually measuring? A review of techniques that
integrate the stress response on distinct time-scales. *Funct. Ecol.* **34**, 2030–2044.
(doi:10.1111/1365-2435.13648)

18. Palme R. 2019 Non-invasive measurement of glucocorticoids: Advances and problems. *Physiol.*
Behav. **199**, 229–243. (doi:10.1016/j.physbeh.2018.11.021)

19. Chapman D, Chapman N. 1997 *Fallow deer: Their history, distribution and biology*. 2nd edn.
Coch-y-bonddu Books.

20. Griffin LL, Haigh A, Amin B, Faull J, Norman A, Ciuti S. 2022 Artificial selection in human-wildlife
feeding interactions. *J. Anim. Ecol.* **91**, 1892–1905. (doi:10.1111/1365-2656.13771)

21. Ciuti S, Bongi P, Vassale S, Apollonio M. 2006 Influence of fawning on the spatial behaviour and
habitat selection of female fallow deer (Dama dama) during late pregnancy and early lactation.
J. Zool. **268**, 97–107. (doi:10.1111/j.1469-7998.2005.00003.x)

22. Palme R, Touma C, Arias N, Dominchin MF, Lepschy M. 2013 Steroid extraction: Get the best out
of faecal samples. *Wien. Tierärztl. Monatsschrift* **100**, 238–248.

23. Möstl E, Maggs JL, Schrötter G, Besenfelder U, Palme R. 2002 Measurement of cortisol
metabolites in faeces of ruminants. *Vetinary Res. Commun.* **26**, 127–139.
(doi:10.1023/A:1014095618125)

24. Konjević D, Janicki Z, Slavica A, Severin K, Krapinec K, Božić F, Palme R. 2011 Non-invasive
monitoring of adrenocortical activity in free-ranging fallow deer (Dama dama L.). *Eur. J. Wildl.*
Res. **57**, 77–81. (doi:10.1007/s10344-010-0401-1)

25. Koren L, Geffen E. 2009 Androgens and social status in female rock hyraxes. *Anim. Behav.* **77**,
233–238. (doi:10.1016/j.anbehav.2008.09.031)

26. R Core Team. 2020 *R: A language and environment for statistical computing*. Vienna, Austria.
See https://www.R-project.org/.

27. Hartig F. 2021 *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed) Regression*
Models. See https://CRAN.R-project.org/package=DHARMa.

28. Wickham H. 2016 *ggplot2: Elegant graphics for data analysis*. New York: Springer.

29. Moore NP, Kelly PF, Cahill JP, Hayden TJ. 1995 Mating strategies and mating success of fallow
(Dama dama) bucks in a non-lekking population. *Behav. Ecol. Sociobiol.* **36**, 91–100.
(doi:10.1007/BF00170713)

30. McElligott AG, Gammell MP, Harty HC, Paini DR, Murphy DT, Walsh JT, Hayden TJ. 2001 Sexual
size dimorphism in fallow deer (Dama dama): Do larger, heavier males gain greater mating
success? *Behav. Ecol. Sociobiol.* **49**, 266–272. (doi:10.1007/s002650000293)

31. Ciuti S, De Cena F, Bongi P, Apollonio M. 2011 Benefits of a risky life for fallow deer bucks (Dama
dama) aspiring to patrol a lek territory. *Behaviour* **148**, 435–460.
(doi:10.1163/000579511X563981)

32. Griffin LL, Haigh A, Amin B, Faull J, Corcoran F, Baker-Horne C, Ciuti S. 2023 Does artificial
feeding impact neonate growth rates in a large free-ranging mammal? *R. Soc. Open Sci.* **10**,
221386. (doi:10.1098/rsos.221386)

33. Kitterman JA, Liggins GC, Campos GA, Clements JA, Forster GC, Lee CH, Creasy RK. 1981
Prepartum maturation in fetal sheep : Relation of the lung to cortisol. *J. Appl. Physiol.* **51**, 384–
390. (doi:10.1152/jappl.1981.51.2.384)

34. Liggins GC. 1994 The role of cortisol in preparing the fetus for birth. *Reprod. Fertil. Dev.* **6**, 141–
150. (doi:10.1071/RD9940141)

35. Auphan N, Didonato JA, Rosette C, Helmberg A, Karin M. 1995 Immunosuppression by
glucocorticoids: Inhibition of NF-κB activity through induction of IκB synthesis. **270**, 286–290.
(doi:10.1126/science.270.5234.286)

36. Meakin AS, Cuffe JSM, Darby JRT, Morrison JL, Clifton VL. 2021 Let's talk about placental sex,
baby: Understanding mechanisms that drive female-and male-specific fetal growth and
developmental outcomes. *Int. J. Mol. Sci.* **22**. (doi:10.3390/ijms22126386)

37. Edwards CRW, Benediktsson R, Lindsay RS, Seckl JR. 1996 11β-Hydroxysteroid dehydrogenases:
Key enzymes in determining tissue-specific glucocorticoid effects. *Steroids* **61**, 263–269.
(doi:10.1016/0039-128X(96)00033-5)

38. Desportes G, Andersen LW, Bloch D. 1994 Variation in foetal and postnatal sex ratios in long-
finned pilot whales. *Ophelia* **39**, 183–196. (doi:10.1080/00785326.1994.10429543)

39. Eriksson JG, Kajantie E, Osmond C, Thornburg K, Barker DJP. 2010 Boys live dangerously in the
womb. *Am. J. Hum. Biol.* **22**, 330–335. (doi:10.1002/ajhb.20995)

40. Kruuk LE, Clutton-Brock TH, Albon SD, Pemberton JM, Guinness FE. 1999 Population density
affects sex ratio variation in red deer. *Nature* **399**, 459–461. (doi:10.1038/20917)

41. Murphy VE, Gibson PG, Giles WB, Zakar T, Smith R, Bisits AM, Kessell CG, Clifton VL. 2003
Maternal asthma is associated with reduced female fetal growth. *Am. J. Respir. Crit. Care Med.*
168, 1317–1323. (doi:10.1164/rccm.200303-374OC)

42. O'Connell BA, Moritz KM, Walker DW, Dickinson H. 2013 Sexually dimorphic placental
development throughout gestation in the spiny mouse (Acomys cahirinus). *Placenta* **34**, 199–
126. (doi:10.1016/j.placenta.2012.11.009)

43. Wieczorek A *et al.* 2019 Sex-specific regulation of stress-induced fetal glucocorticoid surge by
the mouse placenta. *Am. J. Physiol. - Endocrinol. Metab.* **317**, E109–E120.
(doi:10.1152/ajpendo.00551.2018)

44. Rosenfeld CS. 2015 Sex-specific placental responses in fetal development. *Endocrinology* **156**,
3422–3434. (doi:10.1210/en.2015-1227)

45. Tomlinson JW, Stewart PM. 2001 Cortisol metabolism and the role of 11β-hydroxysteroid
dehydrogenase. *Best Pract. Res. Endocrinol. Metab.* **15**, 61–78. (doi:10.1053/beem.2000.0119)

46. Seckl JR, Cleasby M, Nyirenda MJ. 2000 Glucocorticoids, 11β-hydroxysteroid dehydrogenase,
and fetal programming. *Kidney Int.* **57**, 1412–1417. (doi:10.1046/j.1523-1755.2000.00984.x)

47. Braun T *et al.* 2009 Effects of maternal dexamethasone treatment in early pregnancy on
pituitary-adrenal axis in fetal sheep. *Endocrinology* **150**, 5466–5477. (doi:10.1210/en.2009-0086)

48. Miller SL, Sutherland AE, Supramaniam VG, Walker DW, Jenkin G, Wallace EM. 2012 Antenatal
glucocorticoids reduce growth in appropriately grown and growth-restricted ovine fetuses in a
sex-specific manner. *Reprod. Fertil. Dev.* **24**, 753. (doi:10.1071/RD11143)

49. Montano MM, Wang M, vom Saal FS. 1993 Sex differences in plasma corticosterone in mouse
fetuses are mediated by differential placental transport from the mother and eliminated by
maternal adrenalectomy or stress. *J. Reprod. Fertil.* **99**, 283–290. (doi:10.1530/jrf.0.0990283)

50. Fowden A. 2003 The insulin-like growth factors and feto-placental growth. *Placenta* **24**, 803–
812. (doi:10.1016/S0143-4004(03)00080-8)

363