Predator learning can resolve the paradox of local warning signal diversity

Chi-Yun Kuo[1,2]

1. Department of Biomedical Science and Environmental Biology, Kaohsiung Medical

University, 100 Shih-Chuan 1[st] Rd, Kaohsiung City 80707, Taiwan

2. Corresponding author: +886-7-312-1101 ext 2693, kuochiyun@gmail.com

Running title: Explaining local warning signal diversity

Keywords: aposematism, associative learning, classical conditioning, Müllerian mimicry,

Pavlovian learning, positive frequency-dependent selection, predator-prey interaction

Type of article: letter

150 words in the abstract, 6199 words in the main text, 112 references, 3 figures and 1 table

Statement of authorship: CYK conceived the study, performed computer simulations, literature

search and meta-analysis, and wrote the manuscript

Data accessibility statement: The dataset and R codes for the simulation and meta-analysis can

be accessed at https://github.com/thekuolab/warning-signal-diversity

Novelty statement: This paper presents the first widely applicable and testable explanation for

the stable coexistence of distinct warning signals within communities, which has long stood as a

paradox as it defies the expectation of positive frequency-dependent selection acting on

sympatric warning signals. Combining computer simulations and meta-analysis, this paper shows that local warning signal diversity is not paradoxical if prey elicit sufficient predator avoidance and, more importantly, if predators can retain enough learned avoidance within the focal prey's generation time. The meta-analysis also offers new insights into how causes of prey unprofitability, manner of predator-prey encounter, and type of warning signals can influence predator avoidance and the likelihood of local warning signal diversity, representing novel hypotheses to be further tested.

Abstract

Stable coexistence of multiple warning signals within communities has long been a paradox as selection is expected to preserve only the most common signal. This study presents the first generalizable and testable resolution to this paradox by showing that prey displaying rare warning signals can persist if they elicit sufficient predator avoidance and, more importantly, if predators can retain enough learned avoidance within the focal prey's generation time. I validated this hypothesis with empirical learning data that can be readily acquired from laboratory and field experiments. I also performed a meta-analysis examining the variation in predator avoidance. I found that predator avoidance and the likelihood of local warning signal diversity might be higher in communities with the following features – unprofitable prey are toxic instead of merely unpalatable, predators search widely for prey and/or prey aggregate, and when predators could differentiate between unprofitable and profitable prey using only color or pattern.

 Introduction

Unprofitable prey often evolve conspicuous colorations to advertise their unsuitability (i.e.,

aposematism (Poulton 1890). These visual advertisements act as honest warning signals that

predators can use to avoid these prey through associative learning. The convergence of warning

signals among sympatric prey has inspired evolutionary ecologists to formulate hypotheses

regarding the evolution and maintenance of mimicry (Müller 1879; Blakiston & Alexander 1883;

Mallet 2001). Following Müller's pioneer theory, each predator has to attack a fixed number of

prey to fully acquire learned avoidance. This results in a "safety-in-numbers" scenario in which

bearers of rare warning signals are at a disadvantage due to there being fewer individuals to

dilute the mortality cost of predator education, resulting in positive frequency-dependent

selection (hereafter FDS) that will only preserve the most abundant warning signal in a

community (Müller 1879). Müller's theory aptly explains the often-observed signal convergence

among unprofitable prey (i.e., Müllerian mimicry) and predicts signal homogeneity at local

scales. Yet, despite several well-established examples of Müllerian signal convergence, stable

diversity in warning signals within communities does not appear to be a rare occurrence (Ruxton

et al. 2018; Briolat *et al.* 2019). This stark contrast between theoretical expectations and reality

has long stood as a paradox. Several ecological mechanisms have been proposed as potential

resolutions to this paradox (reviewed in (Ruxton *et al.* 2018), but none is both broadly applicable

and testable. It was hypothesized that species displaying distinct warning signals might segregate

along certain microhabitat axes, and predators learn to avoid different signals in each

microhabitat type (Gompert *et al.* 2011; Aubier *et al.* 2017). This possibility can be tested

empirically but so far seems only able to explain local warning signal diversity in neotropical

ithomiine butterflies, in which distinctively colored species fly at different height strata as a

result of host plant distribution (Beccaloni 1997; Willmott *et al.* 2017). Notably, microhabitat

segregation cannot explain the coexistence of distinct warning signals in two prominent systems

of Müllerian mimicry: neotropical *Heliconius* butterflies and *Oophaga* poison frogs (Joron &

Mallet 1998; Dugas *et al.* 2015).

Here I present a novel explanation for local warning signal diversity that directly relates to

predator learning (the unforgetful predator hypothesis), and this explanation begins by re-

examining the notion that number-dependent predator education will lead to positive FDS.

Equating the former to the latter is based on an implicit assumption - the mortality cost of

predator education is imposed on every prey generation, such that prey displaying rare signals

repeatedly suffer from a disproportionally higher mortality rate (Mallet 2010). In other words,

positive FDS results when predators do not retain enough learned avoidance relative to the focal

prey's generation time, and when prey reproduction between generations is not enough to offset

learning-based predation. The conditions for this assumption to hold have never been challenged

and examined in detail from the perspective of predator learning. There have been studies that

reported positive FDS based on the finding that unprofitable prey with novel colorations suffered

higher predation rates in the field (e.g., (Chouteau *et al.* 2016). Yet, given the relatively short

duration of the experiments (a few days) such results could very well be capturing the initial

stage of predator learning, in which prey displaying novel warning signals were incurring the

mortality cost of predator education for the very first time. Without information on forgetting

rate, it cannot be ascertained whether such observations equate the presence of positive FDS. On

the theoretical front, (Mallet & Joron 1999) and (Franks & Noble 2004) used Müller's safety-in-

numbers concept and explored the occurrence of local warning signal diversity. While both

studies each touched on aspects of the unforgetful predator hypothesis, neither fully examined

the relationship between number-dependent predator education, positive FDS, and the

persistence of rare warning signals.

I simulated predator-prey interactions and predator learning in a Pavlovian framework (Turner

1984; Speed 1993, 2000, 2001; Turner & Speed 1996; Speed & Turner 2008; Turner *et al.* 2008),

section I of SI). A main criticism of the Pavlovian learning is that they seem to contradict

observations from laboratory and field studies, in which the number of unprofitable prey attacked

in learning trials increase either proportionally or nearly so with prey abundance (Greenwood *et*

al. 1989; Lindström *et al.* 2001b; Beatty *et al.* 2004; Rowland *et al.* 2010b; Chouteau *et al.* 2016).

However, Pavlovian learning can lead to such observations when the final attack probability is

not exactly zero, which is likely to be the case in these studies where prey were merely

unpalatable. The observations that a proportionately smaller increase in the number of

unprofitable prey with abundance (e.g., (Lindström *et al.* 2001b) can also be explained by a non-

zero final attack probability coupled with predator satiation (i.e., Holling's type II functional

response). Pavlovian learning also has a significant advantage over the other alternative (i.e., the

Bayesian learning framework, (Sherratt 2011; Aubier & Sherratt 2015) in that the two key

parameters - final attack probability and forgetting rate - can be directly measured with learning

experiments (see section II of SI for more detailed explanation).

With the simulations, I examined the conditions under which stable persistence of rare warning

signals can be achieved as a balance between learning-related predation and reproduction. The

predation rate towards prey displaying a particular warning signal is the product of encounter

rate and attack probability, with the former determined by relative abundance and the latter by

predator learning. The role of final attack probability is straightforward; it determines the lower

bound of predation rate and the ever-present baseline mortality after learning is complete. The

role of forgetting rate is twofold. First, forgetful predators may repeatedly impose mortality cost

of predator education on each prey generation, leading to positive FDS. Moreover, when

predators are quick to forget learned avoidance, their attack probabilities towards prey will often

be higher than the final asymptotic value, making baseline predation mortality higher than it is

expected to be. These disadvantages are particularly severe for rare warning signals, as their low

encounter rates lead to more opportunities for predator forgetting. The unforgetful predator

hypothesis posits that when predators can retain learned avoidance for sufficiently long, the

mortality cost for predator education can become trivial compared to the ever-present baseline

predation. Whether and when the unforgetful predator hypothesis could explain local warning

signal diversity hinges on final attack probability and forgetting rate of predators, as well as the

reproductive capability of prey. In addition, I considered the effect of two important phenomena

of predator learning - cognitive generalization and social learning. Cognitive generalization can

occur when a novel warning signal resembles a learned signal. Social learning describes the

phenomenon that avoidance learning can occur from observing the negative responses of other

predators towards an aposematic prey. These two phenomena have been implemented in

Pavlovian learning (Franks & Noble 2004; Thorogood *et al.* 2018), yet their facilitative effects

on local warning signal diversity have never been explicitly tested.

In addition to validating the unforgetful predator hypothesis, another goal of this study is to

examine the variation in predators' final attack probabilities and the implications on the

likelihood of local warning signal diversity in natural communities. For this, I performed a meta-

analysis of the avoidance learning literature, focusing on the roles of the following predictors:

the reason for prey unprofitability, the manner in which predators encountered prey, and the

features that distinguished profitable and unprofitable prey. Specifically, I tested the following

hypotheses and predictions. First, the degree of prey unprofitability determined predator

avoidance; predators would have lower attack probabilities towards more unprofitable prey, such

as those that were highly bitter or toxic. Second, manner of prey encounter would affect the

knowledge of prey distribution and availability. When predators encountered profitable and

unprofitable prey simultaneously, predators would attack unprofitable prey less due to the

knowledge that suitable prey were also present, resulting in lower attack probabilities. Third, the

complexity of signals or cues differentiating prey types would affect their efficacy in predator

education and therefore prey avoidance. Other things being equal, signals or cues that were more

effective in educating predator would result in lower attack probabilities. One could argue that

multi-component features might be more effective for predator education, as these offer more

precise information to help predators differentiate between prey types. If this was true, more

complex signals would result in lower final attack probabilities. Yet, this assumed that predator

had the cognitive capacity to process and remember the information. If this was not true, more

complex features would confuse the predators, resulting in lower educational efficacy and higher

final attack probabilities.

Materials and Methods

All simulations and statistical analyses were performed in R (version 4.2.2, R Core Team).

Simulating predator-prey interactions and predator learning

The basic model

I constructed an individual-based model with discrete time steps and simulated interactions

between Pavlovian predators and aposematic prey in one single habitat. The model did not

consider any spatial segregation or habitat structuring as the intention here was to focus on

explaining local warning diversity with predator learning alone without involving other

mechanisms. The model also did not include a genetic component in which new mutant alleles

arose within populations to produce novel warning signals. The reason for this choice was that in

well studied Müllerian mimetic systems the coexistence of distinct warning signals is mostly the

result of secondary contact between patterns that have evolved allo- or parapatrically, instead of

resulting from sympatric originations of novel phenotypes (Mallet 1993; Posso-Terranova &

Andrés 2016; Ebersbach *et al.* 2020). Under these scenarios, a genetic component describing the

emergence of new warning signals within populations was not essential for capturing the

dynamics of the predator-prey interactions and therefore not considered.

There were 20 predator individuals and two prey types in the simulated community, one

displaying a more abundant warning signal and the other displaying a rare warning signal. The

more abundant warning signal was already familiar to the predators, and the rare signal was

unfamiliar. Even though there could potentially be multiple species within each prey type,

species identity was irrelevant as individuals within each group were treated as phenotypically

indistinguishable. The local prey had an initial abundance of 200, and the new prey had an initial

abundance of 10, 20, and 40 in separate simulations. Each simulation was run for 50,000 time

steps, and in each time step a predator was randomly selected to be the forager. The forager

would encounter prey based on relative abundance and then decide whether to attack the prey

based on the current attack probability towards that prey type. Throughout the simulation,

predators continued to modify their attack probabilities towards the two prey types following the

Pavlovian learning rules (Speed 1993); section II of SI). The attack probability towards the

familiar prey was set at its final asymptotic value at the beginning of the simulation, reflecting

the fact that predators have completed learning. The attack probability towards the unfamiliar

prey was set at 0.5 at the onset of the simulation, signifying predator naivete. There was no

predator mortality in the model; nor was there competition between predators or between prey.

Prey mortality was due solely to predation, jointly determined by encounter rate and attack

probability that changed dynamically between time steps.

Reproduction occurred every 100 time steps in all simulations, and prey were perennial with

overlapping generations. No predation occurred during reproduction and both prey types

replenished their populations following a logistic growth function:

$$\Delta N_i = r N_i \left(1 - \frac{N_i}{K_i}\right) \tag{4}$$

For prey type i, ΔN was the increase in population size, N was the population size at the end of

the previous time step, r was the intrinsic rate of increase, which was set to be the same for both

prey types, and K was the carrying capacity. For both prey types, I set K as equal to their initial

abundance. This constrained the relative abundance of the rare warning signal to be close to the

initial condition throughout the simulation (i.e., a relative abundance of 0.2, 0.1, or 0.05).

Separate K values for each prey group also reflected the fact that individuals bearing different

signals often use different resources (Posso-Terranova & Andrés 2016; Jiggins 2017).

For each simulation, the parameter space of the simulations was a 10 by 10 grid defined by the

final attack probability and forgetting rate, both of which ranged from 0 to 0.5 (Equations S1 and

S2 in section II of SI). I considered final attack probability between 0 and 0.5 as these values

represented unprofitable prey that were avoided. For forgetting rate, the value of 0 denoted no

forgetting, whereas the value of 0.5 represented the most forgetful predators, which would forget

all the learned avoidance towards even the most unprofitable prey within one single time step

without reinforcement. I ran 20 replicate simulations for each combination of the parameter

values. Abundance of the both prey types were recorded at the end of each simulation. There

were three possible simulation outcomes: the extinction of both prey types, the extinction of prey

with the rare warning signal, and the persistence of both prey types. The outcome also depended

on the intrinsic rate of prey increase (r), which in theory could take any positive value. To select

reasonable r values, I used the range of r commonly observed in insects and some mammals

(Schmitz & Lavigne 1984; Hickling & Pekelharling 1989; Puche & Funderburk 1992; Jahn *et al.*

2005; Emmert *et al.* 2008; Damos 2013; Papanikolaou *et al.* 2014).

Generalization between prey types

The above procedure described the basic scenario, in which avoidance learning for the local and

the new prey occurred independently – predator learning with one prey type did not alter their

attack probabilities towards the other prey type. Each predator also learned independently. To

model the effect of generalization, I introduced a parameter G to account for the fact that change

in the attack probability towards prey type i would have a carry-over effect on the attack

probability towards prey type j, such that:

$$\Delta P_{L,j,t} = G \times \Delta P_{L,i,t} \tag{5}$$

In addition,

251
$$P_{new,t=0} = 0.5 - (0.5 - \lambda) \times G \tag{6}$$

The equation above reflected the fact that the initial attack probability towards the new prey was

affected by its similarity to the local prey. Overall, a non-zero G meant that predators' initial

attack probability towards the new prey would be reduced based on its similarity to the local prey

and that change in attack probability from attacking one prey type would simultaneously reduce

the attack probability towards the other prey type. When G equaled one, the attack probability

towards the new prey would already be identical to the final attack probability at the onset of the

simulation, and learning was completely transferable between prey types. A G equaled zero was

equivalent to the basic scenario described above. In simulations with generalization, G was set to

be 0.5.

Social learning

In separate sets of simulations, I incorporated social learning in the Pavlovian learning

framework following the formulation in (Thorogood *et al.* 2018) by introducing two parameters.

The first parameter b determined the percentage of non-forager individuals that were able to

observe the prey interaction of the forager. Larger b values meant that more predators would

benefit from social learning. The other parameter q described the efficacy of social learning.

Similar to how the parameter G functioned, q was a percentage denoting a "discount" in learning

effect relative to first-hand experience, such that

269
$$\Delta P_{L,m,i,t} = q \times \Delta P_{L,n,i,t} \tag{7}$$

At time step t, predators m observed the foraging outcome between the forager n and prey type i.

If q equaled one, social learning would cause the same decrease in attack probability for

observers as it would from direct learning. If q equaled zero, witnessing a foraging outcome

would have no effect on the observers. In simulations with social learning, both b and q were set

to be 0.5.

Model validation with empirical learning data

Extracting final attack probabilities and forgetting rates from published studies

I performed literature search using keyword combinations (topic = *aposemat** or defen* or

avoid* + topic = learning) in the Web of Science database, yielding a total of 387 articles. I then

screened for studies that quantified attack probability in the context of avoidance foraging with

focal animals under normal conditions (e.g., not injured, not sick, not under high stress, not

surgically or genetically modified). This resulted in 112 articles from which I extracted attack

probabilities. These studies all followed a standard experimental design, in which animals were

presented with prey in a series of learning trials. I calculated predators' attack probabilities

towards the unprofitable prey from the last trial as the final attack probability (section III.1 of SI).

Calculating attack probabilities also allowed for the estimations of forgetting rate, the magnitude

of generalization and social learning efficacy. When applicable, forgetting rate was calculated as

the difference in attack probabilities between trials immediately before and after the forgetting

period, divided by the duration of that period. These forgetting rates were then scaled, when

necessary, to be percent changes per day (see the section below). I also extracted the strength of

generalization and social learning from studies that quantified these effects (section III.2 of SI).

Mapping empirical learning data onto model parameter space

For predators whose final attack probabilities and forgetting rates towards unprofitable prey are

quantified, their data can be mapped onto the parameter space of the model to evaluate their

potential for permitting local warning signal diversity. Empirically measured final attack

probability was immediately appropriate for the parameter space in the simulations. Mapping

empirical forgetting rates onto the parameter space required one extra step. In the model, each

time step equaled 0.01 prey generation time. Using a prey generation time of 30 days, each time

step in the model was then equivalent to 0.3 days. An empirical forgetting rate of 0.1 per day

would thus equal to a value of 0.03 in the model. In general, any empirically measured forgetting

rate could be scaled to unit per day and converted to values appropriate for the parameter space

by multiplying the per day value with 0.01 focal prey generation time.

Meta-analysis of the variation in final attack probability

Since final attack probability was an important determinant for the likelihood of local warning

signal diversity, I performed a meta-analysis to examine the roles of three factors in determining

predators' final attack probabilities towards prey: the reason for prey unprofitability, the manner

in which predator encountered prey, and the feature that differentiated profitable prey from

unprofitable ones. To obtain a sufficiently balanced dataset, I focused on 39 of the 112 studies

for the meta-analysis, with a total of 134 final attack probability measurements (Supplementary

Data). These studies shared enough similarities in experimental design while varying in crucial

aspects, allowing for meaningful analyses. The focal predators were either the great tit (*Parus*

major), the blue tit (*Cyanistes caeruleus*), the European starling (*Sturnus vulgaris*), or the

domestic chicken (*Gallus gallus domesticus*). In addition, animals were presented with both

control and unprofitable artificial prey within each learning trial, either simultaneously or

sequentially. The reason for prey unprofitability was either bitterness (using harmless, bitter

chemicals) or toxicity (using toxic prey extracts), and the feature that consistently differentiated

the two prey types was either color alone, pattern alone (i.e., the distribution of colors or the use

of different geometric symbols), both color and pattern, or color and size or odor.

To perform statistical analyses, I used individual measurements of final attack probability as the

response variable, with species and the aforementioned three factors as predictors. Individual

study was originally included as a random-effect factor to account for potential nonindependence

of data points within studies. However, doing so resulted in model overfitting, and I excluded

study from the statistical model. I also did not consider the phylogenetic relationship between

species since the number of species was small. To additionally test whether the effective

concentration of the bitter chemical was associated with variation in final attack probabilities, I

focused on 35 of the 39 studies that used bitter experimental prey. The majority of the studies

used quinine derivatives, whereas four studies used denatonium benzoate (Supplementary Data).

(Skelhorn & Rowe 2005) reported that 2% quinine sulfate and 0.1% denatonium benzoate

(weight percentages) elicited comparable aversion in domestic chicks, signifying a 20-fold

difference in effective concentration between the two. I therefore converted the concentrations of

denatonium benzoate to equivalent concentrations of quinine derivatives by multiplying them by

20. To test whether an increase in the concentration of bitter compounds caused lower attack

probabilities, I performed a gamma regression with manner of prey encounter, the standardized

concentration of bitter compounds, and the interaction between the two as fixed-effect predictors.

More details on model validation are described in section III.3 of SI. When reporting results, I

followed (Muff *et al.* 2021) and used the language of evidence instead of significance dichotomy.

342   **Results**

Unforgetful predators can allow for local warning signal diversity, especially with

generalization and social learning

The persistence of the rare signal depended on its relative abundance, its intrinsic rate of increase

(r), predators' final attack probability, and predator forgetting rate (Figure 1). Across the range

of r, predators with lower final attack probabilities and lower forgetting rates were more

conducive to the persistence of the rare signal, with forgetting rate being a more limiting factor

(Figure 1). Given the r values in the simulations, the rare signal was unlikely to persist when its

relative abundance was below 0.05 under the basic scenario (Figure 1), indicating a threshold of

relative abundance for warning signal diversity if there was no generalization and social learning.

For prey with higher r values, the condition for signal diversity would be more forgiving

regarding predator learning. Both generalization and social learning increased the likelihood of

warning signal diversity (Figure 2). When the new prey was on the brink of ecological

elimination, however, the existence of generalization appeared inconsequential, while the

facilitative effect of social learning was the most striking (bottom graphs in Figure 2). Nine pairs

of final attack probabilities and forgetting rates from six species (one spider, one fish, one lizard,

two birds, and one mammal) were mapped onto the parameter space to evaluate their potential

for facilitating local warning signal diversity (Table S1 and Figure 2). The focus here was not the

species identity *per se*, but the combination of their final attack probabilities and forgetting rates.

There were only nine data points because forgetting rates were rarely quantified in avoidance

learning studies. Even with limited number of cases, the combination of avoidance level and

forgetting rate from most of the predators were at the lower left corner of the parameter space,

suggesting that it might not be a rare occurrence that natural predators could facilitate local

warning signal diversity. Empirical values of generalization and social learning efficacy were

listed in Table S2.

Causes of variation in final attack probabilities

Compared with bitter prey, there was very strong evidence that toxic prey led to a higher

probability of complete avoidance (i.e., an attack probability of zero), but being toxic did not

reduce non-zero attack probabilities (Table 1 and Figure 3A). There was also no evidence that

increasing bitterness led to lower attack probabilities ($z = -0.23$, $P = 0.82$, Figure 3B). There was

very strong evidence that predators avoided unprofitable prey more when profitable prey were

also present (Table 1 and Figure 3C). Finally, there was no evidence that predators avoided

unprofitable prey differently when the feature differentiating prey types involved only color,

only pattern, or color plus size or odor (Table 1 and Figure 3D). However, when the signals or

cues differed in both color and pattern, there was very strong evidence that predators avoided

unprofitable prey less, suggesting that signals with complex color patterns might be less effective

for predator education, at least for the four avian species examined (Table 1 and Figure 3D).

384   **Discussions**

Local warning signal diversity is not paradoxical

When predators can retain learned avoidance for long enough, the persistence of rare signals in communities can be achieved through balanced predation and reproduction and is entirely nonparadoxical. Empirical learning data also suggest that natural predators from a range of major taxa may have sufficiently low forgetting rates to allow for local warning signal diversity. However, local warning signal diversity was unlikely to occur if the novel signal is less than 5% in relative abundance, at least under commonly observed intrinsic rates of increase and without generalization or social learning. Clearly, this value depends on the prey intrinsic rate of increase and should not be taken as a hard threshold. Even though the relative abundance of stably coexisting warning signals has only been quantified for a small number of systems, it appears to be generally true that the rare signals often constitute at least ~ 5% of the population (Myers *et al.* 1995; Chouteau *et al.* 2016; Thurman *et al.* 2019). The result may also help further explain the observation that warning signal diversity is often the result of secondary contact following dispersal (Mallet 1993; Posso-Terranova & Andrés 2016; Ebersbach *et al.* 2020), in which neither signal is likely to be astronomically dominant in number.

The diversity and convergence of warning signals at the regional and local scales has been thought to reflect a shifting balance involving local drift (phase I), local selection (phase II), and the spread of competing, locally adaptive peaks from regional to local scales (phase III)(Mallet & Joron 1999; Mallet 2010). Results in this study offer further ecological insights into the shifting balance process. First, generalization and especially social learning can further flatten

the "adaptive trough" for rare warning signals (Figure 1 in (Mallet & Joron 1999). This can

facilitate the establishment of new waring phenotypes locally through processes such as kin

founding, especially if individuals with the new warning signals have a fitness advantage, for

example, from being more unprofitable or more conspicuous. When all prey are equally

unprofitable, the facilitative effects of generalization and social learning are only apparent when

the prey intrinsic rate of increase is sufficiently high. The fact that number-dependent predator

education does not necessarily lead to positive FDS also points to an alternative outcome during

phase III of the shifting balance, in which two adaptive peaks meet without one replacing the

other. Interestingly, if the number of immigrants with novel warning signals are large enough,

aside from a higher likelihood of local warning signal diversity it also enhances the opportunities

for the local aposematic prey to survive when predators are more forgetful and/or when the prey

themselves are not severely unprofitable (the size and distribution of red areas in the rightmost

column of Figure 1).

Aside from predator forgetting, positive FDS can result when predator turnover rate is

comparable to prey generation time, such that every prey generation faces naïve predators. Even

though the simulations did not directly model predator turnover, the erosion of learned avoidance

over time can be viewed as a consequence of predator turnover in a community that is similar to

forgetting of individual predators. For example, if each predator in the model is treated as a

group that contains multiple individuals, the same process in which attack probabilities increase

towards 0.5 over time can represent the replacement of experienced predator individuals with

naïve ones in each predator group. The only modification required for simulating predator

turnover in the model is that the erosion of learned avoidance will occur at each time step

regardless of whether predators attack the focal prey. In contrast with predator forgetting,

predator turnover is guaranteed to take place at each time step and can therefore have a stronger

limiting effect on the likelihood of local warning signal diversity. Since most Müllerian prey are

insects and frogs, a substantial turnover of predators within one prey generation may be unlikely

(e.g., one month for butterflies and one year for poison frogs, (Duellman & Trueb 1986; Jiggins

2017). For Müllerian prey with longer generation time, such as the putative mimicry rings

between some mustelid, viverrid, and herpestid mammals (Caro 2005), significant predator

turnover could occur between prey generations and thus might play a more pivotal role.

Testing the unforgetful predator hypothesis in natural communities

All key parameters of the unforgetful predator hypothesis - final attack probability, forgetting

rate, strength of generalization, and the efficacy of social learning - can be measured with

learning experiments. Usually, these parameters are measured in a laboratory setting, which

allows data collection from individual predators. Laboratory experiments are particularly useful

in assessing the variation in learning parameters within and among individuals. In communities

where individual predators can be identified and their learning parameters estimated, the model

can be easily modified such that predators differ in their Pavlovian learning parameters in ways

that faithfully reflecting the observed heterogeneity in predator behavior and cognition. Even

though data are still limited, it appears that predators with simpler or less derived nervous

systems (e.g., spiders) can have forgetting rates comparable to those of birds and mammals. This

suggests that predator possessing low enough forgetting rate for allowing local warning signal

diversity might be more common than we might expect. When collecting learning data from

individual predators are not feasible, field learning experiments represent a suitable alternative.

The design of field learning experiments is similar to laboratory experiments, except that prey

are presented to all potential predators in the community instead of targeting specific predator

individuals (e.g., (Noonan & Comeault 2009; Stevens *et al.* 2009; Chouteau & Angers 2011;

Casas☐Cardona *et al.* 2018; Fukuda & Konuma 2019; Hernández-Agüero *et al.* 2020; Yamazaki

et al. 2020; Postema 2021). Final attack probabilities, forgetting rates, and the strength of

generalization can be derived from proportions of different prey being attacked in the same way

as in laboratory experiments. The key difference is that data obtained this way represent values

averaged over all potential predators. Field learning experiments can be very useful in testing

how likely warning signal diversity is being maintained by predator learning in complex

communities with highly heterogeneous predators that belong to multiple species, which is the

norm in nature. The efficacy of social learning can also be estimated in field learning

experiments, in which video playbacks of predator-prey interactions are presented to predators

before measuring their attack probability. The result can then be compared to data obtained from

direct learning conducted in a separate site of the same community to estimate the efficacy of

social learning. So far, no field learning experiment has attempted to quantify forgetting rate, nor

has one quantified the efficacy of social learning. In fact, the vast majority of the avoidance

learning studies has focused on final attack probability as an indicator for overall level of

avoidance, and the lack of data on forgetting rate, the strength of generalization, and social

learning efficacy are major knowledge gaps, as well as fruitful future directions. Filling this

knowledge gap is crucial for more broadly understanding the role of predator learning in the

maintenance of waring signal diversity in natural communities.

Variation in final attack probabilities and the implications for local warning signal diversity

Even though data in the meta-analyses were conducted on a small number of birds under

laboratory settings, findings from the meta-analysis are broadly informative as the experiments

were all designed to capture the essence of predator learning. The analysis revealed that being

toxic can be more effective than unpalatability for dissuading some predators, resulting in

complete avoidance. However, for predators that would consume toxic prey, either because of a

higher toxicity tolerance or more urgent energetic needs, toxicity does not appear to offer more

protection than mere bitterness. Studies that presented toxic food to predators either used

naturally toxic prey (D'Heursel & Haddad 1999; Cock & Matthysen 2001) or prepared artificial

food items with extracted prey toxin (Chouteau *et al.* 2019). These prey have chemical defenses

that dissuade rather than severely harm predators, which is how defense is expected to function

in Müllerian systems. It is therefore reasonable to extrapolate this result to how predators might

avoid Müllerian prey in general. In addition, increasing bitterness does not appear to be more

effective in eliciting higher avoidance, at least within the range of bitterness in the analysis

(weight percentage 0.002 – 0.1). It is possible that the 50-fold difference in the effective

concentration of bitter chemicals does not result in corresponding changes in bitter perception in

birds. Alternatively, this result could indicate that birds respond to prey bitterness in a uniform

fashion, despite continuous bitter perception. Data on taste sensory physiology, coupled with

controlled tests of taste response, would help differentiate these two hypotheses. All in all,

results from the meta-analysis suggest that Müllerian systems with toxic prey could have higher

potential for local warning signal diversity, but only if the toxins cause complete avoidance. On

the other hand, local warning signal diversity is equally likely to occur in Müllerian systems with

highly distasteful species than in those whose prey are less obnoxious in taste. Evasiveness is

another important cause of prey unprofitability but is not include in the meta-analysis because

only one study has quantified attack probabilities towards evasive prey. Using the blue tit as

predators, (Paez *et al.* 2021) showed that evasive prey with conspicuous colorations elicited a

lower level of avoidance but a stronger generalization effect than colorful, chemically defended

prey. Examining predator learning in other Müllerian systems with evasive prey (e.g., the

Amazonian *Morpho* butterflies (Llaurens *et al.* 2021; Roy *et al.* 2021) and more predators will be

important for further testing the effectiveness of being evasive in eliciting predator avoidance

and to examine whether local warning signal diversity is more or less likely to occur in these

systems.

Predators avoid unprofitable prey more when profitable alternatives are present in the

environment. Even though this result comes from laboratory experiments, it has important

implications for predator-prey interactions in natural communities. The simultaneous and

sequential prey presentations in laboratory experiments represent two extremes of the prey-

encounter continuum. In the field, the foraging habits of predators and the density and

distribution of prey can both determine how predators encounter prey. Sit-and-wait predators are

more likely to encounter prey in a sequential manner than widely ranging predators and therefore

might avoid unprofitable prey to a lesser extent. Similarly, predators are more likely to encounter

prey sequentially in habitats where prey density is low and/or where prey do not aggregate.

Under these conditions, predators are also expected to exhibit lower avoidance towards

unprofitable prey. Findings from the meta-analysis suggest that, all else being equal, local

warning signal diversity is more likely to occur in communities when main predators actively

search for prey and when prey are abundant and/or have a concentrated distribution. These

predictions have never been tested. In addition to predator forgetting rate, a comparative analysis

examining the relationship between the presence of local warning signal diversity, predators'

foraging habits, and prey distribution would offer valuable insight.

The nature of the warning signal can affect predators' final attack probabilities; predators

avoided unprofitable prey more when they differed from profitable prey in only color or pattern.

Studies have also shown that different colors vary in their educational efficacy (Marples *et al.*

1998; Skelhorn & Rowe 2006a; Miller & Pawlik 2013; Nokelainen *et al.* 2014; Taylor *et al.*

2014, 2016). Based on this finding, local warning diversity might be more easily achieved in

communities where profitable prey have simple visual appearance and could be differentiated

from unprofitable prey based on color or pattern alone. This does not mean that unprofitable prey

cannot possess visually complex warning signals, as long as predators can prioritize on color or

pattern to select suitable prey. There is ample evidence indicating that predators do focus on

certain features of prey appearance more than others during avoidance learning (unequal feature

salience or overshadow effects; (Kikuchi & Pfennig 2010, 2013; Kazemi *et al.* 2014, 2015;

Kikuchi & Sherratt 2015; Kikuchi *et al.* 2016). However, it is worth noting that this finding was

based on results from a small number of species and might have limited generality. Particularly,

we know next to nothing about avoidance learning abilities in tropical birds, which are the main

predators of the most classic Müllerian prey (e.g., (Langham 2004, 2005; Pinheiro & Campos

2019). Quantifying learning and forgetting in tropical avian predators can be challenging but

would provide crucial information on the maintenance of local warning signal diversity in some

of the most remarkable Müllerian systems.

Acknowledgements

The author thanks Po-Ju Ke, Yi Sun, and the McMillan lab at the Smithsonian Tropical Research Institute for their helpful comments on earlier drafts of this manuscript. The author would also like to thank the Su lab at Kaohsiung Medical University for lending cloud computing resources. This study was supported by the Kaohsiung Medical University Research Foundation (grant number: KMU-Q111001 and KMU-Q112001).

References

Aronsson, M. & Gamberale-Stille, G. (2008). Domestic chicks primarily attend to colour, not pattern, when learning an aposematic coloration. *Anim Behav*, 75, 417–423.

Aronsson, M. & Gamberale-Stille, G. (2009). Importance of internal pattern contrast and contrast against the background in aposematic signals. *Behav Ecol*, 20, 1356–1362.

Aronsson, M. & Gamberale-Stille, G. (2012). Colour and pattern similarity in mimicry: evidence for a hierarchical discriminative learning of different components. *Anim Behav*, 84, 881–887.

Aubier, T.G., Elias, M., Llaurens, V. & Chazot, N. (2017). Mutualistic mimicry enhances species diversification through spatial segregation and extension of the ecological niche space. *Evolution*, 71, 826–844.

Aubier, T.G. & Sherratt, T.N. (2015). Diversity in Müllerian mimicry: The optimal predator

sampling strategy explains both local and regional polymorphism in prey. *Evolution*, 69,

2831–2845.

Barnett, C., Bateson, M. & Rowe, C. (2007). State-dependent decision making: educated

predators strategically trade off the costs and benefits of consuming aposematic prey. *Behav*

Ecol, 18, 645–651.

Barnett, C.A., Bateson, M. & Rowe, C. (2014). Better the devil you know: avian predators find

variation in prey toxicity aversive. *Biol Letters*, 10, 20140533.

Barnett, C.A., Skelhorn, J., Bateson, M. & Rowe, C. (2012). Educated predators make strategic

decisions to eat defended prey according to their toxin content. *Behav Ecol*, 23, 418–424.

Beatty, C.D., Beirinckx, K. & Sherratt, T.N. (2004). The evolution of Müllerian mimicry in

multispecies communities. *Nature*, 431, 63–66.

Beccaloni, G.W. (1997). Vertical stratification of ithomiine butterfly (Nymphalidae: Ithomiinae)

mimicry complexes: the relationship between adult flight height and larval host☐plant height.

Biol J Linn Soc, 62, 313–341.

Beneš, J. & Veselý, P. (2017). The ability of lizards to identify an artificial Batesian mimic.

Zoology, 123, 46–52.

Blakiston, T. & Alexander, T. (1883). Protection by mimicry - a problem in mathematical

zoology. *Nature*, 29, 405–406.

Bosque, R.J., Lawrence, J.P., Buchholz, R., Colli, G.R., Heppard, J. & Noonan, B. (2018).

Diversity of warning signal and social interaction influences the evolution of imperfect

mimicry. *Ecol Evol*, 8, 7490–7499.

Briolat, E.S., Burdfield□Steel, E.R., Paul, S.C., Rönkä, K.H., Seymoure, B.M., Stankowich, T.,

et al. (2019). Diversity in warning coloration: selective paradox or the norm? *Biol Rev*, 94,

388–414.

Caro, T. (2005). *Antipredator Defenses in Birds and Mammals*. University of Chicago Press,

Chicago, IN, USA.

Casas□Cardona, S., Márquez, R. & Vargas□Salinas, F. (2018). Different colour morphs of the

poison frog Andinobates bombetes (Dendrobatidae) are similarly effective visual predator

deterrents. *Ethology*, 124, 245–255.

Chouteau, M. & Angers, B. (2011). The role of predators in maintaining the geographic

organization of aposematic signals. *The American Naturalist*, 178, 810–817.

Chouteau, M., Arias, M. & Joron, M. (2016). Warning signals are under positive frequency-

dependent selection in nature. *Proceedings of the National Academy of Sciences*, 113, 2164–

2169.

Chouteau, M., Dezeure, J., Sherratt, T.N., Llaurens, V. & Joron, M. (2019). Similar predator

aversion for natural prey with diverse toxicity levels. *Anim Behav*, 153, 49–59.

Cock, R.D. & Matthysen, E. (2001). Do Glow□Worm Larvae (Coleoptera: Lampyridae) Use

Warning Coloration? *Ethology*, 107, 1019–1033.

Damos, P. (2013). Demography and Randomized Life Table Statistics for Peach Twig Borer

Anarsia lineatella (Lepidoptera: Gelechiidae). *J Econ Entomol*, 106, 675–682.

D'Heursel, A. & Haddad, C.F.B. (1999). Unpalatability of Hyla semilineata tadpoles (Anura) to

captive and free-ranging vertebrate predators. *Ethol Ecol Evol*, 11, 339–348.

Duellman, W.L. & Trueb, L. (1986). *Biology of Amphibians*. McGraw-Hill Book Company, New

York, NY, USA.

Dugas, M.B., Halbrook, S.R., Killius, A.M., Sol, J.F. & Richards☐Zawacki, C.L. (2015). Colour

and Escape Behaviour in Polymorphic Populations of an Aposematic Poison Frog. *Ethology*,

121, 813–822.

Ebersbach, J., Posso☐Terranova, A., Bogdanowicz, S., Gómez☐Díaz, M., García☐González,

Ma.X., Bolívar☐García, W., *et al.* (2020). Complex patterns of differentiation and gene flow

underly the divergence of aposematic phenotypes in Oophaga poison frogs. *Mol Ecol*, 29,

1944–1956.

Emmert, C.J., Mizell, R.F., Andersen, P.C., Frank, J.H. & Stimac, J.L. (2008). Diet Effects on

Intrinsic Rate of Increase and Rearing of Proprioseiopsis asetus (Acari: Phytoseiidae). *Ann*

Entomol Soc Am, 101, 1033–1040.

Fabricant, S.A., Exnerová, A., Ježová, D. & Štys, P. (2014). Scared by shiny? The value of

iridescence in aposematic signalling of the hibiscus harlequin bug. *Anim Behav*, 90, 315–325.

Franks, D.W. & Noble, J. (2004). Batesian mimics influence mimicry ring evolution. *Proc Royal*

Soc Lond Ser B Biological Sci, 271, 191–196.

Fukuda, S. & Konuma, J. (2019). Using three-dimensional printed models to test for

aposematism in a carabid beetle. *Biol J Linn Soc*, 128, 735–741.

Gagliardo, A. & Guilford, T. (1993). Why do warning-coloured prey live gregariously? *Proc*

Royal Soc Lond Ser B Biological Sci, 251, 69–74.

Gompert, Z., Willmott, K. & Elias, M. (2011). Heterogeneity in predator micro-habitat use and

the maintenance of Müllerian mimetic diversity. *Journal of theoretical biology*, 281, 39–46.

Greenwood, J.J.D., Cotton, P.A. & Wilson, D.M. (1989). Frequency□dependent selection on

aposematic prey: some experiments. *Biol J Linn Soc*, 36, 213–226.

Halpin, C.G., Skelhorn, J. & Rowe, C. (2013). Predators' decisions to eat defended prey depend

on the size of undefended prey. *Anim Behav*, 85, 1315–1321.

Halpin, C.G., Skelhorn, J. & Rowe, C. (2014). Increased predation of nutrient-enriched

aposematic prey. *Proc Royal Soc B Biological Sci*, 281, 20133255.

Hartig, F. (2016). DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression

models. R package version 0.1.0. https://github.com/florianhartig/DHARMa

Hämäläinen, L., Mappes, J., Rowland, H.M., Teichmann, M. & Thorogood, R. (2020). Social

learning within and across predator species reduces attacks on novel aposematic prey. *Journal*

of Animal Ecology, 89, 1153–1164.

Hämäläinen, L., Mappes, J., Rowland, H.M. & Thorogood, R. (2019). Social information use

642   about novel aposematic prey is not influenced by a predator's previous experience with toxins.

643   *Functional Ecology*, 33, 1982–1992.

644 Hernández-Agüero, J.A., Polo, V., García, M., Simón, D., Ruiz-Tapiador, I. & Cayuela, L.

645   (2020). Effects of prey colour on bird predation: an experiment in Mediterranean woodlands.

646   *Anim Behav*, 170, 89–97.

647 Hickling, G.J. & Pekelharling, C.J. (1989). Intrinsic rate of increase for a brushtail possum

648   population in Rata/Kamahi forest, Westland. *New Zealand Journal of Ecology*, 1989, 117–

649   120.

650 Ihalainen, E., Lindström, L. & Mappes, J. (2007). Investigating Müllerian mimicry: predator

651   learning and variation in prey defences. *Journal of Evolutionary Biology*, 20, 780–791.

652 Jahn, G.C., Almazan, L.P. & Pacia, J.B. (2005). Effect of Nitrogen Fertilizer on the Intrinsic

653   Rate of Increase of Hysteroneura setariae (Thomas) (Homoptera: Aphididae) on Rice (Oryza

654   sativa L.). *Environ Entomol*, 34, 938–943.

655 Jiggins, C.D. (2017). *The Ecology and Evolution of Heliconius Butterflies*. Oxford University

656   Press, Oxford, UK.

657 Joron, M. & Iwasa, Y. (2005). The evolution of a Müllerian mimic in a spatially distributed

658   community. *Journal of Theoretical Biology*, 237, 87–103.

659 Joron, M. & Mallet, J. (1998). Diversity in mimicry: paradox or paradigm? *Trends in Ecology &*

660   *Evolution*, 13, 461–466.

Kazemi, B., Gamberale-Stille, G. & Leimar, O. (2015). Multi-trait mimicry and the relative

salience of individual traits. *Proceedings of the Royal Society B: Biological Sciences*, 282,

20152127–7.

Kazemi, B., Gamberale-Stille, G., Tullberg, B.S. & Leimar, O. (2014). Stimulus Salience as an

Explanation for Imperfect Mimicry. *Current Biology*, 24, 965–969.

Kikuchi, D.W., Mappes, J., Sherratt, T.N. & Valkonen, J.K. (2016). Selection for

multicomponent mimicry: equal feature salience and variation in preferred traits. *Behav Ecol*,

27, 1515–1521.

Kikuchi, D.W. & Pfennig, D.W. (2010). Predator Cognition Permits Imperfect Coral Snake

Mimicry. *Am Nat*, 176, 830–834.

Kikuchi, D.W. & Pfennig, D.W. (2013). Imperfect Mimicry and the Limits of Natural Selection.

The Quarterly Review of Biology, 88, 297–315.

Kikuchi, D.W. & Sherratt, T.N. (2015). Costs of Learning and the Evolution of Mimetic Signals.

Am Nat, 186, 321–332.

Langham, G.M. (2004). Specialized avian predators repeatedly attack novel color morphs of

heliconius butterflies. *Evolution*, 58, 2783–2787.

Langham, G.M. (2005). Rufous-tailed jacamars and aposematic butterflies: do older birds attack

novel prey? *Behavioral Ecology*, 17, 285–290.

Lindstedt, C., Lindström, L. & Mappes, J. (2008). Hairiness and warning colours as components

of antipredator defence: additive or interactive benefits? *Anim Behav*, 75, 1703–1713.

Lindström, L., Alatalo, R.V., Lyytinen, A. & Mappes, J. (2001a). Predator experience on cryptic

prey affects the survival of conspicuous aposematic prey. *Proc Royal Soc Lond Ser B*

Biological Sci, 268, 357–361.

Lindström, L., Alatalo, R.V., Lyytinen, A. & Mappes, J. (2001b). Strong antiapostatic selection

against novel rare aposematic prey. *Proc National Acad Sci*, 98, 9181–9184.

Lindström, L., Alatalo, R.V., Lyytinen, A. & Mappes, J. (2004). The effect of alternative prey on

the dynamics of imperfect Batesian and Müllerian mimicries. *Evolution*, 58, 1294–1302.

Lindström, L., Lyytinen, A., Mappes, J. & Ojala, K. (2006). Relative importance of taste and

visual appearance for predator education in Müllerian mimicry. *Anim Behav*, 72, 323–333.

Llaurens, V., Poul, Y.L., Puissant, A., Blandin, P. & Debat, V. (2021). Convergence in sympatry:

Evolution of blue□banded wing pattern in Morpho butterflies. *J Evol Biol*, 34, 284–295.

Mallet, J. (1993). Speciation, raciation, and color pattern evolution in Heliconius butterflies:

evidence from hybrid zones. In: *Hybrid Zones and the Evolutionary Process* (ed. Harrison,

R.G.). Oxford University Press, Oxford, UK, pp. 226–260.

Mallet, J. (2001). Mimicry: An interface between psychology and evolution. *Proc National Acad*

Sci, 98, 8928–8930.

Mallet, J. (2010). Shift happens! Shifting balance and the evolution of diversity in warning

colour and mimicry. *Ecological Entomology*, 35, 90–104.

Mallet, J. & Joron, M. (1999). Evolution of diversity in warning color and mimicry:

polymorphisms, shifting balance, and speciation. *Annual review of ecology and systematics*,

1999, 201–233.

Marples, N.M., Roper, T.J. & Harper, D.G.C. (1998). Responses of wild birds to novel prey:

evidence of dietary conservatism. *Oikos*, 83, 161.

Miller, A.M. & Pawlik, J.R. (2013). Do coral reef fish learn to avoid unpalatable prey using

visual cues? *Anim Behav*, 85, 339–347.

Muff, S., Nilsen, E.B., O'Hara, R.B. & Nater, C.R. (2021). Rewriting results sections in the

language of evidence. *Trends Ecol Evol*, 37, 203–210.

Müller, F. (1879). Ituna and Thyridia; a remarkable case of mimicry in butterflies. (R. Meldola

translation). *Proclamations of the Entomological Society of London*, 20–29.

Myers, C.W., Daly, J.W., Garraffo, H.M., Wisnieski, A. & Jr., J.F.C. (1995). Discovery of the

Costa Rican Poison Frog Dendrobates pumilio, and Comments on Taxonomic Use of Skin

Alkaloids. *American Museum Novitates*, 3144, 1–21.

Nokelainen, O., Valkonen, J., Lindstedt, C. & Mappes, J. (2014). Changes in predator

community structure shifts the efficacy of two warning signals in Arctiid moths. *J Anim Ecol*,

83, 598–605.

Noonan, B.P. & Comeault, A.A. (2009). The role of predator selection on polymorphic

aposematic poison frogs. *Biol Letters*, 5, 51–54.

Paez, E., Valkonen, J.K., Willmott, K.R., Matos-Maravi, P., Elias, M. & Mappes, J. (2021). Hard

to catch: experimental evidence supports evasive mimicrry. *Proceedings of the Royal Society*

B: Biological Sciences, 288, 20213052.

Papanikolaou, N.E., Milonas, P.G., Kontodimas, D.C., Demiris, N. & Matsinos, Y.G. (2014).

Life Table Analysis of Propylea quatuordecimpunctata (Coleoptera: Coccinellidae) at

Constant Temperatures. *Ann Entomol Soc Am*, 107, 158–162.

Pegram, K.V., Fankhauser, K. & Rutowski, R.L. (2021). Variation in predator response to short-

wavelength warning coloration. *Behav Process*, 187, 104377.

Pinheiro, C.E.G. & Campos, V.C. (2019). The responses of wild jacamars (Galbula ruficauda,

Galbulidae) to aposematic, aposematic and cryptic, and cryptic butterflies in central Brazil.

Ecological Entomology, 294, 224–10.

Posso-Terranova, A. & Andrés, J.A. (2016). Complex niche divergence underlies lineage

diversification in Oophaga poison frogs. *Journal of Biogeography*, 43.

Postema, E.G. (2021). The effectiveness of eyespots and masquerade in protecting artificial prey

across ontogenetic and seasonal shifts. *Current Zoology*.

Poulton, E. (1890). *The Colours of Animals: Their Meaning and Use, Especially Considered in*

the Case of Insects. Paul, Trench, Trübner and Co. Ltd., London, UK.

Puche, H. & Funderburk, J. (1992). Intrinsic Rate of Increase of Frankliniella fusca

(Thysanoptera: Thripidae) on Peanuts. *Fla Entomologist*, 75, 185.

R Core Team. (2022). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org

Rönkä, K., Pasqual, C.D., Mappes, J., Gordon, S. & Rojas, B. (2018). Colour alone matters: no

predator generalization among morphs of an aposematic moth. *Animal Behaviour*, 135, 153–

163.

Roper, T.J. (1993). Effects of novelty on taste-avoidance learning in chicks. *Behaviour*, 125,

265–281.

Rowland, H.M., Hoogesteger, T., Ruxton, G.D., Speed, M.P. & Mappes, J. (2010a). A tale of 2

signals: signal mimicry between aposematic species enhances predator avoidance learning.

Behav Ecol, 21, 851–860.

Rowland, H.M., Wiley, E., Ruxton, G.D., Mappes, J. & Speed, M.P. (2010b). When more is less:

the fitness consequences of predators attacking more unpalatable prey when more are

presented. *Biol Letters*, 6, 732–735.

Roy, C.L., Roux, C., Authier, E., Parrinello, H., Bastide, H., Debat, V., *et al.* (2021). Convergent

morphology and divergent phenology promote the coexistence of Morpho butterfly species.

Nat Commun, 12, 7248.

Ruxton, G.D., Sherratt, T.N. & Speed, M.P. (2018). Chapter 9: Evolution and maintenance of

Müllerian mimicry. In: *Avoiding Attack: The Evolutionary Ecology of Crypsis, Warning*

Signals, and Mimicry. Oxford University Press, Oxford, UK, pp. 115–136.

Salva, O.R., Daisley, J.N., Regolin, L. & Vallortigara, G. (2009). Lateralization of social

learning in the domestic chick, Gallus gallus domesticus: learning to avoid. *Anim Behav*, 78,

847–856.

Schmitz, O.J. & Lavigne, D.M. (1984). Intrinsic rate of increase, body size, and specific

metabolic rate in marine mammals. *Oecologia*, 62, 305–309.

Sherratt, T.N. (2011). The optimal sampling strategy for unfamiliar prey. *Evolution*, 65, 2014–

2025.

Siddall, E.C. & Marples, N.M. (2008). Better to be bimodal: the interaction of color and odor on

learning and memory. *Behav Ecol*, 19, 425–432.

Skelhorn, J. (2011). Colour biases are a question of conspecifics' taste. *Anim Behav*, 81, 825–829.

Skelhorn, J. & Rowe, C. (2005). Frequency-dependent taste-rejection by avian predation may

select for defence chemical polymorphisms in aposematic prey. *Biol Letters*, 1, 500–503.

Skelhorn, J. & Rowe, C. (2006a). Do the multiple defense chemicals of visually distinct species

enhance predator learning? *Behav Ecol*, 17, 947–951.

Skelhorn, J. & Rowe, C. (2006b). Predator avoidance learning of prey with secreted or stored

defences and the evolution of insect defences. *Anim Behav*, 72, 827–834.

Skelhorn, J. & Rowe, C. (2007). Automimic frequency influences the foraging decisions of avian

predators on aposematic prey. *Anim Behav*, 74, 1563–1572.

Skelhorn, J. & Rowe, C. (2009). Distastefulness as an antipredator defence strategy. *Anim Behav*,

78, 761–766.

Skelhorn, J. & Rowe, C. (2010). Birds learn to use distastefulness as a signal of toxicity. *Proc

Royal Soc B Biological Sci*, 277, 1729–1734.

Smith, K.E., Halpin, C.G. & Rowe, C. (2014). Body size matters for aposematic prey during

predator aversion learning. *Behav Process*, 109, 173–179.

Smith, K.E., Halpin, C.G. & Rowe, C. (2016). The benefits of being toxic to deter predators

depends on prey body size. *Behav Ecol*, 27, 1650–1655.

Speed, M.P. (1993). Muellerian mimicry and the psychology of predation. *Animal Behaviour*, 45,

571–580.

Speed, M.P. (2000). Warning signals, receiver psychology and predator memory. *Animal

Behaviour*, 60, 269–278.

Speed, M.P. (2001). Can receiver psychology explain the evolution of aposematism? *Anim

Behav*, 61, 205–216.

Speed, M.P. & Turner, J.R.G. (2008). Learning and memory in mimicry: II. Do we understand

the mimicry spectrum? *Biological Journal of the Linnean Society*, 67, 281–312.

Stevens, M., Castor-Perry, S.A. & Price, J.R.F. (2009). The protective value of conspicuous

signals is not impaired by shape, size, or position asymmetry. *Behav Ecol*, 20, 96–102.

Taylor, L.A., Amin, Z., Maier, E.B., Byrne, K.J. & Morehouse, N.I. (2016). Flexible color

learning in an invertebrate predator: Habronattus jumping spiders can learn to prefer or avoid

red during foraging. *Behav Ecol*, 27, 520–529.

Taylor, L.A., Maier, E.B., Byrne, K.J., Amin, Z. & Morehouse, N.I. (2014). Colour use by tiny

predators: jumping spiders show colour biases during foraging. *Anim Behav*, 90, 149–157.

Thorogood, R., Kokko, H. & Mappes, J. (2018). Social transmission of avoidance among

predators facilitates the spread of novel prey. *Nature Ecology & Evolution*, 2, 1–10.

Thurman, T.J., Sigal, A.S. & McMillan, W.O. (2019). Movement of a Heliconius hybrid zone

over 30 years: A Bayesian approach. *Journal of Evolutionary Biology*, 32, 974–983.

Turner, J. & Speed, M.P. (1996). Learning and memory in mimicry. I. Simulations of laboratory

experiments. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 351,

1157–1170.

Turner, J.R.G. (1984). Mimicry: the palatability spectrum and its consequences. In: , The

Biology of Butterflies Vol. 11 (eds. Vane-Wright, R.I. & Ackery, P.R.). Academic Press, pp.

141–161.

Turner, J.R.G., Kearney, E.P. & Exton, L.S. (2008). Mimicry and the Monte Carlo predator: the

palatability spectrum, and the origins of mimicry. *Biological Journal of the Linnean Society*,

23, 247–268.

Webb, J.K., Brown, G.P., Child, T., Greenlees, M.J., Phillips, B.L. & Shine, R. (2008). A native

dasyurid predator (common planigale, Planigale maculata) rapidly learns to avoid a toxic

invader. *Austral Ecol*, 33, 821–829.

Willmott, K.R., Willmott, J.C.R., Elias, M. & Jiggins, C.D. (2017). Maintaining mimicry

diversity: optimal warning colour patterns differ among microhabitats in Amazonian

clearwing butterflies. *Proceedings of the Royal Society B*, 284, 20170744–9.

Yamazaki, Y., Pagani-Núñez, E., Sota, T. & Barnett, C.R.A. (2020). The truth is in the detail:

predators attack aposematic prey with less aggression than other prey types. *Biol J Linn Soc*,

131, 332–343.

Figure legends

Figure 1. Outcomes of simulated predator-prey interactions. In each graph, black cells denote the

extinction of both signals, red cells denote the extinction of the rare signal, and blue cells denote

the persistence of both signals. The vertical panels are simulations under five intrinsic rates of

prey increase (r). The three horizontal panels are simulations under different relative abundances

of the rare signal. The occurrence of local warning signal diversity (blue cells) is more likely

under higher r, lower final attack probability (λ), and lower forgetting rate (β).

Figure 2. Simulation results illustrating the facilitative effect of generalization and social

learning on the likelihood of local warning signal diversity, using $r = 0.28$. The coloring of the

cells are the same as in Figure 1. The vertical panels represent results without generalization or

social learning (left), with generalization (middle), and with social learning (right). The

horizontal panels are results under different relative abundances of the rare prey. White dots are

pairs of empirically measured final attack probability and forgetting rate from predators: 1.

Planigale maculata (Webb *et al.* 2008), 2. *Habronattus pyrrithrix* (Taylor *et al.* 2016), 3-5.

Parus major (Lindström *et al.* 2006), 6. *Chalcides sexlineatus* (Beneš & Veselý 2017), 7. *Gallus*

gallus domesticus (Siddall & Marples 2008), 8-9. *Thalassoma bifasciatum* (Miller & Pawlik

2013)

Figure 3. Factors that influenced predators' final attack probabilities towards unprofitable prey.

(A) Reason for unprofitability, (B) level of bitterness, (C) manner of prey encounter, and (D)

type of signal or cue. Red dots in (A) are attack probabilities that equal zero. Yellow dots and

line in (B) are when predators encounter profitable and unprofitable prey sequentially, and blue

dots and line are when predators encounter the two prey types simultaneously. Data were from

(Gagliardo & Guilford 1993; Roper 1993; D'Heursel & Haddad 1999; Cock & Matthysen 2001;

Lindström *et al.* 2001a, 2004, 2006; Skelhorn & Rowe 2005, 2006a, b, 2007, 2009, 2010;

Barnett *et al.* 2007, 2012, 2014; Ihalainen *et al.* 2007; Aronsson & Gamberale-Stille 2008, 2009,

2012; Lindstedt *et al.* 2008; Siddall & Marples 2008; Salva *et al.* 2009; Rowland *et al.* 2010b, a;

Skelhorn 2011; Halpin *et al.* 2013, 2014; Fabricant *et al.* 2014; Smith *et al.* 2014, 2016; Bosque

et al. 2018; Rönkä *et al.* 2018; Chouteau *et al.* 2019; Hämäläinen *et al.* 2019, 2020; Paez *et al.*

2021; Pegram *et al.* 2021)

864 Table 1. Statistical results of the zero-inflated gamma regression examining the cause of

865 variation in predators' final attack probabilities towards unprofitable prey.

Factor	Coefficient	Standard error	z-value	P-value
Conditional component				
Species				
Gallus gallus domesticus	0.52	0.28	1.86	0.063
Parus major	0.54	0.27	2.00	0.05
Sturnus vulgaris	0.70	0.37	1.91	0.06
Cause of unprofitability				
toxin	-0.29	0.31	-0.92	0.36
Type of signal/cue				
color + other	0.38	0.25	1.52	0.13

color + pattern	0.84	0.19	4.32	< 0.0001
pattern	0.40	0.25	1.60	0.11
Prey encounter				
simultaneous	-0.88	0.23	-3.80	0.0001
Zero-inflated component				
Cause of unprofitability				
toxin	4.37	0.89	4.88	< 0.0001