Connecting higher order interactions with ecological stability in experimental aquatic food webs

Chenyu Shen[1,2], Kimberley Lemmen[1], Jake Alexander[2], Frank Pennekamp[1,*]

[1]Department of Evolutionary Biology and Environmental Studies,
University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
[2]Department of Environmental Systems Science, Institute for Integrative Biology, Swiss Federal Institute of Technology, Zurich, Switzerland
* Corresponding author: frank.pennekamp@ieu.uzh.ch

Abstract

Community ecology is built on theory that represents the strength of interactions between species as pairwise links. Higher order interactions occur when the presence of a third (or more) species changes the pairwise interaction between a focal pair. Recent theoretical work has highlighted the stabilizing role of higher order interactions for communities, yet it remains unclear how important higher order effects are in real communities. Here we used experimental communities of aquatic protists to examine the relationship between higher-order interactions and community stability (i.e., the persistence of species in a community). We cultured a focal pair of consumers in the presence of additional competitors and a predator and collected time series data of their abundances. We then fitted competition models with and without HOIs to measure interaction strength between the focal pair across different community compositions. We used survival analysis to measure the persistence of individual species. We found evidence that additional species positively affected persistence of the focal species and that HOIs were present in most of our communities. However, persistence was only linked to HOIs for one of the focal species. Our results vindicate community ecology theory positing that species interactions may deviate from assumptions of pairwise interactions, opening avenues to consider possible consequences for coexistence and community stability.

Key words: interaction modifications, species interactions, persistence, indirect effects, time series

Introduction

Community ecology seeks to unravel the mechanisms that maintain the tremendous organismal diversity on our planet. Species interactions, such as competition and predation, are crucial to the coexistence of species and hence are important drivers of diversity (Chesson, 2000; HilleRisLambers et al., 2012). Most research on species coexistence has focused on pairwise interactions (Levine et al., 2017). However, in nature, species rarely interact only in a direct pairwise fashion, but rather in large networks of interacting species where indirect effects are likely to be important (Levine et al., 2017). Recently, these indirect effects have gained increasing attention, since in some case they may explain the dynamics and stability of communities better than classic pairwise models (Grilli et al., 2017; Letten & Stouffer, 2019; Mayfield & Stouffer, 2017).

Lotka-Volterra competition or predator-prey models assume that an interaction is an intrinsic property of two species interacting in a given environment, and therefore the parameters that determine the strength of interactions are independent of the community in which these species are embedded (Werner & Peacor, 2003). However, when more than two species co-occur (Worthen & Moore, 1991), indirect effects can arise through chains of pairwise interactions, or through higher-order interactions, that is changes of the per capita effect of one competitor on another in presence of additional species (Levine et al., 2017; Wootton, 1994). Since interaction chains and higher-order interactions emerge from fundamentally different mechanisms, it is important to understand their underlying mechanisms for appropriate modelling and inference (Levine et al., 2017).

Interaction chains emerge when pairwise interactions are embedded in a network of interactions. A series of such direct interactions between species pairs can lead to connections between species that do not directly interact with each other (Wootton, 1993). Despite introducing indirect pathways between species, interaction chains are the result of a series of fixed, strictly pairwise interactions. In contrast, "higher-order interactions" (HOIs) (Levine et al., 2017); also called "interaction modifications"; (Abrams, 1983) are indirect effects that arise when the presence of a third species alters the relationship between two interacting species in a way that

changes the per capita effect of one species on another. Wootton (1994) provides examples of both types of indirect effects in the upper zone of a rocky intertidal community. By consuming limpets (*Lottia digitalis*), bird predators indirectly lead to higher acorn barnacle (*Balanus glandula*) abundance because limpets dislodge or accidentally consume young acorn barnacles, an example of an interaction chain. Because the light-coloured shell of *L. digitalis* is similar in color to barnacle shells, barnacles alter the bird-limpet interaction by changing the ability of birds to find limpets, an example of an interaction modification. While HOIs are most commonly described between species, there are also intraspecific HOIs. These represent the cumulative impacts of interactions among the individuals of each 'competitor species' (that is, intraspecific crowding) on the focal species (Letten & Stouffer, 2019; Mayfield & Stouffer, 2017). For the intertidal community, intraspecific HOIs could result from the bird affecting either the limpet or barnacle foraging behaviour which subsequently influences intraspecific competition (e.g., barnacles will aggregate in areas sheltered from bird predation resulting in higher intraspecific competition). Interaction chains can be predicted with only a knowledge of pairwise species interactions; in contrast, descriptions of HOIs require a knowledge of all species combinations involved (Wootton, 1994).

Theoretical work has suggested that empirical studies of interactions between species should consider HOIs (Grilli et al., 2017) given that simple models of pairwise interactions fail to explain the stable persistence in simulation models of very large ecological communities (Barabás et al., 2016; Clark, 2010; Gibbs et al., 2022; Kleinhesselink et al., 2022; Levine et al., 2017). Recently, many theoretical advances have been made to understand the influence of HOIs on community dynamics (Gibbs et al., 2022; Grilli et al., 2017; Letten & Stouffer, 2019; Mayfield & Stouffer, 2017; Singh & Baruah, 2021). A particular focus has been the effect of HOIs on the stability of communities, specifically the persistence of all species in a community.

Using simulation studies (Bairey et al., 2016) observed that communities in which species only interact in a pairwise fashion displayed a negative relationship between persistence and increasing species richness. However, this pattern was not observed in communities with HOIs (where the interaction strengths across

community members were randomly increased or decreased). These results indicated that the negative relationship between diversity and stability discovered by May (May, 1972) could be inverted when HOIs were considered. Furthermore, simulation studies have demonstrated that both positive and negative three-way HOIs can promote species coexistence by altering the per capita strength of pairwise competition (Singh & Baruah, 2021). The stabilising effect of HOIs in highly diverse communities has been observed in competitive network models of forest dynamics in which new species are constantly introduced through immigration and speciation (Grilli et al., 2017).

Despite this support from theory for the importance of higher order interactions for coexistence and community stability, empirical evidence connecting HOIs with ecological stability remains very scarce (but see Mayfield & Stouffer, 2017; Mickalide & Kuehn, 2019). This is partly due to the challenges of empirically quantifying HOIs (Billick & Case, 1994), which requires the evaluation of the interaction strength between two species in the presence of additional species accounting for changes in density (Adler et al., 2018). Even in simple communities, this can quickly become logistically infeasible. In addition, the quantification of the stability of communities requires time-series to evaluate species persistence over extended periods of time, presenting a second logistical hurdle.

Our goal was to experimentally test the causal links between higher-order interactions and the persistence of species within communities. Microcosms are a convenient tool to investigate concepts in community ecology because they are easily manipulated, can be highly replicated, and have large population sizes (Altermatt et al., 2015). Moreover, a range of analytical approaches is available to estimate intra- and interspecific interaction strengths within these systems (Carrara et al., 2015). We therefore used microcosms to experimentally examine HOIs and community persistence in aquatic microbial communities. We used six different community compositions, which all included a focal pair of ciliate consumers competing for the same resource. We tested the effect of two additional competitors as well as the presence of a predator on the interaction strength between the focal pair. Time series of population dynamics were collected to quantify the strength of

species interactions and examine the effect of HOIs on community stability and address the following questions:

1. Are higher order interactions detectable in our communities?
2. Do HOIs differ depending on the identity and trophic role of the third species?
3. Is there a relationship between the presence of HOIs and community persistence?

We hypothesized that additional competitors would influence the effect of the two focal species on each other, since all species engage in resource competition. For instance, in the presence of a third species, focal species may shift their foraging behaviour and thereby their pairwise effect on one another. We also expected that higher-order effects would arise in communities with a predator whose behaviour is affected by different consumer body sizes which could render one prey susceptible to predation while the other would be attacked but is too large to be consumed. All else being equal, we expected a stabilizing effect on the persistence of the populations of the two focal species for HOIs that decreased their interaction strength.

Methods

Data collection: Community Experiment

Food web construction and culture conditions

To experimentally test higher order interactions, we constructed an experimental food web. The ciliates of our microbial food web were primarily bacterivorous (Pennekamp et al., 2018), thus the experimental communities were sustained on a mix of three species of bacteria (*Bacillus subtilis, Serratia fonticola and Brevibacillus brevis*) decomposing the protozoan pellet medium (PPM; provided by Carolina Biological Supplies, Burlington NC, USA; concentration of 0.55 g/L, see Altermatt et al., 2015).

Four bacterivorous ciliates (*Colpidium striatum, Dexiostoma campylum, Paramecium caudatum* and *Spirostomum teres*) constitute the intermediate

consumers in our experimental microcosms. Both *Colpidium striatum* (50-100 um)

and *Dexiostoma campylum* (35-90 um) are consumed by the top predator

Spathidium sp. (40 - 300 um)*,* which cannot survive only on bacteria as prey

(Woodruff & Spencer, 1921). Due to the large body size of *Paramecium caudatum*

(170–300 um) (Foissner & Berger, 1996) and *Spirostomum teres* (150–400 um)

(Bick & World Health Organization, 1972), we expected these species to interfere

with the predator because they are more difficult or impossible to consume.

We chose these five eukaryotic ciliates (hereafter referred to by the genus

name only) because their pairwise trophic interactions are well documented by

previous studies (Daugaard et al., 2019; Pennekamp et al., 2018; Tabi et al., 2019).

The consumptive and competitive interactions present in our experimental food web

are shown in Figure 1.

Figure 1: Microbial food web in our experiment. Consumptive interactions (in black) among
the top predator (*Spathidium sp.*), the consumers *(Colpidium striatum, Dexiostoma*
campylum, *Paramecium caudatum* and *Spirostomum teres*) and a mix of three bacteria
species (*Bacillus subtilis, Serratia fonticola and Brevibacillus brevis*). All the consumers
compete with each other (red arrows), with the focal pair marked in red.

One week before the start of the community experiment, we prepared the bacterial

medium for prey cultures. We first added 3-4mL DMSZ medium into three 20mL

falcon tubes, then transferred *B. subtilis, B. brevis* and *S. fonticola* to these tubes

using inoculation loops and incubated them at 37°C for 24 hours.

After one day of incubation, we added 100µl of *S. fonticola* and *B. subtilis* and

2ml of *B. brevis* medium (since *B. brevis* grows much slower than *S. fonticola* and *B.*

subtilis) from the falcon tubes to three previously autoclaved 1L glass bottles, each

containing 500ml of ciliates pellet medium. We then incubated them at 20°C for two

days to give the bacteria enough time to reach their carrying capacity. The prey

culture medium was produced by mixing 33.3ml of each bacteria medium from the

1L glass bottles into an autoclaved 250 mL jar (i.e., the microcosm), yielding a total

of 100ml of mixed bacterial culture medium. Two wheat seeds were added to each

jar to provide a source of slow release nutrients that the bacteria could feed on,

leading to stable populations of ciliates during our experiment (Altermatt et al., 2015).

All the consumers were grown at 15° C prior to the experiments for three weeks to

reach their carrying capacities.

Experimental design

To assess how the interaction strength between two focal species is altered

by adding a third or more species, we established the following treatments: (1) two

prey (*Colpidium* and *Dexiostoma*) alone to investigate the direct effects between the

two focal species (CD) (2) two prey and *Paramecium* (CDP). (3) two prey and

Spirostomum (CDS). These communities were also grown in the presence of the top

predator *Spathidium (*CDPd, CDPPd, CDSPd; predator = Pd) (Figure 2), yielding six

treatments in total. For each treatment, we cultured four replicates resulting in 24

microcosms in total.

To start the experiment, we took out 20 or 30 mL of the bacterized PPM

medium (depending on experimental community composition) and replaced it with

10mL of stock culture of each consumer species at their carrying capacity.

Therefore, the starting densities of all consumers were 10% of their carrying

capacities. In the predation treatments, four days after the introduction of the

consumer species ten predator individuals were pipetted from the maintenance plates to the microcosms. To assure establishment in each microcosm, another ten individuals of predator were added after 8 days. The experiments were conducted at 15°C with no light, suitable growing conditions for ciliates (Altermatt et al., 2015).

Video sampling and species classification

Experimental units were sampled every second day for 53 days to capture time series of the dynamical changes in abundance of ciliates. For each sampling event, the microcosm was gently agitated, a subsample of 250µL was mounted onto a glass slide and covered with a glass lid. Three five-second videos (at 25 frames per second) were taken using 25✕ magnification on a stereomicroscope (Leica M205 C) mounted with a digital CMOS camera (Hamamatsu Orca Flash 4.0 C11440, Hamamatsu Photonics, Japan) with dark field illumination. We took three subsamples for each microcosm to get a precise estimate of abundance and processed the videos with the R package BEMOVI (version 1.0.2) (Pennekamp et al., 2015). We took the mean of the three videos as our measure of density. If no individuals were detected, we assigned a zero. The volume lost from the microcosm by sampling during the course of the experiment was replaced with fresh bacterized PPM medium.

For species classification we trained a support vector machine classifier (svm) on 350 to 400 randomly chosen and manually labelled trajectories of each species across the community compositions and time with the R package e1071 (Meyer et al., 2022). We also included a "noise" class in our classifier representing spurious trajectories due to background movement. Twenty morphological and movement features extracted from an established classification pipeline (Pennekamp et al., 2017) were selected to train the support vector machine to distinguish among classes based on information about body size and movement patterns (Table S1). As ciliate phenotypes may change over time (Pennekamp et al., 2017), we included "week number" to enhance the accuracy of the classifier. Further details about the classification can be found in the supplementary material.

Data analyses: interaction strengths and community stability

Abundance

To understand the effect of community composition on the abundance of the species, we calculated the mean abundance over the entire 53-day experiment for each species in each of the six communities. Mean abundance of each of the five species was analysed with two-way ANOVA (predictors: identity of the third species [levels: *Paramecium* or *Spirostomum*] and the presence of the predator [levels: present or absent]).

Estimating species interaction strengths

We estimated the effect of species j on the growth rate of species i within a regression framework, where the growth rate of species i is regressed against the densities of species i and j to determine the intra- and interspecific effects on the growth rate of species i (Pfister, 1995). The *per capita* growth rate of each species i was calculated as $\frac{N_{i,t+1}/N_{i,t}}{(t+1)-t}$ where $N_{i,\,t+1}$ is the abundance at time t+1 and $N_{i,t}$ is abundance at time t. We used the gauseR package to calculate the per capita growth rate (Mühlbauer et al., 2020). To explore how interaction strength and HOIs may differ between communities, we compared the fit of six models to the per capita growth rate of the focal species (either *Dexiostoma* or *Colpidium*). First, we fitted the Lotka-Volterra model with no HOIs:

$$\frac{N_{i,t+1}}{N_{i,t}} = \lambda_i - \sum_{j=1}^{n} \alpha_{ij} N_j$$

For the simple additive model, λ_i is the intrinsic growth rate and α_{ij} represents both the intra- and interspecific interaction coefficients. To test for intra- and interspecific HOIs, we then extended the additive model (Letten & Stouffer, 2019). For the interactive LV model only including intra-specific HOIs, we included the interaction term β_{ijj} for the effect of species j on i, where β_{ijj} captures the cumulative impacts of intra-specific interactions on the focal species:

$$\frac{N_{i,t+1}}{N_{i,t}} = \lambda_i - \sum_{j=1}^{n} \alpha_{ij} N_j - \sum_{j=1}^{n} \beta_{ijj} N_j N_j$$

260 The identity of species j may or may not be the same as species i. Next, we fitted an
261 interactive LV model that only includes interspecific HOIs, where β_{ijk} captures the
262 cumulative impacts of interspecific interactions on the focal species:

$$\frac{N_{i,t+1}}{N_{i,t}} = \lambda_i - \sum_{j=1}^{n} \alpha_{ij} N_j - \sum_{j=1}^{n} \sum_{k=j+1}^{n} \beta_{ijk} N_j N_k$$

263 Here, the identity of species k strictly excludes the focal species i.

264 Finally, we tested the fit of the interactive LV model including both intra- and
265 interspecific HOIs (full HOI model):

$$\frac{N_{i,t+1}}{N_{i,t}} = \lambda_i - \sum_{j=1}^{n} \alpha_{ij} N_j - \sum_{j=1}^{n} \sum_{k=j+1}^{n} \beta_{ijk} N_j N_k - \sum_{j=1}^{n} \beta_{ijj} N_j N_j$$

266 To test whether the shape of the density-dependence deviates from the linear form
267 of the additive Lotka-Volterra model, we fitted the Ricker model with nonlinear
268 density dependence. First, we fitted the additive model without HOIs:

$$\frac{N_{i,t+1}}{N_{i,t}} = \lambda_i e^{-\sum_{j=1}^{n} \alpha_{ij} N_j}$$

269 and interactive form only including interspecific HOIs (terms as previously defined):

$$\frac{N_{i,t+1}}{N_{i,t}} = \lambda_i e^{-\sum_{j=1}^{n} \alpha_{ij} N_j - \sum_{j=1}^{n} \beta_{ijk} N_j N_k}$$

We used generalized linear models with the "identity" link function assuming
gaussian errors to estimate the coefficients of the additive and interactive LV model.
For the Ricker model, we used a GLM with the "log" link function, assuming also
gaussian errors. For the fitted models, the intercept can be interpreted as the
maximum intrinsic growth rate, while the regression coefficients represent the *per*
capita effect of species i on itself and per capita effects of species j on species i.
Interaction terms describe the mediating effects that the density of species j can
have on the effect of species i (and vice versa) on the growth rate of the focal

species (Letten & Stouffer, 2019). We performed a AICc model selection to identify
the model that best explained the growth rate of each focal species. We then
calculated the difference (dAICc) between the model with the lowest AICc and all
other models. Models with a dAIC larger than 2 are considered to be different, thus
the model with the fewest parameters and a dAIC less than 2 is considered the most
parsimonious model. Since the presence of predators drove some prey to early
extinction, we only estimated interaction strengths in communities without predators.

Estimating species survival (persistence)

We measured the persistence of the focal species *Colpidium* and
Dexiostoma, as well as the predator *Spathidium* as a proxy of the population stability
of these species using non-parametric survival analysis. We estimated the time to
event (i.e., extinction), accounting for right-censoring, that is, the possibility of
missing future extinctions since we only recorded the abundance for 53 days. We
tested the effect of community composition on the Kaplan-Meier estimate of the focal
species with log rank tests, followed by pairwise tests of community composition in
case of a significant community wide effect with the survminer package
(Kassambara et al., 2021).

All of the above analyses including the species classification were performed with
the statistical computing environment R (R Core Team, 2022).

Results

Population dynamics over time

The population dynamics of the protists differed depending on the community
context (Figure 2). When *Colpidium* and *Dexiostoma* competed only with each other,
we consistently observed extinction of *Colpidium* towards the end of the experiment,
while *Dexiostoma* persisted. The addition of *Paramecium* or *Spirostomum* led to a
longer period of coexistence of *Dexiostoma* and *Colpidium*. The addition of the top
predator *Spathidium* considerably shortened the persistence of all species. Thus, the
addition of a third competitor increased the period of coexistence of the focal species

pair, whereas the addition of the predator led to the rapid collapse of the whole
community.

Figure 2: Population dynamics for each experimental community. Rows are six different
treatments in our experiment. Each treatment is represented by the species it contains (C for
Colpidium, D for *Dexiostoma*, P for *Paramecium*, S for *Spirostomum* and Pd for predator).
Results for each replicate are shown in columns. Most species in competition treatments
(first three rows) survived to the last day of measurement but rapidly dropped below
detection limits when housed with predators (last three rows).

The addition of the competitor *Paramecium* had a negative effect on the mean

abundance of both *Colpidium* (b=-0.60, 95% CI = -0.89 to -0.32) and *Dexiostoma* (b=

-0.81, 95% CI = -1.0 to -0.59, figure 3). In contrast, adding the competitor

Spirostomum only had a negative effect on the mean abundance of Dexistoma (b= -

0.43, 95% CI = -0.67 to -0.19, figure 3).

321
Figure 3: Density of focal species as a function of community composition. Each panel
shows one of the protist species. Whether the predator was present is shown on the x-axis.
The three competitive communities are shown in color (CD = *Colpidium* & *Dexiostoma*, CDP
= *Colpidium*, *Dexiostoma* & *Paramecium*, CDS = *Colpidium*, *Dexiostoma* & *Spirostomum*).
The solid shapes show the mean and the standard error of the mean (calculated across
duration of the experiment), and the transparent shapes show the actual data points.

The addition of the predator had a negative effect on the mean abundance of

all competing protist species (figure 3). In communities composed of just the focal

pair the presence of the predator resulted in a similar reduction in mean abundance

for both *Dexiostoma*: (b= -1.8, 95% CI = -2.0 to -1.5) and *Colpidium* (b= -1.9, 95% CI

= -2.1 to -1.6, figure 3). In communities with a third competitor the predator had a

greater negative effect on the mean abundance of *Spirostomum* (b= -1.7, 95% CI = -

1.9 to -1.5) than *Paramecium* (b= -1.5, 95% CI = -1.7 to -1.2, figure 3). The addition

of *Spirostomum* had no effect on the mean abundances of *Dexiostoma*, *Colpidium*

and the predator compared to the focal pair cultured with just the predator.

Intriguingly, the presence of both *Paramecium* and the predator had a positive effect

on the density of both *Colpidium* (b= 0.83, 95% CI = 0.43 to 1.2) and *Dexiostoma*

(b= 0.96, 95% CI = 0.62 to 1.3, figure 3) in comparison to when the focal species

were exposed to the predator in isolation. This translated into higher density for the

predator *Spathidium* when *Paramecium*, *Colpidium* and *Dexiostoma* were present

(b= 0.66, 95% CI = 0.48 to 0.84, figure 3).

Higher-order interactions

Model selection revealed that higher order interactions influenced the *per capita*

population growth rates of *Dexiostoma* and *Colpidium* in some communities (Table

1, Figures S1-7). For *Colpidium*, the interactive LV model only including intraspecific

HOIs (dAIC = 0.01 but fewer parameters) was best supported in the two-species

community of *Colpidium* and *Dexiostoma*, indicating that *Dexiostoma* modified the

effect of *Colpidium* on itself. In culture with a third species, the best supported model

differed: when *Colpidium* was present with *Dexiostoma* and *Paramecium*, the

additive Ricker model was the most parsimonious (dAIC = 1.04 but fewer

parameters). In contrast, when *Colpidium* was present with *Dexiostoma* and

Spirostomum, the interactive Ricker model was best supported, suggesting

interspecific HOIs. For the *per capita* growth rate of *Dexiostoma*, the interactive LV

model only including intraspecific HOIs (dAIC = 0.91 but fewer parameters) was

most supported when only *Colpidium* was present, suggesting that *Colpidium*

modified the effect of *Dexiostoma* on itself. When a third species was present, in

both cases the additive Ricker model was best supported, suggesting that neither

intra- nor interspecific HOIs were operating. Since all the best supported models

were either the interactive LV or Ricker models, all intra- and interspecific

interactions were nonlinear.

Table 1: Model selection table showing the AIC values for of the per capita growth rate of

the two focal species across the three community compositions. Additive and interactive

versions of the Lotka-Volterra (LV) and Ricker models were fitted to the data. K is the

number of estimated parameters, dAIC the difference between the model with the lowest

AIC and the AIC of the specific model.

species	community	model	K	AIC	dAIC
Colp	**CD**	**interactive LV (full HOI)**	**7**	**-49.70**	**0.00**
Colp	**CD**	**interactive LV (intra HOI)**	**6**	**-49.69**	**0.01**
Colp	CD	additive Ricker	4	-42.25	7.45
Colp	CD	additive LV	4	-42.05	7.65
Colp	CD	interactive Ricker	5	-40.26	9.44
Colp	CD	interactive LV (inter HOI)	5	-40.06	9.64
Colp	**CDP**	**interactive LV (intra HOI)**	**8**	**61.67**	**0.00**
Colp	**CDP**	**additive Ricker**	**5**	**62.71**	**1.04**
Colp	CDP	interactive LV (full HOI)	11	65.71	4.04
Colp	CDP	interactive Ricker	8	66.57	4.90
Colp	CDP	additive LV	5	74.84	13.17
Colp	CDP	interactive LV (inter HOI)	8	76.32	14.65
Colp	**CDS**	**interactive Ricker (inter HOI)**	**8**	**-77.52**	**0.00**
Colp	**CDS**	**interactive LV (inter HOI)**	**8**	**-75.67**	**1.85**
Colp	CDS	interactive LV (full HOI)	11	-71.82	5.70
Colp	CDS	additive Ricker	5	-67.03	10.49
Colp	CDS	additive LV	5	-66.27	11.24
Colp	CDS	interactive LV (intra HOI)	8	-63.06	14.46
Dexio	**CD**	**interactive LV (full HOI)**	**7**	**-88.23**	**0.00**
Dexio	**CD**	**interactive LV (intra HOI)**	**6**	**-87.32**	**0.91**
Dexio	CD	additive Ricker	4	-80.98	7.25
Dexio	CD	interactive Ricker	5	-80.25	7.98
Dexio	CD	additive LV	4	-75.52	12.71
Dexio	CD	interactive LV (inter HOI)	5	-74.51	13.72
Dexio	**CDP**	**additive Ricker**	**5**	**3.24**	**0.00**
Dexio	CDP	additive LV	5	5.32	2.08
Dexio	CDP	interactive LV (intra HOI)	8	6.22	2.98
Dexio	CDP	interactive Ricker	8	7.22	3.98
Dexio	CDP	interactive LV (inter HOI)	8	8.54	5.30
Dexio	CDP	interactive LV (full HOI)	11	9.03	5.79
Dexio	**CDS**	**additive Ricker**	**5**	**-60.69**	**0.00**
Dexio	CDS	interactive LV (intra HOI)	8	-57.79	2.90
Dexio	CDS	additive LV	5	-57.01	3.67
Dexio	CDS	interactive Ricker	8	-55.90	4.79
Dexio	CDS	interactive LV (full HOI)	11	-54.22	6.47
Dexio	CDS	interactive LV (inter HOI)	8	-52.40	8.29

Population persistence

In the absence of predators, *Colpidium* survival was contingent on the
community composition (log rank test p= 0.0066, figure 4A). Without predators,
adding *Spirostomum* increased *Colpidium* survival (pairwise log-rank test p = 0.02,
figure 4A), whereas adding *Paramecium* did not. In contrast, *Dexiostoma* persisted in
the absence of predators regardless of the community composition (log rank test p =
0.37, figure 4C). In the presence of predators, *Colpidium* survival did not depend on
the identity of the competitor species (figure 4B). In contrast for *Dexiostoma*, the
community composition did impact survival (log rank test p = 0.024), since adding
Paramecium to the community increased *Dexiostoma* survival (figure 4D).
Spathidium survival depended on the prey community composition (log rank test p =
0.011), with *Paramecium* increasing the survival of *Spathidium* (pairwise log rank
test p = 0.018, figure 4E).
387
388
389

Figure 4: Survival curves (lines) of *Colpidium*, *Dexiostoma* and *Spathidium* as a function of community composition (CD = *Colpidium* & *Dexiostoma*, CDP = *Colpidium*, *Dexiostoma* & *Paramecium*, CDS = *Colpidium*, *Dexiostoma* & *Spirostomum*, CDPd = CD + predator, CDPPd = CDP + predator, CDSPd = CDS + predator). 95% confidence intervals are shown by shaded boxes. While for *Colpidium*, survival depends on additional species in the competitive communities (A) but not the communities with predation (B), the opposite is observed for Dexistoma (C-D). *Spathidium* survival depended on the prey community composition (E).

Discussion

Evidence for higher order interactions

Comparing additive and interactive LV models revealed the significant effect of intraspecific HOIs on population growth rate for both *Colpidium* and *Dexiostoma* in the two species communities. Intraspecific HOIs can occur through an " information cascade" in social groups of species, where individuals base their decisions on the actions of others (Bikhchandani et al., 1992; Potts, 1984). Studies have shown that the behavior of neighboring fish influences the direction and speed of the school (Ioannou et al., 2011). If a fish senses a signal of danger and turns, it creates a pressure wave in the water, and other fish respond to this pressure wave by turning as well (Ioannou et al., 2011). Similar examples can also be found in flocks of birds and swarms of bees (Lissaman & Shollenberger, 1970; Parrish & Edelstein-Keshet, 1999). In all cases, an external signal may trigger behavioral changes of most members of the group, thereby altering the interactions among individuals of the same species. Although ciliates do not have elaborate sensory systems such as birds and fish, they are capable of sensing changes in local population density and adjusting their movement strategies accordingly (Fronhofer et al., 2015; Pennekamp et al., 2014). Since movement and foraging are often closely linked (Van Dyck & Baguette, 2005), it is possible that movement-related crowding effects could lead to the emergence of intraspecific HOIs.

Interspecific HOIs were only found on the growth rate of *Colpidium* when *Dexiostoma* and *Spirostomum* were present. *Spirostomum* most likely changed the effect of *Dexiostoma* on *Colpidium*, because both *Dexiostoma* and *Colpidium* strongly competed for the same resources, while the direct effect of *Spirostomum* on *Colpidium* was more limited. It is possible that *Dexiostoma* and *Spirostomum* consume similar sized bacteria, while *Colpidium* uses a different size class. This would also explain why we did not observe an interspecific effect of *Spirostomum* on *Dexiostoma* when *Colpidium* was the focal species. Interestingly, there was no HOI of *Paramecium* on either *Dexiostoma* or *Colpidium*, although *Paramecium* was a stronger competitor than *Spirostomum*. Previous work has found strong evidence that interspecific HOIs can affect coexistence and dynamics of aquatic microbial communities (Mickalide & Kuehn, 2019). Mickalide & Kuehn (2019) observed that *Escherichia coli* can invade cultures of the alga *Chlamydomonas reinhardtii* or the

ciliate *Tetrahymena thermophila* but fails to invade a community where both species are present. The invasion resistance of the algae-ciliate community arises from an HOI caused by the algal inhibition of bacterial aggregation, which leaves bacteria vulnerable to predation.

The prevalence of intra- versus interspecific HOIs is noteworthy since most studies have so far focused on interspecific HOIs. Work on plant communities has shown that the prevalence of intra- or interspecific HOIs can vary across focal species and that neither type is generally more or less important (Mayfield & Stouffer, 2017). Since coexistence will be determined by the relative strengths of intra- and interspecific competition (Gibbs et al., 2022; Singh & Baruah, 2021), changes in intraspecific interaction strength alone can influence the persistence of communities. We therefore recommend that empirical studies investigating HOIs embrace a definition of HOIs that includes both intra- and interspecific effects (e.g., Mayfield & Stouffer, 2017).

The nature of the detected HOIs

Some authors have argued that HOI are not ecological processes in their own right but are instead emergent properties of phenomenological models (Letten & Stouffer, 2019). For example, a Lotka- Volterra competition model imposes a linear relationship between the density of one species and the growth of the other. However, if this relationship is not linear in nature, the addition of HOI terms may improve the fit, but the model would not be accurately describing the species interactions. Properly accounting for nonlinear density dependence is an important prerequisite to estimate interaction strength (Hart et al., 2018) and avoid erroneous conclusions about the presence of HOIs when nonlinearity is present (Kleinhesselink et al., 2022; Letten & Stouffer, 2019). To investigate the potential of nonlinear density dependence we fit models that included both linear and non-linear competition coefficients and well as intra- and interspecific HOI terms. In all communities the presence of nonlinear density-dependence was indicated by either the interactive LV or the Ricker model having the lowest AIC values. The observed HOIs suggests that there are processes resulting in non-linearity which the simple additive LV and the nonlinear Ricker model are missing. Therefore, we believe the HOIs detected are not an artifact of nonlinear density dependence, but true emergent behavioural

responses. A mechanistic model that allows for such non-linearity may provide

further insights into the ecological processes driving community dynamics.

Do HOIs affect abundance and community stability in competitive

communities?

Persistence was different for the two focal prey species. *Dexiostoma* showed

high persistence regardless of the additional species present, while *Colpidium*

became more stable in the presence of a third species. The increased persistence of

Colpidium when cultured with *Paramecium* was not driven by an HOI, since the

additive Ricker model was best supported by the data in the CDP community. But

the highest persistence of *Colpidium* was detected in the presence of *Dexiostoma*

and *Spirostomum*, where the interactive model was best supported. This pattern is

consistent with a stabilizing effect due to interspecific HOIs.

Dexiostoma showed a similar persistence across all combinations of

competitors, despite a non-significant trend to lower persistence when *Paramecium*

was present. *Dexiostoma* also showed variation in the presence of HOIs, but the

pattern is not consistent with a stabilizing role of HOIs: when present with only

Colpidium, intraspecific HOIs were observed and *Dexiostoma* persisted till the end of

the experiment. When *Paramecium* or *Spirostomum* was added, HOIs were no

longer detected but neither did the persistence change. This suggests that changes

in HOIs do not always result in changes in the persistence of focal species. Direct

interactions among species may be much more important.

The predator itself showed the highest persistence when feeding on

communities that contained *Colpidium*, *Dexiostoma* and *Paramecium*. When feeding

on the focal pair without competitors, it showed intermediate persistence times, while

communities of the focal pair with *Spirostomum* persisted least well. Hammill and

colleagues (2015) found a similar effect for non-trophically interacting species in a

food web: if species that do not interact trophically are present, prey persist for

longer in diverse food webs. Since we could not quantify the interaction strength of

the predator on *Colpidium* and *Dexiostoma* in the presence of *Paramecium* or

Spirostomum, we cannot say whether the change in predator persistence was due to

HOIs or simply the direct and indirect effects among the members of the food web.

Quantifying the role of trophic interaction modifications with functional responses is

an exciting area for future research (Terry et al., 2017).

Patterns of population abundance

Overall, species' abundance and coexistence was influenced by community composition. *Colpidium* and *Dexiostoma* grew to lower densities when cultured with *Paramecium*, a sign of competition for shared resources. *Colpidium* abundance was not affected by the presence of *Spirostomum*. However, the mean abundance of *Dexiostoma* was lower in the presence of *Spirostomum*, suggesting competition between the two species, but to a lesser degree than with *Paramecium*. These patterns are in line with known phylogenetic relationships between these species, where *Colpidium* is most closely related to *Dexiostoma*, then to *Paramecium* and finally to *Spirostomum* (Violle et al., 2011). These relationships have been found to predict competitive exclusion due to the similarity in mouth size between species, which in turn defines the feeding niche of species (Violle et al., 2011).

Predation lowered the abundances of the two focal prey species, but this effect could be partially mitigated in the presence of additional species. For instance, *Colpidium* benefitted from the presence of *Paramecium* when predators were present, having higher average abundance than would be expected if effects of both predation and competition were additive. This might be explained by interference of *Paramecium* with the predator, indirectly leading to a decrease in interaction strength between the prey and the predator. Ciliate predators show a preferred predator:prey size ratio of 8:1 (Hansen et al., 1994), explaining why some species may interfere with predator foraging. *Dexiostoma* similarly benefited from *Paramecium* when the predator was present, reaching on higher average abundance than when *Colpidium* and *Spirostomum* were present.

Surprisingly, both *Paramecium* and *Spirostomum* densities decreased in the presence of the predator, indicating that the predator had a negative effect on the two largest protist species. Furthermore, the abundance of the predator was greater when *Paramecium* was present compared to when it was cultured with only the focal species pair, or the focal pair and *Spirostomum*. It thus appears that *Paramecium* was consumed in addition to the two focal prey, and that while *Spirostomum* was attacked, it was potentially too large to be consumed. This observation could be explained by the mode of prey capture employed by the predator *Spathidium*. The cell mouth (cytostome) of *Spathidium* is furnished with a rodlike tip (toxicyst) to paralyze or kill other microorganisms for easier consumption (Fyda et al., 2005). But

while the toxicyst make it possible to paralyze large prey, phagocytosis still requires that prey organisms can be engulfed.

An important limitation of our work is that we were only able to examine the interaction of a single focal pair across a few community compositions, with a limited sample size. To completely disentangle the drivers of community persistence, we would need to measure all interaction strengths between species, which was logistically impossible in our study due to the long and high quality time series. Since our study only investigated HOIs between a focal pair, it is possible that we missed additional interspecific HOIs between the focal pair and the predator.

Conclusions

The role of higher-order interactions in community dynamics remains a key issue in community ecology (Gibbs et al., 2022; Levine et al., 2017). Currently very few empirical studies measure HOIs in communities of real species and study how higher-order interactions affect community stability. Our study demonstrates that HOIs can play a role in community stability, and provides some support for the stabilizing effects of HOIs on competitive communities shown in theoretical studies (Grilli et al., 2017). Our study further provides an example of how to identify HOIs by combining experiments with time-series analyses and hence pave the way for more studies that study the occurrence and consequences of HOIs in natural communities. Our study adds to the growing body of research that shows that species interactions can deviate from the pairwise expectations when additional species are present, and new theory needs to explore the consequences for community coexistence and stability.

Acknowledgements

Chenyu Shen, Kimberley Lemmen and Frank Pennekamp received financial support from the Swiss National Science Foundation (grant 310030_197811 awarded to FP). We thank Uriah Daugaard for inspiring the ciliate illustrations in figure 1.

Data and code accessibility

Data and code are available from the following github repository: [DOI: 10.5281/zenodo.7896204](#).

Competing Interests Statement

None.

References

Abrams, P. A. (1983). Arguments in Favor of Higher Order Interactions. *The American Naturalist*, *121*(6), 887–891. https://doi.org/10.1086/284111

Adler, P. B., Smull, D., Beard, K. H., Choi, R. T., Furniss, T., Kulmatiski, A., Meiners, J. M., Tredennick, A. T., & Veblen, K. E. (2018). Competition and coexistence in plant communities: Intraspecific competition is stronger than interspecific competition. *Ecology Letters*, *21*(9), 1319–1329. https://doi.org/10.1111/ele.13098

Altermatt, F., Fronhofer, E. A., Garnier, A., Giometto, A., Hammes, F., Klecka, J., Legrand, D., Mächler, E., Massie, T. M., Pennekamp, F., Plebani, M., Pontarp, M., Schtickzelle, N., Thuillier, V., & Petchey, O. L. (2015). Big answers from small worlds: A user's guide for protist microcosms as a model system in ecology and evolution. *Methods in Ecology and Evolution*, *6*(2), 218–231. https://doi.org/10.1111/2041-210X.12312

Bairey, E., Kelsic, E. D., & Kishony, R. (2016). High-order species interactions shape ecosystem diversity. *Nature Communications*, *7*(1), Article 1. https://doi.org/10.1038/ncomms12285

Barabás, G., J. Michalska-Smith, M., & Allesina, S. (2016). The Effect of Intra- and Interspecific Competition on Coexistence in Multispecies Communities. *The American Naturalist*, *188*(1), E1–E12. https://doi.org/10.1086/686901

Bick, H., & World Health Organization. (1972). *Ciliated protozoa: An illustrated guide to the species used as biological indicators in freshwater biology*. World Health Organization. https://apps.who.int/iris/handle/10665/40066

Bikhchandani, S., Hirshleifer, D., & Welch, I. (1992). A Theory of Fads, Fashion, Custom,

and Cultural Change as Informational Cascades. *Journal of Political Economy*,

100(5), 992–1026. https://doi.org/10.1086/261849

Billick, I., & Case, T. J. (1994). Higher Order Interactions in Ecological Communities: What

Are They and How Can They be Detected? *Ecology*, *75*(6), 1530–1543.

https://doi.org/10.2307/1939614

Carrara, F., Giometto, A., Seymour, M., Rinaldo, A., & Altermatt, F. (2015). Inferring species

interactions in ecological communities: A comparison of methods at different levels of

complexity. *Methods in Ecology and Evolution*, *6*(8), 895–906.

https://doi.org/10.1111/2041-210X.12363

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of*

Ecology and Systematics, *31*(1), 343–366.

https://doi.org/10.1146/annurev.ecolsys.31.1.343

Clark, J. S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest

Trees. *Science*, *327*(5969), 1129–1132. https://doi.org/10.1126/science.1183506

Daugaard, U., Petchey, O. L., & Pennekamp, F. (2019). Warming can destabilize predator–

prey interactions by shifting the functional response from Type III to Type II. *Journal*

of Animal Ecology, *88*(10), 1575–1586. https://doi.org/10.1111/1365-2656.13053

Foissner, W., & Berger, H. (1996). A user-friendly guide to the ciliates (Protozoa, Ciliophora)

commonly used by hydrobiologists as bioindicators in rivers, lakes, and waste

waters, with notes on their ecology. *Freshwater Biology*, *35*(2), 375–482.

https://doi.org/10.1111/j.1365-2427.1996.tb01775.x

Fronhofer, E. A., Kropf, T., & Altermatt, F. (2015). Density-dependent movement and the

consequences of the Allee effect in the model organism Tetrahymena. *Journal of*

Animal Ecology, *84*(3), 712–722. https://doi.org/10.1111/1365-2656.12315

Fyda, J., Warren, A., & Wolinńska, J. (2005). An investigation of predator-induced defence

responses in ciliated protozoa. *Journal of Natural History*, *39*(18), 1431–1442.

https://doi.org/10.1080/00222930400004396

Gibbs, T., Levin, S. A., & Levine, J. M. (2022). Coexistence in diverse communities with

higher-order interactions. *Proceedings of the National Academy of Sciences*,

119(43), e2205063119. https://doi.org/10.1073/pnas.2205063119

Grilli, J., Barabás, G., Michalska-Smith, M. J., & Allesina, S. (2017). Higher-order

interactions stabilize dynamics in competitive network models. *Nature*, *548*(7666),

210–213. https://doi.org/10.1038/nature23273

Hammill, E., Kratina, P., Vos, M., Petchey, O. L., & Anholt, B. R. (2015). Food web

persistence is enhanced by non-trophic interactions. *Oecologia*, *178*(2), 549–556.

https://doi.org/10.1007/s00442-015-3244-3

Hansen, B., Bjornsen, P. K., & Hansen, P. J. (1994). The size ratio between planktonic

predators and their prey. *Limnology and Oceanography*, *39*(2), 395–403.

https://doi.org/10.4319/lo.1994.39.2.0395

Hart, S. P., Freckleton, R. P., & Levine, J. M. (2018). How to quantify competitive ability.

Journal of Ecology, *106*(5), 1902–1909. https://doi.org/10.1111/1365-2745.12954

HilleRisLambers, J., Adler, P. B., Harpole, W. S., Levine, J. M., & Mayfield, M. M. (2012).

Rethinking Community Assembly through the Lens of Coexistence Theory. *Annual*

Review of Ecology, Evolution, and Systematics, *43*(1), 227–248.

https://doi.org/10.1146/annurev-ecolsys-110411-160411

Ioannou, C. C., Couzin, I. D., James, R., Croft, D. P., & Krause, J. (2011). Social

Organisation and Information Transfer in Schooling Fish. In *Fish Cognition and*

Behavior (pp. 217–239). John Wiley & Sons, Ltd.

https://doi.org/10.1002/9781444342536.ch10

Kassambara, A., Kosinski, M., Biecek, P., & Fabian, S. (2021). *survminer: Drawing Survival*

Curves using "ggplot2" (0.4.9). https://cran.r-

project.org/web/packages/survminer/index.html

Kleinhesselink, A. R., Kraft, N. J. B., Pacala, S. W., & Levine, J. M. (2022). Detecting and

interpreting higher-order interactions in ecological communities. *Ecology Letters*,

25(7), 1604–1617. https://doi.org/10.1111/ele.14022

Letten, A. D., & Stouffer, D. B. (2019). The mechanistic basis for higher-order interactions

and non-additivity in competitive communities. *Ecology Letters*, *22*(3), 423–436.

https://doi.org/10.1111/ele.13211

Levine, J. M., Bascompte, J., Adler, P. B., & Allesina, S. (2017). Beyond pairwise

mechanisms of species coexistence in complex communities. *Nature*, *546*(7656),

56–64. https://doi.org/10.1038/nature22898

Lissaman, P. B. S., & Shollenberger, C. A. (1970). Formation Flight of Birds. *Science*,

168(3934), 1003–1005. https://doi.org/10.1126/science.168.3934.1003

May, R. M. (1972). Will a Large Complex System be Stable? *Nature*, *238*(5364), 413–414.

https://doi.org/10.1038/238413a0

Mayfield, M. M., & Stouffer, D. B. (2017). Higher-order interactions capture unexplained

complexity in diverse communities. *Nature Ecology & Evolution*, *1*(3), 1–7.

https://doi.org/10.1038/s41559-016-0062

Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A., Leisch, F., C++-code), C.-C. C.

(libsvm, & C++-code), C.-C. L. (libsvm. (2022). *e1071: Misc Functions of the*

Department of Statistics, Probability Theory Group (Formerly: E1071), TU Wien (1.7-

11). https://CRAN.R-project.org/package=e1071

Mickalide, H., & Kuehn, S. (2019). Higher-Order Interaction between Species Inhibits

Bacterial Invasion of a Phototroph-Predator Microbial Community. *Cell Systems*,

9(6), 521-533.e10. https://doi.org/10.1016/j.cels.2019.11.004

Mühlbauer, L. K., Schulze, M., Harpole, W. S., & Clark, A. T. (2020). gauseR: Simple

methods for fitting Lotka-Volterra models describing Gause's "Struggle for

Existence." *Ecology and Evolution*, *10*(23), 13275–13283.

https://doi.org/10.1002/ece3.6926

Parrish, J. K., & Edelstein-Keshet, L. (1999). Complexity, Pattern, and Evolutionary Trade-

Offs in Animal Aggregation. *Science*, *284*(5411), 99–101.

https://doi.org/10.1126/science.284.5411.99

Pennekamp, F., Griffiths, J. I., Fronhofer, E. A., Garnier, A., Seymour, M., Altermatt, F., &

Petchey, O. L. (2017). Dynamic species classification of microorganisms across time,

abiotic and biotic environments—A sliding window approach. *PLOS ONE*, *12*(5),

e0176682. https://doi.org/10.1371/journal.pone.0176682

Pennekamp, F., Mitchell, K. A., Chaine, A., & Schtickzelle, N. (2014). Dispersal propensity in

Tetrahymena thermophila ciliates—A reaction norm perspective. *Evolution;*

International Journal of Organic Evolution, *68*(8), 2319–2330.

https://doi.org/10.1111/evo.12428

Pennekamp, F., Pontarp, M., Tabi, A., Altermatt, F., Alther, R., Choffat, Y., Fronhofer, E. A.,

Ganesanandamoorthy, P., Garnier, A., Griffiths, J. I., Greene, S., Horgan, K., Massie,

T. M., Mächler, E., Palamara, G. M., Seymour, M., & Petchey, O. L. (2018).

Biodiversity increases and decreases ecosystem stability. *Nature*, *563*(7729), 109–

112. https://doi.org/10.1038/s41586-018-0627-8

Pennekamp, F., Schtickzelle, N., & Petchey, O. L. (2015). BEMOVI, software for extracting

behavior and morphology from videos, illustrated with analyses of microbes. *Ecology*

and Evolution, *5*(13), 2584–2595. https://doi.org/10.1002/ece3.1529

Pfister, C. A. (1995). Estimating Competition Coefficients from Census Data: A Test with

Field Manipulations of Tidepool Fishes. *The American Naturalist*, *146*(2), 271–291.

https://doi.org/10.1086/285798

Potts, W. K. (1984). The chorus-line hypothesis of manoeuvre coordination in avian flocks.

Nature, *309*(5966), Article 5966. https://doi.org/10.1038/309344a0

R Core Team. (2022). *R: A language and environment for statistical computing*.

https://www.r-project.org/

Singh, P., & Baruah, G. (2021). Higher order interactions and species coexistence.

Theoretical Ecology, *14*(1), 71–83. https://doi.org/10.1007/s12080-020-00481-8

Tabi, A., Petchey, O. L., & Pennekamp, F. (2019). Warming reduces the effects of

enrichment on stability and functioning across levels of organisation in an aquatic

microbial ecosystem. *Ecology Letters*, *22*(7), 1061–1071.

https://doi.org/10.1111/ele.13262

Terry, J. C. D., Morris, R. J., & Bonsall, M. B. (2017). Trophic interaction modifications: An

empirical and theoretical framework. *Ecology Letters*, *20*(10), 1219–1230.

https://doi.org/10.1111/ele.12824

Van Dyck, H., & Baguette, M. (2005). Dispersal behaviour in fragmented landscapes:

Routine or special movements? *Basic and Applied Ecology*, *6*(6), 535–545.

https://doi.org/10.1016/j.baae.2005.03.005

Violle, C., Nemergut, D. R., Pu, Z., & Jiang, L. (2011). Phylogenetic limiting similarity and

competitive exclusion. *Ecology Letters*, *14*(8), 782–787.

https://doi.org/10.1111/j.1461-0248.2011.01644.x

Werner, E. E., & Peacor, S. D. (2003). A Review of Trait-Mediated Indirect Interactions in

Ecological Communities. *Ecology*, *84*(5), 1083–1100. https://doi.org/10.1890/0012-

9658(2003)084[1083:AROTII]2.0.CO;2

Woodruff, L. L., & Spencer, H. (1921). The food reactions of the infusorian Spathidium

spathula. *Proceedings of the Society for Experimental Biology and Medicine*, *18*(6),

183–184. https://doi.org/10.3181/00379727-18-90

Wootton, J. T. (1993). Indirect Effects and Habitat Use in an Intertidal Community:

Interaction Chains and Interaction Modifications. *The American Naturalist*, *141*(1),

71–89.

Wootton, J. T. (1994). Predicting Direct and Indirect Effects: An Integrated Approach Using

Experiments and Path Analysis. *Ecology*, *75*(1), 151–165.

https://doi.org/10.2307/1939391

Worthen, W. B., & Moore, J. L. (1991). Higher-Order Interactions and Indirect Effects: A

Resolution Using Laboratory Drosophila Communities. *The American Naturalist*,

138(5), 1092–1104.

Supplementary material

Species classification

For species classification, we used the videos collected from the first sampling day of every week and randomly chose one video of each replicate for the six community compositions. By observing the morphology and movement patterns of the ciliates in the videos, we labelled 350 to 400 trajectories of each species as training data. We use a similar number of labelled trajectories of each species, which avoids low classification performance for rare classes (e.g. low abundance species) caused by imbalanced numbers of observations (Sommer & Gerlich, 2013). Although the trajectory filtering function from BEMOVI removed most of the background noise, the movements of some tiny, suspended impurities may remain as spurious trajectories within the training data. In order to prevent these trajectories being mistakenly classified as other species, we selected 100 videos that we visually checked to contain no ciliates and labelled all detected trajectories as "noise". Trajectories of class "noise" were then excluded from all subsequent analyses.

Twenty features extracted from an established classification pipeline (Pennekamp et al., 2017) were selected to train the classification algorithm to distinguish among classes based on information about body size and movement patterns (Table S1). As the phenotypes of ciliates may change over time (Pennekamp et al., 2017), the week number was added as the 21st feature into the classification algorithm to enhance the accuracy of the prediction.

As the species included in each treatment were different, we built six customized models for classification that only containing the known species of each community. Prior to the experiment we compared the random forest (RF) and Support Vector Machine (SVM) as our classifiers for their computation efficiency and reliability of results (Fernández-Delgado et al., 2014). We used the randomForest package (Liaw & Wiener, 2002) and e1071 packages that contains the *svm* function (Meyer et al., 2014.). We compared the performances of two classifiers by calculating the classification errors for all training data (Table S2). Since SVM gave

757 us overall lower classification errors, we applied it to our six customized models and

758 generated time series of change density change of species for each community.

759 **Species abundances**

760 **Table S1:** Effects of community composition on the log density of focal species. CI =

761 confidence interval.

	Colpidium			Dexiostoma			Paramecium			Spirostomum			Spathidium		
	Beta	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-value	Beta	95% CI	p-value
Intercept	5.2	5.0, 5.4	<0.001	5.7	5.5, 5.9	<0.001	4.9	4.7, 5.1	<0.001	3.8	3.6, 3.9	<0.001	2.5	2.4, 2.6	<0.001
Predation	-1.9	-2.1, -1.6	<0.001	-1.8	-2.0, -1.5	<0.001	-1.5	-1.7, -1.2	<0.001	-1.7	-1.9, -1.5	<0.001	-	-	-
Paramecium present	-0.60	-0.89, -0.32	<0.001	-0.81	-1.1, -0.57	<0.001	-	-	-	-	-	-	0.66	0.48, 0.84	<0.001
Spirostomum present	0.06	-0.22, 0.34	0.7	-0.43	-0.67, -0.19	0.001	-	-	-	-	-	-	-0.17	-0.36, 0.01	0.062
Predation and *Paramecium* present	0.83	0.43, 1.2	<0.001	0.96	0.62, 1.3	<0.001	-	-	-	-	-	-	-	-	-
Predation and *Spirostomum* present	-0.04	-0.44, 0.36	0.8	0.24	-0.10, 0.58	0.2	-	-	-	-	-	-	-	-	-

763 **Model fitting to estimate interactions**

764 The following section shows the coefficient estimates for the additive and interactive

765 Lotka-Volterra and Ricker competition models.

Figure S1: *Dexiostoma* growth rate in CD community

769
770 Figure S2: *Dexiostoma* growth rate in CDP community

Figure S3: *Dexiostoma* growth rate in CDS community

Figure S4: *Colpidium* growth rate in CD community

776   Figure S5: *Colpidium* growth rate in CDP community

778   Figure S6: *Colpidium* growth rate in CDS community

782
783   Figure S7: Coefficient plots for the best supported models for *Dexiostoma* (upper panel) and
784   *Colpidium* (lower panel) across community compositions. Each panel shows the intra- or
785   interspecific interaction terms as well as the interaction coefficients.
786