An anterior forebrain pathway in parrots is necessary for individual signatures of learned vocalizations

Zhilei Zhao[1], Han Kheng Teoh[2], Julie Carpenter[1], Frieda Nemon[1], Brian Kardon[1], Itai Cohen[2], Jesse H. Goldberg[1,*]

[1]*Department of Neurobiology and Behavior, Cornell University, Ithaca, NY 14853, USA*

[2]*Department of Physics, Cornell University, Ithaca, NY 14853, USA.*

*Corresponding author

Abstract

Parrots have enormous vocal imitation capacities and produce individually unique vocal signatures. Like songbirds, parrots have a nucleated neural song system, but because the two sister groups diverged over 50 million years ago it remains unknown if the song systems operate according to similar principles. Here we established a neuroscience-compatible call-and-response behavioral paradigm to elicit learned contact calls in budgerigars (*Melopsittacus undulatus*). Using variational autoencoder-based machine learning methods, we show that contact calls within affiliated groups converge but that individuals maintain unique acoustic features, or vocal signatures, even after call convergence. Next, we transiently and reversibly inactivated the outputs of the anterior forebrain pathway (AFP) as animals engaged in call-and-response interactions. Contact call pitch modulation, stereotypy and acoustic structure were dramatically degraded, resulting in the loss of individually unique features of the vocal signature. These results are in stark contrast to results from songbirds, where AFP inactivations with identical methods have the opposite result of increasing the stereotypy of learned vocalizations and retaining acoustic structure. Together, our results show that the anterior pathways of parrots and songbirds have different functional organization and that parrot frontal cortical circuits are necessary for individually unique features of vocal communication.

Introduction

The ability to learn vocalizations and use them for social communication was a key milestone in the evolution of human language [1]. This trait, rare among non-human animals, is observed in three clades of birds: oscine songbirds, parrots, and hummingbirds [2–4]. Interestingly, all of these clades evolved nucleated forebrain circuits, or 'song systems' with similar-appearing basal ganglia thalamocortical pathways. But because these clades diverged over 50 million years ago and have non-vocal learners in their lineages, it remains unknown if the principles of song system function are conserved [1,4].

The neural mechanisms of song production and learning have been extensively studied in three songbird species: zebra finches, bengalese finches, and canaries [5–8]. Conceptually similar results across these species support a conserved logic to song system architecture where anatomy maps

clearly to function (Figure 1A). The output of the forebrain song system, RA (robust nucleus of
the arcopallium), is analogous to mammalian primary motor cortex and has only two main inputs:
HVC (used as a proper name) and lateral magnocellular nucleus of the anterior nidopallium
(LMAN) [9,10]. HVC is part of the posterior motor pathway that is necessary for adult song production
but not for vocal babbling in juveniles. HVC lesions or inactivations degrade acoustic structure in
adult song but have little effect on normal vocal babbling in juveniles [8,11–13]. During singing, HVC
neurons in adult birds exhibit stereotyped, temporally precise activity that drives correspondingly
stereotyped syllable sequences in the adult song [10,14,15]. These basic observations of posterior
pathway function were additionally supported in more exotic dueting songbirds, including white-
browed sparrow weavers and plain-tailed wrens, supporting the idea of conserved posterior song
system function across all songbird species studied [16,17].

In contrast to HVC, LMAN is the output of the songbird anterior forebrain pathway (AFP)—a basal
ganglia (BG) thalamocortical loop—necessary for song learning but not for adult song production.
Lesion or inactivation LMAN or its thalamic input [11,12], has little effect on adult song but a
devastating effect on vocal babbling and learning [18–21]. With LMAN inactivated, the normal
variability of juvenile vocal babbling is instantly lost and the song becomes abnormally
stereotyped. Consistent with its role in driving vocal variation, LMAN neurons exhibit variable,
bursty activity during singing that drives the correspondingly variable song components, including
the vocal babbling of juveniles [13,20,21].

The discovery that the AFP functions as a variability-generator necessary for learning is one of
the most important contributions of the songbird field to neuroscience. It shows that variability
characteristic of yet-to-be learned behaviors is not the *de facto* output of an immature motor
system [22], but is instead actively generated by neural circuits for the purpose of learning. Though
a conceptually similar 'variability-generating' circuit has yet to be identified in mammals, frontal
BG-cortical loops may generate variability since catecholamines reduce variability in both humans
and in songbirds [23,24].

To test if frontal cortical variability-generating functions generalize to distantly related vocal
learners or if they are an idiosyncratic feature of songbirds, we studied vocal production in the
budgerigars (*Melopsittacus undulatus*), a member of the parrot family. Anatomical studies show
that the parrot song system resembles the songbird one (Figure 1B), and as a result has been
proposed to have a similar functional organization [25–31]. First, the central nucleus of the anterior
arcopallium (AAC) may be analogous to RA because it is similarly positioned in the posterior
forebrain, sends axons directly to motor neurons in the brainstem, and receives inputs from both
a posterior and frontal pathway. Lesioning AAC or its input from the posteriorly-located nucleus
NLC (central nucleus of the lateral nidopallium) degrades the acoustic structure of learned
vocalizations [28,32,33], supporting the possibility that NLC may be analogous to HVC [25,31]. The
function of the parrot AFP, on the other hand, remains largely unexplored. In contrast to the
songbird, the parrot has two nuclei that constitute AFP outputs: MO (oval nucleus of the
mesopallium) and NAO (oval nucleus of the anterior nidopallium). Like LMAN, they are frontal
cortical nuclei that project to both the basal ganglia and AAC (the RA analogue) [25–31]. Yet to our
knowledge only one study conducted lesions of MO and found that frequency modulation was
gradually lost over a course of several days [28], but only a small number of calls were examined

from one bird and findings were based on visual inspections of calls. Thus it remains unclear if
the frontal song system has LMAN-like functions. Here we revisited the function of AFP in parrots
with reversible brain inactivations and machine learning-enabled quantification of large datasets
of budgerigar vocalizations.

Results

A behavioral paradigm to elicit thousands of learned contact calls per bird

Studying the detailed acoustic structure of vocalizations and how they may change with neural
manipulations requires analysis of large datasets, ideally thousands of calls per bird [34–36]. We set
out to establish a behavioral paradigm that would provide such a dataset even in single-housed
birds with pristine acoustic isolation. We focused on contact calls, a major type of learned
vocalizations in budgerigars, that are used to coordinate flock behavior during flight and to localize
individuals in a colony [37]. Past work showed that contact calls are learned, continuously adjusted
even in adulthood, and may contain unique signatures that allow for individual recognition [38–42].
To sample contact calls efficiently, we simulated its natural use with a trial-by-trial behavioral
paradigm. We first recorded colony noise from a social group (5 males, 3 females) and divided
the recording into two-minute segments. We then isolated individuals in a sound-proof box and
used directional microphones to record their vocalizations (Figure 2A). In a given trial, one
segment of colony noise was randomly selected and played to the bird through a speaker,
followed by a silent inter-trial interval (average 10 mins). Isolated birds responded to colony noise
with the production of contact calls so reliably that we were able to obtain hundreds to thousands
of contact calls per day with this method (average counts range: 180-1612, n=8 birds, Figure 2B).

A variational autoencoder identifies individual signatures even within shared call types

To compare contact calls between individuals, previous studies relied on visual inspection or
descriptive acoustic features [39–42]. However, acoustic features such as pitch, pitch goodness and
entropy can miss important acoustic details [36]. We therefore adopted a recently developed
method [36], variational auto-encoder (VAE), to analyze contact calls in an unsupervised manner
(Figure 2C). Briefly, vocalizations were first converted to spectrograms, then an encoder-decoder
network was trained to reconstruct the original spectrograms from low-dimensional latent
features. We next examined these latent features for downstream visualization using UMAP
(uniform manifold approximation and projection) for dimensionality reduction and with a SVM
(support vector machine) for quantitative classification of uniqueness.

Using these methods, we quantitatively confirmed and extended three previous observations on
budgerigar contact calls. First, individuals produced both individually unique contact calls as well
as shared calls that exhibited highly similar acoustic structures [39–41] (Figure 2B). Second,
quantitative analysis revealed that despite this group convergence individuals maintained unique
signatures even within shared calls [38,42]. This is visualized in the UMAP projection of the latent
features, in which the shared group call appears as a large single cluster, while calls from specific
individuals occupy distinct parts of the large cluster (Figure 2D-E). To quantify the uniqueness of
individual signatures, we trained SVM classifiers to decode individual identity from call latent
features [43]. Unsurprisingly, as we increased the dimension of principal inputs, i.e. used more

information from the spectrogram, the decoding accuracy gradually improved (Figure 2F). With
eight or more principal inputs, the SVM correctly predicted bird identity more than 90% of the time,
even though birds shared highly similar calls. We further restricted the analysis to the two major
call types (Figure 2B, 2E, marked by asterisk and plus symbols) that were shared across
individuals, and found that the SVM could still decode bird identity ~90% of the time (Figure 2G).
This accuracy was significantly higher than a previous study based on summary acoustic metrics
[42] (~40%), suggesting, unsurprisingly, that spectrograms contained extra information for
discrimination not captured by acoustic feature vectors [36].

Group vocalizations dynamically change over time

A proposed function of contact calls is to signal affiliation among birds, which, in the wild, can
change over time [37,44–46]. Call matching would be a more honest signal of affiliation if calls
dynamically changed over time, requiring ongoing social interactions [40,44]. To test if our analyses
could capture this dynamism of budgerigar contact calls, we compared contact calls of each
individual in the group produced at week 3 (Figure 2D) and week 15 (Figure 2E) following the
group assimilation. We found that the group convergence and individual signatures were not static
but exhibited dynamic drift, as shown by past qualitative description [39]. We trained a single VAE
on the group calls produced 12 weeks apart and visualized them in the same UMAP space (Figure
2D-E). We then quantified time-dependent changes in call structures by training an SVM classifier
on calls at week 3, then decode caller identity at week 15, or vice versa. Decoding accuracy
dropped significantly from ~90% to ~40% (Figure 2H).

Although these behavioral phenomena have been described before, here we provided
quantitative characterization under a unified framework. We show that living in a social group,
budgerigars need to constantly monitor contact calls of others, and modify their own calls
accordingly to preserve both convergence and individual signature. Our ability to confirm past
findings with new analytical methods also lays the foundation for studying how brain inactivations
affect call acoustic structure.

AFP-inactivated budgerigars maintain the ability to respond to stimulus calls

We next examined the role of the AFP circuits in contact call production. Since birds do not readily
vocalize in the days after surgery and permanent lesions are associated with experience-
independent circuit remodeling [47], lesions can fail to test the effect of immediate loss of AFP
contribution to vocalizations. We thus used reverse microdialysis to inactivate the output of the
AFP. Reverse microdialysis is an established method in songbirds to transiently and reversibly
inactivate brain regions over hour timescales and is particularly well suited for the study of
nucleated systems [8,48,49]. In our past work in zebra finches using identical homemade probes, we
carefully calibrated inactivated tissue volume as a function of drug concentration and replicated
the observation that LMAN inactivations cause an immediate and reversible loss of vocal
variability [8,12,13,48–50].

Based on our past calibrations, we were not confident that we could specifically target MO or NAO
separately because they are immediately neighboring structures with a shared border (Fig. 1B,
Methods). Thus, we implanted custom microdialysis probes bilaterally at the center of the
MO/NAO complex to inactivate both structures at once (Figure 3A). After birds recovered from

surgery, we infused saline (PBS) or tetrodotoxin (TTX 50 µM) into the probe on alternating days
to compare their effects on contact calls in the call-and-response paradigm. We verified our
targeting with histology and confirmed diffusion from the probes into both MO and NAO but not
the underlying basal ganglia nucleus MMSt (Figure 3B).

To sample contact calls, we used the same call-and-respond behavioral paradigm (Figure 2A),
but played individual calls instead of colony noise to better examine the temporal relationship
between stimulus and response (Figure 3C-D). To test the role of the frontal system in producing
calls with clearly distinct acoustic structures, we chose five birds unknown to one another with
clearly distinguishable individual signatures (Figure 3E-I).

With MO/NAO inactivated, all birds were still able to reliably respond to call stimuli (Figure 3C-I).
Response rate significantly decreased in three birds, but not the other two (Figure 4A). Similarly,
we didn't observe a consistent change of response latency across birds (Figure 4B). Together
these data show that the AFP is not necessary for responding to a stimulus call with appropriate
timing.

Inactivation of budgerigar AFP reduces call acoustic structure and increases call
variability
We next examined the acoustic structure of contact calls before, during, and after MO/NAO
inactivation. The pitch of normal budgerigar contact calls ranges from 2 to 4 kHz, centered near
3 kHz [37] (Figure 4C). We found that when AFP was inactivated, despite a slight decrease the
mean pitch of responded calls still fell into this range (Figure 4C), suggesting that the birds were
able to produce contact-call-like vocalizations. However, two aspects of call acoustic structure
were dramatically disrupted. First, calls produced under the TTX condition exhibited significantly
less pitch modulation compared to vehicle, and virtually all of the rapid pitch modulations were
eliminated (Figure 3E-I), as quantified by coefficient of variation (Figure 4D, PBS:TTX, 1.80±0.58)
and pitch modulation depth (Figure 4E, PBS:TTX, 1.99±0.64, Methods). Second, call durations
were affected. Normal contact calls, and those observed with vehicle in the probes, last between
0.15 to 0.25 sec with a narrow distribution (Figure 4F). When AFP was inactivated, this temporal
control of call duration was reduced, causing the production of abnormally short or long calls
(Figure 3E-I, Figure 4F).

We then quantified call stereotypy using identical statistical methods previously used to analyze
the change in zebra finch syllables following inactivation of LMAN [12,21]. In zebra finches, LMAN
inactivation causes an immediate reduction in syllable duration variability, quantified as the
coefficient of variation of syllable duration distributions [11,12]. Further, the phonological variability
of syllables is also dramatically reduced, as quantified with acoustic feature vectors derived from
the Sound Analysis Pro software package [21]. Using these identical analyses, we found that
inactivation of MO/NAO in budgerigars had the opposite effects of increasing the variability of
syllable duration and phonology (Figure 4G-H).

Notably, budgerigars exhibited defects in pitch and duration in the first calls they produced
immediately after TTX infusion (Figure 3E-I, green boxes). Furthermore, the effects were
reversible: when the drug was washed out, contact calls recovered to normal within hours (Figure

S1). In one bird we also used muscimol to inactivate AFP and observed similar changes in contact
calls (Figure S2), suggesting that the call defects under TTX were caused by inactivation of
MO/NAO, rather than blocking of fibers that pass through this region.

Inactivation of budgerigar AFP reduces the individuality of contact calls

The significant loss of pitch modulation and increase of phonological variability with AFP
inactivation may reduce individual signatures in contact calls. We first checked this possibility by
training a single VAE for calls produced under the PBS and TTX condition of all birds, then
embedding their latent features in the same UMAP space. Since normal calls of these stranger
birds are visually distinct (Figure 3E-I), they should form well-separated clusters in the UMAP.
Changes in acoustic structure induced by AFP inactivation will drive TTX calls to deviate from the
normal clusters. However, the deviation could happen in two ways that lead to different outcomes
on the individual signatures (Figure 5A). If AFP inactivation reduces individuality, calls from
different birds should move away from their normal clusters and converge into one large and
centered cluster (Figure 5A, hypothesis 1). Alternatively, if AFP inactivation affected each bird in
unique ways, individuality may be maintained, in which case the UMAP embeddings of the AFP-
inactivated calls would still reveal clearly distinct clusters associated with different individuals
(Figure 5A, hypothesis 2). Figure 5B shows that our data are more consistent with hypothesis 1:
calls converged to a centered cluster following AFP inactivation, suggesting the loss of individual
identity information.

However, since UMAP is a non-linear method, large distances in UMAP space do not directly
translate to large differences in acoustic space. To more rigorously test the loss of individual
signatures with AFP inactivation, we asked if a potential listener in the social group is still able to
decode caller identity when listening to calls produced under the TTX condition. We simulated
this situation by training a SVM classifier on the normal calls, then using it to decode the caller
identity at the TTX condition. We found that the decoding accuracy dropped significantly from
~95% to ~50% (Figure 5C), demonstrating a dramatic reduction in individual identity information.

Finally, we reasoned that although the original individual signatures in contact calls were greatly
reduced with AFP inactivation, different birds may not produce identical sounds, due to individual
differences in the biomechanics of the vocal organ or individually learned components that may
be driven by the NLC-AAC pathway. This could be seen in the UMAP embedding: although calls
produced under the TTX condition converge into one large central cluster, some level of
separation between individuals is preserved (Figure 5B). For quantification, we trained SVM
classifiers separately on the PBS and TTX datasets. We found that with enough information from
the spectrograms, the SVM could still decode caller identity on the TTX dataset with a decent
level of accuracy (Figure 5D). However, the decoding accuracy at the TTX condition was
significantly lower than the PBS condition across all ranges of principal dimensions (Figure 5D),
suggesting that AFP-inactivated calls were less discernible. We further quantified the separability
of calls produced by different individuals using a Silhouette score that contrasts intra- and inter-
cluster distances (Methods). We found that Silhouette score was significantly smaller on the TTX
dataset (Figure 5E), again suggesting that AFP-inactivated calls were less separable across
individuals.

Discussion

One of the most important contributions of the songbird field to neuroscience is the discovery that frontal cortical circuits, outputs of a basal ganglia-thalamocortical loop, actively inject variability into behavioral output for the purpose of learning [13,18,20,48]. Inactivation of LMAN transforms the variable song of juvenile birds into stereotyped syllables produced in a stereotyped sequence. And in adults, LMAN inactivations similarly reduce natural trial-by-trial variability of learned vocalizations. Here we tested if this result generalizes to budgerigars, a parrot species with a song system that, based on anatomy alone, has been hypothesized to have a similar functional organization [25,31].

Surprisingly, we found that inactivation of the output of the AFP, the NAO/MO complex, in parrots had the opposite effect as observed in songbirds: the acoustic structure of learned calls degraded and became more variable. These findings raise the possibility that the parrot song system functions according to different principles.

In light of these results, we discuss the substantial differences that exist between songbirds and parrots at behavioral, vocal mechanical, and neural circuit levels. Behaviorally, parrot vocalizations retain outsized plasticity even in adulthood and may even convey semantic information [42,51–54]. For example, individuals learn each other's calls during courtship and group assimilation and budgerigars can even be trained to use their calls to label a color of an upcoming choice in a delayed-match-to-sample task [55]. Budgerigars can also, in their warble vocalizations not studied in this paper, learn to produce hundreds to thousands of specific English words [56], and can pronounce specific words differently across trials (e.g. more slowly or quickly, "Hello", versus "Heeeellllllooooooo"). Such trial-by-trail flexibility in a word/syllable's pronunciation is commonplace in human speech but not observed in songbirds such as the zebra finch.

Parrots also differ at the level of vocal mechanics. Like humans, parrots vocalize with a single, bilaterally innervated, centered larynx and use their tongues to pattern sounds [28,57]. In contrast, songbirds pattern their sounds with bilaterally symmetric, unilaterally innervated syringes which are located at the top of each bronchus [58,59]. Finally, though the parrot and songbird song systems are both nucleated and contain distinct frontal and posterior pathways, there are also dramatic differences. First, each 'nucleus' of the parrot song system is recurrently interconnected with surrounding 'shell' tissue [27,31]. In songbirds, shells exist around nuclei, but are not substantially interconnected with the nuclei. Their roles in song learning are unclear, as they appear to be activated not by singing behavior but by general movement, for example, in trained hopping tasks [60,61]. Second, while RA axons specifically target vocalization-associated structures of the brainstem (e.g. respiratory nuclei and syringeal motor neurons) [62], AAC axons in the brainstem are more diffuse, raising the possibility that they control more than just vocalizations [28]. Third, recurrent long range feedback projections (most notably from AAC back to its inputs) abound in parrots (Figure 1B), while songbirds instead have primarily feedforward cortical pathways with recurrence only through a loop with the brainstem and thalamus (Figure 1A) [63]. This omnipresence of both short-range and long-range recurrent cortical feedback loops, a defining feature of the mammalian cortex [64,65], is absent in songbirds. Decades of theoretical work and more recent advances in machine learning suggest that, whereas feedforward networks are good at pattern

recognition and production, recurrent neural nets (RNNs) are suited for short-term memory
storage and complex dynamics [64,65], raising the possibility that the human-like recurrent network
architecture of the parrot vocal system is associated with parrots' outsized vocal capabilities. An
important caveat of our results is that our methods only enabled us to inactivate MO and NAO at
once. It remains possible that inactivation of only one of these structures would, like LMAN
inactivation, preserve the learned acoustic structure of the call and reduce variability. Future
studies with more specific viral or highly targeted lesion methods could in principle address these
possibilities, but it's noteworthy that our experimental approach, including our drug
concentrations, were the same as our past work in finches and yet yielded conceptually distinct
results.

Going forward, these experiments set the stage for two key future directions. First, because frontal
inactivations resulted in the loss of the temporally precise acoustic elements that define a bird's
unique call, our findings suggest that at least one of the frontal nuclei exhibits temporally precise,
call-locked discharge, as has been reported in songbird HVC but not LMAN [10,20]. Future
electrophysiological recordings targeted to either MO or NAO will test if synfire chain-like activity
important for songbird song temporal structure also exists in parrots—but in a distinct brain area
with distinct connectivity. Second, though our study only examined the vocalizations of birds in
isolation, it will be interesting to study the social dynamics of small groups of birds in which some
birds have their vocal signatures degraded with frontal inactivations described here. This would
test the ultimate function of these calls, for example if they are necessary for bond formation,
maintenance, coordinated flight, and overall social cohesion.

Acknowledgements

We thank Michael Sheehan, Lindy McBride, Yumeng Zhang, and members of the Goldberg lab for comments. We thank Jack Bradbury and Sandra Vehrencamp for inspiration for the study.

Author contributions

Z.Z. and J.G. designed the study. Z.Z. collected the data with help from F.N. and B.K.. Z.Z., J.C., F.N., and H.K.T. analyzed the data. H.K.T. received supervision from I.C.. Z.Z. and J.G. wrote the manuscript with help from all other authors.

Declaration of interests

The authors declare no competing interests.

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be answered by the Lead Contact, Jesse H. Goldberg (jessehgoldberg@gmail.com).

Materials availability

This study did not generate any unique biological materials.

Data and code availability

Data and code generated in this study can be obtained through the Lead Contact.

Experimental model and subject details

Animal care and experiments were carried out in accordance with NIH guidelines and were approved by the Cornell Institutional Animal Care and Use Committee. Subjects used in this study were 10 male and 6 female budgerigars (*Melopsittacus undulatus*) that aged between 1 to 3 years. Birds were bred in our aviary or obtained from outside breeders (Magnolia Bird Farm, California; Petsmart, New York; OK Birds, Oklahoma). We used 8 birds for contact call quantification, 3 for determining anatomical coordinates, and 5 for chemical inactivation. Birds were group-housed in flight cages or individually housed in wire cages with a 12-12 light cycle. Food and water were provided *ad libitum*.

Method details

Contact call playback

We conducted two types of playback experiments. For the first type (Figure 2), we recorded the call repertoire of 8 birds (5 males and 3 females) in the same cage by playing colony noise on-and-off in a trial structure. Colony noise was first recorded with a portable recorder (V618, EVIDA) inside their flight cage for 2-3 days. We removed the silent portion of the recording and chopped it into 2-min segments. We then isolated individual birds to a wire cage inside a sound attenuating box (Figure 2A). A flat-spectrum speaker (iLoud Micro Monitor, IK Multimedia) was placed outside the box (~10 cm) and pointed toward an air-exchange hole on the box. We adjusted the speaker

volume so the playback sound measured 53-55 dB inside the box. For each trial, we randomly
chose a colony noise segment and played it to the bird through the speaker. The inter-trial interval
followed an exponential distribution (mean: 10 mins, range: 7.5-12 mins). Playback and response
signals were recorded with directional microphones (Pro 35, Audio-Technica), bandpass-filtered
(30-15,000 Hz) and amplified with a preamplifier (XL48, Midas), and digitized through a DAQ
(USB6210, National Instruments) before sending to the computer. We wrote custom LabView
codes to control the playback and recording. Each bird was recorded for 2 days.

For the second type of playback (Figure 3-4), we recorded responses to individual contact calls
with the same hardware setup. We first constructed the call repertoire of each bird from its colony-
noise responses and normalized the amplitude of all calls to the same value. For each trial, we
randomly selected one caller and one call from its repertoire, then played it three times to the
experimental bird with intervals uniformly sampled between 2-5 sec. The inter-trial interval
followed an exponential distribution (mean: 5 mins, range: 4-6 mins).

Surgery and histology

To perform surgery and histology in the budgerigar, we adopted procedures that are commonly
used in the zebra finch. All surgeries were performed under isoflurane anesthesia. Origin for the
anterior/posterior and medial/lateral axis was established relative to the bifurcate blood vessel
lambda. Brain surface after removing dura mater was used to zero the dorsal/ventral axis. We
adjusted the head angle of the bird until the anterior skull was horizontal, which was referred to
as the zero-degree angle. To access a brain region, we removed the first layer of skull with a
cranial drill, the second layer of skull with a scalpel, and the dura mater with sharp forceps. After
injection or implant, the exposed brain surface was sealed with biocompatible adhesive (World
Precision Instruments, KWIK-CAST). Birds were allowed to recover for at least one week before
any data collection. For histology, brains were fixed in 4% paraformaldehyde solution for 2-3 days,
then cut into 100-µm coronal slices with a vibratome and mounted on slides in polyvinyl alcohol
(Sigma-Aldrich, 10981-100ML). Dark-field and fluorescent images were acquired using a
fluorescence scope (Leica DM4000 B).

Tracer injection

365   To determine the stereotactic coordinates of MO/NAO, we first calculated a rough estimate based
366   on the published budgerigar brain atlas (http://www.brauthlab.umd.edu/atlas.htm). We then
367   injected 100 nL fluorescent cholera toxin subunit B (CTB, C34775, Invitrogen) tracers into the
368   brain and measured the targeting accuracy with histology after 7-9 days. The coordinates were
369   iteratively improved with 6 injections in 3 birds.

Microdialysis probe implant

We made custom microdialysis probes according to the method in Aronov et al 2008. To target
MO/NAO, the inflow polyimide tubing is approximately 2.25 mm long. The cellulose membrane
measures roughly 0.2 mm in width and 1 mm in length, with 0.7 mm exposed to deliver drugs.
The probes were implanted bilaterally at 8.80 AP, 2.90 ML and 1.45 DV (unit: mm), with a 30° AP
angle and 20° ML angle. To secure the implants, we inserted 4 insect pins into the skull and
covered the base of probes with dental cement (C&B Metabond, Parkell, S380). The probes were

flushed regularly with PBS to prevent clogging. After data collection, we infused fluorescein (10
mM) into the probe to visualize implant location and drug diffusion.

Pharmacological inactivation

Compared to direct injection, a higher concentration of drug is needed with the microdialysis
probe, since the solute needs to diffuse passively through the cellulose membrane (Aronov et al
2008). Following previous studies in the zebra finch (Aronov et al 2008), we diluted tetrodotoxin
citrate (Abcam, ab120055-1MG) in PBS to 50 µM. One week after surgery, we started infusing
PBS into probes to acclimate the bird to the procedure. Infusion was conducted roughly 1 hour
before the cage light turned on. For each probe, we infused 1.0 mL solution at a rate of ~0.2
mL/min. For TTX, we left the drug in the probe for 15 mins then washed it out with PBS. The bird
was then placed back to the cage to rest before the light turned on and behavioral trials began.
We used the same procedure for TTX and PBS, and conditions were alternated daily, so a direct
comparison is meaningful. We had 2-3 infusion replicates for each bird under each condition. For
experiments using muscimol, we diluted it to 1.5 mg/mL in PBS (Aronov et al 2008). Since
muscimol degrades faster than TTX, to sample enough contact calls, we kept muscimol in the
probes during data collection for about 6 hours, then washed it out with PBS.

Quantification and statistical analysis

Contact call identification

We wrote custom Matlab scripts to identify contact calls from audio recordings. Noise at low and
high frequency was first removed with a bandpass Butterworth filter (500-10,000 Hz). We then
calculated the spectral flatness (Wiener energy) and set a threshold (0.65) to distinguish vocal
signal from broadband noise. Segments that passed the threshold were considered candidates
and exported for visual inspection. Contact call is characteristic in having a major modulated
frequency band between 2-4 kHz. Cage noise, squawks, harmonic calls, and warble were
removed from downstream analysis.

Acoustic feature analysis

We used the SAT Matlab package from Sound Analysis Pro 2011 to calculate acoustic features
of identified calls (http://soundanalysispro.com/matlab-sat). Pitch modulation depth was defined
as (max - min) / mean of pitch. Spectrograms were calculated using the Short-Time Fourier
Transform (STFT) algorithm in Matlab (window size 512, nfft 512). Variability of acoustic features
was calculated based on the method in Olveczky et al 2005, to facilitate direct comparison to
results in zebra finches. Briefly, we randomly sampled 50 pairs of calls from each bird under each
condition. Since subject F6 had two normal call types (Figure 5B), we only sampled call pairs that
belonged to the same type. We then used the 'Similarity batch' function in Sound Analysis Pro
2011 to calculate pairwise similarity scores with the default parameters except for the 'min
interval', which was set to 25 ms. Following Olveczky et al 2005, values of 'accuracy' were used
as the metric for similarity and further converted to variability scores.

Variational auto-encoder (VAE)

We followed the method in Goffinet et al 2021 to train the VAE network for contact call analysis. Briefly, spectrograms of amplitude-normalized calls were centered in a 128*128 matrix and used as the inputs. We performed resampling so each individual had the same number of calls, then randomly split the dataset for training (80%) and testing (20%). The VAE architecture was adapted from Goffinet et al 2021 with dimension of latent features fixed to 32. The network was trained for 100 epochs at learning rate 0.001 and batch size 32 with the Adam optimizer implemented in Pytorch 1.12.1. We then calculated the latent features of each call by feeding it into the encoder network. Separate networks were trained for the behavioral (Figure 2) and inactivation experiments (Figure 3-5).

Support vector machine (SVM)

In past work, individual signatures in vocal signals were quantified with two major categories of approaches: 1) methods based on information theory [66]; 2) classification models such as support vector machine (SVM) [43,67]. Although information theoretic approaches allow comparison across different systems and quantification of features' contribution to individuality, here our primary goal was to explicitly compare discriminability of the same system across conditions, e.g. train a model on the PBS dataset then check performance on the TTX dataset. We therefore adopted the classification approach to quantify individuality in contact calls. Specifically, we built SVM classifiers to decode individual identity from contact calls using the scikit-learn module in Python. We first conducted principal component analysis (PCA) on the VAE outputs to obtain the most informative features, then used them as inputs for SVM. We varied the dimension of principal inputs and calculated how the decoding accuracy changed accordingly. For each model, we sampled an equal number of calls from each individual to balance the dataset, then split it for training (80%) and testing (20%). Accuracy was calculated only on the testing dataset. The sampling procedure was repeated 50 times for each model. The mean accuracy and standard deviation were plotted in the figures.

Silhouette score calculation

We calculated the Silhouette score to quantify the separability of calls produced by different individuals. We first trained all calls produced on the PBS and TTX condition in a single VAE network, then reduced the dimension of latent features using PCA. We focused on the first 10 principal components, since together they could explain 97.8% of the total variance. We calculated the Silhouette score separately for the PBS and TTX dataset. We sampled 50 calls from each individual, then used the scikit-learn module in Python to calculate the Silhouette score on the first 10 principal components. The sampling was repeated 100 times to obtain a distribution of Silhouette scores. Shuffling of labels was also performed at each sampling to obtain a null distribution.

References

1. Jarvis, E.D. (2019). Evolution of vocal learning and spoken language. Science *366*, 50–54.

2. Petkov, C.I., and Jarvis, E.D. (2012). Birds, primates, and spoken language origins: behavioral phenotypes and neurobiological substrates. Front. Evol. Neurosci. *4*, 12.

3. Tyack, P.L. (2020). A taxonomy for vocal learning. Philos. Trans. R. Soc. Lond. B Biol. Sci. *375*, 20180406.

4. ten Cate, C. (2021). Re-evaluating vocal production learning in non-oscine birds. Philos. Trans. R. Soc. Lond. B Biol. Sci. *376*, 20200249.

5. Sober, S.J., Wohlgemuth, M.J., and Brainard, M.S. (2008). Central contributions to acoustic variation in birdsong. J. Neurosci. *28*, 10370–10379.

6. Warren, T.L., Tumer, E.C., Charlesworth, J.D., and Brainard, M.S. (2011). Mechanisms and time course of vocal learning and consolidation in the adult songbird. J. Neurophysiol. *106*, 1806–1821.

7. Nottebohm, F., Stokes, T.M., and Leonard, C.M. (1976). Central control of song in the canary, Serinus canarius. J. Comp. Neurol. *165*, 457–486.

8. Aronov, D., Veit, L., Goldberg, J.H., and Fee, M.S. (2011). Two distinct modes of forebrain circuit dynamics underlie temporal patterning in the vocalizations of young songbirds. Journal of Neuroscience *31*, 16353–16368.

9. Dugas-Ford, J., Rowell, J.J., and Ragsdale, C.W. (2012). Cell-type homologies and the origins of the neocortex. Proc. Natl. Acad. Sci. U. S. A. *109*, 16974–16979.

10. Fee, M.S., Kozhevnikov, A.A., and Hahnloser, R.H. (2004). Neural mechanisms of vocal sequence generation in the songbird. Ann. N. Y. Acad. Sci. *1016*, 153–170.

11. Chen, J.R., Stepanek, L., and Doupe, A.J. (2014). Differential contributions of basal ganglia and thalamus to song initiation, tempo, and structure. J. Neurophysiol. *111*, 248–257.

12. Goldberg, J.H., and Fee, M.S. (2011). Vocal babbling in songbirds requires the basal ganglia-recipient motor thalamus but not the basal ganglia. J. Neurophysiol. *105*, 2729–2739.

13. Aronov, D., Andalman, A.S., and Fee, M.S. (2008). A specialized forebrain circuit for vocal babbling in the juvenile songbird. Science *320*, 630–634.

14. Picardo, M.A., Merel, J., Katlowitz, K.A., Vallentin, D., Okobi, D.E., Benezra, S.E., Clary, R.C., Pnevmatikakis, E.A., Paninski, L., and Long, M.A. (2016). Population-Level Representation of a Temporal Sequence Underlying Song Production in the Zebra Finch. Neuron *90*, 866–876.

15. Hamaguchi, K., Tschida, K.A., Yoon, I., Donald, B.R., and Mooney, R. (2014). Auditory synapses to song premotor neurons are gated off during vocalization in zebra finches. Elife *3*, e01833.

16. Fortune, E.S., Rodríguez, C., Li, D., Ball, G.F., and Coleman, M.J. (2011). Neural

mechanisms for the coordination of duet singing in wrens. Science *334*, 666–670.

17. Hoffmann, S., Trost, L., Voigt, C., Leitner, S., Lemazina, A., Sagunsky, H., Abels, M.,
Kollmansperger, S., Maat, A.T., and Gahr, M. (2019). Duets recorded in the wild reveal that
interindividually coordinated motor control enables cooperative behavior. Nat. Commun. *10*,
2577.

18. Bottjer, S.W., Miesner, E.A., and Arnold, A.P. (1984). Forebrain lesions disrupt
development but not maintenance of song in passerine birds. Science *224*, 901–903.

19. Scharff, C., and Nottebohm, F. (1991). A comparative study of the behavioral deficits
following lesions of various parts of the zebra finch song system: implications for vocal
learning. J. Neurosci. *11*, 2896–2913.

20. Kao, M.H., Doupe, A.J., and Brainard, M.S. (2005). Contributions of an avian basal ganglia-
forebrain circuit to real-time modulation of song. Nature *433*, 638–643.

21. Olveczky, B.P., Andalman, A.S., and Fee, M.S. (2005). Vocal experimentation in the
juvenile songbird requires a basal ganglia circuit. PLoS Biol. *3*, e153.

22. Edelman, G.M. (1993). Neural Darwinism: selection and reentrant signaling in higher brain
function. Neuron *10*, 115–125.

23. Ben-Pazi, H., Gross-Tsur, V., Bergman, H., and Shalev, R.S. (2003). Abnormal rhythmic
motor response in children with attention-deficit-hyperactivity disorder. Dev. Med. Child
Neurol. *45*, 743–745.

24. Singh Alvarado, J., Goffinet, J., Michael, V., Liberti, W., III, Hatfield, J., Gardner, T.,
Pearson, J., and Mooney, R. (2021). Neural dynamics underlying birdsong practice and
performance. Nature *599*, 635–639.

25. Paton, J.A., Manogue, K.R., and Nottebohm, F. (1981). Bilateral organization of the vocal
control pathway in the budgerigar, Melopsittacus undulatus. J. Neurosci. *1*, 1279–1288.

26. Brauth, S.E., Heaton, J.T., Durand, S.E., Liang, W., and Hall, W.S. (1994). Functional
anatomy of forebrain auditory pathways in the budgerigar (Melopsittacus undulatus). Brain
Behav. Evol. *44*, 210–233.

27. Striedter, G.F. (1994). The vocal control pathways in budgerigars differ from those in
songbirds. J. Comp. Neurol. *343*, 35–56.

28. Brauth, S.E., Heaton, J.T., Shea, S.D., Durand, S.E., and Hall, W.S. (1997). Functional
anatomy of forebrain vocal control pathways in the budgerigar (Melopsittacus undulatus).
Ann. N. Y. Acad. Sci. *807*, 368–385.

29. Durand, S.E., Heaton, J.T., Amateau, S.K., and Brauth, S.E. (1997). Vocal control
pathways through the anterior forebrain of a parrot (Melopsittacus undulatus). J. Comp.
Neurol. *377*, 179–206.

30. Jarvis, E.D., and Mello, C.V. (2000). Molecular mapping of brain areas involved in parrot
vocal communication. J. Comp. Neurol. *419*, 1–31.

31. Chakraborty, M., Walløe, S., Nedergaard, S., Fridel, E.E., Dabelsteen, T., Pakkenberg, B.,

Bertelsen, M.F., Dorrestein, G.M., Brauth, S.E., Durand, S.E., et al. (2015). Core and Shell
Song Systems Unique to the Parrot Brain. PLoS One *10*, e0118496.

32. Heaton, J.T., and Brauth, S.E. (2000). Effects of lesions of the central nucleus of the
anterior archistriatum on contact call and warble song production in the budgerigar
(Melopsittacus undulatus). Neurobiol. Learn. Mem. *73*, 207–242.

33. Lavenex, P.B. (2000). Lesions in the budgerigar vocal control nucleus NLc affect
production, but not memory, of english words and natural vocalizations. J. Comp. Neurol.
421, 437–460.

34. Tchernichovski, O., Lints, T.J., Deregnaucourt, S., Cimenser, A., and Mitra, P.P. (2004).
Studying the song development process: rationale and methods. Ann. N. Y. Acad. Sci.
1016, 348–363.

35. Kollmorgen, S., Hahnloser, R.H.R., and Mante, V. (2020). Nearest neighbours reveal fast
and slow components of motor learning. Nature *577*, 526–530.

36. Goffinet, J., Brudner, S., Mooney, R., and Pearson, J. (2021). Low-dimensional learned
feature spaces quantify individual and group differences in vocal repertoires. Elife *10*.
10.7554/eLife.67855.

37. Farabaugh, S.M., and Dooling, R.J. (1996). Acoustic Communication in Parrots: Laboratory
and Field Studies of Budgerigars, *Melopsittacus undulatus*. In Ecology and Evolution of
Acoustic Communication in Birds.

38. Brown, S.D., Dooling, R.J., and O'Grady, K.E. (1988). Perceptual organization of acoustic
stimuli by budgerigars (Melopsittacus undulatus): III. Contact calls. J. Comp. Psychol. *102*,
236–247.

39. Farabaugh, S.M., Linzenbold, A., and Dooling, R.J. (1994). Vocal plasticity in budgerigars
(Melopsittacus undulatus): evidence for social factors in the learning of contact calls. J.
Comp. Psychol. *108*, 81–92.

40. Hile, A.G., Plummer, T.K., and Striedter, G.F. (2000). Male vocal imitation produces call
convergence during pair bonding in budgerigars, Melopsittacus undulatus. Anim. Behav.
59, 1209–1218.

41. Hile, A.G., and Striedter, G.F. (2000). Call Convergence within Groups of Female
Budgerigars (Melopsittacus undulatus). Ethology *106*, 1105–1114.

42. Dahlin, C.R., Young, A.M., Cordier, B., Mundry, R., and Wright, T.F. (2014). A test of
multiple hypotheses for the function of call sharing in female budgerigars, Melopsittacus
undulatus. Behav. Ecol. Sociobiol. *68*, 145–161.

43. Tachibana, R.O., Oosugi, N., and Okanoya, K. (2014). Semi-automatic classification of
birdsong elements using a linear support vector machine. PLoS One *9*, e92584.

44. Brockway, B.F. (1964). Ethological studies of the budgerigar (Melopsittacus undulatus):
Non-reproductive behavior. Behaviour *22*, 193–222.

45. Brockway, B.F. (1964). Ethological studies of the budgerigar: reproductive behavior.
Behaviour *23*, 294–323.

46. Wyndham, E. (1980). Diurnal cycle, behaviour and social organization of the budgerigar
Melopsittacus undulatus. Emu - Austral Ornithology *80*, 25–33.

47. Otchy, T.M., Wolff, S.B., Rhee, J.Y., Pehlevan, C., Kawai, R., Kempf, A., Gobes, S.M., and
Olveczky, B.P. (2015). Acute off-target effects of neural circuit manipulations. Nature *528*,
358–363.

48. Andalman, A.S., and Fee, M.S. (2009). A basal ganglia-forebrain circuit in the songbird
biases motor output to avoid vocal errors. Proc. Natl. Acad. Sci. U. S. A. *106*, 12518–
12523.

49. Goldberg, J.H., and Fee, M.S. (2012). A cortical motor nucleus drives the basal ganglia-
recipient thalamus in singing birds. Nat. Neurosci. *15*, 620–627.

50. Ölveczky, B.P., Otchy, T.M., Goldberg, J.H., Aronov, D., and Fee, M.S. (2011). Changes in
the neural control of a complex motor sequence during learning. J. Neurophysiol. *106*, 386–
397.

51. Hobson, E.A., Avery, M.L., and Wright, T.F. (2014). The socioecology of Monk Parakeets:
Insights into parrot social complexity. The Auk: Ornithological Advances *131*, 756–775.

52. Ikkatai, Y., Watanabe, S., and Izawa, E.-I. (2016). Reconciliation and third-party affiliation in
pair-bond budgerigars (Melopsittacus undulatus). Behaviour *153*, 1173–1193.

53. Berg, K.S., Delgado, S., Cortopassi, K.A., Beissinger, S.R., and Bradbury, J.W. (2012).
Vertical transmission of learned signatures in a wild parrot. Proc. Biol. Sci. *279*, 585–591.

54. Balsby, T.J.S., Momberg, J.V., and Dabelsteen, T. (2012). Vocal imitation in parrots allows
addressing of specific individuals in a dynamic communication network. PLoS One *7*.

55. Manabe, K., Kawashima, T., and Staddon, J.E.R. (1995). Differential vocalization in
budgerigars: Towards an experimental analysis of naming. J. Exp. Anal. Behav. *63*, 111–
126.

56. Words, Puck the Parakeet speaks 1728 English (1995). Guinness world records (Guinness
World Records).

57. Beckers, G.J.L., Nelson, B.S., and Suthers, R.A. (2004). Vocal-tract filtering by lingual
articulation in a parrot. Curr. Biol. *14*, 1592–1597.

58. Goller, F., and Cooper, B.G. (2004). Peripheral motor dynamics of song production in the
zebra finch. Ann. N. Y. Acad. Sci. *1016*, 130–152.

59. During, D.N., Ziegler, A., Thompson, C.K., Faber, C., Muller, J., Scharff, C., and Elemans,
C.P. (2013). The songbird syrinx morphome: a three-dimensional, high-resolution,
interactive morphological map of the zebra finch vocal organ. BMC Biol. *11*, 1.

60. Bottjer, S.W., Brady, J.D., and Cribbs, B. (2000). Connections of a motor cortical region in
zebra finches: relation to pathways for vocal learning. J. Comp. Neurol. *420*, 244–260.

61. Feenders, G., Liedvogel, M., Rivas, M., Zapka, M., Horita, H., Hara, E., Wada, K.,
Mouritsen, H., and Jarvis, E.D. (2008). Molecular mapping of movement-associated areas
in the avian brain: a motor theory for vocal learning origin. PLoS One *3*, e1768.

62. Wild, J.M. (1993). Descending projections of the songbird nucleus robustus archistriatalis.
J. Comp. Neurol. *338*, 225–241.

63. Schmidt, M.F., Ashmore, R.C., and Vu, E.T. (2004). Bilateral control and interhemispheric
coordination in the avian song motor system. Ann. N. Y. Acad. Sci. *1016*, 171–186.

64. Sussillo, D., and Abbott, L.F. (2009). Generating coherent patterns of activity from chaotic
neural networks. Neuron *63*, 544–557.

65. Sanchez-Vives, M.V., and McCormick, D.A. (2000). Cellular and network mechanisms of
rhythmic recurrent activity in neocortex. Nat. Neurosci. *3*, 1027–1034.

66. Beecher, M.D. (1989). Signaling systems for individual recognition: an information theory
approach. Anim. Behav. *38*, 248–261.

67. Gelfand, D.L., and McCracken, G.F. (1986). Individual variation in the isolation calls of
Mexican free-tailed bat pups (Tadarida brasiliensis mexicana). Anim. Behav. *34*, 1078–
1086.

Figure 1. Vocal learning circuits in songbirds and parrots

(A) Sagittal view showing the posterior motor pathway (green) and anterior forebrain pathway
(pink) in the zebra finch. Orange line indicates the songbird-specific connection from HVC to Area
X. Abbreviations: H, hyperpallium; M, mesopallium; N, nidopallium; A, arcopallium; RA, robust
nucleus of the arcopallium; HVC, proper name; LMAN, lateral magnocellular nucleus of the
anterior nidopallium; DLM, medial portion of the dorsolateral thalamus; VTA, ventral tegmental
area.

(B) Vocal learning circuits in budgerigars also have the posterior (green) and anterior (pink)
pathways. Cyan circles and lines indicate the parrot-specific vocal nuclei and connections. The
shell structures that surround the vocal nuclei are not depicted. Abbreviations: AAC, central
nucleus of the anterior arcopallium; NLC, central nucleus of the lateral nidopallium; MO, oval
nucleus of the anterior mesopallium; NAO, oval nucleus of the anterior nidopallium; MMSt,
magnocellular nucleus of the medial striatum; DMM, magnocellular nucleus of the dorsal
thalamus.

Figure 2. Budgerigar contact calls contain individual signature

(A) Schematics showing the recording setup. Two directional microphones point in the opposite direction to capture sound from inside or outside the box.

(B) Spectrograms of example contact calls from a group of 8 budgerigars (5 males, 3 females) at week 15 since assimilation. All plots have the same scale on the x- and y- axis. Animal and cluster labels above each spectrogram correspond to labels in panel *E*. Asterisk symbol marks a call type that was shared across all 8 individuals, while plus symbol marks one that was shared across 7 individuals.

(C) Pipeline for analyzing contact calls.

(D-E) UMAP embedding of contact calls at week 3 (D) and week 15 (E) since group assimilation.

Two datasets were trained through the same VAE and embedded in the same UMAP space, but

shown separately for clarity. Each dot is a call, colored by caller identity. A maximum of 600 calls

were shown for each individual. Animal and cluster labels in (E) correspond to spectrograms in

panel *B*. Dashed lines in (E) circled the two major clusters that were shared across individuals

and marked by an asterisk or plus symbol in panel *B*.

(F) Accuracy of SVM classifiers that decode individual identity from call latent features as a

function of input dimensions. Solid lines are models trained on real data, while dashed lines are

models trained on shuffled data.

(G) Accuracy of SVM classifiers that were trained and tested solely on the calls that belong to the

two large shared clusters (marked by asterisk and plus symbol in panel B and E).

(H) Accuracy of SVM classifiers that were trained on one time point and used to decode individual

identity at a different time point.

Figure 3. Frontal cortical inactivation abolishes pitch modulation in contact calls

(A) Schematics of the inactivation experiments. Reverse microdialysis probes were implanted in the center of the MO/NAO complex and infused with either 50 micromolar TTX or PBS.

(B) Histological confirmation of implant site in one representative bird. At the end of experiments fluorescein was infused into the probes to visualize diffusion and probe placement. Fluorescein

covered both MO and NAO. Dashed circles at the bottom row indicate the boundaries of
corresponding nuclei.

(C-D) Example trials from one bird under the PBS (C) and TTX (D) condition. Top rows show
spectrograms of the stimuli, bottom rows show time-aligned responses. Y-axis is frequency in
kHz.

(E-I) Example spectrograms from 5 birds under the PBS (top row) and TTX (bottom row) condition.
The first 2-3 spectrograms at each row in the green boxes are the first calls produced during each
PBS or TTX infusion replicate. Other spectrograms were selected randomly. M6-M7, males; F4-
F6, females. Y-axis is frequency in kHz. Numbers above each spectrogram indicate minutes since
drug infusion.

Figure 4. Frontal cortical inactivation increases variability of contact calls

(A-E) Compare behavioral trial statistics (A-B) and acoustic features (C-E) between the PBS (blue) and TTX (red) condition. Each colored dot is one trial (A-B) or one call (C-E). The violins indicate the shape of distribution. The black line and white dot indicate the mean and median. P-value: *<0.05, **<0.01, ***<0.001, ns not significant, with Mann-Whitney U tests.

(F) Histograms of contact call duration under the PBS (blue) and TTX (red) condition.

(G) Variability of call duration quantified by coefficient of variation (CV). Each dot indicates data from one bird.

684 (H) Variability of acoustic features calculated by Sound Analysis Pro (SAP). Each dot indicates
685 the averaged dissimilarity of 50 randomly selected pairs of calls in a bird under the PBS or TTX
686 condition.

Figure 5. Frontal cortical inactivation reduces individual signatures in contact calls.

(A) Hypotheses on how AFP inactivation would change the embedding of contact calls in the UMAP space. Clusters are colored according to individual identity and experimental condition. Gray arrows point from PBS to TTX condition in the same birds.

(B) UMAP embedding of contact calls. Each dot indicates a call, colored by caller identity and experiment condition. Gray arrows point from the centroid of PBS cluster to the centroid of TTX cluster for each bird.

(C) SVM classifier accuracy as a function of input dimensions. The model was trained on the PBS condition, then used to decode individual identity at the TTX condition. Dots indicate the mean accuracy, vertical lines indicate the standard deviation.

(D) Same as (C), but two models were trained separately for PBS and TTX.

(E) Separability of contact calls produced by different individuals at the PBS and TTX condition, as quantified by Silhouette score (Methods). Equal number of calls were sampled from each individual for the analysis. The results of 100 sampling are plotted. Colored violins indicate the shape of Silhouette score distribution. Black lines and white dots indicate the mean and median values, respectively.

Figure S1. Contact calls gradually recover from AFP inactivation

Spectrograms of contact calls from one example TTX infusion trial. The green and red boxes show one call spectrogram from the same bird the day before and after TTX infusion, respectively. Numbers above other spectrograms indicate the time since TTX infusion in the unit of hour. The contact calls began to recover gradually to normal shape after 9-10 hours.

Figure S2. Inactivation by muscimol had a similar effect on contact call acoustic structure

Spectrograms of contact calls from one example muscimol infusion trial. The green box shows one call spectrogram from the same bird the day before muscimol infusion. Numbers above spectrograms at the bottom row indicate the time in hours since muscimol was washed out with saline.