Title: CREID – A ChemoReceptor-Effector Interaction Database

Authors:

Vincent Peta; vincent.peta@usd.edu; Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107; ORCID: 0000-0002-9901-5207

Timothy Hartman; tim.hartman@coyotes.usd.edu; Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107; ORCID: 0000-0002-4127-9672

Shiva Aryal; shiva.aryal@coyotes.usd.edu; Computer Science Department, University of South Dakota, Vermillion, SD 57069; ORCID: 0000-0001-8537-1115

Bichar Shrestha Gurung; Bichar.ShresthaGurun@coyotes.usd.edu; Computer Science Department, University of South Dakota, Vermillion, SD 57069; ORCID: 0000-0002-9818-5108

Ram Singh; ram.singh@sdsmt.edu; Chemical and Biological Engineering Department, South Dakota School of Mines and Technology, Rapid City, SD 57701; ORCID: 0000-0002-3831-6487

Samuel Hass; samuel.haas@coyotes.usd.edu; Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107; ORCID: 0000-0003-2557-2358

Alain Bomgni; alain.bomgni@usd.edu; Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107; ORCID: 0000-0002-3377-7321

Tuyen Do; tuyen.do@usd.edu; Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107; ORCID: 0000-0002-1817-4565

Saurabh Sudha Dhiman; saurabh.dhiman@sdsmt.edu; Department of Chemical and Biological Engineering, South Dakota School of Mines and Technology, Rapid City, SD 57701, ORCID:

Venkataramana Gadhamshetty; venkataramana.gadhamshetty@sdsmt.edu; Civil and Environmental Engineering Department, South Dakota School of Mines and Technology, Rapid City, SD 57701; ORCID: 0000-0002-8418-3515

Etienne Z. Gnimpieba; etienne.gnimpieba@usd.edu; Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107; ORCID: 0000-0002-5338-084X

Affiliations:

Computer Science Department, University of South Dakota, Vermillion, SD 57069

Biomedical Engineering Department, University of South Dakota, Sioux Falls, SD 57107

Department of Chemical and Biological Engineering, South Dakota School of Mines and Technology, Rapid City, SD 57701

Department of Civil and Environmental Engineering, South Dakota School of Mines and Technology, Rapid City, SD 57701

Corresponding author and Lead Contact: Etienne Z. Gnimpieba (etienne.gnimpieba@usd.edu)

Abstract: The ChemoReceptor-Effector Interaction Database (CREID) is a collection of bacterial chemoreceptor and effector protein and interaction data to understand the process that chemoreceptors and effectors play in various environments. Our website includes terms associated with chemosensory pathways to educate users and those involved in collaborative research to help them understand this complex biological network. It includes 2,440 proteins involved in chemoreceptor and effector systems from 7 different bacterial families with 1,996 chemoeffector interactions. It is available at https://react-creid.bicbioeng.org.

Key Highlights

1. CREID links bacterial chemoreceptors with their associated effectors.

2. Researchers interested in what attracts or repels bacteria can use CREID as a comprehensive source for information.

3. Biosensor developers can leverage CREID to discover better interactions for their applications.

4. CREID reveals knowledge gaps in chemoreceptor-effector interactions for both model and non-model organisms.

Chemoreceptors & Effectors in the Environment

Bacterial chemosensory systems are critical in the movement behaviors of bacterial species, known as chemotaxis. Bacterial chemotaxis, movement under the influence of a chemical gradient, either towards (positive chemotaxis) or away (negative chemotaxis) to find optimum conditions for growth and survival due to chemoattractants or chemorepellents respectively. These chemosensory-signaling pathways allow the bacteria to sense the environment around them and travel toward or away from areas of stimuli, such as carbon sources and toxic chemicals. Chemotaxis could be based on bacterial behavioral responses such as energy taxis (stimuli affecting cellular energy), phototaxis (light) (Taylor & Zhulin, 1998), aerotaxis (oxygen concentration in niche) (Taylor, 1983), redox taxis (Armitage, 1997) and taxis to alternative electron acceptors (Grishanin et al., 1997).

The same stimulant could be attractant and repellent for two different bacteria or change in concentration could also reverse its effect from attractant to repellent (Shioi et al., 1987). Chemoreceptors are multi-domain proteins that act as input-signal receivers (that bind to ligands) in the environment and transfer it inside the cell to start the mechanism of response such as chemotaxis (Figure 1). Different stimuli (chemoeffectors) elicit unique responses which results in bacterial movement towards a specific stimulus (Bi & Lai, 2015a; Keegstra et al., 2022).

In marine environments multiple communities such as phytoplankton, zooplankton, and viruses co-exist with microbes (photoautotrophs, photoheterotrophs and heterotrophs) (Figure 2.). These communities have interactions at multiple levels for nutritional exchange. Marine environments are dynamic due to multiple factors such as nutrients which vary depending on the water current, sunlight, upwelling of sea water to various other factors (pH, temperature, salinity, oxygen, pressure) (Brumley et al., 2020). At the same time, nutrient dynamics can also be influenced by the microbial population themselves, which can augment the dissolved organic material (DOM) and inorganic substrates in the water (Blackburn et al., 1997; Jackson, 1980; McCarthy & Goldman, 1979; Smriga et al., 2016). These microenvironments (phycosphere (100 um) and the diffusion boundary layer are rich in significant amounts of simple sugars, amino acids, organic acids, complex polysaccharides, and lipids and are active zones for heterotrophic bacteria (Aaronson, 1978; Fogg, 1977; Hellebust, 1965; Jones & Cannon, 1986; Sharp, 1977). For example, *Synechococcus* sp. CS-94 RRIMP N1 releases 34 nitrogen containing compounds in marine environments and *Marinobacter adhaerens* HP15 metabolize 27 compounds (out of 34) released from strain CS-94 RRIMP N1 (Raina et al., 2023).

In soil systems, plant-microbe interaction is very critical. Plants recruit microbes by releasing root-exudates for microbial nutritional-supply and defense against pathogens. The chemical composition of root exudates has a direct effect on the rhizosphere communities nearest to the plant-host. These include plant growth-promoting rhizobacteria (PGPR) genera such as *Bacillus, Pseudomonas, Enterobacter*, *Burkholderia, Arthrobacter,* and *Paenibacillus* (Sasse et al., 2017; Zhang et al., 2017). For example, the

banana root exudate fumaric acid attracts the Gram-positive *Bacillus subtilis* N11 and stimulates biofilm formation (Zhang et al., 2014); malic acid exuded by *Arabidopsis* stimulates binding to roots and biofilm formation on roots by *B. subtilis* strain FB17 (Rudrappa et al., 2008). Bacterial growth and antifungal activity of certain species of Gram-negative *Pseudomonas* spp. is dependent on organic acids and sugars isolated from tomato root exudates (Kravchenko et al., 2003; Mhlongo et al., 2018). Among PGPR, species of *Pseudomonas* and *Bacillus* are the best studied model organisms for beneficial plant-microbe interaction (Weller et al., 2002; Raaijmakers et al., 2010).

Beneficial bacteria use chemotaxis to move toward plant root chemoeffectors (exudates) by means of chemoreceptor pathways. The plant pathogen *Agrobacterium fabrum* uses chemotaxis to infect dicotyledon plants which includes roses and geraniums (Wang et al., 2021). Wang et al discovered the chemoreceptor protein responsible for this chemotaxis. With these findings the community can leverage this discovery to create new antimicrobial agents for dicotyledons or seeds that have two embryonic leaves. As more microbiologists unveil new chemoreceptor-effector interactions, more microbe- attracting and repelling agents can be discovered—even those that target specific genes that code for chemoreceptors.

Environmental Factors and Conditions

Chemoeffectors occur both naturally and synthetically in an environmental system as any molecule or energy that either attracts or repels microbes (Bi & Lai, 2015b). This includes inorganic elements (metal ions), compounds (organic, aromatic), light, oxygen level and heat. These chemoeffectors could behave differently with variation in micro-niche conditions such as pH, salinity, humidity, barometric pressure, dissolved organic matter, distance between chemoeffectors and microbes, water turbulence, and depth of aquatic environment. Chemosensory pathways drive not just chemotaxis but also biofilm formation in a

crosstalk fashion. Mediation of biofilm formation on can be influenced by various conditions in solid,

fluid, and aerial environments, which can impact how bacterial species congregate and interact in biofilms

(Boyeldieu et al., 2020; Huang et al., 2019). It is well established that chemotaxis play an important role

in biofilm formation(O'Toole & Kolter, 1998; Pratt & Kolter, 1998; Prigent-Combaret et al., 1999; Watnick

& Kolter, 1999) guiding bacterium to swim toward nutrients (chemoeffectors/chemoattractants) adsorbed

to a surface followed by flagellar attachment to initiate the biofilm formation (Stelmack et al., 1999). By

linking chemoreceptor proteins to environmental chemoeffectors, researchers can test environmental

changes to modify microbe behavior in-turn biofilm formation and track proteomic and genomic changes

that could lead to future studies.

Biosensor Selection Criteria

Biosensors are devices or probes that integrates a biological element with an electronic component to

generate a measurable signal. Naresh and Lee outlined six key characteristics of biosensors (Naresh &

Lee, 2021). Table 1 explains these characteristics and how CREID can assist biosensor developers.

Table 1. Biosensor characteristics and how CREID assists biosensor developers match them.		
Characteristic	Definition	How CREID Assists
Selectivity	Important for selecting a chemoreceptor or effector that is integrated into the biosensor. Receptors/effectors attached to a biosensor can detect the analyte of interest in a sample, even if the sample has impurities.	Researchers can link bioreceptors (chemoreceptors) of interest to analytes (chemoeffectors) and vice versa.
Sensitivity	"The minimum amount of analyte that can be correctly detected/identified in a minimum number of steps and in low concentrations (ng/mL or fg/mL) to verify the existence of analyte traces in the sample."	Researchers can find relevant information to synthesize analytes more easily and quickly. CREID can also be used for computational chemistry and retrosynthesis via access to more available data without the need for manual searching.
Linearity	"Contributes to the accuracy of the measured results. The higher the linearity (straight line), the higher the substrate concentration detection."	Populating CREID with new receptor and effector discoveries can improve accuracy through finding the most reliable interaction sets.

Response Time	The time needed to obtain a majority of the targeted result.	Users can leverage CREID in their experimental design to consider ideal environmental conditions for their protein reactions of choice.
Reproducibility	"Characterized by precision (similar output when the sample is measured more than once) and accuracy (capability of a sensor to generate a mean value closer to the actual value when the sample is measured every time). It is the ability of the biosensor to produce identical results whenever the same sample is measured more than once."	Populating CREID with new receptor and effector discoveries can improve accuracy and precision through finding the most reliable interaction pairs/sets.
Stability	Important when continuous monitoring is required. Stability is how vulnerable the biosensor and its components are to environmental disturbances. Stability can be affected by the binding energy of the chemoreceptor to a chemoeffector, and receptor/effector degradation over the biosensor's life.	Data and text mining in CREID can find the best substrates and materials to bind receptors and effectors in place without being washed away by other compounds. Users may find chemoreceptors, effectors, and substrates with the highest molecular binding energies to maximize stability and biosensor life.
Source: Naresh & Lee, 2021		

Data Selection

Data of chemosensory proteins as retrieved from databases: PubMed, Genbank (Benson et al., 2005), Uniprot (UniProt, 2019), Entrez (Maglott et al., 2011) and PROSITE (de Castro et al., 2006). Manual-based search extraction and curation were performed to fully account for chemosensory related proteins which could have previously been submitted as chemoeffector protein, not defined as a chemoattractant or chemorepellent. Confirmation steps were also used to remove any duplicates or low confidence/ predicted protein data. Retrieved data was transformed to structured data that can be searchable and downloaded in a CSV format. Data will also be available for download via API and other web applications.

Database Organization

The focus of the CREID database (Figure 3.) is at the protein level that can be searched by microbial taxa, interaction type, and accession ID for easy data parsing and extraction. The database contains URLs for protein information in Uniprot, ligand binding domain information in the RCSB protein database, chemical information in PubChem, and the article containing the chemoreceptor-effector interaction discovery (Figure 4.)(Figure 5.). Uniprot was also used to confirm protein domains of suspected chemosensory associated proteins. Network visualization option can connect specific receptors to the STRING (Szklarczyk et al., 2019) to show protein-protein interaction network map and connect specific effectors to the STITCH (Kuhn M et al., 2008) to show protein-chemical interaction network map.

Concluding Remarks

CREID is a free webserver and catalogue that contains known protein data for chemosensory proteins that are associated with chemotaxis of various bacterial taxa. It was designed to be used as a tool to collect, organize, and aid in understanding the complexities of chemosensory pathway interactions in different environments that anyone could use and target for future research.

Acknowledgements & Funding

Research reported in this publication was supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number P20GM103443. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. VP and TH are both first co-authors of this manuscript.

Declaration of Interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be constructed as potential conflicts of interest.

References:

Aaronson, S. (1978). Excretion of organic matter by phytoplankton in vitro. *Limnology and Oceanography*, *23*(4), 838.

Armitage, J. (1997). Behavioural responses of bacteria to light and oxygen. *Archives of Microbiology*, *168*, 249–261.

Benson, D., Karsch-Mizrachi, I., Lipman, D., Ostell, J., & Wheeler, D. (2005). GenBank. *Nucleic Acids Res*, *33*. https://doi.org/10.1093/nar/gki063

Bi, S., & Lai, L. (2015a). Bacterial chemoreceptors and chemoeffectors. In *Cellular and Molecular Life Sciences* (Vol. 72, Issue 4, pp. 691–708). Birkhauser Verlag AG. https://doi.org/10.1007/s00018-014-1770-5

Bi, S., & Lai, L. (2015b). Bacterial chemoreceptors and chemoeffectors. In *Cellular and Molecular Life Sciences* (Vol. 72, Issue 4, pp. 691–708). Birkhauser Verlag AG. https://doi.org/10.1007/s00018-014-1770-5

Blackburn, N., Azam, F., & Hagström, Å. (1997). Spatially explicit simulations of a microbial food web. *Limnology and Oceanography*, *42*(4), 613–622. https://doi.org/10.4319/lo.1997.42.4.0613

Boyeldieu, A., Ali Chaouche, A., Ba, M., Honoré, F. A., Méjean, V., & Jourlin-Castelli, C. (2020). The phosphorylated regulator of chemotaxis is crucial throughout biofilm biogenesis in Shewanella oneidensis. *Npj Biofilms and Microbiomes*, *6*(1). https://doi.org/10.1038/s41522-020-00165-5

Brumley, D. R., Carrara, F., Hein, A. M., Hagstrom, G. I., Levin, S. A., & Stocker, R. (2020). Cutting Through the Noise: Bacterial Chemotaxis in Marine Microenvironments. *Frontiers in Marine Science*, *7*. https://doi.org/10.3389/fmars.2020.00527

de Castro, E., Sigrist, C. J. A., Gattiker, A., Bulliard, V., Langendijk-Genevaux, P. S., Gasteiger, E., Bairoch, A., & Hulo, N. (2006). ScanProsite: Detection of PROSITE signature matches and ProRule-associated functional and structural residues in proteins. *Nucleic Acids Research*, *34*(WEB. SERV. ISS.). https://doi.org/10.1093/nar/gkl124

Fogg, G. E. (1977). Excretion of organic matter by phytoplankton. In *Limnology and Oceanography* (Vol. 22, Issue 3, pp. 576–577). https://doi.org/10.4319/lo.1977.22.3.0576

Grishanin, R. N., Gauden, D. E., & Armitage, J. P. (1997). Photoresponses in Rhodobacter sphaeroides: Role of Photosynthetic Electron Transport. In *JOURNAL OF BACTERIOLOGY* (Vol. 179, Issue 1). https://journals.asm.org/journal/jb

Hellebust, J. A. (1965). EXCRETION OF SOME ORGANIC COMPOUNDS BY MARINE PHYTOPLANKTON. *Limnology and Oceanography*, *10*(2), 192–206. https://doi.org/10.4319/lo.1965.10.2.0192

Huang, Z., Wang, Y. H., Zhu, H. Z., Andrianova, E. P., Jiang, C. Y., Li, D., Ma, L., Feng, J., Liu, Z. P., Xiang, H., Zhulin, I. B., & Liu, S. J. (2019). Cross talk between chemosensory pathways that modulate chemotaxis and biofilm formation. *MBio*, *10*(1). https://doi.org/10.1128/mBio.02876-18

Jackson, G. (1980). Phytoplankton growth and Zooplankton grazing in oligotrophic oceans. *Nature*, *284*(3).

Jones, A. K., & Cannon, R. C. (1986). The release of micro-algal photosynthate and associated bacterial uptake and heterotrophic growth. *British Phycological Journal*, *21*(4), 341–358. https://doi.org/10.1080/00071618600650421

Keegstra, J. M., Carrara, F., & Stocker, R. (2022). The ecological roles of bacterial chemotaxis. In *Nature Reviews Microbiology* (Vol. 20, Issue 8, pp. 491–504). Nature Research. https://doi.org/10.1038/s41579-022-00709-w

Maglott, D., Ostell, J., Pruitt, K. D., & Tatusova, T. (2011). Entrez gene: Gene-centered information at NCBI. *Nucleic Acids Research*, *39*(SUPPL. 1). https://doi.org/10.1093/nar/gkq1237

McCarthy, J., & Goldman, J. (1979). Nitrogenous nutrition of marine phytoplankton in nutrient-depleted waters. *Science*, *203*(4381), 670–672.

Naresh, Varnakavi., & Lee, N. (2021). A Review on Biosensors and Recent Development of Nanostructured Materials-Enabled Biosensors. *Sensors*, *21*(4), 1109. https://doi.org/10.3390/s21041109

O'Toole, G. A., & Kolter, R. (1998). Flagellar and twitching motility are necessary for Pseudomonas aeruginosa biofilm development. *Molecular Microbiology*, *30*(2), 295–304. https://doi.org/10.1046/j.1365-2958.1998.01062.x

Pratt, L. A., & Kolter, R. (1998). Genetic analysis of Escherichia coli biofilm formation: Roles of flagella, motility, chemotaxis and type I pili. *Molecular Microbiology*, *30*(2), 285–293. https://doi.org/10.1046/j.1365-2958.1998.01061.x

Prigent-Combaret, C., Vidal, O., Dorel, C., & Lejeune, P. (1999). Abiotic Surface Sensing and Biofilm-Dependent Regulation of Gene Expression in Escherichia coli. In *JOURNAL OF BACTERIOLOGY* (Vol. 181, Issue 19). https://journals.asm.org/journal/jb

Raina, J. B., Giardina, M., Brumley, D. R., Clode, P. L., Pernice, M., Guagliardo, P., Bougoure, J., Mendis, H., Smriga, S., Sonnenschein, E. C., Ullrich, M. S., Stocker, R., & Seymour, J. R. (2023). Chemotaxis increases metabolic exchanges between marine picophytoplankton and heterotrophic bacteria. *Nature Microbiology*, *8*(3), 510–521. https://doi.org/10.1038/s41564-023-01327-9

Sharp, J. H. (1977). Excretion of organic matter by marine phytoplankton: Do healthy cells do it? *Limnology and Oceanography*, *22*(3), 381–399. https://doi.org/10.4319/lo.1977.22.3.0381

Shioi, J., Dang, C. V, & Taylor, B. L. (1987). Oxygen as Attractant and Repellent in Bacterial Chemotaxis. In *JOURNAL OF BACTERIOLOGY* (Vol. 169, Issue 7). https://journals.asm.org/journal/jb

Smriga, S., Fernandez, V. I., Mitchell, J. G., & Stocker, R. (2016). Chemotaxis toward phytoplankton drives organic matter partitioning among marine bacteria. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(6), 1576–1581. https://doi.org/10.1073/pnas.1512307113

Stelmack, P. L., Gray, M. R., & Pickard, M. A. (1999). Bacterial Adhesion to Soil Contaminants in the Presence of Surfactants. In *APPLIED AND ENVIRONMENTAL MICROBIOLOGY* (Vol. 65, Issue 1). https://journals.asm.org/journal/aem

Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva, N. T., Morris, J. H., Bork, P., Jensen, L. J., & Von Mering, C. (2019). STRING v11: Protein-protein association

networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Research*, *47*(D1), D607–D613. https://doi.org/10.1093/nar/gky1131

Taylor, B. L. (1983). How do bacteria find the optimal concentration of oxygen? *Trends in Biochemical Sciences*, *8*(12), 438–441.

Taylor, B. L., & Zhulin, I. B. (1998). In search of higher energy: Metabolism-dependent behaviour in bacteria. In *Molecular Microbiology* (Vol. 28, Issue 4, pp. 683–690). https://doi.org/10.1046/j.1365-2958.1998.00835.x

UniProt, C. (2019). UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res*, *47*(D1), D506–D515. https://doi.org/10.1093/nar/gky1049

Wang, H., Zhang, M., Xu, Y., Zong, R., Xu, N., & Guo, M. (2021). Agrobacterium fabrum atu0526-Encoding Protein Is the Only Chemoreceptor That Regulates Chemoattraction toward the Broad Antibacterial Agent Formic Acid. *Biology*, *10*(12), 1345. https://doi.org/10.3390/biology10121345

Watnick, P. I., & Kolter, R. (1999). Steps in the development of a Vibrio cholerae El Tor biofilm. *Molecular Microbiology*, *34*(3), 586–595. https://doi.org/10.1046/j.1365-2958.1999.01624.x

Kuhn M, von Mering C, Campillos M, Jensen LJ, Bork P. STITCH: interaction networks of chemicals and proteins. Nucleic Acids Res. 2008 Jan;36(Database issue):D684-8. doi: 10.1093/nar/gkm795. Epub 2007 Dec 15. PMID: 18084021; PMCID: PMC2238848.

Bacterial Phospholipid Bilayer Membrane

CREID- ChemoReceptor-Effector Interaction Database

CREID

A database for ChemoReceptor-Effector Interactions

Enter search keyword | 🔍 Search

CREID Dataset Collection:

2020	**508**	**176**
CRE Interaction Table	Chemoreceptors	Chemoeffectors

Visualize the Network Map

Download the dataset

It has 2020 results:

Export CSV

☑ ☐	Id	Accession ID	Chemoreceptor ID	Chemoreceptor Name	Chemoreceptor URL	Chemoreceptor Amino Acid Chain Length	Chemoreceptor Amino Acid Sequence	Microbe Taxa	Microbe Gram Pos Neg
☐	1	Escher00001		Tar	↗			Escherichia coli	(-)
☐	2	Escher00002		Tar	↗			Escherichia coli	(-)

Enter search keyword

🔍 Search

CRE Interaction Table | Chemoreceptors | Chemoeffectors

It has 508 results:

⤓ Export CSV

☑ ☐	Id ⇅	Chemoreceptor ID ⇅	Chemoreceptor Name ⇅	Chemoreceptor URL ⇅	Microbe Taxa ⇅	Chemoreceptor Amino Acid Chain Length ⇅	Chemoreceptor Amino Acid Sequence ⇅
☐	11		CZB domain-containing protein [Citr...		Citrobacter amalonaticus		MLTQEQHHIIMEHMRHSMMNKAVQVHYTAEKCFRL...
☐	12		CZB domain-containing protein [Citr...		Citrobacter braakii		MVPDNTKKLQYIYRLQQSKLVNISHLTENLGNIQE...
☐	13		CZB domain-containing protein [Citr...		Citrobacter cronae		MILTQEQAGAVIAHMRSSLMSKSTHVRYTAEKCFR...
☐	14		CZB domain-containing		Citrobacter europaeus		MKHEILPMENSEMAKFDTVLDQLRIDSSVLMVAFI...

☑ ☐	Id ⬍	Chemoreceptor ID ⬍	Chemoreceptor Name ⬍	Chemoreceptor URL ⬍	Microbe Taxa ⬍	Chemoreceptor Amino Acid Chain Length ⬍	Chemoreceptor Amino Acid Sequence ⬍
			protein [Citr...				
☐	15		CZB domain-containing protein [Citr...		Citrobacter freundii		MIRKIIAKIAKRNNNYDPSTCIIEAEPSVAEQKTE...
☐	16		CZB domain-containing protein [Citr...		Citrobacter sp. L55		MLRDELRDVMVRHIYHSMLTKTIYAEHTINKSLRQ...
☐	17		CZB domain-containing protein [Citr...		Citrobacter sp. RHB35-C21		MIRKIIAKIVKRNSNYNYNTSSEIIEADTPIAEQK...
☐	18		CZB domain-containing protein [Citr...		Citrobacter sp. RHB36-C18		MIRKIIAKIVKRNSNYNYNTSSEIIETDTPIAEQK...
☐	19		CZB domain-containing protein [Citr...		Citrobacter sp. RHBSTW-00017		MIRKIIAKIVKRNSNYNYNTSSDIIEADTPIAEQK...
☐	20		CZB domain-containing protein [Citr...		Citrobacter sp. RHBSTW-00021		MIRKIIAKIVKRNSNYNYNTSSDIIEADTPIAEQK...

1　2　3　4　5　...　51

10 / page

Bacterial Phospholipid Bilayer Membrane

Extracellular space

Attractant

Flagellar motor

Intracellular space

©BiotechfiedBioRender.com

CREID- ChemoReceptor-Effector Interaction Database

	id	Accession ID	Chemoreceptor Name	Chemoreceptor URL	Microbe Taxa	Ligand Binding Domain	Chemoeffector Class	Chemoeffector Name	Chemoeffector URL	Interaction Type	Interaction URL
	1	Tar00001	Tar		Escherichia coli			Methyl-DL-aspartate		attractant	
	2	Tar00002	Tar		Escherichia coli			(+)-Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid		attractant	
	3	Tar00003	Tar		Escherichia coli			L-aspartate		attractant	
	4	Tar00004	Tar		Escherichia coli			Guanidinoacetic acid		attractant	
	5	Tar00005	Tar		Escherichia coli			N-methyl-L-aspartate		attractant	
	6	Tar00006	Tar		Escherichia coli			N-formyl-L-aspartate		attractant	
	7	Tar00007	Tar		Escherichia coli			[2-(nitro-4-oxo-butanoic acid)] amic acid		attractant	
	8	Tar00008	Tar		Escherichia coli			cis-1,2-cyclohexane dicarboxylic acid		attractant	
	9	Tar00009	Tar		Escherichia coli			Phthalic acid		attractant	

Histogram of Microbe taxa