1   **Key role of down-regulated in adenoma (*SLC26A3*) chloride/bicarbonate exchanger in**

2   **linaclotide-stimulated intestinal bicarbonate secretion upon loss of CFTR function**

4   Jessica B. Sarthi[1], Annie M. Trumbull[1], Shayda M. Abazari[1], Vincent van Unen[2], Joshua E.

5   Chan[2], Yanfen Jiang[1], Jesse Gammons[1], Marc O. Anderson[3], Onur Cil[4], Calvin J. Kuo[2], Zachary

6   M. Sellers[1,5,*]

7
8
[1] Department of Pediatrics, Division of Gastroenterology, Hepatology, and Nutrition, Stanford

University, Palo Alto, CA, USA

[2] Department of Medicine, Division of Hematology, Stanford University, Palo Alto, CA, USA

[3] Department of Chemistry and Biochemistry, San Francisco State University, San Francisco,

CA, USA

[4] Department of Pediatrics, University of California San Francisco, San Francisco, CA, USA

[5] Sellers Research and Clinical Development, LLC, Newark, CA, USA

*Corresponding Author: Zachary M. Sellers, MD, PhD. Division of Pediatric Gastroenterology,

Hepatology, and Nutrition, 1701 Page Mill Road, Palo Alto, CA, 94304, USA. E-mail:

zsellers@stanford.edu. Telephone: 650-497-9326.

20
21
22   **Conflicts of Interest:**
23
24   JBS, AMT, SMA, VVU, JEC, YJ, JG, MOA, OC, CJK: The authors have declared no conflict of

25   interest. ZMS: Received financial compensation from AbbVie for professional consultation

26   unrelated to linaclotide. AbbVie did not pay for, contribute, have any special access to data, or

27   have any involvement in any aspect of the work. Also, received consulting fees from 4D

28   Molecular Therapeutics, which had no involvement in the work or access to any data.

ABSTRACT

Duodenal bicarbonate secretion is critical to epithelial protection, nutrient digestion/absorption

and is impaired in cystic fibrosis (CF). We examined if linaclotide, typically used to treat

constipation, may also stimulate duodenal bicarbonate secretion. Bicarbonate secretion was

measured in vivo and in vitro using mouse and human duodenum (biopsies and enteroids). Ion

transporter localization was identified with confocal microscopy and de novo analysis of human

duodenal single cell RNA sequencing (sc-RNAseq) datasets was performed. Linaclotide

increased bicarbonate secretion in mouse and human duodenum in the absence of CFTR

expression (*Cftr* knockout mice) or function (CFTR$_{inh}$-172). NHE3 inhibition contributed to a

portion of this response. Linaclotide-stimulated bicarbonate secretion was eliminated by down-

regulated in adenoma (DRA, SLC26A3) inhibition during loss of CFTR activity. Sc-RNAseq

identified that 70% of villus cells expressed *SLC26A3*, but not *CFTR*, mRNA. Loss of CFTR

activity and linaclotide increased apical brush border expression of DRA in non-CF and CF

differentiated enteroids. These data provide further insights into the action of linaclotide and

how DRA may compensate for loss of CFTR in regulating luminal pH. Linaclotide may be a

useful therapy for CF individuals with impaired bicarbonate secretion.

INTRODUCTION

Proximal duodenal bicarbonate secretion, together with pancreatic bicarbonate secretion, is integral to preventing intestinal mucosal injury from gastric secretions and establishing a non-acidic intraluminal environment for activation of digestive enzymes. This importance is highlighted by cystic fibrosis (CF) intestinal disease and malnutrition, where decreased bicarbonate secretion is central to disease pathogenesis (1, 2). Duodenal bicarbonate secretion is highly regulated, driven by luminal acid and neurohormonal signaling, ultimately resulting in coordinated transepithelial bicarbonate secretion by the cystic fibrosis transmembrane conductance regulator (CFTR), chloride/bicarbonate exchange (e.g., down-regulated in adenoma, DRA; putative anion transporter-1, PAT-1), and/or decreased proton transport through sodium/hydrogen exchange (e.g. NHE3)(3). Despite its physiologic importance, there are limited therapies targeting duodenal bicarbonate secretion.

Linaclotide is a Food and Drug Administration (FDA)-approved medication for the treatment of constipation-type irritable bowel syndrome and chronic idiopathic constipation whose effect relies on its structural similarity to the heat-stable enterotoxin of *Escherichia coli* (STa). Both STa and linaclotide bind guanylyl cyclase C (GC-C) receptors on the apical surface of enterocytes and cause increases in CFTR-dependent chloride secretion and inhibition of NHE3-mediated sodium absorption (4-6). Previously, we identified that STa stimulates duodenal bicarbonate secretion through CFTR and CFTR-independent means (7, 8). Linaclotide has been investigated as a potential therapy for CF-related constipation (6, 9, 10), however, it is unclear if linaclotide stimulates duodenal bicarbonate secretion and if linaclotide could improve impaired intraluminal pH in CF patients by stimulating CFTR-independent bicarbonate secretion.

We undertook a series of experiments utilizing in vivo and in vitro bicarbonate secretion measurements in mice and humans to examine the effect of linaclotide on duodenal bicarbonate secretion. We coupled these experiments with scRNA-seq analysis of acid-based transporters in the human duodenum and confocal microscopy to characterize the brush border expression

of the DRA chloride/bicarbonate exchanger in human duodenum in the presence or absence of

CFTR activity and/or expression.

RESULTS

<u>Linaclotide stimulates bicarbonate secretion in mouse and human duodenum.</u> To determine

linaclotide's ability to stimulate intestinal bicarbonate secretion, we examined its effect on the

duodenal segment of the small intestine, given the robust expression of acid-base transporters

in this segment that function to neutralize the highly acidic gastric contents it encounters. We

first examined the effect of in vivo intestinal perfusion of linaclotide on duodenal bicarbonate

secretion. Linaclotide stimulated sustained increases in bicarbonate secretion at 10^{-7} M and

10^{-5} M (P<0.001 and P=0.004, n=9), with small and transient increases in bicarbonate secretion

over baseline secretory rates at 10^{-9} M (P=0.033, n=9). There was no significant difference

between 10^{-7} M or 10^{-5} M (P=0.926) (Figure 1A, B). These bicarbonate secretory rates were

similar to those observed with STa (Figure 1B, 10^{-9} M: P=0.491 and 10^{-7} M: 0.792, respectively,

n=11). Linaclotide did not stimulate net increases in luminal duodenal fluid secretion (10^{-9} M:

0.01 $\pm$ 0.05, 10^{-7} M: -0.05 $\pm$ 0.06, 10^{-5} M: -0.04 $\pm$ 0.05 ΔmL/cm•h, P=0.856, 0.445, 0.367,

respectively, n=8), similar to what we have previously reported for STa in the duodenum (7). As

linaclotide can decrease proton secretion by inhibiting NHE3 (6, 9), we examined if linaclotide

actively stimulates bicarbonate secretion, or purely inhibits baseline proton secretion to raise

intraluminal pH. Using the same in vivo technique, perfusion with S3226 (10^{-5} M), a selective

inhibitor of NHE3, caused a significant increase in baseline bicarbonate secretion (3.78 $\pm$ 0.73

vs. 6.70 $\pm$ 0.68 μmol/cm•h, n=9-10, P=0.016). In the presence of S3226, linaclotide caused

further increases in bicarbonate secretion (10^{-9} M: 2.22 $\pm$ 0.87, 10^{-7} M: 2.53 $\pm$ 0.82, 10^{-5} M: 2.38

$\pm$ 1.32 $\Delta\mu$mol/cm•h, n=10, P=0.075, P=0.032, P=0.237, respectively)(Figure 1C). In the

presence of S3226, linaclotide-stimulated duodenal bicarbonate secretion was ~1/3 of the

response without S3226 (10^{-7} M: $37.9 \pm 12.3\%$, 10^{-5} M: $35.6 \pm 19.8\%$). Thus, in mouse

duodenum, linaclotide alters duodenal luminal pH through a combination of inhibiting NHE3-

mediating proton secretion and stimulating bicarbonate secretion.

To further characterize linaclotide-stimulated bicarbonate secretion, we performed

combined I_{sc} and bicarbonate secretory measurements in seromuscular-stripped mouse

duodenum mounted in Ussing chambers. As seen in Figure 1D-G, this in vitro method showed

similar results as the in vivo method. Linaclotide stimulated significant increases in duodenal

bicarbonate secretion at 10^{-7} M (P=0.025, n=9) and 10^{-5} M (P=0.047, n=11), with no difference

between them (P=0.875). Short-circuit current analysis revealed that linaclotide stimulated

significant increases in I_{sc} at 10^{-7} M (P<0.001, n=9) and 10^{-5} M (P<0.001, n=11) (Figure 1F, G).

Bicarbonate secretion and I_{sc} were accompanied by small, but significant (P<0.05) changes in

transepithelial resistance (Figure 1H).

We next examined the potential for linaclotide to stimulate bicarbonate secretion in

human duodenum using endoscopically obtained duodenal biopsies from subjects without

known acid-base disturbances or histologic evidence of duodenal disease. Biopsies were

mounted in Ussing chambers and combined bicarbonate secretion and I_{sc} measurements were

performed, similar to Figure 1D-H. As seen in Figure 1I, 10^{-7} M and 10^{-5} M linaclotide stimulated

significant increases in human duodenal bicarbonate secretion (P=0.006, n=11 and P=0.004,

n=10, respectively), at similar magnitudes previously reported for STa in human duodenal

biopsies (11). In contrast to mice, linaclotide did not stimulate a significant change in I_{sc} in

human biopsies (10^{-7} M: P=0.071, n=11; 10^{-5} M: P=0.186, n=10) (Figure 1J) or transepithelial

resistance (Figure 1J, K). Thus, linaclotide stimulates electrogenic bicarbonate secretion in

mouse duodenum and electroneutral bicarbonate secretion in human duodenum.

<u>Linaclotide-stimulated duodenal bicarbonate secretion occurs independent of CFTR expression</u>

<u>or function.</u> Next, given our prior findings that STa can stimulate CFTR-independent duodenal

bicarbonate secretion in mice (7), we studied linaclotide's dependence on CFTR for bicarbonate

secretion. We first repeated the in vivo bicarbonate secretion experiments by perfusing the

duodenum with $CFTR_{inh}$-172, a specific CFTR inhibitor.(12) Linaclotide continued to stimulate

significant bicarbonate secretory responses at 10^{-7} M and 10^{-5} M (P=0.008 and P=0.018,

respectively, n=12), but not 10^{-9} M (P=0.198, n=12), in the presence of $CFTR_{inh}$-172 (2 x 10^{-5} M).

In comparison, in vivo, $CFTR_{inh}$-172 inhibited forskolin-stimulated (10^{-4} M) bicarbonate secretion

by 65.1 $\pm$ 12.5% (P=0.022), preventing significant increases over baseline (P=0.778, n=6-9)

(Supplementary Figure 1A and B). In vitro, sequential addition of CFTR inhibitors with different

mechanisms of action, GlyH-101 (10^{-5} M, mucosal) and glibenclamide (3 x 10^{-4} M, mucosal),

following $CFTR_{inh}$-172 (2 x 10^{-5} M, serosal), resulted in no further increase in the inhibition of

forskolin-stimulated I_{sc} (0.2% and -0.9 %, respectively). To expand upon these pharmacologic

experiments, we performed in vitro Ussing chamber experiments with $CFTR^{tm1UNC}$ knockout

(KO) mice, which have global KO of the *Cftr* gene (13). Given the similar responses between

10^{-7} M and 10^{-5} M in prior experiments, for these studies we specifically focused on 10^{-7} M

linaclotide. We found that linaclotide (10^{-7} M) stimulated significant increases in duodenal

bicarbonate secretion in *Cftr* KO mice (P=0.028 vs. baseline, n=7). These magnitudes were

similar to wildtype (WT) mice (P=0.728) (Figure 2B).   In contrast to mice expressing CFTR,

linaclotide stimulated minimal change in I_{sc} in *Cftr* KO mice (0.77 $\pm$ 0.31 $\Delta\mu A/cm^{2}$, P=0.049, n=8)

with no change in transepithelial resistance (P=0.396) (Figure 2C, D), suggesting that in the

absence of CFTR, linaclotide stimulates electroneutral bicarbonate secretion. To determine if

these findings hold true in humans, we pre-treated human endoscopic duodenal biopsies with

$CFTR_{inh}$-172, which inhibits forskolin-stimulated I_{sc} in human duodenal enteroids

(14)(Supplementary Figure 1C), and then performed repeat Ussing chamber measurements.

Linaclotide (10^{-7} M) continued to stimulate significant increases in duodenal bicarbonate

secretion in biopsies pre-treated with $CFTR_{inh}$-172 (2 x 10^{-5} M, serosal, P=0.024, n=14), at a

magnitude that was comparable to biopsies without $CFTR_{inh}$-172 treatment (P=0.644) (Figure

2E). There was no difference in linaclotide-stimulated I_{sc} with or without $CFTR_{inh}$-172 treatment

(0.27 ± 0.11 vs. 0.13 ± 0.04 $\Delta\mu Eq/cm^2 \bullet h$, P=0.239, n=11-14) (Figure 2F). Transepithelial

resistance in the presence of $CFTR_{inh}$-172 was unchanged before or after linaclotide stimulation

(P=0.889, n=14) (Figure 2G).

<u>Linaclotide-stimulated duodenal bicarbonate secretion upon loss of CFTR function is mediated</u>

<u>by the DRA chloride/bicarbonate exchanger.</u> To investigate the potential mechanism whereby

linaclotide increases duodenal bicarbonate secretion independent of CFTR, we investigated the

effect of DRA_{inh}-A250 (selective inhibitor of DRA), DIDS (DRA-insensitive, non-selective anion

exchanger inhibitor), and S3226 (selective inhibitor of NHE3) on duodenal bicarbonate

secretion. Using our in vivo model, in separate experiments, each of these drugs were

combined with $CFTR_{inh}$-172 and perfused prior to stimulation with linaclotide. DRA_{inh}-A250 (10^{-5}

M) caused a 67 ± 8% reduction in CFTR-independent linaclotide-stimulated duodenal

bicarbonate secretion at 10^{-7} M linaclotide (P=0.044, n=10), and 73 ± 6% inhibition at 10^{-5} M

linaclotide (P=0.038, n=10) (Figure 3A-C). In contrast, DIDS (2 x 10^{-4} M) failed to significantly

impact CFTR-independent linaclotide-stimulated duodenal bicarbonate secretion (10^{-7} M: -7 ±

14%, P=0.993, n=7; 10^{-5} M: -19 ± 14%, P=0.893, n=7). S3226 (10^{-5} M) reduced CFTR-

independent linaclotide-stimulated duodenal bicarbonate secretion by 45 ± 19% (10^{-7} M) and 58

± 16% (10^{-5} M); however, neither of these were statistically significant (P=0.254, P=0.136,

respectively, n=9) (Figure 3A, B). Therefore, it appears that DRA is the primary source of

bicarbonate transport for CFTR-independent linaclotide-stimulated duodenal bicarbonate

secretion.

To determine if DRA's critical role in linaclotide-stimulated duodenal bicarbonate

secretion is unique to when there is functional loss of CFTR, we examined the role of DRA in

mouse duodenal bicarbonate secretion measured in vitro. For these experiments, DRA$_{inh}$-A270,

a near identical chemical compound as DRA$_{inh}$-A250 (Br$^-$ to I$^-$ substitution) with similar DRA

specificity and inhibitory properties as DRA$_{inh}$-A250 at 10^{-5} M (15), was used. DRA$_{inh}$-A270 (10^{-5}

M) had no significant inhibitory impact on linaclotide (10^{-7} M)-stimulated duodenal bicarbonate

secretion (P=0.813, n=5-8) or I$_{sc}$ (P=0.991, n=5-8) in the presence of CFTR function (Figure 3D

and E). In vivo testing also showed no inhibitory effect of DRA$_{inh}$-A270 on linaclotide-stimulated

bicarbonate secretion without CFTR inhibition (DMSO + linaclotide 10^{-7} M: 4.3 $\pm$ 1.1 vs. DRA$_{inh}$-

A270 + linaclotide 10^{-7} M: 3.5 $\pm$ 0.3 μmol.cm^{-1}.h^{-1}, n=4 each, P=0.542). To confirm lack of effect

of DRA inhibition under these conditions, we performed additional in vitro experiments using

DRA$_{inh}$-4a, a DRA specific inhibitor that is structurally and mechanistically distinct from DRA$_{inh}$-

A250 and DRA$_{inh}$-A270 (16). These experiments showed similar results as DRA$_{inh}$-A270; no

inhibition of linaclotide-stimulated duodenal bicarbonate secretion (P=0.899, n=8-16) or I$_{sc}$

(P=0.870, n=8-16) in the presence of CFTR function. Thus, CFTR is sufficient for linaclotide-

stimulated bicarbonate secretion, but upon loss of CFTR function, DRA appears to compensate

for its loss.

<u>DRA expression in human duodenum.</u> DRA and PAT-1 are the primary chloride/bicarbonate

exchangers expressed in mouse intestine, with PAT-1 showing dominant RNA expression in the

small intestine and DRA RNA being more abundant in the colon.(17) There is limited data

regarding DRA expression in human duodenum.(18, 19) We capitalized on two previously

published human duodenum scRNA-seq datasets(20, 21) to examine DRA mRNA expression in

human duodenum. The Elmentaite et al. dataset(21) contained 5,944 cells with an average of

4,164 reads per cell from 5 healthy adult duodenal samples, allowing for high confidence in cell-

type separation. The Busslinger et al. dataset(20) contained 702 crypt cells and 923 villus cells

from 2 adult duodenal samples, with an average of 10,381 reads per cell. Data from one crypt

sample ("Crypt 1") was excluded due to very low sequence reads (2,088 per cell). We verified

accurate classification into crypt and villus cells by examining *LGR5* expression and proliferation

markers *MKI67* and *PCNA*, which are generally restricted to the crypt region. These were all

more abundant in the crypt group, with *LGR5* being exclusively expressed in the crypt, but not

villus, group (Supplementary Figure 2A-C). Using these two datasets, we examined expression

of the key acid-base transporters in the duodenum, DRA (*SLC26A3*), PAT-1 (*SLC26A6*), *CFTR*,

and NHE3 (*SLC9A3*). Our analysis showed that *SLC26A3* mRNA is predominantly expressed

in enterocytes (Figure. 4A), in both the crypt and villus groups (Figure 4E), with higher

expression and expressed in more cells than *SLC26A6*, *CFTR*, or *SLC9A3* (Figures 4A-D and

E-H). Of note, Crypt 2 had lower *SLC26A3* expression than Crypt 3, which correlated with

higher *CFTR, MKI67, PCNA* expression, more *LGR5*+ cells, and less *SLC9A3*+ cells,

suggesting Crypt 2 contained more cells deeper within the crypts than Crypt 3 (Figures 4E, G, H

and Supplementary Figure 2). We confirmed DRA protein expression in human duodenal

biopsies (n=3) using immunofluorescence imaging. The majority of DRA is localized to the

intracellular apical region, with a smaller fraction present at the apical brush border (Figures 4I-

K).

Given DRA's functional independence from CFTR in mediating linaclotide-stimulated duodenal

bicarbonate secretion, we quantified the proportion of *SLC26A3*-expressing cells that co-

expressed *CFTR*. In the Elmentaite et al. dataset, 88.8% of *SLC26A3*-expressing enterocytes

co-expressed *SLC26A3* and *CFTR* (Figure 5a). Using the Busslinger et al. dataset to examine

crypt vs. villus localization, 63.0% of *SLC26A3* positive crypt cells co-expressed *SLC26A3* and

CFTR, whereas in the villus cells, only 29.6% did so (Figures 5B and C). These findings were

not ubiquitous for all chloride/bicarbonate exchangers. *SLC26A6* and *CFTR* co-expression was

relatively low in enterocytes (16.7%) (Supplementary Figure 3A), but examination of the crypt

and villus cell types showed a higher percentage of *CFTR* co-expression with *SLC26A6* than

SLC26A3, with 81.8% of crypt cells and 42.9% of villus cells co-expressing *SLC26A6* and *CFTR*

(Supplementary Figures 3B and C). In the Elmentaite et al. dataset, *SLC9A3* and *CFTR* were

co-expressed in 14.3% of enterocytes, similar to *SLC26A6* (Supplementary Figure 3D). The

Busslinger et al. dataset possessed insufficient numbers of *SLC9A3*-expressing cells to analyze

SLC9A3 and *CFTR* co-expression in crypts and villi.

<u>Linaclotide alters the apical localization of DRA.</u> Ion channel activity can be regulated by

altering membrane expression of channels. We examined if linaclotide alters DRA activity by

changing its apical brush border expression. For these experiments we utilized human derived

three-dimensional duodenal enteroids. Given that the linaclotide receptor GC-C is expressed at

the apical surface, we generated reverse polarity apical-out duodenal enteroids, which allow

facile access to the apical membrane via the enteroid exterior (Supplementary Figures 4A and

B).(22) Using qPCR, we verified that apical-out enteroids express *CFTR*, *SLC26A3, SLC26A6,*

SLC9A3, and GUCY2C in undifferentiated and 3-day differentiated enteroids (Supplementary

Figures 4C and D. 4). As expected, *LGR5* was decreased in differentiated apical-out enteroids.

Using this model of human duodenum, we incubated apical-out differentiated enteroids with

linaclotide (10^{-7} M, 40 minutes) or vehicle control (water, 40 minutes) and performed confocal

immunofluorescence to localize DRA. As seen in Figures 6A-D, linaclotide significantly

increased apical brush border expression of DRA (n=12-16). In contrast, when we performed

the same experiment with NHE3 localization, we found no change in the apical brush border

expression of NHE3 upon linaclotide stimulation (Supplementary Figures 5A-D). We verified

that myosin VI (*MYO6*), a key protein in NHE3 trafficking,(23) mRNA was present in apical-out

enteroids. *MYO6* mRNA was expressed at similar levels in this model as many of the other

transporters we examined (Supplementary Figures 4C and D). To determine if linaclotide also

stimulates DRA localization upon lack of functional CFTR, we repeated the experiments in the

presence of $CFTR_{inh}$-172 (2×10^{-5} M, 40 minutes). Compared to vehicle control (DMSO, 40

1. minutes), CFTR inhibition resulted in an increase in the apical expression of DRA (P=0.032,

2. n=26-31)(Figures 6E-H), but not NHE3 (P=0.633, n=16-30) (Supplementary Figures 5E-H). Of

3. note, DMSO alone increased DRA brush border expression compared to water (MFI means $\pm$

4. SEM: 10.94 $\pm$ 1.32 vs. 3.43 $\pm$ 1.03, n=11-15, P<0.001). Linaclotide stimulation in the presence

5. of CFTR$_{inh}$-172 did not alter DRA or NHE3 brush border expression (P= 0.730, n=31; P=0.535,

6. n=29-30, respectively) (Figures 6E-G and I, Supplementary Figures 5E-G and I).

7.   To determine how our CFTR$_{inh}$-172 studies compare to chronic loss of CFTR expression

8. and function, we examined DRA brush border expression in human duodenal enteroids isolated

9. from a CF subject with F508del homozygous *CFTR* variants. Comparison of baseline (water

10. vehicle control, 40 minutes) DRA brush border expression between non-CF and CF enteroids

11. showed CF enteroids had increased DRA expression at the apical brush border (Figures 6J-M,

12. n=8-11, P=0.006). Linaclotide stimulation (10^{-7} M, 40 minutes) increased DRA brush border

13. expression in F508del homozygous CF enteroids, compared to CF enteroids treated with

14. vehicle control (water, 40 minutes)(Figures 6J-L and N, n=8-10, P=0.010). To examine the

15. discrepancies between CFTR$_{inh}$-172 + linaclotide in non-CF enteroids and linaclotide in the CF

16. enteroids, we examined the absolute mean fluorescent intensities (MFI) of each condition. We

17. found that CFTR$_{inh}$-172 alone caused similar increases in absolute DRA brush border

18. expression as linaclotide in non-CF enteroids (with or without CFTR$_{inh}$-172) or CF enteroids

19. (Figure 6O, n=10-31, P=0.676), suggesting that lack of additional DRA membrane expression in

20. non-CF enteroids treated with CFTR$_{inh}$-172 + linaclotide may be due to maximal DRA brush

21. border expression already being achieved upon CFTR$_{inh}$-172 treatment. Taken together with our

22. bicarbonate secretory measurements, these data show that linaclotide can alter DRA activity

23. and/or brush border expression, in the presence or absence of functional CFTR. In contrast,

24. linaclotide appears to modulate NHE3 activity, rather than brush border expression, regardless

25. of CFTR function.

<u>Inhibition of CFTR transiently increases intracellular pH.</u> To investigate why CFTR inhibition

increases DRA brush border expression, we examined the effect of $CFTR_{inh}$-172 on duodenal

enteroid intracellular pH (pH_i). We hypothesized that loss of CFTR activity would lead to

intracellular bicarbonate trapping (resulting in increased pH_i) and DRA may traffic to the apical

brush border as part of the cell's effort to restore bicarbonate transport and normalize the

altered pH_i. Undifferentiated and 3-day differentiated enteroids grown on coverslips and

$CFTR_{inh}$-172-induced (or DMSO) changes in pH_i were monitored with live cell fluorescence

imaging. $CFTR_{inh}$-172 (2×10^{-5} M) resulted in significant increases in pH_i in undifferentiated and

differentiated enteroids, although the slopes and magnitudes of these increases were different

(Figure 7, n=57-89, $P<0.001$), with undifferentiated enteroids showing more rapid and greater

pH_i changes ($P<0.001$).   DMSO (1:1000) had no effect on pH_i (data not shown). Following

$CFTR_{inh}$-172, all enteroids underwent pH_i recovery. Pre-treatment with DRA_{inh}-A250 (10^{-5} M, 5

min) did not alter the $CFTR_{inh}$-172-induced increase in pH_i but did alter the pH_i recovery. In

undifferentiated enteroids, DRA_{inh}-A250 decreased the slope and magnitude of pH_i recovery

(Figures 7A-C, n=31, $P<0.001$). In differentiated enteroids, the magnitude of pH_i recovery was

not different with DRA_{inh}-A250 (Figure 7F, n=37, $P=0.213$), however, the recovery slope was

altered, albeit in a different way than undifferentiated enteroids (Figure 7D and E). Thus,

inhibition of CFTR function causes an increase in pH_i, the recovery of which is at least in part

influenced by DRA activity.

DISCUSSION

Numerous studies in animal models and humans have shown the critical importance of

duodenal bicarbonate secretion in maintaining epithelial integrity and creating the appropriate

intestinal intraluminal pH for digestion. In CF, with genetic loss of *CFTR*, or *Helicobacter pylori*

infection, with down-regulation of CFTR and PAT-1,(24) decreased duodenal mucosal

bicarbonate secretion contributes to malabsorption and duodenal ulcers, respectively. While

eradication of *Helicobacter pylori* with antibiotics may restore bicarbonate function, the most

common way to improve small intestinal pH in CF is through chronic gastric acid suppression.

Ivacaftor, a CFTR potentiator, was shown to improve proximal intestinal pH in CF patients with

the G551D CFTR gating mutation (accounts for 4-5% of *CFTR* disease-causing variants).(25)

While additional CFTR modulators are FDA-approved, it remains unknown whether these

positively impact small intestinal pH. This gap, coupled with lack of potential bicarbonate-

correcting therapy for CF patients ineligible for CFTR modulator therapies (due to genotype or

medical contraindication), has driven us to seek ways to improve intestinal bicarbonate

secretion independent of CFTR. Our current data are in line with prior mouse and human

duodenum studies with STa (7, 11) and a recent study by Tan et el. showing that linaclotide can

stimulate jejunal bicarbonate secretion in mice.(9) Uniquely, we have shown that linaclotide

stimulates both mouse and human bicarbonate secretion in the duodenum, where bicarbonate

secretion is physiologically important. Furthermore, we have provided evidence for the roles (or

lack of) membrane trafficking of DRA and NHE3 in this process and how decreases in CFTR

function may impact cellular pH.

17   Linaclotide represents an attractive target for restoration of deficient duodenal

bicarbonate secretion in CF. Our data in both mice and humans provide greater confidence in

the translatability of our results to clinical care. While we did not study other segments of the

small intestine, linaclotide stimulates jejunal bicarbonate secretion in F508del CF mice.(9)

Braga Emidio et al. identified that linaclotide's half-life in small intestinal fluid was 48

minutes,(26) suggesting that its effect is likely to be most influential in the proximal small

intestine. Comparing linaclotide-stimulated and acid-stimulated duodenal bicarbonate secretion

using the same methodology,(27, 28) the linaclotide effect is about 75-100% of the stimulatory

effect of acid, indicating linaclotide could produce a clinically meaningful impact for patients.

With existing FDA approval, and the use of clinically relevant dosing,(9) our results may provide

immediate application to patients.

In the absence of CFTR, the logical targets for intestinal pH modulation are either 1)

increasing chloride/bicarbonate exchanger activity, and/or 2) decreasing sodium/hydrogen

exchange. Linaclotide increases intraluminal small intestinal fluid in mice by inhibiting NHE3-

dependent sodium absorption.(6, 9) Another byproduct of NHE3 inhibition is a decrease in

proton secretion, which can raise intraluminal pH. Consistent with this, NHE3 inhibition in our

studies "increased bicarbonate secretion," likely by producing less luminal protons to neutralize

secreted bicarbonate anions. In our NHE3 trafficking studies, we did not see a significant

decrease in NHE3 membrane localization with linaclotide treatment. NHE3 activity may be

impacted by shifting along the microvilli,(23) however, our imaging resolution was unable to

determine these potential nanometer shifts. Nonetheless, our in vivo data supports that NHE3

inhibition alone does not account for linaclotide's ability to increase duodenal pH. Our studies

using specific DRA inhibitors and DRA protein trafficking studies in apical-out human enteroids,

indicate that DRA-mediated chloride/bicarbonate exchange is critical to linaclotide stimulated

duodenal bicarbonate secretion upon loss of CFTR activity. This helps explain the lack of

linaclotide-stimulated short-circuit current in *Cftr* KO mice, human biopsies with $CFTR_{inh}$-172,

and prior studies by McHugh et al. in F508del mice.(6) Interestingly, we found that CFTR

inhibition alone caused an increase in DRA membrane trafficking. There was also increased

apical membrane DRA expression in F508del homozygous duodenal enteroids even without

stimulation. In mouse intestinal crypts, Strubberg et al. identified that pH_i was raised in *cftr* KO

mice or wildtype mice treated with $CFTR_{inh}$-172.(29) In fact, ~20 years ago, Kaunitz et al.

proposed the "CF Paradox" whereby diminished bicarbonate transport may increase

intracellular buffering in CF and prevent duodenal ulcer formation despite diminished

bicarbonate secretion.(30) Thus, we hypothesized that upon CFTR blockade (or genetic lack of

CFTR), the rise in pH_i may trigger an increase in DRA expression and protein localization to at

least partially restore bicarbonate secretion and limit alterations in pH_i. Our current pH_i data

with $CFTR_{inh}$-172, supports this hypothesis. Bijvelds et al. reported a nearly 2,000-fold increase

in *DRA* mRNA in a CF ileal biopsy compared to non-CF ileum (n=1 each).(31) The lack of

additional membrane trafficking of DRA by linaclotide in the presence of CFTR inhibition may

indicate that DRA trafficking requires CFTR function or that a ceiling affect for DRA membrane

expression has already been achieved. Based on our current data we favor the latter

interpretation since linaclotide increased DRA trafficking in F508del homozygous CF enteroids

and a comparison of MFIs across conditions showed that while the relative magnitude of DRA

trafficking may differ between vehicle controls/conditions, the absolute magnitudes are similar.

While CFTR inhibition caused similar pH_i changes in undifferentiated (crypt-like) and

differentiated (villus-like) enteroids, how DRA inhibition impacted pH_i recovery was different.

This may be due to increased expression of a larger variety of acid-base transporters in

differentiated enteroids that may also contribute to pH_i normalization.(19) Further study on how

the expression and activities of DRA and other acid-base transporters are altered in CF

intestine, especially in human samples, may identify new potential therapies for CF intestinal

disease.

Prior to our study, our knowledge on the intestinal expression profile of DRA and other

acid-base transporters was largely based on mouse studies. Wang et al. reported high

expression of *Slc26a3* (DRA) RNA in the colon, but negligible RNA expression in the small

intestine (duodenum, jejunum, ileum). Conversely, *Slc26a6* (PAT-1) RNA expression was high

in mouse duodenum, jejunum, and ileum, with minimal in the colon.(17) Yet, *Slc26a3* KO

mouse studies suggest a prominent role for DRA in basal and stimulated bicarbonate transport.

(32, 33) Human studies are sparse. Yin et al. used human duodenal enteroids to examine

SLC26A3, *SLC26A6*, and *SLC9A3* (NHE3) mRNA expression in undifferentiated and terminally

differentiated human duodenal enteroids.(19) Taking advantage of published scRNA-seq

datasets from the human duodenum, we have provided a comparison on the expression of

SLC26A3, SLC26A6, and *SLC9A3*, from human duodenum tissue. *SLC26A3* mRNA was

generally expressed ~3-4x higher than *SLC26A6, CFTR*, and *SLC9A3.* In the *CFTR* high

expressing *BEST4*+ cells, *CFTR* expression levels approached that of *SLC26A3.* Of note,

SLC26A3 was also highly expressed in *BEST4*+ cells, although enterocytes comprise the

majority of *SLC26A3*-expressing cells. These findings, put in context of prior studies, suggest a

more prominent role for DRA in the human duodenum than may have generally been

appreciated from mouse studies alone. Future work examining not only the mRNA expression,

but also protein expression, and especially pertinent to membrane ion transporters, membrane

localization of DRA, PAT-1, and NHE3 across different segments of human intestine may

advance our knowledge regarding the regional roles of these acid-base transporters in health

and disease.

A potential limitation of our study is the general reliance on pharmacologic inhibitors

throughout our experiments.   In *Xenopus levis* oocytes, $CFTR_{inh}$-172 (2×10^{-5} M) inhibited

~70% of human CFTR current in patch-clamp experiments (34). In our experiments, $CFTR_{inh}$-

172 (2×10^{-5} M) significantly inhibited CFTR ion transport in mouse tissue and human enteroids,

with observed results similar to that by Yin et al.(19) Complementary results in *Cftr* KO mice

and CF enteroids provide further confidence in our findings with $CFTR_{inh}$-172. For DRA, we did

not perform experiments in *Slc26a3* KO animals or cells, however, we did use 3 different DRA

inhibitors, from 2 different structural categories and modes of inhibition. While confidence in

pharmacological inhibitors can wane with time and increased use, we believe the consistency of

results across these different inhibitors provides confidence in the interpretation of the data.

In summary, we have identified that linaclotide may have more pleiotropic ion transport

effects beyond CFTR- and NHE3-mediated chloride and sodium transport, respectively. We

have shown, using mice, duodenal biopsies, and human duodenal enteroids, that linaclotide

stimulates bicarbonate secretion, and does so at least in part, through DRA recruitment to the

apical membrane. Our findings that this remains possible independent of CFTR activity and/or

expression has potentially important implications for CF patients, who may benefit from

linaclotide's bicarbonate stimulatory effect, in addition to its inhibitory effect on NHE3, even if

they can't leverage its pro-chloride secretory properties.(6) Furthermore, our identification of

high DRA expression in the human duodenum may spur interest in identifying DRA-activating

drugs targeting proximal small intestinal disease.

METHODS

Chemicals. $CFTR_{inh}$-172 was purchased from Cayman Chemical (Ann Arbor, MI), 4,4'-

Diisothiocyanatostilbene-2,2'-disulfonate (DIDS) and S3226 were purchased from Sigma-Aldrich

(St. Louis, MO). DRA inhibitors (DRA_{inh}-A250, DRA_{inh}-A270, DRA_{inh}-4a) were synthesized and

purified as previously described.(15, 16, 35) Linaclotide was purchased from Santa Cruz

Biotechnology, Inc (Dallas, TX).

Animals. Adult (8-16 week-old) C57BL/6J mice from Jackson Laboratories (Bar Harbor, ME)

were used as wildtype control mice. $CFTR^{tm1Unc}$ knockout (KO) mice were obtained from Case

Western University CF Mouse Models Core Facility and maintained on 50% PEG-3350 with

electrolytes solution in their drinking water with standard chow. Similar number of male and

female mice were used in experiments.

In Vivo Measurement of Bicarbonate Secretion. In vivo measurement of duodenal bicarbonate

secretion was performed using a well-validated technique.(36) Animals were maintained with

free access to food and water for up to 1 hour before the experiment, then fasted from chow.

Mice were anesthetized with oxygen-delivered isoflurane (1-3%) at 1 L/minute via a vaporizer

(Braintree Scientific, Inc, Braintree, Mass). Mouse temperature was monitored by rectal probe

and maintained at 37°C through automated warming using a controlled warming pad (ATCC

2000, World Precision Instruments, Sarasota, FL). Anesthetic plane was assessed by

respiratory rate and toe pinch reflex. Respiratory and heart rates were determined every 15

minutes. Animals could be sustained for more than 3 hours under these experimental

conditions.

To isolate the proximal duodenum, the abdomen was opened by a central vertical incision, and

the proximal 5–10 mm of duodenum, from the pylorus to just proximal to the entry of the

common bile duct, was isolated in situ without compromising vascular supply. A small

polyethylene tube (PE-50) with a distal flange was advanced to the duodenal bulb via the

stomach, and a ligature was secured around the pylorus. A distal intestinal incision was made,

and PE-50 flanged tubing was advanced to just proximal to the entry of the pancreaticobiliary

duct to prevent entry of the pancreatic or biliary secretions and allow for collection of effluent

from isolated segment. The isolated duodenal segment was gently flushed and then

continuously perfused (Harvard Infusion Pump, Harvard Apparatus, South Natick, MA, USA) at

a rate of 0.21 mL/min with 154 mM NaCl (37°C) $\pm$ stimulatory/inhibitor drugs. Effluents from the

isolated segment were visually free of bile and blood throughout all experiments. After an initial

15-minute washout period, basal bicarbonate secretion (with luminal saline perfusion) was

measured for 30 minutes. Subsequently, linaclotide (10^{-9} M, 10^{-7} M, 10^{-5} M) $\pm$ DRA$_{inh}$-A250

inhibitor (10^{-5} M), CFTR$_{inh}$-172 (2 x 10^{-5} M), or S3226 (10^{-5} M) was perfused intraluminally for 30

minutes. When multiple doses of drugs were studied, they were perfused sequentially in the

same animals. These doses were selected based on previously published data and similarities

with clinical dosing. Forskolin (10^{-4} M) $\pm$ CFTR$_{inh}$-172 (2 x 10^{-5} M) was perfused intraluminally

for 45 minutes. After each experiment, the length of the duodenal test segment was

measured in situ to the nearest 0.5 mm.

Sample volumes were measured by weight to the nearest 0.01 mg. The amount of

bicarbonate in the effluents was quantitated by a validated micro back-titration method.[36]

Briefly, 100 µL of 5×10^{-2} M HCl was added to 2 mL sample with 2 mL of double-distilled water.

Samples were then gassed with N_2, prewashed in $Ba(OH)_2$ to remove all CO_2, and back-titrated

with 2.5×10^{-2} M NaOH to an endpoint of pH 7.0 using an automated pH meter/titration unit (TIM

856 Radiometer, Copenhagen, Denmark). Bicarbonate secretion was determined in 15-minute

periods and expressed as micromoles per centimeter per hour and presented as

bicarbonate output over time or net peak bicarbonate output (peak output minus average basal).

<u>In Vitro Measurement of Ion Transport and Bicarbonate Secretion.</u> For murine measurements,

mice were anesthetized as above, and the proximal 1 cm of duodenum was removed and

placed in ice-cold 3×10^{-1} M mannitol solution with indomethacin (10^{-5} M) to prevent

endogenous prostaglandin formation. The mesentery was dissected off and the duodenum was

opened along the mesenteric edge. The muscular layer was scraped from the mucosa using a

glass slide while being bathed in the indomethacin-containing solution. Stripped duodenal

mucosae were mounted between two 0.1 cm^2 aperture sliders placed in temperature controlled

Ussing chambers maintained at 37°C (P2300, Physiologic Instruments, San Diego, CA). For

human endoscopic biopsy measurements, biopsies were collected in the same iso-osmolar

mannitol solution with indomethacin and then mounted directly between two 0.031 cm^2 aperture

sliders. Tissue was bathed in apical solutions containing (in mM): NaCl 115, KGluconate 5.2,

NaGluconate 25, $MgCl_2$ 1.2, $CaCl_2$ 1.2, Mannitol 10, gassed with 100% O_2 and basolateral

solutions containing (in mM): NaCl 115, K_2HPO_4 2.4, KH_2PO_4 0.4, $NaHCO_3$ 25, $MgCl_2$ 1.2,

$CaCl_2$ 1.2, Glucose 10, gassed with 95% O_2/5% CO_2. Ion transport was measured by short-

circuit current (I_{sc}), where tissues were voltage clamped at 0 mV using a voltage-clamp

apparatus (VCC600, Physiologic Instruments, San Diego, CA). To monitor transepithelial

resistance, the voltage clamp was released, and the open circuit voltage was recorded every 10

minutes. Resistance was calculated using Ohm's Law. To measure bicarbonate secretion, we

utilized the automatic pH-stat method, as previously described.(7, 37) In brief, a pH

microelectrode was placed in the apical bath and set at the pH of the unbuffered apical solution

(typically ~6.8). The titrator (Titrando 902, MetrOhm, Switzerland) was set to titrate 5×10^{-3} M

HCl in 0.2 µL aliquots as the pH of the apical bath rose above the pH set point. Tiamo software

(MetrOhm, Switzerland) was used to control the rate of titration and continuously measure the

amount of HCl titrated into the apical bath and the apical bath pH. The set-up as described

allows for simultaneous measurement of bicarbonate secretion and I_{sc} without electrical

interference. Bicarbonate secretory rates ($\mu mol/cm^2 \bullet h$) were calculated in 5-minute intervals by

noting the amount titrated, the concentration of titrant, and the surface area of the slider

aperture. In a subset of experiments, forskolin (10^{-4} M, serosal) was added at the end of the

experiment to assess tissue viability.

Measurement of Intracellular pH. pH_i levels were measured via epiflorescent microscopy on a

Leica Thunder DMi8 imaging system. Enteroids were grown on collagen-coated 18 mm

coverslips in WENR (undifferentiated) alone or changed to EN for 3 days (differentiated) after

initial seeding in WENR. On day of experiment, enteroids were incubated with the cell-

permeant ratiometric pH indicator SNARF-5F 5-(and-6)-carboxylic acid, acetoxymethyl ester,

acetate (5 µM) (Thermo-Fisher, Waltham, MA) for 30 minutes at room temperature. Following a

15-minute wash period, imaging experiments were performed in physiologic buffer (in mM: NaCl

115, K_2HPO_4 2.4, KH_2PO_4 0.4, $MgCl_2$ 1.2, $CaCl_2$ 1.2, HEPES 10, Glucose 10). Coverslips were

imaged in a heat-controlled (37°C), gassed (95% O_2/5% CO_2) enclosure. Using LasX software,

regions of interest (ROI) were selected by encircling visualized cells for ratiometric

measurements within each ROI. Measurements were taken every 5 seconds. Baseline pH_i was

recorded for 2-3 minutes, followed by application of $CFTR_{inh}$-172 (20×10^{-5} M) or DRA_{inh}-A250

(10^{-5} M, 5 min) then $CFTR_{inh}$-172. Upon full pH_i recovery (typically 4-6 minutes), pH calibration

was performed using serial addition of nigericin + valinomycin in pH 4.5 – 7.5 buffers (Thermo-

Fisher, Waltham, MA) for at least 5 minutes each.

<u>Single cell sequencing analysis.</u> The R toolkit Seurat was employed to conduct an analysis and

quality control of single-cell RNA sequencing (scRNA-seq) data from existing literature.[38] We

clustered the cells according to the Seurat default settings. In brief, we embedded the cells

through a K-nearest neighbor graph, reduced the data with principal component analysis, and

applied the Louvain algorithm for clustering. We visualized these clusters with UMAP (uniform

manifold approximation and projection). To annotate major cell types and compare gene

expression in crypts versus villi, we used the metadata annotations provided by Elmentaite et

al.[21] and Busslinger et al.,[20] respectively. We utilized violin and feature plots to display gene

expression profiles, and FeatureScatter to compare the co-expression of two genes by cell type.

<u>Human duodenal organoids.</u> Duodenal organoids were established and cultured as previously

described.[39] In short, crypt cells were isolated from duodenal endoscopic biopsies using a

series of PBS washes followed by incubation with cold chelation buffer (distilled water with 5.6

mM Na_2HPO_4, 8.0 mM KH_2PO_4, 96.2 mM NaCl, 1.6 mM KCl, 43.4 mM sucrose, 54.9 mM d-

sorbitol, 1 mM dl-dithiothreitol) with 2 μM EDTA for 30 mins on ice. After incubation, EDTA was

removed, and the biopsies were vigorously resuspended in cold chelation buffer (without

EDTA). The resuspension was allowed to sit on ice for 1 minute to allow biopsies to settle by

gravity. The supernatant was collected. This process was repeated 3-5 times until no more

crypts were present in the supernatant. The crypt containing supernatants were pooled and

centrifuged at 600g for 5 minutes, followed by resuspension in 40 µL basement membrane

extract /dome (BME, Culterex Pathclear Type 2 BME) and submerged in WENR complete

media (ADMEM/F12, 1 mM HEPES, 1X Glutamax, 1 mM N-Acetylcysteine, 1X B-27, 0.5 µM

A83-01, 1X Penicillin-Streptomycin-Glutamine, 10 mM Gastrin, 10 µM SB-202190, 50 ng/mL

EGF, 10 mM Nicotinamide, 1X Normocin) with Rho-associated kinase inhibitor (Y-27632 10 µM,

Sigma Aldrich, St. Louis, MO) and CHIR99021 (10µM, Sigma Aldrich, St. Louis, MO) for first two

days, followed by WENR complete media only until ready to passage. Duodenal enteroids were

passaged by disrupting BME with cold (4°C) media, 1X cold (4°C) phosphate-buffer solution

(PBS, without Ca^{2+} and Mg^{2+}), then centrifuged at 600g for 3 minutes at 4°C. Supernatant was

aspirated and single cells were isolated by adding warm (37°C) TrypLE Express (Gibco,

Thermofisher Scientific, Waltham, MA) and incubated for 10 minutes at 37°C. Equal volume of

fetal bovine serum was added to quench trypsin and cells were centrifuged at room temperature

for 5 minutes at 600g. Pellet was resolubilized in BME (40 µL/dome) and plated in a 24-well

tissue culture plate. Generation of apical-out enteroids was performed as previously

described.(22) Briefly, enteroids were dislodged from BME using with 1X cold (4°C) PBS

(without Ca^{2+} and Mg^{2+}) and resuspended in 5 mM EDTA in PBS for 1 hour at 4°C. Enteroids

were then centrifuged at 200g for 3 minutes and resuspended in WENR or EN media (WENR

without Wnt and R-Spondin) and seeded in ultra-low-attachment 24-well plates (Corning

Costar, Thermofisher Scientific, Waltham, MA) for three days. On Day 3, undifferentiated

and/or differentiated enteroids were harvested for experiments.

Quantitative PCR. Quantification of mRNA transcript levels in undifferentiated and differentiated

apical-out enteroids was performed. Briefly, total RNA was extracted from fresh apical-out

enteroids using Zymo Direct-Zol RNA miniprep kit (Zymo Research, Irvine, CA) according to the

manufacturer's instructions. Complementary DNA (cDNA) was synthesized from 1 to 2 µg of

RNA using High-Capacity RNA-to-cDNA™ Kit (Zymo Research, Irvine, CA). Quantitative real-

time polymerase chain reaction (qPCR) was performed on a StepOne Plus real-time PCR

system (Applied Biosystems, Foster City, CA) using PowerUP SYBR Green Master Mix (Applied

Biosystems, Foster City, CA). Each sample was run in triplicate in a 20 µL reaction volume

containing 5 ng cDNA/reaction. PCR was carried out for 40 cycles at 95°C (denature) for 30

seconds, 60°C (anneal/extend) for 30 seconds and amplification data was collected after each

cycle. The sequences of optimized gene-specific primers are listed in Supplemental Table 1.

Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) was used as the reference gene to

obtain delta threshold cycles (ΔCt) values. The relative expression levels of the target genes

were calculated using the $2^{-\Delta\Delta Ct}$ method.

<u>Confocal microscopy.</u>   Treated Day 3 apical-out enteroids were fixed in 2% paraformaldehyde

immediately after treatment and stored at 4°C until processing. All steps were performed in

ultra-low attachment plates. Fixed enteroids were permeabilized for 30 minutes with

permeabilization buffer (3% bovine serum albumin, BSA, ThermoFisher Scientific, Waltham,

MA), 0.5% Triton X-100, and 0.7% Saponin (Sigma-Aldrich, St. Louis, MO) in PBS and blocked

for 60 minutes with gentle shaking in blocking buffer (3% BSA, 0.2% Triton X-100). Samples

were incubated overnight at 4°C with primary antibodies (Supplemental Table 2) diluted in

antibody diluent (3% BSA, 0.2% Triton X-100, and 0.5% Saponin in PBS), washed 4 times, 30

minutes each in antibody diluent buffer, and incubated with secondary antibodies (Table 2) for 1

hour with gentle shaking at room temperature. Samples were washed 4 times for 30 minutes

each in antibody diluent buffer. Samples were then carefully resuspended in SlowFade gold

antifade mounting medium (ThermoFisher Scientific, Waltham, MA), mounted on microscope

slides and sealed with glass coverslips (1.0 mm) (ThermoFisher Scientific, Waltham, MA).

Slides were imaged using Leica Stellaris 8 Confocal microscope. Z-stacks of 0.33 µm

sections/sample were obtained and analyzed using FIJI Software (ImageJ). Following

acquisition, image settings were adjusted to optimize the signal to noise ratio. For quantitative

analysis, the same settings were used across the different conditions compared, and the

protein-of-interest mean fluorescence intensity (MFI) after background subtractions was

analyzed. Villin was used as a marker for the plasma membrane and used to identify the area

for fluorescence intensity measurements. Drug-treated MFI measurements were normalized to

corresponding vehicle control MFI measurements.

Statistics. Results are expressed as means ± standard error of the mean (SEM) for a series of

n experiments. Statistical analysis was performed by using the Student's t-test for unpaired data

or by ANOVA with repeated measures analysis, as appropriate. P values of <0.05 were

considered significant.

Study Approval. All animal protocols were approved by the Institutional Animal Care and Use

Committee at Stanford University (#33183). Children and adults undergoing upper endoscopy

for clinical indications were approached to obtain additional duodenal biopsies for research

under an Institutional Review Board approved protocol (#47536). Subjects without endoscopic

and microscopic evidence of duodenal pathology were included in this study.

AUTHOR CONTRIBUTIONS

JBS: Performed and analyzed experiments, drafted, edited and approved final manuscript.
18
19 AMT: Performed and analyzed experiments, drafted and approved final manuscript.

20 SMA: Performed and analyzed experiments, drafted and approved final manuscript.

21 VVU: Performed and analyzed experiments, drafted and approved final manuscript.
22
23 JEC: Performed and analyzed experiments, approved final manuscript.
24
25 YF: Performed and analyzed experiments, approved final manuscript.
26
27 JG: Performed and analyzed experiments, approved final manuscript.
28
29 MOA: Contributed technical expertise, material support, approved final manuscript.
30
31 OC: Contributed technical expertise, material support, approved final manuscript.
32
33 CJK: Contributed technical expertise, material support, edited and approved final manuscript.

2 ZMS: Study conceptualization, experimental design, funding, drafted, edited and approved final

3 manuscript.

ACKNOWLEDGEMENTS

Drs. Anjaparavanda Naren and Kyu Shik Mun (Cedars Sinai) kindly provided CF enteroids. We

thank Dr. Manuel Amieva (Stanford University) for technical advice on apical-out enteroids.

This research was supported by the National Institute of Diabetes, Digestive, and Kidney

Diseases (K08DK124684 to ZMS, U01DK085527 and R01DK115728 to CJK), R01DK126070

and P30DK072517 to OC, the Cystic Fibrosis Foundation (SELLER19GE0, SELLER20-KB,

SELLER16L0 to ZMS and CIL21G0 to OC), North American Society of Pediatric

Gastroenterology, Hepatology, and Nutrition Foundation (ZMS), Stanford University (ZMS),

Stanford Maternal Child Health Research Institute (ZMS), and Sellers Research and Clinical

Development, LLC (ZMS).

DATA AVAILABILITY

17 The authors declare that the data supporting the findings of this study are available within the

18 paper and its supplementary information files. Any further information on the data in this study

19 are available by reasonable request to the corresponding author.

PREPRINT

This manuscript has been deposited on a public preprint server (doi: https://doi.org/10.1101/

2023.05.05.539132).

FIGURES

Figure 1. Linaclotide stimulates duodenal bicarbonate secretion in mice and humans. A. In vivo measurement
of linaclotide-stimulated (10^{-9} M, 10^{-7} M, 10^{-5} M) duodenal bicarbonate secretion in C57Bl6/J mice (n=9). After 30-
minute baseline measurements, each linaclotide dose was sequentially perfused for 30 minutes. *, P<0.05; **,
P<0.01 vs. baseline by ANOVA. **B.** Net peak (max-baseline) bicarbonate secretion. Linaclotide values were from
1A. For STa (10^{-9} M, 10^{-7} M), separate experiments were performed (n=11). Each point represents a different
mouse. **C.** In vivo effect of linaclotide (10^{-9} M, 10^{-7} M, 10^{-5} M) with NHE3 inhibitor S3226 (10^{-5} M) (n=10). Dotted line
is the mean response to linaclotide (from 1A). *, P<0.05; **, P<0.01 vs. baseline by ANOVA. **D.** Representative trace
for in vitro bicarbonate secretion in mouse duodenum. Y-axis is titrated HCl, which reflects base secreted by the
tissue. During baseline there are few peaks. Upon stimulation, the number of titration events and/or volume of titrant
increases. **E.** In vitro baseline and linaclotide-stimulated (10^{-7}, 10^{-5} M, apical) mouse duodenal mucosal bicarbonate
secretion. Each point (n=8-13) is a separate piece of duodenum from 10 mice. *, P<0.05 vs. baseline by ANOVA. **F.**
Representative short-circuit current (I_{sc}) trace for in vitro mouse experiments. **G.** In vitro baseline and linaclotide-
stimulated (10^{-7}, 10^{-5} M, apical) mouse duodenal I_{sc}. Each point represents a separate piece of duodenum from the
same mice in 1E. ***, P<0.001 vs. baseline by ANOVA. **H.** In vitro transepithelial resistance measurements from 1E
and G. *, P<0.05 vs. baseline by ANOVA. **I-K.** Human duodenal mucosal bicarbonate secretion (I), I_{sc} (J), and
transepithelial resistance (K) from endoscopic biopsies. Experiments were performed and data expressed similar as
murine experiments in 1D-H. Each point (n=9-10) represents a different biopsy. **, P<0.01 vs. baseline by ANOVA.
All data are means ± SEM, unless stated otherwise.

Figure 2. Linaclotide stimulates duodenal bicarbonate secretion independent of CFTR. A. In vivo measurement of duodenal bicarbonate secretion in mice, similar to Figure 1A, with the exception that each perfusate also contained $CFTR_{inh}$-172 (2×10^{-5} M) (n=12). Dotted line is the mean response without $CFTR_{inh}$-172 (from Figure 1A). *, P<0.05 vs. baseline by ANOVA. **B and C.** In vitro duodenal mucosal bicarbonate secretion (B) and I_{sc} (C) in wildtype and *Cftr* KO mice. Data are expressed as fold increase in linaclotide (10^{-7} M) stimulated responses over baseline bicarbonate secretion or I_{sc} in the same mouse. Each point (n=7) is a separate piece of duodenum from 5 mice.   **, P<0.01 by unpaired Student's t-test. **D.** Transepithelial resistance measurements in wildtype (WT) and *Cftr* KO mice. **E-G.** In vitro duodenal mucosal bicarbonate secretion (E), I_{sc} (F), and transepithelial resistance (G) in human endoscopic biopsies with or without $CFTR_{inh}$-172 pre-treatment (2×10^{-5} M, 40-60 minutes). Each point (11-14) represents a different biopsy. All data are means $\pm$ SEM.

Figure 3. Critical role of DRA in CFTR-independent linaclotide-stimulated duodenal bicarbonate secretion. A and B. To determine the source of linaclotide-stimulated bicarbonate transport in the absence of CFTR function, in vivo experiments were repeated, similar to Figure 2A, except in addition to $CFTR_{inh}$-172 (2 x 10^{-5} M, n=12), one of the following was also added to the luminal perfusate: DRA_{inh}-A250 (10^{-5} M, n=10), DIDS (2 x 10^{-4} M, n=7), or S3226 (10^{-5} M, n=9). (A) linaclotide 10^{-7} M, (B) linaclotide 10^{-5} M. Each point represents a different mouse. *, P<0.05 vs. linaclotide + $CFTR_{inh}$-172 (2 x 10^{-5} M) by ANOVA. **C.** Time course with linaclotide dose response (10^{-9} M, 10^{-7} M, 10^{-5} M) in the presence of $CFTR_{inh}$-172 (2 x 10^{-5} M) and DRA_{inh}-A250 (10^{-5} M), as indicated by circles and whiskers (n=12). Dotted line indicates mean response in the presence of $CFTR_{inh}$-172 (2 x 10^{-5} M) only (from Figure 2A). *, P<0.05; **, P<0.01 vs. baseline by ANOVA. **D.** Net peak linaclotide-stimulated (10^{-7} M, apical) mouse duodenal mucosal bicarbonate secretion (D) and I_{sc} (E) from in vitro experiments, with or without DRA inhibition by DRA_{inh}-A270 (10^{-5} M, bilateral, n=5) or DRA_{inh}-4a (10^{-5} M, bilateral, n=14). Each point represents a separate piece of duodenum from 5-10 mice. All data are means ± SEM.

1
Figure 4. Cellular and membrane expression of *SLC26A3* (DRA) in human duodenal enterocytes. A-H.
Cellular mRNA expression of *SLC26A3* (DRA), *SLC26A6* (PAT-1), *CFTR*, and *SLC9A3* (NHE3) based on re-analysis
of Elmentaite et al.(21) (A-D) and Busslinger et al.(20) (E-H). Violin plots represent expression relative to all cells of
that type, with each point representing the expression of individual cells within each type. **I-K.** Representative
confocal immunofluorescence imaging of DRA (I), villin (marker of apical brush border, J), and nucleus (K) in human
duodenum from endoscopic biopsy (n=3). Scale bar = 20 μm.

Figure 5. *SLC26A3* and *CFTR* co-expression in human duodenum by single cell RNA sequencing. A-C. Co-expression of *SLC26A3* (DRA) and *CFTR* mRNA using FeatureScatter based on Elmentaite et al.(21) enterocytes (A) and Busslinger et al.(20) crypt and villi (B and C) datasets that were analyzed in Figure 4. Numbers on top of graphs represent the percentage of *SLC26A3*-expressing cells that express *SLC26A3* only (left) or *SLC26A3* and *CFTR* (right).

Figure 6. Linaclotide increases membrane expression of DRA in apical-out human duodenal enteroids. A. Representative confocal microscopy images of DRA (A), villin (B), and DRA and villin and Hoescht (C) during control conditions (water, 40 minutes, top) or linaclotide (10^{-7} M, 40 minutes, bottom). **D-F.** Quantification of DRA present at the apical brush border using villin to define this cell region. Apical brush border DRA mean fluorescence intensity following linaclotide (10^{-7} M, 40 minutes, n=16) treatment was normalized to vehicle controls for comparison.CFTR$_{inh}$-172 (2×10^{-5} M, 40 minutes) and/or linaclotide (10^{-7} M, 40 minutes) treatment was normalized to vehicle controls for comparison. Vehicle control for linaclotide was water and for CFTR$_{inh}$-172 was DMSO. Columns with whiskers are mean $\pm$ SEM with each dot representing a different enteroid. Enteroids from three different patients were used for each condition. Significance determined by unpaired Student's t-test. For enteroids treated with both CFTR$_{inh}$-172 and linaclotide, enteroids were pretreated with CFTR$_{inh}$-172 for 40 minutes prior to linaclotide treatment. **G and H.** DRA membrane expression in non-CF and CF (F508del homozygous) enteroids during vehicle control conditions (G and H, water) or during linaclotide stimulation (H, 10^{-7} M, 40 minutes). N=8-11 enteroids from 1 patient across 2 different passages. Data were normalized to non-CF enteroids (G) or vehicle control in CF enteroids (H). Significance determined by unpaired Student's t-test. **I.** To compare DRA membrane expression across non-CF and CF enteroids in different situations, absolute DRA membrane mean fluorescence intensity was plotted. Data are from experiments performed in D-H. Significance determined by ANOVA. Arrows provided to assist with identifying regions of interest. All data are means $\pm$ SEM, unless stated otherwise.

Figure 7. Inhibition of CFTR activity induces increases in intracellular pH. A and B. Mean timecourse of CFTR$_{inh}$-172 (2×10^{-5} M) induced changes in pH$_i$ in the absence (A) or presence (B) of DRA$_{inh}$-A250 (10^{-5} M) pre-treatment in undifferentiated human duodenal enteroids. **C.** Quantitative comparison of the peak change (peak – baseline) or recovery (plateau – baseline) in pH$_i$ units for enteroids exposed to CFTR$_{inh}$-172 alone or DRA$_{inh}$-A250 then CFTR$_{inh}$-172. Data are expressed as means ± SEM**. D-F.** Similar experiments and analyses were performed in 3-day differentiated enteroids as A-C. Each circle represents a different cell within an enteroid. Measurements were taken from multiple enteroids across multiple coverslips. ***, $P<0.001$ by paired (within control or DRA$_{inh}$-A250 groups) or unpaired (across control or DRA$_{inh}$-A250 groups) Student's t-test. +++, $P<0.001$ compared to undifferentiated enteroids by unpaired Student's t-test.

REFERENCES

1. Youngberg CA, Berardi RR, Howatt WF, Hyneck ML, Amidon GL, Meyer JH, et al. Comparison of gastrointestinal pH in cystic fibrosis and healthy subjects. *Digestive diseases and sciences.* 1987;32(5):472-80.

2. Gustafsson JK, Ermund A, Ambort D, Johansson ME, Nilsson HE, Thorell K, et al. Bicarbonate and functional CFTR channel are required for proper mucin secretion and link cystic fibrosis with its mucus phenotype. *J Exp Med.* 2012;209(7):1263-72.

3. Seidler UE. Gastrointestinal HCO3- transport and epithelial protection in the gut: new techniques, transport pathways and regulatory pathways. *Curr Opin Pharmacol.* 2013;13(6):900-8.

4. Busby RW, Bryant AP, Bartolini WP, Cordero EA, Hannig G, Kessler MM, et al. Linaclotide, through activation of guanylate cyclase C, acts locally in the gastrointestinal tract to elicit enhanced intestinal secretion and transit. *Eur J Pharmacol.* 2010;649(1-3):328-35.

5. Ahsan MK, Tchernychev B, Kessler MM, Solinga RM, Arthur D, Linde CI, et al. Linaclotide activates guanylate cyclase-C/cGMP/protein kinase-II-dependent trafficking of CFTR in the intestine. *Physiol Rep.* 2017;5(11).

6. McHugh DR, Cotton CU, Moss FJ, Vitko M, Valerio DM, Kelley TJ, et al. Linaclotide improves gastrointestinal transit in cystic fibrosis mice by inhibiting sodium/hydrogen exchanger 3. *Am J Physiol Gastrointest Liver Physiol.* 2018;315(5):G868-G78.

7. Sellers ZM, Childs D, Chow JY, Smith AJ, Hogan DL, Isenberg JI, et al. Heat-stable enterotoxin of Escherichia coli stimulates a non-CFTR-mediated duodenal bicarbonate secretory pathway. *Am J Physiol Gastrointest Liver Physiol.* 2005;288(4):G654-63.

8. Sellers ZM, Mann E, Smith A, Ko KH, Giannella R, Cohen MB, et al. Heat-stable enterotoxin of Escherichia coli (STa) can stimulate duodenal HCO3(-) secretion via a novel GC-C- and CFTR-independent pathway. *FASEB J.* 2008;22(5):1306-16.

9. Tan Q, di Stefano G, Tan X, Renjie X, Romermann D, Talbot SR, et al. Inhibition of Na(+) /H(+) exchanger isoform 3 improves gut fluidity and alkalinity in cystic fibrosis transmembrane conductance regulator-deficient and F508del mutant mice. *Br J Pharmacol.* 2021;178(5):1018-36.

10. Arora K, Huang Y, Mun K, Yarlagadda S, Sundaram N, Kessler MM, et al. Guanylate cyclase 2C agonism corrects CFTR mutants. *JCI Insight.* 2017;2(19).

11. Pratha VS, Hogan DL, Martensson BA, Bernard J, Zhou R, and Isenberg JI. Identification of transport abnormalities in duodenal mucosa and duodenal enterocytes from patients with cystic fibrosis. *Gastroenterology.* 2000;118(6):1051-60.

12. Ma T, Thiagarajah JR, Yang H, Sonawane ND, Folli C, Galietta LJ, et al. Thiazolidinone CFTR inhibitor identified by high-throughput screening blocks cholera toxin-induced intestinal fluid secretion. *J Clin Invest.* 2002;110(11):1651-8.

13. Snouwaert JN, Brigman KK, Latour AM, Malouf NN, Boucher RC, Smithies O, et al. An animal model for cystic fibrosis made by gene targeting. *Science.* 1992;257(5073):1083-8.

14. Tse CM, Yin J, Singh V, Sarker R, Lin R, Verkman AS, et al. cAMP Stimulates SLC26A3 Activity in Human Colon by a CFTR-Dependent Mechanism That Does Not Require CFTR Activity. *Cell Mol Gastroenterol Hepatol.* 2019;7(3):641-53.

15. Lee S, Cil O, Haggie PM, and Verkman AS. 4,8-Dimethylcoumarin Inhibitors of Intestinal Anion Exchanger slc26a3 (Downregulated in Adenoma) for Anti-Absorptive Therapy of Constipation. *J Med Chem.* 2019;62(17):8330-7.

16. Cil O, Anderson MO, de Souza Goncalves L, Tan JA, Haggie PM, and Verkman AS.

Small molecule inhibitors of intestinal epithelial anion exchanger SLC26A3 (DRA) with a

luminal, extracellular site of action. *Eur J Med Chem.* 2023;249:115149.

17. Wang Z, Petrovic S, Mann E, and Soleimani M. Identification of an apical Cl(-)/HCO3(-)

exchanger in the small intestine. *American journal of physiology Gastrointestinal and*

liver physiology. 2002;282(3):G573-9.

18. Foulke-Abel J, In J, Yin J, Zachos NC, Kovbasnjuk O, Estes MK, et al. Human Enteroids

as a Model of Upper Small Intestinal Ion Transport Physiology and Pathophysiology.

Gastroenterology. 2016;150(3):638-49 e8.

19. Yin J, Tse CM, Avula LR, Singh V, Foulke-Abel J, de Jong H, et al. Molecular Basis and

Differentiation-Associated Alterations of Anion Secretion in Human Duodenal Enteroid

Monolayers. *Cellular and Molecular Gastroenterology and Hepatology.* 2018;5(4):591-

609.

20. Busslinger GA, Weusten BLA, Bogte A, Begthel H, Brosens LAA, and Clevers H. Human

gastrointestinal epithelia of the esophagus, stomach, and duodenum resolved at single-

cell resolution. *Cell Rep.* 2021;34(10):108819.

21. Elmentaite R, Kumasaka N, Roberts K, Fleming A, Dann E, King HW, et al. Cells of the

human intestinal tract mapped across space and time. *Nature.* 2021;597(7875):250-5.

22. Co JY, Margalef-Catala M, Li X, Mah AT, Kuo CJ, Monack DM, et al. Controlling

Epithelial Polarity: A Human Enteroid Model for Host-Pathogen Interactions. *Cell Rep.*

2019;26(9):2509-20 e4.

23. Chen T, Hubbard A, Murtazina R, Price J, Yang J, Cha B, et al. Myosin VI mediates the

movement of NHE3 down the microvillus in intestinal epithelial cells. *J Cell Sci.*

2014;127(Pt 16):3535-45.

24. Wen G, Deng S, Song W, Jin H, Xu J, Liu X, et al. Helicobacter pylori infection

downregulates duodenal CFTR and SLC26A6 expressions through TGFbeta signaling

pathway. *BMC Microbiol.* 2018;18(1):87.

25. Gelfond D, Heltshe S, Ma C, Rowe SM, Frederick C, Uluer A, et al. Impact of CFTR

Modulation on Intestinal pH, Motility, and Clinical Outcomes in Patients With Cystic

Fibrosis and the G551D Mutation. *Clin Transl Gastroenterol.* 2017;8(3):e81.

26. Braga Emidio N, Tran HNT, Andersson A, Dawson PE, Albericio F, Vetter I, et al.

Improving the Gastrointestinal Stability of Linaclotide. *J Med Chem.* 2021;64(12):8384-

90.

27. Rao SP, Sellers Z, Crombie DL, Hogan DL, Mann EA, Childs D, et al. A role for

guanylate cyclase C in acid-stimulated duodenal mucosal bicarbonate secretion.

American journal of physiology Gastrointestinal and liver physiology. 2004;286(1):G95-

G101.

28. Hogan DL, Crombie DL, Isenberg JI, Svendsen P, Schaffalitzky de Muckadell OB, and

Ainsworth MA. Acid-stimulated duodenal bicarbonate secretion involves a CFTR-

mediated transport pathway in mice. *Gastroenterology.* 1997;113(2):533-41.

29. Strubberg AM, Liu J, Walker NM, Stefanski CD, MacLeod RJ, Magness ST, et al. Cftr

Modulates Wnt/beta-Catenin Signaling and Stem Cell Proliferation in Murine Intestine.

Cell Mol Gastroenterol Hepatol. 2018;5(3):253-71.

30. Kaunitz JD, and Akiba Y. Duodenal intracellular bicarbonate and the 'CF paradox'. *JOP.*

2001;2(4 Suppl):268-73.

31. Bijvelds MJC, Roos FJM, Meijsen KF, Roest HP, Verstegen MMA, Janssens HM, et al.

Rescue of chloride and bicarbonate transport by elexacaftor-ivacaftor-tezacaftor in

organoid-derived CF intestinal and cholangiocyte monolayers. *Journal of cystic fibrosis :*

official journal of the European Cystic Fibrosis Society. 2021.

32. Walker NM, Simpson JE, Brazill JM, Gill RK, Dudeja PK, Schweinfest CW, et al. Role of down-regulated in adenoma anion exchanger in HCO3- secretion across murine duodenum. *Gastroenterology.* 2009;136(3):893-901.

33. Singh AK, Liu Y, Riederer B, Engelhardt R, Thakur BK, Soleimani M, et al. Molecular transport machinery involved in orchestrating luminal acid-induced duodenal bicarbonate secretion in vivo. *The Journal of physiology.* 2013;591(21):5377-91.

34. Stahl M, Stahl K, Brubacher MB, and Forrest JN, Jr. Divergent CFTR orthologs respond differently to the channel inhibitors CFTRinh-172, glibenclamide, and GlyH-101. *Am J Physiol Cell Physiol.* 2012;302(1):C67-76.

35. Haggie PM, Cil O, Lee S, Tan JA, Rivera AA, Phuan PW, et al. SLC26A3 inhibitor identified in small molecule screen blocks colonic fluid absorption and reduces constipation. *JCI Insight.* 2018;3(14).

36. Hogan DL, Crombie DL, Isenberg JI, Svendsen P, Schaffalitzky de Muckadell OB, and Ainsworth MA. CFTR mediates cAMP- and Ca2+-activated duodenal epithelial HCO3- secretion. *Am J Physiol.* 1997;272(4 Pt 1):G872-8.

37. Pratha VS, Thompson SM, Hogan DL, Paulus P, Dreilinger AD, Barrett KE, et al. Utility of endoscopic biopsy samples to quantitate human duodenal ion transport. *J Lab Clin Med.* 1998;132(6):512-8.

38. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al. Integrated analysis of multimodal single-cell data. *Cell.* 2021;184(13):3573-87 e29.

39. Sato T, Stange DE, Ferrante M, Vries RG, Van Es JH, Van den Brink S, et al. Long-term expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and Barrett's epithelium. *Gastroenterology.* 2011;141(5):1762-72.