Risk and loss aversion and attitude to COVID and vaccines in anxious individuals

Filippo Ferrari, Jesse Alexander, Peggy Seriès

Institute for Adaptive and Neural Computation

School of Informatics, University of Edinburgh

February 2023

Abstract

Anxious individuals are known to show impaired decision-making in economic gambling task and in everyday life decisions. This impairment can be due to aversion to uncertainty about outcomes (risk aversion) and/or aversion to negative outcomes (loss aversion). We investigate how non-clinical individuals with high levels of Generalised Anxiety Disorder (GAD) (N = 54) behave compared to less anxious subjects (N = 61) in a gambling decision-making task delivered online and designed to separate the distinct influences of risk and loss aversion on decision-making. By modelling subjects' choices using computational models derived from Prospect Theory and fitted using Hierarchical Bayesian methods, we estimate individual levels of risk and loss aversion. We also link estimates of these parameters to individual propensity to risk averse behaviours during the COVID pandemic, like wearing safer types of face masks, or completing a COVID vaccination course. We report increased loss aversion in individuals with increased level of GAD compared to less anxious individuals and no differences in risk aversion. We also report no links between risk and loss aversion and attitudes towards COVID and vaccines. These results shed new light on the interplay of anxiety and risk and loss aversion and they can provide useful directions for clinical intervention.

1 Author Summary

In everyday life, we are constantly faced with decisions in which we need to balance between possible losses and risks. In general, humans are believed to weight losses more than gains, i.e. they are loss averse, and they prefer certain

to uncertain outcomes, i.e. they are risk averse. The COVID pandemic made this balance between risks, losses and everyday life even more explicit: should I get a COVID vaccination? What type of mask should I wear? Here, we investigate these constructs using a gambling task delivered online in which participants have to accept or refuse bets involving monetary gains, losses or sure wins. We also capture individual attitudes to COVID and vaccines with a new self-report questionnaire. We find that individuals with high levels of Generalised Anxiety Disorder present higher levels of loss aversion compared to less anxious individuals and no differences in risk aversion. We also report no links between computational measures of risk and loss aversion and attitude to COVID and vaccines, possibly due to limits in using simple tasks to describe behaviours through complex situations like the COVID pandemic.

2 Introduction

Anxiety disorders have long been associated with behavioural differences in decision-making (Bishop and Gagne, 2018). Here we examine economic decisions in which two main factors are at play: risk and loss aversion. A famous behavioural economics theory called Prospect Theory (Kahneman and Tversky, 1979) describes how humans prefer sure outcomes with lower magnitude (low risk) compared to less sure outcomes with higher magnitude (high risk), which results in risk averse behaviours. Prospect Theory also describes how individuals are generally considered loss averse, i.e. humans prefer to avoid losses to acquiring gains of equivalent value, usually with a 2:1 ration between losses and gains.

In this study we use computational models derived from Prospect Theory to quantify the influence of risk and loss aversion on human decision-making. We adapt a gambling task previously used by Charpentier et al. (2017) to be administered online on a non-clinical sample of adults. This task is particularly interesting as it has been one the first of its kind which allows to separate the influence of risk aversion (i.e., aversion to uncertainty about outcomes) and loss aversion (i.e., aversion to negative outcomes) on economic decision-making.

The impact of anxiety on risk aversion has been studied in several studies generally linking increased level of anxiety to increased risk aversion biases in decision-making tasks (Maner et al., 2007; Mueller et al., 2010; Giorgetta et al., 2012; Charpentier et al., 2017; see Hartley and Phelps, 2012; Phelps et al., 2014; Bishop and Gagne, 2018; Wake et al., 2020, for reviews). Despite this, results from studies employing computational models are mixed. As discussed above, Charpentier et al. (2017) reports increased risk aversion in anxious individuals, whereas Xu et al. (2020) reports no difference in risk aversion for low and high

anxiety groups. Two recent studies (Wise et al., 2022; Pike et al., 2023) also failed to observe a relationship between anxiety and differences in risk-taking, despite using different decision-making tasks. A recent meta-analysis (Wake et al., 2020) on the effects of fear (and anxiety) on risk-taking reports a small to moderate effect size linking fear to increased risk aversion, but with high heterogeneity in effect sizes.

Loss aversion, on the other hand, has only been investigated in anxiety in a handful of studies. Charpentier et al. (2017) found that individuals with high anxiety did not show differences in loss aversion compared to the low anxiety individuals. Sokol-Hessner and Rutledge (2019) propose that the effects of affective experiences, like anxiety, on loss aversion are due to very precise temporal structures between decisions and affective responses.

Both risk and loss aversion are important factors in anxiety with major effects, not only on obvious economic decisions, but also on individuals' decisions in everyday life. This is particularly true in crises situation such as observed during the COVID pandemic. When the health of large populations is at stake, it is important to understand what motivates specific attitudes and behaviours toward vaccinations, preemptive measures like wearing face masks and adherence to government guidelines. For example, intention to be vaccinated was associated with more positive beliefs around COVID vaccines and reduced beliefs that these would lead to negative side effects or be unsafe (Sherman et al., 2021). A recent longitudinal study (Shou et al., 2022) showed that risk averse individuals in Australia were more likely to act in a way that would reduce their risk of contracting COVID-19. At the same time, anxious individuals showed increased avoidance in implementing changes in their actions as the pandemic developed, for example continuing to cancel family events or trips even as the pandemic eased off. The first result is also consistent with Wise et al. (2022), where it is reported that higher risk aversion is linked to higher perceived threat from the pandemic. In our study, we are interested in determining whether behaviours like receiving COVID vaccinations, following government's guidelines, wearing more protective types of face masks are modulated by anxiety and individual propensity to taking risks and/or accepting losses. We hypothesise that individual who adopted safer attitudes and behaviours around COVID would show increased risk aversion, and that this difference is modulated by anxiety.

In summary, this work has therefore several goals. First, we replicate and extend, using an online version of the same gambling task, previous work (Charpentier et al., 2017) linking anxiety to increased risk aversion. Second, we investigate how individual levels of anxiety as well as risk and loss aversion relate to decision-

104 making in the context of the COVID pandemic.

3   Results

Reaction Times

Mean reaction times did not differ across low and high anxiety subjects (low
anxiety: 1.254 (0.470) seconds; high anxiety: 1.213 (0.438) s; $t_{113} = -0.791$, $p =$
0.430) or vaccinated and unvaccinated (vaccinated: 1.208 (0.439) s; unvaccinated:
1.264 (0.472) s; $t_{113} = -1.105$, $p = 0.271$). We observe a statistically significant
difference in reaction times between low/high anxiety subjects in mixed gamble
trials in which participants decided not to gamble: the high anxiety group
chooses not to gamble faster compared to the low anxiety group (low anxiety:
1.384 (0.391) s; high anxiety 1.240 (0.245) s; $t_{113} = -2.326$, $p = 0.021$, not
significant under a Bonferroni correction with $\alpha = 0.0027$).

Choice Data Analysis

We report the percentage of gambles across the participants, the low/high anxiety
groups and the vaccinated/unvaccinated groups in table 1. There were no group
differences across percentages of gamble across either the low/high anxiety
groups or the vaccinated/unvaccinated groups (all $p > 0.144$). Percentage of
gambles across mixed gamble and gain-only trials are not significantly correlated
$(r(115) = 0.148$, $p = 0.116$, $\text{BF}_{10} = 0.396)$ indicating that subjects must use
different strategies in these different types of trials.

	p(gamble)	p(gamble-mixed)	p(gamble-gain)
All participants	0.587	0.576	0.609
Low Anxiety	0.597	0.603	0.586
High Anxiety	0.577	0.546	0.636
Vaccinated	0.578	0.553	0.628
Unvaccinated	0.597	0.601	0.589

Table 1: Probability of gamble during trials. Here p(gamble-mixed) and p(gamble-gain) refers to the probability of gambling in, respectively, mixed gamble and gain only trials.

Participants were payed a bonus up to £1, proportional to their final amount
of money collected at the end of the gambling trials. The final bonus won by
participants was on average £0.37 (£0.08) with no significant difference between
vaccination status (vaccinated: £0.369 (£0.082); unvaccinated £0.371 (£0.077);

$t_{113} = 0.131$, $p = 0.896$) or low/high anxiety (low anxiety: £0.318 (£0.087); high
anxiety £0.359 (£0.070); $t_{113} = 1.496$, $p = 0.137$).

Risk and Loss Aversion

The Hierarchical Bayesian (HB) fitting method produced more stable results
compared to Maximum Likelihood Estimation (MLE) during both parameter
recovery and when fitting the participants' data. The 3 parameters model (Model
1 - allP) fitted using HB was thus picked to conduct further analysis.

First, we fit the model with six different hierarchical priors specifications on the
parameters (similarly to Ahn et al., 2014; Aylward et al., 2019): a single prior
for each parameter encompassing all participants, two separate priors for each
parameter for low and high anxiety subjects according to the GAD-7 median
split, two separate priors for each parameter for low and high anxiety subjects
according to the STAI Trait median split, two separate priors for each parameter
for vaccinated and unvaccinated subjects, four different priors for each parameter
for the combinations of GAD-7 low/high anxiety and vaccinated/unvaccinated
subjects and four different priors for each parameter for the combinations of STAI
Trait low/high anxiety and vaccinated/unvaccinated subjects. Throughout the
rest of this work we refer to these models as single prior, GAD-7/Trait anxiety
priors, vaccination priors and GAD-7/Trait anxiety and vaccination priors for
convenience. Our data is best fitted using a single prior for all participants
(table 2), with the model fitted using the GAD-7 median split priors being a
close second.

	LOOIC
Single prior	**13065.12**
GAD-7 Anxiety priors	13068.45
STAI Trait Anxiety priors	13086.08
Vaccination priors	13083.36
GAD-7 Anxiety and vaccination priors	13101.54
STAI Trait Anxiety and vaccination priors	13086.81

Table 2: Model 1 LOOIC scores for different priors specifications.

Due to the negligible difference in LOOIC scores between the single prior model
and the GAD-7 anxiety prior model the rest of the analysis will report results
from the anxiety prior model. This decision is also backed up by previous
research (Valton et al., 2020) which recommends to account for hypothesised
group differences in computational modelling. For completeness, we report
results for the single prior model in appendix D.

156 We report the fitted parameters for the GAD-7 anxiety priors model in table 3
157 and in fig. 1. Consistently with the prospect theory literature (Kahneman and
158 Tversky, 1979), individuals are on average risk averse ($\rho < 1$) and they value
159 losses twice as much as wins ($\lambda \approx 2$). We report differences in loss aversion and in
160 inverse temperature between the low and high anxiety groups using the difference
161 between the 95% Highest Density Interval (HDI) of the two groups. The high
162 anxiety group shows increased loss aversion ($\text{BF}_{10} = 2.156$) and reduced inverse
163 temperature ($\text{BF}_{10} = 2.407$) compared to the low anxiety group. No difference
164 in risk aversion ($\text{BF}_{10} = 0.239$) was found between the low and high anxiety
165 groups. The individual parameters are shown in fig. 2.

	Low Anxiety	High Anxiety	95% Between-group HDI	
Loss Aversion λ	2.126 (0.935)	2.537 (0.966)	**0.0544**	**0.903**
Risk Aversion ρ	0.916 (0.296)	0.949 (0.264)	-0.089	0.155
Inverse Temperature μ	1.889 (1.797)	1.217 (1.117)	**-0.937**	**-0.076**

Table 3: Mean (standard deviation) of the fitted parameters of Model 1 using
the anxiety prior based on the GAD-7 scores median split. Parameters for which
the between-group 95% HDI does not encompass zero (in bold) are considered
different across groups.

Figure 1: Parameters distributions for the anxiety prior model.

166 GAD-7 scores are significantly correlated with loss aversion ($r_{115} = 0.193$,
167 $p = 0.039$, $\text{BF}_{10} = 0.953$) and inverse temperature ($r_{115} = -0.196$, $p = 0.035$,
168 $\text{BF}_{10} = 1.033$), but not risk aversion ($r_{115} = 0.075$, $p = 0.429$, $\text{BF}_{10} = 0.159$).

Figure 2: 95% HDI of the group mean differences for the loss aversion, risk aversion and inverse temperature parameters.

We fitted a logistic regression model with low/high anxiety (based on GAD-7 median split as above) as the dependent variable and loss λ and risk ρ aversion parameters as independent variables. We find that loss aversion was a statistically significant predictor of anxiety group ($z = 2.242$, $p = 0.025$) while risk aversion was not ($z = 0.641$, $p = 0.522$).

Despite previous research pointing toward differences between low/high STAI Trait anxiety scores, our model fails to capture any significant group difference in its parameters. This is surprising given a strong correlation between the STAI Trait Anxiety scores and the GAD-7 scores ($r_{115} = 0.830$, $p < 0.001$, $\mathrm{BF}_{10} > 10^{27}$).

COVID Score analysis

Our data do not support any correlation between fitted parameters and COVID score (or the COVID-Past/Future/Generic scores described in appendix A). We report a positive correlation between State Anxiety and COVID score ($r(115) = 0.194$, $p = 0.037$, $\mathrm{BF}_{10} = 0.993$), State Anxiety and COVID-Future score ($r(115) = 0.193$, $p = 0.038$, $\mathrm{BF}_{10} = 0.972$) and State Anxiety and COVID-Generic score ($r(115) = 0.199$, $p = 0.032$, $\mathrm{BF}_{10} = 1.113$). We also find a near significant correlation between State Anxiety and COVID-Past score ($r(115) = 0.159$, $p = 0.089$, $\mathrm{BF}_{10} = 0.484$)

4 Discussion

In this study we found that individuals with high levels of Generalised Anxiety Disorder show increased loss aversion compared to individuals with lower anxiety levels. No differences in risk aversion were found between the low and high anxiety groups. Our results are consistent with a recent study by Xu et al.

(2020), which found increased levels of loss aversion and no differences in risk aversion in individuals with high trait anxiety. Xu et al. used a prospect theory model similar to the one used in our study but their task design differs in that they only employed mixed gamble trials in which expected value and variance of the gain/loss amounts were manipulated. Our task, on the other hand, like the one used by Charpentier et al. (Charpentier et al., 2017), present different types of trials (mixed gamble and gain-only). Notably, our results do not replicate Charpentier et al.'s differences in risk aversion and instead show differences in loss aversion, where they did not find any. This suggests that more work is needed in this field, both to test the robustness of these results and understand the differences observed across different studies. Several competing factors can explain these mixed results. First, our study and Xu et al.'s study were conducted on subclinical populations, whereas Charpentier et al.'s study was conducted on a clinical population, which may indicate possible behavioural differences between subclinical and clinical populations of anxiety. Charpentier et al.'s study also included an emotional manipulation task which could have had an impact on decision-making that may not be picked up by the models used. Second, Xu et al.'s and Charpentier et al.'s studies employed STAI Trait anxiety scores as a measure of anxiety, whereas in our study results only hold for GAD-7 anxiety scores and not for STAI Trait anxiety scores. This may highlight the use of largely overlapping but different cognitive mechanisms captured by STAI Trait anxiety scores and GAD-7 scores, despite these being highly correlated both in our study ($r_{115} = 0.83$) and more generally in the literature (Doi et al., 2018). Third, our experiment explicitly instructs participants to maximise their final rewards in order to win a larger bonus payment while participants are able to see the result of each gamble (and therefore the amount of money won or lost) at each trial. In Xu et al.'s experiment, participants were only shown the result of a trial selected at random, with the won/lost amount being added/removed from a fixed amount of money received by the participants before the experiment; in Charpentier et al.'s experiment, participants were shown the results of 10 trials selected at random which were then averaged and either added or removed from a fixed amount of money to reward the participant. This difference in experimental design may explain the overall high values (i.e., reduced risk aversion) that our model fitted for the risk aversion ρ parameter (overall: 0.931 (0.281), low anxiety: 0.916 (0.296), high anxiety: 0.949 (0.264)), compared, for example to a lower mean value of ρ reported by Charpentier et al. (overall: 0.713 (0.458), low anxiety: 0.875 (0.537), high anxiety: 0.564 (0.313)). Our study's explicit instructions to maximise rewards could therefore explain our values of ρ close to 1, participants may be motivated to accept more gambles than they normally would in order to increase their in-game scores and their proportional bonus

payments. During the experiment's practice trials, we adjust the trials expected values to each participant's sensibility, and we remove participants for which this procedure was not successful. As a result of this, the average expected value of all trials was positive for each participants. Therefore, increased gambling would lead to net positive wins.

We did not find statistically significant differences in reaction times across anxiety levels, apart from the high anxiety group being faster at choosing not to gamble (i.e., avoiding the chance of losing money) in mixed gamble trials. It is generally believed that high levels of anxiety result in faster evidence accumulation for "threatening" choices in studies employing Drift Diffusion Models (DDMs) (see Roberts and Hutcherson, 2019, for a review). We hypothesise that in our study, given the explicit instruction to maximise rewards, the high anxiety group may give a negative valence to mixed gamble trials – as these may lead to losses – which would then result in faster reaction times, possibly due to higher drift rates. Further work employing DDMs could be used to understand the nature of this difference in reaction times between anxiety groups.

We also highlight how hierarchical Bayesian methods provide better and more reliable fitted parameters compared to Maximum Likelihood estimation, as measured in terms of better parameter recovery.

Previous questionnaires-based research (Millroth and Frey, 2021) has highlighted how anxiety and negative individual disposition towards risk and uncertainty have an important role in predicting strong fear responses related to the COVID-19 pandemic. However, in our study, the hypothesis that individual levels of risk and loss aversion could be linked to decision-making in the context of the COVID pandemic was rejected. Simple economic decision-making tasks like the one we employed might not be able to capture the intrinsic complexity of decision-making in a multifaceted situation like the COVID pandemic in which economic, health and social factors are all at interplay and more complex naturalistic tasks (Schonberg et al., 2011) may be required to better study these relationships. A recent review about predictors of vaccine hesitancy (Hudson and Montelpare, 2021) includes risk aversion as one of the possible individual difference factors which influence vaccine uptake. Risk aversion could impact vaccine hesitancy in two different and opposite ways. A risk averse individual might value the unknown side effects of a vaccine as riskier than the effects of a known disease and would only get vaccinated due to work or travel requirements. At the same time, an individual might find the dangers posed by a vaccine-preventable disease as riskier than the possible side effects of a vaccine. Loss aversion may also contribute to decisions taken during the pandemic in different

ways. For example, workers who would received little protection in case of loss of income due to self isolation following a COVID infection may value the loss of their working hours as more important than the (individual and public) health gains given by self isolating in case of mild COVID symptoms. Therefore, assuming vaccinations and stronger adherence to healthcare guidelines as the risk averse behaviours is a limitation in our study, as some individuals may consider these as more risk- and/or loss-prone actions. Hence, our self-report COVID questionnaire should be expanded to include questions designed to account between these different views.

In conclusion, our results indicate that subjects with high levels of Generalised Anxiety Disorder show increased loss aversion, and no differences in risk aversion, as measured by computational parameters fitted on data from a economic decision-making task. Future computational studies should focus on disentangling the effects of motivation and reward on risky decision-making and to better study the temporal patterns that might affect decision-making. These findings may help clinicians in tailoring behavioural interventions in order to overcome decision biases. A previous study (Lorian et al., 2012) provides evidence for a reduction of risk-averse behaviours following a cognitive behavioural therapy treatment. Therefore, future work could explore the benefits of including loss aversion as a treatment outcome in similar intervention.

5 Methods

5.1 Ethics

The study was approved by the Informatics Research Ethics Process (application number 2019/84385) of the School of Informatics, University of Edinburgh.

5.2 Participants

We recruited 163 participants based in the United Kingdom (80 with no COVID vaccinations, 83 with 2 or more doses of an approve COVID vaccine) in the months of September and October 2022 through Prolific. Participants were selected from the Prolific users residing in the United Kingdom and who received either 0 or 1+ dose of an approved COVID vaccine. Participants were paid £2.67 (the task took on average 20 minutes to complete, corresponding to a pay of £8/hr) with the possibility to win up to £1 proportional to their performances during the task. We performed a power analysis based on the results reported by Charpentier et al. (2017): a sample of 104 participant is required for risk aversion being higher in anxious individuals relative to control subjects in order

³⁰⁶ to achieve an effect size of Cohen's d = 0.72 at a significance level of 0.05 and
³⁰⁷ with power of 0.95. We employed exclusion criteria based on attention checks
³⁰⁸ and on the participants behaviour, these are described in detail in appendix B.

³⁰⁹ After applying these exclusion criteria, we conducted the experiments on 115
³¹⁰ subjects, for whom we report demographic information in table 4, together with
³¹¹ the questionnaire results described in subsection 5.4.

	Vaccinated	Unvaccinated	t_{113}	p-value
Women:Men (Sum)	37:23 (60)	22:33 (55)	-	-
Age in years	40.88 (13.51)	38.38 (11.90)	1.049	0.296
STAI State Anxiety	43.01 (13.83)	38.54 (12.38)	1.819	0.071
STAI Trait Anxiety	47.86 (14.04)	48.52 (13.79)	-0.254	0.799
GAD-7	8.83 (5.71)	7.67 (5.74)	1.085	0.279
COVID Score	50.88 (8.62)	30.47 (9.94)	11.787	**< 0.001**
COVID Score (No #vax)	47.06 (8.42)	29.47 (9.94)	10.267	**< 0.001**

Table 4: Statistics about the sample and questionnaires results. Values are mean (SD). The last row refers to COVID Score group differences between the vaccinated and unvaccinated groups when the number of vaccines is not included in the score.

³¹² Throughout this work we refer as the low/high anxiety group as the lower/higher
³¹³ median split of the data using GAD-7 scores (median GAD-7: 7) and as the
³¹⁴ vaccinated group as subjects who received 1 or more doses of a COVID-19
³¹⁵ approved vaccine (see table 5).

	High Anxiety	Low Anxiety	Total
Vaccinated	30	30	60
Unvaccinated	24	31	55
Total	54	61	115

Table 5: Number of participants for each group. Low/high anxiety groups are split by the median GAD-7 score of 7. Vaccinated participants received 1+ doses of a COVID-19 approved vaccine.

5.3 Task

³¹⁷ The task was adapted from Charpentier et al. (2017) and implemented using
³¹⁸ jsPsych (de Leeuw, 2015). It could only be completed using a computer. During
³¹⁹ the gambling task participants are competing in a game show where, at each
³²⁰ trial, they are asked to make a choice between a gambling wheel and a sure
³²¹ option. Each option on the gambling wheel has a 50% probability to occur.
³²² Example trials are shown in fig. 3. There are two types of trials that are shown

to the participants, *mixed gamble* trials and *gain-only* trials. In mixed gamble trials one option leads to a sure option of £0, whereas the other option consists of a wheel with a 50% chance to win a monetary reward, and a 50% chance to lose some of the money accumulated so far. In gain-only trials the sure option consists of a fixed, small amount of money, whereas the other option consist of a wheel with a 50% chance to win a larger monetary reward, and a 50% chance to win nothing. During gain-only trials, only risk aversion contributes to choosing from the safe option compared to the gambling option, while both risk and loss aversion contribute to choosing from the safe option in mixed gamble trials. At the beginning of the experiment, participants are instructed to win as much money as possible, and they are told that they will receive a bonus payment through Prolific based on their performance during the experiment. A practice section consisting of 40 trials is used to introduce the participants to the task and to adapt, through a double staircase procedure, the amounts shown at each trial to a range of expected values centered on the subject's indifference point, i.e. the expected value corresponding to an accept rate of 50% between the gamble and the sure option. The main section of the experiment consists of 148 trials (98 mixed gamble trials and 50 gain-only trials in random order, the same proportion of mixed gamble/gain-only used in Charpentier et al.), in which participants have to select an option within 4s, followed by a 2s animation showing the sure/win/loss amount. Participants were instructed to use keys [A] and [L] of their keyboards to choose between options.

5.4   Questionnaires

At the end of the gambling task, participants were asked to complete several questionnaires. First, we administered the State Trait Anxiety Inventory (STAI) (Spielberger, 1977) to measure state and trait anxiety scores, two measures of anxiety that have widely been used in computational studies (Raymond et al., 2017), and the GAD-7 (Spitzer et al., 2006) questionnaire to assess Generalised Anxiety Disorder. Following these, participants were asked 13 questions about the COVID pandemic. Four questions asked about the subject vaccination status, the type of mask used throughout the pandemic, how anxious they were feeling at the beginning of the pandemic and how closely they followed government guidelines during the pandemic. We matched these questions with four questions asking about the subject's future attitude towards vaccine boosters, masks, anxiety about attitude towards rules and guidelines in the case of a new surge of COVID cases. The last 5 questions asked about the subject's general attitude towards COVID and vaccines. All these questions are rated on a Likert-scale of [1-5] and, based on the answers, we computed a COVID score

Figure 3: Task structure. (A) The frame shows a mixed gamble trial. The wheels are shown for 4s. If a participant does not make a choice within this time frame, the trial results in a time out. Once a wheel is chosen a 1s animation is shown (B), followed by the amount won/lost shown for 1s (C). The last frame (D) shows a gain-only trial, which has the same trial structure.

where higher values corresponds to more cautious behaviour during the COVID pandemic (for example, completing the vaccination cycle, following government guidelines and wearing N95 masks). The COVID questionnaire and rating scale is reported in full in appendix A.

It can be seen from table 4 that no statistically significant difference were observed between our samples of vaccinated and unvaccinated individuals with respect to state and trait anxiety and GAD-7 scores. However, a statistically significant difference across COVID scores is present - even when the number of vaccination is not included in the score - showing that our questionnaire is able to pick up the different attitude towards COVID between unvaccinated and vaccinated individuals.

5.5 Models

We fitted three different models derived from Prospect Theory to the subjects' data. The models are the same as those used in Charpentier et al. (2017): the first model measures both risk and loss aversion, while the other two are nested models which only measure loss and risk respectively. The models have been fitted using both maximum likelihood estimation (MLE) in MATLAB and using hierarchical Bayesian methods implemented in the hBayesDM R library (Ahn et al., 2017).

The first model (allP) captures both risk and loss aversion and is described by the following equations

$$u(gamble) = 0.5 \times (gain)^\rho + 0.5 \times \lambda \times (-loss)^\rho \tag{1}$$

$$u(sure) = (sure)^\rho \tag{2}$$

$$p(gamble) = \frac{1}{1 + e^{-\mu[u(gamble)-u(sure)]}} \tag{3}$$

where u is the subjective utility of gambling or choosing the sure option, $gain$ and $loss$ are the numeric values shown on the gambling wheel and $sure$ is the value of the sure option. The difference between the subjective utilities is then passed through a softmax function parameterised by inverse temperature μ (eq. (3)); this remains the same across all models. The model has three parameters:

- λ represents loss aversion. When $\lambda > 1$, the subject weights losses more than gains and it is therefore loss aversed; $\lambda < 1$ represents the inverse.

- ρ represents risk aversion. When $\rho < 1$, the subjects is risk aversed, whereas with $\rho > 1$ the subject is risk seeking.

- μ is the inverse temperature of the action selection function where higher values of μ represent less stochastic decisions.

The second model (noRA) has two parameters, one for loss aversion λ and one for the inverse temperature μ:

$$u(gamble) = 0.5 \times (gain) + 0.5 \times \lambda \times (-loss) \tag{4}$$

$$u(sure) = (sure) \tag{5}$$

and the third model (noLA) similarly only includes a parameter for risk aversion

396 ρ and the inverse temperature μ:

$$u(gamble) = 0.5 \times (gain)^\rho + 0.5 \times (-loss)^\rho \tag{6}$$

$$u(sure) = (sure)^\rho \tag{7}$$

5.6 Model fitting procedures

Maximum Likelihood Maximum Likelihood models were fitted using MATLAB R2021b. We run constrained optimisation (`fmincon` with default settings) for $\lambda, \rho, \mu \in [0, \infty]$. For each subject we run optimisation 20 times and we select the best set of parameters according to their negative log likelihood.

Hierarchical Bayes Hierarchical Bayesian models were fitted using R version 3.6.3 and hBayesDM (Ahn et al., 2017) version 1.2.1 which, in turn, uses Stan (Carpenter et al., 2017) version 2.21.7. We fitted 4 chains for each model, with 2000 burn-in samples and 6000 samples (8000 samples in total). We visually inspected traces for each parameter for good mixing and checked that the Gelman–Rubin statistic $\hat{R}$ (Gelman and Rubin, 1992) was close to 1 for all fitted parameters to test for convergence.

5.7 Parameter Recovery

We performed extensive simulation and parameter recovery for the first model using both maximum likelihood estimation (MLE) and Hierarchical Bayesian methods (HB). We tested experimental designs comprising 148, 222 and 296 trials, with different combinations of Indifference Points and low/high anxiety parameters. For each combinations of parameters and Indifference Points(IP) we simulated 100 participants. Simulation parameters for low and high anxiety agents were taken from the existing literature (Charpentier et al., 2017). We report correlations using the more robust Spearman corerlation coefficient, due to outliers resulting from the MLE fitting procedure. Across all combinations of parameters, fitting procedure and number of trials, recovery is successfull, with at least a Spearman $\rho > 0.64$. Parameter recovery results for 148 trials are shown in table 6 and full details of the parameter recovery set up and results for 222 and 296 trials are reported in appendix C. We set the length of the experiment to 148 trials due to the similar results of the parameter recovery across different number of trials.

	MLE			HB		
	λ	ρ	μ	λ	ρ	μ
High Anxiety / Low IP	0.82	0.88	0.88	0.91	0.96	0.90
High Anxiety / High IP	0.71	0.93	0.87	0.92	0.96	0.84
Low Anxiety / Low IP	0.80	0.88	0.91	0.95	0.97	0.79
Low Anxiety / High IP	0.84	0.90	0.84	0.89	0.94	0.75

Table 6: Parameter recovery for 148 trials. We report Spearman Correlation Coefficients. IP = Indifference Point.

5.8 Model Selection

We fitted all three models using both Maximum Likelihood Estimation and Hirarchical Bayesian methods. We compared them using the Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC) for the parameters fitted using MLE and we compared them using Leave One Out Information Criterion (LOOIC) for the HB fitting. As it can be seen in table 7, Model 1 is the best model according to both the AIC, BIC and LOOIC scores and it is the model used for the rest of this analysis.

	MLE		HB
	AIC	BIC	LOOIC
Model 1 (allP)	**46084.59**	**13768.63**	**13065.12**
Model 2 (noRA)	54758.99	21868.35	14234.04
Model 3 (noLA)	52286.11	19395.47	18819.73

Table 7: AIC, BIC and LOOIC scores for the three models. Lower is better.

5.9 Model agnostic analysis

All t-tests are two-sided with a significance level of $\alpha = 0.05$. All correlation coefficients are Pearson's r, unless stated otherwise. Bayesian t-tests and correlations have been carried out using JASP (JASP Team, 2023) version 0.17.0. For multiple comparisons tests, we report "native" p-values together with the Bonferroni corrected p-values. We applied Bonferroni correction for each family of tests we performed: 18 tests related to reaction times (corrected $\alpha = 0.0027$) and 8 tests related to individual choice data (corrected $\alpha = 0.00625$).

A COVID Questionnaire

All items were scored between 1 to 5, we report the questions and the different answers within square brackets. The dots (•) indicate the questions about past behaviours, the asterisks (*) indicate the questions about future behaviours and the dashes (-) indicate the questions about general attitude towards COVID. We group these questions in COVID-Past (•), COVID-Future (*) and COVID-General (-) scores.

- • How many COVID vaccinations have you had? [0, 1, 2, 3, 4+]

- * Would you get another booster dose if offered? [Very Unlikely, ..., Very Likely]

- • What type of mask did you usually wear throughout the pandemic? [No Mask, Scarf/Bandana/Other, Reusable Fabric Mask, Surgical/Disposable Mask, Respirator (N99/N95)]

- * What type of mask would you wear more often in case of a new surge in COVID cases? [No Mask, Scarf/Bandana/Other, Reusable Fabric Mask, Surgical/Disposable Mask, Respirator (N99/N95)]

- • How anxious were you about coronavirus at the beginning of the pandemic in March 2020? [Not at all Anxious, ..., Extremely Anxious]

- * How anxious are you about a new increase in COVID cases in the upcoming Winter? [Not at all Anxious, ..., Extremely Anxious]

- • How closely have you followed the government rules and guidelines during the pandemic? [I have not followed any of the rules, ..., I have followed all of the rules]

- * In case of a hypothetical new pandemic how closely would you follow new government rules and guidelines? [I would not follow any of the rules, ..., I would follow all of the rules]

- - I think Coronavirus vaccinations should be mandatory. [Strongly Disagree, ..., Strongly Agree]

- - During the pandemic I wore a mask and social distanced as often as I could. [Strongly Disagree, ..., Strongly Agree]

- - I think that Coronavirus restrictions introduced in 2020 were necessary and important. [Strongly Disagree, ..., Strongly Agree]

- - I am worried or have been worried about coronavirus affecting my health. [Strongly Disagree, ..., Strongly Agree]

475   - I am worried or have been worried about coronavirus affecting the health
476   of my friends and family. [Strongly Disagree, ..., Strongly Agree]

B   Exclusion Criteria

Different types of exclusion criteria have been used. During the gambling part of the experiment, 4 trials were used as attention checks. In these trials the subjects had to choose between a win of £0 and a loss in the range £[-30,-5] together with a sure option of £0. We discarded participants who chose the gambling option for 2 or more of these 4 attention checks. Another attention check was included in the questionnaire part of the experiment. We removed 22 participants who failed 2 or more gambling attention checks whereas no participant failed the questionnaire attention check. Four participants were removed due to incomplete demographic information.

We also removed participants who did not show any differences in percentage of gambling as a function of the trials' expected values. We remove participants which have a difference in average percentage of gamble lower than 5% between the upper half of expected values and the lower half of expected values. Participants should be able to adapt their gambling to the different expected values of the trials (i.e., higher expected values should lead to higher percentage of gambles). An example of this can be seen in fig. 4. We removed 22 participants using this criteria.

In total we conducted the analysis on 115 participants.

Figure 4: On the left, a participants which correctly adapted their choices to the expected value of the trials. On the right, a participant which incorrectly reduces their percentage of gambles for gambles with higher expected values. The participant on the right has been removed from further analysis.

C Parameter Recovery

Parameter recovery was conducted simulating 100 subjects for each combination of parameters/indifference points. Parameters were sampled from Gaussian distributions using the fitted parameters reported by Charpentier et al. (2017) (reported here in table 8). Indifference points were sampled from Gaussian distributions with SD = 2 and means as follows: High IP mixed gamble = 8, High IP gain only = 4, Low IP mixed gamble = 2, Low IP gain only = 1.

	Low Anxiety	High Anxiety
λ	2.067 (0.752)	2.013 (0.625)
ρ	0.875 (0.537)	0.713 (0.458)
μ	3.857 (4.502)	5.119 (4.124)

Table 8: Parameters from Charpentier et al. (2017) used for the parameter recovery. Mean (Standard deviation).

The results for the parameter recovery are reported in table 9. Due to outliers in the MLE fitting of the data we report the more robust Spearman correlation coefficients.

222 trials	MLE			HB		
	λ	ρ	μ	λ	ρ	μ
High Anxiety / Low IP	0.82	0.87	0.95	0.93	0.99	0.94
High Anxiety / High IP	0.77	0.91	0.78	0.89	0.97	0.92
Low Anxiety / Low IP	0.77	0.90	0.93	0.95	0.98	0.92
Low Anxiety / High IP	0.64	0.87	0.90	0.87	0.95	0.74

296 trials	MLE			HB		
	λ	ρ	μ	λ	ρ	μ
High Anxiety / Low IP	0.85	0.97	0.94	0.94	0.97	0.97
High Anxiety / High IP	0.82	0.96	0.87	0.95	0.98	0.93
Low Anxiety / Low IP	0.81	0.93	0.89	0.98	0.99	0.90
Low Anxiety / High IP	0.89	0.92	0.87	0.86	0.96	0.82

Table 9: Parameter recovery for 222 and 296 trials. We report Spearman Correlation Coefficients. IP = Indifference Point.

D Single Prior Model

Across all participants, the mean (and standard deviation) of the loss aversion parameter λ is 2.323(0.975), the risk aversion parameter ρ is 0.933(0.283) and the inverse temperature μ is 1.568(1.526). We report the fitted parameters for this model in table 10 and in fig. 5.

Single prior	Loss Aversion λ	Risk Aversion ρ	Inverse Temperature μ
All participants	2.323 (0.975)	0.933 (0.283)	1.568 (1.526)
Low Anxiety	2.156 (0.971)	0.924 (0.296)	1.829 (1.748)
High Anxiety	2.510 (0.955)	0.943 (0.270)	1.274 (1.178)

Anxiety prior	Loss Aversion λ	Risk Aversion ρ	Inverse Temperature μ
All participants	2.318 (0.967)	0.931 (0.280)	1.573 (1.546)
Low Anxiety	2.125 (0.935)	0.915 (0.296)	1.889 (1.797)
High Anxiety	2.537 (0.966)	0.949 (0.264)	1.217 (1.116)

Table 10: Mean (standard deviation) of the fitted parameters of Model 1 using a single prior (top table) and a GAD-7 score median split prior.

We find a near significant group difference between the low and high anxiety groups in the single prior model in loss aversion λ ($t_{113} = 1.964$, $p = 0.051$), with the more anxious subjects showing higher levels of loss aversion, and in inverse temperature μ ($t_{113} = -1.970$, $p = 0.051$), with the more anxious subjects having lower values of μ. In the model fitted using the anxiety priors we find the same group differences, but now statistically significant ($\lambda : t_{113} = 2.317$, $p = 0.022$; $\mu : t_{113} = -2.371$, $p = 0.019$). There was no difference in risk aversion across the two groups in either models (single prior $\rho : t_{113} = 0.348$, $p = 0.727$; anxiety prior $\rho : t_{113} = 0.639$, $p = 0.523$).

Risk and loss aversion are not correlated across individuals ($r(115) = 0.004$, $p = 0.962$), while inverse temperature is negatively correlated with both risk ($r(115) = -0.624, p < 0.001$) and loss aversion ($r(115) = -0.333, p < 0.001$). In the single prior model, GAD-7 scores are near significantly correlated to loss aversion ($r_{115} = 0.168$, $p = 0.072$) and inverse temperature ($r_{115} = -0.166$, $p = 0.075$). In the anxiety prior model, GAD-7 scores are significantly correlated with loss aversion ($r_{115} = 0.192$, $p = 0.039$) and inverse temperature ($r_{115} = -0.196$, $p = 0.035$).

Figure 5: Parameters distributions for the single prior model.

References

Ahn, W.-Y., Haines, N., and Zhang, L. (2017). Revealing Neurocomputational Mechanisms of Reinforcement Learning and Decision-Making With the hBayesDM Package. *Computational Psychiatry*, 1:24–57.

Ahn, W.-Y., Vasilev, G., Lee, S.-H., Busemeyer, J. R., Kruschke, J. K., Bechara, A., and Vassileva, J. (2014). Decision-making in stimulant and opiate addicts in protracted abstinence: Evidence from computational modeling with pure users. *Frontiers in Psychology*, 5.

Aylward, J., Valton, V., Ahn, W.-Y., Bond, R. L., Dayan, P., Roiser, J. P., and Robinson, O. J. (2019). Altered learning under uncertainty in unmedicated mood and anxiety disorders. *Nature Human Behaviour*, 3(10):1116–1123.

Bishop, S. J. and Gagne, C. (2018). Anxiety, Depression, and Decision Making: A Computational Perspective. *Annual Review of Neuroscience*, 41(1):371–388.

Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J., Li, P., and Riddell, A. (2017). Stan: A probabilistic programming language. *Journal of statistical software*, 76(1).

Charpentier, C. J., Aylward, J., Roiser, J. P., and Robinson, O. J. (2017). Enhanced Risk Aversion, But Not Loss Aversion, in Unmedicated Pathological Anxiety. *Biological Psychiatry*, 81(12):1014–1022.

de Leeuw, J. R. (2015). jsPsych: A JavaScript library for creating behavioral experiments in a Web browser. *Behavior Research Methods*, 47(1):1–12.

Doi, S., Ito, M., Takebayashi, Y., Muramatsu, K., and Horikoshi, M. (2018). Factorial Validity and Invariance of the 7-Item Generalized Anxiety Disorder Scale (GAD-7) Among Populations With and Without Self-Reported Psychiatric Diagnostic Status. *Frontiers in Psychology*, 9:1741.

Gelman, A. and Rubin, D. B. (1992). Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science*, 7(4).

Giorgetta, C., Grecucci, A., Zuanon, S., Perini, L., Balestrieri, M., Bonini, N., Sanfey, A. G., and Brambilla, P. (2012). Reduced risk-taking behavior as a trait feature of anxiety. *Emotion*, 12(6):1373–1383.

Hartley, C. A. and Phelps, E. A. (2012). Anxiety and Decision-Making. *Biological Psychiatry*, 72(2):113–118.

Hudson, A. and Montelpare, W. J. (2021). Predictors of Vaccine Hesitancy:

Implications for COVID-19 Public Health Messaging. *International Journal of Environmental Research and Public Health*, 18(15):8054.

JASP Team (2023). JASP (Version 0.17)[Computer software].

Kahneman, D. and Tversky, A. (1979). Prospect Theory: An Analysis of Decision under Risk. *Econometrica*, 47(2):263.

Lorian, C. N., Titov, N., and Grisham, J. R. (2012). Changes in risk-taking over the course of an internet-delivered cognitive behavioral therapy treatment for generalized anxiety disorder. *Journal of Anxiety Disorders*, 26(1):140–149.

Maner, J. K., Richey, J. A., Cromer, K., Mallott, M., Lejuez, C. W., Joiner, T. E., and Schmidt, N. B. (2007). Dispositional anxiety and risk-avoidant decision-making. *Personality and Individual Differences*, 42(4):665–675.

Millroth, P. and Frey, R. (2021). Fear and anxiety in the face of COVID-19: Negative dispositions towards risk and uncertainty as vulnerability factors. *Journal of Anxiety Disorders*, 83:102454.

Mueller, E. M., Nguyen, J., Ray, W. J., and Borkovec, T. D. (2010). Future-oriented decision-making in Generalized Anxiety Disorder is evident across different versions of the Iowa Gambling Task. *Journal of Behavior Therapy and Experimental Psychiatry*, 41(2):165–171.

Phelps, E. A., Lempert, K. M., and Sokol-Hessner, P. (2014). Emotion and Decision Making: Multiple Modulatory Neural Circuits. *Annual Review of Neuroscience*, 37(1):263–287.

Pike, A. C., Alves Anet, Á., Peleg, N., and Robinson, O. J. (2023). Catastrophizing and Risk-Taking. *Computational Psychiatry*, 7(1):1.

Raymond, J. G., Steele, J. D., and Seriès, P. (2017). Modeling Trait Anxiety: From Computational Processes to Personality. *Frontiers in Psychiatry*, 8.

Roberts, I. D. and Hutcherson, C. A. (2019). Affect and Decision Making: Insights and Predictions from Computational Models. *Trends in Cognitive Sciences*, 23(7):602–614.

Schonberg, T., Fox, C. R., and Poldrack, R. A. (2011). Mind the gap: Bridging economic and naturalistic risk-taking with cognitive neuroscience. *Trends in Cognitive Sciences*, 15(1):11–19.

Sherman, S. M., Smith, L. E., Sim, J., Amlôt, R., Cutts, M., Dasch, H., Rubin, G. J., and Sevdalis, N. (2021). COVID-19 vaccination intention in the UK: Results from the COVID-19 vaccination acceptability study (CoVAccS),

a nationally representative cross-sectional survey. *Human Vaccines & Immunotherapeutics*, 17(6):1612–1621.

Shou, Y., Smithson, M., Gulliver, A., Murray, K., Banfield, M., Rodney Harris, R. M., McCallum, S. M., Farrer, L. M., Calear, A. L., and Batterham, P. J. (2022). Risk tolerance and changes in coronavirus disease (COVID) related health behaviors: A longitudinal study. *Health Psychology*, 41(8):507–518.

Sokol-Hessner, P. and Rutledge, R. B. (2019). The Psychological and Neural Basis of Loss Aversion. *Current Directions in Psychological Science*, 28(1):20–27.

Spielberger, C. D. (1977). State-Trait Anxiety Inventory for Adults.

Spitzer, R. L., Kroenke, K., Williams, J. B. W., and Löwe, B. (2006). A Brief Measure for Assessing Generalized Anxiety Disorder: The GAD-7. *Archives of Internal Medicine*, 166(10):1092.

Valton, V., Wise, T., and Robinson, O. J. (2020). Recommendations for Bayesian hierarchical model specifications for case-control studies in mental health. *arXiv:2011.01725 [cs, stat]*.

Wake, S., Wormwood, J., and Satpute, A. B. (2020). The influence of fear on risk taking: A meta-analysis. *Cognition and Emotion*, 34(6):1143–1159.

Wise, T., Zbozinek, T. D., Charpentier, C. J., Michelini, G., Hagan, C. C., and Mobbs, D. (2022). Computationally-defined markers of uncertainty aversion predict emotional responses during a global pandemic. *Emotion*.

Xu, P., Van Dam, N. T., van Tol, M.-J., Shen, X., Cui, Z., Gu, R., Qin, S., Aleman, A., Fan, J., and Luo, Y.-j. (2020). Amygdala–prefrontal connectivity modulates loss aversion bias in anxious individuals. *NeuroImage*, 218:116957.