Dysfunctional effector memory CD8 T cells in the bronchoalveolar compartment of people

living with HIV

Maphe Mthembu[1,2], Helgard Claassen[1,2], Sharon Khuzwayo[1,2,3], Valentin Voillet[3], Anneta

Naidoo[3], Kennedy Nyamande[4], Dilshaad F. Khan[4], Priya Maharaj[4], Mohammed Mitha[4], Zoey

Mhlane[1], Farina Karim[1,2], Erica Andersen-Nissen[3], Thumbi Ndung'u[1,2,5,7], Gabriele Pollara*[5,]

and Emily B. Wong*[1,5,10]

* contributed equally to this work

[1]Africa Health Research Institute, Durban, South Africa,

[2]University of KwaZulu-Natal, Medical School, Durban, South Africa,

[3]Cape Town HVTN Immunology Laboratory, Cape Town, South Africa,

[4]Department of Pulmonology, Inkosi Albert Luthuli Central Hospital, Durban, South Africa,

[5]Division of Infection and Immunity, University London College, London, UK,

[6]HIV Pathogenesis Programme, The Doris Duke Medical Research Institute, University of

KwaZulu-Natal, Durban, South Africa,

[7]Ragon Institute of MGH, MIT and Harvard, Cambridge, USA,

[10]Department of Medicine, The University of Alabama at Birmingham, Birmingham, AL, USA.

Correspondence: Dr Emily Wong, Africa Health Research Institute, Durban, South Africa

(emily.wong@ahri.org) and Dr Gabriele Pollara, University London College, London, UK,

(g.pollara@ucl.ac.uk)

Body word count: (1990 words) Target = 2000

Abstract word count: (98 words) Target = 100

Abstract:

Mechanisms by which HIV causes susceptibility to respiratory pathogens remain incompletely understood. We obtained whole blood and bronchoalveolar lavage (BAL) from people with latent TB infection in the presence or absence of antiretroviral-naïve HIV co-infection. Transcriptomic and flow cytometric analyses demonstrated HIV-associated cell proliferation plus type I interferon activity in blood and effector memory CD8 T-cells in BAL. Both compartments displayed reduced induction of CD8 T-cell-derived IL-17A in people with HIV, associated with elevated T-cell regulatory molecule expression. The data suggest that dysfunctional CD8 T-cell responses in uncontrolled HIV contribute to susceptibility to secondary bacterial infections, including tuberculosis.

Keywords:

HIV, BAL, blood, whole transcriptomics, flow cytometry

Introduction

People living with HIV (PLWH), have >18-fold increased risk of respiratory infections, including tuberculosis (TB) (1). Uncontrolled HIV infection results in several immune perturbations, including CD4 T-cell depletion, cytotoxic CD8 T-cell expansion (2) and elevated cytokine activity (3) (e.g. type I interferons (4)). Most studies have focused on immune perturbations in blood but less on target organs for co-infections, such as the lungs. Improved understanding of HIV's impact at the site of host-pathogen interactions may offer insights into immune modulation caused by uncontrolled HIV infection, and in turn, identifying putative therapeutic targets in PLWH. Whole compartment transcriptional profiling permits unbiased assessments of disease-mediated perturbation by HIV at the molecular level, whilst still retaining the ability to deconvolute differences in immune cell frequency, cytokine activity and other functions (5, 6).

In this study, we aimed to explore the lung mucosal immune landscape, through transcriptomic profiling of bronchoalveolar lavage (BAL) cells in individuals with untreated HIV infection. We previously established the feasibility of this technique in a small cohort (7). The current study extends this approach in a population with evidence of *Mycobacterium Tuberculosis* (*Mtb*) infection (i.e. latent TB) in a high TB transmission setting. We test the hypothesis that HIV results in compartment-specific immune perturbations that may underlie the susceptibility to TB disease and other respiratory infections.

Methods

Study subjects and samples collection

We recruited otherwise healthy individuals grouped by HIV serostatus, from a previously

described research bronchoscopy cohort in Durban, South Africa (7). Inclusion criteria included

a positive QuantiFERON-TB Gold Plus for all participants and, for HIV seropositive individuals,

no previous exposure to antiretroviral therapy (ART). Paired blood and BAL samples were

obtained from all study participants, and peripheral blood mononuclear cells (PBMCs) and BAL

mononuclear cells (BLCs) isolated (Supplementary Methods).

Flow cytometry

Mononuclear cells from both compartments were stained with monoclonal antibodies and some

were stimulated with a non-specific mitogen and further stained for intracellular cytokines. All

cells were then acquired, and some subsets sorted through BD FACS™ Aria III (detailed in

Supplementary Methods).

Transcriptional profiling

RNA was extracted from cryopreserved whole compartment BAL and blood samples, as well as

sorted cell populations in BAL. cDNA libraries were prepared and sequenced on an Illumina

NextSeq 500 platform (details in Supplementary Methods).

RNA sequences were quantified, aligned, and annotated to measure and deduce patterns of gene

expression. Whole-genome differential gene expression was analysed. Transcriptional module

expression was quantified as previously described (5), and differential module enrichment

between groups was assessed using Mann-Whitney test (Supplementary Methods).

<u>Results</u>

Thirty-seven participants contributed paired samples, 70% female, with median age 32 (IQR 25-

35, Supplementary Table 1). Sex and age distribution were comparable between PLWH and

those without HIV. Median CD4 count in PLWH was 513 cells/mm^3 (IQR 275-681) and median

VL 25,229 copies/mL (IQR 13,987-33,565).

Unsupervised gene expression analyses identified 139 upregulated and 44 downregulated genes

in blood of PLWH (Figure 1A, left), in contrast to 13 upregulated and 1 downregulated, in BAL

(Figure 1A, right). Of the 152 genes elevated in PLWH across both compartments, only 3 (2%)

were increased in both blood and BAL (Figure 1B), highlighting the compartmentalized nature

of HIV-induced transcriptional changes. HIV-associated changes in blood mapped to cell

proliferation, interferon signaling and antigen presentation pathways, whereas in BAL,

immunoregulatory pathways were enriched (Figure 1C). Cytokines predicted to regulate

upregulated genes in HIV also demonstrated compartment-specific effects, with only 3/12

cytokines (IL-2, IL-15 and IL-12B) upstream of transcripts elevated by HIV in both

compartments (Figure 1D).

BAL was predominantly composed of alveolar macrophages (Supplementary Figure 1), but

several of the 13 genes overexpressed in PLWH were lymphocyte-associated (Figure 1A, right).

Validated transcriptional modules that quantify the relative abundance of immune cells

demonstrated relative depletion of CD4 T cell transcripts in both blood and BAL in PLWH (5)

(Figure 2A, left), an expansion of CD8 T-cells in both compartments and no differences in NK

cell frequency (Figure 2A, left). These modules were derived from peripheral blood cells, and

may not accurately reflect the transcriptional state of BAL lymphocytes (5). Therefore, we

generated an independent set of modules derived from genes differentially expressed in sorted

BAL cells (gating strategy in Supplementary Figure 2). The BAL-derived module specificity was

comparable to those from blood, with better discrimination of CD8 T- cells from MAIT cells

(Supplementary Figure 3). These BAL-derived modules confirmed enrichment of CD8 T-cells,

but not MAIT cells, in PLWH (Figure 2A, right). Flow cytometry confirmed elevated CD8 T-

cell frequency in HIV (Supplementary Figure 4), in turn validating our approach to use

transcriptional modules to quantify cell enrichment from BAL and blood transcriptomes.

Several of the 13 genes with higher expression in BAL of PLWH are associated with CD8 T-cell

effector functions (*GZMK, GZMH, B3GAT1, NKG7*) and we sought to test the hypothesis that

this increased expression reflects an increased frequency of effector memory (EM) CD8 T-cells.

We generated and validated new transcriptional modules from different CD8 T-cell functional

states (Supplementary Figure 5) and they showed enrichment of CD8 T-cells with an EM

phenotype in BAL of PLWH, with no differences in central memory (CM) CD8 T-cells (Figure

2B, left column). To determine whether EM CD8 T-cells were the primary source of

significantly upregulated genes in BAL of PLWH, we sought to adjust for differences in cell

frequency by performing a linear regression at the whole transcriptome level, controlling for

cellular module expression in each sample. This revealed that differential expression of the 13

genes elevated in BAL of PLWH was no longer present following regression against either

blood/BAL CD8 T-cell modules or the CD8 EM module, but this was not the case with CD8

CM, CD4, NK or MAIT cell modules (Figure 2B, right column). We interpreted these data to

suggest transcriptomic perturbations in untreated PLWH in BAL are likely attributable to

increased numbers of EM CD8 T-cells.

Finally, we sought to determine whether the differential frequency of CD8 T-cells in

unstimulated BAL in HIV was also associated with different functional activity of these cells in

HIV. We quantified production of IFN-γ and IL-17A by both blood and BAL CD8 T-cells before

and after stimulation with PMA/ionomycin (Supplementary Figure 6A). Baseline cytokine

expression was low and did not differ by HIV status (Supplementary Figure 6B), but following

stimulation, we observed a decreased frequency of CD8 T-cells expressing IL-17A in both blood

and BAL compartments of PLWH, whereas no such difference was observed for IFN-γ (Figure

2C). Notably, 3/13 of differentially upregulated genes in BAL of PLWH (*KLRG1, EOMES* and

TIGIT), along with *TNFRSF9* and *SLAMF6*, are associated with CD8 T-cell exhaustion or

functional inhibition (8) (Figure 2D). In blood of PLWH, we observed more CD8 T-cell

exhaustion markers, *PDCD1* and additionally *TIGIT, SLAMF6*, also showing increased

expression indicating a compartment-specific effect (Figure 2D). Flow cytometry analysis

confirmed a significant increase in the expression of PD-1 (encoded by the gene *PDCD1*) on

blood CD8 T-cells in PLWH. A higher expression of PD-1 on CD8 T-cells in PLWH was seen in

BAL but this was not statistically significant (Supplementary Figure 6C). Overall, these results

suggest increased immunoregulatory activity in HIV infection that may attenuate IL-17A

responses of CD8 T-cells in both blood and BAL.

Discussion:

HIV infection increases the risk of respiratory infections including TB (1). Of the mechanisms

proposed to date, most have not included systems-level analysis of the immunobiology in lungs.

We demonstrate that uncontrolled HIV infection results in transcriptional changes in the lungs

that differ from those in blood. The BAL of PLWH is enriched for CD8 T-cells that

transcriptionally resemble EM cells but these show deficient IL-17A responses upon stimulation,

a process associated with elevated expression of immunoregulatory molecules. Our data support

a model whereby impaired CD8 T-cell production of IL-17A may contribute to susceptibility to

secondary bacterial infections in HIV infection (9).

We extend our earlier report in which we first reported bulk transcriptomic assessments from

BAL, by including a larger number of participants in whom active lung disease (other than HIV)

was rigorously ruled out, and all of whom had evidence of past exposure to *Mtb*. As expected,

several pathways were upregulated in the peripheral blood compartment of PLWH including cell

proliferation, possibly reflecting a homeostatic response to the depletion of CD4 T-cells in the

HIV infection (10), and type 1 and II IFN signaling antiviral pathways. In contrast, we observed

markedly fewer differentially expressed genes in the BAL compartment of PLWH compared to

HIV-negative controls. This may have reflected the predominance of macrophages in BAL,

which do not show transcriptomic perturbation in PLWH at baseline and may have reduced the

sensitivity to detect effects on other cell types. Nevertheless, the few genes with elevated

expression in PLWH were markedly associated with lymphocyte function, which drew us to

focus on T-cell perturbations in the BAL compartment.

Using module analysis as an approach to bioinformatically de-convolute whole-genome

transcriptomic data, we confirmed the well-known consequences of HIV infection in depleting

CD4 T-cells and enriching CD8 T-cells in both blood and BAL (11). The increased frequency of

CD8 T-cells in BAL is consistent with reports of lymphatic alveolitis (12) and was associated

with elevated expression of several genes linked to cytotoxicity, such as granzymes H and K, and

a transcriptional module reflective of EM CD8 T-cells, but not CM CD8 T-cells, NK or MAIT

cells. Nevertheless, both blood and BAL CD8 T-cells showed diminished IL-17A production

after ex vivo stimulation, but no impairment in IFN-γ production. This IL-17A defect has been

reported in the blood of virally suppressed people with HIV (13), but our current study shows

that this is also evident in the airways of people with untreated HIV.

IL-17A is reported to confer protection against hypervirulent *Mtb* strains such as HN878 (14)

and this may contribute to susceptibility to TB disease in ARV-naïve PLWH (15). Nevertheless,

given that active TB is characterized by exaggerated IL-17A responses that may contribute to

tissue pathology (6), our observations may also provide an explanation for why severe chest-X-

ray abnormalities, such as cavities, are less common in TB patients with advanced HIV (15).

Notably, we observed the defect in IL-17A production in PLWH to be associated with increased

expression of molecules associated with T-cell immunomodulation, suggesting an exhausted

CD8 T-cell phenotype. Indeed, it has been proposed that elevated expression of inhibitory

receptors is responsible for diminished IL-17A responses in HIV (13). Our findings suggest that

modulation of negative regulators of T-cell function, with precisely targeted biologics, could be

one strategy to restore HIV-induced dysregulated immunity in the lung mucosa.

Our study has several limitations. First, this was a single-center study with a limited sample size

that assessed individuals at a single timepoint. Participants with HIV were all ART-naïve, and

we cannot predict if the immune perturbations observed would persist during suppression of HIV

replication by ART. Additionally, assessment of unstimulated transcriptomes at bulk cell level

precluded assessment of the dynamic nature of responses in response to bacterial stimulation.

Finally, we only investigated a limited number of cytokine responses at the protein level in blood

and BAL cells, leaving unanswered whether the observed defects in IL-17A responses are

associated with alterations to a larger set of cytokine responses in HIV still able to leverage.

Our study leveraged unique access to both blood and BAL samples of a well-defined cohort of

individuals to unmask immune dysregulation associated with uncontrolled HIV infection at the

site of pulmonary host-pathogen interactions. We conclude that HIV is associated with divergent

transcriptional effects in the blood and in the bronchoalveolar compartments. Blood was

characterized by elevated cell proliferation and type I/II interferon signaling activity, while BAL

was enriched for CD8 T-cells with an effector phenotype but a defect in inducible IL-17A

production, associated with increased expression of T-cell inhibitory molecules. Future work will

need to assess the ability of ART to reverse this immune dysregulation and to explore the role of

adjunctive immune checkpoint inhibitors to restore mucosal CD8 T-cell function and improve

immune protection against respiratory infections in HIV.

Data availability

Supplementary materials are provided here, consisting of data provided by the authors to benefit

the reader. The original RNA-Seq datasets generated for this study can be found in the Gene

Expression Omnibus (GEO) repository, accession number GSE230738 for whole sample

transcriptomics and GSE231628 for mini-population transcriptomics. The supplementary

materials are not copyedited and are the authors' sole responsibility, so questions or comments

should be addressed to the corresponding authors.

Acknowledgments

Financial Support

This work was supported by the National Institutes of Health (K08 AI118538) and the Burroughs Wellcome Fund Pathogenesis of Infectious Diseases Award (1022002) and a University College London Global Partnership grant for collaboration between University College London and the Africa Health Research Institute. Additional support was received from the Strategic Health Innovation Partnerships (SHIP) Unit of the South African Medical Research Council with funds received from the South African Department of Science and Innovation as part of a bilateral research collaboration agreement with the Government of India. Supplementary support was received through the Sub-Saharan African Network for TB/HIV Research Excellence (SANTHE), a DELTAS Africa Initiative (grant no. DEL-15-006). The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (grant no. 107752/Z/15/Z) and the UK government. Other support was acquired through the South Africa Research Chairs Initiative, the Victor Daitz Foundation, a Burroughs-Wellcome Fund/American Society of Tropical Medicine and Hygiene fellowship (EBW). GP's time was funded by the UCLH NIHR Biomedical Research Centre. Research reported in this

publication was also supported by the National Institute of Allergy and Infectious of the National

Institutes of Health under award number [UM1 AI068618] (EAN). The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health.

Potential conflict of interest

No conflicts were reported by any of the co-authors.

References:

1.   World Health Organisation. Global Tuberculosis Report. 2020.

2.   Smit-McBride Z, Mattapallil JJ, McChesney M, Ferrick D, Dandekar S. Gastrointestinal

T lymphocytes retain high potential for cytokine responses but have severe CD4(+) T-cell

depletion at all stages of simian immunodeficiency virus infection compared to peripheral

lymphocytes. J Virol. 1998;72(8):6646-56.

3.   Hazenberg MD, Otto SA, van Benthem BH, Roos MT, Coutinho RA, Lange JM, et al.

Persistent immune activation in HIV-1 infection is associated with progression to AIDS. Aids.

2003;17(13):1881-8.

4.   Boasso A, Shearer GM. Chronic innate immune activation as a cause of HIV-1

immunopathogenesis. Clin Immunol. 2008;126(3):235-42.

5.   Pollara G, Murray MJ, Heather JM, Byng-Maddick R, Guppy N, Ellis M, et al.

Validation of Immune Cell Modules in Multicellular Transcriptomic Data. PLOS ONE.

2017;12(1):e0169271.

6. Pollara G, Turner CT, Rosenheim J, Chandran A, Bell LCK, Khan A, et al. Exaggerated

IL-17A activity in human in vivo recall responses discriminates active tuberculosis from latent

infection and cured disease. Sci Transl Med. 2021;13(592).

7. Muema DM, Mthembu M, Schiff AE, Singh U, Corleis B, Chen D, et al. Contrasting

Inflammatory Signatures in Peripheral Blood and Bronchoalveolar Cells Reveal Compartment-

Specific Effects of HIV Infection. Frontiers in Immunology. 2020;11(864).

8. Wherry EJ, Kurachi M. Molecular and cellular insights into T cell exhaustion. Nature

reviews Immunology. 2015;15(8):486-99.

9. Mills KHG. IL-17 and IL-17-producing cells in protection versus pathology. Nature

Reviews Immunology. 2023;23(1):38-54.

10. Kim M, Siliciano RF. Reservoir expansion by T-cell proliferation may be another barrier

to curing HIV infection. Proceedings of the National Academy of Sciences. 2016;113(7):1692-4.

11. Catalfamo M, Wilhelm C, Tcheung L, Proschan M, Friesen T, Park JH, et al. CD4 and

CD8 T cell immune activation during chronic HIV infection: roles of homeostasis, HIV, type I

IFN, and IL-7. Journal of immunology (Baltimore, Md : 1950). 2011;186(4):2106-16.

12. Twigg HL, Soliman DM, Day RB, Knox KS, Anderson RJ, Wilkes DS, et al.

Lymphocytic alveolitis, bronchoalveolar lavage viral load, and outcome in human

immunodeficiency virus infection. American journal of respiratory and critical care medicine.

1999;159(5 Pt 1):1439-44.

13. Perdomo-Celis F, Feria MG, Taborda NA, Rugeles MT. A Low Frequency of IL-17-

Producing CD8(+) T-Cells Is Associated With Persistent Immune Activation in People Living

With HIV Despite HAART-Induced Viral Suppression. Front Immunol. 2018;9:2502.

14.   Gopal R, Monin L, Slight S, Uche U, Blanchard E, A. Fallert Junecko B, et al.

Unexpected Role for IL-17 in Protective Immunity against Hypervirulent Mycobacterium

tuberculosis HN878 Infection. PLOS Pathogens. 2014;10(5):e1004099.

15.   Kwan Candice K, Ernst Joel D. HIV and Tuberculosis: a Deadly Human Syndemic.

Clinical Microbiology Reviews. 2011;24(2):351-76.

Figures:

Figure 1: HIV infection associated with different transcriptional profiles in blood

compared to the bronchoalveolar space. **A.** Whole genome differential gene expression

transcriptional analysis from 10 pairs of blood and bronchoalveolar lavage (BAL) samples from

participants with and without HIV (n = 20, HIV-negative (10) and HIV-positive (10)); volcano

plots indicate genes with significantly elevated expression in HIV infection in blood (red) and

BAL (blue) compartments. **B.** Dot plot displaying adjusted P values of genes with significantly

elevated expression in HIV-negative compared to HIV-positive in blood (red), BAL (blue) and

both (magenta) compartments. **C.** Heatmap showing REACTOME levels (white: lowest to black:

highest) of enriched pathways in PLWH transcriptome blood (left column) and BAL (right

column) samples. **D.** Heatmap displaying levels (white: lowest to blue: highest) of upstream

cytokines significantly predicted to regulate differential gene expression in PLWH blood and

BAL samples.

Figure 2: Bronchoalveolar compartment in HIV infection characterized by enrichment of

effector memory CD8 T-cells deficient in IL-17A production. Transcriptional module

analysis of the whole transcriptomics data using **A.** Blood and BAL-derived modules reflecting

immune cell relative frequency, and **B.** effector and central memory CD8 T-cell modules **(left**

column) and a heatmap showing adjusted p-values (blue = lowest, yellow = highest, and

asterisks representing the level of significance) for 13 genes upregulated in the BAL between

PLWH and those without HIV (first column) regressed against 7 transcriptional modules: CD8

T-cells, effector memory CD8 T-cells (EM), central memory CD8 T-cells (CM), CD4 T-cells,

NK cells and MAITs cells **(right column)**. **C.** Expression of IFN-γ and IL-17A following PMA-

Ionomycin stimulation of mononuclear cells from both blood and BAL (top row). **D.** Immune

checkpoint molecule expression in blood and BAL (middle and bottom rows) (n=20, 10 HIV

negative and 10 HIV positive).

Figure 1

Figure 2