The CD4 transmembrane GGXXG and juxtamembrane (C/F)CV+C motifs mediate pMHCII-specific signaling independently of CD4-Lck interactions

Mark S. Lee[1], Peter J. Tuohy[1], Caleb Y. Kim[1], Philip P. Yost[1], Katrina Lichauco[1], Heather L. Parrish[1], Koenraad Van Doorslaer[1, 2, 3, 4, 5, 6], and Michael S. Kuhns[1, 3, 4, 5, 6]

[1] Department of Immunobiology, The University of Arizona College of Medicine, Tucson, AZ 85724, USA.

[2] School of Animal and Comparative Biomedical Sciences, The University of Arizona, Tucson, AZ 85724, USA.

[3] Cancer Biology Graduate Interdisciplinary Program and Genetics Graduate Interdisciplinary Program, The University of Arizona, Tucson, AZ 85724, USA.

[4] The BIO-5 Institute, The University of Arizona, Tucson, AZ 85724, USA.

[5] The University of Arizona Cancer Center, Tucson, AZ, USA

[6] The Arizona Center on Aging, The University of Arizona College of Medicine, Tucson, AZ, USA

*Correspondence: vandoorslaer@arizona.edu, mkuhns@email.arizona.edu

Running title: The CD4 GGXXG and (C/F)CV+C motifs are key determinants of pMHCII-specific signaling.

Abstract
CD4$^+$ T cell activation is driven by 5-module receptor complexes. The T cell receptor
(TCR) is the receptor module that binds composite surfaces of peptide antigens
embedded within MHCII molecules (pMHCII). It associates with three signaling modules
(CD3$\gamma\varepsilon$, CD3$\delta\varepsilon$, and CD3$\zeta\zeta$) to form TCR-CD3 complexes. CD4 is the coreceptor
module. It reciprocally associates with TCR-CD3-pMHCII assemblies on the outside of a
CD4$^+$ T cells and with the Src kinase, Lck, on the inside. Previously, we reported that
the CD4 transmembrane GGXXG motif and cytoplasmic juxtamembrane (C/F)CV+C
motif found in eutherian CD4 (placental mammals) have constituent residues that
evolved under purifying selection. Mutating these motifs together increased CD4-Lck
association but reduced CD3ζ, Zap70, and Plcγ1 phosphorylation levels, as well as IL-2
production, in response to agonist pMHCII. Because these mutants preferentially
localized CD4-Lck pairs to non-raft membrane fractions, one explanation for our results
was that they impaired proximal signaling by sequestering Lck away from TCR-CD3. An
alternative hypothesis is that the mutations directly impacted signaling because the
motifs normally play a Lck-independent role in signaling. The goal of this study was to
discriminate between these possibilities. Our results indicate that: intracellular CD4-Lck
interactions are not necessary for pMHCII-specific signal initiation; the GGXXG and
(C/F)CV+C motifs are key determinants of CD4-mediated pMHCII-specific signal
amplification; the GGXXG and (C/F)CV+C motifs exert their functions independently of
direct CD4-Lck association. These data provide a mechanistic explanation for why
residues within these motifs are under purifying selection, and thus functionally
important for CD4$^+$ T cells in vivo. The results are also important to consider for
biomimetic engineering of synthetic receptors.

Introduction

Delineating the mechanistic principles by which multi-module receptors drive complex biological processes is important both for our broad understanding of receptor biology, and for guiding biomimetic engineering of synthetic receptors for therapeutic purposes. For example, comparing the conventional chimeric antigen receptor (CAR) framework utilized in CAR-T cell therapy, which was informed by an early 1990's understanding of receptor biology, with the multi-module receptors that naturally drive T cell responses suggests that conventional CARs integrate with the T cell intracellular signaling machinery differently than their natural counterparts (Harris and Kranz, 2016; Wu et al., 2020). These differences are likely to underlie differences in signaling output, leading us to posit that fully coopting T cell functions with synthetic receptors requires that they be designed to integrate with the T cell's intracellular signaling machinery in a way that mimics the native receptors (Kobayashi et al., 2020). Achieving this goal requires a more complete understanding of the multi-module pMHC-specific receptors that mediate antigen-specific T cell activation. Understanding how CD4$^+$ T cells respond is particularly important for informing biomimetic engineering of CARs given recent data showing that redirecting CD4$^+$ T cells with CARs is important for long-lived therapeutic efficacy (Melenhorst et al., 2022).

CD4$^+$ T cells are driven by 5-module receptors that recognize and drive responses to peptide antigens embedded within MHCII (pMHCII). Each naïve CD4$^+$ T cell expresses a clonotypic receptor module, called the T cell receptor (TCR), that binds specifically to unique features of composite pMHCII surfaces (Kuhns and Badgandi, 2012; Kuhns and Davis, 2012). The TCR lacks intracellular signaling domains and

instead assembles with three signaling modules (CD3$\gamma\varepsilon$, $\delta\varepsilon$, and $\zeta\zeta$) that have

immunoreceptor tyrosine-based activation motifs (ITAMs), and other motifs, to connect

TCR-pMHCII interactions to the intracellular signaling apparatus. CD4 is the coreceptor

module. It binds pMHCII in a reciprocal fashion with TCR-CD3 on the outside of CD4[+] T

cells and interacts with the Src kinase, Lck, via an intracellular CQC zinc clasp motif and

helix (Huse et al., 1998; Kim et al., 2003). According to the widely accepted TCR

signaling paradigm, co-engagement of pMHCII by TCR-CD3 and CD4 positions Lck and

the CD3 ITAMs in the proper spatial proximity to enable Lck phosphorylation of the CD3

ITAMs to initiate signaling (Rudd, 2021). However, when we tested this model directly,

we found that CD4-Lck interactions are not key determinants of CD3ζ ITAM

phosphorylation (pCD3ζ)(Lee et al., 2022). Based on our work, as well as work from

other labs, we consider the question of how intracellular pMHCII-specific CD4[+] T cell

signaling is initiated to be unresolved (Glassman et al., 2018; Horkova et al., 2023;

Killeen and Littman, 1993; Xu and Littman, 1993).

The answer to this question lies with how the 5 modules of pMHCII receptors

evolved over 435 million years to refine pMHCII-specific signaling. In our recent

publication, we computationally reconstructed CD4 evolution to identify residues and

motifs in the extracellular, transmembrane, and intracellular domains that are of

functional importance (Lee et al., 2022). We then performed structure function analysis

with mutants of key motifs to infer their function by evaluating the phenotypes of the

mutants. When we mutated the intracellular CQC clasp and IKRLL helix motifs that are

known to mediate CD4-Lck interactions, we observed significant reductions in CD4-Lck

association; yet we did not observe the expected impact on CD3ζ ITAM phosphorylation

(pCD3ζ) predicted by the TCR signaling paradigm. In contrast, when we mutated the

transmembrane GGXXG motif and cytoplasmic juxtamembrane (C/F)CV+C

palmitoylation motif, we observed an increased frequency of CD4-Lck pairs that were

preferentially localized to detergent soluble membrane domains. We also found

evidence of reduced CD3ζ, Zap70, and Plcγ1 phosphorylation (pCD3ζ, pZap70, and

pPlcγ1) compared with WT CD4. The reason that constituent residues of these motifs

evolved under purifying selection appears to be due to reduced signaling in response to

pMHCII when the motifs are altered. We took these data as evidence that the GGXXG

and (C/F)CV+C motifs arose in eutherians (placental mammals) to tightly regulate

pMHCII-specific signal initiation and postulated that the mutant CD4 molecules might

sequester Lck away from TCR-CD3 to prevent CD3ζ phosphorylation. However, an

alternative hypothesis is that the GGXXG and (C/F)CV+C motifs impact pCD3ζ levels

independently of CD4-Lck interactions.

Here we took a reductionist approach to test these hypotheses by studying the

impact of GGXXG and (C/F)CV+C motif mutants individually, or together, on proximal

pMHCII-specific signaling in C-terminally truncated CD4 molecules (T1). Working with

T1 eliminated motifs in the intracellular domain, including those that mediate CD4-Lck

association. As expected from prior work, truncating CD4 relieved CD4-Lck interactions

yet only slightly reduced IL-2 production (Lee et al., 2022). Importantly, there was no

impact on average pCD3ζ levels or percent of cells with phosphorylated CD3ζ, when

compared with wild-type (WT) CD4, despite significantly reduced levels of CD4-Lck

pairs. These data provide further evidence that the core tenet of the TCR signaling

paradigm, wherein CD4 recruits Lck to CD3 ITAMs initiates signaling, needs revising.

Importantly, T1 molecules bearing mutations in the GGXXG and (C/F)CV+C motifs

individually or together reduced IL-2 production as well as pCD3ζ levels and other TCR-

proximal signaling events. The simplest interpretation of these data is that the GGXXG

and (C/F)CV+C motifs are key determinants of pMHCII-specific signaling on their own,

independent of CD4-Lck interactions.

Results

The goal of this study was to evaluate if the transmembrane GGXXG and

juxtamembrane palmitoylation ((C/F)CV+C) motifs influence pMHCII-specific proximal

signaling on their own rather than by sequestering CD4-Lck pairs away from membrane

rafts (**Figure 1**). Accordingly, we used 58$\alpha^-\beta^-$ T cell hybridomas transduced to express

the 5c.c7 TCR along with either the WT or mutant CD4 constructs described in **Table 1**.

The well-characterized 5c.c7 TCR recognizes the moth cytochrome c peptide 88-103

(MCC) in the context of mouse MHCII I-E^k. We used this experimental system because

58$\alpha^-\beta^-$ cells were used in seminal work that helped form the foundation of the TCR

signaling paradigm (Glaichenhaus et al., 1991). In that study, a loss of CD4-Lck

interactions due to mutating the intracellular CQC clasp motif was linked with reduced

signaling output as measured by IL-2 production. Furthermore, IL-2 production by these

58$\alpha^-\beta^-$ cells is CD4 dependent, indicating that relevant signaling pathways are intact.

WT and T1 equivalently enhance pMHCII-specific proximal signaling. We

previously reported that the C-terminally truncated mouse CD4-T1 mutant (**T1: Table 1**

and Figure 1 arrow) had only a minor impact on IL-2 production relative to WT CD4

despite lacking most of the intracellular domain, including the CQC clasp and IKRLL

motifs shown to mediate CD4-Lck association (Lee et al., 2022). Because T1 maintains

both the transmembrane GGXXG and juxtamembrane (C/F)CV+C motifs, we reasoned

that it had utility for studying the contributions these motifs make to pMHCII-specific

signaling in the absence of CD4-Lck interactions.

Accordingly, we generated 5c.c7$^+$ 58$\alpha^-\beta^-$ cells expressing WT CD4, T1, or T1

combined with our previously reported Δbind mutant, which disrupts binding to pMHCII

in the CD4 D1 domain as well as signaling output (**Table 1 and Figure 2 – figure**

supplement 1)(Glassman et al., 2016; Glassman et al., 2018; Parrish et al., 2015). The

goals were to determine: 1) if T1 significantly impairs pMHCII-specific proximal signaling

events, relative to WT, as expected based on the TCR signaling paradigm because it

lacks the CQC and IKRLL motifs that mediate CD4-Lck interactions; 2) use T1$^{\Delta bind}$ to

determine the overall contribution of pMHCII-dependent assembly of CD4 with TCR-

CD3 to proximal signaling. For IL-2 production, we observed a slightly reduced

magnitude of the response that was consistent across a range of MCC peptide

concentrations for T1 compared to WT (**Figure 2A**). We also observed reduced

pMHCII-specific endocytosis for T1 compared to WT (**Figure 2 – figure supplement 2)**.

Importantly, the T1$^{\Delta bind}$ mutant did not produce IL-2 or undergo pMHCII-specific

endocytosis. These data indicate that, in our system, extracellular CD4-pMHCII

engagement is critical for IL-2 production while the CD4 intracellular domain C-terminal

of R422 can be deleted.

Next, we evaluated the impact of these mutations on proximal signaling events

measured by flow cytometry in response to antigen presenting cells (APCs) expressing

agonist MCC:I-E^k after background subtraction of signal levels in response to APCs

expressing null Hb:I-E^k (**Figure 2 – figure supplement 3**). Our analysis of five

independently generated pairs of WT and T1 cell lines showed variable difference in

pCD3ζ levels. For three pairs of lines, the T1 response was higher than the WT and for

two it was lower (**Figure 2B**). The averages of the T1 response normalized to the paired

WT controls showed no difference between the population. There were also no

differences for the average percent of T1 cells with response levels to MCC that were

above the null Hb peptide when normalized to their paired WT controls. Similar results

were observed for pZap70 and pPlcγ1 levels (**Figure 2 – figure supplement 4**). These

data, summarized in **Table 2,** provide additional evidence that CD4-Lck interactions via

the CQC clasp and IKRLL motifs are not key determinants of pMHCII-specific early

signal events or IL-2 production in this experimental system.

In contrast, the T1$^{\Delta bind}$ mutant significantly reduced pCD3ζ, pZap70, or pPlcγ1

levels compared to the T1 control but did not obviously impact the percent of cells

responding to agonist MCC:I-E^k (**Figure 2C and Figure 2 – figure supplement 5**).

Because the antibodies used to detect pCD3ζ, pZap70, and pPlcγ1 recognize single

phosphorylated tyrosines, these data indicate that CD4 enhances the number of TCR-

CD3 complexes per cell with phosphorylated CD3ζ molecules at Y142 (same in human

and mouse), as well as the number of phosphorylated Zap70 and Plcγ1 molecules per

cell, in a manner that is dependent on TCR-CD3 and CD4 co-engagement of pMHCII

but independent of the intracellular domain C-terminal of R422. A summary of these

results can be found in **Table 3**.

The GGXXG and (C/F)CV+C motifs influence IL-2 production independently of Lck.

Having established that T1 contributes to early pMHCII-specific signaling (e.g. pCD3ζ) as well as signaling output (i.e. IL-2), we generated 5c.c7$^+$ 58α$^-$β$^-$ cells expressing WT, T1, or the truncated T1 wherein the GGXXG and/or (C/F)CV+C motifs are also mutated (**Table 1 and Figure 3 – figure supplement 1**). To study the function of the GGXXG motif alone, we used our previously described TMD mutant (GGXXG to GVXXL) wherein bulky side chains replace the glycines that compose a flat surface on the transmembrane helix that could mediate protein:protein or protein:cholesterol interactions (Fessler, 2016; Parrish et al., 2015; Song et al., 2014; Teese and Langosch, 2015; Wacker et al., 2013). We called these mutants T1-TMD. To test the function of the palmitoylation motif, which contains the core CVRC (418-421) sequence in mouse and humans, we made note of a broader (C/F)CVRC motif in eutherians wherein the majority of ortholog sequences had CCVRC, as found in mouse, and a minority had FCVRC as found in humans. Importantly, we previously noted that position 417 is under purifying selection for all extant CD4 orthologs we analyzed by the fixed effects likelihood (FEL) method, indicating that this residue evolved under purifying selection due to functional importance (Lee et al., 2022). We therefore made T1-Palm (3C) mutants wherein we mutated all three cysteines to serines as well as T1-Palm (2C) mutants where only the core cysteines common to mouse and humans were mutated to serines, as in our prior analysis of CD4 WT. Finally, we made mutants containing both the TMD and Palm mutants that we called T1-TP(3C) or T1-TP(2C).

To confirm that truncating CD4 relieved CD4-Lck association, and evaluate if the

mutations impacted any residual CD4-Lck interactions that may occur due to

colocalization in particular membrane domains, we performed sucrose gradient analysis

of detergent lysates of WT, T1, T1-TMD, T1-Palm(2C), T1-Palm(3C), T1-TP(2C) and

T1-TP(3C). We observed significantly reduced CD4-Lck association in detergent

resistant membranes (DRMs, aka membrane rafts) and detergent soluble membranes

(DSMs) for T1 compared to WT (**Figure 3 – figure supplement 2**)(Pike, 2006). The

mutants did not impact residual CD4-Lck association in the DRMs or DSMs.

Furthermore, the total Lck signal associated with CD4 was significantly lower for T1

compared with the WT, and the residual association was not impacted by the mutants

(**Table 3 and Figure 3 – figure supplement 3**). We therefore conclude that, in our

experimental system, any functional differences between the T1-TMD, T1-Palm(2C/3C),

or T1-TP(2C/3C) mutants and the T1 control is a function of the motif we are studying in

the absence of direct CD4-Lck interactions.

Next, we measured IL-2 production by WT, T1, and CD4 mutant cells to evaluate

the impact of the mutations on signaling output in response to MCC. Here again, T1

was slightly lower than WT, while the T1-TMD reduced the IL-2 response magnitude

and sensitivity relative to the T1 control (**Figure 3A**). This was expected given our prior

work showing that the TMD mutant reduced IL-2 production driven by a shorter CD4

truncation mutant (CD4T, mutated at R420)(Parrish et al., 2015). Mutating the GGXXG

or (C/F)CV+C motifs individually or together did not impact pMHCII-induced TCR

endocytosis relative to the T1 control, and activation-induced CD4 endocytosis was

inhibited for all truncated CD4 molecules (**Figure 3 – figure supplement 4A and 4B**).

The T1-Palm(2C), T1-TP(2C), T1-Palm(3C), and T1-TP(3C) all reduced IL-2 production

relative to T1, indicating that they also influence the Lck-independent contribution of

CD4 to pMHCII-specific signaling (**Figure 3B**). Of note, the T1-TP(2C) IL-2 response

magnitude was lower than that of T1-Palm(2C) alone, consistent with what we

previously observed with the full-length versions of these CD4 mutants (**Figure 3B**)(Lee

et al., 2022). Furthermore, the T1-Palm(3C) response magnitude was lower than that of

T1-Palm(2C), suggesting that the additional cysteine contributes to signaling. Together,

these data indicate that the GGXXG and (C/F)CV+C motifs contribute to the signaling

cascade that leads to pMHCII-specific IL-2 production when CD4 does not directly

interact with Lck via the intracellular CQC clasp and IKRLL motifs. These data do not

tell us where in the signaling pathway the contributions are made.

The CD4 GGXXG and (C/F)CV+C motifs are key determinants of proximal

signaling.

To evaluate if the GGXXG and (C/F)CV+C motifs influence early signaling events, we

measured pCD3ζ, pZap70, and pPlcγ1 levels for four independently generated T1-TMD

lines, three independently generated T1-Palm(2C) lines, and three independently

generated T1-TP(2C) lines relative to their matched T1 controls in response to APCs

expressing MCC:I-E^k, as introduced above (**Figure 2 – figure supplement 3**). For each

T1-TMD and T1-TP(2C) mutant line, we observed lower pCD3ζ, pZap70, and pPlcγ1

levels than their paired T1 control. For the T1-Palm(2C) lines, all 3 showed reduced

pCD3ζ levels, but only 2 of 3 showed reduced pZap70, and pPlcγ1 levels (**Figure 4A**

and Figure 4 – Figure supplement 1A and 2A). These data indicate that the GGXXG

and (C/F)CV+C motifs contribute to CD4's ability to enhance proximal signaling in the

absence of CD4-Lck interactions.

We also considered the average fold change of the means of the independently

generated mutant cell lines relative to their paired T1 controls (**Figure 4B and Figure 4**

– Figure supplement 1B and 2B). For this analysis we included the mean of the fold

change of the T1$^{\Delta bind}$ mutant cells from Figure 2C relative to their paired T1 controls

(dotted line) as a visual to consider the severity of the impact of the mutations on

proximal signaling relative to the contributions of T1, which can bind pMHCII, as well as

the level of signaling observed with T1$^{\Delta bind}$ that cannot bind pMHCII. The T1-TMD and

T1-Palm(2C) showed reduced pCD3ζ and pZap70 levels midway between T1 and

T1$^{\Delta bind}$, while the T1-TP(2C) mutant closely approximated the dotted line representing

the signaling observed in the T1$^{\Delta bind}$ cells. Differences in the average pPlcγ1 responses

were less pronounced. These data further suggest a role for the GGXXG and

(C/F)CV+C motifs in contributing to CD4-mediated pMHCII-specific signaling in the

absence of CD4-Lck interactions.

Because we had found that T1$^{\Delta bind}$ cells failed to produce IL-2 in response to high

peptide concentrations in our dose response experiments (**Figure 1A**), whereas the T1-

TP(2C) mutant cells made IL-2 in response to high peptide concentrations (**Figure 3B**),

we found it interesting that the early signaling responses we measure to APCs

expressing high ligand densities of tethered pMHCII were similar between the T1$^{\Delta bind}$

and T1-TP(2C) cells. We therefore evaluated IL-2 production of the T1, T1$^{\Delta bind}$, and T1-

TP(2C) in response to the APCs used in these signaling studies to ask if they followed

the same pattern of responses to high peptide doses in our prior experiments. As

expected, we found that the T1-TP(2C) cells made approximately half as much IL-2 as

the T1 controls, whereas IL-2 production by the T1$^{\Delta bind}$ cells was negligible (**Figure 4 –**

Figure supplement 3). These data indicate that the ability of CD4 to interact with

pMHCII can partially overcome the loss of function of the GGXXG and (C/F)CV+C

motifs at high ligand densities with respect to a downstream signaling output, such as

IL-2 production, even if the early signaling events measured at 2 minutes after pMHCII-

engagement were similar.

Finally, we considered the percent of mutant responders normalized to the

controls. We observed small decreases in T1-TP(2C) cells with pCD3ζ and pZap70

levels relative to the control, while pPlcγ1 trended lower for this mutant. The single

mutants were roughly equivalent to their matched T1 controls. These data indicate that

mutating the GGXXG and (C/F)CV+C motifs together in the absence of the intracellular

domain can reduce the number of cells that initiate pMHCII-specific signaling (**Figure**

4C and Figure 4 – Figure supplement 1C and 2C). We did not observe this

phenotype in our prior study when the intracellular domain was present, suggesting this

is unique to the absence of the intracellular domain.

We also measured early signaling for three independently generated T1-

Palm(3C) and T1-TP(3C) mutant cell lines compared to their paired T1 controls. For all

lines, pCD3ζ, pZap70, and pPlcγ1 levels were significantly lower than the controls,

which was reflected in the normalized fold change (**Figure 5A and 5B and Figure 5 –**

figures supplement 1A, 1B, 2A, 2B). The fold change for the T1-TP(3C) levels relative

to the controls was similar to the T1$^{\Delta bind}$ cells (**Figure 5B and Figure 5 – figures**

supplement 1B and 2B). These data provide further evidence that the GGXXG and

(C/F)CV+C motifs together are key determinants of the Lck-independent contributions

that CD4 makes to amplifying pMHCII-specific proximal signaling.

Finally, the percent responders were only lower for pZap70 for both T1-Palm(3C)

and T1-TP(3C) relative to WT, while pCD3ζ and pPlcγ1 trended lower (**Figure 5 and**

Figure 5 – figures supplement 1C and 2C). These data are consistent with the idea

that, in the absence of the intracellular domain, mutating these motifs influences

initiation of key signaling events. See **Table 3** for a summary of the data.

Discussion

Our computational reconstruction of CD4 evolution provided access to a subset of

results from experiments that Nature performed over ~435 million years in a greater

variety of jawed vertebrates than could be achieved with mouse or human studies (Lee

et al., 2022). It allowed us to identify residues that were selected under purifying

selection, and thus functionally significant, because acquiring a mutation at these

residues affects fitness and results in a failure to propagate the change to future

progeny. Our structure-function analysis of these residues in 58$\alpha^-\beta^-$ cells, which we

continued in this study, has started to provide mechanistic insights into why these

residues are functionally important.

The data presented here are inconsistent with the prevailing model in which CD4

recruits Lck to CD3 ITAMs to initiate signaling. The decrease in pMHCII-specific IL-2

production when the CD4 intracellular domain is truncated cannot be attributed to

reductions in early signaling events such as pCD3ζ. Importantly, the pCD3ζ levels were

~2 fold lower in the absence of CD4-pMHCII interactions (T1$^{\Delta bind}$ mutant) under the

experimental conditions tested here. We take these data as evidence that there is a

threshold for the number of CD3ζ molecules phosphorylated, and/or the duration of

signaling, that is required to drive IL-2 production in our system. We also conclude that

CD4 binding to pMHCII functions to increase the number of TCR-CD3 complexes per

cell that experience CD3ζ phosphorylation to reach that threshold and/or sustain

signaling and that it similarly impacts pZap70 levels as well as pPlcγ1 levels, albeit to a

lesser extent for the latter.

A logical extension of these conclusions is that there is a density of agonist

pMHCII below which CD4 binding to pMHCII is essential for signal initiation. This idea is

supported by findings that CD4 is necessary for signal propagation and amplification to

the calcium mobilization step in response to fewer than 25 agonist pMHCII (Irvine et al.,

2002). Attempting to accurately measure the contributions of CD4 to ITAM

phosphorylation, with or without mutant motifs, at very low densities of pMHCII would be

challenging and is technically beyond the scope of this study. Nevertheless, in this and

our prior study we did not observe the expected decreases in pCD3ζ levels when CD4-

Lck interactions were reduced by mutating the CQC clasp motif, the IKRLL motif, or by

eliminating the intracellular domain. Furthermore, we found in our prior study that IL-2

production increased significantly when we mutated the full intracellular helix, the helix

and CQC clasp together, or the IKRLL motif alone. These direct tests of the TCR

signaling paradigm are inconsistent with predictions of the model. As such, they suggest

the contributions of CD4 to signal initiation and amplification at the early timepoint we

measured are independent of direct CD4-Lck interactions.

Because the combined impact of mutating the GGXXG and (C/F)CV+C motifs on

pCD3ζ levels were roughly equivalent to disrupting CD4-pMHCII engagement, even

when direct CD4-Lck interactions were absent, we can conclude that the GGXXG and

(C/F)CV+C motifs are key determinants that help CD4 drive signaling above the

threshold required for IL-2 production. Our data also provide evidence that mutating the

GGXXG and (C/F)CV+C motifs together can reduce the frequency of cells experiencing

CD3ζ and Zap70 phosphorylation in the absence of the intracellular domain. Given that

palmitoylation is rapidly reversable, these data suggest that the switch-like function of

the palmitoylation motif may influence signal initiation. Interestingly, unlike CD4-pMHCII

interactions, these motifs are not essential for signaling output at high ligand densities

as we observed low levels of IL-2 produced in cells bearing mutants of these motifs with

increasing doses of agonist pMHCII. Reciprocal binding of CD4 and TCR-CD3 to

pMHCII can therefore amplify pMHCII-specific signaling without the contributions of the

GGXXG and (C/F)CV+C motifs, or motifs in the intracellular domain, albeit at lower

levels than if these motifs are intact. This could be due to the contributions of additional

unidentified motifs in the transmembrane domain, or to the increase in TCR-CD3 dwell

time on pMHCII that is mediated by the CD4 extracellular domain (Glassman et al.,

2018). Given that these motifs were essential for IL-2 production at low ligand densities,

a logical extension of these data is that there is an agonist pMHCII density below which

these motifs are essential for the initiation of early signaling events.

The data from our current study indicate that the GGXXG and (C/F)CV+C motifs

do not influence early signaling events by regulating the ability of CD4 to either recruit

Lck to, or sequester it away from, the TCR-CD3 ITAMs as we previously postulated

(Lee et al., 2022). Instead, they work independently of CD4-Lck interactions to enhance

pMHCII-specific signaling. It is tempting to speculate that because these motifs can

regulate CD4 membrane domain localization and, in other proteins, palmitate moieties

can sandwich cholesterol against the flat surface of a GG patch on a transmembrane

domain helix, these motifs may allow CD4 to regulate local concentration of cholesterol

or cholesterol sulfate around TCR-CD3, perhaps by taking such molecules away from

TCR-CD3 to stabilize allosteric changes associated with signaling (Chen et al., 2022;

Fessler, 2016; Gil et al., 2002; Lee et al., 2015; Song et al., 2014; Swamy et al., 2016;

Teese and Langosch, 2015; Wacker et al., 2013; Wang et al., 2016). These CD4 motifs

may also impact the accessibility of ITAMs for Lck phosphorylation by changing the

local membrane environment around TCR-CD3 upon reciprocal engagement of pMHCII

(Aivazian and Stern, 2000; Xu et al., 2008). Finally, it is worth considering that, in our

prior study, we found G402 of the GGXXG motif has covaried over evolutionary time

with L438 of the intracellular helix IKRLL motif that has inhibitory function, suggesting

that the function of these motifs are co-evolving (Lee et al., 2022). In support of this

idea, we reported in that study that the signal enhancing activity of the GGXXG and

(C/F)CV+C motifs together counterbalanced the inhibitory activity of the IKRLL motif

with regards to IL-2 production. It is unclear at this point if this counterbalancing is

simply additive of individual motif functions, or if there is a functional interplay of the

motifs when the intracellular domain of CD4 is present. Delineating the mechanistic

details by which the GGXXG and (C/F)CV+C motifs increase signaling on their own, or

when combined, and how their function integrates with those of other motifs will be the

subject of future investigations.

Altogether, the data in this and our preceding study provide direct evidence that

complement indirect results indicating that pMHCII-specific signaling is not initiated, or

dependent on, CD4 recruitment of Lck to the CD3 ITAMs (Glassman et al., 2018;

Horkova et al., 2023; Killeen and Littman, 1993; Lee et al., 2022; Xu and Littman, 1993).

Recent work on CD8 suggest that this coreceptor also does not recruit Lck to

phosphorylate CD3 ITAMs (Casas et al., 2014; Wei et al., 2020). In total, these findings

suggest that the TCR signaling paradigm needs revising as it pertains to coreceptor-Lck

interactions. For CD4, the existing evidence suggest that reciprocal binding of CD4 and

TCR-CD3 enables free Lck and Fyn to initiate signaling by phosphorylating the ITAMs

and Zap70 (Glassman et al., 2016; Glassman et al., 2018; Horkova et al., 2023; Lee et

al., 2022; Salmond et al., 2009; van Oers et al., 1996). The contribution of the current

study is in showing the GGXXG and (C/F)CV+C motifs work together to enhance these

early signaling events independently of direct CD4-Lck interactions.

It is interesting to consider what the primary purpose of CD4-Lck interactions via

the CQC clasp and IKRLL motifs are if not to recruit Lck to the CD3 ITAMs. Data from

two studies suggest that Lck makes a kinase-independent contribution to pMHCII-

specific responses; it is therefore plausible that the phenotype reported for CD4 when

the CQC clasp is mutated is due to relieving the kinase-independent scaffolding function

of Lck (Horkova et al., 2023; Xu and Littman, 1993). In addition, CD4-Lck interaction via

the CQC clasp motif and the intracellular helix will prevent the helix from interacting with

other partners, thus mutating the CQC clasp would favor helix interactions with other

putative partners (Lee et al., 2022). Finally, the CQC clasp is thought to play a role in

positioning CD4 in proximity to Lat, either through direct or indirect interactions, which

could contribute to the phenotype of CQC clasp mutants (Bosselut et al., 1999; Lo et al.,

2018; Lo et al., 2019). If CD4-Lck and CD4-Lat interactions are mutually exclusive, and

both require the CQC clasp, then mutating this motif would have a greater impact than

just relieving CD4-Lck interactions (Bosselut et al., 1999). These questions remain

outstanding, as do questions concerning how the intracellular helix regulates pMHCII-

specific signaling.

The broader implications of the data presented in our current and prior evolution-

structure-function studies provide fertile ground for future directions beyond those

mentioned above. Our approach allows us to identify functionally important

residues/motif and then interrogate basic principles of their function that should hold

true across CD4$^+$ T cell subsets since the biochemical and biophysical bases for

interactions between CD4 and its interacting partners (i.e. other proteins or membrane

components) should be the same in 58$\alpha^-\beta^-$ cells as in thymocytes or in different CD4$^+$ T

cells. However, different CD4$^+$ T cell populations will have different levels of expression

of relevant proteins (e.g. interacting partners or modifying enzymes), and may even

have differences in lipid composition, such that the impact of the motifs we have

identified and studied to date may lead to different outcomes in different CD4$^+$ T cell

subsets (http://immpres.co.uk/)(Tuosto and Xu, 2018). For example, differences in

expression of enzymes with switch-like functions, such as those that add or remove

palmitate to the (C/F)CV+C motif, or those that phosphorylate or dephosphorylate the

serines in the intracellular helix that regulate CD4-Lck interactions and inhibitory activity,

may vary between naïve CD4$^+$ T cells, different Th subsets, or Tregs to differentially

tune the activity of these motifs and their impact on pMHCII-specific signaling. Now that

we have identified residues and broader motifs that have proven to be functionally

important *in vivo* over 435 million years of evolution, while also providing mechanistic

insights as to why the residues are important, there is utility in evaluating the impact of

mutant CD4 conditional knock-in mice on thymocyte development, naïve CD4$^+$ T cell

activation and differentiation, and the execution of Th and Treg effector functions. In so

doing, we will gain additional insights into why there is a fitness cost if a mutation is

acquired in these residues. Such knowledge will increase our fundamental

understanding of CD4$^+$ T cell biology and will also be critical for biomimetic engineering

of synthetic receptors that can redirect CD4$^+$ T cell activity for therapeutic purposes.

Materials and methods

The work performed in this manuscript was conducted according to the materials and

methods of our previous study, which can be found here:

https://elifesciences.org/articles/79508#s4.

Funding

This work was supported by R01AI101053 (MSK), CCSG-CA 023074 (MSK), and AZ

TRIF funds (KVD).

References

Aivazian, D., and Stern, L.J. (2000). Phosphorylation of T cell receptor zeta is regulated by a lipid dependent folding transition. Nat Struct Biol *7*, 1023-1026.

Bosselut, R., Zhang, W., Ashe, J.M., Kopacz, J.L., Samelson, L.E., and Singer, A. (1999). Association of the adaptor molecule LAT with CD4 and CD8 coreceptors identifies a new coreceptor function in T cell receptor signal transduction. J Exp Med *190*, 1517-1526.

Casas, J., Brzostek, J., Zarnitsyna, V.I., Hong, J.S., Wei, Q., Hoerter, J.A., Fu, G., Ampudia, J., Zamoyska, R., Zhu, C., and Gascoigne, N.R. (2014). Ligand-engaged TCR is triggered by Lck not associated with CD8 coreceptor. Nat Commun *5*, 5624.

Chen, Y., Zhu, Y., Li, X., Gao, W., Zhen, Z., Dong, Huang, B., Ma, Z., Zhang, A., Song, X.*, et al.* (2022). Cholesterol inhibits TCR signaling by directly restricting TCR-CD3 core tunnel motility. Mol Cell.

Fessler, M.B. (2016). The Intracellular Cholesterol Landscape: Dynamic Integrator of the Immune Response. Trends Immunol *37*, 819-830.

Gil, D., Schamel, W.W., Montoya, M., Sanchez-Madrid, F., and Alarcon, B. (2002). Recruitment of Nck by CD3 epsilon reveals a ligand-induced conformational change essential for T cell receptor signaling and synapse formation. Cell *109*, 901-912.

Glaichenhaus, N., Shastri, N., Littman, D.R., and Turner, J.M. (1991). Requirement for association of p56lck with CD4 in antigen-specific signal transduction in T cells. Cell *64*, 511-520.

Glassman, C.R., Parrish, H.L., Deshpande, N.R., and Kuhns, M.S. (2016). The CD4 and CD3deltaepsilon Cytosolic Juxtamembrane Regions Are Proximal within a Compact TCR-CD3-pMHC-CD4 Macrocomplex. J Immunol *196*, 4713-4722.

Glassman, C.R., Parrish, H.L., Lee, M.S., and Kuhns, M.S. (2018). Reciprocal TCR-CD3 and CD4 Engagement of a Nucleating pMHCII Stabilizes a Functional Receptor Macrocomplex. Cell Rep *22*, 1263-1275.

Harris, D.T., and Kranz, D.M. (2016). Adoptive T Cell Therapies: A Comparison of T Cell Receptors and Chimeric Antigen Receptors. Trends Pharmacol Sci *37*, 220-230.

Horkova, V., Drobek, A., Paprckova, D., Niederlova, V., Prasai, A., Uleri, V., Glatzova, D., Kraller, M., Cesnekova, M., Janusova, S.*, et al.* (2023). Unique roles of co-receptor-bound LCK in helper and cytotoxic T cells. Nat Immunol *24*, 174-185.

Huse, M., Eck, M.J., and Harrison, S.C. (1998). A Zn2+ ion links the cytoplasmic tail of CD4 and the N-terminal region of Lck. J Biol Chem *273*, 18729-18733.

Irvine, D.J., Purbhoo, M.A., Krogsgaard, M., and Davis, M.M. (2002). Direct observation
of ligand recognition by T cells. Nature *419*, 845-849.

Killeen, N., and Littman, D.R. (1993). Helper T-cell development in the absence of CD4-
p56lck association. Nature *364*, 729-732.

Kim, P.W., Sun, Z.Y., Blacklow, S.C., Wagner, G., and Eck, M.J. (2003). A zinc clasp
structure tethers Lck to T cell coreceptors CD4 and CD8. Science *301*, 1725-1728.

Kobayashi, S., Thelin, M.A., Parrish, H.L., Deshpande, N.R., Lee, M.S., Karimzadeh, A.,
Niewczas, M.A., Serwold, T., and Kuhns, M.S. (2020). A biomimetic five-module
chimeric antigen receptor ((5M)CAR) designed to target and eliminate antigen-specific T
cells. Proc Natl Acad Sci U S A *117*, 28950-28959.

Kuhns, M.S., and Badgandi, H.B. (2012). Piecing together the family portrait of TCR-
CD3 complexes. Immunol Rev *250*, 120-143.

Kuhns, M.S., and Davis, M.M. (2012). TCR Signaling Emerges from the Sum of Many
Parts. Frontiers in immunology *3*, 159.

Lee, M.S., Glassman, C.R., Deshpande, N.R., Badgandi, H.B., Parrish, H.L.,
Uttamapinant, C., Stawski, P.S., Ting, A.Y., and Kuhns, M.S. (2015). A Mechanical
Switch Couples T Cell Receptor Triggering to the Cytoplasmic Juxtamembrane Regions
of CD3zetazeta. Immunity *43*, 227-239.

Lee, M.S., Tuohy, P.J., Kim, C.Y., Lichauco, K., Parrish, H.L., Van Doorslaer, K., and
Kuhns, M.S. (2022). Enhancing and inhibitory motifs regulate CD4 activity. eLife *11*,
e79508.

Lo, W.L., Shah, N.H., Ahsan, N., Horkova, V., Stepanek, O., Salomon, A.R., Kuriyan, J.,
and Weiss, A. (2018). Lck promotes Zap70-dependent LAT phosphorylation by bridging
Zap70 to LAT. Nat Immunol *19*, 733-741.

Lo, W.L., Shah, N.H., Rubin, S.A., Zhang, W., Horkova, V., Fallahee, I.R., Stepanek, O.,
Zon, L.I., Kuriyan, J., and Weiss, A. (2019). Slow phosphorylation of a tyrosine residue
in LAT optimizes T cell ligand discrimination. Nat Immunol *20*, 1481-1493.

Melenhorst, J.J., Chen, G.M., Wang, M., Porter, D.L., Chen, C., Collins, M.A., Gao, P.,
Bandyopadhyay, S., Sun, H., Zhao, Z.*, et al.* (2022). Decade-long leukaemia remissions
with persistence of CD4(+) CAR T cells. Nature *602*, 503-509.

Parrish, H.L., Glassman, C.R., Keenen, M.M., Deshpande, N.R., Bronnimann, M.P., and
Kuhns, M.S. (2015). A Transmembrane Domain GGxxG Motif in CD4 Contributes to Its
Lck-Independent Function but Does Not Mediate CD4 Dimerization. PLoS One *10*,
e0132333.

Pike, L.J. (2006). Rafts defined: a report on the Keystone Symposium on Lipid Rafts
and Cell Function. J Lipid Res *47*, 1597-1598.

Rudd, C.E. (2021). How the Discovery of the CD4/CD8-p56(lck) Complexes Changed
Immunology and Immunotherapy. Front Cell Dev Biol *9*, 626095.

Salmond, R.J., Filby, A., Qureshi, I., Caserta, S., and Zamoyska, R. (2009). T-cell
receptor proximal signaling via the Src-family kinases, Lck and Fyn, influences T-cell
activation, differentiation, and tolerance. Immunol Rev *228*, 9-22.

Song, Y., Kenworthy, A.K., and Sanders, C.R. (2014). Cholesterol as a co-solvent and a
ligand for membrane proteins. Protein Sci *23*, 1-22.

Swamy, M., Beck-Garcia, K., Beck-Garcia, E., Hartl, F.A., Morath, A., Yousefi, O.S.,
Dopfer, E.P., Molnar, E., Schulze, A.K., Blanco, R.*, et al.* (2016). A Cholesterol-Based
Allostery Model of T Cell Receptor Phosphorylation. Immunity *44*, 1091-1101.

Teese, M.G., and Langosch, D. (2015). Role of GxxxG Motifs in Transmembrane
Domain Interactions. Biochemistry *54*, 5125-5135.

Tuosto, L., and Xu, C. (2018). Editorial: Membrane Lipids in T Cell Functions. Frontiers
in immunology *9*, 1608.

van Oers, N.S., Lowin-Kropf, B., Finlay, D., Connolly, K., and Weiss, A. (1996). alpha
beta T cell development is abolished in mice lacking both Lck and Fyn protein tyrosine
kinases. Immunity *5*, 429-436.

Wacker, D., Wang, C., Katritch, V., Han, G.W., Huang, X.P., Vardy, E., McCorvy, J.D.,
Jiang, Y., Chu, M., Siu, F.Y.*, et al.* (2013). Structural features for functional selectivity at
serotonin receptors. Science *340*, 615-619.

Wang, F., Beck-Garcia, K., Zorzin, C., Schamel, W.W., and Davis, M.M. (2016).
Inhibition of T cell receptor signaling by cholesterol sulfate, a naturally occurring
derivative of membrane cholesterol. Nat Immunol *17*, 844-850.

Wei, Q., Brzostek, J., Sankaran, S., Casas, J., Hew, L.S., Yap, J., Zhao, X., Wojciech,
L., and Gascoigne, N.R.J. (2020). Lck bound to coreceptor is less active than free Lck.
Proc Natl Acad Sci U S A *117*, 15809-15817.

Wu, L., Wei, Q., Brzostek, J., and Gascoigne, N.R.J. (2020). Signaling from T cell
receptors (TCRs) and chimeric antigen receptors (CARs) on T cells. Cell Mol Immunol
17, 600-612.

Xu, C., Gagnon, E., Call, M.E., Schnell, J.R., Schwieters, C.D., Carman, C.V., Chou,
J.J., and Wucherpfennig, K.W. (2008). Regulation of T cell receptor activation by
dynamic membrane binding of the CD3epsilon cytoplasmic tyrosine-based motif. Cell
135, 702-713.

Xu, H., and Littman, D.R. (1993). A kinase-independent function of Lck in potentiating
antigen-specific T cell activation. Cell *74*, 633-643.
570

Figure Legends

Figure 1. Computational reconstruction of CD4 evolution.
The maximum likelihood phylogenetic tree clusters mammalian CD4 sequences. The tree highlights ancestral reconstructions of CD4 sequences from marsupials (kangaroo silhouette), Atlantogenata (elephant), and Boreoeutheria (wildcat). The logoplot of extant eutherian (Atlantogenata and Boreoeutheria) CD4 sequences show sequence conservation over evolutionary time. In these plots, the height of symbols indicates the relative frequency of each amino acid at that specific position. The mouse CD4 numbering (uniprot) is used as a reference, and residues are color-coded based on sidechain polarity. Evolutionary insertion or deletion events are indicated by dashes (-). Most recent common ancestor (MRCA) sequences are shown at each node in the tree (Node 1-4). As in our previous study (Lee et al., 2022), the ratio of synonymous (dS) and nonsynonymous (dN) substitution rates was calculated. Black dots indicate dN/dS ratios that are significantly below 1 across the entire dataset. Red dots indicate residues under purifying selection in the mammalian only dataset. Previously identified motifs are indicated by boxes, while the intracellular domain helix is shaded gray. The arrow at position 422 indicates where CD4 was truncated (CD4-T1), while TMD and T1-Palm show the mutations studied in this study.

Figure 2. Truncating CD4 does not reduce CD3ζ phosphorylation.
(**A**) Representative IL-2 production is shown in response to a titration of MCC peptide from one experiment (left). Experiments were performed in triplicate and each symbol equals the mean +/- SEM at that peptide concentration. AUC analysis for the dose response is shown as a measure of the response magnitude for the average of three independent experiments performed with one independently generated set of cell lines (center). The average response to a low dose (41nM) of peptide is shown as a measure of sensitivity for three independent experiments performed with one independently generated set of cell lines (right). The data are representative of those obtained with 4 (WT vs T1) or 2 (WT vs T1$^{\Delta bind}$) independently generated sets of cell lines.
(**B**) Phosphorylation intensity of CD3ζ (pCD3ζ) for WT and CD4-T1 (T1) (left), normalized pCD3ζ intensity for WT and T1 (center), and normalized % responders of pCD3ζ for WT and T1 (right). Five independently generated cell lines (biological replicates) were tested for WT and T1. For phosphorylation intensity, each pair of lines (connected symbols) was tested in three independent experiments. Data were analyzed and collected as in Figure 2 – figure supplement 3. One-way ANOVA was performed with a Dunnett's posttest using GraphPad Prism9. For normalized intensity and % responders, all individual mutant cell line values were normalized to their paired control values. Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific comparisons of normalized values using GraphPad Prism9.
(**C**) Phosphorylation intensity of pCD3ζ for T1 and T1$^{\Delta bind}$ (left), normalized pCD3ζ intensity for T1 and T1$^{\Delta bind}$ (center), and normalized % responders of pCD3ζ for T1 and T1$^{\Delta bind}$ (right) were performed and analyzed as in B, with the exception that the open symbols represent data from a single experiment whereas the closed symbols represent aggregate data from three independent experiments. Two-tailed t tests were performed

616   to compare the single T1 vs T1$^{\Delta bind}$ samples as no other samples were collected in
617   parallel.
618

619   **Figure 2- figure supplement 1.**
620   Flow cytometry analysis of CD4 (left) and TCR (right) expression on 58$\alpha^-\beta^-$ hybridoma
621   cells. Parental 58$\alpha^-\beta^-$ hybridoma cells served as negative control for surface expression
622   (open black histogram trace).
623

Figure 2- figure supplement 2.
TCR (left) and CD4 (right) endocytosis after pMHCII engagement is shown for the
indicated cell lines after 16 hours coculture with APCs in the presence of 10μM MCC
peptide. The change in TCR and CD4 gMFI, as measured by flow cytometry, is shown
for each cell line relative to an equivalent sample cultured with APCs in the absence of
MCC peptide. Each data point represents the mean ± SEM for three independent
experiments. For endocytosis measurements were performed in triplicate for each
experiment. One-way ANOVA was performed with a Dunnett's posttest.
632

Figure 2-figure supplement 3.
Example of intracellular signaling analysis workflow.
(**A**) Flow cytometry analysis of WT 58$\alpha^-\beta^-$ hybridoma:M12 cell couples. Representative
dot plots are shown for TCRβ^{GFP+} CD4$^+$ 58$\alpha^-\beta^-$ hybridoma cells coupled to Tag-it Violet-
labeled M12s expressing the indicated tethered pMCHII Hb:I-E^k (left) and MCC:I-E^k
(center). Representative histograms of WT 58$\alpha^-\beta^-$ hybridoma cells coupled to M12 cells
transduced to express the indicated tethered pMHCII are shown for pCD3ζ intensity
(right). 10,000 coupled cells were collected per individual experiment.
(**B**) Flow cytometry analysis of T1 58$\alpha^-\beta^-$ hybridoma-M12 cell couples. Representative
dot plots are shown for TCRβ^{GFP+} CD4$^+$ 58$\alpha^-\beta^-$ hybridoma cells coupled to Tag-it Violet-
labeled Hb:I-E^{k+} (left) and MCC:I-E^{k+} (center) M12 cells. Representative histograms of
T1 58$\alpha^-\beta^-$ hybridoma cells coupled to M12 cells expressing the indicated tethered
pMHCII are shown for pCD3ζ intensity (right). 10,000 coupled cells were collected per
individual experiment.
(**C**) A representative smoothed overlapping histogram of pCD3ζ intensity is shown for
58$\alpha^-\beta^-$ cells coupled to Hb:I-E^{k+} (cyan) or MCC:I-E^{k+} (black) M12 cells. Histogram of
pCD3ζ intensity for 58$\alpha^-\beta^-$ cells coupled to MCC:I-E^{k+} M12 cells subtracted from Hb:I-
E^{k+} M12 cell couples show the difference in pCD3ζ intensity on a bin-by-bin basis after
stimulation with agonist MCC:I-E^k compared with null Hb:I-E^k for WT (left) and T1
(center) cells. Overlapping pCD3ζ histogram (right) of cells responding to MCC:I-E^k
after Hb:I-E^k subtraction shows the responding populations for the WT and T1 cell lines.
Data represent the aggregate from three individual experiments (30,000 couples
analyzed). One-way ANOVA was performed with a Dunnett's posttest for comparison
with the WT sample because other mutants were simultaneously analyzed in this
experiment (not shown).
(**D**) Concatenated pCD3ζ average intensity ± SEM of WT and T1 cells (left) and the
percent of responding WT and T1 cells (right).
660

Figure 2-figure supplement 4.
(**A**) Zap70 (left), and Plcγ1 (right) phosphorylation intensity for WT and T1 are shown for paired (connecting line) WT and T1 cell lines. Five independently generated cell lines were tested (biological replicates). For phosphorylation intensity, each pair of lines (connected symbols) was tested in three independent experiments (technical replicates). The data from those experiments was aggregated, and the symbols represent the mean intensity of the aggregated pCD3ζ intensity values. One-way ANOVA was performed with a Dunnett's posttest.
(**B**) Average T1 phosphorylation for Zap70 (left) and Plcγ1 (right) from A are shown normalized to the paired WT controls. Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific comparisons.
(**C**) Average T1 % responders for pZap70 (left), and pPlcγ1 (right) are shown normalized to the paired WT controls for the cell lines shown in A. Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific comparisons.

Figure 2-figure supplement 5.
(**A**) Phosphorylation intensity for T1 and T1$^{\Delta bind}$ Zap70 (left), and Plcγ1 (right) are shown for paired (connecting line) T1 and T1$^{\Delta bind}$ cell lines. Three independently generated cell lines were tested (biological replicates). Analysis was performed as in Figure 2— figure supplement 4 with the exception that the open symbol represent data from a single experiment whereas the closed symbols represent the average of aggregated data from three independent experiments. For the open symbol comparisons we performed a two-tailed t test as no other samples were collected in parallel.
(**B**) Average T1$^{\Delta bind}$ cell line phosphorylation intensity for Zap70 (left), and Plcγ1 (right) are shown normalized to the average intensity of the paired T1 control cells shown in A. Analysis was performed as in Figure 2— figure supplement 4.
(**C**) Average T1$^{\Delta bind}$ cell line % responders for pZap70 (left) and pPlcγ1 (right) are shown normalized to their paired T1 control. Analysis was performed as in Figure 2— figure supplement 4.

Figure 3. The GGXXG and CV+C motifs are key determinants of CD4 function.
(**A, B**) Representative IL-2 production is shown in response to a titration of MCC peptide from one experiment (left). Experiments were performed in triplicate and each symbol equals the mean +/- SEM at that peptide concentration. AUC analysis for the dose response is shown as a measure of the response magnitude for the average of three independent experiments performed with one independently generated set of cell lines (center). The average response to a low dose (41nM) of peptide is shown as a measure of sensitivity for three independent experiments performed with one independently generated set of cell lines (right). Results are representative of those obtained with at least three independently generated sets of cell lines. One-way ANOVA was performed with a Dunnett's posttest for comparisons with WT and T1 samples, and a Sidak's posttest for comparisons between selected samples.

Figure 3- figure supplement 1.

706 (**A, B**) Flow cytometry analysis of CD4 (left) and TCR (right) expression on 58$\alpha^-\beta^-$
707 hybridoma cells. Parental 58$\alpha^-\beta^-$ hybridoma cells served as negative control for surface
708 expression (open black histogram trace).
709
Figure 3- figure supplement 2.
(**A, B**) Lck signal is shown for each sucrose fraction normalized to the CD4 signal
detected in the corresponding fraction (left). The AUC is shown for the normalized Lck
signal in the DRM (center) and DSM (right) fractions. Each data point represents the
mean +/- SEM for three independent experiments with the same cell line. Data are
representative of experiments performed with three independently generated sets of
lines for TMD, Palm(3C), and TP(3C) mutants. Analysis was performed with two set of
lines for the Palm(2C) and TP(2C) mutants. One-way ANOVA was performed with a
Dunnett's posttest for comparisons with WT and T1 samples, and a Sidak's posttest for
comparisons between selected samples.
720
Figure 3- figure supplement 3.
(**A, B**) Total Lck signal (total AUC for sucrose gradient) normalized to CD4 signal is
shown for the indicated cell lines. Each data point represents the mean +/- SEM for
three independent experiments with the same cell line (experimental replicates). Data
are representative of experiments performed with three independently generated sets of
lines for TMD, Palm(3C), and TP(3C) mutants (biological replicates). Analysis was
performed with two set of lines for the Palm(2C) and TP(2C) mutants. One-way ANOVA
was performed with a Dunnett's posttest for comparisons with WT and T1 samples, and
a Sidak's posttest for comparisons between selected samples.
730
Figure 3- figure supplement 4.
(**A, B**) TCR (left) and CD4 (right) endocytosis after pMHCII engagement is shown for
the indicated cell lines after 16 hours coculture with APCs in the presence of 10μM
MCC peptide. The change in TCR and CD4 gMFI, as measured by flow cytometry, is
shown for each cell line relative to an equivalent sample cultured with APCs in the
absence of MCC peptide. Each data point represents the mean ± SEM for three
independent experiments (experimental replicates). Data are representative of those
acquired with at least two independently generated sets of cell lines (biological
replicates). Endocytosis measurements were performed in triplicate (technical
replicates) for each experiment. One-way ANOVA was performed with a Dunnett's
posttest for comparisons with WT and T1 samples.
742
Figure 4. The GGXXG and CV+C motifs reduce pCD3ζ levels.
(**A**) Phosphorylation intensity of CD3ζ for T1 and T1-TMD (left), T1 and T1-Palm (2C)
(center), and T1 and T1-TP (2C) (right) are shown for independently generated pairs of
(connecting line) T1 and T1-TMD (left), T1 and T1-Palm (2C) (center), and T1 and T1-
TP (2C) (right) cell lines. Each pair of lines (connected closed symbols) was tested in
three independent experiments (technical replicates). The data from those experiments
was aggregated, and the symbols represent the mean intensity of the aggregated
pCD3ζ intensity values. One-way ANOVA was performed with a Dunnett's posttest
using GraphPad Prims 9.

(**B**) Data for each cell line in **A** are shown as the average pCD3ζ intensity for all T1-
TMD (left), T1-Palm (2C) (center), and T1-TP (2C) (right) cell lines normalized to their
paired T1 control. Dotted line is the normalized pCD3ζ intensity for T1$^{\Delta bind}$ as a visual
reference point for the contributions of CD4-pMHCII interactions. Bars represent the
mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific
comparisons using GraphPad Prims 9.
(**C**) The average % responders for phosphorylation of CD3ζ is shown for T1-TMD (left),
T1-Palm (2C) (center), and T1-TP (2C) (right) normalized to their paired T1 control line.
Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's
posttest for specific comparisons using GraphPad Prims 9.
762
Figure 4- figure supplement 1.
(**A**) Phosphorylation intensity of Zap70 for T1 and T1-TMD (left), T1 and T1-Palm (2C)
(center), and T1 and T1-TP (2C) (right) are shown for paired (connecting line) cell lines.
Analysis was performed as in Figure 4A.
(**B**) Normalized phosphorylation intensity of Zap70 for T1-TMD (left), T1-Palm (2C)
(center), and T1-TP (2C) (right) are shown as bars. Analysis was performed as in Figure
4B.
(**C**) Normalized % responders of the phosphorylation Zap70 for T1-TMD (left), T1-Palm
(2C) (center), and T1-TP (2C) (right) are shown as bars for T1 and T1$^{\Delta bind}$ cell lines.
Analysis was performed as in Figure 4C.
773
Figure 4- figure supplement 2.
(**A**) Phosphorylation intensity of Plcγ1 for T1 and T1-TMD (left), T1 and T1-Palm (2C)
(center), and T1 and T1-TP (2C) (right) are shown for paired (connecting line) cell lines.
Analysis was performed as in Figure 4A.
(**B**) Normalized phosphorylation intensity of Plcγ1 for T1-TMD (left), T1-Palm (2C)
(center), and T1-TP (2C) (right) are shown as bars. Analysis was performed as in Figure
4B.
(**C**) Normalized % responders of phosphorylated pPlcγ1 for T1-TMD (left), T1-Palm (2C)
(center), and T1-TP (2C) (right) are shown as bars for T1 and T1$^{\Delta bind}$ cell lines. Analysis
was performed as in Figure 4C.
784
Figure 4- figure supplement 3. Representative IL-2 production is shown in response to
M12 cells transduced to express tethered MCC:I-E^k constructs as used in
phosphorylation analysis. Experiments were performed in triplicate and each bar equals
the mean +/- SEM at that peptide concentration. Results are representative of those
obtained with two independently generated matched sets of cell lines. One-way ANOVA
was performed with a Dunnett's posttest for comparisons with WT and T1 samples, and
a Sidak's posttest for comparisons between selected samples.
792
Figure 5. The (C/F)CVRC motifs reduces proximal TCR-CD3 signaling.
(**A**) Phosphorylation intensity of CD3ζ for T1 and T1-Palm (3C) (left) and T1 and T1-TP
(3C) (right) are shown for independently generated pairs of (connecting lines) T1 and
T1-Palm (3C) (left) and T1 and T1-TP (3C) (right) cell lines. Analysis was performed as
in Figure 4A.

(**B**) Data for each mutant cell line in **A** are shown as the average pCD3ζ intensity values
for T1-Palm (3C) (left) and T1-TP (3C) (right) normalized to their paired T1 controls. The
dotted line represents the normalized phosphorylation CD3ζ intensity for T1$^{\Delta bind}$ as a
visual reference for the contributions of CD4-pMHCII interactions. Analysis was
performed as in Figure 4C.
(**C**) Average % responders for phosphorylation of CD3ζ is shown for T1-Palm (3C) (left)
and T1-TP (3C) (right) normalized to their paired T1 control. Analysis was performed as
in Figure 4C.
806
Figure 5- figure supplement 1.
(**A**) Phosphorylation intensity of Zap70 for T1 and T1-Palm (3C) (left) and T1 and T1-TP
(3C) (right) are shown for paired (connecting line) cell lines. Analysis was performed as
in Figure 5A.
(**B**) Normalized phosphorylation intensity of Zap70 for T1-Palm (3C) (left) and T1-TP
(3C) (right) are shown as bars. Analysis was performed as in Figure 5B.
(**C**) Normalized % responders of phosphorylated Zap70 for T1-Palm (3C) (left) and T1-
TP (3C) (right) are shown as bars. Analysis was performed as in Figure 5C.
815
Figure 5- figure supplement 2.
(**A**) Phosphorylation intensity of Plcγ1 for T1 and T1-Palm (3C) (left) and T1 and T1-TP
(3C) (right) are shown for paired (connecting line) cell lines. Analysis was performed as
in Figure 5A.
(**B**) Normalized phosphorylation intensity of Plcγ1 for T1-Palm (3C) (left) and T1-TP (3C)
(right) are shown as bars. Analysis was performed as in Figure 5B.
(**C**) Normalized % responders of phosphorylated Plcγ1 for T1-Palm (3C) (left) and T1-
TP (3C) (right) are shown as bars. Analysis was performed as in Figure 5C.

Figure 2

Figure 2 — figure supplement 1

Figure 2 — figure supplement 3

A

pZap70

pZap70 intensity

p=< 0.001
p=0.037
p=< 0.001
p=< 0.001
p=0.014

WT   T1

pPlcγ1

pPlcγ1 intensity

p=< 0.001
p=< 0.001
p=> 0.999
p=0.414
p=0.909

WT   T1

B

pZap70

Normalized pZap70 intensity

p= 0.320

WT   T1

pPlcγ1

Normalized pPlcγ1 intensity

p= 0.532

WT   T1

C

pZap70

Normalized % responders of pZap70

p= 0.800

WT   T1

pPlcγ1

Normalized % responders of pPlcγ1

p= 0.196

WT   T1

Figure 2 — figure supplement 4

A

pZap70

pPlcγ1

B

pZap70

pPlcγ1

C

pZap70

pPlcγ1

Figure 2 — figure supplement 5

Figure 3

A

% Max

CD4 TCR.jo

9.4.22 CD4 TCR.jo

Layout

Layout

CD4

TCR

Live Cells

WT
T1
T1-TMD

B

% Max

16 PM

8:16 PM

wJo v9.9.6)

CD4

TCR

Live Cells

WT
T1
T1-Palm (2C)
T1-TP (2C)
T1-Palm (3C)
T1-TP (3C)

Figure 3 — figure supplement 1

A

B

A

TCR Endocytosis

CD4 Endocytosis

B

TCR Endocytosis

CD4 Endocytosis

Figure 3 — figure supplement 4

pPlcγ1

A

B

pPlcγ1

C

pPlcγ1

Figure 4 — figure supplement 2

MCC:I-E^k co-culture

A pCD3ζ intensity

B Normalized pCD3ζ intensity

C Normalized % responders of pCD3ζ

pPlcγ1

A

B

C

Table 1. Motifs and mutants analyzed in this study.

Motif location/known function	Mutant Names	Mutated Motif	Residue Mutations
C-terminally truncated CD4	T1	Ends at R422	R422 is the last residue
Extracellular D1 C"-strand (binds pMHCII)	T1$^{\Delta bind}$	G<u>KGVLIR</u>	K68D, V70D, L71S, I72D, R73S
TMD/Protein or cholesterol interactions	T1-TMD	G<u>G</u>xx<u>G</u>	G403V, G406L
Juxtamembrane/Palmitoylation	T1-Palm (2C)	<u>C</u>V+<u>C</u>	C418S, C421S
Juxtamembrane/Palmitoylation	T1-Palm (3C)	<u>CC</u>V+<u>C</u>	C471S,C418S, C421S
TMD+Palm/raft localization	T1-TP (2C)	G<u>G</u>xx<u>G</u>, <u>C</u>V+<u>C</u>	G403V, G406L, C418S, C421S
TMD+Palm/raft localization	T1-TP (3C)	G<u>G</u>xx<u>G</u>, <u>CC</u>V+<u>C</u>	G403V, G406L, C417S, C418S, C421S

Table 2. All values presented as percent of WT control (truncation average/WT average x 100) for normalized values.

Mutant Names	IL-2 magnitude	IL-2 sensitivity	Normalized pCD3ζ intensity	Normalized % Responders pCD3ζ	Normalized pZap70 intensity	Normalized % Responders pZap70	Normalized pPlcγ1 intensity	Normalized % Responders pPlcγ1	Lck DRM AUC	Lck Total AUC	Lck DSM AUC	CD4 expression	CD4 endocytosis
WT	100	100	100	100	100	100	100	100	100	100	100	100	100
T1	61.46	13.52	106.30	95.20	106.10	102.20	104.40	112.60	28.81	19.22	16.08	110.09	14.30

Table 3. All values presented as percent of T1 control (mutant average/T1 average x 100) for normalized values.

Mutant Names	IL-2 magnitude	IL-2 sensitivity	Normalized pCD3ζ intensity	Normalized % Responders pCD3ζ	Normalized pZap70 intensity	Normalized % Responders pZap70	Normalized pPlcγ1 intensity	Normalized % Responders pPlcγ1	Lck DRM AUC	Lck Total AUC	Lck DSM AUC	CD4 expression	CD4 endocytosis
T1	100	100	100	100	100	100	100	100	100	100	100	100	100
T1$^{\Delta bind}$	1.53	0.00	54.86	91.39	71.96	111.9	74.32	106.8	ND	ND	ND	73.93	38.71
TMD	53.54	17.05	76.68	94.23	88.36	98.77	88.24	107.40	62.36	92.97	110.94	131.10	3.63
Palm (2C)	57.99	116.07	76.35	94.09	86.17	93.03	92.09	98.09	274.03	139.40	130.91	80.30	-98.00
TP (2C)	33.11	0.19	62.06	85.61	79.79	75.58	82.57	83.34	121.39	113.03	114.98	96.54	47.50
Palm (3C)	27.84	0.19	71.09	94.76	86.32	76.82	80.37	88.62	137.57	94.34	92.07	87.58	6.99
TP (3C)	31.19	45.53	62.49	84.60	80.16	66.92	70.91	82.81	86.84	101.20	102.74	79.59	-54.69

Figure Legends

Figure 1. Computational reconstruction of CD4 evolution.
The maximum likelihood phylogenetic tree clusters mammalian CD4 sequences. The tree highlights ancestral reconstructions of CD4 sequences from marsupials (kangaroo silhouette), Atlantogenata (elephant), and Boreoeutheria (wildcat). The logoplot of extant eutherian (Atlantogenata and Boreoeutheria) CD4 sequences show sequence conservation over evolutionary time. In these plots, the height of symbols indicates the relative frequency of each amino acid at that specific position. The mouse CD4 numbering (uniprot) is used as a reference, and residues are color-coded based on sidechain polarity. Evolutionary insertion or deletion events are indicated by dashes (-). Most recent common ancestor (MRCA) sequences are shown at each node in the tree (Node 1-4). As in our previous study (Lee et al., 2022), the ratio of synonymous (dS) and nonsynonymous (dN) substitution rates was calculated. Black dots indicate dN/dS ratios that are significantly below 1 across the entire dataset. Red dots indicate residues under purifying selection in the mammalian only dataset. Previously identified motifs are indicated by boxes, while the intracellular domain helix is shaded gray. The arrow at position 422 indicates where CD4 was truncated (CD4-T1), while TMD and T1-Palm show the mutations studied in this study.

Figure 2. Truncating CD4 does not reduce CD3ζ phosphorylation.
(**A**) Representative IL-2 production is shown in response to a titration of MCC peptide from one experiment (left). Experiments were performed in triplicate and each symbol equals the mean +/- SEM at that peptide concentration. AUC analysis for the dose response is shown as a measure of the response magnitude for the average of three independent experiments performed with one independently generated set of cell lines (center). The average response to a low dose (41nM) of peptide is shown as a measure of sensitivity for three independent experiments performed with one independently generated set of cell lines (right). The data are representative of those obtained with 4 (WT vs T1) or 2 (WT vs T1$^{\Delta bind}$) independently generated sets of cell lines.
(**B**) Phosphorylation intensity of CD3ζ (pCD3ζ) for WT and CD4-T1 (T1) (left), normalized pCD3ζ intensity for WT and T1 (center), and normalized % responders of pCD3ζ for WT and T1 (right). Five independently generated cell lines (biological replicates) were tested for WT and T1. For phosphorylation intensity, each pair of lines (connected symbols) was tested in three independent experiments. Data were analyzed and collected as in Figure 2 – figure supplement 3. One-way ANOVA was performed with a Dunnett's posttest using GraphPad Prism9. For normalized intensity and % responders, all individual mutant cell line values were normalized to their paired control values. Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific comparisons of normalized values using GraphPad Prism9.
(**C**) Phosphorylation intensity of pCD3ζ for T1 and T1$^{\Delta bind}$ (left), normalized pCD3ζ intensity for T1 and T1$^{\Delta bind}$ (center), and normalized % responders of pCD3ζ for T1 and T1$^{\Delta bind}$ (right) were performed and analyzed as in B, with the exception that the open symbols represent data from a single experiment whereas the closed symbols represent aggregate data from three independent experiments. Two-tailed t tests were performed

612 to compare the single T1 vs T1$^{\Delta bind}$ samples as no other samples were collected in
613 parallel.
614
615 **Figure 2- figure supplement 1.**
616 Flow cytometry analysis of CD4 (left) and TCR (right) expression on 58$\alpha^-\beta^-$ hybridoma
617 cells. Parental 58$\alpha^-\beta^-$ hybridoma cells served as negative control for surface expression
618 (open black histogram trace).
619
Figure 2- figure supplement 2.
TCR (left) and CD4 (right) endocytosis after pMHCII engagement is shown for the
indicated cell lines after 16 hours coculture with APCs in the presence of 10μM MCC
peptide. The change in TCR and CD4 gMFI, as measured by flow cytometry, is shown
for each cell line relative to an equivalent sample cultured with APCs in the absence of
MCC peptide. Each data point represents the mean ± SEM for three independent
experiments. For endocytosis measurements were performed in triplicate for each
experiment. One-way ANOVA was performed with a Dunnett's posttest.
628
Figure 2-figure supplement 3.
Example of intracellular signaling analysis workflow.
(**A**) Flow cytometry analysis of WT 58$\alpha^-\beta^-$ hybridoma:M12 cell couples. Representative
dot plots are shown for TCRβ^{GFP+} CD4$^+$ 58$\alpha^-\beta^-$ hybridoma cells coupled to Tag-it Violet-
labeled M12s expressing the indicated tethered pMCHII Hb:I-E^k (left) and MCC:I-E^k
(center). Representative histograms of WT 58$\alpha^-\beta^-$ hybridoma cells coupled to M12 cells
transduced to express the indicated tethered pMHCII are shown for pCD3ζ intensity
(right). 10,000 coupled cells were collected per individual experiment.
(**B**) Flow cytometry analysis of T1 58$\alpha^-\beta^-$ hybridoma-M12 cell couples. Representative
dot plots are shown for TCRβ^{GFP+} CD4$^+$ 58$\alpha^-\beta^-$ hybridoma cells coupled to Tag-it Violet-
labeled Hb:I-E^{k+} (left) and MCC:I-E^{k+} (center) M12 cells. Representative histograms of
T1 58$\alpha^-\beta^-$ hybridoma cells coupled to M12 cells expressing the indicated tethered
pMHCII are shown for pCD3ζ intensity (right). 10,000 coupled cells were collected per
individual experiment.
(**C**) A representative smoothed overlapping histogram of pCD3ζ intensity is shown for
58$\alpha^-\beta^-$ cells coupled to Hb:I-E^{k+} (cyan) or MCC:I-E^{k+} (black) M12 cells. Histogram of
pCD3ζ intensity for 58$\alpha^-\beta^-$ cells coupled to MCC:I-E^{k+} M12 cells subtracted from Hb:I-
E^{k+} M12 cell couples show the difference in pCD3ζ intensity on a bin-by-bin basis after
stimulation with agonist MCC:I-E^k compared with null Hb:I-E^k for WT (left) and T1
(center) cells. Overlapping pCD3ζ histogram (right) of cells responding to MCC:I-E^k
after Hb:I-E^k subtraction shows the responding populations for the WT and T1 cell lines.
Data represent the aggregate from three individual experiments (30,000 couples
analyzed). One-way ANOVA was performed with a Dunnett's posttest for comparison
with the WT sample because other mutants were simultaneously analyzed in this
experiment (not shown).
(**D**) Concatenated pCD3ζ average intensity ± SEM of WT and T1 cells (left) and the
percent of responding WT and T1 cells (right).
656

Figure 2-figure supplement 4.
(**A**) Zap70 (left), and Plcγ1 (right) phosphorylation intensity for WT and T1 are shown for paired (connecting line) WT and T1 cell lines. Five independently generated cell lines were tested (biological replicates). For phosphorylation intensity, each pair of lines (connected symbols) was tested in three independent experiments (technical replicates). The data from those experiments was aggregated, and the symbols represent the mean intensity of the aggregated pCD3ζ intensity values. One-way ANOVA was performed with a Dunnett's posttest.
(**B**) Average T1 phosphorylation for Zap70 (left) and Plcγ1 (right) from A are shown normalized to the paired WT controls. Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific comparisons.
(**C**) Average T1 % responders for pZap70 (left), and pPlcγ1 (right) are shown normalized to the paired WT controls for the cell lines shown in A. Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific comparisons.

Figure 2-figure supplement 5.
(**A**) Phosphorylation intensity for T1 and T1$^{\Delta bind}$ Zap70 (left), and Plcγ1 (right) are shown for paired (connecting line) T1 and T1$^{\Delta bind}$ cell lines. Three independently generated cell lines were tested (biological replicates). Analysis was performed as in Figure 2— figure supplement 4 with the exception that the open symbol represent data from a single experiment whereas the closed symbols represent the average of aggregated data from three independent experiments. For the open symbol comparisons we performed a two-tailed t test as no other samples were collected in parallel.
(**B**) Average T1$^{\Delta bind}$ cell line phosphorylation intensity for Zap70 (left), and Plcγ1 (right) are shown normalized to the average intensity of the paired T1 control cells shown in A. Analysis was performed as in Figure 2— figure supplement 4.
(**C**) Average T1$^{\Delta bind}$ cell line % responders for pZap70 (left) and pPlcγ1 (right) are shown normalized to their paired T1 control. Analysis was performed as in Figure 2— figure supplement 4.

Figure 3. The GGXXG and CV+C motifs are key determinants of CD4 function.
(**A, B**) Representative IL-2 production is shown in response to a titration of MCC peptide from one experiment (left). Experiments were performed in triplicate and each symbol equals the mean +/- SEM at that peptide concentration. AUC analysis for the dose response is shown as a measure of the response magnitude for the average of three independent experiments performed with one independently generated set of cell lines (center). The average response to a low dose (41nM) of peptide is shown as a measure of sensitivity for three independent experiments performed with one independently generated set of cell lines (right). Results are representative of those obtained with at least three independently generated sets of cell lines. One-way ANOVA was performed with a Dunnett's posttest for comparisons with WT and T1 samples, and a Sidak's posttest for comparisons between selected samples.

Figure 3- figure supplement 1.

702 (**A, B**) Flow cytometry analysis of CD4 (left) and TCR (right) expression on $58\alpha^-\beta^-$
703 hybridoma cells. Parental $58\alpha^-\beta^-$ hybridoma cells served as negative control for surface
704 expression (open black histogram trace).
705
Figure 3- figure supplement 2.
(**A, B**) Lck signal is shown for each sucrose fraction normalized to the CD4 signal
detected in the corresponding fraction (left). The AUC is shown for the normalized Lck
signal in the DRM (center) and DSM (right) fractions. Each data point represents the
mean +/- SEM for three independent experiments with the same cell line. Data are
representative of experiments performed with three independently generated sets of
lines for TMD, Palm(3C), and TP(3C) mutants. Analysis was performed with two set of
lines for the Palm(2C) and TP(2C) mutants. One-way ANOVA was performed with a
Dunnett's posttest for comparisons with WT and T1 samples, and a Sidak's posttest for
comparisons between selected samples.
716
Figure 3- figure supplement 3.
(**A, B**) Total Lck signal (total AUC for sucrose gradient) normalized to CD4 signal is
shown for the indicated cell lines. Each data point represents the mean +/- SEM for
three independent experiments with the same cell line (experimental replicates). Data
are representative of experiments performed with three independently generated sets of
lines for TMD, Palm(3C), and TP(3C) mutants (biological replicates). Analysis was
performed with two set of lines for the Palm(2C) and TP(2C) mutants. One-way ANOVA
was performed with a Dunnett's posttest for comparisons with WT and T1 samples, and
a Sidak's posttest for comparisons between selected samples.
726
Figure 3- figure supplement 4.
(**A, B**) TCR (left) and CD4 (right) endocytosis after pMHCII engagement is shown for
the indicated cell lines after 16 hours coculture with APCs in the presence of $10\mu M$
MCC peptide. The change in TCR and CD4 gMFI, as measured by flow cytometry, is
shown for each cell line relative to an equivalent sample cultured with APCs in the
absence of MCC peptide. Each data point represents the mean ± SEM for three
independent experiments (experimental replicates). Data are representative of those
acquired with at least two independently generated sets of cell lines (biological
replicates). Endocytosis measurements were performed in triplicate (technical
replicates) for each experiment. One-way ANOVA was performed with a Dunnett's
posttest for comparisons with WT and T1 samples.
738
Figure 4. The GGXXG and CV+C motifs reduce pCD3ζ levels.
(**A**) Phosphorylation intensity of CD3ζ for T1 and T1-TMD (left), T1 and T1-Palm (2C)
(center), and T1 and T1-TP (2C) (right) are shown for independently generated pairs of
(connecting line) T1 and T1-TMD (left), T1 and T1-Palm (2C) (center), and T1 and T1-
TP (2C) (right) cell lines. Each pair of lines (connected closed symbols) was tested in
three independent experiments (technical replicates). The data from those experiments
was aggregated, and the symbols represent the mean intensity of the aggregated
pCD3ζ intensity values. One-way ANOVA was performed with a Dunnett's posttest
using GraphPad Prims 9.

(**B**) Data for each cell line in **A** are shown as the average pCD3ζ intensity for all T1-
TMD (left), T1-Palm (2C) (center), and T1-TP (2C) (right) cell lines normalized to their
paired T1 control. Dotted line is the normalized pCD3ζ intensity for T1$^{\Delta bind}$ as a visual
reference point for the contributions of CD4-pMHCII interactions. Bars represent the
mean +/- SEM. One-way ANOVA was performed with a Sidak's posttest for specific
comparisons using GraphPad Prims 9.
(**C**) The average % responders for phosphorylation of CD3ζ is shown for T1-TMD (left),
T1-Palm (2C) (center), and T1-TP (2C) (right) normalized to their paired T1 control line.
Bars represent the mean +/- SEM. One-way ANOVA was performed with a Sidak's
posttest for specific comparisons using GraphPad Prims 9.
758
Figure 4- figure supplement 1.
(**A**) Phosphorylation intensity of Zap70 for T1 and T1-TMD (left), T1 and T1-Palm (2C)
(center), and T1 and T1-TP (2C) (right) are shown for paired (connecting line) cell lines.
Analysis was performed as in Figure 4A.
(**B**) Normalized phosphorylation intensity of Zap70 for T1-TMD (left), T1-Palm (2C)
(center), and T1-TP (2C) (right) are shown as bars. Analysis was performed as in Figure
4B.
(**C**) Normalized % responders of the phosphorylation Zap70 for T1-TMD (left), T1-Palm
(2C) (center), and T1-TP (2C) (right) are shown as bars for T1 and T1$^{\Delta bind}$ cell lines.
Analysis was performed as in Figure 4C.
769
Figure 4- figure supplement 2.
(**A**) Phosphorylation intensity of Plcγ1 for T1 and T1-TMD (left), T1 and T1-Palm (2C)
(center), and T1 and T1-TP (2C) (right) are shown for paired (connecting line) cell lines.
Analysis was performed as in Figure 4A.
(**B**) Normalized phosphorylation intensity of Plcγ1 for T1-TMD (left), T1-Palm (2C)
(center), and T1-TP (2C) (right) are shown as bars. Analysis was performed as in Figure
4B.
(**C**) Normalized % responders of phosphorylated pPlcγ1 for T1-TMD (left), T1-Palm (2C)
(center), and T1-TP (2C) (right) are shown as bars for T1 and T1$^{\Delta bind}$ cell lines. Analysis
was performed as in Figure 4C.
780
Figure 4- figure supplement 3. Representative IL-2 production is shown in response to
M12 cells transduced to express tethered MCC:I-E^k constructs as used in
phosphorylation analysis. Experiments were performed in triplicate and each bar equals
the mean +/- SEM at that peptide concentration. Results are representative of those
obtained with two independently generated matched sets of cell lines. One-way ANOVA
was performed with a Dunnett's posttest for comparisons with WT and T1 samples, and
a Sidak's posttest for comparisons between selected samples.
788
Figure 5. The (C/F)CVRC motifs reduces proximal TCR-CD3 signaling.
(**A**) Phosphorylation intensity of CD3ζ for T1 and T1-Palm (3C) (left) and T1 and T1-TP
(3C) (right) are shown for independently generated pairs of (connecting lines) T1 and
T1-Palm (3C) (left) and T1 and T1-TP (3C) (right) cell lines. Analysis was performed as
in Figure 4A.

(**B**) Data for each mutant cell line in **A** are shown as the average pCD3ζ intensity values
for T1-Palm (3C) (left) and T1-TP (3C) (right) normalized to their paired T1 controls. The
dotted line represents the normalized phosphorylation CD3ζ intensity for T1$^{\Delta bind}$ as a
visual reference for the contributions of CD4-pMHCII interactions. Analysis was
performed as in Figure 4C.
(**C**) Average % responders for phosphorylation of CD3ζ is shown for T1-Palm (3C) (left)
and T1-TP (3C) (right) normalized to their paired T1 control. Analysis was performed as
in Figure 4C.
802
Figure 5- figure supplement 1.
(**A**) Phosphorylation intensity of Zap70 for T1 and T1-Palm (3C) (left) and T1 and T1-TP
(3C) (right) are shown for paired (connecting line) cell lines. Analysis was performed as
in Figure 5A.
(**B**) Normalized phosphorylation intensity of Zap70 for T1-Palm (3C) (left) and T1-TP
(3C) (right) are shown as bars. Analysis was performed as in Figure 5B.
(**C**) Normalized % responders of phosphorylated Zap70 for T1-Palm (3C) (left) and T1-
TP (3C) (right) are shown as bars. Analysis was performed as in Figure 5C.
811
Figure 5- figure supplement 2.
(**A**) Phosphorylation intensity of Plcγ1 for T1 and T1-Palm (3C) (left) and T1 and T1-TP
(3C) (right) are shown for paired (connecting line) cell lines. Analysis was performed as
in Figure 5A.
(**B**) Normalized phosphorylation intensity of Plcγ1 for T1-Palm (3C) (left) and T1-TP (3C)
(right) are shown as bars. Analysis was performed as in Figure 5B.
(**C**) Normalized % responders of phosphorylated Plcγ1 for T1-Palm (3C) (left) and T1-
TP (3C) (right) are shown as bars. Analysis was performed as in Figure 5C.