Omicron Spike confers enhanced infectivity and interferon resistance to SARS-CoV-2 in human nasal tissue

Guoli Shi[1,*], Tiansheng Li[2,*], Kin Kui Lai[1], Reed F. Johnson[2], Jonathan W Yewdell[2], Alex A Compton[1,#]

[1]HIV Dynamics and Replication Program, Center for Cancer Research, National Cancer Institute, Frederick, MD
[2]Laboratory of Viral Diseases, National Institute of Allergy and Infectious Diseases, Bethesda, MD
[*]These authors contributed equally.
[#]corresponding author (alex.compton@nih.gov)

Summary

Omicron emerged following COVID-19 vaccination campaigns, displaced previous SARS-CoV-2 variants of concern worldwide, and gave rise to lineages that continue to spread. Here, we show that Omicron exhibits increased infectivity in primary adult upper airway tissue relative to Delta. Using recombinant forms of SARS-CoV-2 and nasal epithelial cells cultured at the liquid-air interface, enhanced infectivity maps to the step of cellular entry and evolved recently through mutations unique to Omicron Spike. Unlike earlier variants of SARS-CoV-2, Omicron enters nasal cells independently of serine transmembrane proteases and instead relies upon metalloproteinases to catalyze membrane fusion. This entry pathway unlocked by Omicron Spike enables evasion of constitutive and interferon-induced antiviral factors that restrict SARS-CoV-2 entry following attachment. Therefore, the increased transmissibility exhibited by Omicron in humans may be attributed not only to its evasion of vaccine-elicited adaptive immunity, but also to its superior invasion of nasal epithelia and resistance to the cell-intrinsic barriers present therein.

Introduction

Efficient infection of human cells by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) requires the interaction of SARS-CoV-2 Spike with its receptor at the cell surface, angiotensin-converting enzyme 2 (ACE2)[1,2]. In addition, Spike has been described to bind to various cellular factors to promote coronavirus attachment to the cell surface. 78-kilodalton glucose-regulated protein, neuropilin-1, high-density lipoprotein scavenger receptor B type 1, CD209/CD299, Axl, sialic acid, and heparan sulfate have all been reported to interact with Spike and promote the virus entry process[3-9]. In general, it is thought that factors aiding virus attachment enable subsequent ACE2 binding. Engagement of ACE2 alters Spike conformation and facilitates its processing by cellular proteases, such as serine transmembrane proteases like TMPRSS2, matrix metalloproteinases (MMP), or a disintegrin and metalloproteinases (ADAM)[2,10-14]. Protease cleavage enables Spike to trigger fusion between viral and cellular membranes to complete cellular entry[15].

The Omicron B.1.1.529.1 (BA.1) variant emerged in late 2021 and, as a result of increased transmissibility between humans[16,17], quickly replaced the Delta variant of concern as the dominant form of SARS-CoV-2 circulating worldwide (https://www.cdc.

gov/coronavirus/2019-ncov/variants/variant-classifications.html). Compared to previous SARS-
CoV-2 variants of concern, Spike from BA.1 contained a plethora of unique mutations, and its
descendants (including the BA.2, BA.4, BA.5, BQ.1, and XBB subvariants) contained additional
mutations in Spike (https://covid.cdc.gov/covid-data-tracker/#datatracker-home). The functional
characterization of Omicron has revealed that said Spike mutations, including those found in the
receptor binding domain, allow for evasion of neutralizing antibodies induced by COVID-19
vaccination[18-26]. Furthermore, some studies have shown that Omicron Spike displays increased
affinity for human ACE2 compared to the early D614G variant[23,27,28], but this was not supported
by others[29]. Therefore, it remains unclear whether antibody evasion and ACE2 affinity are the
only factors explaining the improved transmissibility of Omicron subvariants.
It has also been reported that Omicron utilizes a distinct cellular entry pathway compared
to ancestral or early forms of SARS-CoV-2[30-34]. Specifically, the proteolytic activation of
Omicron Spike is less dependent on TMPRSS2 but may be more dependent on cathepsin activity
in endolysosomes or metalloproteinase activity at the plasma membrane[11]. However, the precise
characterization of the entry pathways used by Omicron in physiologically relevant primary cells
is lacking, and whether they are sensitive to cell-intrinsic antiviral barriers has not been
determined. The apparent loss of TMPRSS2 dependence may have consequences for the cellular
tropism of Omicron in vivo. Some reports suggest that Omicron BA.1 exhibits a decreased
propensity for lower respiratory tract (lung) infection, which may explain the decreased
pathogenicity associated with Omicron infections[35-40]. Since the upper respiratory tract, and
specifically the nasal epithelium, represents the initial site of SARS-CoV-2 infection and the
initial protective innate immune response[41,42], we compared the ability of Omicron to invade
human nasal epithelia relative to ancestral SARS-CoV-2 and the preceding variant of concern,
Delta, using a primary human nasal tissue culture model. We found that Omicron BA.1 and
BA.2 exhibited a markedly enhanced infectivity in primary nasal epithelia compared to early
SARS-CoV-2, and using recombinant virus, we show that Spike from Omicron, but not Delta,
controlled this phenotype. Furthermore, despite Omicron eliciting a type-I interferon response in
nasal tissue, Omicron Spike used a metalloproteinase-mediated entry pathway that was
associated with evasion of interferon-inducible factors targeting virus entry. Collectively, these
findings may explain the increased transmissibility of Omicron as well as its ability to displace
more interferon-sensitive variants of SARS-CoV-2.
Overall, our findings suggest that the global spread of Omicron was fueled not only by
increased ACE2 affinity and resistance to vaccine-induced adaptive immunity, but also by
resistance to nasal cell-intrinsic entry inhibitors.
81
82 **Results**
83
84 **Omicron BA.1 exhibits superior replicative fitness relative to early SARS-CoV-2 and Delta**
85 **in primary human nasal epithelial cells**
86
Primary human nasal epithelial cells isolated from three donors were pooled and cultured
as submerged monolayers, which allows for the propagation of cells in the basal
(undifferentiated) state. Immunofluorescence staining of acetylated tubulin indicated that mature
cilia are lacking in cells cultured in this manner, although they express detectable levels of ACE2
(Figure 1A). Nasal monolayers were inoculated with early/ancestral SARS-CoV-2 (USA-
WA1/2020 strain, herein referred to as WA1) at a multiplicity of infection (MOI) of 0.05, and

RT-qPCR of viral ORF1a RNA at multiple time points were suggestive of virus replication over
time (**Figure 1B**). Productive infection was confirmed by demonstrating that cell culture
supernatants contained infectious virus that could be titered on Vero E6-ACE2-TMPRSS2 cells
by ORF1a qPCR and anti-nucleocapsid immunostaining (**Supplemental Figure 1A**). By
comparison, inoculation of nasal monolayers with the same MOI of BA.1 resulted in far greater
virus replication at all time points (**Figure 1B**). In contrast, BA.1 did not replicate in submerged
monolayers of human small airway (lung) epithelial cells (three pooled donors), while WA1
replicated with similar kinetics in nasal and lung cells (**Figure 1C**). These data suggest that BA.1
displays a growth advantage specific to the upper airway of the human respiratory tract. BA.1
exhibited an approximately 100-fold greater replication capacity in nasal monolayers compared
to WA1, either when measuring absolute (**Supplemental Figure 1B**) or relative (**Supplemental
Figure 1C**) copy numbers of viral ORF1a. We found that this replication advantage by BA.1
was accompanied by an approximately 30-fold superior capacity to adhere to the nasal cell
surface compared to WA1 (**Supplemental Figure 1D**). We excluded that WA1 and BA.1
inoculants (initially titered on Vero E6-TMPRSS2 cells) contained different quantities of virus
particles by inoculating HEK293T-ACE2 and Vero E6-TMPRSS2 cells in parallel. In contrast to
our observations in nasal monolayers, WA1 and BA.1 infected these permissive cell lines to
equal extents (**Supplemental Figure 1E and 1F**). Even when virus inoculants were calculated
using an alternative virus titering strategy (based on viral ORF1a RT-qPCR instead of Vero E6-
TMPRSS2 infection), BA.1 maintained a significant replicative advantage in nasal monolayers
relative to WA1 (**Supplemental Figure 1G**). Therefore, BA.1 exhibits enhanced replicative
potential in primary human nasal cells relative to WA1 and this is associated with improved cell
surface adhesion by BA.1 Spike. Since BA.1 emerged in humans while the preceding variant of
concern, Delta, was widespread, we compared the capacity for BA.1 and Delta to infect human
nasal monolayers. In contrast to the orders of magnitude improvement exhibited by BA.1, Delta
infected nasal cells only modestly better than WA1, and the difference was not statistically
significant (**Figure 1D**). Therefore, the enhanced ability for BA.1 to infect primary human nasal
epithelia emerged in the Omicron ancestor, not in Delta. Importantly, remdesivir, an inhibitor of
the viral RNA polymerase, significantly reduced ORF1a levels in nasal cells inoculated with
WA1, Delta, and BA.1, demonstrating that measurement of ORF1a by RT-qPCR is reflective of
virus replication (**Figure 1E**).
During the preparation of this manuscript, it was reported that Omicron displays greater
adhesion to human nasal cells, and this was attributed to enhanced binding to cilia in
differentiated nasal epithelia[43]. Our results here suggest that, besides cilia, other factors intrinsic
to nasal cells must govern the improved replicative fitness of BA.1 compared to early SARS-
CoV-2.
129
130 **Omicron Spike enables enhanced SARS-CoV-2 infectivity in primary nasal epithelia
131 cultured at the air-liquid interface**
132
To recreate the three-dimensional, pseudostratified architecture of nasal epithelia in vivo,
primary human nasal epithelial cells were pooled from 14 donors and differentiated at the air-
liquid interface (ALI). In addition to ample ACE2 expression, differentiation status was
confirmed by stratification of nuclei and the presence of mature cilia at the tissue surface (**Figure
2A**). Relative to nasal monolayers (**Supplemental Figure 1B**), nasal ALI were more permissive
to SARS-CoV-2 infection, as assessed by absolute ORF1a copy number (**Supplemental Figure**

2A). In this tissue, replication of Omicron BA.1 and BA.2 exceeded that of WA1 by multiple
orders of magnitude at 48 hours post-inoculation, with BA.2 exhibiting the greatest replicative
potential **(Figure 2B)**. To confirm that nasal ALI support repeated rounds of productive
infection, we collected culture medium supernatants of nasal ALI inoculated with WA1, BA.1,
or BA.2 and quantified infectious virus levels by titrating on Vero E6-TMPRSS2 cells. In
agreement with RT-qPCR results, BA.1 and BA.2 achieved higher infectious titers in nasal ALI
relative to WA1 **(Figure 2C)**. In accordance with enhanced virus replication of BA.1 and BA.2,
we detected elevated type-I interferon induction (*IFNB*) at 48 hours post-inoculation **(Figure
2D)**.
Given the degree to which Omicron BA.1 and BA.2 replicate in primary nasal ALI, we
also tested the infective potential of an Omicron sublineage more recently circulating in humans
(BQ.1) as well as Delta and the D614G variant, the ancestor of Pango lineage B that gave rise to
all SARS-CoV-2 variants of concern. D614G replicated to a similar extent as WA1, while Delta
achieved a significantly higher degree of replication (23-fold). However, replication of Omicron
lineages BA.1 and BQ.1 in nasal ALI far exceeded that of all variants tested (282- and 88-fold,
respectively) **(Figure 2E)**. While all viruses triggered the expression of the interferon-stimulated
gene *IFITM3* (**Supplemental Figure 2B**), BA.1 and BQ.1 induced more *IFNB* expression
compared to the other variants (**Figure 2F**). We also inoculated nasal ALI with Omicron XBB,
the ancestor of sublineages currently circulating in humans as of October 2023, and it replicated
to a significantly greater extent than D614G (**Supplemental Figure 2C**).
To determine whether the enhanced nasal tropism of Omicron BA.1 can be attributed to
its Spike protein, we generated mCherry-expressing, recombinant SARS-CoV-2 WA1 encoding
Spike from WA1, Delta, or BA.1 (referred to as WA1 (WA.1 Spike), WA1 (Delta Spike), or
WA1 (BA.1 Spike). We found that the presence of Spike from BA.1, but not Delta, resulted in a
significant gain in nasal cell infectivity relative to WA1 (20-fold by mCherry fluorescence
(**Figure 2G**) or 100-fold by viral ORF1a RT-qPCR at 24 hours post-inoculation) **(Supplemental
Figure 2D)**. Results were similar regardless of how we calculated virus titers to prepare
inoculants (infection of Vero E6-TMPRSS2 cells (**Supplemental Figure 2D**) or viral ORF1a
RT-qPCR (**Supplemental Figure 2E**). When earlier timepoints were examined, WA1 (BA.1
Spike)-infected cells were observed as early as 6 hours post-inoculation and numbers increased
substantially between 9 and 12 hours post-inoculation, indicative of rapid virus spread in nasal
ALI (**Supplemental Figure 2F**). These results formally indicate that the elevated nasal tissue
infectivity of BA.1 is governed by its Spike protein, and more specifically, by the mutations that
distinguish it from Delta Spike. Therefore, enhanced nasal cell tropism evolved recently in the
Omicron ancestor as a result of unique mutations in Spike.
174
175 **Entry of Omicron BA.1 into primary nasal epithelial cells is mediated by proteases of the
176 MMP/ADAM families and occurs independently of endocytosis**
177
Since BA.1 Spike encodes for an unprecedented ability to enter human nasal epithelia,
we characterized the subcellular location at which Omicron Spike-mediated entry occurs in nasal
epithelia and determined the cellular proteases that enable it. Several reports claim that Omicron
enters cells via cathepsin-dependent fusion in endosomes[31,32,36], but the Omicron entry pathway
may vary in transformed cell lines[44]. Therefore, we examined the entry determinants of WA1,
Delta, and BA.1 in primary human nasal ALI. We used E64d to disrupt cathepsin-mediated entry
in endolysosomes and found that, while WA1 replication was reduced by 25%, that of Delta and

BA.1 were not **(Figure 3A)**. Interestingly, Delta and BA.1 infections were enhanced
approximately 2-fold by E64d treatment, despite the same dose being previously shown to
partially inhibit Omicron infection in a transformed lung cell line[31]. However, addition of
camostat mesylate (an inhibitor of serine transmembrane proteases including TMPRSS2) alone
or in combination with E64d strongly blocked infection of WA1 and Delta in nasal ALI **(Figure
3A)**. These data suggest that WA1 and Delta enter nasal ALI primarily through a TMPRSS2-
dependent entry route at the plasma membrane. In contrast, infection of nasal ALI by BA.1 was
completely insensitive to blockade by camostat **(Figure 3A)**, suggesting the use of a divergent
entry pathway that is TMPRSS2-independent. The TMPRSS2-independence of Omicron
infection has been reported by multiple groups, and as a result, it has been inferred that Omicron
uses an endosomal, cathepsin-dependent route to enter human cells[31,32,36]. However, a recently
described metalloproteinase-mediated entry pathway for SARS-CoV-2 that may enable virus
fusion at the plasma membrane[11,45] prompted us to test the effect of incyclinide, a broad-
spectrum inhibitor of MMP and ADAM metalloproteinases. WA1, Delta, and BA.1 infections
were reduced by incyclinide treatment in primary ALI, but BA.1 was particularly sensitive
(inhibited by 27-fold compared to 4-fold inhibition of WA1 and 5-fold inhibition of Delta)
(Figure 3A). These findings suggest that processing of BA.1 Spike by metalloproteinases, but
not by TMPRSS2, enables BA.1 entry into primary nasal epithelia. To address whether a
metalloproteinase-dependent route of entry allows for fusion at the plasma membrane or at
endolysosomal membranes, we blocked endosomal trafficking with the endolysosomal
acidification inhibitor bafilomycin A1. WA1 and Delta infections were inhibited by bafilomycin
A1, suggesting that these viruses can utilize an endocytic pathway for entry (**Figure 3A**). In
contrast, bafilomycin A1 enhanced BA.1 infection 4-fold, suggesting that BA.1 utilizes an entry
pathway that is non-endocytic **(Figure 3A)**. We also tested the inhibitor sensitivity of
recombinant WA1 (Delta Spike) and WA1 (BA.1 Spike) to further clarify the entry pathway
unlocked by BA.1 Spike in nasal ALI. As seen with the full-length clinical isolates, WA1 (Delta
Spike) was inhibited by bafilomycin A1 while WA1 (BA.1 Spike) was not (**Figure 3B**).
However, WA1 (BA.1 Spike) was inhibited 10-fold by combined treatment of bafilomycin A1
and incyclinide. For WA1 (Delta Spike), treatment with bafilomycin A1 and incyclinide
inhibited infection to the same extent as bafilomycin A1 alone (**Figure 3B**). These results may
suggest that metalloproteinases at the cell surface promote entry mediated by BA.1 Spike, while
those in endosomes may promote entry mediated by Delta Spike. Overall, our data support a
model whereby Delta Spike may be processed by TMPRSS2 at the plasma membrane or by
metalloproteinases in endosomes, allowing for fusion at both sites in nasal ALI. In contrast,
BA.1 Spike may be processed primarily by metalloproteinases at or near the cell surface and
fusion ensues in this compartment, not in endosomes.
To further substantiate a role for metalloproteinases in the entry process driven by
different Spikes in nasal ALI, we tested the sensitivity of WA (Delta Spike) and WA1 (BA.1
Spike) to additional metalloproteinase inhibitors. Apratastat is an inhibitor of ADAM10,
ADAM17 and MMP-13, and batimastat is an inhibitor of MMP-1, MMP-2, MMP-3, MMP-7,
MMP-9, ADAM10, and ADAM17. While incyclinide inhibited WA1 (Delta Spike) infection,
apratastat and batimastat did not inhibit but instead modestly boosted infection (**Figure 3C**). In
contrast, both apratastat and batimastat reduced WA (BA.1 Spike) infection by 60% by (**Figure
3C**). These results reveal a functional divergence between Delta Spike and BA.1 Spike in terms
of their processing by metalloproteinases. Our results suggest that, at the concentrations used,
apratastat and batimastat inhibit a narrower range of metalloproteinases than incyclinide.

Nonetheless, the elevated sensitivity of WA1 (BA.1 Spike) to inhibition by apratastat and
batimastat suggests that BA.1 Spike can be processed by metalloproteinases targeted by both
compounds—such as ADAM10 and ADAM17. Furthermore, since apratastat and batimastat
decrease BA.1 Spike-mediated infection, ADAM10 and ADAM17 may be elements of a
productive entry pathway for BA.1 in nasal ALI. In contrast, processing of Delta Spike by these
metalloproteinases appears to have deleterious consequences for Delta entry, since apratastat and
batimastat boosted infection rather than inhibited it (**Figure 3C**). Therefore, BA.1 Spike has
evolved to utilize a unique entry route that is, while TMPRSS2-independent, dependent upon
ADAM10, ADAM17, and other metalloproteinases in the nasal epithelia. To assess whether
preferential usage of neuropilin-1 at the cell surface by BA.1 Spike may explain the increased
adherence of BA.1 to nasal cell epithelia and adoption of a unique entry pathway, we tested the
impact of the neuropilin-1 inhibitor EG00229 on nasal ALI infection by WA1 (WA1 Spike),
WA1 (Delta Spike), and WA1 (BA.1 Spike). WA1-Spike was most sensitive to inhibition by
EG00229, while Delta Spike was completely resistant, and BA.1 Spike conferred modest
sensitivity (**Supplemental Figure 3**). These results indicate that the extent of neuropilin-1
dependency does not explain the differential ability for WA1 Spike, Delta Spike, and BA.1 Spike
to mediate entry into nasal epithelia. Therefore, factors other than neuropilin-1 may be
responsible for the improved infectivity of BA.1 in nasal tissue.
249
250 **Omicron Spike enables an entry pathway into primary nasal epithelial cells that is resistant**
251 **to inhibition by type-I and type-III interferons**
252
Next, we explored whether the increased nasal cell infectivity exhibited by Omicron is
related to evasion of or resistance to cell-intrinsic antiviral defenses. During the preparation of
this manuscript, it was demonstrated that SARS-CoV-2 variants of concern displayed an
increasing resistance to inhibition by interferon (IFN) treatment, with Omicron exhibiting the
greatest degree of resistance [46]. However, the role of Spike in conferring IFN resistance was not
assessed, and it was unknown whether an Omicron-specific cellular entry pathway enabled IFN-
resistance in nasal epithelia. Therefore, we tested whether the unique entry requirements of
Omicron Spike allow for evasion of the IFN-induced antiviral state in primary nasal ALI (**Figure**
4A). Pre-treatment with IFN-beta or IFN-lambda inhibited WA1 in a dose-dependent manner (up
to two orders of magnitude) as measured by viral ORF1a RT-qPCR, while BA.1 or BA.2
infections were significantly less impacted **(Figure 4B)**. We also quantified infectious virus yield
under these conditions and confirmed that BA.1 and BA.2 achieved higher titers and were
inhibited by IFN to a significantly lesser extent than WA1 **(Figure 4C)**.   When we compared
the sensitivity of full-length clinical virus isolates with recombinant WA1 encoding different
Spike proteins, we found that the identity of Spike governed sensitivity to IFN. Compared to
WA1, which was inhibited 6-fold by low amounts of IFN-beta or IFN-lambda, WA1 (Delta
Spike) exhibited a slight decrease in sensitivity to IFN (**Figure 4D**). However, WA1 (BA.1
Spike) was completely insensitive to the same doses of IFN, and this closely resembled the
resistance profile of full-length BA.1 (**Figure 4D**). Therefore, BA.1 Spike is sufficient to confer
IFN-resistance to an otherwise IFN-sensitive virus, while Delta Spike is not. Moreover, these
results reveal the presence of host factors intrinsic to nasal cells that are interferon-inducible and
target the Spike-mediated entry step.
We also examined the IFN sensitivity of WA1, BA.1 and BA.2 in submerged primary
nasal monolayers **(Supplemental Figure 4A)**. While IFN-beta or IFN-lambda exposure strongly

blocked WA1 infection by more than 100-fold, the same amount of IFN inhibited BA.1 and
BA.2 to a much lesser extent (less than 2-fold) (**Supplemental Figure 4B**). To confirm that
Spike is a major determinant of IFN sensitivity in these cells, we generated pseudovirus
consisting of vesicular stomatitis virus (VSV) decorated with Spike from WA1, BA.1, or BA.2.
VSV-WA1 Spike was exquisitely sensitive to inhibition by IFN-beta or IFN-lambda in a dose-
dependent manner, with inhibition up to two orders of magnitude observed (**Supplemental
Figure 4C**). In contrast, VSV-BA.1 Spike and VSV-BA.2 Spike were much less sensitive to
inhibition by IFN-beta or IFN-lambda (**Supplemental Figure 4C**). Therefore, Spike is the major
determinant of the IFN resistance displayed by Omicron sublineages, and this feature can be
transferred to other viruses by pseudotyping. Additionally, we determined that cellular
attachment of WA1 and BA.1 in nasal monolayers was not affected by IFN-beta or IFN-lambda,
indicating that nasal cell-intrinsic host defenses act at a post-attachment step of the virus entry
process (**Supplemental Figure 4D**). Therefore, Omicron Spike enables evasion of restriction by
nasal cell-intrinsic host factors inhibiting the post-attachment step of virus entry.
291
292 **Amphotericin B partially relieves a block to infection mediated by WA1 Spike and Delta
293 Spike in primary nasal epithelia**
294
A limited number of interferon-stimulated genes are known to block the cellular entry
step of enveloped viruses[47]. Among the best characterized are the interferon-inducible
transmembrane (IFITM) proteins. IFITM proteins provide constitutive and interferon-induced
antiviral protection in many cell types by inhibiting the virus-cell membrane fusion step required
for entry[47,48]. While it is established that murine IFITM3 restricts SARS-CoV-2 infection and
limits viral disease in vivo[49,50], the roles played by human IFITM proteins during SARS-CoV-2
infection have remained controversial. For example, it was demonstrated that IFITM proteins can
inhibit SARS-CoV-2 infection in endosomes but can promote infection at the cell surface[51,52].
Furthermore, it has been suggested that endogenous human IFITM proteins in lung epithelial
cells enable SARS-CoV-2 infection by possibly acting as a receptor or attachment factor[53]. To
address whether IFITM proteins act as a barrier to SARS-CoV-2 infection in primary human
nasal epithelia, we tested the impact of the antifungal Amphotericin B on infection with
recombinant WA1 (WA1 Spike), WA1 (Delta Spike), and WA1 (BA.1 Spike). Amphotericin B
was previously reported to counteract the antiviral properties of human IFITM2 and IFITM3[54-56]
by restoring membrane fluidity in cellular membranes where IFITM3 is present[57]. In nasal ALI
pretreated with Amphotericin B or left untreated, we inoculated with WA1 (WA1 Spike), WA1
(Delta Spike), or WA1 (BA.1 Spike) and scored infection by mCherry fluorescence at 24 hours
post-inoculation. As described above, WA1 (BA.1 Spike) displayed a far superior capacity to
infect untreated nasal ALI relative to WA1 (WA1 Spike) and WA1 (Delta Spike) (**Figure 5**).
However, Amphotericin B treatment led to marked increases in WA1 (WA1 Spike) and WA1
(Delta Spike) infections, boosting them by approximately 7-fold. By comparison, Amphotericin
B had a negligible impact on WA1 (BA.1 Spike) infection at the same time point (**Figure 5**). We
also determined whether Amphotericin B impacted WA1 (BA.1 Spike) infection at earlier time
points when infected cells were less abundant. At 9 and 12 hours post-inoculation, Amphotericin
B marginally boosted infection of WA1 (BA.1 Spike) (less than 2-fold) (**Supplemental Figure
5**). Therefore, these results suggest that human IFITM proteins restrict WA1 (WA1 Spike) and
WA1 (Delta Spike) in primary nasal epithelia, while BA.1 Spike confers escape from this

restriction. Therefore, antiviral IFITM proteins form a part of the nasal cell-intrinsic defenses
that limit infection by early/ancestral SARS-CoV-2.
Overall, by combining experiments with authentic SARS-CoV-2, recombinant virus, and
Spike-decorated pseudovirus, we show that Spike is the major determinant of the enhanced
infectivity and interferon resistance exhibited by Omicron in primary human nasal epithelia, two
phenotypes which are associated with a unique dependence on metalloproteinases for cell entry
at or near the cell surface.
329
330 **Discussion**
331
Using entirely primary nasal epithelia from human adult donors, we demonstrate that
Omicron (including BA.1, BA.2, BQ.1, and XBB) exhibits drastically increased infectivity in
nasal tissue compared to early SARS-CoV-2, the D614G variant, and the Delta variant of
concern. This mapped to Omicron Spike and, at least in part, to an increased ability for Omicron
virions to adhere to the surface of nasal epithelia, which occurred independently of the presence
of cilia. Furthermore, we show that Omicron utilizes an entry route into cells that depends on
cellular metalloproteinases but that is independent of the transmembrane serine protease
TMPRSS2. Since we also demonstrate that Omicron Spike-mediated infection enables evasion of
the antiviral state induced by type-I and type-III IFN in this tissue, these results suggest that the
metalloproteinase-dependent entry pathway utilized by Omicron may promote escape from
constitutive and IFN-induced antiviral factors targeting virus entry.
During the preparation of this manuscript, it was reported that BA.1 exhibits a growth
advantage in primary human nasal epithelia compared to the Delta variant[31]. Here, using
recombinant SARS-CoV-2, we extend this finding by demonstrating that BA.1 Spike is
responsible for increased nasal cell infectivity. However, in another study, while BA.1 exhibited
a gain in nasal cell infectivity compared to Alpha, no significant differences in nasal cell
infectivity were observed between Delta and BA.1 [32]. This discrepancy is likely the result of
utilization of distinct cell culture models to compare relative infectivity. Our use of recombinant,
fluorogenic SARS-CoV-2 in nasal ALI derived from pools of 14 human adult donors shows that
Spike from BA.1 promotes a gain in nasal cell infectivity compared to Delta Spike. Therefore,
the emergence and evolution of Omicron involved unique mutations in Spike that contributed to
enhanced nasal cell binding and entry, and this may explain how Omicron replaced previously
circulating variants of concern and continues to persist in human populations.
It was previously reported that human nasal airways express relatively low levels of
TMPRSS2, and combined with the demonstration that Omicron exhibits TMPRSS2-
independence in transformed cell lines, others have inferred that Omicron is more reliant on
endosomal cathepsins for entry [32]. However, our findings with primary nasal epithelia reveal that
TMPRSS2-independence does not necessarily imply the use of an endosomal entry route by
Omicron. Metalloproteinases such as MMP-14, MMP-16, ADAM10, and ADAM17 are also
highly expressed in upper airways and have been shown to activate Omicron Spike-mediated
fusion at the surface of several transformed cell lines [11]. Here, we demonstrate that the primary
route of entry of Omicron in primary nasal epithelia is mediated by metalloproteinases. The pan-
metalloproteinase inhibitor incyclinide strongly blocked BA.1 infection in nasal ALI, but the
endosomal cathepsin inhibitor E64d and the endolysosomal acidification inhibitor bafilomycin
A1 did not. Instead, these endosomal inhibitors boosted infection of BA.1 in nasal epithelia,
suggesting that the endosomal entry route is deleterious for BA.1 in this tissue. Bafilomycin A1

inhibits vacuolar-ATPase to prevent endolysosomal acidification, and furthermore, endosomal
maturation and transport between early and late endosomes is repressed [58]. Therefore, our results
suggest that processing of Spike by metalloproteinases enables fusion at the plasma membrane
and/or in early or recycling endosomes, while late endosomes or lysosomes and the cathepsins
found therein are unlikely to be involved.
One possibility explaining the enhanced capacity for Omicron Spike to adhere to and
enter nasal cells (either in the presence or absence of ciliated cells) is its increase in net overall
positive charge compared to previous variants of concern[59]. It has also been proposed that
Omicron Spike may adhere more strongly to attachment factors present at the epithelial cell
surface, including heparan sulfate[60,61] and neuropilin-1[62]. In this study, sensitivity to neuropilin-1
inhibitor did not correlate with the nasal cell infectivity of WA1, Delta, or BA.1, suggesting that
other cellular factors may play a role (**Supplemental Figure 3**). A significant aim of future
research will be to establish whether mutations in Omicron Spike promote increased residence
time at the cell surface and whether this influences its dependence on certain cellular proteases
for cleavage and the triggering of fusion. It is possible that longer residency at the cell surface
may negatively interfere with processing by TMPRSS2 and steer processing towards
metalloproteinases.
Another factor that likely contributed to the spread and dominance of Omicron lineages is
their resistance to the antiviral properties of IFN. Early/ancestral SARS-CoV-2 was previously
reported to antagonize IFN signaling in cells, and nearly every viral protein encoded by SARS-
CoV-2 has been shown to contribute to evasion of IFN through various mechanisms[63-65]. While
the relative insensitivity of Omicron to exogenous administration of multiple IFN types has been
reported previously[46,66], the viral proteins responsible for rendering IFN ineffective against
Omicron were unknown. Other studies showed that Omicron exhibits a decreased capacity to
interfere with IFN production and signaling, relative to early/ancestral SARS-CoV-2 or
Delta[44,67-69], and our measurements of type-I IFN production by Omicron-infected nasal ALI
affirm this finding. However, in one case, Omicron was purported to induce a relatively inferior
IFN response compared to Delta [66], and this discrepancy underlines the importance of assessing
this activity in primary human upper airway epithelia. Overall, we postulate that Omicron
exhibits resistance to IFN not because of its ability to interfere with the production or function of
IFN from infected cells, but rather, because Omicron Spike enables passage into nasal cells in a
manner which is insensitive to IFN-stimulated proteins. In addition, it is possible that Omicron
displays enhanced infectivity in nasal cells even in the absence of IFN because Omicron Spike
confers resistance to antiviral factors that are constitutively expressed in nasal epithelia and
further upregulated by type-I or type-III IFN. Our results indicating that Amphotericin B
partially relieves a constitutive barrier to infection mediated by IFITM proteins provides direct
evidence for this possibility. Furthermore, our findings provide a new explanation for why
Omicron exhibits decreased IFN antagonism—if the Spike-mediated entry process of Omicron is
resistant to type-I and type-III IFN, the selective advantage to antagonize IFN pathways may be
lost. Furthermore, the improved capacity for Omicron lineages to infect nasal cells despite the
induction of an antiviral state therein provides a plausible scenario for how Omicron displaced
previous variants of concern. It was demonstrated that IFN produced from Omicron-infected
cells can inhibit infection by Influenza A virus [67], so it is plausible that the IFN response elicited
by Omicron infection may render nasal epithelia resistant to infection by previous SARS-CoV-2
variants of concern.

Our description of the IFN resistance conferred by Omicron Spike has implications for
the ongoing development of type-I and type-III IFN as antiviral therapeutics for the prevention or
treatment of SARS-CoV-2 infection and COVID-19. A case has been made for the deployment
of type-III IFN in the context of SARS-CoV-2 infection, since the receptor for type-III IFN is
expressed in the respiratory tract but is not widely distributed elsewhere (resulting in fewer
unintended side effects such as inflammation)[63,70]. As an important proof-of-concept, nasal
administration of IFN-lambda was protective against SARS-CoV-2 variants, including Omicron,
in mice [71]. However, clinical trials thus far in humans have reported conflicting findings. One
such study reported that participants who were administered a single injection of IFN-lambda
following symptom onset showed no significant reductions in viral shedding or symptom
severity compared to placebo[72], while another reported that a single injection upon symptom
onset resulted in reduced viral loads[73] and reduced COVID-19-related hospitalizations[74].
However, in neither case was it determined how IFN-lambda fared in individuals infected with
Omicron variants. Since Omicron displays a greater degree of IFN-resistance compared to earlier
variants, additional clinical studies are needed to test the suitability of IFN-lambda in the fight
against contemporaneous Omicron and additional variants that arise in the future.

While some mutations in Omicron Spike may have conferred Omicron with superior
evasion of vaccine-elicited neutralizing antibodies compared to the Delta variant, our findings
indicate that mutations in Omicron Spike also allow for efficient seeding of infection in the nasal
epithelium and resistance to the interferon-induced antiviral state therein. Our findings provide
mechanistic insight into the basis for the efficient transmission and persistence of Omicron in
vaccinated and unvaccinated human populations. Future efforts aimed at understanding how
individual Spike mutations contribute to these different phenotypes may enable a better
understanding of the selective forces driving their emergence.

Figures and Figure Legends

440
Figure 1. Omicron BA.1 exhibits superior replicative fitness relative to early SARS-CoV-2
and Delta in primary human nasal epithelial cells.

(A) Primary human nasal epithelial cells (pooled from 3 human donors) were cultured as
undifferentiated, submerged monolayers and challenged with SARS-CoV-2. Acetylated tubulin
levels were determined by anti-AcTub immunofluorescence, ACE2 levels were determined by
anti-ACE2 immunofluorescence, and DAPI was used to stain nuclei. 3D reconstructions of *xy*
and *yz* fields are shown. Cartoon made with Biorender.com. (B) Primary human nasal epithelial
cells (cells from three human donors, pooled) or (C) human small air way (lung) epithelial cells

(cells from three human donors, pooled) were challenged with replication competent SARS-
CoV-2 WA1 or Omicron BA.1 at an MOI of 0.05. Total cellular RNA was extracted and viral
ORF1a was quantified by RT-qPCR at the indicated time points. Relative viral RNA abundance
compared to actin was determined by the 2^(-ΔΔCT) method. ORF1a abundance of WA1 at 24
hours post inoculation was set to 1. (D) Virus replication in primary human nasal epithelial cells
was measured by detecting ORF1a levels with RT-qPCR at 24 hours post inoculation with WA1,
Delta, or BA.1 at an MOI of 0.05. (E) As in (D), except that primary human nasal epithelial cells
were treated with remdesivir (10 µM) prior to and during inoculation with WA1, Delta, or BA.1.
ORF1 abundance of each virus treated with DMSO was set to 1. All results are represented as
means plus standard error from three independent infections (symbols represent biological
replicates). Statistically significant differences (* $P < 0.05$) between the indicated condition and
the corresponding data point of WA.1 were determined by one-way ANOVA.
462
463
464

Figure 2. Omicron Spike enables enhanced SARS-CoV-2 infectivity in primary nasal epithelia cultured at the air-liquid interface.

(A) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface to mimic the human respiratory tract and challenged with SARS-CoV-2. Acetylated tubulin levels were determined by anti-AcTub immunofluorescence, ACE2 levels were determined by anti-ACE2 immunofluorescence, and DAPI was used to stain nuclei. 3D reconstructions of *xy* and *yz* fields are shown. Scale bars = 10 μm. Cartoon made with Biorender.com. (B) Cells were subjected to total cellular RNA extraction at 48 hours post inoculation with 10000 plaque forming units of WA1, BA.1, or BA.2, and RT-qPCR of viral ORF1a was performed. Relative ORF1a transcript abundance compared to actin was determined

by the 2^(-ΔΔCT) method. ORF1a abundance of WA1 at 48 hours post inoculation was set to 1.
(C) Infectious virus titers from cell culture medium collected at 48 hours post-inoculation were
measured by challenging Vero E6-TMPRSS2 cells followed by fixation and immunostaining
with anti-N antibody. Infectious units were quantified by measuring focus forming units with
high-content imaging. Symbols represent results from three independent infections of Vero E6-
TMPRSS2 cells. (D) RT-qPCR of cellular *IFNB* was performed at 48 hours post inoculation.
Relative *IFNB* transcript abundance compared to actin was determined by the 2^(-ΔΔCT)
method. *IFNB* abundance of WA1 at 48 hours post inoculation was set to 1. (E) Viral ORF1a
RT-qPCR was performed at 48 hours post-inoculation with 10000 plaque forming units of WA1,
D614G, Delta, BA.1, or BQ.1. Relative ORF1a transcript abundance compared to actin was
determined by the 2^(-ΔΔCT) method. ORF1a abundance of WA1 at 48 hours post inoculation
was set to 1. (F) RT-qPCR of cellular *IFNB* was performed at 48 hours post inoculation. Relative
IFNB transcript abundance compared to actin was determined by the 2^(-ΔΔCT) method. *IFNB*
abundance of WA1 at 48 hours post inoculation was set to 1. (G) Recombinant WA.1 encoding
mCherry and Spike protein from WA1, Delta, or BA.1 (WA1-mCherry (WA1 Spike), WA1-
mCherry (Delta Spike), and WA1-mCherry (BA.1 Spike)) were used to inoculate primary human
nasal epithelial cells (pooled from 14 human donors) cultured at the air-liquid interface, and
infection was measured by mCherry fluorescence at 24 hours post-inoculation. Summary
mCherry fluorescence from three independent infections is shown. The fluorescence intensity of
WA1 (WA1 Spike)-mCherry was set to 1. Representative fields of view are shown. Scale bars =
300 µm. All results are represented as means plus standard error from three independent
infections (symbols represent biological replicates). Statistically significant differences (* $P <$
0.05) between the indicated condition and the corresponding data point of WA.1 were
determined by one-way ANOVA.
502
503

504
Figure 3. Entry of Omicron BA.1 into primary nasal epithelial cells is mediated by proteases of the MMP/ADAM families and occurs independently of endocytosis.

507
(A) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface and pretreated with 10 µM E64d, 10 µM camostat mesylate, a combination of 10 µM E64d and 10 µM camostat mesylate, 10 µM incyclinide, or 1 µM bafilomycin A1 for 2 hours. Afterwards, cells were challenged with WA.1, Delta, or BA.1 at an MOI of 0.05. At 48 hours post inoculation, cells were subjected to total cellular RNA extraction and ORF1a levels were measured by RT-qPCR. Relative ORF1a transcript abundance compared to actin was determined by the 2^(-ΔΔCT) method. For each virus, ORF1a abundance in the DMSO condition was set to 1. All results are represented as means plus standard error from three independent infections (symbols represent biological replicates). Statistically significant differences (* $P <$ 0.05) between the indicated condition and the corresponding condition of WA.1 were determined by one-way ANOVA. ORF1a abundance in the DMSO condition at 48 hours post inoculation was set to 1. (B) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface and pretreated with 1 µM bafilomycin A1 or 1 µM bafilomycin A1 and 10 µM incyclinide for 2 hours. Afterwards, cells were challenged with WA1-based recombinant viruses encoding Delta Spike or BA.1 Spike at an MOI of 0.05. At 48

hours post-inoculation, ORF1a levels were measured by RT-qPCR. For each virus, ORF1a
abundance in the DMSO condition was set to 1. (C) Primary human nasal epithelial cells (pooled
from 14 human donors) were cultured at the air-liquid interface and pretreated with 10 µM
incyclinide, 10 µM aprastatat, or 10 µM batimastat for 2 hours. Afterwards, cells were
challenged with WA1-based recombinant viruses encoding Delta Spike or BA.1 Spike at an MOI
of 0.05. At 48 hours post-inoculation, ORF1a levels were measured by RT-qPCR. For each
virus, ORF1a abundance in the DMSO condition was set to 1. Statistically significant differences
(* $P < 0.05$) between the indicated condition of BA.1 and the corresponding condition of Delta
were determined by one-way ANOVA. All results are represented as means plus standard error
from three independent infections (symbols represent biological replicates).

Figure 4. Omicron Spike enables an entry pathway into primary nasal epithelial cells that is resistant to inhibition by type-I and type-III interferons.

(A) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface, pre-treated for 18 hours with the indicated amounts of IFN-beta or IFN-lambda, and challenged with 10000 plaque forming units of WA1, BA.1, or BA.2. Cartoon made with Biorender.com. (B) At 48 hours post inoculation, total RNA was extracted from cells and subjected to RT-qPCR. ORF1a levels were measured compared to actin by the $2^{\wedge}(-\Delta\Delta CT)$ method. For each virus, ORF1a abundance in the No IFN condition was set to 1. (C) At 48 hours post inoculation, infectious virus titers from cell culture medium were measured by challenging Vero E6-TMPRSS2 cells followed by fixation and immunostaining with anti-N antibody. Infectious units were quantified by measuring focus forming units by high-content imaging. Symbols represent independent infections of Vero E6-TMPRSS2 cells. (D) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface, pre-treated for 18 hours with the indicated amounts of IFN-beta or IFN-lambda, and challenged with 10000 plaque forming units of WA1, WA1-mCherry (Delta Spike), WA1-mCherry (BA.1 Spike), or BA.1. ORF1a levels were measured by RT-qPCR and compared to actin by the $2^{\wedge}(-\Delta\Delta CT)$ method. For each virus, ORF1a abundance in the No IFN condition was set to 1. All results are represented as means plus standard error from three independent infections (symbols represent biological replicates). Statistically significant differences (* $P < 0.05$) between the indicated condition and the corresponding data point of WA.1 were determined by one-way ANOVA.

557
Figure 5. Amphotericin B partially relieves a block to infection mediated by WA1 Spike and Delta Spike in primary nasal epithelia.

560
Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-
liquid interface, pre-treated with 1 µM Amphotericin B for two hours or untreated, and
inoculated with 10000 plaque forming units of WA1-mCherry (WA1 Spike), WA1-mCherry
(Delta Spike), or WA1-mCherry (BA.1 Spike). mCherry fluorescence was measured at 24 hours
post-inoculation by high-content imaging. The fluorescence intensity of the untreated condition
for each virus was set to 1. Scale bar = 300 µm. Ampho B; amphotericin B.

567
568 **Supplemental Figure 1.**
569
(A) Primary human nasal epithelial cells (pooled from 3 human donors) were cultured as
undifferentiated, submerged monolayers and challenged with WA1 at an MOI of 0.05. Cell
culture supernatants were collected at 24, 48, and 72 hours post-inoculation and added to Vero
E6-ACE2-TMPRSS2 cells. 24 hours later, Vero E6-ACE2-TMPRSS2 cells were subjected to
total RNA extraction and viral ORF1a RT-qPCR was performed. In addition, productive
infection of Vero E6-ACE2-TMPRSS2 by the 24 hours post-inoculation supernatant was
confirmed by anti-N immunofluorescence microscopy. (B) Primary human nasal epithelial cells
(cells from three human donors, pooled) were inoculated with WA1 or BA.1 at an MOI of 0.05.

Total cellular RNA was extracted and viral ORF1a was quantified by RT-qPCR at 24 hours post-
inoculation. Absolute ORF1a RNA copy numbers were calculated by comparison to an ORF1a
standard curve. (C) Relative viral ORF1a RNA abundance compared to actin was determined by
the $2\char`^(-\Delta\Delta CT)$ method. ORF1a abundance of WA1 was set to 1. (D) Primary human nasal
epithelial cells (cells from three human donors, pooled) were inoculated with WA1 or BA.1 at an
MOI of 0.05 on ice. At 1 hour post-inoculation, total cellular RNA was extracted and viral
ORF1a was quantified by RT-qPCR to measure virus adherence to cells. Relative viral ORF1a
RNA abundance compared to actin was determined by the $2\char`^(-\Delta\Delta CT)$ method. ORF1a
abundance of WA1 was set to 1. (E) HEK293T-ACE2 cells were inoculated with WA1 or BA.1
at an MOI of 0.05. 24 hours post-inoculation, total cellular RNA was extracted and viral ORF1a
was quantified by RT-qPCR. Relative viral ORF1a RNA abundance compared to actin was
determined by the $2\char`^(-\Delta\Delta CT)$ method. ORF1a abundance of WA1 was set to 1. (F) Vero E6-
TMPRSS2 cells were inoculated with WA1 or BA.1 at an MOI of 0.05. 24 hours post-
inoculation, cells were fixed, stained with anti-N antibody, and infection was scored by flow
cytometry. (G) Primary human nasal epithelial cells (cells from three human donors, pooled)
were inoculated with WA1 or BA.1 ($5\times10\char`^7$ copies of absolute ORF1a RNA used as input). 24
hours post-inoculation, total cellular RNA was extracted and viral ORF1a was quantified by RT-
qPCR. Relative viral ORF1a RNA abundance compared to actin was determined by the $2\char`^(-$
$\Delta\Delta CT)$ method. ORF1a abundance of WA1 was set to 1. All results are represented as means
plus standard error from three independent infections (symbols represent biological replicates).
Statistically significant differences (* $P < 0.05$) between the indicated condition of BA.1 and the
corresponding condition of WA.1 were determined by one-way ANOVA.

600
Supplemental Figure 2.
(A) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the
air-liquid interface and inoculated with 10000 plaque forming units of WA1 or BA.1. Total
cellular RNA was extracted and viral ORF1a was quantified by RT-qPCR at 24 hours post-
inoculation. Absolute ORF1a RNA copy numbers were calculated by comparison to an ORF1a
standard curve. (B) Primary human nasal epithelial cells (pooled from 14 human donors) were
cultured at the air-liquid interface and inoculated with 10000 plaque forming units of WA1,
D614G, Delta, BA.1, or BQ.1. RT-qPCR of cellular *IFITM3* was performed at 48 hours post
inoculation. Relative *IFITM3* transcript abundance was compared to actin using the 2^(-ΔΔCT)

method. *IFITM3* abundance in non-inoculated (NI) cells was set to 1. Statistically significant differences (* $P < 0.05$) between the indicated condition and the NI condition were determined by one-way ANOVA. (C) Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface and inoculated with 10000 plaque forming units of D614G, BA.1, BQ.1, or XBB. At 48 hours post-inoculation, total cellular RNA was extracted and viral ORF1a was quantified by RT-qPCR. Relative ORF1a abundance was determined by comparing to actin using the $2^{\wedge}(-\Delta\Delta CT)$ method. Statistically significant differences (* $P < 0.05$) between the indicated condition and D614G were determined by one-way ANOVA. (D) 10000 plaque forming units of recombinant WA.1 encoding mCherry and Spike protein from WA1, Delta, or BA.1 (WA1-mCherry (WA1 Spike), WA1-mCherry (Delta Spike), and WA1-mCherry (BA.1 Spike)) were used to inoculate primary human nasal epithelial cells (pooled from 14 human donors) cultured at the air-liquid interface. At 48 hours post-inoculation, total cellular RNA was extracted and viral ORF1a was quantified by RT-qPCR. Relative ORF1a abundance was determined by comparing to actin using the $2^{\wedge}(-\Delta\Delta CT)$ method. Statistically significant differences (* $P < 0.05$) between the indicated condition and WA1-mCherry (WA1 Spike) were determined by one-way ANOVA. (E) As in (D), except that $5 \times 10^{\wedge}7$ copies of absolute ORF1a RNA were used as input. At 48 hours post-inoculation, infection was measured by viral ORF1a RT-qPCR. Relative ORF1a abundance was determined by comparing to actin using the $2^{\wedge}(-\Delta\Delta CT)$ method. Statistically significant differences (* $P < 0.05$) between the indicated condition WA.1 were determined by one-way ANOVA. All results are represented as means plus standard error from three independent infections (symbols represent biological replicates). (F) 10000 plaque forming units of recombinant WA.1 encoding mCherry and Spike protein from Delta or BA.1 (WA1-mCherry (Delta Spike), and WA1-mCherry (BA.1 Spike)) were used to inoculate primary human nasal epithelial cells (pooled from 14 human donors) cultured at the air-liquid interface. At 6, 9, and 12 hours post-inoculation, infection was measured by mCherry fluorescence. Scale bar = 300 microns.

Supplemental Figure 3.

Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-liquid interface, pre-treated with 100 µM EG00229 or DMSO for two hours, and inoculated with

642 10000 plaque forming units of WA1, Delta, or BA.1. mCherry fluorescence was measured at 24
643 hours post-inoculation by high-content imaging. The fluorescence intensity of the DMSO-treated
644 condition for each virus was set to 1. Scale bar = 300 µm. Results are represented as means plus
645 standard error from one infection (symbols represent three RT-qPCR runs).
646

647
648 **Supplemental Figure 4.**
649
(A) Primary human nasal epithelial cells (pooled from 3 human donors) were cultured as
undifferentiated, submerged monolayers, treated with IFN-beta or IFN-lambda for 18 hours, and
challenged with SARS-CoV-2. Cartoon made with Biorender.com. (B) Cells were pre-treated

with 2 units of IFN-beta or 2 units of FN-lambda for 18 hours, and inoculated with WA1, BA.1,
or BA.2 at an MOI of 0.05. Total RNA was extracted from cells at 24 hours post inoculation, and
ORF1a levels were measured by RT-qPCR. Relative ORF1a abundance was determined by
comparing to actin using the 2^(-ΔΔCT) method. For each virus, ORF1a levels in the absence of
IFN were set to 1. (C) Primary human nasal epithelial cells (pooled from 3 human donors) were
pre-treated with the indicated amounts of IFN-beta or IFN-lambda for 18 hours and challenged
with VSV-based pseudovirus decorated with Spike from WA1, BA.1, or BA.2. At 24 hours post
inoculation, luciferase activity was measured from lysed cells. Luciferase activity of WA1,
BA.1, and BA.2 pseudoviruses in the absence of IFN were set to 1. (D) Primary human nasal
epithelial cells (pooled from 3 human donors) were pre-treated with 2 units of IFN-beta or 5
ng/mL IFN-lambda for 18 hours, and inoculated with WA1 or BA.1 at an MOI of 0.05 on ice.
Total RNA was extracted from cells at 1 hour post inoculation, and ORF1a levels were measured
by RT-qPCR. Relative ORF1a abundance was determined by comparing to actin using the 2^(-
ΔΔCT) method. ORF1a levels of WA.1 in the absence of IFN were set to 1. All results are
represented as means plus standard error from three independent infections (symbols represent
biological replicates). Statistically significant differences (* $P < 0.05$) between the indicated
condition and the corresponding No IFN condition were determined by one-way ANOVA.

670
671   **Supplemental Figure 5.**

672
Primary human nasal epithelial cells (pooled from 14 human donors) were cultured at the air-
liquid interface, pre-treated with 1 μM Amphotericin B for two hours or untreated, and
inoculated with 10000 plaque forming units of WA1-mCherry (BA.1 Spike). mCherry
fluorescence was measured at 9 and 12 hours post-inoculation by high-content imaging. The

fluorescence intensity of the untreated condition was set to 1. Scale bar = 300 µm. Ampho B;
amphotericin B.
679
680 **Materials and Methods**
681
682 **Tissue culture**
683
Frozen primary human nasal airway epithelial cells (hNAEC) were purchased from Epithelix and
cultured as submerged monolayers with hAEC Culture Medium (Epithelix). hNAEC from three
human donors were thawed and pooled, and cells were passed every 5 days. Fresh primary
human nasal airway epithelial cells cultured at the liquid-air interface (nasal ALI) were obtained
from Epithelix (MucilAir, Pool of Donors) and cultured with MucilAir culture medium
(Epithelix) in trans-well plates provided by the company; culture medium was replaced every 2
days. Nasal ALI cultures were maintained for 2-5 weeks in our laboratories prior to inoculation
with SARS-CoV-2. hNAEC and Nasal ALI were fixed with 4% paraformaldehyde,
permeabilized with 0.1% Triton-X, and stained with anti-ACE2 (Abcam; ab15348), anti-Acetyl-
a-Tubulin (Lys40) (D20G3) (Cell Signaling; 5335), anti-phalloidin (Cell Signaling; 8878), and
DAPI to assess ACE2 levels, the presence of ciliated cells, and pseudostratification of nuclei,
respectively. Images were obtained using the Zeiss LSM 880 workstation and AiryScan
capability and were processed into 3D reconstructions by volume rendering using Imaris
software. Primary human small airway (lung) epithelial cells (hSAEC) were purchased from
ATCC (PCS-301-010) and were cultured with Airway Epithelial Cell Basal Medium (ATCC,
PCS-300-030) and the Bronchial Epithelial Growth Kit (ATCC, PCS-300-400). hSAEC from
three human donors were thawed and pooled, and cells were passed every 5 days. Vero E6 were
purchased from ATCC (CRL-1586) and Vero E6-TMPRSS2 cells were generated from them by
introducing the TMPRSS2 open reading frame using the Sleeping Beauty transposon system.
Vero E6-ACE2-TMPRSS2 cells were obtained from BEI Resources (NR-54970). HEK293T-
ACE2 cells were obtained from BEI Resources (NR-52511).
705
706 **Production of SARS-CoV-2 variants and infections**
707
SARS-CoV-2 isolate USA/WA1/2020 (WA1; NR-52281), isolate hCoV-19/USA/HI-CDC-
4359259-001/2021 (Lineage B.1.1.529; Omicron BA.1 variant; NR-56475), and isolate hCoV-
19/USA/NY-MSHSPSP-PV56475/2022 (Lineage BA.2.12.1; Omicron BA.2 variant; NR-
56782), isolate hCoV-19/USA/CA-Stanford-109_S21/2022 (Lineage XBB; Omicron XBB
variant; NR-58927) and hCoV-19/USA/MDHP38960/2022 (Lineage BQ; Omicron BQ.1
variant) were obtained from BEI Resources. hCoV-19/New York-PV08410/2020 (Lineage B.1;
D614G variant; NR-53514) was a gift from John Dye (USAMRIID). Isolate hCoV-19/USA/MD-
HP05285/2021 VOC Delta G/478K.V1 (Lineage B.1.617.2 + AY.1 + AY.2; Delta variant) was
obtained from Andrew Pekosz (Johns Hopkins University). Recombinant WA1-mCherry virus
was rescued from a SARS-CoV-2 cDNA construct[75], and mCherry was inserted at the N-
terminus of the *N* gene. A P2A linker was placed between mCherry and N. To generate WA1-
mCherry (BA.1 Spike) and WA1-mCherry (Delta Spike), the sequence encoding WA1 Spike
was replaced with Spike from BA.1 or Delta, respectively. All viruses were grown in Vero E6-
TMPRSS2 cells and harvested cell culture supernatants were titrated in Vero E6-TMPRSS2 cells
to calculate infectious titers. Infectious titers were determined by focus-forming unit assay

whereby foci of infection were measured using anti-nucleocapsid (N) antibody (Invitrogen;
MA1-7403). Where indicated, ORF1a RNA titers were also calculated for some viruses to
approximate virus particle quantity.
Infections of hNAEC or hSAEC monolayers were performed at a multiplicity of infection (MOI)
of 0.05 as follows: cells were inoculated with virus suspension for 2 hours at 37°C; inoculum
was removed; cells were washed with PBS and returned to 37°C for the time indicated. For
detection of virus attachment, hNAEC were inoculated with virus suspension for 1 hour on ice;
inoculum was removed; cells were washed three times with cold PBS.
For detection of infection or virus attachment by RT-qPCR, cells were lysed with Trizol (Sigma).
Viral replication was measured using RT-qPCR amplification of viral ORF1a, as previously
described[50]. Cells lysed with Trizol were mixed with chloroform (Sigma) at a 5:1
(Trizol:chloroform) ratio. Mixed samples were mixed thoroughly and incubated at room
temperature for 10 minutes, followed by centrifugation at 12000 x G for 5 minutes to allow
separation of the aqueous and organic phases. Equal volumes of 70% ethanol were added to the
aqueous phases, mixed thoroughly, and incubated at room temperature for 5 minutes. RNA
purification was performed using the PureLink RNA Mini Kit (Invitrogen) according to
manufacturer's instructions. Purified RNA product was immediately used with the One-step
PrimeScript RT-PCR Kit (Takara). Primers and probes were obtained from IDT. The primers
and probes used to amplify and quantify ORF1a are as follows (5'-3'): ORF1a-F
AGAAGATTGGTTAGATGATGATAGT; ORF1a-R TTCCATCTCTAATTGAGGTTGAACC;
ORF1a-P FAM/TCCTCACTGCCGTCTTGTTGACCA/BHQ13. The primers and probes used to
amplify and quantify beta actin (ACTB) are as follows (5'-3'): ACTB-F
ACAGAGCCTCGCCTTTG; ACTB-R CCTTGCACATGCCGGAG; ACTB-P 56-
FAM/TCATCCATG/ZEN/GTGAGCTGGCGG/31ABkFQ. The primers and probes used to
amplify and quantify IFITM3 are as follows (5'-3'): IFITM3-F
ACCATGAATCACACTGTCCAAACCTT; IFITM3-R CCAGCACAGCCACCTCG; IFITM3-P
FAM/ZEN-CTCTCCTGTCAACAGTGGCCAGCCCC-IBFQ. The primers and probes used to
amplify and quantify IFNB are as follows (5'-3'): IFNB-F GAAACTGAAGATCTCCTAGCCT;
IFNB-R GCCATCAGTCACTTAAACAGC; IFNB-P 56-
FAM/TGAAGCAAT/ZEN/TGTCCAGTCCCAGAGG/3IABkFQ. Reaction mixtures of 20 µL
(including 2.2 µL total RNA, 0.2 µM forward and reverse primers, and 0.1 µM probe) were
subjected to reverse transcription (5 min at 45°C, followed by 10 s at 95°C) and 40 cycles of
PCR (5 s at 95°C followed by 20 s at 60°C) in a CFX Opus 96 Real-Time PCR System
(BioRad). Results were analyzed by the Comparative Ct Method (ΔΔCt Method) [76]. RNA levels
for viral ORF1a, IFNB, and IFITM3 were normalized to cellular ACTB.
For infections of primary nasal epithelial cells cultured at the air-liquid interface, the apical (air-
exposed) surface was gently rinsed with 100 µL cell culture medium to partially remove mucous
layers. Virus suspension in a volume of 50 µL was added to the apical surface and cells were
incubated for 2 hours at 37°C. Inoculum was then removed and 100 µL PBS was used to gently
wash the apical surface. Cells were then returned to 37°C. At 48 hours post inoculation, cells
were lysed with Trizol and subjected to RNA extraction and RT-qPCR as indicated above. To
quantify the infectious virus particles produced into the cell culture supernatant, medium was
removed, 200 µL PBS was added to the apical surface, and cells were incubated at room
temperature for 20 minutes before the PBS was recovered. A focus-forming units assay was
performed by inoculating Vero E6-TMPRSS2 cells with the recovered volume. At 7 hours post
inoculation, Vero E6-TMPRSS2 cells were fixed with 4% paraformaldehyde and stained with

anti-N (Invitrogen; MA1-7403). The number of fluorescent foci was measured using a Cytation 5
Cell Imaging Multimode Reader (BioTeK).
For detection of recombinant WA-mCherry infection, the Cytation 5 Cell Imaging Multimode
Reader (BioTeK) was used to measure mCherry fluorescence at the indicated hours post
inoculation. Quantification of mCherry fluorescence was performed by measuring integrated
signal intensities in Fiji.
775
776 **Production of VSV-Spike pseudoviruses and infections**
777
Spike sequences from WA.1, BA.1, or BA.2 were codon-optimized for expression in human
cells, synthesized with a 6xHis tag on the amino-terminus, and cloned into pcDNA3.1 (+) by
GenScript. HEK293T cells were seeded in a 10 cm dish and transfected with 12 μg pcDNA3.1
Spike plasmid using Lipofectamine 2000 (Thermo Fisher). Forty-eight hours after transfection,
culture medium was removed from cells, and 1 mL VSV-luc/GFP plus VSV-G (seed particles)
was added. Twenty-four hours after infection, virus supernatants were collected, clarified by
centrifugation at $500g$ for 5 minutes followed by filtration with a 45 μm filter, and stored at -
80°C. A total of 50 μL virus supernatants was added to submerged hNAEC from three pooled
human donors, and 24 hours post inoculation, cells were lysed with Passive Lysis Buffer
(Promega). Luciferase activity was measured on a Perkin Elmer MicroBeta 2450 microplate
luminometer using the Luciferase Assay System (Promega). 50 μL volumes of VSV-WA1,
VSV-BA.1, and VSV-BA.2 were found to infect Vero E6-TMPRSS2 cells to similar extents,
suggestive of similar infectious titers.
791
792 **Interferons and Inhibitors**
793
Recombinant human IFN-β (beta) (300-02BC) and human IFN-λ (lambda) (300-02L) were
obtained from PeproTech and were used to test IFN sensitivity of full-length, authentic SARS-
CoV-2 variants (IFN was added 18 hours prior to inoculation and removed prior to virus
addition). Recombinant human IFN-β (beta) 1a (11415-1) and human IFN-λ (lambda) (11725-1)
were obtained from PBL Assay Science and were used to test IFN sensitivity of VSV-based
pseudoviruses (IFN was added 18 hours prior to inoculation and removed prior to virus
addition). The following inhibitors were reconstituted in DMSO: E64d (Sigma; E8640), camostat
mesylate (Sigma; SML0057), incyclinide (MedChemExpress; HY-13648), apratastat
(MedChemExpress; HY-119307), batimastat (SelleckChem: S7155), EG00229 (SelleckChem;
E1119). Bafilomycin A1 (Sigma; SML1661) was received as a ready-made solution in DMSO
and amphotericin B (Sigma; A2942) was received as a ready-made solution in deionized water.
805
806 **Biosafety Approval**
807
808 This study was conducted in compliance with all relevant local, state, and federal regulations.
809 Approval for the generation and use of recombinant SARS-CoV-2 WA1-mCherry and its
810 variants using Biosafety Level 3 practices was provided by the NIAID Institutional Biosafety
811 Committee following evaluation by the Dual Use Research of Concern Institutional Review
812 Entity (case number RD-22-X1-11; PI: Jonathan W Yewdell).
813
814 **Disclaimer**

815
816 The content of this publication does not necessarily reflect the views or policies of the
817 Department of Health and Human Services, nor does mention of trade names, commercial
818 products, or organizations imply endorsement by the US Government.
819
820 **Acknowledgements**
821
We would like to thank Brett Eaton, Elena Postnikova, Michael Murphy, Sushma Bhosle, Julie
Tran, and Jillian Geiger (NIAID, Integrated Research Facility) for reagents and protocols to
quantify viral ORF1a by RT-qPCR, Vincent Munster (NIAID) and Michael Letko (Washington
State University) for reagents and protocols to produce VSV-Spike pseudoviruses, and Kim
Peifley (NCI) for assistance in confocal immunofluorescence microscopy. Work in the
laboratory of AAC was funded by the Intramural Research Program, NIH, NCI, Center for
Cancer Research, and an Intramural Targeted Anti-COVID-19 award from NIAID. Work in the
laboratory of JWY was funded by the Division of Intramural Research, NIH, NIAID.
830

831 **References**
832
1   Letko, M., Marzi, A. & Munster, V. Functional assessment of cell entry and receptor
usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nature Microbiology* **11**,
1-17 (2020). https://doi.org:10.1038/s41564-020-0688-y
2   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
Blocked   by   a   Clinically   Proven   Protease   Inhibitor.   *Cell*,   1-19   (2020).
https://doi.org:10.1016/j.cell.2020.02.052
3   Chu, H. *et al.* Host and viral determinants for efficient SARS-CoV-2 infection of the
human lung. *Nat Commun* **12**, 134 (2021). https://doi.org:10.1038/s41467-020-20457-w
4   Clausen, T. M. *et al.* SARS-CoV-2 Infection Depends on Cellular Heparan Sulfate and
ACE2. *Cell* **183**, 1043-1057 e1015 (2020). https://doi.org:10.1016/j.cell.2020.09.033
5   Carlos, A. J. *et al.* The chaperone GRP78 is a host auxiliary factor for SARS-CoV-2 and
GRP78 depleting antibody blocks viral entry and infection. *J Biol Chem* **296**, 100759
(2021). https://doi.org:10.1016/j.jbc.2021.100759
6   Cantuti-Castelvetri, L. *et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry and
infectivity. *Science* **370**, 856-+ (2020). https://doi.org:10.1126/science.abd2985
7   Daly, J. L. *et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection. *Science* **370**,
861-865 (2020). https://doi.org:10.1126/science.abd3072
8   Wei, C. *et al.* HDL-scavenger receptor B type 1 facilitates SARS-CoV-2 entry. *Nat*
Metab **2**, 1391-1400 (2020). https://doi.org:10.1038/s42255-020-00324-0
9   Nguyen, L. *et al.* Sialic acid-containing glycolipids mediate binding and viral entry of
SARS-CoV-2. *Nat Chem Biol* **18**, 81-90 (2022). https://doi.org:10.1038/s41589-021-
00924-1
10   Koch, J., Uckeley, Z. M., Doldan, P., Stanifer, M., Boulant, S. & Lozach, P. Y.
TMPRSS2 expression dictates the entry route used by SARS-CoV-2 to infect host cells.
EMBO J **40**, e107821 (2021). https://doi.org:10.15252/embj.2021107821
11   Chan, J. F. *et al.* Altered host protease determinants for SARS-CoV-2 Omicron. *Sci Adv*
9, eadd3867 (2023). https://doi.org:10.1126/sciadv.add3867

12 Zang, R. *et al.* TMPRSS2 and TMPRSS4 promote SARS-CoV-2 infection of human
small intestinal enterocytes. *Sci Immunol* **5** (2020).
https://doi.org:10.1126/sciimmunol.abc3582
13 Laporte, M. *et al.* The SARS-CoV-2 and other human coronavirus spike proteins are fine-
tuned towards temperature and proteases of the human airways. *PLoS Pathog* **17**,
e1009500 (2021). https://doi.org:10.1371/journal.ppat.1009500
14 Kishimoto, M. *et al.* TMPRSS11D and TMPRSS13 Activate the SARS-CoV-2 Spike
Protein. *Viruses* **13** (2021). https://doi.org:10.3390/v13030384
15 V'Kovski, P., Kratzel, A., Steiner, S., Stalder, H. & Thiel, V. Coronavirus biology and
replication: implications for SARS-CoV-2. *Nat Rev Microbiol* **19**, 155-170 (2021).
https://doi.org:10.1038/s41579-020-00468-6
16 Karim, S. S. A. & Karim, Q. A. Omicron SARS-CoV-2 variant: a new chapter in the
COVID-19 pandemic. *Lancet* **398**, 2126-2128 (2021). https://doi.org:10.1016/S0140-
6736(21)02758-6
17 Pulliam, J. R. C. *et al.* Increased risk of SARS-CoV-2 reinfection associated with
emergence of Omicron in South Africa. *Science* **376**, eabn4947 (2022).
https://doi.org:10.1126/science.abn4947
18 Dejnirattisai, W. *et al.* SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape from
neutralizing antibody responses. *Cell* **185**, 467-484 e415 (2022).
https://doi.org:10.1016/j.cell.2021.12.046
19 Tuekprakhon, A. *et al.* Antibody escape of SARS-CoV-2 Omicron BA.4 and BA.5 from
vaccine and BA.1 serum. *Cell* **185**, 2422-2433 e2413 (2022).
https://doi.org:10.1016/j.cell.2022.06.005
20 Nutalai, R. *et al.* Potent cross-reactive antibodies following Omicron breakthrough in
vaccinees. *Cell* **185**, 2116-2131 e2118 (2022). https://doi.org:10.1016/j.cell.2022.05.014
21 VanBlargan, L. A. *et al.* An infectious SARS-CoV-2 B.1.1.529 Omicron virus escapes
neutralization by therapeutic monoclonal antibodies. *Nat Med* **28**, 490-495 (2022).
https://doi.org:10.1038/s41591-021-01678-y
22 Planas, D. *et al.* Considerable escape of SARS-CoV-2 Omicron to antibody
neutralization. *Nature* **602**, 671-675 (2022). https://doi.org:10.1038/s41586-021-04389-z
23 Wang, Q. *et al.* Antibody evasion by SARS-CoV-2 Omicron subvariants BA.2.12.1,
BA.4 and BA.5. *Nature* **608**, 603-608 (2022). https://doi.org:10.1038/s41586-022-05053-
w
24 Lubin, J. H. *et al.* Structural models of SARS-CoV-2 Omicron variant in complex with
ACE2 receptor or antibodies suggest altered binding interfaces. *bioRxiv* (2021).
https://doi.org:10.1101/2021.12.12.472313
25 Huo, J. *et al.* A delicate balance between antibody evasion and ACE2 affinity for
Omicron BA.2.75. *Cell Rep* **42**, 111903 (2023).
https://doi.org:10.1016/j.celrep.2022.111903
26 Jiang, X. L. *et al.* Omicron BQ.1 and BQ.1.1 escape neutralisation by omicron subvariant
breakthrough infection. *Lancet Infect Dis* **23**, 28-30 (2023).
https://doi.org:10.1016/S1473-3099(22)00805-2
27 Cao, Y. *et al.* BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron
infection. *Nature* **608**, 593-602 (2022). https://doi.org:10.1038/s41586-022-04980-y
28 Hong, Q. *et al.* Molecular basis of receptor binding and antibody neutralization of
Omicron. *Nature* **604**, 546-552 (2022). https://doi.org:10.1038/s41586-022-04581-9

29 Han, P. *et al.* Receptor binding and complex structures of human ACE2 to spike RBD
from omicron and delta SARS-CoV-2. *Cell* **185**, 630-640 e610 (2022).
https://doi.org:10.1016/j.cell.2022.01.001
30 Peacock, T. P. *et al.* The altered entry pathway and antigenic distance of the SARS-CoV-
2 Omicron variant map to separate domains of spike protein. *bioRxiv*,
2021.2012.2031.474653 (2022). https://doi.org:10.1101/2021.12.31.474653
31 Willett, B. J. *et al.* SARS-CoV-2 Omicron is an immune escape variant with an altered
cell entry pathway. *Nat Microbiol* **7**, 1161-1179 (2022). https://doi.org:10.1038/s41564-
022-01143-7
32 Meng, B. *et al.* Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts infectivity
and fusogenicity. *Nature* **603**, 706-714 (2022). https://doi.org:10.1038/s41586-022-
04474-x
33 Hui, K. P. Y. *et al.* SARS-CoV-2 Omicron variant replication in human bronchus and
lung ex vivo. *Nature* **603**, 715-720 (2022). https://doi.org:10.1038/s41586-022-04479-6
34 Qu, P. *et al.* Determinants and Mechanisms of the Low Fusogenicity and High
Dependence on Endosomal Entry of Omicron Subvariants. *mBio* **14**, e0317622 (2023).
https://doi.org:10.1128/mbio.03176-22
35 Yuan, S. *et al.* Pathogenicity, transmissibility, and fitness of SARS-CoV-2 Omicron in
Syrian hamsters. *Science* **377**, 428-433 (2022). https://doi.org:10.1126/science.abn8939
36 Suzuki, R. *et al.* Attenuated fusogenicity and pathogenicity of SARS-CoV-2 Omicron
variant. *Nature* **603**, 700-705 (2022). https://doi.org:10.1038/s41586-022-04462-1
37 Zhao, H. *et al.* SARS-CoV-2 Omicron variant shows less efficient replication and fusion
activity when compared with Delta variant in TMPRSS2-expressed cells. *Emerg*
Microbes Infect **11**, 277-283 (2022). https://doi.org:10.1080/22221751.2021.2023329
38 Halfmann, P. J. *et al.* SARS-CoV-2 Omicron virus causes attenuated disease in mice and
hamsters. *Nature* **603**, 687-692 (2022). https://doi.org:10.1038/s41586-022-04441-6
39 Kozlov, M. Omicron's feeble attack on the lungs could make it less dangerous. *Nature*
601, 177 (2022). https://doi.org:10.1038/d41586-022-00007-8
40 Maslo, C., Friedland, R., Toubkin, M., Laubscher, A., Akaloo, T. & Kama, B.
Characteristics and Outcomes of Hospitalized Patients in South Africa During the
COVID-19 Omicron Wave Compared With Previous Waves. *JAMA* **327**, 583-584
(2022). https://doi.org:10.1001/jama.2021.24868
41 Sposito, B. *et al.* The interferon landscape along the respiratory tract impacts the severity
of COVID-19. *Cell* **184**, 4953-4968 e4916 (2021).
https://doi.org:10.1016/j.cell.2021.08.016
42 Reggio, A. *et al.* Role of FAM134 paralogues in endoplasmic reticulum remodeling, ER-
phagy, and Collagen quality control. *EMBO reports* **22**, e52289 (2021).
https://doi.org:10.15252/embr.202052289
43 Wu, C. T. *et al.* SARS-CoV-2 replication in airway epithelia requires motile cilia and
microvillar reprogramming. *Cell* **186**, 112-130 e120 (2023).
https://doi.org:10.1016/j.cell.2022.11.030
44 Bojkova, D., Widera, M., Ciesek, S., Wass, M. N., Michaelis, M. & Cinatl, J., Jr.
Reduced interferon antagonism but similar drug sensitivity in Omicron variant compared
to Delta variant of SARS-CoV-2 isolates. *Cell Res* **32**, 319-321 (2022).
https://doi.org:10.1038/s41422-022-00619-9

45 Jocher, G. *et al.* ADAM10 and ADAM17 promote SARS-CoV-2 cell entry and spike
protein-mediated lung cell fusion. *EMBO Rep* **23**, e54305 (2022).
https://doi.org:10.15252/embr.202154305
46 Guo, K. *et al.* Interferon resistance of emerging SARS-CoV-2 variants. *Proc Natl Acad*
Sci U S A **119**, e2203760119 (2022). https://doi.org:10.1073/pnas.2203760119
47 Majdoul, S. & Compton, A. A. Lessons in self-defence: inhibition of virus entry by
intrinsic immunity. *Nature Reviews Immunology* (2021). https://doi.org:10.1038/s41577-
021-00626-8
48 Shi, G., Schwartz, O. & Compton, A. A. More than meets the I: the diverse antiviral and
cellular functions of interferon-induced transmembrane proteins. *Retrovirology* **14**, 1-11
(2017). https://doi.org:10.1186/s12977-017-0377-y
49 Kenney, A. D. *et al.* Interferon-induced transmembrane protein 3 (IFITM3) limits
lethality of SARS-CoV-2 in mice. *EMBO Rep* **24**, e56660 (2023).
https://doi.org:10.15252/embr.202256660
50 Shi, G. *et al.* Rapalogs downmodulate intrinsic immunity and promote cell entry of
SARS-CoV-2. *J Clin Invest* **132** (2022). https://doi.org:10.1172/JCI160766
51 Shi, G. *et al.* Opposing activities of IFITM proteins in SARS☐CoV☐2 infection. *The*
EMBO journal **3**, e201900542-201900512 (2020).
https://doi.org:10.15252/embj.2020106501
52 Winstone, H. *et al.* The polybasic cleavage site in the SARS-CoV-2 spike modulates viral
sensitivity to Type I interferon and IFITM2. *Journal of Virology* (2021).
https://doi.org:10.1128/JVI.02422-20
53 Prelli Bozzo, C. *et al.* IFITM proteins promote SARS-CoV-2 infection and are targets for
virus inhibition in vitro. *Nat Commun* **12**, 4584 (2021). https://doi.org:10.1038/s41467-
021-24817-y
54 Lin, T.-Y. *et al.* Amphotericin B Increases Influenza A Virus Infection by Preventing
IFITM3-Mediated Restriction. *Cell Reports* **5**, 895-908 (2013).
https://doi.org:10.1016/j.celrep.2013.10.033
55 Wrensch, F. *et al.* Interferon-Induced Transmembrane Protein-Mediated Inhibition of
Host Cell Entry of Ebolaviruses. *The Journal of Infectious Diseases*, jiv255 (2015).
https://doi.org:10.1093/infdis/jiv255
56 Suddala, K. C. *et al.* Interferon-induced transmembrane protein 3 blocks fusion of
sensitive but not resistant viruses by partitioning into virus-carrying endosomes. *PLoS*
Pathogens **15**, e1007532-1007535 (2019). https://doi.org:10.1371/journal.ppat.1007532
57 Rahman, K., Coomer, C. A., Majdoul, S., Ding, S. Y., Padilla-Parra, S. & Compton, A.
A. Homology-guided identification of a conserved motif linking the antiviral functions of
IFITM3 to its oligomeric state. *eLife* **9**, 975-925 (2020).
https://doi.org:10.7554/eLife.58537
58 Huotari, J. & Helenius, A. Endosome maturation. *EMBO J* **30**, 3481-3500 (2011).
https://doi.org:10.1038/emboj.2011.286
59 Nie, C., Sahoo, A. K., Netz, R. R., Herrmann, A., Ballauff, M. & Haag, R. Charge
Matters: Mutations in Omicron Variant Favor Binding to Cells. *Chembiochem* **23**,
e202100681 (2022). https://doi.org:10.1002/cbic.202100681
60 Shi, D. *et al.* Structural Characteristics of Heparin Binding to SARS-CoV-2 Spike Protein
RBD of Omicron Sub-Lineages BA.2.12.1, BA.4 and BA.5. *Viruses* **14** (2022).
https://doi.org:10.3390/v14122696

61 Gelbach, A. L. *et al.* Interactions between heparin and SARS-CoV-2 spike glycoprotein
RBD from omicron and other variants. *Front Mol Biosci* **9**, 912887 (2022).
https://doi.org:10.3389/fmolb.2022.912887
62 Baindara, P., Roy, D., Mandal, S. M. & Schrum, A. G. Conservation and Enhanced
Binding of SARS-CoV-2 Omicron Spike Protein to Coreceptor Neuropilin-1 Predicted by
Docking Analysis. *Infect Dis Rep* **14**, 243-249 (2022).
https://doi.org:10.3390/idr14020029
63 Beyer, D. K. & Forero, A. Mechanisms of Antiviral Immune Evasion of SARS-CoV-2. *J
Mol Biol* **434**, 167265 (2022). https://doi.org:10.1016/j.jmb.2021.167265
64 Lei, X. *et al.* Activation and evasion of type I interferon responses by SARS-CoV-2. *Nat
Commun* **11**, 3810 (2020). https://doi.org:10.1038/s41467-020-17665-9
65 Xia, H. *et al.* Evasion of Type I Interferon by SARS-CoV-2. *Cell Rep* **33**, 108234 (2020).
https://doi.org:10.1016/j.celrep.2020.108234
66 Shalamova, L. *et al.* Omicron variant of SARS-CoV-2 exhibits an increased resilience to
the antiviral type I interferon response. *PNAS Nexus* **1**, pgac067 (2022).
https://doi.org:10.1093/pnasnexus/pgac067
67 Bojkova, D. *et al.* Omicron-induced interferon signalling prevents influenza A virus
infection. *bioRxiv*, 2022.2009.2006.506799 (2022).
https://doi.org:10.1101/2022.09.06.506799
68 Singh, J. *et al.* BA.1, BA.2 and BA.2.75 variants show comparable replication kinetics,
reduced impact on epithelial barrier and elicit cross-neutralizing antibodies. *PLoS Pathog*
19, e1011196 (2023). https://doi.org:10.1371/journal.ppat.1011196
69 Alfi, O. *et al.* SARS-CoV-2 Omicron Induces Enhanced Mucosal Interferon Response
Compared to other Variants of Concern, Associated with Restricted Replication in
Human Lung Tissues. *Viruses* **14** (2022). https://doi.org:10.3390/v14071583
70 Prokunina-Olsson, L. *et al.* COVID-19 and emerging viral infections: The case for
interferon lambda. *J Exp Med* **217** (2020). https://doi.org:10.1084/jem.20200653
71 Chong, Z. *et al.* Nasally delivered interferon-lambda protects mice against infection by
SARS-CoV-2 variants including Omicron. *Cell Rep* **39**, 110799 (2022).
https://doi.org:10.1016/j.celrep.2022.110799
72 Jagannathan, P. *et al.* Peginterferon Lambda-1a for treatment of outpatients with
uncomplicated COVID-19: a randomized placebo-controlled trial. *Nat Commun* **12**, 1967
(2021). https://doi.org:10.1038/s41467-021-22177-1
73 Feld, J. J. *et al.* Peginterferon lambda for the treatment of outpatients with COVID-19: a
phase 2, placebo-controlled randomised trial. *Lancet Respir Med* **9**, 498-510 (2021).
https://doi.org:10.1016/S2213-2600(20)30566-X
74 Reis, G. *et al.* Early Treatment with Pegylated Interferon Lambda for Covid-19. *N Engl J
Med* **388**, 518-528 (2023). https://doi.org:10.1056/NEJMoa2209760
75 Ye, C. *et al.* Rescue of SARS-CoV-2 from a Single Bacterial Artificial Chromosome.
mBio **11** (2020). https://doi.org:10.1128/mBio.02168-20
76 Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C(T)
method. *Nat Protoc* **3**, 1101-1108 (2008). https://doi.org:10.1038/nprot.2008.73
1039