1
12 Temporal Dynamics of Brain Activity Predicting Illusory Agency over Involuntary Movements
13
14
15
16

17 John P. Veillette[1†], Pedro Lopes[2], Howard C. Nusbaum[1]
18
19 [1]Department of Psychology, University of Chicago
20 [2]Department of Computer Science, University of Chicago
21
22 [†]Correspondence should be addressed to John P. Veillette; E-mail: johnv@uchicago.edu

Author Contributions (CRediT Taxonomy)

J.P.V.: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Software, Validation, Visualization, Writing - original draft, and Writing - review & editing. **P.L.**: Conceptualization, Funding acquisition, Methodology, Resources, Software, and Writing - review & editing. **H.C.N.**: Conceptualization, Funding acquisition, Resources, Supervision, and Writing - review & editing.

Acknowledgements

P.L. and H.C.N. were supported by NSF NCS 2024923, and J.P.V. was supported by NSF GRFP DGE 1746045 and a Neubauer Family Distinguished Doctoral Fellowship.

Conflict of Interest Statement

The authors declare no competing interests.

47   Abstract
48
49
Our muscles are the primary means through which we affect the external world, and the sense of
agency (SoA) over the action through those muscles is fundamental to our self-awareness.
However, SoA research to date has focused almost exclusively on agency over action outcomes
rather than over the musculature itself, as it was believed that SoA over the musculature could not
be manipulated directly. Drawing on methods from human-computer interaction and adaptive
experimentation, we use human-in-the-loop Bayesian optimization to tune the timing of electrical
muscle stimulation so as to robustly elicit an illusion of agency over electrically-actuated muscle
movements. We use time-resolved decoding of subjects' EEG to estimate the time course of neural
activity which predicts illusory agency on a trial-by-trial basis. Like paradigms which assess SoA
over action consequences, we found that the late (post-conscious) neural activity predicts SoA.
Unlike typical paradigms, however, we also find patterns of early (sensorimotor) activity with
distinct temporal dynamics predicts agency over muscle movements, suggesting that the "neural
correlates of agency" may depend on the level of abstraction (i.e., direct sensorimotor feedback
vs. downstream consequences) most relevant to a given agency judgement. Moreover, fractal
analysis of the EEG suggests that SoA-contingent dynamics of neural activity may modulate the
sensitivity of the motor system to external input.
66
67
68

1. Introduction

Voluntary movements tend to be accompanied by a subjective experience of volition – that is, the feeling that "I did that." This feeling is referred to as the sense of agency (SoA), which is considered one of the most basic building blocks of conscious selfhood (Gallagher, 2000; Haggard, 2008). Pathologies affecting SoA – including schizophrenia (C. Frith, 2012), alien hand syndrome (Panikkath et al., 2014), perceived (non-)control of a phantom limb (Ramachandran & Hirstein, 1998), automatic "utilization behavior" (Lhermitte et al., 1986), and learned paralysis (Wolf et al., 1989) – are often characterized by anomalies in the experience of control over the body itself (i.e., the musculature) rather than over external action outcomes per se.

SoA research in healthy populations, however, has focused primarily on external sensory consequences of action (Haggard, 2008, 2017). Indeed, recent discussions have noted the lack of experimental paradigms for investigating SoA over movements – "narrow" SoA over the musculature as opposed to "broad" SoA encompassing action outcomes (Christensen & Grünbaum, 2018). Indeed, the field tends to generalize too readily to other classes of agency judgments that were observed in specific settings, such as from subjects experiencing an auditory feedback (that could be delayed or manipulated) following a button press. As such, the literature tends to treat SoA as a homogenous phenomenon always accompanied by the same neural correlates. However, an alternative hypothesis is that the neural correlates of SoA may vary as a *function of the modality* (e.g., proprioceptive vs. auditory) or of the *level of abstraction appropriate* for a given agency judgement (Charalampaki et al., 2022). In particular, it has been noted recently that current models may not generalize to SoA over the musculature (Christensen & Grünbaum, 2018) or over one's stream of thoughts (C. Frith, 2012). This potential discrepancy bears on a fundamental scientific question of whether the mechanisms that give rise to the experience of an agentic self are common across scales of biological organization – and if not, how and why do we assign agency to the same unified self at these different scales?

One reason for the lack of paradigms assessing SoA over the musculature is that control over one's own muscles is normally unambiguous. Indeed, previous attempts to elicit SoA for experimenter-evoked (e.g., by TMS) movements have not succeeded (Christensen & Grünbaum, 2018; Haggard & Clark, 2003), leading to the conclusion that "such involuntary movements are never accompanied by a sense of agency" (Haggard, 2017). However, since cognitive scientists favor indirect measures of agency such as intentional binding (i.e., the perceived delay) between an action (button press) and outcome (tone), these findings still primarily reflect broad SoA over external outcomes rather than narrow SoA over the muscles (Haggard et al., 2002). Meanwhile, human-computer interaction researchers have recently begun investigating SoA in interfaces that use electrical-muscle stimulation (EMS) to drive users' muscles. They find, in contrast, that subjects self-report EMS-caused movements (that is, the movements themselves) as self-caused, so long as stimulation temporally aligns with users' endogenous intention to move (Kasahara et al., 2019, 2021; Tajima et al., 2022). The cognitive neuroscience literature has yet to fully embrace these findings, in part because they have yet to be verified using an implicit measure of agency, such as intentional binding or sensory attenuation (Dewey & Knoblich, 2014).

Thus, in the present work, we "preempt" subjects' endogenous muscle movements with EMS during a cue-response reaction time task, using Bayesian optimization to guide stimulation timing so as to control the proportion of EMS-caused movements perceived as self-caused. Using time-resolved multivariate pattern analysis ("neural decoding") of subjects' trial-by-trial EEG, we show that the neural response to EMS predicts subjects' agency judgement about the resulting

muscle movement as early as 83 ms following stimulation onset, showing that subjects' self-report
has a basis in early, *pre-conscious sensorimotor processing* – not just a response bias. Notably,
this finding differs from those obtained using typical button-tone paradigm, in which early evoked
responses (to tones) have repeatedly failed to predict subjective agency judgments (Kühn et al.,
2011; Timm et al., 2016). This early activity shows distinct temporal dynamics from the neural
patterns that predict SoA later in the epoch. Finally, an exploratory analysis shows that the fractal
dimension of the EEG also predicts SoA, suggesting that the complexity of sensory processing
may differ for sensations perceived as movement feedback.

122
2. Materials and Methods

2.1 Participants and Ethics Statement

25 University of Chicago undergraduate students participated in the study; however, two
subjects were subsequently excluded for noncompliance with task instructions (i.e., one admitted
to letting the electrical stimulator perform the task without attempting a volitional button press,
and another pressed the button continually to speed through the task instead of when cued).
Participants were recruited through the University of Chicago's human subject recruitment system,
SONA Systems. All subjects gave written, informed consent before participating. All of the
methods performed in the study were in accordance with relevant safety and ethics guidelines for
human subject research and were approved by the Social and Behavioral Sciences Institutional
Review Board at the University of Chicago (IRB19-1302). This study was not a clinical trial.

2.2 Experimental Design

Subjects completed three blocks of trials. In the first and final blocks (30 trials each), each
trial was as follows: (a) a fixation cross was displayed on the screen; (b) after a random wait period
(2-4 seconds, uniformly distributed), a cue to press the response button as quickly as possible was
displayed on the screen ("Go!"); after which, (c) the subject pressed a button (on a *Cedrus RB-620*
button box; California, United States) voluntarily and without any electrical stimulation; finally,
(d) the trial ended with another fixation cross displayed on the screen.

In the middle (EMS) block (300 trials), (a), (b), (c) and (d) from the above remained the
same, but this time, electrical muscle stimulation (EMS) was applied to the forearm in order to
move the subjects' finger (see *2.4* for stimulation details), causing their finger to press the button
involuntarily after the cue to move. A typical trial in the stimulation block is shown in Figure 1.

The latency of the EMS relative to the cue to move (b) was guided by a Bayesian
Optimization procedure (see 2.3 below) to be as near as possible to the latency at which subjects
will experience 50% of movements to be self-caused, which prior studies have shown is
substantially before the subjects' average reaction time (Kasahara et al., 2019, 2021). (However,
the Bayesian optimization procedure only considered latencies between 50 and 600 ms, so
stimulation did always occur in that interval.) As a result, trials in which the subject pressed the
button before stimulation was delivered were rare (see *Results 3.1*). If the subject pressed the
button before stimulation was delivered, stimulation occurred immediately upon the button press
so that the onset of the electrical stimulation, which causes a perceptible though painless tingling
sensation on the skin, was always temporally confusable with that of the movement.

As depicted in Figure 1, after each button press in the stimulation block, the subject was
asked to judge whether they caused the button press or if the electrical stimulation caused the
button press. This response was fed into the Bayesian Optimization algorithm to inform the timing
of stimulation in subsequent trials (see below). The experiment interface was implemented using
PsychoPy (Peirce, 2007).

Figure 1: Graphical depiction of a typical trial in the stimulation block. Stimulation is hard-coded to occur before
600 ms after the cue to move, but a Bayesian Optimization procedure biases it to occur at the latency the subject is
estimated to perceive agency over the resulting muscle movement with 50% probability. In practice, this almost always
occurs before the subject is able to move on their own.

2.3 Bayesian Optimization

We adaptively fit a Bayesian logistic regression, updated after each trial, to estimate the
stimulation latency (relative to the movement cue) at which the subject would judge that they
caused the muscle movement on that trial with 50% probability. We parametrized the logistic
regression as in Formula (1) so that the parameter α would correspond to that 50% threshold.

170
$$\begin{aligned} \text{logit}(p) &= \beta\,(x-\alpha) \\ y &\sim \text{Bernoulli}(p) \end{aligned} \qquad (1)$$

Using the 50 reaction time trials from the first block of the experiment to compute each subject's
mean reaction time μ_{RT} with standard deviation σ_{RT}, we began the stimulation block with
informative priors on α and β to start our model in a good position to learn the true threshold
quickly. For α, we used the log-normal distribution with mean $\mu_{\text{RT}} - 40$ ms and standard deviation
σ_{RT}, since the "preemptive gain" estimated in previous studies is roughly 40 ms (Kasahara et al.,
2021). For β, the slope parameter, we used the log-normal distribution with mean 0.017 and
standard deviation 0.005, which generously encompasses the slopes in the subject-specific logistic
regressions reported by a previous study (Kasahara et al., 2019).

On each trial, the stimulation latency is chosen by taking a draw from the current posterior
distribution for α, truncated between 50 and 600 ms after the movement cue. After each trial, the
subject judges whether they caused the muscle movement, and their response is used to update the
posterior, which is approximated via variational inference. (However, if the subject pressed the
button before stimulation was delivered, and thus actually caused the movement, the trial was

counted as being judged self-caused regardless of how the subject responded.) In this way, as the
posterior becomes more precise (i.e., we are more certain about the true threshold), the stimulation
times will hone in on that threshold. But while we are still uncertain, the posterior we are sampling
from is wide, and we will explore a wider range of latencies. Thus, this algorithm navigates the
exploration-exploitation tradeoff naturally; it is a modified version of Thompson sampling, a
Bayesian Optimization approach which benefits from strong regret-minimization guarantees
(Agrawal & Goyal, 2012).

This procedure was implemented in *Python* using the probabilistic programming language
Pyro for variational inference (Bingham et al., 2019), and all associated code is available (see *Data*
and Code Availability).

2.4 Electrical Muscle Stimulation

Before the experiment began, two EMS electrodes were applied to the skin above the *flexor*
digitorum profundus muscle on the right (dominant) forearm, which is an easily accessible muscle
that moves the ring finger, which subjects used to press the button during the experiment.

Stimulation was performed with a *RehaStim 1* device (*HASOMED* GmbH, Mageburg,
Germany). On each trial, muscle actuation consisted of a single, biphasic pulse of constant current
stimulation lasting 900 microseconds (400 μs positive, 400 μs negative, separated by 100 μs),
since biphasic pulses result in less perceptible tingling of the skin than monophasic pulses.

Before beginning the experiment, we calibrated the stimulation amplitude to the minimum
intensity required to reliably move the subject's finger. The calibration procedure was as follows:
(a) The subject placed their ring finger on the button that would be used during the experiment and
was instructed not to move their hand. (b) Starting at an intensity of 1 mA, we stimulated the
subjects' arm 10 times. If fewer than 10 button presses were registered, then we iterated the
intensity by 1 mA and repeated. (c) We stopped increasing the intensity upon achieving 10
consecutive actuated button presses, or if a conservative safety limit of 25 mA was reached. (In
practice, electrical stimulation intensities can safely go much higher, but might eventually become
uncomfortable, which we avoided using this conservative lower bound.) (d) If stopping in (c) was
due to reaching the safety limit, the electrode positions were adjusted, and the procedure was
repeated until stimulation elicited button presses reliably.

2.6 Manipulation Check and Outlier Removal

We performed a brief analysis to demonstrate the effectiveness of our Bayesian optimization
procedure at preempting subjects' movements.

First, we identified trials in which stimulation was ineffective at eliciting a muscle
movement, which were excluded from subsequent behavioral analysis, as well as from EEG
analysis. To identify such trials, we (a) removed trials in which subjects pressed the button prior
to stimulation, since such movements were obviously caused by the subject, (b) removed trials in
which the recorded response time was greater than 600 ms, outside the stimulation time window,
then (c) fit a log-normal distribution to the lag between stimulation and the time of the subsequent
button press and removed the trials in the most extreme 5% of that fit distribution. Briefly, (a) and
(b) removed an average of 18.4 trials and (c) removed an average of 12.4 trials per subject. The
result of outlier removal is visualized, for convenience, in Figure 3. After this step, measured
"reaction" time in the stimulation block should be a nearly linear function of stimulation latency.

We then fit a Bayesian multilevel model to the trial-by-trial reaction times in each
experiment block using the *Bambi* package (Capretto et al., 2022) as an interface to (Patil et al.,
2010) using the default (weakly informative) priors, which are conservative in the sense that they
should shrink estimates of the RT in each condition toward each other (Westfall, 2017). This is
done in the interest of replicating the finding that EMS-induced movements can substantially
preempt subjects' natural movements in this paradigm, in which case the measured "reaction" time
should be lowest in the stimulation block (Kasahara et al., 2019, 2021). We report the empirical
percentage of stimulation trials that subjects perceived as self-caused, which should be close to
50% if our Bayesian Optimization procedure was successful.

2.6 EEG Acquisition

EEG was recorded with 64 active Ag/AgCl electrodes (*actiCHamp*, Brain Products, Munich,
Germany) placed according to the extended 10-20 system. At the time of recording, the electrodes
were referenced to Cz and sampled at 10,000 Hz. Two of the 64 electrodes (which would have
been AF7 and AF8 on the typical *actiCAP* layout) were dropped below the left and right eyes so
that they could later be re-referenced to become EOG channels.

Experiment events were marked in the EEG recording using TTL triggers, delivered using
an *RTBox* (Li et al., 2010) in lieu of a built-in parallel port on the experiment control computer.

We additionally placed a photo-sensor (*Brain Products*, Munich, Germany) on the
subjects' screen to record screen brightness at the same sampling rate as the EEG. The timestamps
of the TTL triggers corresponding to the onset of the visual cue to move (see section 2.2) was
corrected after the recording using the ground-truth event time measured using the photo sensor.
The difference between the TTL trigger, which was sent in parallel with the start of the reaction
time trial, and the actual time of the screen flip recorded with the photo sensor was used to correct
reaction time measurements taken with the response box as well.

Before beginning the experiment, we tested the hardware latency between the EMS pulses
and the TTL pulses marking them in the EEG recording; to this end, we used a custom cable that
allowed us to record the output voltage of the stimulation device directly with the EEG amplifier.
Over 500 repetitions, we measured a latency of mean 1.23 ms with a jitter of standard deviation
0.04 ms. This constant 1.23 ms delay in the event timestamps was corrected offline after the EEG
recordings. All timestamps found in the open dataset associated with this article (see ***2.9 Data and***
Code Availability) have already been corrected for convenience.

Additionally, the precise subject-specific positions of the 62 head electrodes were
measured at the end of each recording using a *CapTrak* (*Brain Products*, Munich, Germany).

2.6 EEG Preprocessing

EEG was preprocessed in *Python* using *MNE-Python* package (Gramfort et al., 2014). First, the
electrodes below the left and right eye, which were referenced to Cz at the time of recording, were
re-referenced against forehead electrodes Fp2 and Fp1, respectively, to become bipolar EOG
channels montaged to capture both horizontal and vertical eye movements.

Subsequently, the electrical artifact from the EMS needed to be removed. The EMS pulse
was brief (less than a millisecond), so it could be removed by linearly interpolating over the
interval starting 5 ms before and ending 10 ms after the event timestamp. However, since by the
end of the stimulation block, stimulation was delivered at a narrow distribution of latencies after
the cue to move, which occurs at the 60 Hz screen refresh rate of the subjects' monitor, stimulation

did not occur at a random phase of the background 60 Hz line noise. Thus, we found data quality
was improved by removing line noise prior to interpolating over the stimulation artifact. However,
applying an FIR notch filter before removing the pulse-like (flat power spectrum) stimulus would
result in distortion of the filtered signal. To this end, we fit a multi-taper estimation of the
sinusoidal components at the line noise frequency and its first two harmonics and removed those
components (as implemented in MNE). Then, after performing linear interpolation, we applied an
FIR notch filter at 60 Hz and its harmonics up to the upper edge of the bandpass filter that would
be applied subsequently (see below) to thoroughly clean the data of line noise, and then resampled
the data to 5,000 Hz to improve the speed of computation for subsequent preprocessing steps.

Next, we applied common preprocessing operations in adherence with the standardized
PREP preprocessing pipeline for EEG data (Bigdely-Shamlo et al., 2015) using the
implementation in the *PyPREP* package (Appelhoff et al., 2022). This pipeline robustly re-
references the EEG signal to the average of all electrodes and interpolates electrodes it determines
have poor signal quality; since data quality is sensitive to the order in which these operations are
performed (i.e., bad channel identification depends on the reference, and the average reference
should not include bad channels), the pipeline applies the two steps iteratively until convergence.
Criteria for identifying bad channels includes extreme amplitudes, lack of correlation with other
channels, lack of predictability by neighboring channels, and unusual high frequency noise; see
the PREP paper for a full description (Bigdely-Shamlo et al., 2015). Precise, subject-specific
electrode positions were used for the interpolation of bad channels, and a record of which channels
were interpolated are available in subject-specific preprocessing reports (see 2.9).

Usually, line noise is also removed as part of the PREP pipeline, but since we removed line
noise when interpolating over the stimulation artifact, we did not perform this operation again
during PREP. (Note that removal of the stimulation artifact needed to be performed before PREP,
since PREP include a detrending step that uses a highpass filter, which would result in signal
distortion if the pulse-like stimulation artifact was still present.)

Next, we filtered the data to the frequency band used for analysis. We used a low cutoff of
1 Hz to remove low-frequency drift, and we used two separate high cutoffs, 30 Hz and 70 Hz, for
the two types of analysis to follow preprocessing (see 2.7 and 2.8 below). Thus, the following
preprocessing steps were applied twice, once for each copy of the data.

We then removed EOG contamination of the EEG signal. We decomposed to EEG data
into 15 independent components (ICs) using the *FastICA* algorithm implemented in MNE
(Hyvarinen, 1999). Then, we correlated each IC with the EOG channels, *z*-scored the correlation
coefficients, and deemed an IC to contain eye artifact if the absolute value of its *z*-score exceeded
1.96. Those ICs were zeroed out to remove them from the original data. Plots of the scalp
topographies of removed ICs for each subject can be found in their preprocessing reports (see 2.9).

Subsequently, we segmented the data into epochs starting 100 ms before the onset of
stimulation and ending 500 ms after stimulation. We then estimated the peak-to-peak rejection
threshold that would be used to identify trials containing unwanted artifacts using the *Autoreject*
package (Jas et al., 2017), which estimates the optimal threshold as that which minimizes the 5-
fold cross-validated root-mean-squared difference between the mean of the training folds and the
median of the testing fold, a robust proxy metric for signal-to-noise. The resulting per-subject
rejection thresholds are recorded in each subjects' preprocessing report (see 2.9).

Since the visual evoked response to the movement cue is unlikely to be over by the time of
stimulation, we attempted to remove the visual evoked response from our epoched data to
minimize confounds. To do so, we computed evoked responses to both the visual and electrical
stimuli simultaneously using a regression-based overlap correction on the continuous (non-
epoched) data, excluding second-long chunks of the data in which peak-to-peak amplitude exceeds
the rejection threshold (Smith & Kutas, 2015); conceptually, this is very similar to the way
generalized linear models (GLMs) are used to deconvolve hemodynamic responses in fMRI. Then,
the overlap-corrected visual evoked response was aligned with the epoched version of the data and
subtracted out. Thus, the average visual response to the movement cue was removed from the
stimulation-locked epochs. Subject-level evoked responses can be found in our open dataset and
are visualized in the subject-specific preprocessing reports (see 2.9).

Finally, the rejection threshold was applied to the cleaned and overlap-corrected epochs,
removing trials still contaminated by artifacts. The surviving epochs were down-sampled to twice
their high-cutoff frequency for computational expediency and saved for further analysis. This
epoched data is available in our open dataset, and subject-level trial yields are recorded in the
accompanying quality check reports (see 2.9).

2.7 EEG Decoding: Multivariate Pattern Analysis

For MVPA, we used the copy of the epoched data filtered from 1-30 Hz, since a conservative high-
cutoff smooths signals of interest, which may occur at slightly different times on a per-epoch or
per-subject basis, such that they can be aligned across subjects.

To assess the temporal dynamics of patterns which differentiate trial perceived as self- and
EMS-caused, we employ the temporal generalization method (King & Dehaene, 2014). In this
approach, a linear classifier is trained on the pattern of voltages at each timepoint (using a training
set), and then its classification performance is quantified at every other timepoint (using a test set).
In our case, we quantify classification performance using the area under the receiver-operator
curve (ROC-AUC), a non-parametric metric of class separation that is more robust than accuracy.
This results in a $n_{\text{times}} \times n_{\text{times}}$ matrix M, where M_{ij} is the performance of the classifier trained at
time i evaluated at time j. For our linear classifier, we use a logistic regression fit with generalized
estimating equations (GEE) as implemented in the *statsmodels* package (Liang & Zeger, 1986;
Seabold & Perktold, 2010). GEE is a semi-parametric fitting procedure for linear models that does
not assume the data comes from a certain distribution, resulting in parameter estimates that are
robust to heterogeneity; moreover, it allows the fitting of the marginal (i.e., population) parameters
of a multilevel model without explicitly computing the subject-level model, which allows us to
account for subject-level random effects without the computational burden of a multilevel model,
which is in general too high for MVPA.

Transient patterns, which predict agency but only appear for a short time, are characterized
by strong classification performance only on the diagonal of M; sustained patterns show strong
decoding performance off the diagonal as well, indicating that the same neural process persists
over time. This method has been argued to provide insight into the duration of and dynamic
transitions between mental representations as a cognitive process unfolds in time (King &
Dehaene, 2014).

We employ a 10-fold stratified cross-validation scheme where subjects are randomly split
into non-overlapping trials across folds (i.e., stratification entails an attempt to ensure an equal
proportion of agency and non-agency trials across folds). Thus, classifiers are only evaluated on

trials from subjects that did not appear in their training set, ensuring identified patterns generalize
across subjects. Specifically, we used this cross-validation scheme as implemented in the *scikit-*
learn package (Pedregosa et al., 2011). This 10-fold cross-validation scheme was repeated 10
times, with 10 different random splits into folds so that results are robust against random effects
of particular splits of the data. Then, ROC-AUC scores obtained across the 10-by-10 splits were
compared to the chance performance (ROC-AUC = 0.5) using a one-sided, one-sample *t*-test at
every (train-time, test-time) pair M_{ij}. However, the independence assumption of the standard *t*-test
are violated due to observations being repeated across training and test sets (Dietterich, 1998); to
account for this, we use the variance correction proved by Nadeau & Bengio to contain false
positive rates in this context when computing our *t*-statistics (Bouckaert & Frank, 2004; Nadeau
& Bengio, 1999).

Since the "shape" and onset of classification performances in the matrix *M* is of interest,
then cluster-level inference (e.g., on the size of contiguous subsets of M_{ij}), rather than inference
on the individual M_{ij}'s, is desirable in principle. However, the cluster-level tests used in
neuroimaging do not control the false-positive rate for claims about the extent of effects, such as
how far they extend off the diagonal, which are critical to the interpretation of the time-by-time
generalization results (Sassenhagen & Draschkow, 2019). Instead, we use All-Resolutions
Inference (Rosenblatt et al., 2018), as implemented in our *MNE-ARI* package (Veillette, 2022), to
compute simultaneous lower bounds on the true positive proportion in each cluster across all
$n_{\text{times}} \times n_{\text{times}}$ possible clustering thresholds. This approach to dealing with the multiple
comparisons problem adequately conveys uncertainty about the localization of true effects within
clusters, while controlling the false-positive rate in the strong sense. For instance, if the proportion
of true positives in a cluster is low, then it is unclear which M_{ij} in the cluster are the true positives;
conversely, if the proportion is high (e.g., >95%), then the localization is quite certain.

2.8 EEG Decoding: Fractal Analysis (Exploratory)

To compute fractal measures of the EEG data, we use the copy of the epoched data filtered between
1 Hz and 70 Hz, since fractal metrics are fundamentally about how the behavior of a time series
varies across timescales, and filtering out high frequencies removes more granular timescales.
Nonetheless, we still filter at 70 Hz because it is a topic of active debate whether high gamma band
(>70 Hz) can be reliably detected using scalp EEG (Darvas et al., 2010; Fries et al., 2008). To
improve spatial specificity of this analysis (since the time domain is no longer of interest given the
properties of these fractal measures), we computed the current source density transformation of
our scalp-recorded voltages to reduce the effect of volume conduction before computing the
following fractal metrics. Both simulations and empirical results indicate that this method is
effective at reducing the radius of cortical tissue to which each electrode's signal is sensitive to a
few centimeters (Kayser & Tenke, 2015).

The first metric of interest is the Higuchi fractal dimension, which is an efficiently
computable approximation of the fractal dimension (Higuchi, 1988). Generally speaking, this
quantity is a measure of the complexity or roughness of a time series (or of its underlying
dynamical attractor), which is directly related to the entropy of a signal (i.e., an upper bound on
the amount of information it can encode) but can be computed more reliably from small time series.
The fractal dimension of both the background EEG and the EEG response to perturbation is highly
predictive of states of consciousness (Kesić & Spasić, 2016; Ruiz de Miras et al., 2019), consistent
with information theoretic accounts of conscious awareness (Oizumi et al., 2014). Moreover, some

evidence suggests that neural activity corresponding to internally generated percepts (e.g., mind
wandering) is of a higher fractal dimension than that resulting from sensory stimulation, but it is
unclear whether this difference is accounted for by the different modalities (i.e., mind wandering
vs. external sensation) or to the self-attribution of the percept, which our analysis should clarify.

The second metric of interest is the Hurst exponent, which is a measure of long-range
temporal dependencies in a time series (Qian & Rasheed, 2004). While, in a self-affine (i.e.,
strongly self-similar) time series, the fractal dimension D and the Hurst exponent H are related by
the deterministic relationship $D + H = 2$, these quantities can vary independently in time series
that depart from strict self-similarity (Gneiting & Schlather, 2004), and these metrics have been
reported to diverge in EEG data (Martis et al., 2015). As such, fractal dimension is a local property
of the signal, whereas Hurst is a global property which quantifies how sensitive the larger time
series is to local changes (Gneiting & Schlather, 2004). In the cognitive neuroscience literature,
this quantity has been interpreted as corresponding to how responsive the underlying neural system
is to external input (Churchill et al., 2016; Kardan et al., 2020; Zhuang et al., 2022), and it is
notably suppressed in those suffering from psychiatric conditions associated with an impaired
sense of agency (Sokunbi et al., 2014; Stier et al., 2021).

It is important to note, as a caveat, that the Hurst exponent cannot be reliably estimated
from time-series as small as our single-trials. While Hurst is in principle a scale-invariant metric,
in practice, time series as long as 10,000 time points are required to produce reliable estimates
with conventional estimates (Eke et al., 2000, 2002; Oliver & Ballester, 1998), and while our time
series were in principle sampled at 10,000 Hz, filtering reduces the effective sampling rate to twice
the upper passband edge in practice. Despite the unreliability of the Hurst estimates at the trial
level, however, *within*-subject variation in trial-by-trial Hurst exponents estimated from EEG are
sensitive to cognitive functions (Kardan et al., 2020). In other words, while our individual trial-
by-trial Hurst estimates may not be reliable indices of the mathematically-defined Hurst exponent,
their variation may still be meaningful.

We estimate fractal dimension using Higuchi's algorithm (Higuchi, 1988) and the Hurst
exponent using rescaled range analysis (Qian & Rasheed, 2004), using the implementations
provided in the MNE-Features package (Schiratti et al., 2018). Then, for each electrode and
subject, we compute the ROC-AUC between each metric and subjects' self-reported agency. Since
these metrics are determined a priori and not from the data, no cross-validation scheme is
necessary; a one-sample, two-sided *t*-test is used compare subject-level decoding performance to
chance (ROC-AUC = 0.5).

2.9 Data and Code Availability

Code for running the experiment can be found on *GitHub* (github.com/apex-lab/agency-
experiment) and in a permanent archive on *Zenodo* (doi.org/10.5281/zenodo.7894011). Similarly,
all data analysis code, including EEG preprocessing code, can be found at github.com/apex-
lab/agency-analysis and https://doi.org/10.5281/zenodo.7894007. All data, including both raw
data, preprocessed derivatives, and post-preprocessing quality check reports for each subject, can
be found on *OpenNeuro* (doi.org/10.18112/openneuro.ds004561.v1.0.0).

3. Results

3.1 Bayesian optimization effectively controls the proportion of trials perceived as self-caused.

The Bayesian Optimization procedure resulted in trial-by-trial stimulation latencies honing in on
some threshold estimate throughout the stimulation block. A representative time course is shown
in Figure 2.

Figure 2: Trial-by-trial stimulation latency over the course of the stimulation block for a representative subject.
(a) Stimulation latency hones in on a stable value over time. (b) A logistic regression computed after the experiment
and shows that stimulation times are close to the retro-actively estimated 50% threshold, even though that threshold
was not known in advance. The subject featured here is "sub-07" in the associated dataset.

After removing trials in which stimulation failed to produce a muscle movement (and
therefore the "reaction" time was not a function of stimulation latency), our multilevel model of
the recorded reaction times estimated a 99.9% posterior probability that button presses occurred
earlier in the stimulation block than in either of the other blocks. In particular, we estimate that
"reaction" times resulting from EMS-actuated movements were between 17.5 ms and 65.0 ms
faster than true reaction times in the first (pre) block with 95% probability, and between 13.8 ms
and 43.9 ms faster than those in the final (post) block. A nominal speedup between the pre and
post blocks was observed with 90.6% probability (95% HDI: [-6.7 ms, 32.8 ms]), suggesting that
subjects may have improved their reaction times by the end of the task, but not enough to account
for the much lower reaction times in the stimulation block. Posterior distributions for the (group)
mean response times in each condition are shown in Figure 3.

Figure 3: Electrical muscle stimulation consistently preempted subjects' volitional movements. (a) Posterior distributions of the mean reaction times in each condition show that EMS-induced muscle movements occur earlier than subjects' natural muscle movements. (b) After outlier removal, measured "reaction" times (shown for all trials and subjects) are a nearly linear function of the stimulation latency, indicating that movements in the remaining trials are, in fact, EMS-actuated.

While EMS-caused movements were substantially faster than subject-caused movements, and the time of button press was clearly a function of stimulation latency (see Figure 3), subjects still reported that they caused roughly half of the movements. Overall, after outlier removal (see Methods and Figure 3), 51.98% of all trials across all subjects were judged as self-caused. On average, subjects reported that they caused 50.99% (SD: 14%) of movements. In other words, the Bayesian Optimization procedure was effective at controlling the proportion of trials in which movements were experienced as self-caused, generating a roughly 50-50 split of agency vs. non-agency trials.

3.2 Distinct early and late neural processes predict agency judgments.

Figure 4: The grand-average evoked EEG response to muscle stimulation.

Our MVPA procedure showed above-chance decoding performance *across subjects* (more often
such decoding is performed within subjects), reaching up to ROC-AUC = 0.587. While we report
the true-positive proportion within clusters across all clustering threshold (see Figure 5), we will
focus primarily on the clusters in which the true positive proportion exceeds 95%, since these
clusters are where we are sufficiently certain about the localization of the effect (Veillette, 2022).

Figure 5: Temporal generalization of neural patterns predicting sense of agency. (a) Classification performance
(ROC-AUC) for decoding subjects' judgement of agency for individual muscle movements, cross-validated across
subjects. Results are shown for all (train-time, test-time) pairs to visualize the temporal dynamics of patterns that
predict SoA. (b) Lower bounds on the true-positive proportion (TPP) within clusters, computed across all clustering
thresholds. The value represented at each (train-time, test-time) pair is the highest TPP of any cluster in which that
pair is included, thus larger values reflect greater certainty in the localization of effects.

Figure 6: Voltage patterns that predict sense of agency. (a) The EEG topographies that the linear classifiers trained
at each timepoint select for, reconstructed by inverting the trained classifier parameters using Haufe's trick (Haufe et

al., 2014). (b) The decoding performance when testing at each train time (identical to the values on the diagonal in
Fig. 5a). Training times are highlighted in yellow if included in a cluster with TPP > 0.95 at any test time (Fig. 5b).

The earliest such cluster occurs 83 ms after the onset of muscle stimulation (adjusted
threshold: $p < 4.5 \times 10^{-6}$). When comparing the patterns our decoding model selects for (see
Figure 6) to the average evoked response (see Figure 4), one notes that the polarity of the pattern
that predicts SoA is opposite the average response, indicating that the classifier would predict a
self-agency judgment as the result when the sensory response is suppressed—a finding consistent
with sensory attenuation (Voss et al., 2006). However, classifiers trained at earlier times do not
generalize to predict SoA at later times (see Figure 5). Later in the epoch, however, the temporal
dynamics of the predictive patterns change to reflect a single, sustained neural signature that
predicts SoA starting by at least 250 ms after stimulation and persisting at least until the end of the
epoch ($p < 0.003$).

3.3 Fractal complexity of brain activity predicts agency judgements.

Notably, trial-by-trial fractal dimension predicted SoA at almost every electrode (see Figure 7),
reaching an ROC-AUC of 0.614 at electrode C1 (adjusted threshold: p < 0.027), even after the
current source density transformation of the EEG signal was applied to attenuate the effects of
volume conduction (see Methods). This suggest that the (local) complexity of the brain activity is
increased uniformly throughout cortex following muscle movement when that movement is
perceived as self-caused (as compared to when it is not perceived as self-caused).

515
Figure 7: Classification performance for predicting trial-by-trial sense of agency from single-electrode fractal
metrics. (a) Classification performance for Higuchi fractal dimension. (b) Classification performance for the Hurst
exponent. Electrodes included in clusters in which the true positive proportion exceeds 95% are marked with white.

On the other hand, the Hurst exponent only predicted SoA at a single electrode at position
FC1 (ROC-AUC = 0.559, p = 0.0006), located above cortical regions involved in motor control
and planning, contralateral to the arm in which stimulation occurred (though we did not vary the
arm used for stimulation, so we would caution against interpreting this as a strictly contralateral
effect, though it is suggestive). This finding suggests a much more selective modulation of long-
range temporal dependencies, such that the activity of specific frontocentral cortical regions
becomes globally less to local perturbations in the absence of SoA—or, conversely, frontocentral
areas are more sensitive to inputs in the presence of SoA. However, it should be noted that this
analysis using fractal measures was exploratory, and consequently it should be interpreted as

motivating hypotheses about the dynamical characteristics of neural activity that may give rise to
the sense of agency, rather than as providing confirmatory evidence about such hypotheses.

4. Discussion

Our findings advance our understanding of how the sense of agency (SoA) is generated in
the brain, with important implications for the relationship between conscious self-awareness and
unconscious self-referential processing. In particular, the time course neural activity predicting
SoA in response to muscle stimulation is more consistent with classical sensorimotor monitoring
accounts (Blakemore et al., 2000; Wolpert et al., 1995) than previous studies have shown
comparing the neural responses to self- and other-caused tones (Kühn et al., 2011; Timm et al.,
2016). While our results still leave room for common downstream correlates of agency, they
suggest that early responses differentiating self and other may be more *modality specific* than
previously thought.

In the classic comparator model of SoA, originally imported from the motor control
literature (Wolpert et al., 1995), sensations are compared to the intended or predicted sensory
consequences of actions, and then congruent feedback is deemed self-caused and incongruent
feedback externally caused (Feinberg, 1978; C. D. Frith, 1987; Gallagher, 2000). Since it is well
documented that early sensory responses, especially those that are predictable, are attenuated
during movement (Blakemore et al., 2000), it seemed plausible that the same machinery could
parsimoniously account for conscious self-other discrimination. While this simple model is still
the basis of most modern accounts of SoA, it is now understood that the mechanisms of conscious
SoA diverge from low-level sensorimotor monitoring (C. Frith, 2012; Press et al., 2023; Synofzik
et al., 2008; Zaadnoordijk et al., 2019).

To this effect, recent studies using the typical paradigm, which probes the perception of
causality between a button press and subsequent tone (i.e., "broad" sense of agency over action
outcomes), have failed to find a relationship between the neural processes which would be affected
by low-level sensorimotor monitoring – that is, the early, pre-conscious (< 200 ms) response to
sensory stimulation – and conscious SoA (Kühn et al., 2011; Ohata et al., 2020; Timm et al., 2016).
Voss and colleagues report a strong dissociation, showing that comparator-model-like suppression
of early responses to self-caused sensation occur in both the presence and absence of SoA (Voss
et al., 2006). Since decades of research tell us that these early (< 200 ms) sensory responses reflect
preconscious, rather than conscious, processing of the sensory stimulus (Dehaene & Changeux,
2011; Libet et al., 1967; Sergent et al., 2005), these findings have been interpreted as meaning that
temporally early "exogeneous" neural responses (i.e., those that are a direct consequence sensory
input) do not inform agency judgements, but later "endogenous" neural responses (e.g., P3
component) associated with conscious attention do (Kühn et al., 2011). None of these authors
argue against the general idea of the comparator model, but rather suggest that the comparison
takes place at a higher level of abstraction than in the low-level sensorimotor monitoring used to
guide motor learning (Wolpert et al., 2011).

In contrast to Voss and colleagues' strong dissociation, in which an unconscious effect of
self-causation on early sensory processing is seen in the absence of conscious SoA, patterns in the
early neural response to sensation predicts   SoA even when that sensation was not actually self-
caused, as we exclusively analyzed trials in which movements were caused by EMS. The critical
difference is that we measured the neural response to muscle stimulation, and subjects made
agency judgements about *the muscle movement itself* rather than a downstream consequence of

movement. Thus, the mechanisms that give rise to narrow SoA over the musculature may overlap
with low-level sensorimotor monitoring mechanisms more than those mechanisms that give rise
to SoA over action outcomes more far removed from a subject's motor intention (Charalampaki et
al., 2022). In this vein, our data support the view that the earliest (usually pre-conscious) correlates
of conscious SoA may differ based on context (i.e., what is one being asked to make a judgement
about?), modality (e.g., proprioceptive or auditory), and level of abstraction.

However, it is worth noting that the earliest neural correlates of agency are not the end of
the story. Indeed, the comparator model for SoA has largely been usurped by dual-process models
in which the outcome of an initial comparator process is integrated with prospective, prior
information to produce a final agency judgement (Haggard, 2017; Legaspi & Toyoizumi, 2019;
Synofzik et al., 2008), and there is no clear theoretical for why or how multiple comparator
processes taking place at multiple levels of abstraction may not be integrated into a single agency
judgement. In fact, the shift we observe from transient to sustained patterns of neural activity
predicting agency is quite consistent with that predicted by dual-process models of action
processing (Charles et al., 2014; Del Cul et al., 2009). Specifically, the sustained nature of the
predictive voltage patterns is consistent with a previously observed signature of high-level
novelty/error detection that has been argued to require conscious awareness (Dehaene & King,
2016) and previously proposed to inform agency judgements (Kühn et al., 2011). An intriguing
possibility then, which hybridizes the competing views proposed in the introduction, is that pre-
conscious (roughly < 200 ms) predictors of SoA judgements will be context specific, but post-
consciousness "neural correlates of-self-awareness" integrate across modality-specific
comparators. We do not manipulate awareness of action and outcomes here, so it is up to future
work to test this hypothesis directly. Such investigations, which can compare sense of agency over
actions with SoA over those actions' downstream outcomes, are made possible by extending the
paradigm we introduce here.

Lastly, though departing somewhat from our main research question, our exploratory
analysis too yielded important findings to motivate future SoA research. We find that the fractal
dimension of the EEG measured across almost the whole scalp is higher on trials in which the
subject experiences movements as self-caused. The fractal dimension is just one complexity
measure that has been associated with conscious awareness; indeed, the fractal dimension of both
the background EEG and the EEG response to perturbation is highly predictive of state of
consciousness (Kesić & Spasić, 2016; Ruiz de Miras et al., 2019), motivating the argument that
fractal dimension of neural responses is related to the depth of conscious processing. Importantly,
in a strictly self-similar time series, the fractal dimension should be inversely related to the Hurst
exponent (Gneiting & Schlather, 2004), which is a measure of long-range temporal correlation
often interpreted as reflecting a neural time series' global sensitivity to local perturbation
(Churchill et al., 2016; Kardan et al., 2020; Zhuang et al., 2022). If Hurst exponent varied with
fractal dimension in the expected direction, this would suggest that the brain was less sensitive to
feedback on agentic trials (which would perhaps suggesting that the illusion of agency occurred
when the brain was "ignoring" incongruent feedback). Interestingly, Hurst exponent only predicts
SoA after multiple comparisons correction in a single electrode found over sensorimotor cortex,
in the *opposite* of the expected direction. This analysis and its interpretation are post hoc additions
to our study, and the Hurst exponent is unreliable when estimates from such short time series as
ours, so this latter result may not be robust. However, the observed divergence from the expected
relationship between fractal dimension and the Hurst exponent may suggest that neural activity
departs from fractal self-similarity in an important way, which allows the complexity of brain

activity to scale with agency while selectively reducing sensorimotor systems' sensitivity to local
perturbation when sensation is deemed externally caused. Perhaps an explanation for the known
effects of SoA on motor learning may be lurking in measurable properties of the intrinsic dynamics
of neural time series (Kasahara et al., 2021; Russo et al., 2020; Wen et al., 2021). Exploring these
measures further may be of interest to both SoA and motor control researchers.

In summary, while SoA has become a topic of increased attention in recent decades, most
research in the area has focused on the experience of agency over downstream consequences of
one's actions as they affect the external world rather than the more basal experience of directing
one's own muscles (Haggard, 2008, 2017). We introduce the use of human-in-the-loop Bayesian
optimization, in combination with electrical muscle stimulation, to experimentally manipulate the
subjective experience of controlling the musculature. As we showcase here, this approach enables
novel behavioral and neuroimaging investigations into the substrate of embodied self-awareness.
Our results provide confirmatory evidence for the predictive relationship between low-level
sensorimotor processes and SoA for muscle movements, which seems not to hold for the sensory
response to action consequences (Dewey & Knoblich, 2014; Timm et al., 2016). While our
findings suggest that the early neural correlates of SoA may differ by context and modality, the
transition from transient to sustained neural patterns that predict SoA in our data suggest at least
two distinct neural processes contributing to agency judgments, as posited by dual-process theories
of action selection and monitoring (Del Cul et al., 2009). This leaves open the possibility that
modality-specific, pre-conscious predictors of SoA are still integrated in a single agency judgment
downstream. Such a possibility could explain how information from multiple scales of biological
organization are integrated into a unified experience of self even the mechanism of self-other
differentiation differs across scales. We suggest that this hypothesis is a fruitful avenue of research
for the emerging science of self-awareness.

<center>References</center>

Agrawal, S., & Goyal, N. (2012). Analysis of Thompson Sampling for the Multi-armed Bandit Problem. *Proceedings of the 25th Annual Conference on Learning Theory*, 39.1-39.26. https://proceedings.mlr.press/v23/agrawal12.html

Appelhoff, S., Hurst, A., Lawrence, A., Li, A., Mantilla, R., Yorguin, J., O'Reilly, C., Xiang, L., & Dancker, J. (2022). *PyPREP: A Python implementation of the preprocessing pipeline (PREP) for EEG data*. Zenodo. https://doi.org/10.5281/zenodo.6363575

Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K.-M., & Robbins, K. A. (2015). The PREP pipeline: Standardized preprocessing for large-scale EEG analysis. *Frontiers in Neuroinformatics*, *9*. https://www.frontiersin.org/articles/10.3389/fninf.2015.00016

Bingham, E., Chen, J. P., Jankowiak, M., Obermeyer, F., Pradhan, N., Karaletsos, T., Singh, R., Szerlip, P., Horsfall, P., & Goodman, N. D. (2019). Pyro: Deep Universal Probabilistic Programming. *Journal of Machine Learning Research*, *20*(28), 1–6.

Blakemore, S.-J., Wolpert, D., & Frith, C. (2000). Why can't you tickle yourself? *NeuroReport*, *11*(11), R11.

Bouckaert, R. R., & Frank, E. (2004). Evaluating the Replicability of Significance Tests for Comparing Learning Algorithms. In H. Dai, R. Srikant, & C. Zhang (Eds.), *Advances in Knowledge Discovery and Data Mining* (pp. 3–12). Springer. https://doi.org/10.1007/978-3-540-24775-3_3

Capretto, T., Piho, C., Kumar, R., Westfall, J., Yarkoni, T., & Martin, O. A. (2022). Bambi: A Simple Interface for Fitting Bayesian Linear Models in Python. *Journal of Statistical Software*, *103*, 1–29. https://doi.org/10.18637/jss.v103.i15

668 Charalampaki, A., Ninija Karabanov, A., Ritterband-Rosenbaum, A., Bo Nielsen, J., Roman

669   Siebner, H., & Schram Christensen, M. (2022). Sense of agency as synecdoche: Multiple

670   neurobiological mechanisms may underlie the phenomenon summarized as sense of

671   agency. *Consciousness and Cognition*, *101*, 103307.

672   https://doi.org/10.1016/j.concog.2022.103307

673 Charles, L., King, J.-R., & Dehaene, S. (2014). Decoding the Dynamics of Action, Intention, and

674   Error Detection for Conscious and Subliminal Stimuli. *Journal of Neuroscience*, *34*(4),

675   1158–1170. https://doi.org/10.1523/JNEUROSCI.2465-13.2014

676 Christensen, M. S., & Grünbaum, T. (2018). Sense of agency for movements. *Consciousness and*

677   *Cognition*, *65*, 27–47. https://doi.org/10.1016/j.concog.2018.07.002

678 Churchill, N. W., Spring, R., Grady, C., Cimprich, B., Askren, M. K., Reuter-Lorenz, P. A., Jung,

679   M. S., Peltier, S., Strother, S. C., & Berman, M. G. (2016). The suppression of scale-free

680   fMRI brain dynamics across three different sources of effort: Aging, task novelty and task

681   difficulty. *Scientific Reports*, *6*(1), Article 1. https://doi.org/10.1038/srep30895

682 Darvas, F., Scherer, R., Ojemann, J. G., Rao, R. P., Miller, K. J., & Sorensen, L. B. (2010). High

683   gamma mapping using EEG. *NeuroImage*, *49*(1), 930–938.

684   https://doi.org/10.1016/j.neuroimage.2009.08.041

685 Dehaene, S., & Changeux, J.-P. (2011). Experimental and Theoretical Approaches to Conscious

686   Processing. *Neuron*, *70*(2), 200–227. https://doi.org/10.1016/j.neuron.2011.03.018

687 Dehaene, S., & King, J.-R. (2016). Decoding the Dynamics of Conscious Perception: The

688   Temporal Generalization Method. In G. Buzsáki & Y. Christen (Eds.), *Micro-, Meso- and*

689   *Macro-Dynamics of the Brain*. Springer.

690   http://www.ncbi.nlm.nih.gov/books/NBK435757/

Del Cul, A., Dehaene, S., Reyes, P., Bravo, E., & Slachevsky, A. (2009). Causal role of prefrontal

cortex in the threshold for access to consciousness. *Brain*, *132*(9), 2531–2540.

https://doi.org/10.1093/brain/awp111

Dewey, J. A., & Knoblich, G. (2014). Do implicit and explicit measures of the sense of agency

measure the same thing? *PLoS ONE*, *9*. https://doi.org/10.1371/journal.pone.0110118

Dietterich, T. G. (1998). Approximate Statistical Tests for Comparing Supervised Classification

Learning Algorithms. *Neural Computation*, *10*(7), 1895–1923.

https://doi.org/10.1162/089976698300017197

Eke, A., Hermán, P., Bassingthwaighte, J. B., Raymond, G. M., Percival, D. B., Cannon, M., Balla,

I., & Ikrényi, C. (2000). Physiological time series: Distinguishing fractal noises from

motions. *Pflugers Archiv European Journal of Physiology*, *439*(4), 403–415. Scopus.

https://doi.org/10.1007/s004240050957

Eke, A., Herman, P., Kocsis, L., & Kozak, L. R. (2002). Fractal characterization of complexity in

temporal physiological signals. *Physiological Measurement*, *23*(1), R1–R38. Scopus.

https://doi.org/10.1088/0967-3334/23/1/201

Feinberg, I. (1978). Efference Copy and Corollary Discharge: Implications for Thinking and Its

Disorders. *Schizophrenia Bulletin*, *4*, 636–640. https://doi.org/10.1093/schbul/4.4.636

Fries, P., Scheeringa, R., & Oostenveld, R. (2008). Finding Gamma. *Neuron*, *58*(3), 303–305.

https://doi.org/10.1016/j.neuron.2008.04.020

Frith, C. (2012). Explaining delusions of control: The comparator model 20 years on.

Consciousness and Cognition, *21*(1), 52–54. https://doi.org/10.1016/j.concog.2011.06.010

Frith, C. D. (1987). The positive and negative symptoms of schizophrenia reflect impairments in

the perception and initiation of action. *Psychological Medicine*, *17*(3), 631–648.

https://doi.org/10.1017/S0033291700025873

Gallagher, S. (2000). Philosophical conceptions of the self: Implications for cognitive science.

Trends in Cognitive Sciences, *4*(1), 14–21. https://doi.org/10.1016/S1364-6613(99)01417-

5

Gneiting, T., & Schlather, M. (2004). Stochastic models which separate fractal dimension and

Hurst effect. *SIAM Review*, *46*(2), 269–282. https://doi.org/10.1137/S0036144501394387

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Parkkonen,

L., & Hämäläinen, M. S. (2014). MNE software for processing MEG and EEG data.

NeuroImage, *86*, 446–460. https://doi.org/10.1016/j.neuroimage.2013.10.027

Haggard, P. (2008). Human volition: Towards a neuroscience of will. *Nature Reviews*

Neuroscience, *9*(12), Article 12. https://doi.org/10.1038/nrn2497

Haggard, P. (2017). Sense of agency in the human brain. *Nature Reviews Neuroscience*, *18*(4),

Article 4. https://doi.org/10.1038/nrn.2017.14

Haggard, P., & Clark, S. (2003). Intentional action: Conscious experience and neural prediction.

Consciousness and Cognition, *12*(4), 695–707. https://doi.org/10.1016/S1053-

8100(03)00052-7

Haggard, P., Clark, S., & Kalogeras, J. (2002). Voluntary action and conscious awareness. *Nature*

Neuroscience, *5*(4), Article 4. https://doi.org/10.1038/nn827

Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J.-D., Blankertz, B., & Bießmann, F.

(2014). On the interpretation of weight vectors of linear models in multivariate

neuroimaging. *NeuroImage*, *87*, 96–110.

https://doi.org/10.1016/j.neuroimage.2013.10.067

Higuchi, T. (1988). Approach to an irregular time series on the basis of the fractal theory. *Physica*

D: Nonlinear Phenomena, *31*(2), 277–283. https://doi.org/10.1016/0167-2789(88)90081-

4

Hyvarinen, A. (1999). Fast and robust fixed-point algorithms for independent component analysis.

IEEE Transactions on Neural Networks, *10*(3), 626–634.

https://doi.org/10.1109/72.761722

Jas, M., Engemann, D. A., Bekhti, Y., Raimondo, F., & Gramfort, A. (2017). Autoreject:

Automated artifact rejection for MEG and EEG data. *NeuroImage*, *159*, 417–429.

https://doi.org/10.1016/j.neuroimage.2017.06.030

Kardan, O., Adam, K. C. S., Mance, I., Churchill, N. W., Vogel, E. K., & Berman, M. G. (2020).

Distinguishing cognitive effort and working memory load using scale-invariance and alpha

suppression in EEG. *NeuroImage*, *211*, 116622.

https://doi.org/10.1016/j.neuroimage.2020.116622

Kasahara, S., Nishida, J., & Lopes, P. (2019). Preemptive action: Accelerating human reaction

using electrical muscle stimulation without compromising agency. *Proceedings of the 2019*

CHI Conference on Human Factors in Computing Systems, 1–15.

https://dl.acm.org/doi/abs/10.1145/3290605.3300873

Kasahara, S., Takada, K., Nishida, J., Shibata, K., Shimojo, S., & Lopes, P. (2021). Preserving

Agency During Electrical Muscle Stimulation Training Speeds up Reaction Time Directly

After Removing EMS. *Proceedings of the 2021 CHI Conference on Human Factors in*

Computing Systems, 1–9. https://doi.org/10.1145/3411764.3445147

Kayser, J., & Tenke, C. E. (2015). On the benefits of using surface Laplacian (current source

density) methodology in electrophysiology. *International Journal of Psychophysiology*,

97(3), 171–173. https://doi.org/10.1016/j.ijpsycho.2015.06.001

Kesić, S., & Spasić, S. Z. (2016). Application of Higuchi's fractal dimension from basic to clinical

neurophysiology: A review. *Computer Methods and Programs in Biomedicine*, *133*, 55–

70. https://doi.org/10.1016/j.cmpb.2016.05.014

King, J.-R., & Dehaene, S. (2014). Characterizing the dynamics of mental representations: The

temporal generalization method. *Trends in Cognitive Sciences*, *18*(4), 203–210.

https://doi.org/10.1016/j.tics.2014.01.002

Kühn, S., Nenchev, I., Haggard, P., Brass, M., Gallinat, J., & Voss, M. (2011). Whodunnit?

Electrophysiological Correlates of Agency Judgements. *PLOS ONE*, *6*(12), e28657.

https://doi.org/10.1371/journal.pone.0028657

Legaspi, R., & Toyoizumi, T. (2019). A Bayesian psychophysics model of sense of agency. *Nature*

Communications, *10*(1), Article 1. https://doi.org/10.1038/s41467-019-12170-0

Lhermitte, F., Pillon, B., & Serdaru, M. (1986). Human autonomy and the frontal lobes. Part I:

Imitation and utilization behavior: A neuropsychological study of 75 patients. *Annals of*

Neurology, *19*(4), 326–334. https://doi.org/10.1002/ana.410190404

Li, X., Liang, Z., Kleiner, M., & Lu, Z.-L. (2010). RTbox: A device for highly accurate response

time measurements. *Behavior Research Methods*, *42*(1), 212–225.

https://doi.org/10.3758/BRM.42.1.212

Liang, K.-Y., & Zeger, S. (1986). Longitudinal data analysis using generalized linear models.

Biometrika, *73*(1), 13–22. https://doi.org/10.1093/biomet/73.1.13

Libet, B., Alberts, W. W., Wright, E. W., & Feinstein, B. (1967). Responses of Human

Somatosensory Cortex to Stimuli below Threshold for Conscious Sensation. *Science*,

158(3808), 1597–1600. https://doi.org/10.1126/science.158.3808.1597

Martis, R. J., Tan, J. H., Chua, C. K., Loon, T. C., Yeo, S. W. J., & Tong, L. (2015). Epileptic eeg

classification using nonlinear parameters on different frequency bands. *Journal of*

Mechanics   in   Medicine   and   Biology,   *15*(03),   1550040.

https://doi.org/10.1142/S0219519415500402

Nadeau, C., & Bengio, Y. (1999). Inference for the Generalization Error. *Advances in Neural*

Information   Processing   Systems,   *12*.

https://proceedings.neurips.cc/paper/1999/hash/7d12b66d3df6af8d429c1a357d8b9e1a-

Abstract.html

Ohata, R., Asai, T., Kadota, H., Shigemasu, H., Ogawa, K., & Imamizu, H. (2020). Sense of

Agency Beyond Sensorimotor Process: Decoding Self-Other Action Attribution in the

Human Brain. *Cerebral Cortex*, *30*(7), 4076–4091. https://doi.org/10.1093/cercor/bhaa028

Oizumi, M., Albantakis, L., & Tononi, G. (2014). From the Phenomenology to the Mechanisms

of Consciousness: Integrated Information Theory 3.0. *PLOS Computational Biology*,

10(5), e1003588. https://doi.org/10.1371/journal.pcbi.1003588

Oliver, R., & Ballester, J. L. (1998). Is there memory in solar activity? *Physical Review E -*

Statistical Physics, Plasmas, Fluids, and Related Interdisciplinary Topics, *58*(5), 5650–

5654. Scopus. https://doi.org/10.1103/PhysRevE.58.5650

Panikkath, R., Panikkath, D., Mojumder, D., & Nugent, K. (2014). The alien hand syndrome.

Proceedings (Baylor University. Medical Center), *27*(3), 219–220.

Patil, A., Huard, D., & Fonnesbeck, C. J. (2010). PyMC: Bayesian Stochastic Modelling in Python.

Journal of Statistical Software, *35*(4), 1–81.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,

Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D.,

Brucher, M., Perrot, M., & Duchesnay, É. (2011). Scikit-learn: Machine Learning in

Python. *The Journal of Machine Learning Research*, *12*, 2825–2830.

Peirce, J. W. (2007). PsychoPy—Psychophysics software in Python. *Journal of Neuroscience*

Methods, *162*(1), 8–13. https://doi.org/10.1016/j.jneumeth.2006.11.017

Press, C., Thomas, E. R., & Yon, D. (2023). Cancelling cancellation? Sensorimotor control,

agency, and prediction. *Neuroscience & Biobehavioral Reviews*, *145*, 105012.

https://doi.org/10.1016/j.neubiorev.2022.105012

Qian, B., & Rasheed, K. (2004, November). Hurst exponent and financial market predictability.

Proceedings of the IASTED International Conference. IASTED conference on Financial

Engineering and Applications, Cambridge, MA.

Ramachandran, V. S., & Hirstein, W. (1998). The perception of phantom limbs. The D. O. Hebb

lecture. *Brain*, *121*(9), 1603–1630. https://doi.org/10.1093/brain/121.9.1603

Rosenblatt, J. D., Finos, L., Weeda, W. D., Solari, A., & Goeman, J. J. (2018). All-Resolutions

Inference   for   brain   imaging.   *NeuroImage*,   *181*,   786–796.

https://doi.org/10.1016/j.neuroimage.2018.07.060

Ruiz de Miras, J., Soler, F., Iglesias-Parro, S., Ibáñez-Molina, A. J., Casali, A. G., Laureys, S.,

Massimini, M., Esteban, F. J., Navas, J., & Langa, J. A. (2019). Fractal dimension analysis

of states of consciousness and unconsciousness using transcranial magnetic stimulation.

Computer Methods and Programs in Biomedicine, *175*, 129–137. https://doi.org/10.1016/j.cmpb.2019.04.017

Russo, A. A., Khajeh, R., Bittner, S. R., Perkins, S. M., Cunningham, J. P., Abbott, L. F., & Churchland, M. M. (2020). Neural Trajectories in the Supplementary Motor Area and Motor Cortex Exhibit Distinct Geometries, Compatible with Different Classes of Computation. *Neuron*, *107*(4), 745-758.e6. https://doi.org/10.1016/j.neuron.2020.05.020

Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology*, *56*(6), e13335. https://doi.org/10.1111/psyp.13335

Schiratti, J.-B., Le Douget, J.-E., Le Van Quyen, M., Essid, S., & Gramfort, A. (2018). An Ensemble Learning Approach to Detect Epileptic Seizures from Long Intracranial EEG Recordings. *2018 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, 856–860. https://doi.org/10.1109/ICASSP.2018.8461489

Seabold, S., & Perktold, J. (2010). *Statsmodels: Econometric and Statistical Modeling with Python*. 92–96. https://doi.org/10.25080/Majora-92bf1922-011

Sergent, C., Baillet, S., & Dehaene, S. (2005). Timing of the brain events underlying access to consciousness during the attentional blink. *Nature Neuroscience*, *8*(10), Article 10. https://doi.org/10.1038/nn1549

Smith, N. J., & Kutas, M. (2015). Regression-based estimation of ERP waveforms: II. Nonlinear effects, overlap correction, and practical considerations. *Psychophysiology*, *52*(2), 169–181. https://doi.org/10.1111/psyp.12320

Sokunbi, M. O., Gradin, V. B., Waiter, G. D., Cameron, G. G., Ahearn, T. S., Murray, A. D.,

Steele, D. J., & Staff, R. T. (2014). Nonlinear Complexity Analysis of Brain fMRI Signals

in Schizophrenia. *PLOS ONE*, *9*(5), e95146. https://doi.org/10.1371/journal.pone.0095146

Stier, A., Cardenas-Iniguez, C., Kardan, O., Moore, T., Meyer, F., Rosenberg, M., Kaczkurkin, A.,

Lahey, B., & Berman, M. (2021). *A Scale-Free Gradient of Cognitive Resource*

Disruptions in Childhood Psychopathology. https://doi.org/10.1101/2021.08.24.457554

Synofzik, M., Vosgerau, G., & Newen, A. (2008). Beyond the comparator model: A multifactorial

two-step account of agency. *Consciousness and Cognition*, *17*(1), 219–239.

https://doi.org/10.1016/j.concog.2007.03.010

Tajima, D., Nishida, J., Lopes, P., & Kasahara, S. (2022). Whose Touch is This?: Understanding

the Agency Trade-Off Between User-Driven Touch vs. Computer-Driven Touch. *ACM*

Transactions on Computer-Human Interaction, *29*(3), 24:1-24:27.

https://doi.org/10.1145/3489608

Timm, J., Schönwiesner, M., Schröger, E., & SanMiguel, I. (2016). Sensory suppression of brain

responses to self-generated sounds is observed with and without the perception of agency.

Cortex, *80*, 5–20. https://doi.org/10.1016/j.cortex.2016.03.018

Veillette, J. (2022). *MNE-ARI: All-resolutions Inference for M/EEG in Python* (0.1.2). Zenodo.

https://doi.org/10.5281/zenodo.5813807

Voss, M., Ingram, J. N., Haggard, P., & Wolpert, D. M. (2006). Sensorimotor attenuation by

central motor command signals in the absence of movement. *Nature Neuroscience*, *9*(1),

Article 1. https://doi.org/10.1038/nn1592

Wen, W., Ishii, H., Ohata, R., Yamashita, A., Asama, H., & Imamizu, H. (2021). Perception and

control: Individual difference in the sense of agency is associated with learnability in

sensorimotor adaptation. *Scientific Reports*, *11*(1), Article 1.

https://doi.org/10.1038/s41598-021-99969-4

Westfall, J. (2017). Statistical details of the default priors in the Bambi library. In *ArXiv e-prints*.

https://doi.org/10.48550/arXiv.1702.01201

Wolf, S. L., Lecraw, D. E., Barton, L. A., & Jann, B. B. (1989). Forced use of hemiplegic upper

extremities to reverse the effect of learned nonuse among chronic stroke and head-injured

patients. *Experimental Neurology*, *104*(2), 125–132. https://doi.org/10.1016/S0014-

4886(89)80005-6

Wolpert, D. M., Diedrichsen, J., & Flanagan, J. R. (2011). Principles of sensorimotor learning.

Nature Reviews Neuroscience, *12*(12), Article 12. https://doi.org/10.1038/nrn3112

Wolpert, D. M., Ghahramani, Z., & Jordan, M. I. (1995). An Internal Model for Sensorimotor

Integration. *Science*, *269*(5232), 1880–1882. https://doi.org/10.1126/science.7569931

Zaadnoordijk, L., Besold, T. R., & Hunnius, S. (2019). A match does not make a sense: On the

sufficiency of the comparator model for explaining the sense of agency. *Neuroscience of*

Consciousness, *2019*(1), niz006. https://doi.org/10.1093/nc/niz006

Zhuang, C., Meidenbauer, K. L., Kardan, O., Stier, A. J., Choe, K. W., Cardenas-Iniguez, C.,

Huppert, T. J., & Berman, M. G. (2022). Scale invariance in fNIRS as a measurement of

cognitive load. *Cortex*, *154*, 62–76. https://doi.org/10.1016/j.cortex.2022.05.009