A transmission chain linking *Mycobacterium ulcerans* with *Aedes notoscriptus*

mosquitoes, possums and human Buruli ulcer cases in southeastern Australia

Peter T. Mee[1,*,§], Andrew H. Buultjens[2, §], Jane Oliver[3, §], Karen Brown[1], Jodie C. Crowder[1], Jessica L. Porter[2],

Emma C. Hobbs[3], Louise M. Judd[2], George Taiaroa[3], Natsuda Puttharak[1], Deborah A. Williamson[3,4], Kim R.

Blasdell[5], Ee Laine Tay[6], Rebecca Feldman[6], Mutizwa Odwell Muzari[7], Chris Sanders[2], Stuart Larsen[2], Simon

R. Crouch[8], Paul D. R. Johnson[9], John R. Wallace[10], David J. Price[3,11], Ary A. Hoffmann[12,^], Katherine B.

Gibney[3,^], Timothy P. Stinear[3,13,^,*] and Stacey E. Lynch[1&,^]

[1]Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria, Australia.

[2]Department of Microbiology and Immunology, Doherty Institute for Infection and Immunity, University of
Melbourne, Melbourne, Victoria, Australia.

[3]Department of Infectious Diseases, University of Melbourne, Doherty Institute for Infection and Immunity, Victoria,
3000, Australia

[4]Victorian Infectious Diseases Reference Laboratory, Melbourne Health, Doherty Institute for Infection and Immunity,
Victoria, Australia

[5]Australian Centre for Disease Preparedness, CSIRO, Geelong, Victoria, Australia.

[6]Department of Health, Melbourne, Victoria, Australia.

[7]Medical Entomology, Tropical Public Health Services Cairns, Cairns and Hinterland Hospital & Health Services, Cairns,
Queensland, Australia.

[8]South East Public Health Unit, Monash Health, Clayton, Victoria, Australia.

[9]North East Public Health Unit, Austin Health, Heidelberg, Victoria, Australia.

[10]Department of Biology, Millersville University, Millersville, PA, USA.

[11]Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, University of
Melbourne, Parkville, Victoria, Australia.

[12]Pest and Environmental Adaptation Research Group, School of BioSciences, Bio21 Institute, University of
Melbourne, Parkville, Victoria, Australia.

[13]WHO Collaborating Centre for *Mycobacterium ulcerans*, Doherty Institute for Infection and Immunity, Melbourne,
Victoria, Australia

Corresponding authors: *peter.mee@agriculture.vic.gov.au; tstinear@unimelb.edu.au

§Contributed equally

^Joint senior authors

& Current affiliations Australian Centre for Disease Preparedness, CSIRO, Geelong, Victoria, Australia

Keywords: *Mycobacterium ulcerans*, Buruli ulcer, *Aedes notoscriptus, disease transmission*

Abstract

In temperate southeastern Australia over the past two decades there has been a marked progressive increase in human cases of Buruli ulcer, an infection of subcutaneous tissue caused by *Mycobacterium ulcerans*. Native possums are the major local environmental reservoir of *M. uclerans* as they not only develop Buruli lesions but they also shed *M. ulcerans* in their excreta. However the way humans acquire *M. ulcerans* from possums has not been determined. Previous case-control studies, insect field surveys and vector competence studies have suggested a role for mosquitoes in *M. ulcerans* transmission between possums and humans. To explore these links we conducted an extensive, 4-month structured mosquito field survey and four *ad hoc* field surveys across an area of 350km^2 on the Mornington Peninsula, an area endemic for Buruli ulcer. We then compared spatial and temporal patterns of *M. ulcerans*-positive mosquito occurrence with *M. ulcerans*-positive possums (established by previous possum excreta surveys) and human Buruli ulcer cases across the region. We used metabarcoding to assess mosquito blood-feeding host preference and to reconstruct *M. ulcerans* genomes from positive mosquitoes to test epidemiological inferences. We collected 66,325 mosquitoes spanning 26 different species from 180 repeatedly sampled traps over a 4-month period. *Culex molestus* and *Aedes notoscriptus* were the dominant species (42% and 35% of trapped mosquitoes, respectively). PCR screening 25% of trapped mosquitoes revealed a significant association between *M. ulcerans* and *Ae. notoscriptus* (p<0.0001) with a maximum likelihood estimate (MLE) of 5.88 *M. ulcerans* positive mosquitoes per 1,000 tested. Using spatial scanning statistics, we also observed significant overlap between clusters of *M. ulcerans*-positive *Ae. notoscriptus*, *M. ulcerans*-positive possum excreta and human Buruli ulcer cases. Metabarcoding analyses of blood-fed *Ae. notoscriptus* showed individual mosquitoes had fed both on humans and native possums. Enrichment genome sequencing from PCR-positive mosquitoes confirmed identical *M. ulcerans* genome single-nucleotide polymorphism (SNP) profiles between insects, possums and clinical human isolates within the same regions. These findings indicate that certain mosquito species likely transmit *M. ulcerans* in southeastern Australia and highlight mosquito control as a plausible means to control the Buruli ulcer epidemic in our region.

Introduction

Mycobacterium ulcerans is the causative agent of a neglected tropical skin disease called Buruli ulcer, a necrotising infection of skin and subcutaneous tissue [1]. Buruli ulcer is rarely a fatal condition but can cause severe tissue destruction if not diagnosed and managed effectively [2]. Buruli ulcer has been described in more than 32 countries worldwide [3] and is an ongoing public health issue in west and central Africa [4]. Buruli ulcer has also been unexpectedly surging in temperate southeastern Australia (Figure 1) and encroaching on the major metropolitan centres of Melbourne (population 5.1 million) and Geelong (population 274,000), with >250 cases routinely notified each year since 2017 to the Victorian State Government

Department of Health [5].

Figure 1. Location of the Mornington Peninsula and the Bellarine Peninsula of Victoria, Australia. Population density is represented on the map from low to high (white to red) based on the human population per square kilometre.

How humans contract Buruli ulcer is a central question that has intrigued scientists and confounded public

health control efforts since the discovery of *M. ulcerans* from patients in the Bairnsdale region of Australia

in the 1930s and across Africa shortly after [1,6,7]. Buruli ulcer epidemiology can be unpredictable, with

outbreaks emerging in specific geographical areas and then disappearing over a number of years. *M.*

ulcerans   has   a

4-5 month median incubation period. It is very challenging to isolate the bacterium in pure culture from the

environment, presumably due to its very slow growth, although it can be isolated from human skin lesions.

These factors combined have made it incredibly challenging to establish how *M. ulcerans* is spread to

humans, despite global efforts to investigate this over more than 80 years [8].

The discovery that Buruli ulcer is a zoonosis and that Australian native possums are a major wildlife

source of *M. ulcerans* that is intimately linked with disease transmission has addressed one key component

of the transmission enigma [9-14]. The role of mosquitoes as vectors spreading *M. ulcerans* from possums to

humans (and between possums) is the subject of the research presented in this current study. The first

indications that mosquitoes might be vectors of *M. ulcerans* from possums to humans in Australia came

from a series of entomological field surveys in the southeast of the country in response to an increase in

Buruli ulcer cases in the seaside township of Point Lonsdale, located on the Bellarine Peninsula [15]. Among

the 12 species identified from a trapping effort that collected 11,500 mosquitoes, five different species

were IS2404 PCR positive for *M. ulcerans,* including *Aedes camptorhynchus*, *Aedes notoscriptus*,

Coquillettidia linealis, Culex australicus, and *Anopheles annulipes* (maximum likelihood estimate (MLE) was

4.11/1000 mosquitoes) [15]. Note that IS2404 is a *M. ulcerans* specific insertion sequence and molecular

target for the gold-standard diagnostic PCR for Buruli ulcer [16].

A concurrent case control study performed in the same geographic area identified only two factors

associated with the odds of being diagnosed with Buruli ulcer: insect repellent use (OR 0.37, 95% CI 0.19–

0.69) and being bitten by mosquitoes on the lower legs (OR 2.60, 95% CI 1.22–5.53). A variety of outdoor

activities were also surveyed but were not independently predictive suggesting that mosquito exposure

specifically rather than environmental exposure generally might be the main mode of MU transmission to

humans [17]. In Africa, two case control studies conducted in Cameroon both found use of bed nets as a

protective factor against Buruli ulcer, (OR 0.4, 95% CI 0.2-0.9, p=0.04) [18] and (OR 0.1, 95% CI 0.03-0.3,

p<0.001) [19].

However, case control studies and entomological surveys alone are insufficient to indict biological

agents as vectors of pathogens. There are formal frameworks used in biomedicine, such as the Barnett

criteria [3] that build hierarchies of evidence to implicate a candidate disease vector. Here, we build on this

aforementioned research to formally address the Barnett criteria, and test the hypothesis that mosquitoes

are vectors of *M. ulcerans* to humans [20-22]. A summary of our findings is shown in Table 1, including the new

data presented in this study that specifically address criteria 1-3.

Table 1: Summary of evidence to indict mosquitoes as vectors of *M. ulcerans*

Barnett Criterion	Evidence	Reference
1. Acquire from source reservoir	*M. ulcerans* genome sequences from possum excreta matches genomes recovered from mosquitoes in same area	This study
2. Close links with infected hosts	*M. ulcerans* genome sequences from humans matches genome recovered from mosquitoes in same area. Mosquito bloodmeal analysis confirms human-possum link.	This study
3. Repeat detections from candidate vector	*M. ulcerans* detected with mosquitoes in seven separate field surveys	[15,21], this study
4. Experimental demonstration of transmission	Mosquito-*M. ulcerans* mechanical transmission demonstrated using murine contaminated skin model	[22]

114   In the research presented here, based on a very substantial field survey of >65,000 mosquitoes

115 undertaken over four months in 2019 and 2020, and four smaller *ad hoc* surveys conducted between 2016

116 and 2021, we employed spatial clustering analyses, bacterial enrichment genome sequencing and mosquito

117 blood meal metabarcoding to continue building a hierarchy of evidence that implicates mosquitoes as

118 vectors of *M. ulcerans* from local wildlife sources to humans.

Material and methods

Study Site

Insects were collected from the Mornington Peninsula suburbs of Rye (population 8,416), Blairgowrie (population 2,313), Tootgarook (population 2,869) and Capel Sound (population 4,930) [23]. The study area encompasses an area of 350 km^2 and is located 90 km south of Melbourne, the capital city of Victoria [10] (Figure 1). The area was originally covered in low-lying coastal vegetation [10], receives an average annual rainfall of 740 mm, and an elevation of 60 metres above sea-level [24]. The Mornington Peninsula region continues to maintain among the highest incidences of Buruli ulcer in the world [25] with a conservative local incidence estimate of 55 human cases/100,000 population in 2022 [5,26].

Arthropod collection and identification

Mosquito trapping campaigns used Biogent Sentinal (BGS) traps (Biogent) that were baited with dry ice pellets to provide a source of carbon dioxide over an intended 12-hour period. BGS traps were set out at dusk and collected at dawn in shaded locations on the grassed edges of the roadways in the study area. GPS locations for all traps were recorded. Trapped mosquitoes were knocked down with CO_2 by placing the catch-bag in dry ice before being transported back to the laboratory and kept at -20°C until processing. Mosquito species were morphologically identified using a stereo dissecting microscope (Nikon, SMZ800N) and reference to taxonomic keys [27-29].

Other (non-mosquito) arthropods were collected using Yellow Sticky Traps (YST) and Sticky Ovitraps (SO). Two YST and SO were placed in residential properties where householders had previously noted insect activity to the researchers [9]. The SO traps were placed on the ground and had hay grass infusion (3 Jack Rabbit (clover/lucerne) pellets (Laucke Mills, Barossa Valley, SA, Australia) in 500 mL water) added to them, while the YST (Bugs for Bugs, Toowoomba, QLD, Australia) were placed on the ground with a 14 cm plant tag plastic T-support (Garden City Plastics, Dandenong, VIC, Australia). Within three to four days of

144 being set, residents were asked to pack up the YST and SO by covering the sticky card with a plastic film,

145 and return the sealed traps to the laboratory, where they were stored at -20°C. Non-mosquito arthropods

146 were morphologically identified to family level and, if PCR positive for *M. ulcerans,* were DNA barcoded for

147 species confirmation by targeting the cytochrome c oxidase subunit I (COI) gene [30].

DNA extraction from mosquitoes

Mosquitoes were sorted by species per trap and by sex and then pooled in 2 mL o-ring tubes, with a

maximum of 15 individuals in each pool. A subset of *Ae. notoscriptus* mosquitoes were also screened

individually. Mosquitoe(s) were homogenised with 10 x 1.0 mm zirconia-silica beads (BioSpec Products),

with 597 µL of Buffer RLT and 2.8 µL of carrier RNA. Homogenisation was performed using a TissueLyser II

(Qiagen) at 30 oscillations/sec for 100 sec, repeated twice. Tubes were then centrifuged at 16,000 g for 3

mins. A 550 µL volume of supernatant was transferred into a 96 well deep well plate, with extraction

performed as is the protocol for the BioSprint 96 One-For-All Vet Kit (Qiagen). Every eleventh of twelve

wells in a 96-well plate was a blank DNA extraction control (seven in total) and a synthetic IS2404 positive

control was spiked into one of these seven wells to act as a positive extraction control. Extraction was

performed on a KingFisher™ Flex Magnetic Particle Processors (Thermo Scientific).

DNA extraction from arthropods

Arthropods other than mosquitoes collected on YST or SO were removed from the sticky cards and placed

in 1.5 mL microtubes (Eppendorf). Insects were separated by family and trap location, and were pooled

with a maximum of 10 individuals from each family per 1.5 mL tube. Samples were extracted non-

destructively to allow species confirmation if positive detections occurred. DNA was extracted using the

ISOLATE II Genomic DNA Kit (Bioline). Briefly, 25 µL of Proteinase K and 180 µL of Lysis Buffer GL was added

to each tube with samples incubated overnight at 56°C. Following incubation, the insects were removed

and stored to allow for further morphological identification if required, with the DNA extraction completed

on the incubation solution as per manufacturer instructions.

Synthetic PCR positive control

A synthetic PCR positive control DNA molecule was designed to discriminate false positives due to

contamination with positive control DNA versus the authentic IS2404 amplicon. The synthetic positive

control was designed to have an amplicon size of 120 bp to easily differentiate from a true IS2404 PCR

positive (59 bp) [16]. The synthetic positive was added at the DNA extraction stage on all 96-well plates, as a

positive control for this step and for the subsequent qPCR. The additional DNA sequence used to construct

the synthetic positive control was randomly selected from a DNA sequence unlikely to be in the laboratory,

in this case, Irrawaddy Dolphin (MK032252).

The synthetic positive control had the sequence: 5' –

TCCT<u>AAAGCACCACGCAGCATCT</u>ATCGCGAGCTTAATCACCATGCCG<u>CGTCCAACGCGATC</u>CCCGCTCGGCAGGGATC

CCTCTTCTCGCACCGGGCCA<u>CAATCCACTGGGGTCGCT</u>ATGA – 3' and was synthesised as an ssDNA oligo

(Sigma-Aldrich). The synthesised IS2404 synthetic positive was resuspended in nuclease-free water and

diluted to 0.001 pM, with 2 µL being used for extraction and positive controls. To confirm the presence of a

true positive as opposed to contamination, 5 µL of the qPCR product was added to 1 µL of DNA Gel Loading

Dye 6X (Thermo Scientific™) and run on a 2 % agarose gel (TopVision Agarose Tablets, Thermo Scientific™),

with 1% SYBR Safe DNA Gel Stain (Invitrogen™). The size of any positive IS2404 detection was assessed

against 2 µL of 100 bp DNA Ladder (Promega) and run at 50 V for 1.5 hrs before being visualised with an EZ

Gel Documentation System (Bio-Rad). Before the screening of insects commenced, the synthetic positive

control for IS2404 was successfully designed and tested. By running the amplified PCR products on an

agarose gel with the synthetic positive control and a real positive control, visual differentiation between a

synthetic positive occurring at 120 bp and a true positive at 59 bp (Figure S1).

Screening insects by qPCR for *M. ulcerans*

The qPCR screening was performed using three independent assays IS2404, IS2606 and KR [16]. All samples were first screened with the IS2404 qPCR; if a positive was detected, additional confirmation was attempted with IS2606 and KR qPCR assays. Reactions were performed using 7.5 µL TaqMan™ Fast Universal PCR Master Mix (2X), no AmpErase™ UNG (Applied Biosystems™), 1 µL of the primer-probe mix, 2 µL of DNA and 4.5 µL of nuclease-free water. A final primer-probe concentration for the IS2404 assay was as follows: 250:650:450nM for the forward primer, reverse primer and probe and 800:800:220nM for the forward primer, reverse primer and probe for the IS2606 and KR assay. A 2 µL volume of the synthetic positive control was added for the IS2404 reactions, whereas 2 µL of *M. ulcerans* DNA was used for IS2606 and KR. All reactions included six no-template extraction controls and were run in a 96 well plate format. Cycling conditions were as follows: denaturation at 95°C for 2 mins followed by 45 cycles at 95°C for 10 sec and 60°C for 30 sec, with qPCR performed on a QuantStudio™ 5 Real-Time PCR System (Applied Biosystems™). Positives were classified as reactions that produced a cycle quantification (Cq) value less than 40. Data were analysed using the QuantStudio™ Design and Analysis Software v1.4.3 with the Delta Rn threshold set at 0.04 for IS2404 and IS2606 and Delta Rn threshold of 0.1 for KR. The maximum likelihood estimate (MLE) per 1,000 mosquitoes tested (bias corrected MLE for point estimation of infection rate and a skew-corrected score confidence intervals) was calculated from the pooled samples as described [31]. Fisher's Exact test for assessing the significance of differences in IS2404 PCR positivity between mosquito species was calculated in R 4.0.2 [32].

All qPCR screening was performed blind with mixed-species 96-well plates. The synthetic positive control was added at the DNA extraction phase to one well of each plate and to qPCR plates to check that both extraction and qPCR detections were successful. All no-template controls (extraction and qPCR stage) were checked to ensure they remained negative, and that synthetic positive controls were detected for both the DNA extraction and qPCR stage in each run. Positive samples were run on agarose gels to confirm they were true positives and not contamination from the synthetic positive control.

An IS2404 qPCR standard curve was prepared using 10-fold serial dilutions of *M. ulcerans* genomic DNA,

with quadruplicate testing of each dilution. The DNA was extracted from *M. ulcerans* JKD8049 as described,

and quantified using fluorimetry (Qubit, dsDNA HS ThermoFisher Scientific) [22]. A limit-of-detection was

defined as the lowest dilution that returned a positive signal for all four replicates. Genome equivalents

were calculated based on the estimated mass of the *M. ulcerans* genome of 5.7 femtograms [22]. IS*2404* cycle

threshold (Ct) values were converted to genome equivalents (GE) to estimate bacterial load within a

sample by reference to a standard curve (r^2 = 0.9956, y = [-3.829Ln(x)+37.17]*Z, where y = Ct and x =

amount of DNA [fg] and Z = the dilution factor]) (Figure S2). An IS2404 qPCR standard curve was fitted using

non-linear regression in GraphPad Prism (v9.5.1).

Mycobacterium ulcerans genome sequencing

Whole-genome sequencing was performed directly on DNA extracted from selected PCR positive mosquito

samples and possum excreta specimens using a hybridisation capture approach, based on 120 nucleotide

RNA baits spanning the 5.8 Mbp chromosome of the the *M. ulcerans* JKD8049 reference genome

[Bioproject ID: PRJNA771185] (Datafile S1) (SureSelect Target enrichment system, Agilent, Santa Clara, CA,

USA) and the Illumina *Nextera Flex for Enrichment with RNA Probes* protocol [33]. Resulting sequence reads

were submitted to NCBI GenBank and are available under Bioproject PRJNA943595 (Table S2).

Mycobacterium ulcerans phylogenetic analysis

To compare genomic variations between clinical isolates and sequence capture datasets, we mapped the

sequence reads against a fully assembled *M. ulcerans* chromosome reconstructed from a Victorian clinical

isolate (JKD8049; GenBank accession: NZ_CP085200.1) using the Snippy pipeline (v4.4.5)

(https://github.com/tseemann/snippy). While standard parameters, including a minimum coverage of 10x

were used for the clinical isolates and single possum sequence capture dataset, the mosquito sequence

capture datasets had lower coverage, necessitating the adjustment of parameters. Thus, we lowered the

minimum coverage threshold to 1x to facilitate SNP calling for the mosquito sequence capture datasets.

The resulting SNPs were combined with those from the clinical isolates to generate an alignment of core

genome SNPs, which was used to infer a maximum likelihood tree with the GTR model of nucleotide

substitution using FastTree (v2.1.10) [34].

Aedes notoscriptus typing and species confirmation sequencing

Mosquito genotyping was performed by sequence comparisons of a partial fragment of the cytochrome c

oxidase subunit I (COI) gene [30].   DNA was extracted using the above protocols. PCR was performed using 5

µL of 5x MyFi Reaction Buffer, 1 µL of MyFi DNA polymerase, 5 µL of DNA, and primer concentrations as

described [35], with reaction made up to 25 µL with nuclease-free water. Reaction conditions were as follows

for COI, initial denaturation at 95 °C for 1 min, followed by 35 cycles at 95°C for 20 sec, 46°C for 20 sec, and

72°C for 60 sec before a final extension at 72°C for 5 mins. A 5µL volume of the amplified PCR product was

added to 1 µL of DNA Gel Loading Dye 6X (Thermo Scientific[TM]) and run on a 1 % agarose gel (TopVision

Agarose Tablets, Thermo Scientific[TM]), with 1% SYBR Safe DNA Gel Stain (Invitrogen[TM]). 2 µL of 100 bp DNA

Ladder (Promega) was added to confirm amplicons size and run at 100 V for 45 mins. PCR products that

produced bands of the correct size were purified using the ISOLATE II PCR and Gel Kit (Bioline) as per

manufacturer's protocol and submitted for sequencing using an Applied Biosystems 3730xl capillary

analyser (Macrogen), with sequencing occurring on both strands.

Sequences were analysed in Geneious Prime (v2019.2.1) and trimmed to high-quality bases, aligned using

ClustalW v2.1 and trimmed to a consensus region, *Ae. notoscriptus* COI (874 bp) and for species

identification COI (816-882 bp). Sequences were analysed using blastn against the NCBI database. COI

sequences generated for species identification are available under accession numbers OQ600123-4 and COI

sequences for *Ae. notoscriptus* phylogenetics under accession numbers OQ588831-67 (Table S2).

Mosquito bloodmeal analysis

Ninety blood-fed mosquitoes were identified as having an engorged abdomen and stilling having a red pigment indicating a fresh bloodmeal and were dissected with a sterile scalpel blade. Blood from the dissected abdomen was absorbed onto a 3 x 20 mm piece of a Whatman FTA™ card (Merck) and placed in a 2 mL tube. DNA was extracted from the FTA™ card using an ISOLATE II Genomic DNA Kit (Bioline) with a pre-lysis in 180 µL of Lysis Buffer GL and 25 µL Proteinase K for 2 hours before completing the extraction as per manufacturer's protocol. Extracted DNA was amplified for Cyt *b* using primers previously described (Townzen et al. 2008), with MyTaq™ HS Red Mix (Bioline), thermocycling conditions used were as follows: 95°C for 1 min, 30 cycles of 95°C for 15 secs, 50°C for 20 sec, 72°C for 20 sec, with a final extension at 72°C for 2 mins. Negative extraction and negative PCR controls were included with each PCR reaction. 5 µL of the amplified products was run on a 1% agarose gel containing 0.1% SYBR Safe DNA gel stain (Invitrogen) and visualised with a G:BOX Syngene blue light visualisation instrument. If a band was visualised at ~480 bp the PCR product was purified using an ISOLATE II PCR and Gel Kit (Bioline).

PCR products were quantified using a Qubit (Invitrogen) Fluorometer with an HS dsDNA kit. Sequencing libraries were prepared using 10 ng of purified PCR product with a NEXTFLEX® Rapid XP DNA-Seq Kit (PerkinElmer) barcoded using NEXTFLEX® UDI Barcodes (PerkinElmer). As a result, 70 bloodmeal libraries were sequenced, as well as 6 PCR negative control and 3 extraction negatives on a NovaSeq 6000 (Illumina), with 2 Gb requested per sample.

Sequence data were analysed by identifying poor-quality reads using Rcorrector (Song & Florea, 2015) and removed with TrimGalore v0.6.5 (Krueger 2019). De novo assembly was performed on the remaining sequence reads using Trinity v2.8.6 (Grabherr et al., 2011). The resulting contigs were filtered to be between 400-480bp and analysed using BLASTN (Altschul et al. 1990) against the nucleotide database publicly available on the National Centre for Biotechnology Information (NCBI) website. The resulting hits were filtered to exclude those sequences which had < 97% sequence identity to the database. Contigs identified as a host bloodmeal were confirmed by mapping raw reads using BWA-MEM v0.7.17 (Li & Durbin, 2009). Sequence reads were submitted to Genbank (Table S2).

Mosquito phylogenetic analysis

Phylogenetic analysis was performed on trimmed consensus regions of *Ae. notoscriptus* COI. The substitution model was selected using jModelTest2 v2.1.10, with the topology taking the best of nearest neighbour interchange, subtree pruning and regrafting [36]. The most appropriate substitution model was selected based on the Akaike information criterion. Maximum-likelihood trees were constructed in PhyML v3.3.2 with 1,000 bootstrap replicates; the gamma distribution parameter was used to estimate rate variation across sites [37]. The Hasegawa-Kishino-Yano (HKY) substitution model was selected for the COI tree.

Geographical data acquisition and spatial cluster analysis

The population map was created in qGIS v2.18.20 [38], using a 1 km^2 population grid [39] with 2011 Victorian mesh block data. Since 2004, Buruli ulcer has been a notifiable condition in Victoria, requiring Health Department reporting by doctors and laboratories. De-identified case notification data of Buruli ulcer patients who had laboratory-confirmed *M. ulcerans* infection and who lived on the Mornington Peninsula during the years 2019-2020 were provided by the Victorian Department of Health. The cases were defined as patients with a clinical lesion that was diagnosed using IS2404 qPCR and culture [16]. To conduct high-resolution spatial analyses, the data were aggregated at the mesh block level, the smallest geographical census units which typically contain 30-60 dwellings. The 2011 Victorian mesh block boundaries and the Victorian mesh block census population counts datasets were obtained from the Australian Bureau of Statistics (ABS) website [40]. The datasets were joined using the unique mesh block IDs using the QGIS v.3.16.7 geographic information system software [41]. The latitude and longitude (projected in GDA94) were derived from the centroids of the mesh block polygon. The dataset was then down-sampled to include only the Mornington Peninsula study area, specifically the Point Nepean and Rosebud-McCrae ABS level 2 Statistical Area.

SaTScan version 10.1.0 [42] was utilised to identify spatial clusters among trapped mosquitoes positive for *M.*

ulcerans, possum excreta positive for *M. ulcerans*, and human Buruli ulcer cases. The software searches for

instances where the observed number of spatial incidences exceeds the expected number within a circular

window of varying size across a defined study area. A log likelihood ratio statistic is calculated for each

window by comparing the number of observed and expected cases inside and outside the circle against the

assumption of randomly distributed cases. In addition to the most likely cluster, there are usually secondary

clusters with almost as high likelihood that substantially overlap with the primary clusters. These secondary

clusters can be indicative of sub-clusters within the primary cluster or potentially distinct clusters that are

spatially adjacent to the primary cluster. The Mornington Peninsula surveillance data used in these analyses

consisted of trapped mosquitoes (177 traps screened for IS2404 collected: 12/11/19 to 20/03/20), *M.*

ulcerans detected in possum excreta collected during the summer (December to February) of 2019 using

data from a previous study [43] and notified human Buruli ulcer cases from the study area in the years 2019-

2020. The use of possum excreta collected during the summer was appropriate as Buruli ulcer transmission

is most likely to occur during that time of year [44,45]. For each of the three data sources (trapped mosquitoes,

possum excreta, and human cases), the null hypothesis assumes that *M. ulcerans* detections or Buruli ulcer

cases are uniformly distributed across the study area, where the alternative hypothesis suggests that there

may be certain locations with higher rates than expected if the risk was evenly distributed. Primary and

secondary clusters were accepted only if the secondary clusters did not overlap with previously reported

clusters with a higher likelihood. Given that the trapped mosquito and possum excreta IS2404 PCR results

were binary (positive or negative), a Bernoulli model was used to scan for spatial clusters, with the

maximum cluster size set to 50% of the population size. The human Buruli ulcer case data aggregated at the

mesh block level varied in number, with some mesh blocks having zero cases and others having one or

more. We applied the Poisson probability model to the notified Buruli ulcer case counts, using a

background population at risk that was derived from the 2011 population census. The maximum cluster

size was limited to 14,481 individuals, 10% of the total population at risk. To determine the likelihood of a

triple-cluster overlap between the three SaTScan analyses occurring by chance, we conducted a

permutation test (https://github.com/abuultjens/BU-3-way-SatScan). In each of 10,000 iterations, the

geographical coordinates for each variable were randomly shuffled. The number of SaTScan clusters with

triple overlap was determined using the 'sf' package [46] in the R statistical programming language [47].

Results

Mosquito surveys of the Mornington Peninsula

A primary goal of this research was to test the hypothesis that mosquitoes are associated with *M. ulcerans*

transmission, as reflected by IS2404 PCR positivity at a certain frequency in areas of the Mornington

Peninsula with human Buruli ulcer cases. A total of 73,580 mosquitoes were collected, consisting of 72,263

females (Table 2) and 1,317 males (Table S2). The majority (90%) of these insects were collected in the

large survey of 2019-2020 (Datafile S2). Across all five surveys, 26 different mosquito species were collected

covering six genera (Table 2). The most dominant species identified during the 2019-2020 survey was *Cx.*

molestus accounting for 42% of mosquitoes, followed by *Ae. notoscriptus* (35%) and *Cx. australicus* (8%).

Twenty-three other species comprised the remaining 15% of mosquitos. The distribution of the two

dominant species across the survey area is shown in Figure 2. This mapping revealed an asymmetric

distribution of each species, with *Ae. notoscriptus* dominant to the eastern end and *Cx. molestus* dominant

towards the western end of the Peninsula (Figure 2).

Figure 2. Dominant mosquito species distribution across the Mornington Peninsula. *Map showing the proportional distribution of the two dominant mosquito species trapped during 2019/2020. The pie charts are an aggregation of the 180 different trap sites. Trap groups containing mosquitoes that were PCR positive for M. ulcerans are also indicated. Shown too, are the meshblock statistical areas, with those in red containing at least one human* Buruli ulcer *case diagnosed in 2019-2020.*

M. ulcerans *PCR positive mosquitoes were predominantly* Aedes notoscriptus

The IS2404 qPCR assay was used to infer the presence of *M. ulcerans* in association with mosquitoes. Of the 73,580 mosquitoes trapped across all years, 18,610 (25%) were screened by IS2404 qPCR for the presence of *M. ulcerans* (Table 2). *Mycobacterium ulcerans* qPCR positives were observed in 53 pools or individuals of *Ae. notoscriptus*, with detections occurring in each year across all survey events (Figure 2, Table 2). Only one other mosquito species tested IS*2404* positive, which was a two-insect pool of *Ae. camptorhynchus* (Table 3). The positive association between *M. ulcerans* and *Ae. notoscriptus* compared to other

mosquitoes, in particular the other abundant mosquito species, *Culex molestus*, was highly significant (P

<0.0001, Fisher's exact test).

Subsets of *Ae. notoscriptus* were screened individually to better estimate the prevalence of *M. ulcerans*

positive mosquitoes in this species. Individually tested mosquitoes included all of the 2017/2018 collection

(367/367 individuals with 8 positives), 448/4,330 of the 2019/2020 collection with 3 positives, and all 1,247

individuals of the 2021 collection with 7 positives (Table S3). Thus, based on screening individual insects,

18/2,062 (0.87%) *Ae. notoscriptus* were IS2404 PCR positive. All other *Ae. notoscriptus* and other mosquito

species in this study were screened in pools (up to 15 insects per pool) with 24/747 *Ae. notoscriptus* pools

(3%), PCR positive (Table S4).

Table 2. Female mosquitoes trapped on the Mornington Peninsula and screened by IS2404 PCR.

Species	No. of mosquitoes/No. mono-species pools screened for *M. ulcerans* (No. pools positive for *M. ulcerans*, IS*2404**)					
	Dec 2016 - Apr 2017	Nov 2017- May 2018	Dec 2018- May 2019	Nov 2019- Mar 2020	Feb 2021- Mar 2021	Total
Ae. alboannulatus	9/9	4/4	3/3	183/96	21/21	220/133
Ae. australis	0	0	0	1/0	0	1/0
Ae. bancroftianus	0	0	0	3/2	0	3/2
Ae. camptorhynchus	258/258	5/5	0	121/64 **(1)**	8/8	392/335 **(1)**
Ae. clelandi	0	0	0	41/32	0	41/32
Ae. flavifrons	1/1	0	0	31/26	13/13	45/40
Ae. imperfectus	0	0	0	266/226	0	226/226
Ae. notoscriptus	174/174 **(2)**	367/367 **(8)**	1,793/1,779 **(15)**	23,360/4,330 **(14)**	1,247/1,247 **(7)**	26,941/7,897 **(46)**
Ae. rubrithorax	9/9	27/27	230/230	1,402/335	45/45	1,713/646
Ae. sagax	0	0	0	2/2	0	2/2
Ae. theobaldi	0	0	0	1/1	0	1/1
Ae. vigilax	0	0	0	2/0	0	2/0
Ae. vittiger	0	0	0	2/0	0	2/0
An. annulipes	1/1	24/24	4/4	751/376	1/1	781/406
An. atratipes	0	0	0	2/0	0	2/0

Cq. linealis	89/89	142/142	2/2	1,254/258	11/11	1,498/502
Cu. inconspicua	0	0	0	2/2	3/3	5/5
Cx. annulirostris	0	3/3	0	42/1	0	45/4
Cx. australicus	3/3	198/198	17/17	4,910/534	33/33	5,161/785
Cx. cylindricus	0	0	2/2	160/81	0	162/83
Cx. globocoxitus	0	24/24	0	3,207/854	9/9	3,240/887
Cx. molestus	61/61	29/29	384/377	28,107/3,308	1,465/1,465	30,046/5,240
Cx. orbostiensis	0	0	0	131/69	0	131/69
Cx. quinquefasciatus	3/3	257/257	37/37	1,098/401	163/163	1,558/861
Tp. atripes	0	0	0	18/0	12/12	30/12
Tp. tasmaniensis	0	0	0	15/13	0	15/13
Total	**608/608 (2)**	**1,080/1,080 (8)**	**2,512/2,451 (15)**	**65,112/11,011 (15)**	**3,031/3,031 (7)**	**72,263/18,181 (47)**

* pool size ranged from 1 individual up to 15 individuals

Of the 46 *Ae. notoscriptus* that tested positive for IS2404, 26 (56%) were confirmed positive for IS2606 and

31 (67%) for KR, with 24 (52%) pools positive by all three qPCR assays (Table S3). Of the 46 IS2404 positive

Ae. notoscriptus, eight (17%) pools were not tested using the IS2606 assay and one (2%) pool using the KR

assay due to limited template DNA available. The average Ct value for IS2404 positive *Ae. notoscriptus* was

36.00 (range 29.74-39.65). The average Ct value for IS2606 was 37.73 (range 32.49-45.00), and the average

Ct value for KR was 34.19 (range 28.52-39.26). With reference to an IS2404 qPCR standard curve (Figure

S2), we estimated the *M. ulcerans* burden per mosquito. The mean bacterial genome equivalents (GE) per

insect was 294 GE (range: 11 – 4200) (Table S3). The MLE of estimated infection rate for all *Ae. notoscriptus*

was 5.88 based on the IS2404 *M. ulcerans* qPCR-positive mosquitoes per 1,000 tested (95% CI 4.37-7.76) for

all *Ae. notoscriptus* tested over the years and 2.96 for *Ae. camptorhynchus* (95% CI 0.17-14.19) (Table 3).

IS2404 PCR positive Ae. notoscriptus and possum excreta yield M.

ulcerans genomic SNP profiles that match human patient isolates

Genomic epidemiological studies have shown there are unique SNP signatures associated with *M. ulcerans*

clinical isolates from specific endemic areas in southeastern Australia [48]. To test if the *M. ulcerans* genotype

present in mosquitoes from our study area matched that found in possums and human Buruli ulcer cases in

the same region, we conducted genome sequence enrichment using *M. ulcerans* custom RNA baits

(Datafile S1). This method facilitates whole genome sequencing of target pathogen genomes in complex

microbial samples or when competing background DNA precludes sequencing the target directly. As

reported in other pathogen sequence enrichment studies, we observed decreasing genome sequence

recovery with decreasing pathogen load, as indicated by increasing IS2404 Ct values (Figure 3A) [49]. While

DNA sequence reads were obtained from three mosquitoes, only DNA from one *Ae. notoscriptus* extract

(sample ID: 5675, Ct 32.66) had a high enough concentration of *M. ulcerans* DNA to yield sufficient reads

for SNP-calling following read mapping. Nevertheless, the fact that sequence reads mapped across the

length of the *M. ulcerans* chromosome for all three insects, despite low coverage for two of them, confirms

the presence of the pathogen on (or in) *Ae. notoscriptus* (Figure 3B). We also included sequence capture

enrichment results for two IS2404 positive possum excreta specimens, collected as part of a large field

survey of *M. ulcerans* in Australian native possums in the region [43]. More than 500,000 reads were obtained

from these samples (primary IS2404 Ct values <23). We then inferred a phylogeny, using a 5.6Mbp core-

genome alignment that identified 23 SNP positions from mapping the reads from these possum excreta

samples, the *Ae. notoscriptus* sample (ID: 5675) and a reference panel of sequence read sets from 36

published *M. ulcerans* genomes from southeastern Australia [48]. These 36 genomes were selected because

they span the population structure of *M. ulcerans* in this locale (Table S1). The resulting phylogeny (tree

tips aligned by sample/isolate geographic origin) showed possum and mosquito *M. ulcerans* genotypes co-

clustering with each other and with human *M. ulcerans* isolates from the Mornington Peninsula (north)

area, with no SNP differences between them.

Figure 3. Comparison of M. ulcerans genome from human Buruli ulcer cases compared with sequences

recovered from possum excreta and mosquitoes on the Mornington Peninsula. *(A) Summary of IS2404*

qPCR screening of primary samples and the sequence capture libraries pre- and post-enrichment. Note that

possum excreta were enriched as barcoded sequence library pools so share the same pre- and post-

enrichment Ct values; (B) Artemis coverage plots showing sequence capture reads mapped to the M.

ulcerans JKD8409 chromosome from possum excreta samples and three qPCR positive mosquitoes; (C)

Maximum-likelihood phylogeny inferred from an alignment of 23 core-genome SNPs using reads mapped to

the JKD8049 reference chromosome, and with tips aligned with environmental sample or patient origin.

Dataset includes reads from the five environmental samples listed in panel (A) and a reference collection of

36 M. ulcerans *genomes representing the genomic diversity of the bacterial population in southeastern*

Australia [48]. *The shortest vertical branch length represents a single SNP difference.*

Genetic characterisation of Ae. notoscriptus *collected on the* Mornington Peninsula

To assess if *M. ulcerans* presence was associated with a particular clade of *Ae. notoscriptus* on the

Mornington Peninsula we compared COI gene sequences for 18 *M. ulcerans* positive (confirmed by all

IS2404, IS2606 and KR qPCR assays) *Ae. notoscriptus* and 19 *M. ulcerans* negative *Ae. notoscriptus* (Figure

2). Based on the COI phylogenetic tree, *Ae. notoscriptus* from the Mornington Peninsula spanned the three

previously identified clades and there was no association between *M. ulcerans* and a particular mosquito

lineage (Figure S3) [35].

Table 3. The number *of M. ulcerans* **positive pools tested by insertion sequence IS2404 qPCR and**

Maximum likelihood estimate (MLE) of *M. ulcerans* **per 1,000 mosquitoes trapped in the Mornington**

Peninsula.

Species	No. mosquitoes	No. positive/No. pools tested	MLE‡	95% Confidence interval
Ae. alboannulatus	133	0/97	0	0-27.71
Ae. bancroftianus	2	0/2	0	0-657.62
Ae. camptorhynchus	335	1/85	2.96	0.17-14.19
Ae. clelandi	32	0/3	0	0-67.97
Ae. flavifrons	40	0/16	0	0-78.10
Ae. imperfectus	226	0/27	0	0-14.96
Ae. notoscriptus	7897	46/2937	5.88	4.37-7.76
Ae. rubithorax	646	0/220	0	0-5.84

Ae. sagax	2	0/1	0	0-545.52
Ae. theobaldi	1	0/1	0	0-793.45
An. annulipes	406	0/148	0	0-9.16
Cq. linealis	502	0/139	0	0-7.43
Cu. Inconspicua	5	0/4	0	0-408.11
Cx. annulirostris	4	0/3	0	0-450.75
Cx. australicus	785	0/150	0	0-4.76
Cx. cylindricus	83	0/38	0	0-41.60
Cx. globocoxitus	887	0/150	0	0-4.22
Cx. molestus	5240	0/609	0	0-0.73
Cx. orbostiensis	69	0/9	0	0-42.95
Cx. quinquefasciatus	861	0/237	0	0-4.39
Tp. atripes	12	0/9	0	0-230.35
Tp. tasmaniensis	13	0/10	0	0-215.26

450 ‡MLE bias was corrected when ≥ 1 pool was positive.

IS2404 PCR screening of arthropods other than mosquitoes

We also investigated the association of *M. ulcerans* with arthropods other than mosquitoes in the region

using yellow sticky traps (YST) and sticky ovitraps (SO). A total of 21,000 specimens were collected and

sorted from 278 YST and 33 SO traps. We were able to classify 2,696 as insects that may bite or pierce

human skin (Table 4). Flies were the largest group collected on the sticky traps. YSTs collected more insects

than the SOs, but this was proportional to the number of traps set (Table 4). Of the 2,696 insects screened

by PCR, only two flies tested positive for *M. ulcerans* (Table S3). Both flies had high Ct values for IS2404 (Ct

35.92 and 37.54) and each sample was only confirmed for either IS2606 or KR assay (Table S3), but not

with all three assays. Both insects were blowflies (*Calliphora hilli* Patton) based on morphology and

sequencing of the COI region with sequences having >99.86 nt identity to *C. hilli* (Table S1).

Table 4. Number of arthropods collected on Yellow Sticky Traps (YST) and Sticky Ovitraps (SO) on the

Mornington Peninsula that might bite or pierce human skin.

Trap	Ants	Biting midges	Flies	Mosquitoes	Spiders	Ticks	Wasps	Total
YST	28	58	1,562	10	76	21	562	2,317
SO	70	1	144	11	141	4	8	379
Total	98	59	1,706	21	217	25	570	2,696

Mosquito bloodmeal analysis

Of the mosquitoes collected, a proportion of individuals identified as being engorged (bloodfed) were PCR

screened and the resulting amplicon sequenced for Cytochrome B (CytB) to identify host bloodmeal

sources. A total of 90 individual engorged mosquitoes were extracted, with 70 DNA preparations producing

high quality amplicons that were of sufficient concentration to permit Illumina amplicon sequencing. After

quality filtering, 36 individuals were identified as having had a recent bloodmeal: 14 *Cx. molestus*, 13 *Ae.*

notoscriptus, 2 *Ae. rubrithorax*, 2 *Cx. globocoxitus*, 2 *Cx. quinquefasciatus*, 2 *Cq. linealis* and 1 *Cx.*

australicus. Of the bloodmeals detected, common ringtail possum was the most commonly identified with

20 detections across the 36 samples, followed by 17 blackbirds, 13 humans, 11 red wattle birds, and 5 little

wattle birds; a further 16 bloodmeals identified 10 minor host species (Figure 4, Figure S4). Dual

bloodmeals were commonly identified, with 55% (20/36) of individuals having more than one bloodmeal

identified. Additionally, two mosquitoes had evidence of three different bloodmeal sources within an

individual insect. Three individuals (2 x *Ae. notoscriptus* and 1 x *Ae. rubrithorax*) were also identified as

having dual bloodmeals from ringtail possum and human origins (Figure 4).

Figure 4. Mosquito blood meal analysis. *Summary of cytochrome B (Cytb) gene sequences from the 36*

bloodfed mosquitoes to identify host bloodmeal sources . A blue circle indicates positive for host blood

source; the larger the circle, the more individual mosquitoes with an identical bloodmeal profile. Red boxes

indicate individual mosquitoes which have dual bloodmeals for both humans and ringtail possum sources.

Spatial clustering links M. ulcerans *positive mosquitoes with positive possum excreta and human Buruli ulcer cases*

In our previous research, we demonstrated a spatial association between possum excreta containing *M. ulcerans*, as detected by structured surveys, and clusters of human Buruli ulcer cases [50]. To expand on this finding, we investigated whether further spatial clustering associations could be detected by analysing qualitative qPCR (IS2404) data of trapped mosquitoes using SaTScan. Here, three separate analyses were conducted: (i) human Buruli ulcer case data from individuals reported to have acquired the disease in the study area during 2019-2020, (ii) qPCR data from possum excreta collected in a previous investigation (2018-2019), and (iii) qPCR data from trapped mosquitoes (2019-2020). These analyses identified a single mosquito cluster, four possum clusters and six human Buruli ulcer clusters (Figure 4A-B). Notably, one human Buruli ulcer cluster and two possum excreta clusters had higher numbers of observed Buruli ulcer cases or *M. ulcerans* detections, respectively, than expected if uniformly distributed (p-value <0.05) (Figure 4B). Importantly, the analyses revealed an instance of triple cluster overlap (mosquito/possum excreta/human) in the Mornington Peninsula suburb of Rye, where all three SaTScan analyses had an overlapping cluster (Figure 4A).

	CLUSTER	P_VALUE	OBSERVED	EXPECTED	LLR
🦟	M1	0.915	2	0.26	4.16
	M2	0.472	3	0.52	4.17
👪	H1	2.06E-05*	31	9.35	17.75
	H2	0.148	13	3.56	7.80
	H3	0.265	15	4.95	7.03
	H4	0.702	4	0.42	5.53
	H5	0.820	4	0.46	5.14
	H6	0.999	2	0.12	3.74
🐀	P1	9.21E-05*	41	17.54	17.05
	P2	0.002*	9	1.350	13.58
	P3	0.804	4	0.562	6.42
	P4	0.997	3	0.450	4.46

Figure 4. *SaTScan analyses of trapped mosquitoes, possum excreta, and human* Buruli ulcer *cases on the*

Mornington Peninsula, revealing spatial clustering. A) Map illustrating the clusters identified by the three

separate SaTScan analyses: (I) trapped mosquitoes (177 traps screened for IS2404 collected: 12.11.19 to

20.03.20), (ii) M. ulcerans detected in possum scats collected during the summer of 2019 (Dec 2018 – Feb

2019) using data from a previous study, and (iii) notified human Buruli ulcer *cases from the study area in*

the years 2019-2020. The zoomed insert highlights an instance where all three analyses had overlapping

clusters in the suburb of Rye. B) Table summarising the SaTScan results for all identified clusters. Log

likelihood ratio is abbreviated as "LLR." Clusters with p-values <0.05 are marked with asterisks.

The results of our permutation test to explore the probability of these three categories (mosquitoes,

possums and humans) overlapping showed that a triple-cluster overlap occurred with randomly rearranged

location labels at a low frequency of 8.9%. This analysis adds support to a causal relationship between the

presence of *M. ulcerans* in possums and mosquitoes and humans contracting Buruli ulcer in the

Mornington Peninsula suburb of Rye.

Discussion

Using insect field surveys augmented with genomics we have addressed *Barnett Criteria* 1 – 3 (Table 1) to

add to a hierarchy of evidence indicting mosquitoes as principal vectors in the spread of *M. ulcerans* from

the environment to humans [3,15,17,21,22]. Our survey area was the Mornington Peninsula, where human Buruli

ulcer cases have increased progressively over the past 20 years [12]. Our key findings were that *M. ulcerans*

was almost exclusively associated with one mosquito species in this region, *Aedes notoscriptus*, a

mammalian host feeder, at a frequency of 0.87%. The estimated pathogen burden in each insect (Table S3)

was consistent with the reported low infectious dose for *M. ulcerans* [22,51]. Pathogen genomics provided

confirmation of a linked transmission chain between mosquitoes, native possums and humans in this

region. The metabarcoding bloodmeal analysis of trapped mosquitoes revealed dual feeding by individual

mosquitoes on both possums and humans, providing an example of a potential transmission pathway

between infected possums and humans. Finally, spatial clustering analysis showed a striking overlap

between clusters of possums shedding *M. ulcerans*, mosquitoes harbouring *M. ulcerans* and human Buruli

ulcer cases, reinforcing the importance of possums and mosquitoes in the spread of *M. ulcerans* to humans.

Indications that invertebrates might be playing a role in the spread of Buruli ulcer first came from field

surveys in West Africa in the late 1990s, where surveys in Benin identified water bugs in the genera

Naucoris and *Diplonychus* as IS2404 PCR positive for *M. ulcerans* [15,16,52]. Subsequent culture isolation of *M.*

ulcerans from an aquatic insect in Benin (*Gerris* sp.) and potentially from *Naucoris* sp. in neighbouring Cote

d'Ivoire further suggested a role for water bugs in disease transmission, at least in west Africa [53,54]. A role

for mosquitoes in the transmission of *M. ulcerans* anywhere was first revealed from entomological surveys

in the mid-2000s on the Bellarine Peninsula and then more recently in Far North Queensland [15,21,55]. Other

observations that supported a role for mosquitoes in transmission included an analysis of the Buruli ulcer

lesion location on the human body, which included more than 600 human Buruli ulcer cases, and showed a

preponderance of lesions on ankles, elbows and backs of legs. These areas are frequently exposed and

insects may preferentially bloodfeed there [56]. Laboratory studies under one transmission scenario have also

shown the competence of *Ae. notoscriptus* as mechanical vectors of *M. ulcerans* (Barnett criteria No. 4,

Table 1) [22].

Of the ~18,600 mosquitoes tested by IS2404 PCR from five different survey periods in the present study,

Ae. notoscriptus was consistently positively associated with *M. ulcerans*. Where and how *M. ulcerans* is

contaminating (or infecting) these mosquitoes remains to be determined. However, we made the

somewhat unexpected observation that the most abundant mosquito species trapped on the Mornington

Peninsula, *Culex molestus,* which predominately feeds on avian species [57] was consistently IS2404 PCR

negative. This observation might be explained by a difference in the ecology of these two mosquito species,

such as how (or where) they are encountering *M. ulcerans.* The apparent *Ae. notoscriptus*/*M. ulcerans*

tropism might also indicate a specific biological interaction between this insect and the mycobacterium, or

alternatively, a *Cx. molestus* antagonism to mycobacterial carriage.

MLE is a parameter used in vector ecology to estimate the proportion of infected mosquitoes that are

pathogen positive, based on a pooling-based screening assay, where the proportion is the parameter of a

binomial distribution [58] . The MLE of *Ae. notoscriptus* over all 5 years of our surveys was 5.88 *M. ulcerans*

positive mosquitoes/1,000 tested (95% CI 4.37-7.76). The MLE for the only other positive mosquito species,

Ae. camptorhynchus (consisting of two insects that tested positive for *M. ulcerans*), was 2.96/1,000 tested

(95% CI 0.17-14.19). These estimates are consistent with those previously reported for *Ae. camptorhynchus*

(10,558 mosquitoes tested, MLE 3.98 per 1,000) and *Ae. notoscriptus* (221 mosquitoes tested, MLE of 4.47

per 1,000) during the Bellarine Peninsula survey of that Buruli ulcer endemic area [15]. Interestingly, these

values are significantly higher than the MLE values from an *M. ulcerans* mosquito survey conducted in

tropical far north Queensland, Australia which reported an MLE value of 0.13 /1,000 (95% CI 0.01–0.61) for

different species of mosquito [55]. To note though, there were few reported cases of Buruli ulcer in this area

of Queensland during the field survey period, consistent with the suggestion that mosquito surveys for *M.*

ulcerans will be useful for predicting Buruli ulcer risk in humans.

In tropical far north Queensland, Australia, host-seeking mosquitoes that harboured *M. ulcerans* collected

through similar carbon-dioxide light traps to those used in this study were within the genera *Verallina,*

Coquillettidia and *Mansonia* [55]. These comparisons highlight that the mosquitoes encountering *M. ulcerans*

appear to be region/environment specific, and these observations support the idea that mosquitoes are

mechanical vectors of *M. ulcerans*. The major mosquito species associated with detection of *M. ulcerans* in

Victoria, *Ae. notoscriptus, Ae. camptorhynchus, An. annulipes* and *Cq. linealis,* are all diverse opportunistic

feeders.

To explore host sources for mosquitoes trapped in our study, we used metagenomic amplicon sequencing

of engorged mosquitoes [59-64]. Of the 90 blood-fed mosquitoes processed, 36 individual insects successfully

had their host blood source identified. The 54 bloodmeals that were unidentified were probably due to

degradation of the host DNA that occurs approximately 36 hours post-feeding [65]. As the mosquitoes

analysed in this study were collected from baited mosquito traps, the exact time post-feeding they were

collected is unknown. Of the host blood sources identified, common ringtail possum was the most

commonly identified bloodmeal, followed by birds and humans, with dual bloodmeals observed in 55% of

mosquitoes (Fig. 4, Fig. S4). Most notably, three insects were determined to have bloodmeals from both

common ringtail possums and humans. These insects consisted of two *Ae. notoscriptus* and one *Ae.*

rubrithorax. Common ringtail possums are major wildlife reservoirs of *M. ulcerans* [11], and in this study we

identified *M. ulcerans* in frequent association with *Ae. notoscriptus*. The identification of these dual

bloodmeals provides evidence that *Ae. notoscriptus* mosquitoes are bloodfeeding between possums and

humans within a relatively short timeframe. Dual bloodmeals may provide an opportunity for a mechanical

transmission event of *M. ulcerans* from a possum with a Buruli ulcer teeming with *M. ulcerans,* from which

a mosquito bloodfeeds and then moves to a nearby human. However, the observed positive association

specifically between *Ae. notoscriptus* and *M. ulcerans* is not explained by our bloodmeal analysis, which

showed several other mosquito species that were not *M. ulcerans* positive, such as *Cx. molestus*, *Ae.*

rubrithorax and *Cq. Linealis,* also fed on possums. An alternative source of *M. ulcerans* acquisition for *Ae.*

notoscriptus might be from possum excreta contaminating the *Ae. notoscriptus* breeding sites, as this

species breeds in small artificial containers [59]. Previous studies have shown that mosquito larvae can ingest

M. ulcerans and test positive for *M. ulcerans* during the larval stage, but *M. ulcerans* is not detected in the

pupa or adult, which is largely thought to be a result of the larval midgut being purged [66]. However, if

possum excreta, which can have a high concentration of *M. ulcerans* [11], falls into these artificial breeding

sites, then this may provide a means to contaminate the adult *Ae. notoscriptus* during eclosion and

emergence on top of the water's surface.

In common with other arthropod-borne bacterial pathogens, *M. ulcerans* also has the genomic hallmarks of

niche adaptation [67,68]. For instance, the bacterial diseases vectored by blood-feeding arthropods, such as

bubonic plague, spread to humans through the bite of fleas harbouring the bacterium *Yersinia pestis* [69], the

tick-borne infections such as Lyme disease, Rocky Mountain Spotted Fever, ehrlichiosis (among others)

caused by *Borrelia burgdorferi*, *Rickettsia rickettsii* and *Ehrlichia* sp., respectively [70], or tularemia, caused by

infection with *Francisella tularensis* spread through the bite of infected ticks (with some subspecies also

spread by mosquitoes) [71-73], are all caused by bacteria that have degenerating genomes. That is, like M.

ulcerans, their genomes bear the distinctive hallmarks of evolutionary bottlenecks and niche-adaptation

(plasmid acquisition, pseudogene accumulation, insertion sequence expansion), a genomic pattern thought

indicative of a shared trajectory towards symbiosis with the arthropod host [74]. In addition, the absence of

recombination and highly clonal population structure of *M. ulcerans* is aligned with the population

structure of bacteria in 'closed symbiosis' with their host animal [75].

Mosquitoes were not the only arthropods from which *M. ulcerans* were detected in the Mornington

Peninsula. With over 20,500 arthropods collected using sticky traps (YST and SO), this study screened other

insects that might broadly act as a mechanical vector, that is, an insect that is involved in the accidental

transport of a pathogen [76 22]. *M. ulcerans* was successfully identified on two of ~1,800 flies tested, identified

as blowflies (*Calliphora hilli*) . Previous studies have screened other species of flies, such as March flies

(Tabanidae), with no positive detections [55]. *C. hilli* is a native species occurring along the Australian east

coast and in South Australia [77]. This species is a carrion breeder occurring year-round throughout Victoria,

with the female laying her eggs in decaying flesh where the larvae emerge [77]. Possum carcases are readily

infested by *C. hilli* [78], which may explain how this fly species is becoming contaminated with *M. ulcerans*, as

possums are wildlife reservoirs of *M. ulcerans* [10]. The likelihood of *C. hilli* carrying *M. ulcerans* to humans is

relatively low due to this species rarely biting humans [79].

Interestingly, all species that have tested positive for *M. ulcerans* share quite different breeding

environments and feeding preferences, although most feed on humans. *Aedes camptorhynchus* and

Verrallina sp. are saltmarsh mosquitoes [80,81], while *Ae. notoscriptus* is a freshwater container breeder

occurring in close association to humans [82]. *Anopheles annulipes* and *Culex australicus* are also freshwater

breeding mosquitoes [83], although *Culex australicus* typically feeds on avian species and is not considered to

feed on humans [84]. This diversity in mosquito species, habitats, flight range from breeding sites and host-

preference feeding raises interesting questions about how these mosquitoes are coming into contact with

M. ulcerans and how potential vector mosquito species may change based on the study area. Additionally,

the ecology of the mosquito sub-species are generally poorly understood, including *Ae. Notoscriptus,*

which occurs as a series of genetically differentiated clades within Australia [35].

In this study, we also deployed for the first time targeted enrichment genome capture using the Agilent

RNA bait system to demonstrate beyond reasonable doubt the presence of *M. ulcerans* in association with

mosquitoes [49]. The IS2404, IS2606 and KR PCR assays have provided robust evidence for the presence of *M.*

ulcerans in the environment, but detecting pathogen genomic DNA and using those sequences to

forensically dissect transmission chains by matching pathogen genome SNP profiles is a powerful

technological advance [49].   While the method lost sensitivity for IS2404 qPCR Ct values >32, we generated

sufficient genome coverage from one IS2404 PCR positive mosquito and from two possum excreta

specimens to show that the *M. ulcerans* genotypes of the captured genomes were identical to those

associated with human Buruli ulcer cases in the study area, rather than Buruli ulcer cases linked to other

regions (Fig. 3). These findings provide strong support for mosquitoes and possums playing a role in the

transmission of *M. ulcerans* in the Mornington Peninsula. These results also add to the growing body of

evidence demonstrating the utility of pathogen genomics in specifically identifying the sources and

transmission pathways of environmental pathogens and suggest that continued surveillance of *M. ulcerans*

genotypes in mosquitoes and possums could be useful in guiding public health interventions that seek to

control the spread of Buruli ulcer in the region. We are now exploring the use of genome sequence

enrichment to dissect the genomic epidemiology from the perspective of environmental sources, in

particular to track the spread of Buruli ulcer in Australian native possums across the region [50].

The SaTScan clustering analyses provide compelling evidence for the spatial association between the

presence of *M. ulcerans* in possums, mosquitoes, and human Buruli ulcer cases in the study area (Figure 4).

The triple-cluster overlap observed in our SaTScan analyses suggests that the spatial distribution of these

three factors is likely to be closely linked and may be contributing to the persistence of Buruli ulcer in the

Mornington Peninsula. Our finding of a low frequency of triple-cluster overlap among 10,000

randomisations, with only 8.9% of replicates showing this phenomenon, provides a level of confidence that

our observations have not occurred randomly. This further supports the validity of our findings and

reinforces the idea that the presence of *M. ulcerans* in possums and mosquitoes may play a crucial role in

the transmission of *M. ulcerans* to humans in the study area. Our findings highlight the importance of

ongoing surveillance of possum and mosquito populations, which may provide critical insights into the

epidemiology of Buruli ulcer in the region and inform targeted public health interventions to control the

disease.

Our research has some limitations. Although we found a spatial association between the presence of *M.*

ulcerans in possums and mosquitoes, and human Buruli ulcer cases, we cannot absolutely conclude

causality or directionality. In addition, the very specific set of circumstances that have led to the rise of

Buruli ulcer in temperate southeastern Australia restricts the generalisability of our results. One must be

cautious to draw parallels with African Buruli ulcer endemic countries for instance, where a highly

susceptible mammalian reservoir equivalent to the Australian native possum is yet to be identified. Further

investigations are needed to better understand the underlying mechanisms that drive the association

between the presence of *M. ulcerans* in these different species and the development of human Buruli ulcer

cases.

Despite these caveats, our collective research over more than 15 years makes it very clear that mosquitoes

are likely vectors and native possums are major wildlife reservoirs of *M. ulcerans* in southeastern Australia.

Mosquito surveillance with *M. ulcerans* screening coupled with mosquito control and public health

messaging to avoid mosquito bites are practical interventions that would be expected to reduce the

incidence of human Buruli ulcer.

Funding

TPS: National Health and Medical Research Council of Australia (GNT1152807, GNT1196396). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Acknowledgments

We would like to acknowledge the residents of the Mornington Peninsula for assisting in collecting and sending sticky cards to our laboratory as part of our surveillance and to Mary Tachedjian, Michael Dunn, Simone Clayton, Julie Gaburro and Victoria Boyd for assistance during field work.

Ethics

Ethical approval for the use in this study of de-identified human BU case location, aggregated at mesh block level, was obtained from the Victorian Government Department of Health Human Ethics Committee under HREC/54166/DHHS-2019-179235(v3), 'Spatial risk map of Buruli ulcer infection in Victoria.'.

Author Contributions

Conceptualisation PTM, JO, RF, SRC, PDRJ, JRW, AAH, KBG, TPS, SEL; Data curation PTM, KB, JO; Formal Analysis PTM, AHB, TPS; Funding acquisition SRC, PDRJ, KBG, JRW, AH, KG, TPS, SEL; Methodology PTM, AHB, KB, JCC, JRW, TPS, SEL; Writing – original draft PTM, AHB, SEL, TPS; Writing – review & editing all authors.

References

1   MacCullum, P., Tolhurst, J. C., Buckle, G. & Sissons, H. A. A new Mycobacterial infection in man. *J Pathol Bacteriol* **60**, 93-102 (1948).
2   World Health Organization.   (World Health Organization, Geneva, 2004).
3   Merritt, R. W. *et al.* Ecology and transmission of Buruli ulcer disease: a systematic review. *PLoS Negl Trop Dis* **4**, e911-e911, doi:10.1371/journal.pntd.0000911 (2010).
4   Organization, W. H. *Buruli ulcer (Mycobacterium ulcerans infection)*, <https://www.who.int/news-room/fact-sheets/detail/buruli-ulcer-(mycobacterium-ulcerans-infection)> (2022).
5   Department of Health and Human Services (Victoria). *Interactive infectious disease surveillance reports [internet]*, <https://www2.health.vic.gov.au/public-health/infectious-diseases/infectious-diseases-surveillance/interactive-infectious-disease-reports> (2021).
6   Janssens, P. G., Quertinmont, M. J., Sieniawski, J. & Gatti, F. Necrotic tropical ulcers and mycobacterial causative agents. *Trop Geogr Med* **11**, 293-312 (1959).
7   Clancey, J. K., Dodge, O. G., Lunn, H. F. & Oduori, M. L. Mycobacterial skin ulcers in Uganda. *Lancet (London, England)* **2**, 951-954, doi:10.1016/s0140-6736(61)90793-0 (1961).
8   Muleta, A. J., Lappan, R., Stinear, T. P. & Greening, C. Understanding the transmission of Mycobacterium ulcerans: A step towards controlling Buruli ulcer. *PLoS Negl Trop Dis* **15**, e0009678, doi:10.1371/journal.pntd.0009678 (2021).
9   Blasdell, K. R. *et al.* Environmental risk factors associated with the presence of Mycobacterium ulcerans in Victoria, Australia. *PLoS One* **17**, e0274627, doi:10.1371/journal.pone.0274627 (2022).
10 Carson, C. *et al.* Potential wildlife sentinels for monitoring the endemic spread of human buruli ulcer in South-East australia. *PLoS Negl Trop Dis* **8**, e2668, doi:10.1371/journal.pntd.0002668 (2014).
11 Fyfe, J. A. M. *et al.* A major Role for mammals in the ecology of *Mycobacterium ulcerans*. *PLoS Negl Trop Dis* **4**, e791, doi:10.1371/journal.pntd.0000791 (2010).
12 Johnson, P. D. R. in *Buruli Ulcer: Mycobacterium Ulcerans Disease*   (eds G. Pluschke & K. Roltgen) 61-76 (2019).
13 O'Brien, C. R. *et al.* Clinical, microbiological and pathological findings of Mycobacterium ulcerans infection in three Australian Possum species. *PLoS Negl Trop Dis* **8**, e2666, doi:10.1371/journal.pntd.0002666 (2014).
14 Xu, R. W., Stinear, T. P., Johnson, P. D. & O'Brien, D. P. Possum bites man: case of Buruli ulcer following possum bite. *Med J Aust* **216**, 452-453, doi:10.5694/mja2.51505 (2022).
15 Johnson, P. D. R. *et al. Mycobacterium ulcerans* in mosquitoes captured during outbreak of Buruli ulcer, southeastern Australia. *Emerging Infectious Diseases* **13**, 1653-1660, doi:10.3201/eid1311.061369 (2007).
16 Fyfe, J. A. *et al.* Development and application of two multiplex real-time PCR assays for the detection of *Mycobacterium ulcerans* in clinical and environmental samples. *Applied and environmental microbiology* **73**, 4733-4740 (2007).
17 Quek, T. Y. *et al.* Risk factors for Mycobacterium ulcerans infection, southeastern Australia. *Emerg Infect Dis* **13**, 1661-1666, doi:10.3201/eid1311.061206 (2007).
18 Landier, J. *et al.* Adequate wound care and use of bed nets as protective factors against Buruli Ulcer: results from a case control study in Cameroon. *PLoS Negl Trop Dis* **5**, e1392, doi:10.1371/journal.pntd.0001392 (2011).
19 Pouillot, R. *et al.* Risk factors for buruli ulcer: a case control study in Cameroon. *PLoS Negl Trop Dis* **1**, e101, doi:10.1371/journal.pntd.0000101 (2007).
20 Johnson, P. D. *et al.* Consensus recommendations for the diagnosis, treatment and control of Mycobacterium ulcerans infection (Bairnsdale or Buruli ulcer) in Victoria, Australia. *Med J Aust* **186**, 64-68, doi:10.5694/j.1326-5377.2007.tb00802.x (2007).
21 Lavender, C. J. *et al.* Risk of Buruli ulcer and detection of Mycobacterium ulcerans in mosquitoes in southeastern Australia. *PLoS Negl Trop Dis* **5**, e1305, doi:10.1371/journal.pntd.0001305 (2011).

OFFICIAL

22 Wallace, J. R. *et al.* Mycobacterium ulcerans low infectious dose and mechanical transmission
support insect bites and puncturing injuries in the spread of Buruli ulcer. *PLoS Negl Trop Dis* **11**,
e0005553, doi:10.1371/journal.pntd.0005553 (2017).
23 Australian Bureau of Statistics. *2016 Census QuickStats*,
<https://quickstats.censusdata.abs.gov.au/census_services/getproduct/census/2016/quickstat/SSC
22187?opendocument> (2021).
24 Australian Bureau of Meteorology. *Summary statisitcs Mornington*,
<http://www.bom.gov.au/climate/averages/tables/cw_086079.shtml> (2022).
25 World Health Organization. *Number of new reported cases of Buruli ulcer*,
<https://apps.who.int/gho/data/node.main.A1631> (2023).
26 Mornington Peninsula Shire. *Mornington Peninsula Shire population forecasts*,
<https://forecast.id.com.au/mornington-peninsula> (2023).
27 Russell, R. C. *A colour photo atlas of mosquitoes of Southeastern Australia.* (University of Sydney
Printing Service, NSW, 1996).
28 Dobrotworsky, N. V. *The mosquitoes of Victoria (Diptera, Culicidae).* (Melbourne University Press,
1965).
29 Liehne, P. F. S. *An atlas of the mosquitoes of Western Australia.* (Health Department of Western
Australia, 1991).
30 Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. DNA primers for amplification of
mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol
Biotechnol* **3**, 294-299 (1994).
31 PooledInfRate, Version 4.0: a Microsoft[(R)] Office Excel[(C)] Add-In to compute prevalence estimates
from pooled samples. (Fort Collins, CO, USA, 2009).
32 R: A language and environment for statistical computing. R Foundation for Statistical Computing,
Vienna, Austria (2020).
33 Illumina. *Nextera Flex for Enrichment with RNA Probes*,
<https://support.illumina.com/content/dam/illumina-
support/documents/documentation/chemistry_documentation/samplepreps_nextera/nextera-
flex-enrichment/nextera-flex-for-enrichment-rna-probes-demonstrated-protocol-1000000070581-
01.pdf > (2023).
34 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: computing large minimum evolution trees with
profiles instead of a distance matrix. *Mol Biol Evol* **26**, 1641-1650, doi:10.1093/molbev/msp077
(2009).
35 Endersby, N. M. *et al.* Evidence of cryptic genetic lineages within *Aedes notoscriptus* (Skuse).
*Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in
infectious diseases* **18**, 191-201, doi:10.1016/j.meegid.2013.04.035 (2013).
36 Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new heuristics and
parallel computing. *Nature Methods* **9**, 772, doi:10.1038/nmeth.2109 (2012).
37 Guindon, S. & Gascuel, O. A simple, fast, and accurate algorithm to estimate large phylogenies by
maximum likelihood. *Systematic biology* **52**, 696-704, doi:10.1080/10635150390235520 (2003).
38 QGIS Geographic Information System. Open Source Geospatial Foundation Project. (2022).
39 Statistics, A. B. o. *1270.0.55.007 - Australian Population Grid, 2011*,
<https://www.abs.gov.au/ausstats/abs@.nsf/lookup/1270.0.55.007main+features12011> (2011).
40 Australian Bureau of Statistics. <https://www.abs.gov.au> (2023).
41 QGIS Geographic Information System (2021).
42 Kulldorff, M. A spatial scan statistic. *Commun Stat-Theor M* **26**, 1481-1496, doi:Doi
10.1080/03610929708831995 (1997).
43 Vandelannoote, K. *et al.* Structured surveys of Australian native possum excreta predict Buruli ulcer
occurrence in humans. *Biorxiv*, doi:https://doi.org/10.1101/2022.11.16.516821 (2022).
44 Loftus, M. J. *et al.* The incubation period of Buruli ulcer (Mycobacterium ulcerans infection) in
Victoria, Australia - Remains similar despite changing geographic distribution of disease. *PLoS Negl
Trop Dis* **12**, e0006323, doi:10.1371/journal.pntd.0006323 (2018).

45   Trubiano, J. A., Lavender, C. J., Fyfe, J. A., Bittmann, S. & Johnson, P. D. The incubation period of
810   Buruli ulcer (Mycobacterium ulcerans infection). *PLoS Negl Trop Dis* **7**, e2463,
811   doi:10.1371/journal.pntd.0002463 (2013).
812   46   Pebesma, E. Simple Features for R: Standardized Support for Spatial Vector Data. *R J* **10**, 439-446
813   (2018).
814   47   R: A language and environment for statistical computing (R Foundation for Statistical Computing,
815   Vienna, Austria., 2021).
816   48   Buultjens, A. H. *et al.* Comparative Genomics Shows That Mycobacterium ulcerans Migration and
817   Expansion Preceded the Rise of Buruli Ulcer in Southeastern Australia. *Appl Environ Microbiol* **84**,
818   doi:10.1128/AEM.02612-17 (2018).
819   49   Taouk, M. L. *et al.* Characterisation of Treponema pallidum lineages within the contemporary
820   syphilis outbreak in Australia: a genomic epidemiological analysis. *Lancet Microbe* **3**, e417-e426,
821   doi:10.1016/S2666-5247(22)00035-0 (2022).
822   50   Vandelannoote, K. *et al.* Statistical modelling based on structured surveys of Australian native
823   possum excreta harbouring Mycobacterium ulcerans predicts Buruli ulcer occurrence in humans.
824   *Elife* **12**, doi:10.7554/eLife.84983 (2023).
825   51   Mangas, K. M. *et al.* Vaccine-Specific Immune Responses against Mycobacterium ulcerans Infection
826   in a Low-Dose Murine Challenge Model. *Infect Immun* **88**, doi:10.1128/IAI.00753-19 (2020).
827   52   Portaels, F., Elsen, P., Guimaraes-Peres, A., Fonteyne, P. A. & Meyers, W. M. Insects in the
828   transmission of *Mycobacterium ulcerans* infection. *Lancet (London, England)* **353**, 986,
829   doi:10.1016/s0140-6736(98)05177-0 (1999).
830   53   Marsollier, L. *et al.* Aquatic plants stimulate the growth of and biofilm formation by Mycobacterium
831   ulcerans in axenic culture and harbor these bacteria in the environment. *Appl Environ Microbiol* **70**,
832   1097-1103, doi:10.1128/AEM.70.2.1097-1103.2004 (2004).
833   54   Portaels, F. *et al.* First cultivation and characterization of *Mycobacterium ulcerans* from the
834   environment. *PLoS Negl Trop Dis* **2**, e178-e178, doi:10.1371/journal.pntd.0000178 (2008).
835   55   Singh, A., McBride, W. J. H., Govan, B., Pearson, M. & Ritchie, S. A. A survey on *Mycobacterium*
836   *ulcerans* in mosquitoes and march flies captured from endemic areas of Northern Queensland,
837   Australia. *PLoS Negl Trop Dis* **13**, e0006745, doi:10.1371/journal.pntd.0006745 (2019).
838   56   Yerramilli, A. *et al.* The location of Australian Buruli ulcer lesions—Implications for unravelling
839   disease transmission. *PLoS Negl Trop Dis* **11**, e0005800, doi:10.1371/journal.pntd.0005800 (2017).
840   57   Gomes, B. *et al.* Feeding patterns of molestus and pipiens forms of Culex pipiens (Diptera:
841   Culicidae) in a region of high hybridization. *Parasit Vectors* **6**, 93, doi:10.1186/1756-3305-6-93
842   (2013).
843   58   Bustamante, D. M. & Lord, C. C. Sources of error in the estimation of mosquito infection rates used
844   to assess risk of arbovirus transmission. *The American journal of tropical medicine and hygiene* **82**,
845   1172-1184, doi:10.4269/ajtmh.2010.09-0323 (2010).
846   59   Kay, B. H., Boyd, A. M., Ryan, P. A. & Hall, R. A. Mosquito feeding patterns and natural infection of
847   vertebrates with Ross River and Barmah Forest viruses in Brisbane, Australia. *The American journal*
848   *of tropical medicine and hygiene* **76**, 417-423 (2007).
849   60   Jansen, C. C. *et al.* Blood sources of mosquitoes collected from urban and peri-urban environments
850   in eastern Australia with species-specific molecular analysis of avian blood meals. *The American*
851   *journal of tropical medicine and hygiene* **81**, 849-857, doi:10.4269/ajtmh.2009.09-0008 (2009).
852   61   Flies, E. J., Flies, A. S., Fricker, S. R., Weinstein, P. & Williams, C. R. Regional comparison of mosquito
853   bloodmeals in South Australia: implications for Ross River virus ecology. *Journal of Medical*
854   *Entomology* **53**, 902-910, 909 (2016).
855   62   Muller, M., Murray, M. & Edwards, J. Blood-sucking midges and mosquitoes feeding on mammals
856   at Beatrice Hill, N.T. *Australian Journal of Zoology* **29**, 573-588,
857   doi:https://doi.org/10.1071/ZO9810573 (1981).
858   63   Lee, D. J., Clinton, K. J. & O'Gower, A. K. The blood sources of some Australian mosquitoes.
859   *Australian journal of biological sciences* **7**, 282-301, doi:10.1071/bi9540282 (1954).
860   64   Johansen, C. A., Power, S. L. & Broom, A. K. Determination of mosquito (Diptera: Culicidae)
861   bloodmeal sources in western australia: implications for arbovirus transmission. *Journal of Medical*
862   *Entomology* **46**, 1167-1175, doi:10.1603/033.046.0527 (2009).

65 Oshaghi, M. A. *et al.* Effects of post-ingestion and physical conditions on PCR amplification of host
blood meal DNA in mosquitoes. *Exp Parasitol* **112**, 232-236, doi:10.1016/j.exppara.2005.11.008
(2006).
66 Wallace, J. R. *et al.* Interaction of Mycobacterium ulcerans with mosquito species: implications for
transmission and trophic relationships. *Appl Environ Microbiol* **76**, 6215-6222,
doi:10.1128/AEM.00340-10 (2010).
67 Doig, K. D. *et al.* On the origin of Mycobacterium ulcerans, the causative agent of Buruli ulcer. *BMC*
Genomics **13**, 258, doi:10.1186/1471-2164-13-258 (2012).
68 Stinear, T. P. *et al.* Reductive evolution and niche adaptation inferred from the genome of
Mycobacterium ulcerans, the causative agent of Buruli ulcer. *Genome Res* **17**, 192-200,
doi:10.1101/gr.5942807 (2007).
69 Eisen, R. J. & Gage, K. L. Transmission of flea-borne zoonotic agents. *Annu Rev Entomol* **57**, 61-82,
doi:10.1146/annurev-ento-120710-100717 (2012).
70 Kernif, T., Leulmi, H., Raoult, D. & Parola, P. Emerging Tick-Borne Bacterial Pathogens. *Microbiol*
Spectr **4**, doi:10.1128/microbiolspec.EI10-0012-2016 (2016).
71 Ryden, P. *et al.* Outbreaks of tularemia in a boreal forest region depends on mosquito prevalence. *J*
Infect Dis **205**, 297-304, doi:10.1093/infdis/jir732 (2012).
72 Thelaus, J. *et al.* Francisella tularensis subspecies holarctica occurs in Swedish mosquitoes, persists
through the developmental stages of laboratory-infected mosquitoes and is transmissible during
blood feeding. *Microb Ecol* **67**, 96-107, doi:10.1007/s00248-013-0285-1 (2014).
73 Abdellahoum, Z., Maurin, M. & Bitam, I. Tularemia as a Mosquito-Borne Disease. *Microorganisms* **9**,
doi:10.3390/microorganisms9010026 (2020).
74 Hendry, T. A. *et al.* Ongoing Transposon-Mediated Genome Reduction in the Luminous Bacterial
Symbionts of Deep-Sea Ceratioid Anglerfishes. *mBio* **9**, doi:10.1128/mBio.01033-18 (2018).
75 Perreau, J. & Moran, N. A. Genetic innovations in animal-microbe symbioses. *Nat Rev Genet* **23**, 23-
39, doi:10.1038/s41576-021-00395-z (2022).
76 Drucker, M. & Then, C. Transmission activation in non-circulative virus transmission: a general
concept? *Current Opinion in Virology* **15**, 63-68, doi:https://doi.org/10.1016/j.coviro.2015.08.006
(2015).
77 Archer, M. S. *The ecology of invertebrate associations with vertebrate carrion in Victoria, with*
reference to forensic entomology Doctor of Philosophy thesis, The University of Melbourne, (2002).
78 Lang, M. D., Allen, G. R. & Horton, B. J. Blowfly succession from possum (*Trichosurus vulpecula*)
carrion in a sheep-farming zone. *Medical and Veterinary Entomology* **20**, 445-452,
doi:https://doi.org/10.1111/j.1365-2915.2006.00654.x (2006).
79 Skopyk, A. D., Forbes, S. L. & LeBlanc, H. N. Recognizing the inherent variability in Dipteran
colonization and decomposition rates of human donors in Sydney, Australia. *Journal of Clinical and*
Health Sciences **6**, 102-119 (2021).
80 C A. Van Schie, Spafford, H., Carver, S. S. & Weinstein, P. Salinity tolerance of Aedes
camptorhynchus (Diptera: Culicidae) from two regions in southwestern Australia. *Austral*
Entomology **48**, 293-299 (2009).
81 Jeffery, J. A. L., Kay, B. H. & Ryan, P. A. Development time and survival of *Verrallina funerea*
(Theobald) (Diptera: Culicidae) immatures and other brackish water mosquito species in southeast
Queensland, Australia. *Australian Journal of Entomology* **44**, 226-232,
doi:https://doi.org/10.1111/j.1440-6055.2005.00479.x (2005).
82 Hamlyn-Harris, R. Notes on the breeding-places of *Aedes* (Finlaya) *notoscriptus*, Skuse, in
Queensland. *Bulletin of Entomological Research* **19**, 405-409, doi:10.1017/S0007485300020782
(1928).
83 Russell, R. C. *Mosquitoes and mosquito-borne disease in southeast Australia.* (Department of
Medical Entomology, Westmead Hospital and Department of Medicine, University of Sydney, NSW,
1993).
84 Russell, R. C. A review of the status and significance of the species within the *Culex pipiens* group in
Australia. *Journal of the American Mosquito Control Association* **28**, 24-27 (2012).