DNA repair and anti-cancer mechanisms in the longest-living mammal: the

bowhead whale

Denis Firsanov[1]*, Max Zacher[1]*, Xiao Tian[1], Yang Zhao[1], John C. George[2], Todd L. Sformo[2], Greg

Tombline[1], Seyed Ali Biashad[1], Abbey Gilman[1], Nicholas Hamilton[1], Avnee Patel[1], Maggie

Straight[1], Minseon Lee[1], J. Yuyang Lu[1], Ena Haseljic[1], Alyssa Williams[1], Nalani Miller[1], Vadim N.

Gladyshev[3], Zhengdong Zhang[4], Jan Vijg[4, #], Andrei Seluanov[1,5, #], Vera Gorbunova[1,5, #]

[1]Department of Biology, University of Rochester, Rochester, NY, USA.

[2]Department of Wildlife Management, North Slope Borough, Utqiaġvik (Barrow), AK 99723, USA

[3]Division of Genetics, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School,

Boston, MA 02115, USA

[4]Department of Genetics, Albert Einstein College of Medicine, Bronx, NY 10461, USA

[5]Department of Medicine, University of Rochester Medical Center, Rochester, NY, USA

*These authors contributed equally to this work.

[#]Corresponding authors:

Vera Gorbunova

University of Rochester

Email: vera.gorbunova@rochester.edu

Andrei Seluanov

University of Rochester

Email: andrei.seluanov@rochester.edu

Jan Vijg

Albert Einstein College of Medicine

Email: jan.vijg@einsteinmed.edu

Abstract

At over 200 years, the maximum lifespan of the bowhead whale exceeds that of all other mammals. The bowhead is also the second-largest animal on Earth, reaching over 80,000 kg[1]. In spite of its very large number of cells, the bowhead is not highly cancer-prone, an incongruity termed Peto's Paradox[2]. This has been explained by the evolution of additional tumor suppressor genes in larger animals, which is supported by research on elephants demonstrating expansion of the p53 gene[3–5]. However, we show here that bowhead whale fibroblasts undergo oncogenic transformation after disruption of fewer tumor suppressors than required for human fibroblasts. Instead, analysis of DNA repair revealed that bowhead cells repair double-strand breaks with uniquely high efficiency and accuracy compared to other mammals. Further, we identified two proteins, CIRBP and RPA2, that are present at high levels in bowhead fibroblasts and increase the efficiency and fidelity of DNA repair in human cells. These results suggest that rather than possessing additional tumor suppressor genes as barriers to oncogenesis, the bowhead whale relies on more accurate and efficient DNA repair to preserve genome integrity. This strategy that does not eliminate cells but repairs them may be critical for the long and cancer-free lifespan of the bowhead whale. Our work demonstrates the value of studying long-lived organisms in identifying novel longevity mechanisms and their potential for translation to humans.

Introduction

The Alaskan Iñupiat Inuit, who carry on a long tradition of subsistence hunting of the bowhead whale (*Balaena mysticetus*), maintain that these animals "live two human lifetimes"[6]. Remarkably, a series of bowhead whales captured in the late-twentieth and early-twenty-first centuries lent new credence to these claims, as embedded in their bodies were traditional stone harpoon points and bomb lance fragments dating to the Victorian era[7]. Subsequent scientific study and age estimation through quantification of

ovarian corpora, baleen dating, and eye lens aspartic acid racemization analysis

supported a maximum lifespan exceeding 200 years in the bowhead whale[7–12]. Thus,

the range of mammalian lifespans covers roughly 2 orders of magnitude, with the model

organism *Mus musculus* living for 2-3 years while the bowhead whale lives 100 times as

long.

An increased number of cells and cell divisions in larger organisms might be expected

to contribute to increased cancer incidence and shorter lifespans. The apparent

contradiction between expected and observed cancer rates in relation to species body

mass has been noted for decades and is known as Peto's Paradox.[2,13–15] It is theorized

to result from compensatory evolutionary adaptations driven by reduced extrinsic

mortality in larger species[2]. The bowhead whale exceeds 80,000 kg in mass and 200

years in lifespan. Both of these factors predispose it to accumulating large numbers of

DNA mutations throughout life. To remain alive for so long it must possess uniquely

potent genetic mechanisms to prevent cancer and other age-related diseases.

However, primary research publications on genetic and molecular mechanisms of aging

in the bowhead whale are scarce, consisting primarily of genome and transcriptome

analysis[16–18].

The multi-stage model of carcinogenesis posits that the transition from a normal cell to a

cancer cell involves multiple distinct genetic "hits," or mutations[19]. Larger and longer-

living species might require greater numbers of "hits" for oncogenic transformation,

given their greater cell number and increased lifespan. Indeed, there is experimental

evidence to support this hypothesis. Rangarajan et al. found that while mouse

fibroblasts require perturbation of 2 pathways for tumorigenic transformation (p53 and

Ras), human fibroblasts require 5 hits (p53, pRb, PP2A, telomerase and Ras)[20]. A

human should thus have a dramatically lower per-cell incidence of malignant

transformation than a mouse, and can maintain a larger number of cells for a longer

period of time.

Species that are larger and longer-lived than humans may be expected to have even

more layers of protection against oncogenic transformation than humans and common

model species. In support of this hypothesis, recent studies have identified copy number

expansion and functional diversification of multiple tumor suppressor genes, such as

TP53 and *LIF*, in elephants and other taxa[3,5,21–23]. These studies identified multiple

copies of *TP53* in the elephant genome, several of which were confirmed to be

transcribed and translated in elephant fibroblasts and contributed to an enhanced

apoptotic response to genotoxic stress.

However, additional copies of p53 genes are unlikely to slow down aging. One

promising mechanism that could help explain both cancer resistance and slower aging

in long-lived mammals is more accurate or efficient DNA repair. Genetic mutations have

been identified as causal factors in carcinogenesis for over a century[24]. Perhaps one of

the most compelling lines of evidence supporting the role of DNA repair in the

pathogenesis of aging and cancer comes from studies of mutants with accelerated

aging phenotypes. Remarkably, most such mutants have defects in DNA repair

enzymes[25–29]. Across species, several studies have also pointed toward improved DNA

repair capacity and reduced mutation accumulation as characteristics associated with

species longevity[30–34]. Here, we identify specific cellular and molecular traits

characterizing bowhead cancer resistance and longevity and distinguishing the

bowhead from shorter-lived mammals including humans. We show that bowhead whale

cells are not more prone to apoptosis, and require fewer genetic hits for malignant

transformation than human cells. Instead, the bowhead whale relies on more accurate

and efficient DNA double strand break (DSB) repair mediated by CIRBP and RPA2.

This more "conservative" strategy that does not needlessly eliminate cells but repairs

them may be critical for the long and cancer-free lifespan of the bowhead whale.

Results

Growth characteristics, cellular senescence, and cell death in bowhead whale

Most human somatic cells lack telomerase activity and as a result undergo replicative

senescence with serial passaging in culture[35]. Replicative and stress-induced

senescence are important mechanisms for preventing cancer. Using TRF and TRAP

assays to measure telomere length and telomerase activity, we found that bowhead

whale skin fibroblasts, like human fibroblasts, lack telomerase activity and experience

telomere shortening followed by replicative senescence with passaging in culture

(Figure 1a, b). In both species, nearly all cells stained positive for senescence-

associated β-galactosidase upon terminal growth arrest (Figure 1c, d). As in human

fibroblasts, stable overexpression of human telomerase reverse transcriptase (*hTERT*)

to maintain telomere length prevented replicative senescence in bowhead cells (Figure

1a). Senescence can also be induced by DNA damage. Like human cells, bowhead

whale skin fibroblasts efficiently entered senescence but did not significantly induce cell

death in response to 10 or 20 Gy of γ-irradiation (Figure 1c-e).

Interestingly, transcriptome analysis of human and bowhead whale senescent

fibroblasts showed reduced induction of senescence-associated secretory phenotype

(SASP) factors in bowhead whale (Figure 1f) relative to human. Paracrine effects of

SASP on surrounding cells are thought to contribute to age-related diseases and

carcinogenesis. These transcriptomic differences may indicate that senescence is able

to preserve its anti-cancer function in the bowhead with reduced harmful paracrine

signaling.

To test whether increased p53 activity could contribute to cancer resistance in the

bowhead whale, we transiently transfected primary mouse, cow, human and bowhead

whale skin fibroblasts with a luciferase reporter vector containing a p53-response

element. The bowhead whale cells had the lowest p53 activity of the species tested

(Figure 1g). Additionally, we did not observe any differences in the induction of

apoptosis in response to UVC between species (Figure 1h). Together, our results argue

against the idea that increased clearance of damaged cells through apoptosis

contributes to cancer resistance in the bowhead whale.

Requirements for oncogenic transformation of bowhead whale cells

We initially identified a minimal combination of oncogene and tumor suppressor hits

required for *in vitro* malignant transformation of bowhead whale skin fibroblasts using

the soft agar assay, which measures anchorage-independent growth, a hallmark of

cancer. While normal cells undergo growth arrest or programmed cell death (anoikis) in

soft agar, malignant cells continue to grow without substrate adhesion and form visible

colonies[36]. We introduced constructs targeting oncogene and tumor suppressor

pathways into primary skin fibroblasts with PiggyBac (PB) transposon vectors, which

integrate into the genome and drive stable expression. Since bowhead whale primary

fibroblasts, like human fibroblasts, exhibit progressive telomere shortening and lack

telomerase activity (Figure 1b), we used cell lines expressing (*hTERT*) to bypass

replicative senescence.

In agreement with published findings, transformation of human *hTERT+* fibroblasts

required combined expression of H-Ras^{G12V}, SV40 Large T (LT) antigen (which binds

and inactivates p53 and the Rb family of tumor suppressors), and SV40 Small T (ST)

antigen (which binds and inactivates PP2A) (Figure 2a)[20]. Rather than requiring hits to

additional pathways, however, bowhead *hTERT+* fibroblasts were transformed by H-

Ras^{G12V} and SV40 LT alone, suggesting that bowhead cells may require fewer genetic

mutations to become cancerous compared to human cells (Figure 2a). These findings

were supported by mouse xenograft assays, in which the number of hits needed for

tumor growth matched the findings from soft agar (Figure 2b).

We next sought to confirm these findings at the genetic level, through CRISPR editing

of individual tumor suppressor genes in bowhead fibroblasts. While the sequenced

bowhead genome has not revealed copy number expansion of canonical tumor

suppressor genes[16,17], CRISPR knockout allows for more precise quantification of the

number of genetic mutations required for oncogenesis. While the most important target

of SV40 LT is thought to be Rb (*RB1* gene), it is also known to inactivate p130 and

p107, two additional members of the Rb-family, providing some level of functional

redundancy. Using CRISPR, we were able to introduce targeted mutations into the

bowhead *RB1* gene specifically, along with *TP53* (the other target of LT), and *PTEN* (an

upstream inhibitor of Akt signaling commonly mutated in human cancers, and operating

in the same pathway as PP2A). Following transfection of *hTERT+* bowhead fibroblasts

with Cas9-guide RNA ribonucleoprotein complexes targeting each of the

aforementioned genes, we screened clonally isolated colonies for loss of the targeted

protein by Western blot (Figure 2c, d, Extended Data Figure 1a, b). We additionally

screened the colonies with luciferase reporter assays to confirm loss of protein function

and activity (Extended Data Figure 1c, d). For each selected clone, we sequenced the

CRISPR-targeted genes to confirm homozygous knockout at the genetic level and

determine the causal mutations (Supplementary Figure 1, 2). Through this strategy, we

generated single and compound homozygous knockout bowhead whale fibroblasts for

TP53, *RB1*, and *PTEN*. In agreement with our initial findings, genetic inactivation of

TP53 and *RB1* in bowhead whale fibroblasts expressing *hTERT* and H-Ras^{G12V} was

sufficient for malignant transformation in both soft agar and mouse xenograft assays

(Figure 2a, b). These findings suggest that despite its large size and long lifespan, the

cells of the bowhead whale unexpectedly require fewer mutational hits for malignant

transformation than human cells.

Excision repair and mutagenesis in the bowhead whale

To understand if improved DNA repair might reduce cancer risk in the bowhead whale,

we first assessed the efficiency of nucleotide excision repair (NER) and base excision

repair (BER) in bowhead whale cells. NER is primarily responsible for removing helix-

distorting DNA lesions. To quantify NER activity, we utilized a host cell plasmid

reactivation assay and quantitation of cyclobutane pyrimidine dimers by ELISA to

measure repair of UVC-induced DNA damage. NER efficiency was similar between

bowhead and human (Extended Data Figure 2a, b). BER is responsible for ameliorating

many types of spontaneous DNA base damage such as oxidation and deamination. The

efficiency of BER, as measured by plasmid reactivation assay, trended toward higher

BER activity in bowhead whale compared to human cells, but this difference was not

statistically significant (Extended Data Figure 2c). Additionally, bowhead fibroblasts

displayed dramatically higher PARP activity in response to H_2O_2 treatment (Extended

Data Figure 2d, e). PARP proteins play important roles in the DNA damage response

and are recruited to sites of DNA damage where they participate in the DNA damage

response and in repair.

We next compared the point mutation frequency in the bowhead whale and human

using the HPRT mutagenesis assay, which relies on loss of HPRT activity after

mutagen treatment[37]. The *HPRT* gene exists as a single copy on the X chromosome in

male mammalian cells, a feature we found to be true for the bowhead as well (see

Methods). We treated primary fibroblast lines from male bowhead whale and human

with the potent mutagen and alkylating agent N-ethyl-N-nitrosourea (ENU), and then

plated cells in selective media containing 6-thioguanine, which kills cells with functional

HPRT. Despite a slightly higher sensitivity to ENU in bowhead whale as indicated by

colony-forming efficiency in non-selective media, the rate of HPRT mutant colony

formation was markedly lower in bowhead whale than human fibroblasts, an effect

which remained significant after adjusting for plating efficiency (Extended Data Figure

2f, g). This result suggests that bowhead whale has more accurate DNA repair.

High efficiency and fidelity of bowhead whale double-strand break repair

DNA DSBs are toxic if not repaired and may lead to mutations through inaccurate

repair. DSBs are repaired by two major pathways: non-homologous end joining (NHEJ)

and homologous recombination (HR). To assess relative NHEJ efficiency, we integrated

fluorescent GFP-based reporter cassettes[38] into fibroblasts from mouse, cow, human

and bowhead whale. Following DSB induction with I-SceI, we observed markedly

elevated NHEJ efficiency in bowhead whale relative to other species (Figure 3a,

Extended Data Figure 3a-c). Additionally, bowhead whale fibroblasts accumulated fewer

micronuclei, a marker of chromosomal instability, after 2Gy γ-irradiation (Figure 3b,

Extended Data Figure 3d). The kinetics of radiation-induced DSB rejoining as measured

by pulsed-field gel electrophoresis were similar between bowhead and human cells,

with almost all breaks repaired within 48h in both species (Extended Data Figure 3e, f).

As mutations resulting from inaccurate DSB repair promote cancer development, we

next sought to assess the fidelity of DSB repair in the bowhead whale. Sequencing and

analysis of repair junctions from integrated NHEJ reporter (Figure 3) and extra-

chromosomal NHEJ reporter (Extended Data Figure 3a, c) assays showed higher

fidelity of NHEJ in bowhead whale cells: compared to human, the bowhead whale is

less prone to producing deletions during the repair of incompatible DNA termini and far

more frequently joins ends without deleting any bases beyond the small overhang

region.

We also measured the fidelity of repair at an endogenous genomic locus. Initial

sequencing results from CRISPR tumor suppressor knockout experiments suggested a

strong bias in bowhead whale cells toward DSB repair products with single-base

insertion mutations, in contrast to the greater diversity and size of repair outcomes

commonly observed in human and mouse cells. To systematically compare mutational

outcomes of CRISPR break repair in the bowhead whale to those of humans and

shorter-living mammals, we performed CRISPR transfections in multiple primary

fibroblast lines from bowhead whale, human, cow, and mouse, and used deep amplicon

sequencing of the targeted locus to generate detailed profiles of repair outcomes. We

took advantage of the fact that exon 1 of the *PTEN* tumor suppressor gene is highly

conserved across mammals, with 100% sequence identity across included species

(Extended Data Figure 4a). We were therefore able to examine species-specific DSB

repair outcomes at an endogenous genomic locus while minimizing intra-species

variation in the break-proximal sequence context.

Analysis of sequencing data revealed species-specific repair outcomes, which were

consistent across cell lines derived from multiple animals of each species (Figure 3d-f).

In human, cow, and mouse, the most common mutational outcomes were deletions. In

contrast, the bowhead was the only species that preferentially repaired breaks with a

single-base insertion. The frequency of unmodified alleles, which are known to occur

after error-free repair of CRISPR DSBs[39,40], was the highest in bowhead whale (Figure

3e). No insertions or deletions were detected in untreated control samples (Extended

Data Figure 4b). As analysis of CRISPR RNP transfection efficiency by flow cytometry

and cleavage efficiency by digital droplet PCR showed similar CRISPR efficiencies

across species (Extended Data Figure 5a, Supplementary Figure 3a, b), differences in

the unmodified allele fraction most likely result from differences in repair fidelity. While

small indels predominated in all species, we observed a marked inverse correlation

between the frequency of large deletions and species lifespan, with the bowhead

producing fewer large deletions than human, cow, and mouse (Figure 3d-f). When we

assigned frequency-based percentile ranks from most negative to most positive indel

size (largest deletions to largest insertions), we observed a strong correlation between

species lifespan and 5th percentile indel size, corresponding to large deletions

(Pearson's r=0.85, p=0.0009) (Extended Data Figure 6). The results from these

experiments suggest a greater efficiency and accuracy of DSB repair in the bowhead

whale relative to humans and other mammals.

CIRBP contributes to high DSB repair efficiency in the bowhead whale

To identify mechanisms contributing to the efficiency and accuracy of DSB repair in the

bowhead whale, we measured and compared expression of DNA repair proteins in the

bowhead to other mammalian species by Western blot, quantitative MS proteomics, and

RNA seq (Figure 4a, Extended Data Figure 7a-d). Bowhead cells showed strikingly

higher abundance of cold-inducible RNA-binding protein (CIRBP) than other

mammalian species. (Figure 4a, Extended Data Figure 7a, d). Levels of PARP1, a

functional partner of CIRBP during DNA repair[41], were also increased relative to human.

Unexpectedly, we found that levels of several important canonical NHEJ proteins (e.g.

Ku70, Ku80, DNA-PKcs) are substantially higher in human cells than any other species

tested (Figure 4a, Extended Data Figure 7a).

CIRBP is an RNA- and PAR-binding protein whose expression is induced by a variety of

cellular stressors including cold shock, hypoxia, and UV irradiation[41–44]. It binds the 3'

UTR of mRNAs that encode proteins involved in cellular stress and DNA damage

responses and promotes their stability and translation: known targets include ATR,

RPA2, thioredoxin, and p27[45–48]. There is also evidence for a more direct role of CIRBP

in DNA repair: PARP-1-dependent localization of CIRBP to sites of DNA damage

promotes DSB repair and antagonizes micronucleus formation[41].

To test whether CIRBP contributes to efficient NHEJ in bowhead whale cells, we

overexpressed human (hCIRBP) and bowhead whale (bwCIRBP) in human NHEJ

reporter cells. Overexpression of bwCIRBP, but not hCIRBP, enhanced NHEJ efficiency

in human cells ~1.6-fold (p=0.002) and in mouse embryonic fibroblasts ~3-fold (Figure

4b, Extended Data Figure 8a). Overexpression of bwCIRBP, but not hCIRBP, also

increased HR efficiency in human cells ~2-fold (Figure 4c, e). Conversely, CIRBP

depletion in bowhead whale cells by siRNA significantly reduced NHEJ efficiency

(p=0.032) (Figure 4d, e). Consistent with published observations, overexpression of

bwCIRBP with nine arginines in the repeated RGG motif mutated to alanine (9R/A),

which impairs CIRBP's ability to bind to PAR-polymers[41], failed to stimulate HR and

reduced stimulation of NHEJ (Figure 4b, c, e). To test the effects of bwCIRBP

overexpression on chromosomal stability, we quantified formation of micronuclei in

human cells after γ-irradiation. We found that overexpression of bwCIRBP significantly

decreased the percentage of cells with micronuclei, reflecting an improvement in

genomic stability (Figure 4f).

Surprisingly, we observed during our experiments that CIRBP protein abundance in the

bowhead remained high regardless of whether cells were cultured at 37°C, 33°C, or

lower temperatures (the core body temperature of the bowhead has been measured at

33.8°C)[49,50], suggesting constitutive expression in contrast to its strong temperature

dependence in other mammals (Extended Data Figure 8b)[42]. At no temperature did

human cells produce similarly high CIRBP levels. In contrast, RNA-seq analysis

(verified by qPCR) showed nearly identical CIRBP transcript levels between human and

bowhead whale fibroblasts (Extended Data Figure 7b, 8c). The discrepancy between

mRNA and protein levels in tested species agrees with previous findings of CIRBP

regulation at the post-transcriptional level[51,52] .

The human and bowhead CIRBP proteins differ by only 5 C-terminal amino acids

(Extended Data Figure 9a, b). Substitution of these 5 codons in hCIRBP with bowhead

codons increased protein expression, while mutation of bwCIRBP with the 5 hCIRBP

codons decreased it (Extended Data Figure 9c). This was not accompanied by changes

in mRNA transcript levels, consistent with post-transcriptional regulation (Extended Data

Figure 9d). Interestingly, although CIRBP abundance increased following introduction of

the 5 bowhead substitutions to hCIRBP, it did not achieve the expression of bowhead

CIRBP, suggesting that synonymous changes to the mRNA coding sequence contribute

to higher translation efficiency of bowhead whale CIRBP. Consistent with this notion,

bwCIRBP has a higher codon adaptation index (CAI)[53] than hCIRBP; this difference

remains after introduction of the five reciprocal mutations (Extended Data Figure 9e).

To test whether a hypothermia-mediated increase in human CIRBP can also affect DSB

repair efficiency, we kept human cells for two days at 33°C before transfection with

NHEJ reporter plasmid pre-digested with I-Sce-1. Indeed, cells exposed to hypothermia

showed a ~2-fold (p=0.0017) increase in NHEJ efficiency (Figure 4g).

Mechanisms of bowhead whale DNA repair fidelity

LC-MS proteomics data and Western blots with multiple antibodies suggested increased

abundance of the single-stranded DNA binding protein RPA2 in the bowhead whale

(Figure 4a, Extended Data Figure 7a, c). This was paralleled by elevated RPA2

transcript abundance (Extended Data Figure 7b). RPA is a conserved heterotrimeric

ssDNA-binding protein complex required for eukaryotic DNA replication, which plays a

critical role in DNA repair and DNA damage signaling[54,55]. RPA deficiency increases the

frequency of DSBs under both basal[56] and stressed[57] conditions. Conversely, RPA

overexpression increases resistance to genotoxic insults[57–60]. RPA promotes NHEJ *in*

vitro[61], can protect ssDNA overhangs at Ku-bound DSBs[62,63], and antagonizes

microhomology-mediated end joining[6448,49].

It has previously been shown that RPA2 is an unstable protein highly prone to oxidative

damage, and that the abundance of this subunit affects the abundance of RPA1 and

RPA3[65,66]. We therefore sought to determine whether the high abundance of RPA2 in

bowhead whale cells may contribute to DNA repair fidelity. Because RPA is required for

cell viability, and can cause toxicity when not expressed with appropriate subunit

stoichiometry[67], we selected methods aimed at transiently increasing or decreasing

RPA activity in the period surrounding break induction to specifically isolate effects on

DSB repair fidelity. Treatment of cells transfected with CRISPR to induce DSBs with the

small-molecule RPA DNA-binding inhibitor TDRL-505[68] significantly increased indel

rates in both bowhead and human fibroblasts (Figure 5a, b). In bowhead, RPA inhibition

also increased the frequency of larger deletions, although this difference did not reach

significance (Figure 5b). Conversely, co-transfection of trimeric recombinant human

RPA protein during CRISPR significantly decreased the frequency of mutated alleles in

human cells without affecting CRISPR RNP transfection efficiency (Figure 5a,

Supplementary Figure 3c). We additionally tested the effect of bwCIRBP

overexpression on repair fidelity in human cells and observed a modest but significant

reduction in indel rates (Figure 5c). These results support the notion that variations in

CIRBP and RPA2 abundance, affect the fidelity of DSB repair and promote genomic

stability in the bowhead whale.

Discussion

By studying a mammal capable of maintaining its health and avoiding death from cancer

for over 2 centuries, we are offered a unique glimpse behind the curtain of a global

evolutionary experiment that tested more mechanisms affecting cancer and aging than

humans could ever hope to approach. Through experiments using primary fibroblasts

from the bowhead whale, we experimentally determined genetic requirements for

oncogenic transformation in the world's longest living mammal, and provide evidence

that additional tumor suppressors are not the only solution to Peto's Paradox. Instead,

we find that the bowhead whale solution lies upstream of tumor suppressor loss, and is

defined by a capacity for highly accurate DNA DSB repair. We also present evidence

that two proteins, RPA2 and CIRBP, are highly expressed in the bowhead relative to

other mammals and contribute to more efficient and accurate DSB repair.

When cells experience DNA breaks, they are faced with a choice between several

different fates, which cooperate to balance short-term risks to organismal viability with

the long-term risks of genome instability. One option is resolution of the DNA damage

response and resumption of cell function. While an unrepaired DSB is sufficient to kill a

cell, an inaccurately repaired DSB can give rise to a cancer that kills the entire

organism. The magnitude of this threat is reflected in the diversity of checkpoint

mechanisms mammals have evolved to arrest or eliminate cells with DNA damage.

Thus, persistent or excessive damage signaling can trigger a second category of DNA

damage outcomes: apoptosis or senescence.

Prompt and efficient DSB repair promotes the survival and continued function of

somatic cells. Ideally, this results in faithful restoration of the original genomic

sequence. However, aberrant outcomes such as loss of chromosome fragments (e.g.

micronuclei), chromosomal rearrangements, or large deletions cause massive changes

to the original DNA sequence[69]. These forms of genome instability are hallmarks of

cancer and core drivers of malignant transformation. In quiescent cell populations, there

is increased potential for accumulation of persistent unrepaired DSBs. Efficient repair of

DSBs is particularly important for permanently post-mitotic cells like neurons and

cardiomyocytes, which often are not replaced after cell death. Ultimately, the value of

DSB repair to an organism's survival is a function of both accuracy and efficiency. Upon

excessive damage, it is often better for organismal survival and longevity to eliminate

cells at risk of genomic instability through apoptosis or senescence.

Senescence is an important mechanism to control cells with potential genome instability

through permanent growth arrest. Telomere shortening functions to induce senescence

by limiting the number of times a cell can replicate. This mechanism appears to be

functional in bowhead whale fibroblasts, in line with the theory that telomere shortening

is prevalent as a cancer-prevention mechanism in large-bodied mammals[70]. An

additional such mechanism was recently reported for the bowhead whale in the form of

a *CDKN2C* checkpoint gene duplication[71]. Senescence in general appears to represent

a tradeoff between cancer and other forms of age-related dysfunction. There is

substantial evidence for harmful paracrine effects of senescence through the pro-

inflammatory SASP[72]. Here we demonstrate that bowhead whale has reduced

inflammatory components of the SASP, which could reduce the negative impact of

senescence.

Like senescence, apoptosis is an efficient and critical mechanism to reduce the number

of cells with dangerous levels of genome instability. Its importance as a cancer

prevention mechanism is evident from the prevalence of p53 mutations in cancer[73] and

supported by findings of enhanced p53 function in the elephant[3–5]. However, high rates

of apoptosis are likely to accelerate age-related decline through permanent cell loss[74],

especially in post-mitotic and irreplaceable cell populations in tissues such as brain and

heart. Because excessive cell death or arrest is incompatible with organismal survival,

baseline levels of DNA damage must be repaired. Given that most cells in mammalian

bodies experience multiple endogenous DSBs per day[75], faithful repair of these breaks

is critical.

Why might the bowhead whale have evolved improved DNA repair, as opposed to the

increased tumor suppressor copy number and elevated apoptotic response found in the

elephant and often proposed as a solution to Peto's Paradox? One possible explanation

is that tumor suppressors, apoptosis, and senescence all appear to pose costs to the

organism and represent tradeoffs between cancer and stem cell depletion leading to

age-related degeneration. Simply shifting the balance from apoptosis/senescence to

survival and repair could be detrimental if not also coupled with increased fidelity, as

evidenced by the frequent upregulation of DNA repair pathways in cancer cells[76].

However, evolutionary improvements that couple high efficiency with high fidelity, as

found in the bowhead whale, would promote long-term tissue function and maintenance

at both the cellular and genomic levels. Efficient repair could reduce loss of valuable

post-mitotic cells with age and reduce mutations in dividing cells. Maintenance of

genome stability would reduce cancer risk and - as suggested by a rapidly growing body

of evidence implicating age-related somatic mosaicism as a ubiquitous feature and

functional driver of aging[77–81]- likely also protect against numerous other aspects of age-

related decline. Thus, the lower accuracy and efficiency of DSB repair observed in

mammals with shorter lifespans may simply reflect an absence of sufficient selective

pressure. One potential drawback of a very accurate DNA repair system is a reduction

in diversity and slower rate of evolution of new traits. However, species living in safe

and stable environments have no evolutionary pressure to rapidly evolve new

adaptations. Interestingly, genome analysis of long-lived rockfishes living in deep ocean

revealed positive selection in DNA repair pathways[82].

The evolution of constitutively high CIRBP expression in the bowhead whale is likely to

have been driven, at least in part, by the unique physiological stresses this Arctic

cetacean must endure. This naturally includes the extremely cold water in which it is

constantly immersed, but also the ischemia/reperfusion-like fluctuations in tissue

oxygen concentrations experienced by diving mammals[83].

Interestingly, beneficial effects of cold as a therapeutic agent have been known for a

long time with brief cold-water immersion believed to promote health and hardening in

Nordic cultures. Currently, whole body cryotherapy is widely used in sports medicine to

reduce inflammation and facilitate recovery after exercise or injury[84]. While molecular

mechanisms responsible for the beneficial effects of cryotherapy are largely unknown,

we speculate that increased CIRBP expression may contribute to health benefits by

facilitating DNA repair.

Cold showers provide a mild and transient reduction in body temperature, whereas

artificial reduction of body temperature, or therapeutic hypothermia, is used clinically

during surgery[85]. Interestingly, recovery from therapeutic hypothermia is compromised

in CIRBP-KO rats and improved in rats with transgenic CIRBP overexpression[86]. Similar

protective effects of CIRBP during hypothermia have been reported in brain[87], kidney[88]

intestine[89], and cold-preserved hearts[90]. Given the dramatically enhanced protein

production driven by bwCIRBP and hCIRBP mutants, local delivery of such CIRBP

variants (e. g. through mRNA/DNA vector injection) could hold promise as strategies to

improve tissue recovery from surgery or organ transplant.

There are currently no approved therapeutics which aim to bolster DNA repair for the

prevention of cancer or age-related decline, and it has been suggested that DNA repair

would be difficult or impossible to improve[91]. However, the bowhead whale provides

evidence that this notion is incorrect. Therapeutics based on increasing the activity or

abundance of proteins like CIRBP or RPA2 could one day enable the treatment of

genome instability as a modifiable disease risk factor.

Acknowledgements

We would like to thank the researchers at the North Slope Borough Department of

Wildlife Management, the Alaska Eskimo Whaling Commission, and the Iñupiaq

community of Barrow for generously sharing bowhead whale samples, time, resources,

skill, and knowledge, and without whom the above work would not have been possible.

This work was supported by grants from US National Institutes on Aging to VNG, ZZ,

JV, AS and VG, and by an award from the Milky Way Research Foundation to VG.

Author contributions

VG, AS, DF, MZ designed research; DF analyzed senescence, DNA repair, CIRBP with
help from SAB, AP, EH, AW, NM; MZ conducted tumor suppressor CRISPR
experiments, RPA experiments, HPRT assays, and assessed DNA repair fidelity using
CRISPR with help from SB, MS, NH; XT analyzed tumorigenicity; YZ and AG assisted
with mouse tumor studies; JCG, TLS, MZ, DF collected bowhead specimens; MZ, GT
performed proteomics; ML analyzed micronuclei; JYL analyzed SASP; VG, ZZ, JV
contributed data analysis and conceptualization; VG and AS obtained funding and
supervised the study; MZ, DF, AS and VG wrote the manuscript with input from all
authors.

Competing interests

487   The authors declare no competing interests.

Methods

Reagents

Detailed information on reagents, such as antibodies and sequences of primers, probes, CRISPR guides, and siRNAs, is provided in Supplementary Methods.

Animal experiments

All animal experiments were approved and performed under pre-approved protocols and in accordance with guidelines set by the University of Rochester Committee on Animal Resources (UCAR).

Whale sample collection

Bowhead whale tissues were obtained from adult bowhead whales (*Balaena mysticetus*) captured during 2014 and 2018 Iñupiaq subsistence harvests in Barrow (Utqiaġvik), AK, in collaboration with the North Slope Borough Department of Wildlife Management (NSB DWM) and Alaska Eskimo Whaling Commission after signing a Memorandum of Understanding (September 2014 and March 2021). Tissues were sampled immediately after bowhead whales were brought ashore, after permission to sample was given by the whaling captain, and explants kept in culture medium on ice or at 4°C through initial processing and shipping until arrival at the University of Rochester (UR) for primary fibroblast isolation from skin and lung. Transfer of bowhead whale samples from NSB DWM to UR was under NOAA/NMFS permit 21386.

Primary cell cultures

Primary skin fibroblasts were isolated from skin (dermal) tissues as previously

described.[92] Briefly, skin tissues were shaved and cleaned with 70% ethanol. Tissues

were minced with a scalpel and incubated in DMEM/F-12 medium (ThermoFisher) with

Liberase™ (Sigma) at 37ºC on a stirrer for 15-90 min. Tissues were then washed and

plated in DMEM/F-12 medium containing 12% fetal bovine serum (GIBCO) and

Antibiotic-Antimycotic (GIBCO). All subsequent maintenance culture for fibroblasts from

bowhead and other species was in EMEM (ATCC) supplemented with 12% fetal bovine

serum (GIBCO), 100 units/mL penicillin, and 100 mg/mL streptomycin (GIBCO). All

primary cells were cultured at 37ºC with 5% CO_2 and 3% O_2 except bowhead whale

cells, which were cultured at 33ºC with 5% CO_2 and 3% O_2 based on published field

measurements of bowhead body temperature, which measured a core temperature of

33.8 ºC and a range of lower temperatures in muscle and peripheral tissue.[49,50] Prior to

beginning experiments with bowhead whale fibroblasts, optimal growth and viability

conditions were empirically determined through testing of alternative temperatures,

serum concentrations, and cell culture additives, with optimal culture medium found to

be the same for bowhead and other species. Following isolation, low population-

doubling (PD) primary cultures were preserved in liquid nitrogen, and PD was

continually tracked and recorded during subsequent use for experiments.

Soft agar assay

Fibroblast culture medium as described above was prepared at 2X concentration using

2X EMEM (Lonza). To prepare the bottom layer of agar plates, 2X medium was mixed

with a sterile autoclaved solution of 1.2% Noble Agar (Difco) at a 1:1 volumetric ratio,

and 3 mL of 1X medium/0.6% agar was pipetted into each 6-cm cell culture dish and

allowed to solidify at room temperature in a tissue culture hood. To plate cells into the

upper layer of soft agar, cells were harvested and washed, and immediately prior to

plating were resuspended in 2X medium at 20,000 cells/1.5 mL and diluted twofold in

0.8% Noble Agar pre-equilibrated to 37°C. The cells in 0.4% agar/1X medium were

pipetted gently to ensure a homogeneous single cell suspension, and 3 mL (20,000

cells) per 6 cm dish were layered on top of the solidified lower layer. After solidifying in

tissue culture hoods for 20-30 min, additional medium was added to ensure the agar

layers were submerged, and dishes were moved into cell culture incubators. Fresh

medium was added onto the agar every 3 days. 4 weeks after plating, viable colonies

were stained overnight with nitro blue tetrazolium chloride (Thermo Fisher) as

previously described.[36] All cell lines were plated in triplicate.

Mouse xenograft assay

NIH-III nude mice (Crl:NIH-Lystbg-J Foxn1nuBtkxid) were purchased from Charles

River Laboratories Inc. (Wilmington, MA, USA). Seven-week-old female mice were used

to establish xenografts. For each injection, 2×10^6 cells were harvested and

resuspended in 100 µl of ice-cold 20% matrigel (BD Bioscience, Franklin Lakes, NJ) in

PBS (Gibco). Mice were anesthetized with isoflurane gas, and 100 µl solution per

injection was injected subcutaneously into the right and left flanks of each mouse with a

22 gauge needle. 3 mice were injected bilaterally, for a total of 6 injections, per cell line

tested. Tumor length and width were measured and recorded every 3-4 days. Mice

were euthanized after reaching a predetermined humane tumor burden endpoint of a

maximum tumor dimension of 20mm in diameter, determined by the longest dimension

of the mouse's largest tumor. For mice that did not reach tumor burden endpoints,

experiments were terminated and mice euthanized after a maximum of 60 days.

Euthanized mice were photographed and tumors were excised, photographed, and

weighed to determine the mass of each tumor. Sections of each tumor were frozen at -

80°C and preserved in formalin.

Telomere lengths

Telomere length was analyzed by Southern blot using the TRF method. Genomic DNA

was extracted from cultured fibroblasts at different population doublings, digested with a

mixture of AluI, HaeIII, RsaI, and HinfI restriction enzymes that do not cut within

telomeric repeat sequences, separated using pulsed-field gel electrophoresis, and

hybridized with a radiolabeled oligonucleotide containing telomeric sequence

(TTAGGG)$_4$. Pulsed field gels were run using a CHEF-DR II apparatus (Bio-Rad) for

22h at a constant 45 V, using ramped pulse times from 1 to 10 s.

Telomeric repeat amplification protocol

Telomeric repeat amplification protocol assay was performed using the TRAPeze kit

(Chemicon, Temecula, CA, USA) according to manufacturer instructions. Briefly, in the

first step of the TRAP assay, radiolabeled substrate oligonucleotide is added to 0.5 µg

of protein extract. If telomerase is present and active, telomeric repeats (GGTTAG) are

added to the 3′ end of the oligonucleotide. In the second step, extended products are

amplified by PCR. Telomerase extends the oligonucleotide by multiples of 6 bp,

generating a ladder of products of increasing length. A human cancer cell line

overexpressing telomerase as well as rodent cells were used as a positive control.

CRISPR ribonucleoprotein transfection

CRISPR RNP complexes were formed in vitro by incubating Alt-R™ S.p.Cas9 Nuclease

V3 (Integrated DNA technologies) with tracRNA annealed to target-specific crRNA

(Integrated DNA Technologies) according to manufacturer instructions. For generation

of tumor suppressor knockouts, 3 RNP complexes with crRNAs targeting different sites

in a single target gene were combined and Alt-R Cas9 Electroporation Enhancer

(Integrated DNA Technologies) was added to transfection mixes prior to electroporation.

For comparative analysis of repair fidelity, 3 µg of pmaxGFP plasmid (Lonza) was

added to transfection mixes to monitor transfection efficiency. Cells were trypsinized

and washed with PBS, and 1×10^6 cells were resuspended in 100 µL of NHDF

Nucleofector Solution (Lonza). The cell suspension was then combined with the

CRISPR transfection solution and gently mixed prior to electroporation on an Amaxa

Nucleofector 2b (Lonza) using program U-23. For RPA inhibition, human and bowhead

fibroblasts were treated with 50 uM TDRL-505 (Sigma) (diluted 1000x into culture

medium from 50mM stock solution in DMSO) for 3 hours prior to CRISPR transfection

and kept in medium with 50 uM TDRL-505 for 18 hours after transfection. For RPA co-

transfection, 1 µg of recombinant human RPA heterotrimer (Enzymax) was added to

CRISPR transfection solution immediately prior to electroporation.

Isolation of clonal cell colonies and screening for tumor suppressor knockout

Following CRISPR transfection, cells were plated at low density in 15-cm dishes to

allow for the formation of isolated colonies. Once clonal colonies of sufficient size had

formed, positions of well-isolated colonies were visually marked on the bottom of the

cell culture dish while under a microscope using a marker. Dishes were aspirated and

washed with PBS. Forceps were used to dip PYREX® 8x8 mm glass cloning cylinders

in adhesive Dow Corning® high-vacuum silicone grease (Millipore Sigma) and one glass cylinder was secured to the dish over each marked colony. 150 µL of trypsin was added to each cylinder and returned to the incubator. When cells had rounded up from the plate, the trypsin in each cylinder was pipetted to detach cells and each colony was added to a separate well in a 6-cm culture dish containing culture medium. After colonies were expanded and split into 2 wells per colony, one well was harvested for Western blot screening for absence of target proteins, while the remaining well was kept for further experiments.

Luciferase reporter assays for knockout verification

For p53 activity measurement, 1×10^6 cells of control (WT) and clonally isolated p53 KO cell lines were electroporated with 3 µg p53 firefly luciferase reporter plasmid pp53-TA-Luc (Clontech/Takara) and 0.3 ug renilla luciferase control plasmid pRL-CMV (Promega) on an Amaxa Nucleofector 2b (Lonza). 24h later, cells were treated with 200 uM etoposide (Sigma) to induce p53 activity. 24h following etoposide treatment, cells were harvested and luciferase activity of cell lysates was measured using the Dual-Luciferase Reporter Assay System (Promega) in a GloMax 20/20 Luminometer (Promega) according to manufacturer instructions.

For Rb activity measurement, 2 different reporters were tested in parallel: pE2F-TA-Luc (Clontech/Takara) to measure E2F transcriptional activity (repressed by Rb), and pRb-TA-Luc (Clontech/Takara) (promoter element directly suppressed by Rb). 1×10^6 cells of control (WT) and clonally isolated Rb KO cell lines were electroporated with 3 µg of either pE2F-TA-luc or pRb-TA-luc and 0.3 ug renilla luciferase plasmid on an Amaxa Nucleofector 2b (Lonza). Following transfection, cells were grown in complete medium

623 for 24h followed by serum-free medium for 24h. Cells were then harvested and

624 luciferase activity measured as described above.

Next-generation sequencing of CRISPR repair products

72h after transfection, cells were harvested and genomic DNA was isolated with the

Wizard Genomic DNA Purification Kit (Promega). DNA concentration was measured on

a Nanodrop spectrophotometer and 100 ng of DNA per sample was PCR-amplified with

KAPA2G Robust HotStart ReadyMix (Roche) based on findings of low PCR bias for

KAPA polymerase[9394]. Primers targeted a conserved region surrounding *PTEN* exon 1

(Extended Data Figure 4a). PCR was performed according to manufacturer instructions,

with an annealing temperature of 66°C for 30 cycles. To purify samples for next-

generation sequencing, PCR products were electrophoresed on a 0.8% agarose gel

and post-stained with SYBR Gold Nucleic Acid Gel Stain (Thermo Fisher). Gels were

visualized on a blue light tray (BioRad) to minimize damage to DNA. A gel slice for each

lane was excised using a scalpel, and each slice was cut to include the region ranging

from just above the prominent *PTEN* PCR band down to and including the "primer

dimer" region to ensure inclusion of any deletion alleles. DNA was extracted from gel

slices using the QiaQuick Gel Extraction Kit (Qiagen), and triplicate PCR reaction

eluates per sample were pooled for sequencing. Sample concentrations were measured

by Nanodrop and adjusted as necessary prior to submission for 2x250 bp paired-end

Illumina MiSeq sequencing with target depth of >40,000 reads/sample (Genewiz).

Analysis of CRISPR NGS data

FASTQ files from each sequenced sample were analyzed with both CRISPResso2,[95]

which uses an alignment-based algorithm, and CRISPRPic,[96] which uses a kmer-based

algorithm. CRISPResso2 was run using the following parameters: window size = 30,

maximum paired-end overlap = 500, bp excluded from left and right ends = 15,

minimum alignment score = 50, minimum identity score = 50, plot window size = 20. For

CRISPRPic analysis, SeqPrep[97] was used to merge overlapping read pairs and trim

adapter sequences. CRISPRPic was run on merged FASTQ sequences for each

sample with the following parameters: index size = 8, window size = 30.

HPRT mutation assay

For the HPRT mutation assay, cells used were low-passage primary dermal fibroblasts

from multiple species that were known to originate from male animals, to ensure single

copy-number of the X-linked *HPRT* gene. Each species was tested with 3 different cell

lines from 3 individual animals. The bowhead *HPRT* coding sequence was BLASTed

against bowhead genome scaffolds[16] and neighboring gene sequences were analyzed

to confirm mammal-typical localization of *HPRT* on the bowhead X-chromosome. Cells

were cultured in standard fibroblast growth medium, but with FBS being replaced with

dialyzed FBS (Omega Scientific, Inc.) and supplemented with Fibroblast Growth Kit

Serum-Free (Lonza) to improve growth and viability in dialyzed FBS. Dialyzed FBS was

found in optimization experiments to be necessary for efficient 6-thioguanine selection.

Prior to mutagenesis, cells were cultured for 7 days in medium containing HAT

Supplement (Gibco) followed by 4 days in HT Supplement (Gibco) to eliminate any pre-

existing HPRT mutants. To induce mutation, cells were incubated for 3 hours in serum-

free MEM containing 150 µg/mL N-ethyl-N-nitrosourea (ENU) (Sigma). Cells were then

maintained in ENU-free medium for 9 days to allow mutations to establish and existing

HPRT to degrade. 1×10^6 cells from each cell line were harvested and plated in

dialyzed FBS medium containing 5 µg/mL 6-thioguanine (Chem-Impex), in parallel with

1×10^6 untreated control cells for each cell line. Cells were plated at a density of 1×10^5

cells per 15-cm dish to allow for efficient selection and colony separation, and to prevent

potential "metabolic cooperation"[37]. In tandem, for each cell line 200 cells from

untreated and control conditions were plated in triplicate 10-cm dishes in non-selective

medium to calculate plating efficiency. After 3 weeks of growth, surviving colonies were

fixed and stained with a crystal violet/glutaraldehyde solution as previously described[98].

Colonies were counted, and HPRT mutation rate was calculated as plating-efficiency

adjusted number of HPRT-negative colonies containing >50 cells. Appropriate

concentrations of ENU and 6-TG, as well as optimal plating densities and growth

conditions, were determined prior to the experiment described above through

optimization and dose titration experiments.

Digital droplet PCR measurement of CRISPR cleavage rate

A ddPCR assay similar to a previously published method[99] was used for time-course

quantification of CRISPR DSB induction across species. qPCR primers at conserved

sites flanking the guide RNA target site in the *PTEN* gene were designed such that

cleavage would prevent PCR amplification. As an internal copy number reference

control, a second set of previously validated qPCR primers targeting an ultraconserved

element present in all mammals as a single copy per genome (UCE.359) was designed

based on published sequences[100]. To allow for multiplexing and copy number

normalization of *PTEN* within each ddPCR reaction, 5' fluorescent hydrolysis probes

(FAM for *PTEN* and HEX for UCE.359) targeting conserved sequences were designed,

with 3' Iowa Black® and internal ZEN™ quenchers (Integrated DNA Technologies). All

primers and probes were checked for specificity by BLAST against each species'

genome.[100] Fibroblasts were transfected with *PTEN* CRISPR RNP as described in

"Next-generation sequencing of CRISPR repair products" and returned to cell culture

incubators. At the indicated times post-transfection, cells were harvested, flash frozen,

and genomic DNA was isolated with the Wizard Genomic DNA Purification Kit

(Promega). During isolation, newly lysed cells were treated with Proteinase K and

RNase A for 30 minutes each at 37°C to minimize the possibility of residual CRISPR

RNP activity. DNA concentration was measured on a Nanodrop spectrophotometer, and

genomic DNA was pre-digested with BamHI-HF (NEB) and XhoI (NEB), which do not

cut within target amplicons, to maximize PCR efficiency and distribution across droplets.

15 ng of genomic DNA per sample was added to duplicate PCR reactions using the

ddPCR™ Supermix for Probes (No dUTP) master mix (Bio-Rad). Droplets were

prepared and measured according to manufacturer instructions. Briefly, each 20 µL

reaction was mixed with 70 µL Droplet Generation Oil for Probes (Bio-Rad) and droplets

were formed in a QX100 Droplet Generator (Bio-Rad). 40 µL of droplets per reaction

were transferred to 96-well PCR plates and sealed with a PX1 PCR Plate Sealer (Bio-

Rad). The sealed plates were then subjected to PCR using a pre-optimized cycling

protocol. Following PCR, the plates were loaded into a QX100™ Droplet Reader (Bio-

Rad) and each droplet measured on both FAM and HEX channels. *PTEN* copy number

normalized to UCE.359 reference copy number within each well was determined with

QuantaSoft™ software (Bio-Rad). For each species, positive/negative gates in mock-

transfected control samples were adjusted as necessary to compensate for differences

in multiplex PCR efficiency/specificity and "rain" droplets between species and bring

normalized *PTEN* copy number closer to 1. The control gates were then applied across

all samples/time points within the same species and used for *PTEN* copy number

calculation.

Flow cytometric measurement of CRISPR RNP transfection efficiency

CRISPR RNP transfections were performed as described above, but with ATTO-550

fluorescently-labeled tracRNA (Integrated DNA Technologies). At 0h and 24h post-

transfection, cells were harvested, pelleted, and analyzed by flow cytometry on a

CytoFlex S Flow Cytometer (Beckman Coulter). Gain and ATTO-550 positive gates

were set based on mock-transfected control cells included in each experiment.

Senescence-associated β-galactosidase (SA-β-gal) staining

SA-β-gal staining was performed as previously described.[101,102] Cells were washed

twice with PBS and fixed in a solution containing 2% formaldehyde and 0.2%

glutaraldehyde in PBS for 5 min at room temperature. After fixation, cells were

immediately washed twice with PBS and stained in a solution containing 1 mg/mL 5-

bromo-4-chloro-3-indolyl P3-D-galactoside (X-Gal), 40 mM citric acid/sodium phosphate

buffer, pH 6.0, 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 150 mM

NaCl, and 2 mM $MgCl_2$. Plates were incubated at 37°C for 16 h without CO_2.

Colorimetric images were taken from different areas of each plate and quantified.

Cell survival assay

Percentage of live cells was quantified using the Annexin-V FLUOS Staining Kit (Roche)

following the manufacturer's instructions. After staining, cells were analyzed on a

CytoFlex S flow cytometer (Beckman Coulter).

p53 activity

To test p53 activity in cultured primary fibroblasts, 150,000 cells were seeded in 6-well

plates 1 day before transfection with 1 µg pp53-TA-Luc vector (Clontech) and 0.015 µg

pRL-CMV-Renilla (Promega) to normalize for transfection efficiency. Transfections were

performed using PEI MAX Transfection Grade Linear Polyethylenimine Hydrochloride

(MW 40,000) (Polysciences) according to manufacturer instructions. 24h after

transfections cells were lysed using 50µl passive lysis buffer (Promega) per 10^5 cells

and flash frozen/thawed two times in liquid nitrogen and a 37°C water bath. Luciferase

assays were performed using the Dual-Luciferase Reporter Assay System (Promega)

and program DLR-2-INJ on a Glomax 20/20 Luminometer (Promega) with 20µl cell

extract as the input.

Generation of NHEJ and HR reporter cell lines

NHEJ and HR reporter constructs[38] were digested with NheI restriction enzyme and

purified with the QIAEX II gel extraction kit (QIAGEN). The same plasmid DNA

preparation was used for generating all reporter cell lines of the studied species. Cells

PD < 15 were recovered from liquid nitrogen and passaged once before the integration

of the constructs. 0.25 µg of linearized NHEJ and HR constructs were electroporated

into one million cells for each cell line. Two days after transfection, media was

refreshed, and G418 was applied to select stable integrant clones. Triplicates of each

reporter in each cell line were prepared to obtain an adequate number of stable clones.

Clones from triplicate plates were pooled to get at least 50 clones per reporter per cell

line.

DSB repair assays and flow cytometry analysis

DSB repair assays were performed as previously described[103]. Briefly, growing cells

were co-transfected with 3 µg of plasmid encoding I-SceI endonuclease and 0.03 µg of

plasmid encoding DsRed. The same batch of I-SceI and DsRed mixture was used

throughout all species to avoid batch-to-batch variation. To test the effect of CIRBP on

DSB repair, 3 µg of CIRBP plasmids were co-transfected with I-SceI and DsRed

plasmids. Three days after transfection, the numbers of GFP+ and DsRed+ cells were

determined by flow cytometry on a CytoFlex S Flow Cytometer (Beckman Coulter). For

gating strategy see Supplementary Figure 4 in Supplementary Information. For each

sample, a minimum of 50,000 cells was analyzed. DSB repair efficiency was calculated

by dividing the number of GFP+ cells by the number of DsRed+ cells.

For knockdown experiments, bowhead whale cells containing the NHEJ reporter were

transfected with 120 pmol of anti-bwCIRBP or control siRNAs (Dharmacon) 3 days

before I-Sce1/DsRed transfections using an Amaxa Nucleofector (U-023 program). The

efficiency of knockdown was determined by Western blot.

For the extrachromosomal assay and fidelity analysis, NHEJ reporter plasmid was

digested with I-Sce1 for 6h and purified using a QIAEX II Gel Extraction Kit (QIAGEN).

Exponentially growing cells were transfected using an Amaxa nucleofector with the U-

023 program. In a typical reaction, 10^6 cells were transfected with 0.25 µg of

predigested NHEJ reporter substrate along with 0.025 μg of DsRed to serve as a

transfection control. 72h after transfection, cells were harvested and analyzed by flow

cytometry on a BD LSR II instrument. At least 20,000 cells were collected for each

sample. Immediately after FACS, genomic DNA was isolated from cells using the

QIAGEN Blood & Tissue kit. DSB repair sites in the NHEJ construct were amplified by

PCR using Phusion polymerase (NEB), cloned using the TOPO Blunt cloning kit (NEB),

and sent for Sanger sequencing. At least 100 sequenced clones were aligned and

analyzed using the ApE software.

Western blotting

All antibodies were checked for conservation of the target epitope in the protein

sequence of each included species, and only those targeting regions conserved across

these species were used. For a limited number of proteins where the available

antibodies with specific epitope information disclosed did not target conserved regions,

we selected antibodies based on demonstrated reactivity across a broad range of

mammal species and always confirmed these results with multiple antibodies.

Information on antibodies is provided in Supplementary Methods.

Exponentially growing cells were harvested with trypsin and counted, and 10^6 cells were

resuspended in 100 μL of PBS containing protease inhibitors. 100μL of 2x Laemmli

buffer (Bio-Rad) was added, and samples were boiled at 95ºC for 10 minutes. Samples

were separated with 4-20% gradient SDS-PAGE, transferred to a PVDF membrane,

and blocked in 5% milk-TBS-T for 2 hours at room temperature. Membranes were

incubated overnight at +4ºC with primary antibodies in 5% milk-TBS-T. After 3 washes

for 10 minutes with TBS-T, membranes were incubated for 1 hour at room temperature

with secondary antibodies conjugated with HRP or a fluorophore. After 3 washes with

TBS-T signal was developed for HRP secondaries with Clarity Western ECL Substrate

(Bio-Rad). CIRBP and RPA2 expression were each measured with 3 different

antibodies targeting conserved epitopes (Extended Data Figure 7c, d).

PARP activity

PARP activity was measured in cell nuclear extracts with the PARP Universal

Colorimetric Assay Kit (Trevigen) according to the manufacturer's instructions. Nuclear

extracts were prepared using EpiQuik Nuclear Extraction Kit (EpigenTek) following

manufacturer protocol. 2.5µg of total nuclear extract was added to measure PARP

activity.

For measurement of PARylation efficiency, cells were treated with 400µM H_2O_2 for 15

and 30 min. At the end of incubation, cells were placed on ice, washed once with PBS,

and lysed directly on a plate with 2x Laemmli buffer. Samples were boiled for 10min at

95ºC and processed by Western Blot.

Pulsed-field gel electrophoresis and analysis of DSBs

After irradiation and repair incubation, confluent human and bowhead whale skin

fibroblasts were harvested, ~400,000 cells were resuspended in PBS, mixed with an

equal volume of 1.4% low gelling temperature agarose and embedded into agarose

plugs. Plugs were kept for 1h at +4ºC and incubated in lysis solution 1 (0.5M EDTA, 2%

sodium sarcosine, 0.5 mg/ml Proteinase K) for 24h at +4ºC. Subsequently plugs were

placed into lysis solution 2 (1.85M NaCl, 0.15M KCl, 5mM $MgCl_2$, 2mM EDTA, 4mM Tris

pH 7.5, 0.5% TritonX100) for 40h at +4ºC. Plugs were then washed two times for 1h in

TE buffer and stored in TE buffer at +4ºC. PFGE was carried out with a CHEF DRII

system (Bio-Rad) in 0.8% agarose gels. The gels were run at 14°C with linearly

increasing pulse times from 50 to 5,000 s for 66 h at a field strength of 1.5 V/cm. Gels

were stained with 0.5mg/ml ethidium bromide for 4h, washed with TBE buffer and

imaged. Quantitative analysis was performed with Image Lab software (Bio-Rad). The

fraction of DNA entering the gel was quantified. Samples irradiated with various doses

and not incubated for repair served as a calibration to determine the percentage of

remaining DSBs in the repair samples from the fraction of DNA entering the gel.

Construction of lentiviral overexpression vectors and lentivirus production

The coding sequence of bwCIRBP was amplified by PCR using Phusion polymerase

(NEB), digested with EcoRI and NotI, and cloned in between EcoRI and NotI sites of the

Lego-iC2 plasmid. The sequence was verified by Sanger sequencing.

Lentiviral particles were produced in Lenti-X 293T cells (Takara). Approximately 10×10^6

cells were transfected with a mixture of pVSV-G (1.7 µg), psPAX2 (3.4 µg) and Lego-

iC2-bwCIRBP (6.8 µg) using PEI MAX (Polysciences). The day after transfection,

DMEM culture medium (ThermoFisher) was replaced with fresh medium and lentiviral

particles were harvested from the supernatant for the next 3 days.

Quantification of micronuclei

Two different approaches were used. To analyze binucleated cells containing

micronuclei (MN), 20,000 cells were plated per chamber slide 2d before ionizing

radiation. Immediately after irradiation (for human cells) or 2d after irradiation (for

bowhead whale cells), cytochalasin B was added to cell culture media at a final

concentration 0.5 µg/ml, and cells were incubated for an additional 72h. At the end of

incubation, cells were washed with PBS, incubated in 75mM KCl for 10min at RT, fixed

with ice-cold methanol for 1.5 min, air-dried, and stored. Immediately before the

analysis, cells were stained with 100 µg/ml acridine orange for 2min, washed with PBS,

mounted in PBS and analyzed by fluorescence microscopy. At least 1000 binucleated

cells were analyzed per sample.

For analysis of MN in single cells overexpressing bwCIRBP, exponentially growing cells

were transduced with LEGO-iC2-bwCIRBP lentivirus in the presence of 10 µg/ml

polybrene. Two days after transduction, cells were passaged, and irradiated the

following day. 3 days after irradiation, cells were fixed with 2% formaldehyde for 15 min,

washed with PBS, permeabilized with 0.2% TritonX100 for 2min, washed in PBS, and

mounted in VECTASHIELD Antifade Mounting Medium with DAPI. Cells were analyzed

with a fluorescent microscope. At least 1000 cells were analyzed per condition.

Host cell reactivation assay

A host cell reactivation assay was used to measure the repair of UV-induced DNA

damage through nucleotide excision repair as previously described.[33]

To measure the repair of oxidative DNA damage (base excision repair), 20 µg of firefly

luciferase (FFL) plasmid was mixed with 20-200 µM methylene blue (MB), and water

was added to a final volume of 0.4ml. DNA-MB mixture was dropped onto a petri dish

and placed on ice. An additional petri dish containing water was placed on top. The

DNA-MB mixture was exposed to visible light for 15 min using a 100W lamp at an 11cm

distance. Damaged DNA was purified, and the host cell reactivation assay was

performed as described for UV-induced DNA damage.

Cyclobutane pyrimidine dimer (CPD) ELISA

Human and bowhead whale skin fibroblasts were grown until they reached confluency

before UVC radiation. Cells were irradiated in PBS at doses of 0, 5, 10, 20 and 30J/m^2

and harvested immediately for building an induction curve. To assess DNA repair, cells

were irradiated at 30 J/m^2 and incubated for 6, 24 and 48h before harvesting. Genomic

DNA was isolated using QIAamp Blood Kit (Qiagen). Samples of DNA were diluted in

PBS to a final concentration of 2 µg/ml, denatured at 100ºC for 10min and incubated in

an ice bath for 15 min. 100 ng of denatured DNA solution was applied to ELISA plate

wells precoated with protamine sulfate (Cosmo Bio) and dried O/N at 37ºC. Plates were

washed five times with PBS supplemented with 0.05% Tween-20 (PBS-T) and blocked

in 2% FBS in PBS-T for 30min at 37ºC. After 5 washes with PBS-T, plates were

incubated with mouse monoclonal anti-CPD antibodies (Clone TDM-2, 1:1000) in PBS

for 30 min at 37ºC. Plates were then sequentially incubated with goat-anti mouse biotin

IgG (Invitrogen, 1:1000) and streptavidin-HRP (Invitrogen, 1:5000) in PBS for 30 min at

37ºC each with 5 washes with PBS-T before and after incubations. Plates were washed

with citrate buffer and incubated with substrate solution (citrate buffer/o-

phenylenediamine/hydrogen peroxide) for 30 min at 37ºC. Finally, the reaction was

stopped with 2M H_2SO_4 and the absorbance was measured at 492 nm with a plate

reader.

CIRBP variant sequence analysis

Identification of rare codons (<10% usage for the corresponding amino acid in human

CDS sequences) was performed on CIRBP coding sequences using the Benchling

Codon Optimization Tool.[104] Codon adaptation index (CAI) was calculated with human

codon frequencies using the E-CAI web server.[53]

RNA isolation and RNA-seq analysis

RNA from exponentially growing or senescent human and bowhead whale primary skin

fibroblasts was isolated using the Quick-RNA MiniPrep kit (Zymo Research) according

to manufacturer instructions.

The RNA-seq reads were first processed using Trim_Galore (version 0.6.6). Cleaned

RNA-seq reads were used to quantify gene expression with Salmon (version 1.4.0).[105]

Transcript FASTA files of species were from our published de novo transcriptome

assemblies.[106] Transcript FASTA files were indexed with the command "Salmon index"

with default parameters. Cleaned RNA-seq reads were aligned to the indexed transcript

with the specific parameters (--useVBOpt --seqBias --gcBias), which were set for

sequence-specific bias correction and fragment GC bias correction. The read counts of

genes obtained from Salmon were further normalized by trimmed mean of M-values

(TMM)[107] and median of ratios methods.[108]

LC-MS proteomic analysis of fibroblasts

2 15-cm dishes of growing primary fibroblasts from 2 cell lines for each species were

harvested for protein. Cells were washed with PBS and pellets were snap frozen and

stored in liquid nitrogen until processing. Cells were solubilized with 5% SDS; 50 mM

TEAB pH 7, and sonicated at 8°C with 10x 45s pulses using 30% power with 15 s rest

between each pulse with a cup horn Q800R3 Sonicator System (Qsonica; Newtown

CT). Soluble proteins were reduced with 10 mM DTT for 30 min at 55°C, followed by

alkylation with 15 mM iodoacetamide at 25°C in the dark for 30 min. S-trap micro

columns (Protifi; Farmingdale, NY) were employed after this step for overnight tryptic

digestion and peptide isolation according to manufacturer instructions. All solvents were

MS-grade. Resulting tryptic peptides were resuspended in MS-grade water and were

quantified using a Pierce™ Quantitative Fluorometric Peptide Assay (Thermo Fisher cat

#23290). Prior to MS, peptides were mixed with a common internal retention time

standards[109] (CiRT) peptide mix (50 fmol CiRT/2ug total tryptic peptides) and

acetonitrile (ACN) and formic acid were added to concentrations of 5% and 0.2%

respectively. The final concentration of the peptide mix was 0.5µg/µl. 2ug (4ul) of each

were resolved by nano-electrospray ionization on an Orbitrap Fusion Lumos MS

instrument (Thermo) in positive ion mode. A 30 cm home-made column packed with 1.8

µm C-18 beads was employed to resolve the peptides. Solvent A was 0.1% formic acid

and solvent B was 80% acetonitrile with 0.1% formic acid and flow rate was 300 nl/min.

The length of the run was 3 h with a 155 min gradient from 10-38% B. HCD (30%

collision energy) was used for MS2 fragmentation and dynamic exclusion was operative

after a single time and lasted for 30s. Other details of the run parameters may be found

in the embedded run report of the RAW data file uploaded to the ProteomeXchange

database. Peptide assignments and quantitation were done using the label-free quant

match between runs (LFQ-MBR) workflow of MSFragger[110–112]. MaxLFQ with a

minimum of two ions was implemented and normalization was selected. Additional

details are available in MSFragger log files. Searches were performed within the

Philosopher/Fragpipe pipeline that incorporates PeptideProphet and ProteinProphet

filtering steps to increase the likelihood of correct assignments.[113] The databases used

for searches were predicted proteins from the published bowhead genome[114] as well as

our custom proteome derived from our de novo sequenced and Trinity[115,116]-assembled

pool of transcriptomes from whale tissues. Human (UP000005640), mouse

(UP000000589), and bovine (UP000009136) databases were from the latest build

available from Uniprot.[117] For the searches, databases also included a reverse

complement form of all peptides as well as common contaminants to serve as decoys

for false discovery rate (FDR) calculation by the target/decoy approach (decoy present

at 50%).[118] Final FDR was below 1%.

Statistical analyses

Statistical comparisons were performed as indicated in the figure legends. Unless

otherwise specified in the text or legend, *n* refers to separate biological replicate cell

lines, isolated from different individuals for a given species. Exceptions include specific

genetically modified cell lines or clones, e.g. tumor suppressor knockout lines and Ku-

deficient MEFs. In such cases, *n* refers to technical replicates and indicates the number

of times the experiment was repeated with the specified cell line. Details for

comparisons done by ANOVA are included in Supplementary Information.

Data Availability

Proteomics data are accessible through ProteomeXchange [URL to be added]. DNA

sequence (CRISPR repair fidelity analysis) and RNA sequence (transcriptome) data are

accessible through NCBI Sequence Read Archive (SRA) [URL to be added].

References

1.   Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic Acids*
Res **46**, D1083–D1090 (2018).

2.   Peto, R. Epidemiology, multistage models, and short-term mutagenicity tests. in *The origins of*
human cancer (eds. Hiatt, H., Watson, J. & Winsten, J.) vol. 4 1403–1428 (Cold Spring Harbor
Conferences on Cell Proliferation, 1977).

3.   Abegglen, L. M. *et al.* Potential Mechanisms for Cancer Resistance in Elephants and Comparative
Cellular Response to DNA Damage in Humans. *JAMA* **314**, 1850 (2015).

4.   Vazquez, J. M., Sulak, M., Chigurupati, S. & Lynch, V. J. A Zombie LIF Gene in Elephants Is
Upregulated by TP53 to Induce Apoptosis in Response to DNA Damage. *Cell Rep* **24**, 1765–1776
(2018).

5.   Sulak, M. *et al.* TP53 copy number expansion is associated with the evolution of increased body
size and an enhanced DNA damage response in elephants. *Elife* **5**, (2016).

6.   TEK & Bowhead Life History and Longevity - The North Slope Borough. https://www.north-
slope.org/departments/wildlife-management/studies-research-projects/bowhead-
whales/traditional-ecological-knowledge-of-bowhead-whales/tek-bowhead-life-history-and-
longevity/.

7.   George, J. C. & Bockstoce, J. R. Two historical weapon fragments as an aid to estimating the
longevity and movements of bowhead whales. *Polar Biol* **31**, 751–754 (2008).

8.   Rosa, C. *et al.* Age estimates based on aspartic acid racemization for bowhead whales (Balaena
mysticetus) harvested in 1998-2000 and the relationship between racemization rate and body
temperature. *Mar Mamm Sci* **29**, 424–445 (2013).

9.   Philo, L. M., Shotts, E. B. & George, J. C. Morbidity and mortality- The bowhead whale. *Soc. Mar.*
Mammal., Spec. Publ 275–312 (1993).

10.   George, J. C. *et al.* Age and growth estimates of bowhead whales (Balaena mysticetus) via
aspartic acid racemization. *Can J Zool* **77**, 571–580 (1999).

11.   Wetzel, D. *et al.* Age estimation for bowhead whales, Balaena mysticetus, using aspartic acid
racemization with enhanced hydrolysis and derivatization procedures. *Paper SC/65b/BRG05*
presented to the Scientific Committee of the International Whaling Commission (2014).

12.   George, J. C. *et al.* A new way to estimate the age of bowhead whales (Balaena mysticetus) using
ovarian corpora counts. *https://doi.org/10.1139/z11-057* **89**, 840–852 (2011).

13.   Boddy, A. M. *et al.* Lifetime cancer prevalence and life history traits in mammals. *Evol Med Public*
Health **2020**, 187–195 (2020).

14.   Vincze, O. *et al.* Cancer risk across mammals. *Nature* **601**, 263–267 (2022).

15.   Tollis, M., Boddy, A. M. & Maley, C. C. Peto's Paradox: how has evolution solved the problem of
cancer prevention? *BMC Biol* **15**, (2017).

16. Keane, M. *et al.* Insights into the evolution of longevity from the bowhead whale genome. *Cell*
Rep **10**, 112–122 (2015).

17. Tollis, M. *et al.* Return to the Sea, Get Huge, Beat Cancer: An Analysis of Cetacean Genomes
Including an Assembly for the Humpback Whale (Megaptera novaeangliae). *Mol Biol Evol* **36**,
1746–1763 (2019).

18. Seim, I. *et al.* The transcriptome of the bowhead whale Balaena mysticetus reveals adaptations of
the longest-lived mammal. *Aging* **6**, 879–899 (2014).

19. Armitage, P. & Doll, R. The age distribution of cancer and a multi-stage theory of carcinogenesis.
Br J Cancer **8**, 1–12 (1954).

20. Rangarajan, A., Hong, S. J., Gifford, A. & Weinberg, R. A. Species- and cell type-specific
requirements for cellular transformation. *Cancer Cell* **6**, 171–183 (2004).

21. Vazquez, J. M., Sulak, M., Chigurupati, S. & Lynch, V. J. A Zombie LIF Gene in Elephants Is
Upregulated by TP53 to Induce Apoptosis in Response to DNA Damage. *Cell Rep* **24**, 1765–1776
(2018).

22. Vazquez, J. M. & Lynch, V. J. Pervasive duplication of tumor suppressors in Afrotherians during
the evolution of large bodies and reduced cancer risk. *Elife* **10**, 1–45 (2021).

23. Vazquez, J. M. *et al.* Parallel evolution of reduced cancer risk and tumor suppressor duplications
in Xenarthra. *Elife* **11**, (2022).

24. Boveri, T. Concerning the origin of malignant tumours by Theodor Boveri. Translated and
annotated by Henry Harris. *J Cell Sci* **121**, 1–84 (2008).

25. Hasty, P., Campisi, J., Hoeijmakers, J., van Steeg, H. & Vijg, J. *Aging and Genome Maintenance:*
Lessons from the Mouse? http://science.sciencemag.org/.

26. Kashiyama, K. *et al.* Malfunction of nuclease ERCC1-XPF results in diverse clinical manifestations
and causes Cockayne syndrome, xeroderma pigmentosum, and Fanconi anemia. *Am J Hum Genet*
92, 807–819 (2013).

27. Navarro, C. L., Cau, P. & Lévy, N. Molecular bases of progeroid syndromes. *Human Molecular*
Genetics vol. 15 R151–R161 Preprint at https://doi.org/10.1093/hmg/ddl214 (2006).

28. Varon, R. *et al.* Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen
breakage syndrome. *Cell* **93**, 467–476 (1998).

29. Liu, B. *et al.* Genomic instability in laminopathy-based premature aging. *Nat Med* **11**, 780–785
(2005).

30. Lorenzini, A. *et al.* Significant Correlation of Species Longevity with DNA Double Strand Break-
Recognition but not with Telomere Length. *Mech Ageing Dev* **130**, 784 (2009).

31. Zhang, L. *et al.* Maintenance of genome sequence integrity in long- And short-lived rodent
species. *Sci Adv* **7**, (2021).

32. Cagan, A. *et al.* Somatic mutation rates scale with lifespan across mammals. *Nature 2022* 1–8
(2022) doi:10.1038/s41586-022-04618-z.

33. Tian, X. *et al.* SIRT6 Is Responsible for More Efficient DNA Double-Strand Break Repair in Long-
Lived Species. *Cell* **177**, 622-638.e22 (2019).

34. Fortunato, A., Fleming, A., Aktipis, A. & Maley, C. C. Upregulation of DNA repair genes and cell
extrusion underpin the remarkable radiation resistance of Trichoplax adhaerens. *PLoS Biol* **19**,
(2021).

35. Tian, X. *et al.* Evolution of telomere maintenance and tumour suppressor mechanisms across
mammals. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**, (2018).

36. Borowicz, S. *et al.* The soft agar colony formation assay. *J Vis Exp* e51998 (2014)
doi:10.3791/51998.

37. Johnson, G. E. Mammalian Cell HPRT Gene Mutation Assay: Test Methods. in 55–67 (2012).
doi:10.1007/978-1-61779-421-6_4.

38. Seluanov, A., Mao, Z. & Gorbunova, V. Analysis of DNA Double-strand Break (DSB) Repair in
Mammalian Cells. *J Vis Exp* 43 (2010) doi:10.3791/2002.

39. Guo, T. *et al.* Harnessing accurate non-homologous end joining for efficient precise deletion in
CRISPR/Cas9-mediated genome editing. *Genome Biol* **19**, (2018).

40. Richardson, C. D., Ray, G. J., Bray, N. L. & Corn, J. E. Non-homologous DNA increases gene
disruption efficiency by altering DNA repair outcomes. *Nature Communications 2016 7:1* **7**, 1–7
(2016).

41. Chen, J. K., Lin, W. L., Chen, Z. & Liu, H. wen. PARP-1–dependent recruitment of cold-inducible
RNA-binding protein promotes double-strand break repair and genome stability. *Proc Natl Acad*
Sci U S A **115**, E1759–E1768 (2018).

42. Nishiyama, H. *et al.* A glycine-rich RNA-binding protein mediating cold-inducible suppression of
mammalian cell growth. *J Cell Biol* **137**, 899–908 (1997).

43. Sheikh, M. S. *et al.* Identification of several human homologs of hamster DNA damage-inducible
transcripts. Cloning and characterization of a novel UV-inducible cDNA that codes for a putative
RNA-binding protein. *J Biol Chem* **272**, 26720–26726 (1997).

44. Wellmann, S. *et al.* Oxygen-regulated expression of the RNA-binding proteins RBM3 and CIRP by
a HIF-1-independent mechanism. *J Cell Sci* **117**, 1785–1794 (2004).

45. Yang, C. & Carrier, F. The UV-inducible RNA-binding protein A18 (A18 hnRNP) plays a protective
role in the genotoxic stress response. *J Biol Chem* **276**, 47277–47284 (2001).

46. Yang, R., Weber, D. J. & Carrier, F. Post-transcriptional regulation of thioredoxin by the stress
inducible heterogenous ribonucleoprotein A18. *Nucleic Acids Res* **34**, 1224–1236 (2006).

47.   Yang, R. *et al.* Functional significance for a heterogenous ribonucleoprotein A18 signature RNA
1061   motif in the 3'-untranslated region of ataxia telangiectasia mutated and Rad3-related (ATR)
1062   transcript. *J Biol Chem* **285**, 8887–8893 (2010).

1063   48.   Roilo, M., Kullmann, M. K. & Hengst, L. Cold-inducible RNA-binding protein (CIRP) induces
1064   translation of the cell-cycle inhibitor p27Kip1. *Nucleic Acids Res* **46**, 3198–3210 (2018).

1065   49.   Elsner, R., Meiselman, H. J. & Baskurt, O. K. TEMPERATURE-VISCOSITY RELATIONS OF BOWHEAD
1066   WHALE BLOOD: A POSSIBLE MECHANISM FOR MAINTAINING COLD BLOOD FLOW. *Mar Mamm Sci*
1067   **20**, 339–344 (2004).

1068   50.   George, J. C. Growth, morphology and energetics of bowhead whales (Balaena mysticetus).
1069   (University of Alaska Fairbanks, 2009).

1070   51.   Gotic, I. *et al.* Temperature regulates splicing efficiency of the cold-inducible RNA-binding protein
1071   gene Cirbp. *Genes Dev* **30**, 2005–2017 (2016).

1072   52.   Al-Fageeh, M. B. & Smales, C. M. Cold-inducible RNA binding protein (CIRP) expression is
1073   modulated by alternative mRNAs. *RNA* **15**, 1165–1176 (2009).

1074   53.   Puigbò, P., Bravo, I. G. & Garcia-Vallvé, S. E-CAI: a novel server to estimate an expected value of
1075   Codon Adaptation Index (eCAI). *BMC Bioinformatics* **9**, (2008).

1076   54.   Zou, Y., Liu, Y., Wu, X. & Shell, S. M. Functions of Human Replication Protein A (RPA): From DNA
1077   Replication to DNA Damage and Stress Responses. *J Cell Physiol* **208**, 267 (2006).

1078   55.   Wold, M. S. Replication protein A: a heterotrimeric, single-stranded DNA-binding protein
1079   required for eukaryotic DNA metabolism. *Annu Rev Biochem* **66**, 61–92 (1997).

1080   56.   Paulsen, R. D. *et al.* A genome-wide siRNA screen reveals diverse cellular processes and pathways
1081   that mediate genome stability. *Mol Cell* **35**, 228–239 (2009).

1082   57.   Toledo, L. I. *et al.* ATR prohibits replication catastrophe by preventing global exhaustion of RPA.
1083   *Cell* **155**, 1088 (2013).

1084   58.   Velegzhaninov, I. O. *et al.* Radioresistance, DNA Damage and DNA Repair in Cells With Moderate
1085   Overexpression of RPA1. *Front Genet* **11**, 855 (2020).

1086   59.   Guven, M., Brem, R., Macpherson, P., Peacock, M. & Karran, P. Oxidative damage to RPA limits
1087   the nucleotide excision repair capacity of human cells. *J Invest Dermatol* **135**, 2834 (2015).

1088   60.   Graziano, S. *et al.* Lamin A/C recruits ssDNA protective proteins RPA and RAD51 to stalled
1089   replication forks to maintain fork stability. *Journal of Biological Chemistry* **297**, 101301 (2021).

1090   61.   Perrault, R., Cheong, N., Wang, H., Wang, H. & Iliakis, G. RPA facilitates rejoining of DNA double-
1091   strand breaks in an in vitro assay utilizing genomic DNA as substrate. *int. j. radiat. biol* **77**, 593–
1092   607 (2001).

1093   62.   Krasner, D. S., Daley, J. M., Sung, P. & Niu, H. Interplay between Ku and Replication Protein A in
1094   the Restriction of Exo1-mediated DNA Break End Resection. *J Biol Chem* **290**, 18806–18816
1095   (2015).

1096   63.   Chen, H., Lisby, M. & Symington, L. S. RPA coordinates DNA end resection and prevents
1097   formation of DNA hairpins. *Mol Cell* **50**, 589–600 (2013).

1098   64.   Deng, S. K., Gibb, B., de Almeida, M. J., Greene, E. C. & Symington, L. S. RPA Antagonizes
1099   Microhomology-Mediated Repair of DNA Double-Strand Breaks. *Nat Struct Mol Biol* **21**, 405
1100   (2014).

1101   65.   Guven, M., Brem, R., Macpherson, P., Peacock, M. & Karran, P. Oxidative damage to RPA limits
1102   the nucleotide excision repair capacity of human cells. *J Invest Dermatol* **135**, 2834 (2015).

1103   66.   Kim, K. *et al.* Novel Checkpoint Response to Genotoxic Stress Mediated by Nucleolin-Replication
1104   Protein A Complex Formation. *Mol Cell Biol* **25**, 2463 (2005).

1105   67.   Bélanger, F. *et al.* Mutations in Replicative Stress Response Pathways Are Associated with S
1106   Phase-specific Defects in Nucleotide Excision Repair *. (2015) doi:10.1074/jbc.M115.685883.

1107   68.   Shuck, S. C. & Turchi, J. J. Targeted inhibition of RPA reveals cytotoxic activity, synergy with
1108   chemotherapeutic DNA damaging agents and insight into cellular function. *Cancer Res* **70**, 3189
1109   (2010).

1110   69.   White, R. R. *et al.* Double-Strand Break Repair by Interchromosomal Recombination: An In Vivo
1111   Repair Mechanism Utilized by Multiple Somatic Tissues in Mammals. *PLoS One* **8**, 84379 (2013).

1112   70.   Seluanov, A. *et al.* Telomerase activity coevolves with body mass not lifespan. *Aging Cell* **6**, 45–52
1113   (2007).

1114   71.   Vazquez, J. M., Kraft, M. & Lynch, V. J. A CDKN2C retroduplication in Bowhead whales is
1115   associated with the evolution of extremely long lifespans and alerted cell cycle dynamics. *bioRxiv*
1116   2022.09.07.506958 (2022) doi:10.1101/2022.09.07.506958.

1117   72.   Coppé, J. P., Desprez, P. Y., Krtolica, A. & Campisi, J. The Senescence-Associated Secretory
1118   Phenotype: The Dark Side of Tumor Suppression. *Annu Rev Pathol* **5**, 99 (2010).

1119   73.   Mendiratta, G. *et al.* Cancer gene mutation frequencies for the U.S. population. *Nature
1120   Communications 2021 12:1* **12**, 1–11 (2021).

1121   74.   Wright, K. M. & Deshmukh, M. Restricting Apoptosis for Postmitotic Cell Survival and its
1122   Relevance to Cancer. *http://dx.doi.org/10.4161/cc.5.15.3129* **5**, 1616–1620 (2006).

1123   75.   Lieber, M. R., Gu, J., Lu, H., Shimazaki, N. & Tsai, A. G. Nonhomologous DNA end joining (NHEJ)
1124   and chromosomal translocations in humans. *Subcell Biochem* **50**, 279–296 (2014).

1125   76.   Yoshioka, K. I., Kusumoto-Matsuo, R., Matsuno, Y. & Ishiai, M. Genomic Instability and Cancer
1126   Risk Associated with Erroneous DNA Repair. *International Journal of Molecular Sciences 2021,
1127   Vol. 22, Page 12254* **22**, 12254 (2021).

1128   77.   Yokoyama, A. *et al.* Age-related remodelling of oesophageal epithelia by mutated cancer drivers.
1129   *Nature* **565**, 312–317 (2019).

1130   78.   Martincorena, I. *et al.* Somatic mutant clones colonize the human esophagus with age. *Science*
1131   **362**, 911–917 (2018).

79. Jaiswal, S. *et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J*
Med **371**, 2488–2498 (2014).

80. Martincorena, I. *et al.* High burden and pervasive positive selection of somatic mutations in
normal human skin. *Science (1979)* **348**, 880–886 (2015).

81. Mitchell, E. *et al.* Clonal dynamics of haematopoiesis across the human lifespan. *Nature 2022*
606:7913 **606**, 343–350 (2022).

82. Kolora, S. R. R. *et al.* Origins and evolution of extreme life span in Pacific Ocean rockfishes.
Science **374**, 842–847 (2021).

83. Allen, K. N. & Vázquez-Medina, J. P. Natural Tolerance to Ischemia and Hypoxemia in Diving
Mammals: A Review. *Front Physiol* **10**, 1199 (2019).

84. Lombardi, G., Ziemann, E. & Banfi, G. Whole-Body Cryotherapy in Athletes: From Therapy to
Stimulation. An Updated Review of the Literature. *Front Physiol* **8**, 258 (2017).

85. Gocoł, R. *et al.* The Role of Deep Hypothermia in Cardiac Surgery. *Int J Environ Res Public Health*
18, (2021).

86. Liu, Y. *et al.* Chronic hypoxia–induced Cirbp hypermethylation attenuates hypothermic
cardioprotection via down-regulation of ubiquinone biosynthesis. *Sci Transl Med* **11**, (2019).

87. Wu, L. *et al.* Therapeutic Hypothermia Enhances Cold-Inducible RNA-Binding Protein Expression
and Inhibits Mitochondrial Apoptosis in a Rat Model of Cardiac Arrest. *Mol Neurobiol* **54**, 2697–
2705 (2017).

88. Yu, L., Gu, T., Liu, Y., Jiang, X. & Shi, E. Cold-inducible ribonucleic acid-binding protein attenuates
acute kidney injuries after deep hypothermic circulatory arrest in rats. *Interact Cardiovasc Thorac*
Surg **26**, 124–130 (2018).

89. Li, Y. *et al.* Cold-inducible RNA-binding protein maintains intestinal barrier during deep
hypothermic circulatory arrest. *Interact Cardiovasc Thorac Surg* **29**, 583–591 (2019).

90. Pan, H. Z. *et al.* Cold-inducible RNA binding protein agonist enhances the cardioprotective effect
of UW solution during extended heart preservation. *Artif Organs* **44**, E406–E418 (2020).

91. Chen, Y. *et al.* Fight to the bitter end: DNA repair and aging. *Ageing Res Rev* **64**, 101154 (2020).

92. Seluanov, A., Vaidya, A. & Gorbunova, V. Establishing Primary Adult Fibroblast Cultures From
Rodents. *J Vis Exp* **44**, (2010).

93. Quail, M. A. *et al.* Optimal enzymes for amplifying sequencing libraries. *Nat Methods* **9**, 10–11
(2012).

94. Blackburn, M. C. Development of new tools and applications for high-throughput sequencing of
microbiomes in environmental or clinical samples. (2010).

95. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence analysis.
Nature Biotechnology 2019 37:3 **37**, 224–226 (2019).

96.   Lee, H. J., Chang, H. Y., Cho, S. W. & Ji, H. P. CRISPRpic: fast and precise analysis for CRISPR-
1168   induced mutations via prefixed index counting. *NAR Genom Bioinform* **2**, (2020).

1169   97.   GitHub - jstjohn/SeqPrep: Tool for stripping adaptors and/or merging paired reads with overlap
1170   into single reads. https://github.com/jstjohn/SeqPrep.

1171   98.   Franken, N. A. P., Rodermond, H. M., Stap, J., Haveman, J. & van Bree, C. Clonogenic assay of cells
1172   in vitro. *Nat Protoc* **1**, 2315–2319 (2006).

1173   99.   Rose, J. C. *et al.* Rapidly inducible Cas9 and DSB-ddPCR to probe editing kinetics. *Nat Methods* **14**,
1174   891 (2017).

1175   100.   Hudon, S. F. *et al.* Primers to highly conserved elements optimized for qPCR-based telomere
1176   length measurement in vertebrates. *Mol Ecol Resour* **21**, 59–67 (2021).

1177   101.   Dimri, G. P. *et al.* A biomarker that identifies senescent human cells in culture and in aging skin in
1178   vivo. *Proc Natl Acad Sci U S A* **92**, 9363 (1995).

1179   102.   Debacq-Chainiaux, F., Erusalimsky, J. D., Campisi, J. & Toussaint, O. Protocols to detect
1180   senescence-associated beta-galactosidase (SA-βgal) activity, a biomarker of senescent cells in
1181   culture and in vivo. *Nature Protocols 2009 4:12* **4**, 1798–1806 (2009).

1182   103.   Mao, Z. *et al.* SIRT6 promotes DNA repair under stress by activating PARP1. *Science (1979)* **332**,
1183   1443–1446 (2011).

1184   104.   Benchling. https://www.benchling.com/.

1185   105.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-
1186   aware quantification of transcript expression. *Nature Methods 2017 14:4* **14**, 417–419 (2017).

1187   106.   Lu, J. Y. *et al.* Comparative transcriptomics reveals circadian and pluripotency networks as two
1188   pillars of longevity regulation. *Cell Metab* **34**, 836-856.e5 (2022).

1189   107.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
1190   expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

1191   108.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
1192   seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).

1193   109.   Escher, C. *et al.* Using iRT, a normalized retention time for more targeted measurement of
1194   peptides. *Proteomics* **12**, 1111–1121 (2012).

1195   110.   Kong, A. T., Leprevost, F. v., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I. MSFragger:
1196   ultrafast and comprehensive peptide identification in mass spectrometry–based proteomics.
1197   *Nature Methods 2017 14:5* **14**, 513–520 (2017).

1198   111.   Teo, G. C., Polasky, D. A., Yu, F. & Nesvizhskii, A. I. Fast Deisotoping Algorithm and Its
1199   Implementation in the MSFragger Search Engine. *J Proteome Res* **20**, 498–505 (2021).

1200 112. Yu, F., Haynes, S. E. & Nesvizhskii, A. I. IonQuant enables accurate and sensitive label-free
1201    quantification with FDR-controlled match-between-runs. *Molecular and Cellular Proteomics* **20**,
1202    100077 (2021).

1203 113. da Veiga Leprevost, F. *et al.* Philosopher: a versatile toolkit for shotgun proteomics data analysis.
1204    *Nature Methods 2020 17:9* **17**, 869–870 (2020).

1205 114. The Bowhead Whale Genome Resource. http://www.bowhead-whale.org/.

1206 115. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity
1207    platform for reference generation and analysis. *Nature Protocols 2013 8:8* **8**, 1494–1512 (2013).

1208 116. Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a reference
1209    genome. *Nature Biotechnology 2011 29:7* **29**, 644–652 (2011).

1210 117. Bateman, A. *et al.* UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res* **49**,
1211    D480–D489 (2021).

1212 118. Elias, J. E. & Gygi, S. P. Target-decoy search strategy for increased confidence in large-scale
1213    protein identifications by mass spectrometry. *Nature Methods 2007 4:3* **4**, 207–214 (2007).

1214

Figure 1. Bowhead whale fibroblasts exhibit senescence with reduced SASP and low p53 activity. a, Growth curves of primary and hTERT-immortalized skin fibroblasts

(n=2 for each cell line). **b**, Telomerase activity and telomere length in skin fibroblasts. **c**, Quantification of β-gal–positive human and bowhead skin fibroblasts in response to γ-irradiation (12 days) and replicative senescence (n=3 for each species). **d**, Representative images of SA-β-gal staining of human and bowhead skin fibroblasts in response to γ-irradiation and replicative senescence. The bar is 100 μm. **e**, Apoptosis of human and bowhead whale fibroblasts in response to γ-irradiation. Three days after γ-irradiation, cells were harvested and subjected to an Annexin V apoptosis assay using flow cytometry (n=3 for each species). **f**, Log fold change of SASP mRNA expression in human and bowhead whale skin fibroblasts 12 days after γ-irradiation. **g**, Relative luciferase expression in mouse, cow, human and bowhead whale fibroblasts transfected with the p53 reporter vector. Data are shown as ratios of firefly/renilla luciferase (to normalize for transfection efficiency) expression 24 h after transfections (n=3 for mouse, human, BW; n=2 for cow). **h**, Apoptosis of mouse, cow, human and bowhead whale fibroblasts in response to UVC. Two days after UVC, cells were harvested and subjected to an Annexin V apoptosis assay using flow cytometry. Error bars represent mean ± SD. * $p<0.05$, ** $p<0.01$, *** $p<0.001$. Two-tailed t-test was used to quantify the significance. Whale, bowhead whale; NMR, naked mole-rat.

hTERT	+	+	+	+	+	+
H-Ras^{G12V}	-	-	+	+	+	+
p53	WT	-	-	WT	-	-
Rb	WT	-	WT	WT	-	-
PP2A/PTEN	WT	WT	WT	WT	WT	-

a

SV40 LT LTK1 LT$^{Δ434-444}$ LT LT + ST

TP53$^{-/-}$RB1$^{-/-}$ *TP53$^{-/-}$* *TP53$^{-/-}$* + LT$^{Δ434-444}$ LT + ST

TP53$^{-/-}$RB1$^{-/-}$

LT

b

Tumor Volume (mm^3)

Days Post-Injection

LT^{K1} LT$^{Δ434-444}$ TP53$^{-/-}$ LT$^{Δ434-444}$ LT   TP53$^{-/-}$ RB$^{-/-}$ TP53$^{-/-}$ RB$^{-/-}$ PTEN$^{-/-}$

c

	WT	p53$^{-/-}$	p53$^{-/-}$
p53			
β-actin			

d

	WT	Rb$^{-/-}$		Rb$^{-/-}$
Rb				
β-actin				

Figure 2. Fewer hits are required for oncogenic transformation of bowhead whale fibroblasts than for human fibroblasts. a, Images of representative fibroblast colonies for tested cell lines after 4 weeks of growth in soft agar. The top panel indicates whether the cell lines in the column below have the indicated protein overexpressed (+), inactivated (-), or expressed in the active endogenous form (WT). Text above individual images indicate for that cell line whether tumor suppressors are inactivated through genetic knockout or SV40 Large T (or LT mutants) or Small T (ST) antigen. Icons in corners of images indicate species. Scale bar is 250 µm. **b**, Volumetric growth curves for the indicated bowhead whale fibroblast cell lines in mouse xenograft assays. All cell lines shown stably express H-Ras^{G12V} and *hTERT* in addition to the genotype indicated in the figure legend. Data points represent averages from 3 immunodeficient nude mice injected bilaterally (6 injections) for each cell line, except for *TP53$^{-/-}$RB1$^{-/-}$* double knockouts, for which 2 independent cell lines were tested, for a total of 6 mice/12 injections. Experiments were terminated based on endpoints for maximum tumor size or duration of experiment as described in Methods. Images in the legend show a representative mouse for the indicated cell line at the final measured time point. Error bars show SEM. **c**, Western blot for p53 protein in clonally isolated fibroblast colonies following CRISPR targeting of *TP53*. Underlined lanes indicate colonies selected for further validation and experiments. **d**, Western blot for Rb protein in fibroblast clones following CRISPR targeting of *RB1* on an p53 knockout background.

Figure 3. The bowhead whale repairs DSBs with higher accuracy and efficiency than other mammals. a, NHEJ efficiency was measured using fluorescent reporter construct where GFP gene is interrupted by a "killer" exon flanked by I-SceI sites (Extended Data Figure 3a). DSB induction by I-SceI removes the exon and successful NHEJ event leads to reactivation of the GFP gene. NHEJ reporter construct was integrated into primary, low passage skin fibroblasts. NHEJ repair efficiency was assayed by transfecting cells with I-SceI expression vector and a DsRed plasmid as a transfection control. The repair efficiency was calculated as the ratio of GFP+/DsRed+ cells. Experiments were repeated at least 3 times for each cell line. **b**, % binucleated cells containing micronuclei in human and bowhead whale fibroblasts after 2Gy γ-irradiation. Error bars represent SD (n=2). **c**, Allele plot of Sanger sequencing products resulting from repair of integrated NHEJ reporter cassette after I-SceI cleavage. **d**, Histograms of CRISPR indel size distribution by species. Data for biological replicates are superimposed and partially transparent with lines connecting data points for each sample. Unmodified alleles and alleles with substitutions only are excluded from this analysis. **e**, Distribution of sequenced PTEN allele variants by species after CRISPR DSB induction at a conserved region of the endogenous PTEN gene. Data are averages from multiple primary dermal fibroblast lines isolated from different individual animals for bowhead whale (n=3), human (n=3), cow (n=2), and mouse (n=3). Error bars represent SEM. **f**, Allele plots showing 15 most frequent allele types after CRISPR for one representative cell line per species. Sequences are displayed within a window centered on the cleavage site and extending 20 bp in each direction. Data bars and values indicate proportion of total alleles represented by each row. For the purposes of this display and quantification, all individual alleles with identical sequences in the 40-bp window have been pooled, so rows represent composites of alleles that may differ outside the display window. Error bars represent SD. * $p<0.05$, ** $p<0.01$ (two-tailed t-test). Whale, bowhead whale.

Figure 4. CIRBP is highly expressed in bowhead whale fibroblasts and promotes genomic stability. a, Western blots of DNA repair proteins in primary fibroblasts from different species. **b-c**, bwCIRBP promotes NHEJ and HR in human cells as measured by

flow cytometric GFP-reporter assays (see Methods). In these assays DSBs are induced within inactive NHEJ or HR reporter cassettes by expressing I-SceI endonuclease. Successful NHEJ or HR events lead to reactivation of the fluorescent GFP reporters that are scored by flow cytometry. All experiments in these figures were repeated at least 3 times. **d**, Knockdown of CIRBP in bowhead whale fibroblasts decreases NHEJ efficiency. Cells were transfected with siRNA 3d before I-Sce1 transfection. **e**, Western blot of human fibroblasts overexpressing human CIRBP, whale CIRBP or 9R/A mutated whale CIRBP (left panel); and bowhead whale fibroblasts with knockdown of CIRBP (right panel). **f**, Overexpression of bwCIRBP decreases the percentage of human cells containing micronuclei 3d after 2Gy γ-irradiation (n=2). **g**, Hypothermia promotes NHEJ efficiency in primary human fibroblasts. Cells were pre-incubated at 33°C for 2 days, co-transfected with I-SceI-digested NHEJ reporter and DsRed, and returned to the 33°C incubator. NHEJ efficiency was measured by flow cytometry 3 days following transfection (n=3). Error bars represent mean ± SD. * $p<0.05$, ** $p<0.01$, *** $p<0.001$ (two-tailed t-test).

Figure 5. RPA and CIRBP contribute to increased DNA repair fidelity. a, Distribution of sequenced PTEN allele variants in human primary fibroblasts treated with RPA inhibitor TDRL-505 or transfected with human RPA complex after CRISPR DSB induction at a

conserved region of the endogenous PTEN gene. **b**, Distribution of sequenced PTEN allele variants by species in bowhead whale primary fibroblasts treated with TDRL-505 after CRISPR DSB induction in PTEN gene. **c**, Distribution of sequenced PTEN allele variants by species in human fibroblasts with lentiviral overexpression of luciferase or bwCIRBP after CRISPR DSB induction in endogenous PTEN gene. (**a-c**) Data are averages from experiments performed in triplicate. Error bars represent SEM. * $p < 0.05$, **** $p < 0.0001$. All charts analyzed by two-way ANOVA with Fisher's LSD.

d, Graphical summary. Bowhead whale has evolved very efficient and accurate DSB repair that is mediated by high protein levels of CIRBP, RPA2 and high PARP activity. This enhanced DNA repair may help bowhead whale achieve cancer resistance despite fewer mutational hits required for malignant transformation of whale cells as compared to human cells. The strategy of enhanced repair rather than enhanced elimination of damaged cells through apoptosis or senescence may benefit the extreme longevity of the bowhead whale.

Extended Data Figure 1. Characterization of p53 and Rb knockout cell lines. a, Western blot for p53 protein in fibroblasts clones following CRISPR targeting of *TP53*. Labeled clones were for further validation and experiments. **b**, Western blot for Rb protein in fibroblast clones following CRISPR targeting of *RB1* on p53 knockout background. **c**, Ratio of firefly:renilla luciferase luminescence in fibroblasts transfected with firefly luciferase reporter of p53 transcriptional activity and renilla luciferase control. Cells were treated with etoposide to induce p53 activity. **d**, Ratio of firefly:renilla luciferase luminescence in fibroblasts transfected with firefly luciferase reporter of E2F transcriptional activity and renilla luciferase control. Transfected cells were serum starved for 24h and returned to complete medium for 24h before luminescence measurement. Higher E2F activity results from reduced Rb activity. Error bars represent SD. ****p<0.001 (two-tailed t test), n=3.

Extended Data Figure 2. Analysis of NER and BER in bowhead whale cells. a, NER efficiency was measured by host cell reactivation assay where a plasmid containing luciferase reporter is UV-irradiated *in vitro* to induce DNA damage, transfected into cells, and reactivation of the reporter is measured (n=3 for each cell line). **b**, Kinetics of cyclobutane pyrimidine dimer repair after 30 J/m^2 UVC. Confluent human and whale skin fibroblasts were subjected to UVC, harvested at different time-points, genomic DNA was isolated and analyzed for cyclobutene dimers as described in Methods (n=2 for each cell line). **c**, BER efficiency was measured by host cell reactivation where luciferase reporter plasmid is treated with methylene blue and light to induce oxidative DNA damage, transfected into cells, and luciferase activity measured as described in Methods. **d**, Bowhead whale cells show greater poly-ADP-ribosylation in response to hydrogen peroxide treatment. **e**, Nuclear extracts of bowhead whale fibroblasts exhibit higher basal PARP activity (n=3). **f**, HPRT mutagenesis assay, adjusted by plating efficiency measured for each cell line. **g**, Colony forming efficiency for HPRT mutagenesis assay. Error bars represent mean ± SD. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, ns=not significant (two-tailed t-test).

Extended Data Figure 3. Analysis of DSB repair in bowhead whale cells. a, Possible repair outcomes after induction of DSBs with incompatible ends by I-SceI in NHEJ reporter construct. **b**, NHEJ efficiency in extrachromosomal assay. NHEJ reporter construct was pre-digested with I-SceI, purified and co-transfected with DsRed into human and bowhead skin fibroblasts. Three days after transfection cells were harvested and subjected to flow cytometry to calculate NHEJ efficiency (n=3). **c**, NHEJ fidelity in

extrachromosomal assay. Immediately after FACS/flow cytometry, genomic DNA was isolated, subjected to PCR, cloned and analyzed by Sanger sequencing. At least 100 clones were analyzed for each species. **d**, Representative images of human and bowhead whale binucleated cells containing micronuclei after 2 Gy of γ-irradiation. Scale bar is 20 µm. **e**, Pulse-field gel stained with ethidium bromide, showing chromosomal DNA fragmentation in human and bowhead confluent skin fibroblasts immediately after different doses of γ-irradiation 0.7, 1.5, 3 and 6h after 40 Gy of γ-irradiation. **f**, Kinetics of DSB repair measured by PFGE in confluent human and bowhead fibroblasts after 40 Gy of γ-irradiation. n=2 for each species. Error bars represent mean ± SD. *** p<0.001 (two-tailed t-test).

Extended Data Figure 4. *PTEN* **sequences used for CRISPR NHEJ assays. a**, Alignment of NGS amplicon sequences for each species used to measure NHEJ repair fidelity, showing *PTEN* Exon 1 and guide RNA target site. Amplicon/primer pair was designed based on size appropriate for paired-end sequencing, conservation of internal sequence and primer binding sites, and optimal positioning to detect large deletions. Mismatches to consensus sequence are highlighted in red. Alignments are shown in Benchling with Clustal Omega. **b**, Allele plots from untreated (wild-type) control samples for each species. Plots are outputs from CRISPResso2.

Extended Data Figure 5. Analysis of CRISPR cutting efficiency in different species.
Time course of CRISPR cleavage measured by digital droplet PCR (ddPCR). PTEN copy
number at varying time points after CRISPR RNP transfection was measured with ddPCR
using primers flanking the predicted cleavage site and normalized within each sample to
a single-copy genomic ultra-conserved element as described in Methods. Error bars show
confidence intervals of Poisson distribution calculated in QuantaSoft. Data are averages
from experiments performed in triplicate.

a

	Mouse				Cow			Human				Bowhead whale			
	WTMSF9	WTMSF10	BL6MSF2	Avg.	BT1SF	BT2SF	Avg.	NHDF	HDF637	YAHSF	Avg.	18B9SF	14B10SF	14B11SF	Avg.
1%	-142.0	-140.0	-134.0	**-138.7**	-154.0	-121.0	**-137.5**	-104.0	-126.0	-133.0	**-121.0**	-38.0	-30.0	-68.0	**-45.3**
2%	-110.0	-113.0	-105.0	**-109.3**	-134.0	-82.0	**-108.0**	-63.0	-88.0	-102.0	**-84.3**	-15.0	-14.0	-24.0	**-17.7**
3%	-84.0	-92.0	-80.0	**-85.3**	-112.0	-43.0	**-77.5**	-41.0	-60.0	-71.0	**-57.3**	-10.0	-10.0	-16.0	**-12.0**
4%	-60.0	-80.0	-55.0	**-65.0**	-93.0	-34.0	**-63.5**	-34.0	-37.0	-46.0	**-39.0**	-7.0	-8.0	-11.0	**-8.7**
5%	-44.0	-57.0	-43.0	**-48.0**	-76.0	-31.0	**-53.5**	-29.0	-30.0	-35.0	**-31.3**	-6.0	-6.0	-8.0	**-6.7**
6%	-34.0	-44.0	-34.0	**-37.3**	-54.0	-26.0	**-40.0**	-25.0	-25.0	-30.0	**-26.7**	-4.0	-5.0	-7.0	**-5.3**
7%	-26.0	-38.0	-29.0	**-31.0**	-42.0	-25.0	**-33.5**	-22.0	-23.0	-25.0	**-23.3**	-3.0	-4.0	-6.0	**-4.3**
8%	-21.0	-34.0	-22.0	**-25.7**	-36.0	-22.0	**-29.0**	-21.0	-21.0	-23.0	**-21.7**	-3.0	-3.0	-4.0	**-3.3**
9%	-16.0	-29.0	-18.0	**-21.0**	-31.0	-21.0	**-26.0**	-18.0	-18.0	-22.0	**-19.3**	-3.0	-3.0	-3.0	**-3.0**
10%	-12.0	-25.0	-14.0	**-17.0**	-26.0	-19.0	**-22.5**	-16.0	-16.0	-21.0	**-17.7**	-3.0	-3.0	-3.0	**-3.0**
15%	-4.0	-14.0	-5.0	**-7.7**	-18.0	-12.0	**-15.0**	-10.0	-10.0	-13.0	**-11.0**	-2.0	-3.0	-3.0	**-2.7**
20%	-3.0	-6.0	-3.0	**-4.0**	-12.0	-9.0	**-10.5**	-6.0	-6.0	-10.0	**-7.3**	-1.0	-2.0	-2.0	**-1.7**
25%	-3.0	-4.0	-3.0	**-3.3**	-10.0	-5.0	**-7.5**	-3.0	-3.0	-6.0	**-4.0**	1.0	-1.0	-1.0	**-0.3**
30%	-3.0	-3.0	-3.0	**-3.0**	-7.0	-3.0	**-5.0**	-3.0	-3.0	-4.0	**-3.3**	1.0	-1.0	1.0	**0.3**
35%	-2.0	-3.0	-3.0	**-2.7**	-5.0	-3.0	**-4.0**	-2.0	-2.0	-3.0	**-2.3**	1.0	-1.0	1.0	**0.3**
40%	-2.0	-3.0	-2.0	**-2.3**	-3.0	-3.0	**-3.0**	-1.0	-1.0	-3.0	**-1.7**	1.0	-1.0	1.0	**0.3**
50%	-1.0	-2.0	-1.0	**-1.3**	-3.0	-2.0	**-2.5**	-1.0	-1.0	-3.0	**-1.7**	1.0	1.0	1.0	**1.0**
60%	-1.0	-1.0	-1.0	**-1.0**	-2.0	-1.0	**-1.5**	-1.0	-1.0	-1.0	**-1.0**	1.0	1.0	1.0	**1.0**
70%	-1.0	-1.0	-1.0	**-1.0**	-1.0	-1.0	**-1.0**	-1.0	-1.0	-1.0	**-1.0**	1.0	1.0	1.0	**1.0**
85%	-1.0	-1.0	-1.0	**-1.0**	-1.0	1.0	**0.0**	1.0	-1.0	1.0	**0.3**	1.0	1.0	1.0	**1.0**
90%	-1.0	-1.0	-1.0	**-1.0**	1.0	1.0	**1.0**	1.0	1.0	1.0	**1.0**	1.0	1.0	1.0	**1.0**
95%	1.0	1.0	1.0	**1.0**	1.0	1.0	**1.0**	1.0	1.0	1.0	**1.0**	1.0	1.0	1.0	**1.0**

b

$r = 0.8508$
p = 0.0008989

Extended Data Figure 6. CRISPR indel size percentiles. a, Indel size at the specified percentile rank (left column) by cell line and species, based on frequency of each indel size in sequencing data, ranked from most negative (largest deletions) to most positive (largest insertions), with unmodified or substitution-only alleles excluded. **b**, Pearson correlation between 5th percentile indel size and species lifespan (r=0.8508, 95% CI = 0.5125 to 0.9605, p=0.0009, n=11).

Extended Data Figure 7. Proteomic and transcriptomic quantification of DNA repair proteins. a, Heatmap of LC-MS protein abundance for primary fibroblasts of the indicated species and proteins. Color scale corresponds to $\log_{10}$ ion intensity. **b**, Heatmap of transcript expression from RNA-seq of primary fibroblasts for the indicated species and transcripts. BW, bowhead whale; H, human. Numbers represent different cell lines. Color scale corresponds to normalized transcript abundance as described in Methods. **c**, Western blot of RPA2 in cultured skin fibroblasts, using 2 different monoclonal primary antibodies targeting conserved epitopes and normalized to histone H3. Each lane is a primary fibroblast line from a different adult individual. Fluorescent secondary antibodies were used to increase linear dynamic range for higher quantitative accuracy. **d**, Western blot for CIRBP with 3 different antibodies in 3 fibroblast lines per species. mAb, monoclonal antibody; pAb, polyclonal antibody.

Extended Data Figure 8. Bowhead whale CIRBP stimulates NHEJ in MEFs and effect of temperature on CIRBP expression. a, Effect of bwCIRBP expression on NHEJ in MEFs. CIRBP was expressed in mouse embryonic fibroblasts containing chromosomally integrated NHEJ reporter cassettes. (N=3). MEF, mouse embryonic fibroblasts. **b**, Western blot of CIRBP in human and bowhead whale primary skin fibroblasts at different temperatures. **c**, Relative endogenous CIRBP mRNA expression by qRT-PCR in 3 different human and 3 different bowhead whale cell lines. Error bars represent mean ± SD. ** $p<0.01$. *** $p<0.001$ (two-tailed t-test).

Extended Data Figure 9. Effect of CIRBP mutations on mRNA and protein expression. a, Human and whale CIRBP differ by only 5 amino acids. Alignment of bowhead whale and human CIRBP amino acid sequences (Clustal Omega). **b**, Alignment of bowhead whale and human CIRBP coding sequences (Clustal Omega). **c**, Western blot of bwCIRBP, hCIRBP, and reciprocal amino acid mutants overexpressed in human cells. **d**, Total CIRBP mRNA levels in human cells analyzed by qRT-PCR 24h after transfection of plasmids containing either wild type (hCIRBP, bwCIRBP) or mutants encoding hCIRBP with 5 amino acids mutated to whale sequence (H5BW) or bwCIRBP with 5 amino acids from human sequence (BW5H). β-Actin was used for data normalization. Units shown on the graph are relative to human basal endogenous CIRBP mRNA abundance (n=3). **e**, Calculated codon adaptation index (CAI)[53] for CIRBP coding sequence variants. Error bars are mean ± SD.