DNA repair and anti-cancer mechanisms in the long-lived bowhead whale

Denis Firsanov[1]*, Max Zacher[1]*, Xiao Tian[1], Todd L. Sformo[2], Yang Zhao[1], Greg Tombline[1], J. Yuyang Lu[1], Zhizhong Zheng[1], Luigi Perelli[3], Enrico Gurreri[3], Li Zhang[3], Jing Guo[1], Anatoly Korotkov[1], Valentin Volobaev[1], Seyed Ali Biashad[1], Zhihui Zhang[1], Johanna Heid[4], Alex Maslov[4], Shixiang Sun[4], Zhuoer Wu[1], Jonathan Gigas[1], Eric Hillpot[1], John Martinez[1], Minseon Lee[1], Alyssa Williams[1], Abbey Gilman[1], Nicholas Hamilton[1], Ena Haseljic[1], Avnee Patel[1], Maggie Straight[1], Nalani Miller[1], Julia Ablaeva[1], Lok Ming Tam[1], Chloé Couderc[1], Michael Hoopman[5], Robert Moritz[5], Shingo Fujii[6], Dan J. Hayman[7], Hongrui Liu[8,9], Yuxuan Cai[8], Anthony K. L. Leung[8,10,11,12], Mirre J. P. Simons[7], Zhengdong Zhang[4], C. Bradley Nelson[13], Lisa M. Abegglen[14,15], Joshua D. Schiffman[14,15], Vadim N. Gladyshev[16], , Mauro Modesti[6], Giannicola Genovese[3], Jan Vijg[4, #], Andrei Seluanov[1,17, #], Vera Gorbunova[1,17, #]

[1]Department of Biology, University of Rochester, Rochester, NY, USA.

[2]Department of Wildlife Management, North Slope Borough, Utqiaġvik (Barrow), AK 99723, USA

[3]Department of Genitourinary Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

[4]Department of Genetics, Albert Einstein College of Medicine, Bronx, NY 10461, USA

[5]Institute for Systems Biology, Seattle, WA USA

[6]Cancer Research Center of Marseille, Department of Genome Integrity, CNRS UMR7258, Inserm U1068, Institut Paoli-Calmettes, Aix Marseille Univ, Marseille, France

[7]School of Biosciences, University of Sheffield, UK

[8]Department of Biochemistry and Molecular Biology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD 21205, USA

[9]Cross-Disciplinary Graduate Program in Biomedical Sciences, School of Medicine, Johns Hopkins University, Baltimore, MD 21205, USA

[10]McKusick-Nathans Institute of the Department of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[11]Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[12]Department of Oncology, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[13]Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, USA

[14]Department of Pediatrics & Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, USA

[15]Peel Therapeutics, Inc., Salt Lake City, UT, USA

[16]Division of Genetics, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School,

Boston, MA 02115, USA

[17]Department of Medicine, University of Rochester Medical Center, Rochester, NY, USA

*These authors contributed equally to this work.

[#]Corresponding authors:

Vera Gorbunova
University of Rochester
Email:
vera.gorbunova@rochester.edu

Andrei Seluanov
University of Rochester
Email:
andrei.seluanov@rochester.edu

Jan Vijg
Albert Einstein College of
Medicine
Email: jan.vijg@einsteinmed.edu

Abstract

At over 200 years, the maximum lifespan of the bowhead whale exceeds that of all

other mammals. The bowhead is also the second-largest animal on Earth, reaching

over 80,000 kg[1]. Despite its very large number of cells and long lifespan, the bowhead

is not highly cancer-prone, an incongruity termed Peto's Paradox[2]. This phenomenon

has been explained by the evolution of additional tumor suppressor genes in other

larger animals, supported by research on elephants demonstrating expansion of the p53

gene[3–5]. Here we show that bowhead whale fibroblasts undergo oncogenic

transformation after disruption of fewer tumor suppressors than required for human

fibroblasts. However, analysis of DNA repair revealed that bowhead cells repair double

strand breaks (DSBs) and mismatches with uniquely high efficiency and accuracy

compared to other mammals. The protein CIRBP, implicated in protection from

genotoxic stress, was present in very high abundance in the bowhead whale relative to

other mammals. We show that CIRBP and its downstream protein RPA2, also present

at high levels in bowhead cells, increase the efficiency and fidelity of DNA repair in

human cells. These results indicate that rather than possessing additional tumor

suppressor genes as barriers to oncogenesis, the bowhead whale relies on more

accurate and efficient DNA repair to preserve genome integrity. This strategy which

does not eliminate damaged cells but repairs them may be critical for the long and

cancer-free lifespan of the bowhead whale.

Introduction

The Alaskan Iñupiat Inuit, who carry on a long tradition of subsistence hunting of the

bowhead whale (*Balaena mysticetus*), maintain that these animals "live two human

lifetimes"[6]. A series of bowhead whales captured in the late-twentieth and early-twenty-

first centuries lent new credence to these claims, as embedded in their bodies were

traditional stone harpoon points and bomb lance fragments dating to the Victorian era[7].

Subsequent scientific study and age estimation through quantification of ovarian

corpora, baleen dating, and eye lens aspartic acid racemization analysis supported a

maximum lifespan exceeding 200 years in the bowhead whale[7–12]. Thus, the range of

mammalian lifespans covers roughly 2 orders of magnitude, with the model organism

Mus musculus living for 2-3 years while the bowhead whale lives 100 times as long.

The increased number of cells and cell divisions in larger organisms does not lead to

increased cancer incidence and shorter lifespans[13]. The apparent contradiction between

expected and observed cancer rates in relation to species body mass has been noted

for decades and is known as Peto's Paradox[2,14–16]. Cancer resistance and longer

lifespans in larger species are theorized to result from compensatory evolutionary

adaptations driven by reduced extrinsic mortality[2]. The bowhead whale exceeds 80,000

kg in mass and 200 years in lifespan. Both factors predispose it to accumulating large

numbers of DNA mutations throughout life. To remain alive for so long it must possess

uniquely potent genetic mechanisms to prevent cancer and other age-related diseases.

However, primary research publications on genetic and molecular mechanisms of aging

in the bowhead whale are scarce, consisting primarily of genome and transcriptome

analysis[17–19].

The multi-stage model of carcinogenesis posits that the transition from a normal cell to a

cancer cell involves multiple distinct genetic "hits," or mutations[20]. Larger and longer-

living species might require greater numbers of "hits" for oncogenic transformation,

given their greater cell number and increased lifespan. Indeed, there is experimental

evidence to support this hypothesis. Rangarajan et al. found that while mouse

fibroblasts require perturbation of 2 pathways for tumorigenic transformation (p53 and

Ras), human fibroblasts require 5 hits (p53, pRb, PP2A, telomerase and Ras)[21]. A

human should thus have a dramatically lower per-cell incidence of malignant

transformation than a mouse, and as a result can maintain a larger number of cells for a

longer period of time.

Species that are large-bodied and long-lived may be expected to have even more layers

of protection against oncogenic transformation than humans. In support of this

hypothesis, recent studies have identified copy number expansion and functional

diversification of multiple tumor suppressor genes, such as *TP53* and *LIF*, in elephants

and other taxa[3,5,22–24]. These studies identified multiple copies of *TP53* in the elephant

genome, several of which were confirmed to be transcribed and translated in elephant

fibroblasts and contributed to an enhanced apoptotic response to genotoxic stress[25].

However, additional copies of p53 genes are unlikely to slow down aging[26,27]. One

promising mechanism that could explain both cancer resistance and slower aging in

long-lived mammals is more accurate or efficient DNA repair. Genetic mutations have

been identified as causal factors in carcinogenesis for over a century[28]. Perhaps one of

the most compelling lines of evidence supporting the role of DNA repair in the

pathogenesis of aging and cancer comes from studies of mutants with accelerated

aging phenotypes. Remarkably, most such mutants have defects in DNA repair

enzymes[29–33]. Across species, several studies have also pointed toward improved DNA

repair capacity and reduced mutation accumulation as characteristics associated with

species longevity[34–38]. Here, we identify specific cellular and molecular traits

characterizing bowhead whale cancer resistance and longevity that distinguish it from

shorter-lived mammals including humans. We show that bowhead whale cells are not

more prone to apoptosis and do not require additional genetic hits for malignant

transformation relative to human cells. Instead, the bowhead whale relies on more

accurate and efficient DNA double strand break (DSB) repair promoted by CIRBP and

RPA2, as well as more efficient mismatch repair. This more "conservative" strategy that

does not needlessly eliminate cells but repairs them may be critical for the long and

cancer-free lifespan of the bowhead whale.

Results

Growth characteristics, cellular senescence, and cell death in the bowhead whale

Most human somatic cells lack telomerase activity and as a result undergo replicative

senescence with serial passaging in culture[39]. Replicative and stress-induced

senescence are important mechanisms for preventing cancer. Using TRF and TRAP

assays to measure telomere length and telomerase activity, we found that bowhead

whale skin fibroblasts, like human fibroblasts, lack telomerase activity and experience

telomere shortening followed by replicative senescence with serial passaging in culture

(Figure 1a, b). In both species, nearly all cells stained positive for senescence-

associated β-galactosidase upon terminal growth arrest (Figure 1c, d). As in human

fibroblasts, stable overexpression of human telomerase reverse transcriptase (*hTERT)*

to maintain telomere length prevented replicative senescence in bowhead cells (Figure

1a). Senescence can also be induced by DNA damage. Like human cells, bowhead

whale skin fibroblasts readily entered senescence but did not significantly induce cell

death in response to 10 or 20 Gy of γ-irradiation (Figure 1c-e).

Interestingly, transcriptome analysis of human and bowhead whale senescent

fibroblasts showed reduced induction of senescence-associated secretory phenotype

(SASP) factors in bowhead whale fibroblasts (Figure 1f) relative to human cells.

Paracrine effects of SASP on surrounding cells are thought to contribute to age-related

diseases and carcinogenesis. These transcriptomic differences may indicate that

senescence is able to preserve its anti-cancer function in the bowhead with reduced

harmful paracrine signaling.

To test whether increased p53 activity could contribute to cancer resistance in the

bowhead whale, we transiently transfected primary mouse, cow, human and bowhead

whale skin fibroblasts with a luciferase reporter vector containing a p53-response

element. The bowhead whale cells had the lowest basal p53 activity of the species

tested (Figure 1g). Additionally, we did not observe any differences in the induction of

apoptosis in response to UVC between species (Figure 1h). Together, our results argue

against the idea that increased clearance of damaged cells through apoptosis

contributes to cancer resistance in the bowhead whale.

Requirements for oncogenic transformation of bowhead whale cells

We initially identified a minimal combination of oncogene and tumor suppressor hits

required for *in vitro* malignant transformation of bowhead whale skin fibroblasts using

the soft agar assay, which measures anchorage-independent growth, a hallmark of

cancer. While normal cells undergo growth arrest or programmed cell death (anoikis) in

soft agar, malignant cells continue to grow without substrate adhesion and form visible

colonies[40]. We introduced constructs targeting oncogene and tumor suppressor

pathways into primary skin fibroblasts with PiggyBac (PB) transposon vectors, which

integrate into the genome and drive stable expression. Since bowhead whale primary

fibroblasts, like human fibroblasts, exhibit progressive telomere shortening and lack

telomerase activity (Figure 1b), we used cell lines expressing (*hTERT*) to bypass

replicative senescence.

In agreement with published findings, malignant transformation of human *hTERT+*

fibroblasts required combined expression of H-Ras^{G12V}, SV40 Large T (LT) antigen

(which binds and inactivates p53 and the Rb family of tumor suppressors), and SV40

Small T (ST) antigen (which binds and inactivates PP2A) (Figure 2a)[21]. Rather than

requiring hits to additional pathways, however, bowhead whale *hTERT+* fibroblasts

were transformed by H-Ras^{G12V} and SV40 LT alone, suggesting that bowhead cells may

require fewer genetic mutations to become cancerous compared to human cells (Figure

2a). These findings were supported by mouse xenograft assays, in which the number of

hits needed for tumor growth matched findings from soft agar (Figure 2b).

We next sought to confirm these findings at the genetic level, through CRISPR editing

of individual tumor suppressor genes in bowhead fibroblasts. While the sequenced

bowhead genome has not revealed copy number expansion of canonical tumor

suppressor genes[17,18], CRISPR knockout allows for more precise quantification of the

number of genetic mutations required for oncogenesis. While the most important target

of SV40 LT is thought to be Rb (*RB1* gene), it is also known to inactivate p130 and

p107, two additional members of the Rb-family, providing some level of functional

redundancy. Using CRISPR, we introduced targeted mutations into the bowhead *RB1*

gene, along with *TP53* (the other target of LT), and *PTEN* (an upstream inhibitor of Akt

signaling commonly mutated in human cancers and operating in the same pathway as

PP2A). Following transfection of *hTERT+* bowhead fibroblasts with Cas9-guide RNA

ribonucleoprotein complexes targeting each of the aforementioned genes, we screened

clonally isolated colonies for loss of the targeted protein by Western blot (Figure 2c, d,

Extended Data Figure 1a, b). We additionally screened the colonies with luciferase

reporter assays to confirm loss of protein function and activity (Extended Data Figure

1c, d). For each selected clone, we sequenced the CRISPR-targeted genes to confirm

homozygous knockout at the genetic level and determine the causal mutations

(Supplementary Figures 1, 2). Through this strategy, we generated single and

compound homozygous knockout bowhead whale fibroblasts for *TP53*, *RB1*, and

PTEN. In agreement with our initial findings, genetic inactivation of *TP53* and *RB1* in

bowhead whale fibroblasts expressing *hTERT* and H-Ras^{G12V} was sufficient for

malignant transformation in both soft agar and mouse xenograft assays (Figure 2).

These findings suggest that despite its larger size and longer lifespan, the cells of the

bowhead whale unexpectedly require fewer mutational hits for malignant transformation

than human cells.

Mismatch repair, excision repair and mutagenesis in the bowhead whale

As defects in mismatch repair genes are well-characterized drivers of oncogenesis, we

assessed the efficiency of mismatch repair (MMR) in bowhead whale cells using a

reporter assay that measures cellular correction of a targeted G/T mismatch introduced

to a plasmid *in vitro*[41]. We found that correction of the mismatch was significantly more

efficient in whale cells than in mouse, cow, and human cells (Extended Data Figure 2a).

We next assessed the efficiency of nucleotide excision repair (NER) and base excision

repair (BER) repair in bowhead whale cells. NER is primarily responsible for removing

helix-distorting DNA lesions. To quantify NER activity, we utilized a host cell plasmid

reactivation assay[33] and quantified clearance of cyclobutane pyrimidine dimers (CPDs)

by ELISA to measure repair of UVC-induced DNA damage. NER efficiency by plasmid

reactivation was similar between bowhead and human cells (Extended Data Figure 2b),

but the kinetics of CPD removal tended to be slower in whale cells (Extended Data

Figure 2c). BER is responsible for ameliorating many types of spontaneous DNA base

damage, such as oxidation and deamination. The efficiency of BER, as measured by

the plasmid reactivation assay, trended toward higher BER activity in bowhead whales

compared to human cells, but this difference was not statistically significant (Extended

Data Figure 2d).

We found that PARP activity in bowhead fibroblasts exposed to H_2O_2, and γ-irradiation

was dramatically higher than in human cells (Extended Data Figure 3a, b). Basal PARP

activity was also much higher in untreated bowhead whale nuclear extracts (Extended

Data Figure 3c). PARP proteins are recruited to sites of DNA damage, where they

participate in the DNA damage response and repair. Bowhead whale cells also

displayed higher survival rates after H_2O_2 treatment in comparison to human cells

(Extended Data Figure 3d). When we measured the kinetics of damage repair after

H_2O_2 treatment by alkaline comet assay, repair was slightly accelerated in bowhead

whale relative to human cells, which may relate to its increased PARP activity

(Extended Data Figure 3e).

To determine whether whale cells might accumulate fewer mutations after DNA

damage, we measured mutation frequency following treatment with the potent mutagen

and alkylating agent N-ethyl-N-nitrosourea (ENU) using single-molecule, quantitative

detection of low-abundance somatic mutations by high-throughput sequencing (SMM-

seq)[42]. ENU treatment resulted in a statistically significant increase in somatic mutation

frequency in fibroblasts from all tested species (Extended data Figure 2e). Specifically,

we found that mouse cells showed the greatest increase in ENU-induced single

nucleotide variants, while bowhead whale cells experienced the lowest mutation

induction. The levels of induced mutational load in cow and human cells were

intermediate, in line with their relative lifespans. This suggests a correlation between the

rate of mutation induction and maximum lifespan among the included species

(Extended Data Figure 2e), confirming previous findings by us and others[43][44].

Importantly, the excessive mutational burden observed in ENU-treated cells

predominantly comprised an increased fraction of A to T transversions (Extended Data

Figure 2f), the type of mutation preferentially induced by ENU[42].

We additionally compared mutation induction in response to chemical mutagen

treatment in the bowhead whale and human using the HPRT mutagenesis assay, which

relies on loss of HPRT activity after mutagen treatment[45]. The *HPRT* gene exists as a

single copy on the X chromosome in male mammalian cells, a feature we found to be

true for the bowhead (see Methods). We treated primary fibroblast lines from male

bowhead whale and human with ENU and then plated cells in selective media

containing 6-thioguanine, which kills cells with functional HPRT. Despite a slightly

higher sensitivity to ENU in bowhead whale cells as indicated by colony-forming

efficiency in non-selective media, the rate of HPRT mutant colony formation was

markedly lower in bowhead whale than human fibroblasts, an effect which remained

significant after adjusting for plating efficiency (Extended Data Figure 2g, h). This result

supports the above SMM-seq data that bowhead whale cells may possess more

accurate DNA repair than humans. There was no difference in the rate of apoptosis in

response to ENU treatment in human and bowhead whale fibroblasts, suggesting that

apoptosis is not responsible for the observed differences in ENU sensitivity (Extended

Data Figure 2i). To further validate these findings, we also measured HPRT mutant

colony formation after γ-irradiation. As with ENU, we observed markedly lower HPRT

mutant colony formation in bowhead whale cells (Extended Data Figure 2j, k).

Double-strand break repair and chromosomal stability in the bowhead whale

DNA DSBs are toxic if not repaired and may lead to mutations through inaccurate

repair. DSBs are repaired through two major pathways: non-homologous end joining

(NHEJ) and homologous recombination (HR). To assess relative NHEJ and HR

efficiencies, we integrated fluorescent GFP-based reporter cassettes[46] (Extended Data

Figure 5a) into fibroblasts from mouse, cow, human and bowhead whale. Following DSB

induction with I-SceI, we observed markedly elevated NHEJ efficiency in bowhead

whales relative to other species (Figure 3a, Extended Data Figure 4a). We also found

that HR efficiency is significantly higher in whale cells than in human cells (Figure 3b).

To examine whether this more efficient DSB repair could promote chromosomal

stability, we measured formation of micronuclei induced by γ-irradiation and by I-SceI

cleavage. One potential outcome resulting from an unrepaired DSB in mitotic cells is the

loss of an acentric chromosome fragment, which can be measured as the formation of a

micronucleus. We found that bowhead whale fibroblasts accumulated fewer micronuclei

than human fibroblasts after 2 Gy γ-irradiation (Figure 3c, Extended Data Figure 4b).

We also observed that DSB induction with I-SceI increased the rate of micronucleus

formation, likely reflecting acentric fragment loss, and that this rate was reduced in the

bowhead whale compared to human (Extended Data Figure 4c). Thus, the more

efficient rejoining of DSB ends observed in bowhead whale cells appears to guard

against chromosomal instability.

We also measured resolution of γH2AX and 53BP1 foci, which mark cellular DSBs. We

found that endogenous levels of these foci are significantly lower in whale cells,

suggesting reduced baseline burden of DSBs (Figure 3d). We observed that the kinetics

of DSB repair after γ-irradiation are not faster in the bowhead whale than in human cells

(Extended Data Figure 4d, e)[33]. We further tested the ability of bowhead whale cells to

resolve DSBs after treatment with the DSB-inducing drug bleomycin. We observed

similar induction of foci one hour after bleomycin treatment in human and whale cells;

however, fewer foci remained in cells from the bowhead whales than in those from

humans after 24 hours (Figure 3e, f), indicating a reduced burden of residual unrepaired

damage in the bowhead whale.

Fidelity of DSB repair in the bowhead whale

As mutations resulting from inaccurate DSB repair can promote cancer development,

we next sought to assess the fidelity of DSB repair in the bowhead whale. Sequencing

and analysis of repair junctions from integrated NHEJ reporter (Extended Data Figure

5a, b) and extra-chromosomal NHEJ reporter (Extended Data Figure 5a, c) assays

suggested higher fidelity of NHEJ in bowhead whale cells: compared to human, the

bowhead whale is less prone to producing deletions during the repair of incompatible

DNA termini and far more frequently joins ends without deleting any bases beyond the

small overhang region.

We also measured the fidelity of repair at an endogenous genomic locus. To

systematically compare mutational outcomes of CRISPR break repair in the bowhead

whale to those of humans and shorter-living mammals, we performed CRISPR

transfections in primary fibroblast lines, 2-3 individual animals per species, from

bowhead whale, human, cow, and mouse, and used deep amplicon sequencing of the

targeted locus to generate detailed profiles of repair outcomes. We took advantage of

the fact that exon 1 of the *PTEN* tumor suppressor gene is highly conserved across

mammals, with 100% sequence identity across included species (Supplementary Figure

3). We were therefore able to examine species-specific DSB repair outcomes at an

endogenous genomic locus while minimizing intra-species variation in the break-

proximal sequence context.

Analysis of sequencing data revealed species-specific repair outcomes, which were

consistent across cell lines derived from multiple individual animals of each species

(Figure 3g-i). In human, cow, and mouse, the most common mutational outcomes were

deletions. In contrast, the bowhead was the only species for which a single-base

insertion was the most common mutational event. The frequency of unmodified alleles,

which are known to occur after error-free repair of CRISPR DSBs[47,48], was the highest

in bowhead whale (Figure 3h). Sequencing of untreated control samples confirmed that

the detected insertions and deletions were CRISPR-induced (Supplementary Table 1).

As analysis of CRISPR RNP transfection efficiency by flow cytometry and cleavage

efficiency by digital droplet PCR showed similar CRISPR efficiencies across species

(Extended Data Figure 5d, Supplementary Figure 4a, b), differences observed in the

unmodified allele fraction most likely result from differences in repair fidelity. While small

indels predominated in all species, we observed a marked inverse correlation between

the frequency of large deletions and species lifespan, with the bowhead producing

fewer large deletions than human, cow, and mouse (Figure 3g-i). Intriguingly, this

reduction in large deletions was not accompanied by reduced microhomology usage

(Extended Data Figure 5f, g). When we assigned frequency-based percentile ranks from

most negative to most positive indel size (largest deletions to largest insertions), we

observed a strong correlation between species lifespan and 5th percentile indel size,

corresponding to large deletions (Pearson's r=0.85, p=0.0009) (Extended Data Figure

5e, Supplementary Table 2). The results of these experiments suggest a greater fidelity

of DSB repair in the bowhead whale relative to humans and other mammals.

To determine whether these differences in repair outcomes of targeted DSBs might

predict the types of genomic changes accumulated through spontaneous cellular DNA

damage, we performed whole genome sequencing (WGS)[49] of bowhead whale, human

and mouse fibroblast-derived tumor xenografts and assessed somatic mutations

through comparison to parental non-transformed primary fibroblast cultures sequenced

in tandem (Extended Data Figure 1e). The frequency of spontaneous de novo somatic

single nucleotide variants (SNVs) was significantly lower in bowhead whale tumor

xenografts than in human and mouse (Extended Data Figure 1g). Intriguingly, we

observed no differences in the relative proportions of each type of SNV across species,

suggesting shared underlying mutational drivers during tumor evolution despite

differences in overall mutation rate (Extended Data Figure 1f). We further assessed

WGS data for small indels (Extended Data Figure 1h, i) and large structural variants

(SVs) (Extended Data Figure 1j-l) across species; strikingly, whale tumors were

characterized by a significant reduction in both small and large deletions, as well as

small insertions, large duplications and inversions (Extended Data Figures 1g-l,

Supplementary Table 3). SV size distributions were remarkably different in bowhead

whale tumors in comparison to human and mouse tumors: whale tumors showed a

significant reduction in the proportion of large SVs (>500Kb, $p < 0.0001$, Extended Data

Figure 1m, n). Altogether, these data demonstrate consistent reductions in both the

frequency and size of inserted and deleted bases in bowhead whale cells relative to

those of shorter-lived mammals in response to both nuclease-induced and endogenous

DNA breaks. These differences are likely to reduce the accumulation of deleterious

genomic instability over time.

CIRBP contributes to high DSB repair efficiency and chromosomal stability in the

bowhead whale

To identify mechanisms contributing to the efficiency and accuracy of DSB repair in the

bowhead whale, we compared expression of DNA repair proteins in the bowhead whale

to other mammalian species by Western blot, quantitative mass spectrometry, and

transcriptome sequencing (Figure 4a, Extended Data Figures 6-7). Unexpectedly, we

found that levels of three canonical NHEJ proteins- Ku70, Ku80, and DNA-PKcs- are

substantially higher in human cells than any other species tested (Figure 4a, Extended

Data Figure 6f), while their abundance in the bowhead whale appears to be at more

typical mammalian levels. We speculate that the unusually high levels of Ku/DNA-PKcs

in humans may be a human-specific adaptation to promote DSB repair and genome

stability.

However, we consistently observed a strikingly higher abundance of cold-inducible

RNA-binding protein (CIRBP) in cells and tissues of the bowhead whale than in other

mammalian species (Figure 4a, Extended Data Figure 6a, b, d, f). Levels of PARP1, a

functional partner of CIRBP during DNA repair[50], were also increased relative to human

and showed even greater enrichment at an *in vitro* DSB substrate (Extended Data

Figure 6g). Interestingly, we also found that the CtIP protein, which is required for

efficient HR[51], is more abundant in whale cells compared to humans. It is possible that

CtIP upregulation in the whale contributes to better HR compared to humans.

CIRBP is an RNA- and PAR-binding protein whose expression is induced by a variety of

cellular stressors including cold shock, hypoxia, and UV irradiation[50,52–54]. CIRBP has

been shown to bind the 3' UTR of mRNAs that encode proteins involved in cellular

stress and DNA damage responses and promote their stability and translation[55–58].

There is also evidence for a more direct role of CIRBP in DNA repair: PARP-1-

dependent localization of CIRBP to sites of DNA damage promotes DSB repair and

antagonizes micronucleus formation[50].

To test whether CIRBP contributes to efficient NHEJ and HR in bowhead whale cells,

we overexpressed human (hCIRBP) and bowhead whale (bwCIRBP) in human reporter

cells. Overexpression of bwCIRBP, but not hCIRBP, enhanced NHEJ and HR

efficiencies in human cells (Figure 4b, c, f). Conversely, CIRBP depletion in bowhead

whale cells by siRNA significantly reduced NHEJ and HR efficiency (Figure 4d, e, g).

Consistent with published observations, overexpression of bwCIRBP with nine arginines

in the repeated RGG motif mutated to alanine (9R/A), which impairs CIRBP's ability to

bind to PAR-polymers[50], failed to stimulate HR and reduced stimulation of NHEJ (Figure

4b, c, f).

To test the effects of CIRBP overexpression on chromosomal stability, we quantified the

formation of micronuclei in human cells after γ-irradiation. In cells overexpressing wild-

type hCIRBP, a codon-optimized hCIRBP, and bwCIRBP, we observed a significant

decrease in both basal and γ-irradiation-induced micronuclei in all CIRBP-

overexpressing cells relative to control vector (Figure 4h). Micronucleus formation

appeared to decrease as CIRBP protein expression increased. To probe the

relationship between high NHEJ efficiency, high CIRBP expression, and genome

stability, we also overexpressed human and bowhead whale CIRBP in human NHEJ

reporter cells and observed a reduction in I-SceI-induced micronuclei in cells

overexpressing CIRBP (Extended Data Figure 8f).

We next examined the effect of CIRBP overexpression on formation of gross

chromosomal aberrations in human cells exposed to γ-irradiation. We observed a

markedly decreased frequency of chromosomal aberrations after irradiation in cells

overexpressing hCIRBP and bwCIRBP (Extended Data Figure 8g). This further

suggests a protective effect of high CIRBP expression on genomic stability. Basal

γH2AX/53BP1 foci were reduced by bwCIRBP overexpression, consistent with

improved genome stability (Figure 4i).

The human and bowhead CIRBP proteins differ by only 5 C-terminal amino acids, which

do not overlap with any residues of known functional significance (Extended Data

Figure 9a). Substitution of these 5 codons in hCIRBP with bowhead codons increased

protein expression, while substitution of bwCIRBP with the 5 hCIRBP codons decreased

it (Extended Data Figure 9d). Interestingly, although CIRBP abundance increased

following introduction of the 5 bowhead substitutions, it did not achieve the expression

levels of bwCIRBP, suggesting that synonymous changes to the mRNA coding

sequence contribute to higher translation efficiency of bwCIRBP. Consistent with this

notion, bwCIRBP has a higher codon adaptation index (CAI)[59] than hCIRBP (Extended

Data Figure 9e). We also conducted a phylogenetic analysis of the CIRBP variant

present in the bowhead whale. Serine 126 appears to be ancient, already present in

bovids and bats. The unique cluster of 4 amino acids starting at position 147 is more

recent, only appearing in *Balaenopteridae*, the baleen whales (Extended Data Figure

9b). All baleen whales are very large and long-lived. Interestingly, analysis of primary

fibroblasts from other marine mammals and hippopotamus (who share a common

ancestor with whales), showed that CIRBP is similarly abundant in the humpback whale

but not in sea lions or hippos, while dolphins showed a very mild increase in CIRBP

compared to other mammals (Extended Data Figure 7e).

We did not observe significant upregulation of canonical CIRBP targets in the bowhead

whale following DNA damage. However, CIRBP knockdown showed a trend towards

reducing RPA2 levels (Extended Data Figure 8b). Conversely, bwCIRBP

overexpression in human cells showed a trend towards upregulation of RPA2 levels

(Extended Data Figure 8c).

When we compared the PAR-binding affinity of bwCIRBP to that of hCIRBP through

fluorescence polarization (FP) measurements of labeled PAR, we observed similar K_D

values for both proteins, indicating similar affinity for PAR. K_D values were lower for

longer PAR polymers (PAR_{28} or PAR_{16}) than shorter PAR chains (PAR_8), indicating

higher-affinity binding of CIRBP to the longer polymers (Extended Data Figure 8j). As

CIRBP is present in over 10-fold excess in the whale, there is likely to be greater overall

PAR-binding capacity in whale cells. Intriguingly, despite similar affinity for PAR,

bwCIRBP produced a greater increase in FP of labeled PAR than did hCIRBP,

indicating a stronger effect on PAR hydrodynamics. This raises the possibility that

amino acid differences between the two proteins might lead to differences in binding

conformation or stoichiometry of the CIRBP-PAR complex.

We next investigated the direct involvement of CIRBP in the DSB repair process. We

observed that in bowhead whale cells, the majority of CIRBP is in the nuclear soluble

fraction, but some is always associated with chromatin. This association with chromatin

appeared to be in large part RNA-dependent (Extended Data Figure 8a).

Chen et al [50] demonstrated CIRBP recruitment to laser induced DSBs. To confirm this

result using a different method, we treated whale cells with the DSB-inducing agent

neocarzinostatin (NCS) and tested CIRBP enrichment in the chromatin fraction. Within

minutes after the addition of NCS, CIRBP became transiently enriched in the chromatin

fraction (Extended Data Figure 8d). Damage-induced CIRBP enrichment was sensitive

to RNase A treatment, suggesting that local RNA binding contributes to the association

of CIRBP with chromatin upon DNA damage (Extended Data Figure 8e).

Upon *in vitro* incubation with various nucleic acid substrates, recombinant human

CIRBP produced a concentration-dependent electrophoretic mobility decrease for both

RNA and DNA (Extended Data Figure 8k, l), lending support to prior findings of DNA

binding by CIRBP[60,61]. With sufficient CIRBP, nearly all sheared RNA and DNA

fragments were retained in the well, suggesting an ability of CIRBP to physically tether

or aggregate nucleic acid fragments.

We next investigated whether CIRBP directly facilitates *in vitro* end joining of linearized

plasmid incubated *in vitro* with human XRCC4-Ligase IV complex. We observed that the

ligation of cohesive DNA ends was markedly enhanced by CIRBP in a concentration-

dependent manner (Figure 4j). In contrast, the addition of the accessory protein PAXX[62]

failed to stimulate ligation. Interestingly, without addition of Ku70/Ku80, ligation

stimulation by CIRBP was more pronounced and almost comparable to stimulation by

XLF, which directly interacts with the XRCC4-Ligase IV complex and is considered a

core component of the NHEJ machinery[63]. This result suggests that CIRBP directly

stimulates NHEJ and can promote NHEJ in the absence of Ku70/Ku80. In the whale,

abundant CIRBP may compensate for lower levels of Ku70/Ku80 relative to human.

We additionally tested the effect of bwCIRBP overexpression on repair fidelity in human

cells and observed a reduction in indel rates (Figure 5c, Extended Data Figure 8h). We

also knocked down CIRBP in whale cells harboring an integrated NHEJ reporter and

assessed the mutation spectrum after I-SceI-induced DSBs by sequencing. We

observed an increase in deletions, upon CIRBP knockdown (Extended Data Figure 8i),

suggesting that CIRBP also contributes to repair fidelity.

We additionally observed that human fibroblasts with integrated NHEJ reporters

displayed an increase in NHEJ efficiency when cultured at 33°C rather than 37°C. This

increase in NHEJ efficiency was accompanied by an increase in CIRBP protein

abundance in the cells cultured at 33°C (Extended Data Figure 8m).

Effect of bowhead whale CIRBP on anchorage-independent cell growth

We investigated whether CIRBP overexpression affects the anchorage-independent

growth of transformed cells. bwCIRBP was overexpressed in human fibroblasts

containing SV40 LT, SV40 ST, H-Ras^{G12V}, and hTERT. These cells showed delayed

formation of colonies in soft agar compared to control cells (Extended Data Figure 11a,

b). Importantly, there was no difference in the proliferation rate and cell viability between

CIRBP-expressing and control cells in 2D culture, as assessed by MTT and trypan blue

exclusion assays (Extended Data Figure 11c, d). Furthermore, there was no significant

change in the expression of SV40 LT, H-Ras^{G12V}, and cyclin-dependent kinase

inhibitors (p16INK4a and p21) (Extended Data Figure 11e). Strikingly, we also observed

a lower frequency of chromosomal aberrations in human transformed cells

overexpressing bwCIRBP (Extended Data Figure 11f), suggesting that a possible

explanation for the delay in colony formation is a reduction in chromosomal instability.[65]

Role of RPA2 in bowhead whale DNA repair fidelity

LC-MS proteomics data, Western blots, and transcriptome analysis suggested

increased abundance of the single-stranded DNA binding protein RPA2 in bowhead

whale cells and tissues relative to other mammals (Figure 4a, Extended Data Figure 6c,

e-g, Extended Data Figure 7a-d). RPA is a conserved heterotrimeric ssDNA-binding

protein complex required for eukaryotic DNA replication, which plays a critical role in

DNA repair and DNA damage signaling[64,65]. RPA deficiency increases the frequency of

DSBs in human cells under both basal[66] and stressed[67] conditions. Conversely, RPA

overexpression increases resistance to genotoxic insults[67–70]. RPA promotes NHEJ *in*

vitro[71] and protects ssDNA overhangs at DSBs[72,73].

Treatment of cells transfected with CRISPR to induce DSBs with the small-molecule

RPA DNA-binding inhibitor TDRL-505[74] significantly increased indel rates in both

bowhead and human fibroblasts (Figure 5a, b; Supplementary Table 5, 6). In bowhead,

RPA inhibition also increased the frequency of larger deletions, although this difference

did not reach significance (Figure 5b; Supplementary Table 6). Conversely, co-

transfection of trimeric recombinant human RPA protein during CRISPR significantly

decreased the frequency of mutated alleles in human cells without affecting CRISPR

RNP transfection efficiency (Figure 5a, Supplementary Figure 4c).

In summary, these results suggest that increased abundance of CIRBP and RPA2

positively affects the fidelity of DSB repair, promoting genomic stability in the bowhead

whale.

Discussion

By studying a mammal capable of maintaining its health and avoiding death from cancer

for over two centuries, we are offered a unique glimpse behind the curtain of a global

evolutionary experiment that tested more mechanisms affecting cancer and aging than

humans could hope to approach. Through experiments using primary fibroblasts from

the bowhead whale, we experimentally determined genetic requirements for oncogenic

transformation in the world's longest living mammal and provide evidence that additional

tumor suppressors are not the only solutions to Peto's Paradox. Instead, we find that

the bowhead whale solution lies upstream of tumor suppressor loss and is defined by a

capacity for highly accurate and efficient DNA DSB repair, as well as improved

mismatch repair. We also present evidence that two proteins highly expressed in the

bowhead relative to other mammals, RPA2 and CIRBP, contribute to more efficient and

accurate DSB repair.

CIRBP is highly abundant in the bowhead whale compared to most other mammals. We

speculate that baleen whales evolved this high abundance of CIRBP as an adaptation

to life in cold water and CIRBP was subsequently co-opted to facilitate genome

maintenance. We show that CIRBP directly stimulates NHEJ, and in a purified system

can promote NHEJ in the absence of Ku70/80. A potential mechanism by which CIRBP

could promote end joining *in vivo* is through recruitment to damaged DNA in a manner

enhanced by PAR and RNA, and by promoting the ligation of DNA ends either through

direct interaction with DNA and/or through interaction with DNA repair proteins within a

synaptic complex, tethering broken ends. Another potential mechanism by which CIRBP

could promote ligation of DNA ends might involve liquid-liquid phase separation (LLPS).

Recent evidence suggests that RNA-binding proteins can promote phase separation

around DSB sites. For example, the FUS RNA-binding protein was found to maintain

genomic stability and promote formation of droplet- like compartments in response to

DNA damage[75]. Furthermore, it has been shown that long non-coding RNAs

synthesized at DSBs are necessary to drive molecular crowding of 53BP1 into foci that

exhibit LLPS condensate properties[76]. CIRBP has also been shown to undergo LLPS *in*

vitro[77]. CIBRP may attract DNA into condensates through its affinity for nucleic acids,

increasing rates of intermolecular interaction and ligation kinetics.

Cytosine deamination is an important endogenous source of age-related point

mutations. Our mismatch repair assay measured repair of a G-T mismatch, one

possible intermediate during cytosine deamination. The markedly improved efficiency of

repair of this type of mismatch which we observed in the bowhead whale may hint at

improved repair of this important source of age-related mutation.

While the source of the single-base insertion preference in the bowhead whale remains

an open question, we speculate that this may be related to differential employment of

bypass polymerases during repair synthesis. While mammals have over a dozen

different DNA polymerases, many being specialized error-prone enzymes involved in

lesion bypass, polymerase mu (Pol μ) was specifically found to be lost in the cetacean

lineage[78]. Pol μ is one of the primary error-prone polymerases involved in NHEJ, along

with polymerase lambda (Pol λ) and in some cases terminal deoxynucleotidyl

transferase (TdT)[79]. Pol μ, in comparison with other repair polymerases such as Pol λ,

has a unique propensity to "skip" ahead of 3' unpaired bases, generating deletions[79,80].

The absence of Pol μ may lead to increased reliance on other repair polymerases,

many of which are able to add a single terminal templated or untemplated nucleotide to

a 3' end and could explain the 1-bp insertion bias observed in the bowhead whale.

While we did not identify a signature of positive selection on amino acids in bowhead

whale RPA2, we did identify a difference of possible functional significance in an N-

terminal phosphorylation domain. Previous research has shown that

hyperphosphorylation of multiple S/T residues within this domain drives important

alterations in RPA2's function. Bowhead whale RPA2 has more N-terminal S/T residues

(13) than human RPA2 (9). It has been shown that RPA2 N-terminal

hyperphosphorylation inhibits DNA end resection[81], prevents localization to replication

centers but not to damage sites[82], and is specifically required for the maintenance of

genome stability during replication stress[83].

Prior work has identified positive selection in cancer-related genes such as *CXRC2,*

ADAMTS8, and ANXA1 in cetaceans, as well as cetacean-specific evolutionary

changes to multiple FGF genes[84,85]. Expansion of the eukaryotic initiation factor 2,

polyadenylate binding protein, and 60S ribosomal L10 gene families have also been

identified in the genome of the bowhead whale, potentially implying substantial

alterations to translational regulation[86]. An additional anti-cancer mechanism was

recently reported for the bowhead whale in the form of a *CDKN2C* checkpoint gene

duplication[87]. It has further been suggested that the low body temperature and low

metabolic rate of the bowhead whale may also contribute in part to its extended

lifespan[88]. Indeed, it is likely that numerous individual factors combine to modify cancer

risk and produce the longest mammalian lifespan.

Improvements in DNA repair pathways have also been previously implicated in the

evolution of mammalian longevity. Expression of DNA repair genes was found to be

positively correlated with longevity in a transcriptomic study of 26 mammalian species.[89]

Interestingly, a prior study found higher levels of PAR synthesis and higher PARP1

recruitment to a DNA probe in vitro in the long-lived naked mole rat relative to the

mouse[90], which mirror cellular phenotypes we observed in the bowhead whale relative

to human. While this study also noted higher NER activity in the naked mole rat relative

to mouse, a subsequent study using additional rodent species found that the efficiency

of DSB repair correlates more strongly with longevity across rodent species[91].

One potential drawback of a very accurate DNA repair system could be a reduction in

standing genetic variation and thus a slower rate of evolution of new traits. However,

species living in safe and stable environments have less evolutionary pressure to

rapidly evolve new adaptations. A genome analysis of long-lived rockfishes living in

deep ocean revealed positive selection in DNA repair pathways[92]. Interestingly, a recent

analysis of germline mutations in baleen whales based on analysis of pedigrees

concluded that germline mutation frequencies are similar to those in primates, in

contrast to prior studies finding reduced germline mutation rates in whales[93]. Thus, it

appears that germline and somatic mutation rates are not inherently linked and could

respond to selection independently.

Why would improved DNA repair have evolved in the bowhead whale, as opposed to

the increased tumor suppressor copy number and elevated apoptotic response found in

the elephant and often proposed as a solution to Peto's Paradox? One possible

explanation is that tumor suppressors, apoptosis, and senescence all appear to pose

costs to the organism and force tradeoffs between cancer and cell depletion leading to

age-related degeneration. Simply shifting the balance from apoptosis/senescence to

survival and repair could be detrimental if not also coupled with increased fidelity, as

evidenced by the frequent upregulation of DNA repair pathways in cancer cells[94].

However, evolutionary improvements that couple high efficiency with high fidelity, as

found in the bowhead whale, would promote long-term tissue function and maintenance

at both the cellular and genomic levels. Maintenance of genome stability would reduce

cancer risk and, as suggested by a rapidly growing body of evidence implicating age-

related somatic mosaicism as a ubiquitous feature and functional driver of aging[95–99],

likely protect against numerous other aspects of age-related decline. Thus, the lower

accuracy and efficiency of DNA repair observed in mammals with shorter lifespans may

simply reflect the absence of sufficient selective pressure[88]. Indeed, there is little

selective advantage of DNA repair capacity to last far beyond the age of first

reproduction.

There are currently no approved therapeutics which aim to bolster DNA repair for the

prevention of cancer or age-related decline, and it has been suggested that DNA repair

would be difficult or impossible to improve[100]. However, the bowhead whale provides

evidence that this notion is incorrect. Expression of bwCIRBP in human cells improves

the efficiency and accuracy of DSB repair. Therapeutics based on the evolutionary

strategy of the bowhead whale, including trying to increase activity or abundance of

proteins like CIRBP or RPA2, could one day enable the treatment of genome instability

as a modifiable disease risk factor (for discussion of therapeutic hypothermia see

Supplementary Note). Improving DNA repair could be especially important for patients

with increased genetic predisposition for cancer, or more generally, for aging

populations at increased risk for cancer development.

Acknowledgements

We would like to thank the researchers at the North Slope Borough Department of Wildlife Management, the Alaska Eskimo Whaling Commission, and the Iñupiaq community of Barrow for generously sharing bowhead whale samples, time, resources, skill, and knowledge, and without whom the above work would not have been possible. We would like to give a special thanks to John Craighead "Craig" George, whose pioneering field work established the remarkable longevity of the bowhead whale, and whose kind collaboration and insights helped initiate this project, but who sadly was unable to see its completion. We would like to thank Carlo Maley for suggesting collaboration between Schiffman and Gorbunova groups during a memorable meeting in Arcachon. We would like to thank Aaron Rogers, Mallory Wilmot, and Ryan Kennington for technical support. We would like to thank Tara Harrison, Leigh Duke, the Exotic Species Research Alliance, and Georgia Aquarium for facilitating the collection of the bottlenose dolphin sample. We thank the Marine Mammal Care Center Los Angeles for collecting samples from California sea lions. We would like to thank San Diego Zoo Wildlife Alliance for providing cells from hippopotamus, common dolphin, and humpback whale. Experiments on in vitro ligation were supported by the French National Research Agency. Experiments on PAR-binding were supported by National Institutes of Health R01GM104135 to A.K.L.L. We thank V.K. Thomas in the URMC Center for Advanced Light Microscopy and Nanoscopy (RRID:SCR_023177) for images acquired on the Abberior 3D STED instrument (S10 OD023440). This work was supported by grants from US National Institutes on Aging to VNG, ZZ, JV, AS and VG, and by an award from The Milky Way Research Foundation to VG.

Author contributions

VG, AS, DF, MZ designed research; DF conducted molecular cloning, lentivirus production, immunofluorescence experiments, PFGE, senescence experiments, cell survival experiments, DNA repair assays, micronuclei assays, PARP experiments, CIRBP experiments, and assessed DNA repair fidelity using NHEJ reporter with help from SAB, AP, EH, AW, NM, ML; MZ conducted tumor suppressor CRISPR experiments, RPA experiments, HPRT assays, comet assays, micronuclei analysis, EMSA, and assessed DNA repair fidelity using CRISPR with help from SAB, MS, NH; XT analyzed tumorigenicity; DF and XT conducted cell growth curves and telomere experiments; VV analyzed chromosomal aberrations; AK conducted MMR assay; YZ, CC, Zhihui Zhang and AG assisted with mouse tumor studies; MM and SF conducted NHEJ ligation assay; JCG, TLS, MZ, DF collected bowhead specimens; LP, EG, LZ, and GG performed tumor xenograft sequencing and analysis with help from MZ; JH, AM, SS, and JV performed SMM-Seq of ENU-treated cells with help from JA; ZW assisted with LC-MS and micronuclei; JG assisted with micronuclei and the alkaline comet assay; JM assisted with the HPRT assay; JG performed STED imaging; MH, RM, and GT did LC-MS of liver tissue; LMT did nuclear extractions from liver; MZ, GT performed cell proteomics; JYL and Zhizhong Zheng analyzed RNAseq with help from DF; HL, YC, AKLL performed PAR-binding assays; VG, Zhengdong Zhang, JV contributed data analysis and conceptualization; VG and AS obtained funding and supervised the study; MZ, DF, AS and VG wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

Methods

Reagents

Detailed information on reagents, such as antibodies and sequences of primers, probes, CRISPR guides, and siRNAs, is provided in Supplementary Methods.

Animal experiments

All animal experiments were approved and performed under pre-approved protocols and in accordance with guidelines set by the University of Rochester Committee on Animal Resources (UCAR).

Whale sample collection

Bowhead whale tissues were obtained from adult bowhead whales (*Balaena mysticetus*) captured during 2014 and 2018 Iñupiaq subsistence harvests in Barrow (Utqiaġvik), AK, in collaboration with the North Slope Borough Department of Wildlife Management (NSB DWM) and Alaska Eskimo Whaling Commission after signing a Memorandum of Understanding (September 2014 and March 2021). Tissues were sampled immediately after bowhead whales were brought ashore, after permission to sample was given by the whaling captain, and explants kept in culture medium on ice or at 4°C through initial processing and shipping until arrival at the University of Rochester (UR) for primary fibroblast isolation from skin and lung. Transfer of bowhead whale samples from NSB DWM to UR was under NOAA/NMFS permit 21386.

Primary cell cultures

Primary skin fibroblasts were isolated from skin (dermal) tissues as previously

described.[101] Briefly, skin tissues were shaved and cleaned with 70% ethanol. Tissues

were minced with a scalpel and incubated in DMEM/F-12 medium (ThermoFisher) with

Liberase™ (Sigma) at 37°C on a stirrer for 15-90 min. Tissues were then washed and

plated in DMEM/F-12 medium containing 12% fetal bovine serum (GIBCO) and

Antibiotic-Antimycotic (GIBCO). All subsequent maintenance culture for fibroblasts from

bowhead and other species was in EMEM (ATCC) supplemented with 12% fetal bovine

serum (GIBCO), 100 units/mL penicillin, and 100 mg/mL streptomycin (GIBCO). All

primary cells were cultured at 37°C with 5% CO_2 and 3% O_2 except bowhead whale

cells, which were cultured at 33°C with 5% CO_2 and 3% O_2 based on published field

measurements of bowhead body temperature, which measured a core temperature of

33.8 $^{\circ}$C and a range of lower temperatures in muscle and peripheral tissue.[102,103] Prior

to beginning experiments with bowhead whale fibroblasts, optimal growth and viability

conditions were empirically determined through testing of alternative temperatures,

serum concentrations, and cell culture additives, with optimal culture medium found to

be the same for bowhead and other species. Following isolation, low population-

doubling (PD) primary cultures were preserved in liquid nitrogen, and PD was

continually tracked and recorded during subsequent use for experiments.

Established, primary fibroblasts from mammals were obtained from San Diego Zoo

Wildlife Alliance (hippopotamus, common dolphin, and humpback whale) or generated

at Huntsman Cancer Institute from bottlenose dolphin tissues collected by Georgia

Aquarium through Tara Harrison (Exotic Species Cancer Research Alliance) and

California sea lion tissues collected by the Marine Mammal Care Center Los Angeles

under Institutional Animal Care and Use Committee oversight and National Marine

Fisheries Service permit number 21636.

Soft agar assay

Fibroblast culture medium as described above was prepared at 2X concentration using

2X EMEM (Lonza). To prepare the bottom layer of agar plates, 2X medium was mixed

with a sterile autoclaved solution of 1.2% Noble Agar (Difco) at a 1:1 volumetric ratio,

and 3 mL of 1X medium/0.6% agar was pipetted into each 6-cm cell culture dish and

allowed to solidify at room temperature in a tissue culture hood. To plate cells into the

upper layer of soft agar, cells were harvested and washed, and immediately prior to

plating were resuspended in 2X medium at 20,000 cells/1.5 mL and diluted twofold in

0.8% Noble Agar pre-equilibrated to 37°C. The cells in 0.4% agar/1X medium were

pipetted gently to ensure a homogeneous single cell suspension, and 3 mL (20,000

cells) per 6 cm dish were layered on top of the solidified lower layer. After solidifying in

tissue culture hoods for 20-30 min, additional medium was added to ensure the agar

layers were submerged, and dishes were moved into cell culture incubators. Fresh

medium was added onto the agar every 3 days. 4 weeks after plating, viable colonies

were stained overnight with nitro blue tetrazolium chloride (Thermo Fisher) as

previously described.[40] All cell lines were plated in triplicate.

Images of colonies in soft agar were captured using the ChemiDoc MP Imaging System

(Bio-Rad). Colony quantification was performed using ImageJ software (NIH). Initially,

images were converted to 8-bit format. Subsequently, the threshold function was

adjusted to eliminate any red pixels highlighting non-colony objects. Following threshold

adjustment, images were converted to binary. Colony counting was executed using the

'Analyze Particles' function with the following parameters: Size (pixel^2) = 1 to infinity;

Circularity = 0.5 to 1.

Mouse xenograft assay

NIH-III nude mice (Crl:NIH-Lystbg-J Foxn1nuBtkxid) were purchased from Charles

River Laboratories Inc. (Wilmington, MA, USA). Seven-week-old female mice were used

to establish xenografts. For each injection, 2×10^6 cells were harvested and

resuspended in 100 μl of ice-cold 20% matrigel (BD Bioscience, Franklin Lakes, NJ) in

PBS (Gibco). Mice were anesthetized with isoflurane gas, and 100 μl solution per

injection was injected subcutaneously into the right and left flanks of each mouse with a

22 gauge needle. 3 mice were injected bilaterally, for a total of 6 injections, per cell line

tested. Tumor length and width were measured and recorded every 3-4 days. Mice

were euthanized after reaching a predetermined humane tumor burden endpoint of a

maximum tumor dimension of 20mm in diameter, determined by the longest dimension

of the mouse's largest tumor. For mice that did not reach tumor burden endpoints,

experiments were terminated, and mice euthanized after a maximum of 60 days.

Euthanized mice were photographed, and tumors were excised, photographed, and

weighed to determine the mass of each tumor. Sections of each tumor were frozen at -

80°C and preserved in formalin.

MTT Assay

Cell metabolic activity was determined using Thiazolyl Blue Tetrazolium Bromide (MTT)

(Sigma). Cells were seeded in 24-well plates at a density of 20,000 cells per well one

day before the assay. An MTT solution in PBS was added to the growth medium to

achieve a final concentration of 0.5 mg/mL, and cells were then incubated for 4 hours in

a CO2 incubator. Following incubation, the growth medium was discarded, and 0.5 mL

of DMSO was added to each well to solubilize the purple formazan crystals completely.

The plate was further incubated until the crystals were fully dissolved.

Spectrophotometric absorbance of the samples was measured at a wavelength of 570

nm using a Tecan Spark 20M plate reader.

Telomere lengths

Telomere length was analyzed by Southern blot using the TRF method. Genomic DNA

was extracted from cultured fibroblasts at different population doublings, digested with a

mixture of AluI, HaeIII, RsaI, and HinfI restriction enzymes that do not cut within

telomeric repeat sequences, separated using pulsed-field gel electrophoresis, and

hybridized with a radiolabeled oligonucleotide containing telomeric sequence

$(TTAGGG)_4$. Pulsed field gels were run using a CHEF-DR II apparatus (Bio-Rad) for

22h at a constant 45 V, using ramped pulse times from 1 to 10 s.

Telomeric repeat amplification protocol

Telomeric repeat amplification protocol assay was performed using the TRAPeze kit

(Chemicon, Temecula, CA, USA) according to manufacturer instructions. Briefly, in the

first step of the TRAP assay, radiolabeled substrate oligonucleotide is added to 0.5 µg

of protein extract. If telomerase is present and active, telomeric repeats (GGTTAG) are

added to the 3′ end of the oligonucleotide. In the second step, extended products are

amplified by PCR. Telomerase extends the oligonucleotide by multiples of 6 bp,

generating a ladder of products of increasing length. A human cancer cell line

overexpressing telomerase as well as rodent cells were used as a positive control.

CRISPR ribonucleoprotein transfection

CRISPR RNP complexes were formed in vitro by incubating Alt-R™ S.p.Cas9 Nuclease

V3 (Integrated DNA technologies) with tracRNA annealed to target-specific crRNA

(Integrated DNA Technologies) according to manufacturer instructions. For generation

of tumor suppressor knockouts, 3 RNP complexes with crRNAs targeting different sites

in a single target gene were combined and Alt-R Cas9 Electroporation Enhancer

(Integrated DNA Technologies) was added to transfection mixes prior to electroporation.

For comparative analysis of repair fidelity, 3 μg of pmaxGFP plasmid (Lonza) was

added to transfection mixes to monitor transfection efficiency. Cells were trypsinized

and washed with PBS, and 1×10^6 cells were resuspended in 100 μL of NHDF

Nucleofector Solution (Lonza). The cell suspension was then combined with the

CRISPR transfection solution and gently mixed prior to electroporation on an Amaxa

Nucleofector 2b (Lonza) using program U-23. For RPA inhibition, human and bowhead

fibroblasts were treated with 50 uM TDRL-505 (Sigma) (diluted 1000x into culture

medium from 50mM stock solution in DMSO) for 3 hours prior to CRISPR transfection

and kept in medium with 50 uM TDRL-505 for 18 hours after transfection. For RPA co-

transfection, 1 μg of recombinant human RPA heterotrimer (Enzymax) was added to

CRISPR transfection solution immediately prior to electroporation.

Isolation of clonal cell colonies and screening for tumor suppressor knockout

Following CRISPR transfection, cells were plated at low density in 15 cm dishes to allow

for the formation of isolated colonies. Once clonal colonies of sufficient size had formed,

positions of well-isolated colonies were visually marked on the bottom of the cell culture

dish while under a microscope using a marker. Dishes were aspirated and washed with

PBS. Forceps were used to dip PYREX® 8x8 mm glass cloning cylinders in adhesive

Dow Corning® high-vacuum silicone grease (Millipore Sigma) and one glass cylinder

was secured to the dish over each marked colony. 150 µL of trypsin was added to each

cylinder and returned to the incubator. When cells had rounded up from the plate, the

trypsin in each cylinder was pipetted to detach cells and each colony was added to a

separate well in a 6-cm culture dish containing culture medium. After colonies were

expanded and split into 2 wells per colony, one well was harvested for Western blot

screening for absence of target proteins, while the remaining well was kept for further

experiments.

Luciferase reporter assays for knockout verification

For p53 activity measurement, 1 x 10^6 cells of control (WT) and clonally isolated p53 KO

cell lines were electroporated with 3 µg p53 firefly luciferase reporter plasmid pp53-TA-

Luc (Clontech/Takara) and 0.3 µg renilla luciferase control plasmid pRL-CMV

(Promega) on an Amaxa Nucleofector 2b (Lonza). 24h later, cells were treated with 200

µM etoposide (Sigma) to induce p53 activity. 24h following etoposide treatment, cells

were harvested, and luciferase activity of cell lysates was measured using the Dual-

Luciferase Reporter Assay System (Promega) in a GloMax 20/20 Luminometer

(Promega) according to manufacturer instructions.

For Rb activity measurement, 2 different reporters were tested in parallel: pE2F-TA-Luc

(Clontech/Takara) to measure E2F transcriptional activity (repressed by Rb), and pRb-

TA-Luc (Clontech/Takara) (promoter element directly suppressed by Rb). 1 x 10^6 cells

of control (WT) and clonally isolated Rb KO cell lines were electroporated with 3 µg of

either pE2F-TA-luc or pRb-TA-luc and 0.3 ug renilla luciferase plasmid on an Amaxa

Nucleofector 2b (Lonza). Following transfection, cells were grown in complete medium

for 24h followed by serum-free medium for 24h. Cells were then harvested, and

luciferase activity measured as described above.

Error-corrected sequencing by SMM-seq of ENU-mutated cells

Skin fibroblasts from mouse, cow, human and whale were isolated and cultured as

described before. Confluent cells were treated with 20mg/ml ENU overnight. Then cells

were split 1:4 and grown until confluence for harvest.

Genomic DNA (gDNA) was isolated from frozen cell pellets using the Quick DNA/RNA

Microprep Plus Kit (Zymo D7005). Three hundred ng were used for library preparation

as described before[104]: in brief, DNA was enzymatically fragmented, treated for end

repair before adapter ligation and exonuclease treatment. A size selection step was

performed using a 1.5% cassette on a PippinHT machine prior pulse rolling circle

amplification (RCA) and indexing PCR. Library quality was determined with a Tape

Station (Agilent) and quantified with Qubit (Thermo Fisher). All libraries were sequenced

by Novogene Corp. Inc (CA) on an Illumina platform.

Sequencing analysis and mutation calling was performed as described[104] . After

trimming and alignment to the corresponding reference genome, additional filtering was

performed to distinguish germline mutations from somatic mutations.

Graphs were generated and statistical testing was performed using GraphPad Prism.

Next-generation sequencing of CRISPR repair products

72h after transfection, cells were harvested, and genomic DNA was isolated with the

Wizard Genomic DNA Purification Kit (Promega). DNA concentration was measured on

a Nanodrop spectrophotometer and 100 ng of DNA per sample was PCR-amplified with

KAPA2G Robust HotStart ReadyMix (Roche) based on findings of low PCR bias for

KAPA polymerase[105106]. Primers targeted a conserved region surrounding *PTEN* exon 1

(Extended Data Figure 4a). PCR was performed according to manufacturer instructions,

with an annealing temperature of 66°C for 30 cycles. To purify samples for next-

generation sequencing, PCR products were electrophoresed on a 0.8% agarose gel

and post-stained with SYBR Gold Nucleic Acid Gel Stain (Thermo Fisher). Gels were

visualized on a blue light tray (BioRad) to minimize damage to DNA. A gel slice for each

lane was excised using a scalpel, and each slice was cut to include the region ranging

from just above the prominent *PTEN* PCR band down to and including the "primer

dimer" region to ensure inclusion of any deletion alleles. DNA was extracted from gel

slices using the QiaQuick Gel Extraction Kit (Qiagen), and triplicate PCR reaction

eluates per sample were pooled for sequencing. Sample concentrations were measured

by Nanodrop and adjusted as necessary prior to submission for 2x250 bp paired-end

Illumina MiSeq sequencing with target depth of >40,000 reads/sample (Genewiz).

Analysis of CRISPR NGS data

FASTQ files from each sequenced sample were analyzed with both CRISPResso2,[107] which uses an alignment-based algorithm, and CRISPRPic,[108] which uses a kmer-based algorithm. CRISPResso2 was run using the following parameters: window size = 30, maximum paired-end overlap = 500, bp excluded from left and right ends = 15, minimum alignment score = 50, minimum identity score = 50, plot window size = 20. For CRISPRPic analysis, SeqPrep[109] was used to merge overlapping read pairs and trim adapter sequences. CRISPRPic was run on merged FASTQ sequences for each sample with the following parameters: index size = 8, window size = 30.

HPRT mutation assay

For the HPRT mutation assay, cells used were low-passage primary dermal fibroblasts from multiple species that were known to originate from male animals, to ensure single copy-number of the X-linked *HPRT* gene. Each species was tested with 3 different cell lines from 3 individual animals. The bowhead *HPRT* coding sequence was BLASTed against bowhead genome scaffolds[17] and neighboring gene sequences were analyzed to confirm mammal-typical localization of *HPRT* on the bowhead X-chromosome. Cells were cultured in standard fibroblast growth medium, but with FBS being replaced with dialyzed FBS (Omega Scientific, Inc.) and supplemented with Fibroblast Growth Kit Serum-Free (Lonza) to improve growth and viability in dialyzed FBS. Dialyzed FBS was found in optimization experiments to be necessary for efficient 6-thioguanine selection. Prior to mutagenesis, cells were cultured for 7 days in medium containing HAT Supplement (Gibco) followed by 4 days in HT Supplement (Gibco) to eliminate any pre-existing HPRT mutants. To induce mutation, cells were incubated for 3 hours in serum-

free MEM containing 150 µg/mL N-ethyl-N-nitrosourea (ENU) (Sigma), or exposed to 2

Gy γ-irradiation. Cells were then maintained in ENU-free medium for 9 days to allow

mutations to establish and existing HPRT to degrade. 1×10^6 cells from each cell line

were harvested and plated in dialyzed FBS medium containing 5 µg/mL 6-thioguanine

(Chem-Impex), in parallel with 1×10^6 untreated control cells for each cell line. Cells

were plated at a density of 1×10^5 cells per 15-cm dish to allow for efficient selection

and colony separation, and to prevent potential "metabolic cooperation"[45]. In tandem, for

each cell line 200 cells from untreated and control conditions were plated in triplicate

10-cm dishes in non-selective medium to calculate plating efficiency. After 3 weeks of

growth, surviving colonies were fixed and stained with a crystal violet/glutaraldehyde

solution as previously described[110]. Colonies were counted, and HPRT mutation rate

was calculated as plating-efficiency adjusted number of HPRT-negative colonies

containing >50 cells. Appropriate concentrations of ENU and 6-TG, as well as optimal

plating densities and growth conditions, were determined prior to the experiment

described above through optimization and dose titration experiments.

Digital droplet PCR measurement of CRISPR cleavage rate

A ddPCR assay similar to a previously published method[111] was used for time-course

quantification of CRISPR DSB induction across species. qPCR primers at conserved

sites flanking the guide RNA target site in the *PTEN* gene were designed such that

cleavage would prevent PCR amplification. As an internal copy number reference

control, a second set of previously validated qPCR primers targeting an ultraconserved

element present in all mammals as a single copy per genome (UCE.359) was designed

based on published sequences[112]. To allow for multiplexing and copy number

normalization of *PTEN* within each ddPCR reaction, 5'fluorescent hydrolysis probes

(FAM for *PTEN* and HEX for UCE.359) targeting conserved sequences were designed,

with 3' Iowa Black® and internal ZEN™ quenchers (Integrated DNA Technologies). All

primers and probes were checked for specificity by BLAST against each species'

genome [112]. Fibroblasts were transfected with *PTEN* CRISPR RNP as described in

"Next-generation sequencing of CRISPR repair products" and returned to cell culture

incubators. At the indicated times post-transfection, cells were harvested, flash frozen,

and genomic DNA was isolated with the Wizard Genomic DNA Purification Kit

(Promega). During isolation, newly lysed cells were treated with Proteinase K and

RNase A for 30 minutes each at 37°C to minimize the possibility of residual CRISPR

RNP activity. DNA concentration was measured on a Nanodrop spectrophotometer, and

genomic DNA was pre-digested with BamHI-HF (NEB) and XhoI (NEB), which do not

cut within target amplicons, to maximize PCR efficiency and distribution across droplets.

15 ng of genomic DNA per sample was added to duplicate PCR reactions using the

ddPCR™ Supermix for Probes (No dUTP) master mix (Bio-Rad). Droplets were

prepared and measured according to manufacturer instructions. Briefly, each 20 µL

reaction was mixed with 70 µL Droplet Generation Oil for Probes (Bio-Rad) and droplets

were formed in a QX100 Droplet Generator (Bio-Rad). 40 µL of droplets per reaction

were transferred to 96-well PCR plates and sealed with a PX1 PCR Plate Sealer (Bio-

Rad). The sealed plates were then subjected to PCR using a pre-optimized cycling

protocol. Following PCR, the plates were loaded into a QX100™ Droplet Reader (Bio-

Rad) and each droplet measured on both FAM and HEX channels. *PTEN* copy number

normalized to UCE.359 reference copy number within each well was determined with

QuantaSoft™ software (Bio-Rad). For each species, positive/negative gates in mock-

transfected control samples were adjusted as necessary to compensate for differences

in multiplex PCR efficiency/specificity and "rain" droplets between species and bring

normalized *PTEN* copy number closer to 1. The control gates were then applied across

all samples/time points within the same species and used for *PTEN* copy number

calculation.

Flow cytometric measurement of CRISPR RNP transfection efficiency

965 CRISPR RNP transfections were performed as described above, but with ATTO-550

966 fluorescently labeled tracRNA (Integrated DNA Technologies). At 0h and 24h post-

967 transfection, cells were harvested, pelleted, and analyzed by flow cytometry on a

968 CytoFlex S Flow Cytometer (Beckman Coulter). Gain and ATTO-550 positive gates

969 were set based on mock-transfected control cells included in each experiment.

Senescence-associated β-galactosidase (SA-β-gal) staining

SA-β-gal staining was performed as previously described [113,114]. Cells were washed

twice with PBS and fixed in a solution containing 2% formaldehyde and 0.2%

glutaraldehyde in PBS for 5 min at room temperature. After fixation, cells were

immediately washed twice with PBS and stained in a solution containing 1 mg/mL 5-

bromo-4-chloro-3-indolyl P3-D-galactoside (X-Gal), 40 mM citric acid/sodium phosphate

buffer, pH 6.0, 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 150 mM

NaCl, and 2 mM MgCl$_2$. Plates were incubated at 37°C for 16 h without CO$_2$.

Colorimetric images were taken from different areas of each plate and quantified.

Cell survival assay

Percentage of live cells was quantified using the Annexin-V FLUOS Staining Kit (Roche)

and Annexin V Apoptosis Kit [FITC] (Novus Biologicals) following the manufacturer's

instructions. After staining, cells were analyzed on a CytoFlex S flow cytometer

(Beckman Coulter).

Where indicated cell viability was assessed using a trypan blue exclusion assay. All

cells (both floated and attached to the culture dish) were collected into the same tube,

centrifuged, and resuspended in PBS. The cells were then mixed in a 1:1 ratio with

0.4% trypan blue solution, and approximately 3 minutes later, the percentage of dead

cells was assessed using the Countess 3FL instrument (ThermoFisher) according to the

manufacturer's instructions.

p53 activity

To test p53 activity in cultured primary fibroblasts, 150,000 cells were seeded in 6-well

plates 1 day before transfection with 1 µg pp53-TA-Luc vector (Clontech) and 0.015 µg

pRL-CMV-Renilla (Promega) to normalize for transfection efficiency. Transfections were

performed using PEI MAX Transfection Grade Linear Polyethylenimine Hydrochloride

(MW 40,000) (Polysciences) according to manufacturer instructions. 24h after

transfections cells were lysed using 50µl passive lysis buffer (Promega) per 10^5 cells

and flash frozen/thawed two times in liquid nitrogen and a 37°C water bath. Luciferase

assays were performed using the Dual-Luciferase Reporter Assay System (Promega)

and program DLR-2-INJ on a Glomax 20/20 Luminometer (Promega) with 20µl cell

extract as the input.

Generation of NHEJ and HR reporter cell lines

NHEJ and HR reporter constructs[46] were digested with NheI restriction enzyme and

purified with the QIAEX II gel extraction kit (QIAGEN). The same plasmid DNA

preparation was used for generating all reporter cell lines of the studied species. Cells

PD < 15 were recovered from liquid nitrogen and passaged once before the integration

of the constructs. 0.25 µg of linearized NHEJ and HR constructs were electroporated

into one million cells for each cell line. Two days after transfection, media was

refreshed, and G418 was applied to select stable integrant clones. Triplicates of each

reporter in each cell line were prepared to obtain an adequate number of stable clones.

Clones from triplicate plates were pooled to get at least 50 clones per reporter per cell

line.

DSB repair assays and flow cytometry analysis

DSB repair assays were performed as previously described[115]. Briefly, growing cells

were co-transfected with 3 µg of plasmid encoding I-SceI endonuclease and 0.03 µg of

plasmid encoding DsRed. The same batch of I-SceI and DsRed mixture was used

throughout all species to avoid batch-to-batch variation. To test the effect of CIRBP on

DSB repair, 3 µg of CIRBP plasmids were co-transfected with I-SceI and DsRed

plasmids. Three days after transfection, the numbers of GFP+ and DsRed+ cells were

determined by flow cytometry on a CytoFlex S Flow Cytometer (Beckman Coulter). For

gating strategy see Supplementary Figure 5 in Supplementary Information. For each

sample, a minimum of 50,000 cells was analyzed. DSB repair efficiency was calculated

by dividing the number of GFP+ cells by the number of DsRed+ cells.

For NHEJ knockdown experiments, bowhead whale cells containing the NHEJ reporter

were transfected with 120 pmol of anti-bwCIRBP or control siRNAs (Dharmacon) three

days before I-SceI/DsRed transfections using an Amaxa Nucleofector (U-023 program).

For HR knockdown experiments, bowhead whale cells containing the HR reporter were

transfected twice every three days with a final concentration of 10 nM anti-bwCIRBP or

negative control siRNAs (Silencer Select, Thermo Fisher) using Lipofectamine

RNAiMAX transfection reagent (Thermo Fisher) following the manufacturer's

instructions. Cells were further transfected with I-SceI/DsRed plasmids using a 4D-

Nucleofector (P2 solution, DS150 program). The efficiency of knockdown was

determined by Western blot.

For the extrachromosomal assay and fidelity analysis, NHEJ reporter plasmid was

digested with I-Sce1 for 6h and purified using a QIAEX II Gel Extraction Kit (QIAGEN).

Exponentially growing cells were transfected using an Amaxa nucleofector with the U-

023 program. In a typical reaction, 10^6 cells were transfected with 0.25 µg of

predigested NHEJ reporter substrate along with 0.025 µg of DsRed to serve as a

transfection control. 72h after transfection, cells were harvested and analyzed by flow

cytometry on a BD LSR II instrument. At least 20,000 cells were collected for each

sample. Immediately after FACS, genomic DNA was isolated from cells using the

QIAGEN Blood & Tissue kit. DSB repair sites in the NHEJ construct were amplified by

PCR using Phusion polymerase (NEB), cloned using the TOPO Blunt cloning kit (NEB),

and sent for Sanger sequencing. At least 100 sequenced clones were aligned and

analyzed using the ApE software.

Western blotting

All antibodies were checked for conservation of the target epitope in the protein

sequence of each included species, and only those targeting regions conserved across

these species were used. For a limited number of proteins where the available

antibodies with specific epitope information disclosed did not target conserved regions,

we selected antibodies based on demonstrated reactivity across a broad range of

mammal species and always confirmed these results with multiple antibodies.

Information on antibodies is provided in Supplementary Methods.

Exponentially growing cells were harvested with trypsin and counted, and 10^6 cells were

resuspended in 100 µL of PBS containing protease inhibitors. 100µL of 2x Laemmli

buffer (Bio-Rad) was added, and samples were boiled at 95ºC for 10 minutes. Samples

were separated with 4-20% gradient SDS-PAGE, transferred to a PVDF membrane,

and blocked in 5% milk-TBS-T for 2 hours at room temperature. Membranes were

incubated overnight at +4ºC with primary antibodies in 5% milk-TBS-T. After 3 washes

for 10 minutes with TBS-T, membranes were incubated for 1 hour at room temperature

with secondary antibodies conjugated with HRP or a fluorophore. After 3 washes with

TBS-T signal was developed for HRP secondaries with Clarity Western ECL Substrate

(Bio-Rad). CIRBP and RPA2 expression were each measured with 3 different

antibodies targeting conserved epitopes (Extended Data Figure 7b, c).

For detecting chromatin-bound proteins, cells were lysed in 1 mL of CSK buffer (10 mM

Pipes pH 6.8, 100 mM NaCl, 300 mM sucrose, 3 mM MgCl2, 1 mM EGTA, 0.2% Triton

X-100) or CSK+R buffer (10 mM Pipes pH 6.8, 100 mM NaCl, 300 mM sucrose, 3 mM

MgCl2, 1 mM EGTA, 0.2% Triton X-100, and 0.3 mg/mL RNAse A) at +4°C for 30 min

with gentle rotation. Samples were centrifuged for 10 min at 10,000 × g at 4°C, and the

supernatant was discarded. Pellets were washed twice with 1 mL of CSK/CSK+R buffer,

resuspended in PBS, and an equal volume of 2x Laemmli buffer (Bio-Rad) was added.

Samples were boiled at 95°C for 10 minutes and subjected to Western blotting as

described above.

For analyzing CIRBP expression in mice and bowhead whale tissues, tissues were

pulverized using the cell crusher. For each 5 mg of tissue, 300 μL of 4x Laemmli buffer

(Bio-Rad) was added, samples were extensively vortexed, and boiled at 95°C with 1000

rpm for 10 minutes.

Expression and purification of human and bowhead whale CIRBP protein

N-terminal histidine-tagged (6xHis) CIRBP was cloned into a pET11a expression vector.

The plasmid was transformed into Rosetta gami B (DE3) pLysS competent E. coli for

protein expression. Bacteria were grown at 37oC to an optical density (OD600) of 2.0

and protein expression was induced by adding 0.4mM isopropyl β-d-1-

thiogalactopyranoside (IPTG) for 20 h at 23oC. Bacteria were collected by centrifugation

and pellets were flash frozen on liquid nitrogen and stored at -80oC. In Bacteria were

resuspended in lysis buffer consisting of 50mM Tris pH 7.5, 2.0M NaCl, 50mM

imidazole, 10mg lysozyme, 0.1% Triton X-100, 1mM DTT and protease inhibitors. The

bacterial pellets were sonicated, rotated for 1 h at 4oC, and sonicated again. The

bacterial lysate was clarified by centrifugation at 22,000g for 20 min at 4oC and the

supernatant passed through a 0.45μm filter. The clarified lysate was purified using Ni-

NTA agarose beads (Qiagen) washed with 20 column volumes of water and 20 column

volumes of buffer containing 50mM Tris pH 7.5, 2.0M NaCl, 1mM DTT, and 50mM

imidazole (Wash buffer 1). The lysate was placed onto the washed beads and

transferred to a 50mL conical tube and rotated 3 hr at 4oC. The suspended beads were

pelleted by centrifugation and washed with 40 column volumes wash buffer 1 and 10

column volumes with buffer containing 50mM Tris pH 7.5, 150mM NaCl, 1mM DTT, and

50mM imidazole. CIRBP was eluted by adding 5 column volumes of buffer containing

50mM Tris pH 7.5, 150mM NaCl, 1mM DTT, and 500mM imidazole and rotated the

conical tube for 15 minutes at 4oC. The supernatant was collected by centrifugation and

filtered before adding 5% glycerol. The protein was aliquoted, and flash frozen on liquid

nitrogen and stored at -80oC.

NHEJ ligation in vitro assay

The assay was performed essentially as described [116,117]. Reaction mixtures (10 μl)

contained 20 mM Tris-HCl (pH 7.5), 8 mM MgCl2, 0.1 mM ATP, 2 mM DTT, 0.1 M KCl,

2% Glycerol, 4% PEG 8000, 1 nM linearized pUC19 (with cohesive ends via XbaI; 17.3

ng), 10 nM XRCC4/Ligase IV complex, and 0.5 or 1 μM human CIRBP. When indicated,

reaction mixtures also contained 10 nM Ku70/80 heterodimer, 1 μM XLF dimer, or 1 μM

PAXX dimer. The reaction mixtures were incubated for 1 hr at 30°C, followed by the

addition of 2 μl of Gel Loading Dye, Purple (6X) (NEB), and incubation for 5 min at

65°C. Subsequently, 4 μl of each sample was loaded onto a 0.7% agarose gel and

subjected to gel electrophoresis (50 V, 50 min). The gel was stained with ethidium

bromide, and DNA bands were visualized using a ChemiDoc MP (Bio-Rad).

PARP activity

PARP activity was measured in cell nuclear extracts with the PARP Universal

Colorimetric Assay Kit (Trevigen) according to the manufacturer's instructions. Nuclear

extracts were prepared using EpiQuik Nuclear Extraction Kit (EpigenTek) following

manufacturer protocol. 2.5µg of total nuclear extract was added to measure PARP

activity.

For measurement of PARylation efficiency, cells were treated with 400µM H_2O_2 for 15

and 30 min or subjected to 20 Gy γ-radiation. At the end of incubation, cells were placed

on ice, washed once with PBS, and lysed directly on a plate with 2x Laemmli buffer.

Samples were boiled for 10min at 95ºC and processed by Western Blot.

Preparation of fluorescent ligands, binding assays and fluorescence polarization

measurements

Poly(ADP-ribose) (PAR) oligomers of different lengths (PAR_8, PAR_{16}, and PAR_{28}) were

synthesized, purified, fractionated, and labeled with Alexa Fluor 488 (AF488) dye at the

1'' end, following as described previously[118,119].

To investigate the binding of human and bowhead whale CIRBPs to the fluorescently

labeled PAR and RNA oligomers, titration experiments were conducted. CIRBP proteins

were 4:3 serially diluted and titrated into solutions containing a fixed concentration (3

nM) of the fluorescently labeled PAR. The binding reactions were performed in triplicate

in a buffer comprising 50 mM Tris-HCl pH 7.5, 100 mM KCl, 2 mM MgCl2, 10 mM ß-

mercaptoethanol, and 0.1 mg/mL BSA. The reactions were incubated in dark at room

temperature for 30 minutes in a Corning 384-well Low Flange Black Flat Bottom

Polystyrene NBS Microplate (3575).

After incubation, fluorescence polarization (FP) measurements were performed on a

CLARIOstar Plus Microplate Reader from BMG LABTECH equipped with polarizers and

Longpass Dichroic Mirror 504 nm. The excitation wavelength was set at 482 nm with 16

nm bandwidth, and emission was monitored at 530 nm with 40 nm bandwidth. The FP

values were measured three times, the means of which were analyzed to determine

binding affinities. The binding curves were fitted using a nonlinear regression model to

determine dissociation constants (K_D). The FP increase was quantified to indicate the

hydrodynamic differences upon proteins binding to ligands. Data analysis and curve

fitting were performed using GraphPad Prism.

Pulsed-field gel electrophoresis and analysis of DSBs

After irradiation and repair incubation, confluent human and bowhead whale skin

fibroblasts were harvested, ~400,000 cells were resuspended in PBS, mixed with an

equal volume of 1.4% low gelling temperature agarose and embedded into agarose

plugs. Plugs were kept for 1h at +4ºC and incubated in lysis solution 1 (0.5M EDTA, 2%

sodium sarcosine, 0.5 mg/ml Proteinase K) for 24h at +4ºC. Subsequently plugs were

placed into lysis solution 2 (1.85M NaCl, 0.15M KCl, 5mM $MgCl_2$, 2mM EDTA, 4mM Tris

pH 7.5, 0.5% TritonX100) for 40h at +4ºC. Plugs were then washed two times for 1h in

TE buffer and stored in TE buffer at +4ºC. PFGE was carried out with a CHEF DRII

system (Bio-Rad) in 0.8% agarose gels. The gels were run at 14°C with linearly

increasing pulse times from 50 to 5,000 s for 66 h at a field strength of 1.5 V/cm. Gels

were stained with 0.5 mg/ml ethidium bromide for 4h, washed with TBE buffer and

imaged. Quantitative analysis was performed with Image Lab software (Bio-Rad). The

fraction of DNA entering the gel was quantified. Samples irradiated with various doses

and not incubated for repair served as a calibration to determine the percentage of

remaining DSBs in the repair samples from the fraction of DNA entering the gel.

Immunofluorescence

Exponentially growing cells from humans and bowhead whales were cultured on Lab-

Tek II Chamber Slides (ThermoFisher Scientific), followed by treatment with bleomycin

(BLM) at a final concentration of 5 µg/mL for 1 hour. DNA damage foci were stained

with γH2AX and 53BP1 antibodies and quantified at 1 hour and 24 hours. Considering

the potential non-specificity of γH2AX and 53BP1 antibodies across species, we used

co-localized foci as a more reliable indication of DNA damage.

After BLM treatment, cells were washed twice in PBS, fixed with 2% formaldehyde for

20 minutes at room temperature (RT), washed three times in PBS, and incubated in

chilled 70% ethanol for 5 minutes. After three additional washes in PBS, fixed cells were

permeabilized with 0.2% Triton X-100 for 15 minutes at RT, washed twice for 15

minutes in PBS, and blocked in 8% BSA diluted in PBS supplemented with 0.1%

Tween20 (PBST) for 2 hours at RT. Cells were then incubated with mouse monoclonal

anti-γH2AX (Millipore, 05-636, 1:1000) and rabbit polyclonal anti-53BP1 antibodies

(Abcam, ab172580, 1:1000) diluted in 1% BSA-PBST at +4°C overnight. After

incubation with primary antibodies, cells were washed in PBST three times for 10

minutes and incubated with goat anti-rabbit (Alexa Fluor 488) (Abcam, 1:1500) and goat

anti-mouse antibodies (Alexa Fluor 568) (Thermo Fisher Scientific, 1:1000) for 1 hour at

room temperature. After four washes for 15 minutes in PBST, slides were mounted in

VECTASHIELD Antifade Mounting Medium with DAPI.

For chromatin CIRBP association, cells were pre-incubated with CSK/CSK+R buffer for

3 minutes at RT, washed once in PBS, and subjected to the procedure described above

using rabbit monoclonal anti-CIRBP antibodies (Abcam, 1:1000).

1182   Images were captured using the Nikon Confocal system. Confocal images were

1183   collected with a step size of 0.5 µm covering the depth of the nuclei. Foci were counted

1184   manually under 60x magnification.

1185   Construction of lentiviral overexpression vectors and lentivirus production

The coding sequences of hCIRBP and bwCIRBP were amplified by PCR using Phusion

polymerase (NEB), digested with EcoRI and NotI, and cloned between the EcoRI and

NotI sites of the Lego-iC2 plasmid. The sequence was verified by Sanger sequencing.

Lentiviral particles were produced in Lenti-X 293T cells (Takara). Approximately 10×10^6

cells were transfected with a mixture of pVSV-G (1.7 µg), psPAX2 (3.4 µg), and Lego-

iC2-bwCIRBP (6.8 µg) using PEI MAX (Polysciences). The day after transfection, the

DMEM culture medium (ThermoFisher) was replaced with fresh medium, and lentiviral

particles were harvested from the supernatant for the next 3 days.

1194   Quantification of micronuclei

To analyze binucleated cells containing micronuclei (MN), 10,000-20,000 cells were

plated per chamber slide before irradiation or I-SceI transfection. Immediately after

treatment, cytochalasin B was added to the cell culture media at a final concentration of

0.5-1 µg/ml, and cells were incubated for an additional 72-120 hours. At the end of the

incubation period, cells were washed with PBS, incubated in 75 mM KCl for 10 minutes

at RT, fixed with ice-cold methanol for 1.5-3 minutes, air-dried, and stored. Immediately

before analysis, cells were stained with 100 µg/ml acridine orange for 2 minutes,

washed with PBS, mounted in PBS, and analyzed by fluorescence microscopy.

Alternatively, cells were mounted in VECTASHIELD Antifade Mounting Medium with

DAPI. At least 100 binucleated cells were analyzed per sample.

Chromosomal aberration analysis

Metaphase spreads were prepared according to a standard protocol. Briefly, 0.06 µg/ml

colchicine (Sigma) was added to the growth medium for 4 hours, and cells were

harvested with a 0.25% solution of trypsin/EDTA, treated for 10 minutes with a

hypotonic solution (0.075 M KCl/1% sodium citrate) at 37°C, and fixed with three

changes of pre-cooled (-20°C) methanol/acetic acid mixture (3:1) at -20°C. Cells were

dropped onto pre-cleaned microscope glass slides and air-dried. Metaphase spreads

were stained with Giemsa Stain (Sigma) solution in PBS. For each variant, 100

metaphases were analyzed.

Mismatch repair assay

pGEM5Z(+)-EGFP was a gift from LuZhe Sun (Addgene plasmid #65206;

http://n2t.net/addgene:65206; RRID:Addgene_65206). p189 was a gift from LuZhe Sun

(Addgene plasmid #65207; http://n2t.net/addgene:65207; RRID:Addgene_65207).

Preparation of the heteroduplex EGFP plasmid was following a published method [41]

Briefly, pGEM5Z(+)-EGFP plasmid was nicked with Nb.Bpu10I (Thermo Scientific). After

phenol/chloroform extraction and ethanol precipitation, the nicked plasmid was digested

with Exonuclease III (Thermo Scientific) for 10 minutes at 30°C. p189 was linearized

with restriction enzyme BstXI (NEB) and mixed with the purified circular ssDNA at a

ratio of 1.0:1.5 to generate a heteroduplex EGFP plasmid containing a G/T mismatch

and a nick. The heteroduplex EGFP plasmid with high purity was recovered using a

DNA cleanup kit.

Exponentially growing cells were transfected using a 4D-nucleofector (Lonza) with the

P1 solution using the DS120 program. In a typical reaction, 2×10^5 cells were transfected

with 50 ng of heteroduplex EGFP plasmid along with 50 ng of DsRed2 to serve as a

transfection control. After transfection (48 hours), cells were harvested and analyzed by

flow cytometry on a CytoFlex S flow cytometer (Beckman Coulter).

Host cell reactivation assay

A host cell reactivation assay was employed to assess the repair of UV-induced DNA

damage via nucleotide excision repair, following previously described methods[120].

To evaluate the repair of oxidative DNA damage (base excision repair), a mixture of 20

µg of firefly luciferase (FFL) plasmid and 20-200 µM methylene blue (MB) was

prepared, with water added to reach a final volume of 0.4 ml. The DNA-MB mixture was

dropped onto a petri dish and placed on ice, with another petri dish containing water

positioned on top. Subsequently, the DNA-MB mixture was exposed to visible light for

15 minutes using a 100W lamp positioned at an 11 cm distance. Damaged DNA was

then purified, and the host cell reactivation assay was performed as described for UV-

induced DNA damage[33].

Cyclobutane pyrimidine dimer (CPD) ELISA

Human and bowhead whale skin fibroblasts were cultured until they reached confluency

before UVC radiation. Cells were irradiated in PBS at doses of 0, 5, 10, 20, and 30 J/m2

and immediately harvested to construct an induction curve. To assess DNA repair, cells

were irradiated at 30 J/m2 and then incubated for 6, 24, and 48 hours before harvesting.

Genomic DNA was isolated using the QIAamp Blood Kit (Qiagen). DNA samples were

diluted in PBS to a final concentration of 2 µg/ml, denatured at 100°C for 10 minutes,

and then incubated in an ice bath for 15 minutes. Next, 100 ng of denatured DNA

solution was applied to ELISA plate wells precoated with protamine sulfate (Cosmo Bio)

and dried overnight at 37°C. Plates were washed five times with PBS supplemented

with 0.05% Tween-20 (PBS-T) and then blocked in 2% FBS in PBS-T for 30 minutes at

37°C. After five washes with PBS-T, plates were incubated with mouse monoclonal anti-

CPD antibodies (Clone TDM-2, 1:1000) in PBS for 30 minutes at 37°C. Subsequently,

plates were sequentially incubated with goat-anti-mouse biotin IgG (Invitrogen, 1:1000)

and streptavidin-HRP (Invitrogen, 1:5000) in PBS for 30 minutes at 37°C each, with five

washes with PBS-T before and after each incubation. Plates were then washed with

citrate buffer and incubated with a substrate solution (citrate buffer/o-

phenylenediamine/hydrogen peroxide) for 30 minutes at 37°C. Finally, the reaction was

stopped with 2M H2SO4, and the absorbance was measured at 492 nm using a plate

reader.

CIRBP variant sequence analysis

Identification of rare codons (<10% usage for the corresponding amino acid in human

CDS sequences) was performed on CIRBP coding sequences using the Benchling

Codon Optimization Tool[121]. Codon adaptation index (CAI) was calculated with human

codon frequencies using the E-CAI web server[59].

RNA isolation and RNA-seq analysis

RNA from exponentially growing or senescent mouse, cow, human and bowhead whale

primary skin fibroblasts was isolated using the RNeasy Plus Mini Kit (Qiagen) according

to manufacturer instructions.

Raw reads were demultiplexed using configurebcl2fastq.pl (v1.8.4). Adapter sequences

and low-quality base calls (threshold: Phred quality score <20) in the RNA-seq reads

were first trimmed using Fastp (0.23.4)[122]. For all species, the clean reads were aligned

using Salmon (v1.5.1)[123] to longest coding sequence (CDS) of each gene extracted

from corresponding genome assembly based on human-referenced TOGA annotations.

The values of read count and effective gene lengths for each gene were collected and

integrated into gene-sample table according to their orthologous relationship. Salmon

transcript counts were used to perform differential expression analysis. Only human

genes with orthologs in all species were kept for the downstream species. To filter out

low expressed genes, only gene with all sample read counts sum >10 were retained.

The filtered count matrix was normalized using median of ratios method[124] implemented

in DESeq2 package[125]. The matrix of effective lengths for each gene in each sample

was delivered to the DESeq2 'DESeqDataSet' object to avoid biased comparative

quantifications resulting from species-specific transcript length variation. Differential

expression analysis was performed using DESeq2 and log transformed fold changes

were used for gene set enrichment analysis to assess the differential expression of DNA

repair pathways in bowhead whale, cow, and mouse compared to human. Genes of

DNA repair pathways were compiled from 3 resources: MsigDB database, GO ontology,

and a curated gene list (www.mdanderson.org/documents/Labs/Wood-

Laboratory/human-dna-repair-genes.html)[126,127].

Nanopore sequencing

72h after transfection, cells were harvested and genomic DNA was isolated with the Wizard Genomic DNA Purification Kit (Promega). DNA concentration was measured on a Nanodrop spectrophotometer and 100 ng of DNA per sample was PCR-amplified with Q5 High-Fidelity 2x Master Mix (NEB). PCR products were prepared for multiplexed Nanopore sequencing using the Native Barcoding Kit 96 V14 SQK-NBD114.96 (Oxford Nanopore Technologies). Following end prep, barcoding, and adapter ligation, samples were cleaned up using AMPure XP Beads and loaded onto a R10.4.1 flow cell on a MinION Mk1C (Oxford Nanopore Technologies) for sequencing. Raw data was basecalled in Super-High accuracy mode with barcode and adapter trimming enabled, demultiplexed, and aligned to the NHEJ reporter construct reference sequence FASTA in Dorado. A custom Python script was used to parse CIGAR strings from the resulting BAM files and quantify indels.

In vitro Cas9 pulldown

The conserved *PTEN* target amplicon used for assessing DSB repair fidelity was PCR-amplified from untreated control human fibroblasts as described in "Next-generation sequencing of CRISPR repair products." The *PTEN* target site was cloned into a plasmid using the TOPO TA Cloning Kit (Thermo Fisher Scientific). CRISPR RNP complexes with guide RNAs specific to the target were prepared as follows: 7.2 µL of 200 µM Alt-R™ tracRNA was combined with 7.2 µL of 200 µM Alt-R™ crRNA specific to the *PTEN* genomic target. The tracRNA/crRNA mixture was heated to 95°C and then cooled to room temperature. To the guide duplex were added 25.2 µL of PBS and 20.4 µL of either Alt-R™ S.p. Cas9 Nuclease V3 (Integrated DNA technologies) or Alt-R™

S.p. dCas9 Nuclease V3 (Integrated DNA technologies). RNPs were incubated at room

temperature for 20 minutes. Next, 10 μL of Cas9 and dCas9 RNP were combined with

75 μg of *PTEN* TOPO plasmid and incubated at 37°C for 1 hour in EMEM (ATCC).

Following incubation, 100 ng of soluble nuclear protein extract from either human or

bowhead whale primary fibroblasts, which had been extracted using the NE-PER

Nuclear and Cytoplasmic Extraction Kit (Thermo Fisher Scientific), was added to the

RNP-plasmid complexes to produce cleavage reactions with Cas9 and control reactions

with dCas9 for each species. Nuclear protein extracts were incubated with RNP-plasmid

complexes at 37°C for 45 min. To pull down nuclear proteins associated with

Cas9/dCas9-plasmid complexes (by the His-tag on Cas9/dCas9), 100 μL Ni-NTA

Agarose (Qiagen) was added to each reaction and incubated at 4°C with rotation for 30

min. Ni-NTA Agarose was washed 3x with EMEM + pMaxGFP plasmid to remove

proteins with non-specific plasmid binding, and washed 2x with EMEM + 40 mM

imidazole + pMaxGFP. To aid elution of bound proteins, the *PTEN* plasmid bound by

Cas9/dCas9 was digested by adding 1 μL Benzonase for 5 min. Elution was completed

by adding 1x SDS lysis buffer for S-trap with 250 mM imidazole and eluted proteins

were analyzed by LC-MS as described in "LC-MS proteomic analysis of fibroblasts." To

distinguish between non-quantifiable and non-detected proteins for figure displays,

proteins detected but below the limit of quantification were imputed to an abundance of

10^4, and proteins not detected were imputed to an abundance of 0. For the figure

display, the absolute abundance for each protein was normalized to a value of 1 for the

maximum abundance detected for that protein in any of the Cas9 pulldowns.

Genomic DNA extraction and whole genome sequencing of tumor xenografts

Matching primary cell lines, transformed cell lines, and tumor xenograft samples were

prepared as described above. Samples included 1 mouse cell line, 2 human cell lines,

and 2 bowhead whale cell lines. 1 fresh cell pellet was prepared for each primary and

transformed cell line. For frozen tumor samples, 1 tumor for mouse, 1 tumor for each

human cell line (2 tumors total), 4 tumors for whale cell line 14B11SF, and 5 tumors for

whale cell line 18B2SF were included in the analysis. Genomic DNA extraction and

whole genome sequencing were performed as previously described with minor

modifications[49,128]. Briefly, DNA was extracted from samples using the QIamp DNA Mini

Kit, per manufacturer's recommendations. Isolated genomic DNA was quantified with

Qubit 2.0 DNA HS Assay (ThermoFisher, Massachusetts, USA) and quality assessed

by agarose gel. Library preparation was performed using KAPA Hyper Prep kit (Roche,

Basel, Switzerland) per manufacturer's recommendations. gDNA was sheared to

approximately 400bp using Covaris LE220-plus, adapters were ligated, and DNA

fragments were amplified with minimal PCR cycles. Library quantity and quality were

assessed with Qubit 2.0 DNA HS Assay (ThermoFisher, Massachusetts, USA),

Tapestation High Sensitivity D1000 Assay (Agilent Technologies, California, USA), and

QuantStudio ® 5 System (Applied Biosystems, California, USA). Illumina® 8-nt dual-

indices were used. Equimolar pooling of libraries was performed based on QC values

and sequenced on an Illumina® NovaSeq X Plus (Illumina, California, USA) with a read

length configuration of 150 PE for 60M PE reads (30M in each direction) per sample.

Bioinformatic analysis of tumor xenograft whole genome sequencing

The bioinformatic processing pipeline of raw whole-genome (WGS) high throughput

sequencing data was adapted for human, murine and bowhead whale data [49].

Sequencing FASTA files were applied to FastQC[129] for quality control, adapters were

trimmed by Trimmomatic[130], and the genomic fragments were aligned to the human,

mouse, and whale genome reference (hg19, mm10 and the published bowhead whale

genome assembly[17]) using Burrows-Wheeler Aligner (BWA)[131], then sorted and indexed

by Samtools[132]. Somatic mutations were detected from tumor samples using MuTect

v2[133] to call somatic single-nucleotide variants (SNVs) and small indels (<10bp). Tumor

samples from WGS were compared to their respective matched healthy tissue. All

mutations were also filtered for depth (tumor sample coverage > 30x, normal sample

coverage > 30x) and variant allele frequency (VAF ≥ 0.1). Structural variations were

called by Manta applying default settings and structural variant length > 6000bp were

used for downstream analysis [134].

Alkaline Comet Assay

For the alkaline comet assay, we adapted the Alkaline CometAssay protocol provided

by TREVIGEN based on a published in-gel comet assay method[135] to increase the

number of cell lines and time points assessed and minimize assay variation introduced

during sample harvesting and processing. Slides were pre-coated with a base layer 50µl

of 1% SeaKem LE Agarose (Lonza) to enhance adhesion. We cultured cells to near

100% confluency and then resuspended them in CometAssay LMAgarose (R&D

Systems). We applied 500 cells suspended in 100 µL LMAgarose onto each slide. The

slides were then placed in the dark at 4°C for 10 minutes to allow the agarose to solidify.

After that, slides with live cells were incubated in tissue culture incubators in fibroblast

culture medium containing 700 µM freshly diluted H_2O_2 for 30 minutes, followed by

washing with PBS and incubation for various recovery periods (ranging from 0 minutes

to 12 hours) in culture medium. Slides were collected at each time point, washed with

PBS, and immersed in CometAssay Lysis Solution (R&D Systems). Before

electrophoresis, slides were placed in alkaline unwinding solution prepared according to

TREVIGEN's protocol for 10 minutes. After electrophoresis at 22V for 30 minutes, the

slides were placed in a DNA precipitation buffer following the TREVIGEN protocol for 10

minutes and subsequently washed three times with distilled water. The slides were then

immersed in 70% ethanol for 10 minutes and allowed to air dry in the dark. Before

imaging, each sample was stained with 50 µl of 1x SYBR Gold (Thermo Fisher

Scientific) for 5 minutes before being washed three times with distilled water. Comet

images were acquired through fluorescent microscopy. For scoring, we used profile

analysis in OpenComet[136] within ImageJ. Outliers automatically flagged by OpenComet

were excluded from analysis and remaining incorrectly demarcated comets were further

systematically filtered out according to two criteria: a comet area greater than 5000 or

head area greater than 500.

Tissue processing

Tissues obtained from wild-caught animals were assumed to be of younger/middle age

since predation normally precedes aging in the wild. Postmortem interval was

minimized, and, in all cases, samples were kept on ice and frozen in less than 24h. At

the earliest opportunity after dissection, tissues from representative animals from each

species were flash frozen in liquid nitrogen and stored at -80°C. Tissues were

pulverized to a fine powder within a Biosafety cabinet under liquid nitrogen using a

stainless-steel pulverizer Cell Crusher (Fisher Scientific) chilled in liquid nitrogen and

delivered to storage tubes with a scoop that had also been pre-chilled in liquid nitrogen

and kept on dry ice. Similarly, when sampled for various "omics" processing, pulverized

tissues were removed with a stainless-steel spatula that was pre-chilled in liquid

nitrogen. Samples were never thawed after initial freezing until extractions were

performed.

Cross-species tissue proteomics

We employed a "shotgun" style untargeted data-dependent acquisition (DDA) label-free

quantitative (LFQ) approach. Approximately 5 mg of tissue was mixed with 250µl of

50mM TEAB pH7.6; 5% SDS, mixed by pipetting, and briefly vortexed. Samples were

sonicated in a chilled cup-horn Q800R3 Sonicator System (Qsonica; Newtown, CT) for

a total of 15min at 30% output and duration of 30 x 30 sec pulses (with 30 sec in

between pulses) at 6°C using a chilled circulating water bath. When nuclear proteomes

were analyzed, nuclei were first isolated using a hypotonic lysis approach as in the

preparation of histones[137]. Isolated nuclei were lysed and processed as indicated above

with SDS and sonication and then handled similarly for the rest of the prep. Samples

were heated to 90°C for 2min and allowed to cool to room temperature (RT). Next,

samples were centrifuged at 14,000xg for 10min to pellet insoluble debris and the

supernatants were transferred to clean tubes. Total protein was quantified by the BCA

assay and 100µg was reduced with 5 mM dithiothreitol (DTT) for 30min at 60°C.

Samples were cooled to RT and then alkylated with 10mM iodoacetamide (from a

freshly prepared stock) for 30 min at RT in the dark. Samples were processed using the

standard S-trap mini column method (Protifi; Farmingdale, NY). Samples were digested

with 4µg trypsin overnight at 37°C. Elution fractions were pooled and dried using a

Speedvac (Labconco). Peptides were resuspended in 100µl MS-grade water

(resistance ≥18MΩ) and quantified using the Pierce Quantitative Fluorometric Peptide

Assay (Thermo). Common internal Retention Time standards (CiRT) peptide mix was

added (50 fmol mix/2 µg tryptic peptides) and 2 µg (in 4 µL) of tryptic peptides were

injected/analyzed by mass spectrometry (MS) on a Orbitrap Tribrid Fusion Lumos

instrument (Thermo) equipped with an EASY-Spray HPLC Column (500mm x 75um

2um 100A P/N ES803A, Nano-Trap Pep Map C18 100A; Thermo). Buffer A was 0.1%

formic acid and buffer B was 100% acetonitrile (ACN) with 0.1% FA. Flow rate was

300nl/min and runs were 150 min: 0-120 min, 5% B to 35% B; then from 120-120.5 min,

35-80% B; followed by a 9-minute 80% B wash until 130min. From 130-130.5min B was

decreased to 5% and the column was re-equilibrated for the remaining 20-min at 5% B.

the instrument was run in data dependent analysis (DDA) mode. MS2 fragmentation

was with HCD (30% energy fixed) and dynamic exclusion was operative after a single

time and lasted for 30sec. Additional instrument parameters may be found in the

Thermo RAW files.

Computational proteomics analysis

Raw files were analyzed directly with the MSFragger/Philosopher pipeline[138,139] and

included Peptide and Protein Prophet modules[140] for additional quality control.

Quantitation at the level of MS1 was performed with the "LFQ-MBR; label-free quant

match-between-runs" workflow using default parameters. This allows for alignment of

chromatographic peaks between separate runs. Methionine oxidation and N-terminal

acetylation were set as variable modifications. MaxLFQ with a minimum of two ions was

implemented and normalization of intensity across runs was selected[141].

LC-MS proteomic analysis of fibroblasts

2 15-cm dishes of growing primary fibroblasts from 2 cell lines for each species were

harvested for protein. Cells were washed with PBS and pellets were snap frozen and

stored in liquid nitrogen until processing. Cells were solubilized with 5% SDS; 50 mM

TEAB pH 7 and sonicated at 8°C with 10x 45s pulses using 30% power with 15 s rest

between each pulse with a cup horn Q800R3 Sonicator System (Qsonica; Newtown

CT). Soluble proteins were reduced with 10 mM DTT for 30 min at 55°C, followed by

alkylation with 15 mM iodoacetamide at 25°C in the dark for 30 min. S-trap micro

columns (Protifi; Farmingdale, NY) were employed after this step for overnight tryptic

digestion and peptide isolation according to manufacturer instructions. All solvents were

MS-grade. Resulting tryptic peptides were resuspended in MS-grade water and were

quantified using a Pierce™ Quantitative Fluorometric Peptide Assay (Thermo Fisher cat

#23290). Prior to MS, peptides were mixed with a common internal retention time

standards115 (CiRT) peptide mix (50 fmol CiRT/2μg total tryptic peptides) and

acetonitrile (ACN) and formic acid were added to concentrations of 5% and 0.2%

respectively. The final concentration of the peptide mix was 0.5μg/μl. 2 μg (4μl) of each

were resolved by nano-electrospray ionization on an Orbitrap Fusion Lumos MS

instrument (Thermo) in positive ion mode. A 30 cm home-made column packed with 1.8

μm C-18 beads was employed to resolve the peptides. Solvent A was 0.1% formic acid

and solvent B was 80% acetonitrile with 0.1% formic acid and flow rate was 300 nl/min.

The length of the run was 3 h with a 155 min gradient from 10-38% B. HCD (30%

collision energy) was used for MS2 fragmentation and dynamic exclusion was operative

after a single time and lasted for 30s. Other details of the run parameters may be found

in the embedded run report of the RAW data file uploaded to the ProteomeXchange

database. Peptide assignments and quantitation were done using the label-free quant

match between runs (LFQ-MBR) workflow of MSFragger[138–140]. MaxLFQ with a

minimum of two ions was implemented and normalization was selected. Additional

details are available in MSFragger log files. Searches were performed within the

Philosopher/Fragpipe pipeline that incorporates PeptideProphet and ProteinProphet

filtering steps to increase the likelihood of correct assignments[140]. The databases used

for searches were predicted proteins from the published bowhead genome[17] as well as

our custom proteome derived from our de novo sequenced and Trinity[89,142]-assembled

pool of transcriptomes from whale tissues. Human (UP000005640), mouse

(UP000000589), and bovine (UP000009136) databases were from the latest build

available from Uniprot[143]. For the searches, databases also included a reverse

complement form of all peptides as well as common contaminants to serve as decoys

for false discovery rate (FDR) calculation by the target/decoy approach (decoy present

at 50%). Final FDR was below 1%. To distinguish between non-quantifiable and non-

detected proteins in figure displays, proteins detected but below the limit of

quantification were imputed to an abundance of 10^4, and proteins not detected were

imputed to an abundance of 0.

LC-MS identification of proteins upregulated after DNA damage

3 primary fibroblast lines each for human and bowhead whale were prepared under 3

conditions: untreated control, H_2O_2-treated, and UV treated. For H_2O_2 treatment, culture

medium was replaced with medium containing 400 µM H_2O_2 that had been diluted into

the medium immediately prior to use. For UV treatment, culture medium was aspirated

and replaced with a thin layer of PBS. Cells were exposed to 6 J/m^2 UVC in a UV

Crosslinker (Fisher Scientific) with the culture dish lid removed. 4h after treatments,

cells were harvested by washing with PBS and lysed directly in-dish by addition of 2x

SDS lysis buffer for S-trap. Cells were subsequently processed for LC-MS as described

in "LC-MS proteomic analysis of fibroblasts." Data were acquired on an Orbitrap Astral

Mass Spectrometer (Thermo) equipped with an EASY-Spray HPLC Column (500mm x

75μm 2μm 100A P/N ES803A, Nano-Trap Pep Map C18 100A; Thermo). Samples were

run in DIA mode. Computational- Raw data files were converted to mzML files using

ProteoWizard with peak picking set to 1-n, and demultiplexing selected[144,145]. The mzML

files were searched using DIANN[146] with along with FASTA file databases described

above for tissue extracts using library-free search/library generation with deep learning-

based spectra retention time (RT) and IMs prediction criteria selected. Mass accuracy

was set to 20 and MS1 accuracy set to 5.0, and oxidized methionine (Ox(M)) was also

selected. Nearest human homologs for each species' protein (determined by BLAST as

previously described[89]) were added to the protein group matrix (DIANN output). As

before, the human annotation was used to facilitate cross-species comparisons. Missing

values for protein groups were imputed using deterministic minimum imputation[147,148].

Stimulated emission depletion (STED) microscopy

Immortalized primary fibroblast cultures were plated on 10mm diameter glass coverslips

embedded in 35mm plates (Mattek, P35G-1.5-10-C) at a concentration of 10,000 cells

per plate. 48 hours after plating, plates were washed once with 1x PBS and fixed in

3.7% formaldehyde for 15min on a shaking platform. Plates were washed 3x 10min in

PBS and permeabilized in 0.5% Triton X-100 for 15 minutes on a shaking platform.

Plates were washed again 3x 10min in PBS before blocking for 1h at RT with 5% BSA

in PBS. After blocking, primary antibody was added at a concentration of 1:50 diluted in

100µL blocking buffer and incubated overnight at 4°C. Plates were washed in wash

buffer (0.1% Triton X-100 in PBS) 4x 10min before adding Alexa Fluor 594-labelled

secondary antibody (Abcam, ab150080) at a concentration of 1:50 diluted in 100µL

blocking buffer and incubating for 1h at RT. Plates were washed 4x 10min with wash

buffer and incubated with 100 µL of 1 µg/mL DAPI in PBS for 5min. Plates were washed

once with PBS for 5min before adding mountant (Invitrogen, P10144) and placing a

cover slip. Gated Stimulated Emission Depletion microscopy was performed in the

URMC Center for Advanced Light Microscopy and Nanoscopy (RRID:SCR_023177) on

an Abberior) Göttingen, Germany) 3D STED instrument (S10 OD023440) equipped with

an Olympus UPlanSApo 100x/1.4NA objective. Images were acquired with a pixel size

of 20nm and 775nm depletion laser at 30% power with 15% directed toward 3D

resolution. Single-channel images were pseudocolored red (Alexa-Fluor 594) and blue

(DAPI), merged, and scale bars added using ImageJ software.

Doxycycline-inducible I-SceI NHEJ reporter

The plasmid was assembled from several parts. The backbone was amplified from a

pN1 plasmid without f1 bacteriophage origin of replication and modified by the addition

of short insulator sequences[149] (E2, A2 and A4) purchased from Integrated DNA

Technologies (IDT). The GFP reporter gene with I-SceI endonuclease sites was

amplified from the reporter described above and fused via the P2A self-cleaving peptide

with TetOn transactivator, amplified from Lenti-X™ Tet-One™ Inducible Expression

System Puro (Takara, 631847). A bi-directional promoter sequence featuring hPGK and

TRE3GS was amplified from the same plasmid and cloned upstream of the GFP

reporter, in the orientation for TetOn-P2A-reporter to be driven by the constitutive hPGK

promoter. Downstream of the Tre3GS promoter was closed codon-optimized sequence

for intron-encoded endonuclease I (I-SceI) with SV40 nuclear localization sequence

(NLS) at the N-terminus and nucleoplasmin NLS at the C-terminus fused to the

enhanced blue fluorescent protein (eBFP2) via P2A. The fusion was purchased from

IDT. EBFP2 sequence was derived from eBFP2-N2 plasmid (Addgene #54595).

Cloning was done with In-fusion Snap assembly kit (Takara 638947), NEBuilder HiFi

DNA Assembly (NEB, E5520) and T4 DNA ligase (NEB, M0202). The efficiency of non-

homologous end-joining (NHEJ) double-strand break (DSB) repair was analyzed in

immortalized normal human dermal fibroblasts (NHDF2T). The expression cassette

containing a GFP reporter gene under hPGK promoter and an I-SceI endonuclease

under doxycycline-inducible Tre3GS promoter was inserted into the genome by random

integration method. The positive clones were selected by G418 for 10 days and the

clones were pooled together. The GFP reporter had a short adeno-exon flanked by two

I-SceI recognition sites (in inverted orientation) surrounded by the rat Pem1 intron.

Upon stimulation with doxycycline (100 ng/ml), I-SceI produced two non-ligatable

double strand breaks, resulting in excision of the adeno-exon and reconstitution of the

functional GFP.

EMSA

Recombinant human CIRP protein (Abcam AB106903) was incubated in the indicated

amounts with the indicated nucleic acid substrates in 20 µL EMEM (ATCC) at 37°C for 1

hour. Subsequently, reactions were mixed with 4 µL sucrose loading dye (2M sucrose +

0.2% Orange G) and loaded into agarose gels immersed in 0.5x TAE buffer followed by

electrophoresis at 30V. Following electrophoresis, gels were stained in 1x SYBR Gold

(Thermo Fisher Scientific) and imaged. Extraction of genomic DNA from human primary

fibroblasts was with the Monarch HMW DNA Extraction Kit for Cells & Blood (NEB

#T3050L). To produce the damaged DNA samples and induce PAR formation, cells

were treated with H_2O_2 and UV prior to genomic DNA extraction. For H_2O_2 treatment,

culture medium was replaced with medium containing 400 μM H_2O_2 that had been

diluted into the medium immediately prior to use. For UV treatment, culture medium was

aspirated and replaced with a thin layer of PBS. Cells were exposed to 6 J/m^2 UVC in a

UV Crosslinker (Fisher Scientific) with the culture dish lid removed. During genomic

DNA extraction from damaged chromatin, Proteinase K was added per manufacturer

instructions, but RNase A was omitted and Protector Rnase inhibitor (Sigma-Aldrich)

was added to the extraction buffers and eluate. Nucleic acids used in reactions were

sonicated to uniform size in a QSONICA Sonicator.

Statistical analyses

Statistical comparisons were performed as indicated in the figure legends. Unless

otherwise specified in the text or legend, *n* refers to separate biological replicate cell

lines, isolated from different individuals for a given species. Exceptions include specific

genetically modified cell lines or clones, e.g. tumor suppressor knockout lines and Ku-

deficient MEFs. In such cases, *n* refers to technical replicates and indicates the number

of times the experiment was repeated with the specified cell line. Details for

comparisons done by ANOVA are included in Supplementary Information.

Data Availability

Proteomics data are accessible through ProteomeXchange [URL to be added]. DNA

and RNA sequencing data are accessible through NCBI Sequence Read Archive (SRA)

[URL to be added].

References

1. Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic Acids Res* **46**, D1083–D1090 (2018).

2. Peto, R. Epidemiology, multistage models, and short-term mutagenicity tests. in *The origins of human cancer* (eds. Hiatt, H., Watson, J. & Winsten, J.) vol. 4 1403–1428 (Cold Spring Harbor Conferences on Cell Proliferation, NY, 1977).

3. Abegglen, L. M. *et al.* Potential Mechanisms for Cancer Resistance in Elephants and Comparative Cellular Response to DNA Damage in Humans. *JAMA* **314**, 1850 (2015).

4. Vazquez, J. M., Sulak, M., Chigurupati, S. & Lynch, V. J. A Zombie LIF Gene in Elephants Is Upregulated by TP53 to Induce Apoptosis in Response to DNA Damage. *Cell Rep* **24**, 1765–1776 (2018).

5. Sulak, M. *et al.* TP53 copy number expansion is associated with the evolution of increased body size and an enhanced DNA damage response in elephants. *Elife* **5**, (2016).

6. TEK & Bowhead Life History and Longevity - The North Slope Borough. https://www.north-slope.org/departments/wildlife-management/studies-research-projects/bowhead-whales/traditional-ecological-knowledge-of-bowhead-whales/tek-bowhead-life-history-and-longevity/.

7. George, J. C. & Bockstoce, J. R. Two historical weapon fragments as an aid to estimating the longevity and movements of bowhead whales. *Polar Biol* **31**, 751–754 (2008).

8. Rosa, C. *et al.* Age estimates based on aspartic acid racemization for bowhead whales (Balaena mysticetus) harvested in 1998-2000 and the relationship between racemization rate and body temperature. *Mar Mamm Sci* **29**, 424–445 (2013).

9. Philo, L. M., Shotts, E. B. & George, J. C. Morbidity and mortality- The bowhead whale. *Soc. Mar. Mammal., Spec. Publ* 275–312 (1993).

10. George, J. C. *et al.* Age and growth estimates of bowhead whales (Balaena mysticetus) via aspartic acid racemization. *Can J Zool* **77**, 571–580 (1999).

11. Wetzel, D. *et al.* Age estimation for bowhead whales, Balaena mysticetus, using aspartic acid racemization with enhanced hydrolysis and derivatization procedures. *Paper SC/65b/BRG05 presented to the Scientific Committee of the International Whaling Commission* (2014).

12. George, J. C. *et al.* A new way to estimate the age of bowhead whales (Balaena mysticetus) using ovarian corpora counts. *https://doi.org/10.1139/z11-057* **89**, 840–852 (2011).

13. Vincze, O. *et al.* Cancer risk across mammals. *Nature* **601**, 263–267 (2022).

14. Boddy, A. M. *et al.* Lifetime cancer prevalence and life history traits in mammals. *Evol Med Public Health* **2020**, 187–195 (2020).

15. Vincze, O. *et al.* Cancer risk across mammals. *Nature* **601**, 263–267 (2022).

1627   16.   Tollis, M., Boddy, A. M. & Maley, C. C. Peto's Paradox: how has evolution solved the problem of
1628   cancer prevention? *BMC Biol* **15**, (2017).

1629   17.   Keane, M. *et al.* Insights into the evolution of longevity from the bowhead whale genome. *Cell*
1630   *Rep* **10**, 112–122 (2015).

1631   18.   Tollis, M. *et al.* Return to the Sea, Get Huge, Beat Cancer: An Analysis of Cetacean Genomes
1632   Including an Assembly for the Humpback Whale (Megaptera novaeangliae). *Mol Biol Evol* **36**,
1633   1746–1763 (2019).

1634   19.   Seim, I. *et al.* The transcriptome of the bowhead whale Balaena mysticetus reveals adaptations of
1635   the longest-lived mammal. *Aging* **6**, 879–899 (2014).

1636   20.   Armitage, P. & Doll, R. The age distribution of cancer and a multi-stage theory of carcinogenesis.
1637   *Br J Cancer* **8**, 1–12 (1954).

1638   21.   Rangarajan, A., Hong, S. J., Gifford, A. & Weinberg, R. A. Species- and cell type-specific
1639   requirements for cellular transformation. *Cancer Cell* **6**, 171–183 (2004).

1640   22.   Vazquez, J. M., Sulak, M., Chigurupati, S. & Lynch, V. J. A Zombie LIF Gene in Elephants Is
1641   Upregulated by TP53 to Induce Apoptosis in Response to DNA Damage. *Cell Rep* **24**, 1765–1776
1642   (2018).

1643   23.   Vazquez, J. M. & Lynch, V. J. Pervasive duplication of tumor suppressors in Afrotherians during
1644   the evolution of large bodies and reduced cancer risk. *Elife* **10**, 1–45 (2021).

1645   24.   Vazquez, J. M. *et al.* Parallel evolution of reduced cancer risk and tumor suppressor duplications
1646   in Xenarthra. *Elife* **11**, (2022).

1647   25.   Preston, A. J. *et al.* Elephant TP53-RETROGENE 9 induces transcription-independent apoptosis at
1648   the mitochondria. *Cell Death Discov* **9**, 66 (2023).

1649   26.   García-Cao, I. *et al.* 'Super p53' mice exhibit enhanced DNA damage response, are tumor
1650   resistant and age normally. *EMBO J* **21**, 6225–6235 (2002).

1651   27.   Tyner, S. D. *et al.* p53 mutant mice that display early ageing-associated phenotypes. *Nature* **415**,
1652   45–53 (2002).

1653   28.   Boveri, T. Concerning the origin of malignant tumours by Theodor Boveri. Translated and
1654   annotated by Henry Harris. *J Cell Sci* **121**, 1–84 (2008).

1655   29.   Hasty, P., Campisi, J., Hoeijmakers, J., Van Steeg, H. & Vijg, J. *Aging and Genome Maintenance:*
1656   *Lessons from the Mouse?* http://science.sciencemag.org/.

1657   30.   Kashiyama, K. *et al.* Malfunction of nuclease ERCC1-XPF results in diverse clinical manifestations
1658   and causes Cockayne syndrome, xeroderma pigmentosum, and Fanconi anemia. *Am J Hum Genet*
1659   **92**, 807–819 (2013).

1660   31.   Navarro, C. L., Cau, P. & Lévy, N. Molecular bases of progeroid syndromes. *Human Molecular*
1661   *Genetics* vol. 15 R151–R161 Preprint at https://doi.org/10.1093/hmg/ddl214 (2006).

32.   Varon, R. *et al.* Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen
breakage syndrome. *Cell* **93**, 467–476 (1998).

33.   Liu, B. *et al.* Genomic instability in laminopathy-based premature aging. *Nat Med* **11**, 780–785
(2005).

34.   Lorenzini, A. *et al.* Significant Correlation of Species Longevity with DNA Double Strand Break-
Recognition but not with Telomere Length. *Mech Ageing Dev* **130**, 784 (2009).

35.   Zhang, L. *et al.* Maintenance of genome sequence integrity in long- And short-lived rodent
species. *Sci Adv* **7**, (2021).

36.   Cagan, A. *et al.* Somatic mutation rates scale with lifespan across mammals. *Nature 2022* 1–8
(2022) doi:10.1038/s41586-022-04618-z.

37.   Tian, X. *et al.* SIRT6 Is Responsible for More Efficient DNA Double-Strand Break Repair in Long-
Lived Species. *Cell* **177**, 622-638.e22 (2019).

38.   Fortunato, A., Fleming, A., Aktipis, A. & Maley, C. C. Upregulation of DNA repair genes and cell
extrusion underpin the remarkable radiation resistance of Trichoplax adhaerens. *PLoS Biol* **19**,
(2021).

39.   Tian, X. *et al.* Evolution of telomere maintenance and tumour suppressor mechanisms across
mammals. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**, (2018).

40.   Borowicz, S. *et al.* The soft agar colony formation assay. *J Vis Exp* e51998 (2014)
doi:10.3791/51998.

41.   Zhou, B. *et al.* Preparation of heteroduplex enhanced green fluorescent protein plasmid for in
vivo mismatch repair activity assay. *Anal Biochem* **388**, 167–169 (2009).

42.   Kucab, J. E. *et al.* A Compendium of Mutational Signatures of Environmental Agents. *Cell* **177**, 821
(2019).

43.   Cagan, A. *et al.* Somatic mutation rates scale with lifespan across mammals. *Nature 2022* 1–8
(2022) doi:10.1038/s41586-022-04618-z.

44.   Zhang, L. *et al.* Maintenance of genome sequence integrity in long- And short-lived rodent
species. *Sci Adv* **7**, (2021).

45.   Johnson, G. E. Mammalian Cell HPRT Gene Mutation Assay: Test Methods. in 55–67 (2012).
doi:10.1007/978-1-61779-421-6_4.

46.   Seluanov, A., Mao, Z. & Gorbunova, V. Analysis of DNA Double-strand Break (DSB) Repair in
Mammalian Cells. *J Vis Exp* 43 (2010) doi:10.3791/2002.

47.   Guo, T. *et al.* Harnessing accurate non-homologous end joining for efficient precise deletion in
CRISPR/Cas9-mediated genome editing. *Genome Biol* **19**, (2018).

48.   Richardson, C. D., Ray, G. J., Bray, N. L. & Corn, J. E. Non-homologous DNA increases gene
disruption efficiency by altering DNA repair outcomes. *Nature Communications 2016 7:1* **7**, 1–7
(2016).

49.   Perelli, L. *et al.* Interferon signaling promotes tolerance to chromosomal instability during
metastatic evolution in renal cancer. *Nat Cancer* **4**, 984–1000 (2023).

50.   Chen, J. K., Lin, W. L., Chen, Z. & Liu, H. wen. PARP-1–dependent recruitment of cold-inducible
RNA-binding protein promotes double-strand break repair and genome stability. *Proc Natl Acad*
Sci U S A **115**, E1759–E1768 (2018).

51.   Sartori, A. A. *et al.* Human CtIP promotes DNA end resection. *Nature* **450**, 509–514 (2007).

52.   Nishiyama, H. *et al.* A glycine-rich RNA-binding protein mediating cold-inducible suppression of
mammalian cell growth. *J Cell Biol* **137**, 899–908 (1997).

53.   Sheikh, M. S. *et al.* Identification of several human homologs of hamster DNA damage-inducible
transcripts. Cloning and characterization of a novel UV-inducible cDNA that codes for a putative
RNA-binding protein. *J Biol Chem* **272**, 26720–26726 (1997).

54.   Wellmann, S. *et al.* Oxygen-regulated expression of the RNA-binding proteins RBM3 and CIRP by
a HIF-1-independent mechanism. *J Cell Sci* **117**, 1785–1794 (2004).

55.   Yang, C. & Carrier, F. The UV-inducible RNA-binding protein A18 (A18 hnRNP) plays a protective
role in the genotoxic stress response. *J Biol Chem* **276**, 47277–47284 (2001).

56.   Yang, R., Weber, D. J. & Carrier, F. Post-transcriptional regulation of thioredoxin by the stress
inducible heterogenous ribonucleoprotein A18. *Nucleic Acids Res* **34**, 1224–1236 (2006).

57.   Yang, R. *et al.* Functional significance for a heterogenous ribonucleoprotein A18 signature RNA
motif in the 3'-untranslated region of ataxia telangiectasia mutated and Rad3-related (ATR)
transcript. *J Biol Chem* **285**, 8887–8893 (2010).

58.   Roilo, M., Kullmann, M. K. & Hengst, L. Cold-inducible RNA-binding protein (CIRP) induces
translation of the cell-cycle inhibitor p27Kip1. *Nucleic Acids Res* **46**, 3198–3210 (2018).

59.   Puigbò, P., Bravo, I. G. & Garcia-Vallvé, S. E-CAI: a novel server to estimate an expected value of
Codon Adaptation Index (eCAI). *BMC Bioinformatics* **9**, (2008).

60.   Spruijt, C. G. *et al.* Dynamic readers for 5-(hydroxy)methylcytosine and its oxidized derivatives.
Cell **152**, 1146–1159 (2013).

61.   Huang, Z. L. *et al.* Identification of G-Quadruplex-Binding Protein from the Exploration of RGG
Motif/G-Quadruplex Interactions. *J Am Chem Soc* **140**, 17945–17955 (2018).

62.   Seif-El-Dahan, M. *et al.* PAXX binding to the NHEJ machinery explains functional redundancy with
XLF. *Sci Adv* **9**, (2023).

63.   Ahnesorg, P., Smith, P. & Jackson, S. P. XLF Interacts with the XRCC4-DNA Ligase IV Complex to
Promote DNA Nonhomologous End-Joining. *Cell* **124**, 301–313 (2006).

64.   Zou, Y., Liu, Y., Wu, X. & Shell, S. M. Functions of Human Replication Protein A (RPA): From DNA
1731   Replication to DNA Damage and Stress Responses. *J Cell Physiol* **208**, 267 (2006).

1732   65.   Wold, M. S. Replication protein A: a heterotrimeric, single-stranded DNA-binding protein
1733   required for eukaryotic DNA metabolism. *Annu Rev Biochem* **66**, 61–92 (1997).

1734   66.   Paulsen, R. D. *et al.* A genome-wide siRNA screen reveals diverse cellular processes and pathways
1735   that mediate genome stability. *Mol Cell* **35**, 228–239 (2009).

1736   67.   Toledo, L. I. *et al.* ATR prohibits replication catastrophe by preventing global exhaustion of RPA.
1737   *Cell* **155**, 1088 (2013).

1738   68.   Velegzhaninov, I. O. *et al.* Radioresistance, DNA Damage and DNA Repair in Cells With Moderate
1739   Overexpression of RPA1. *Front Genet* **11**, 855 (2020).

1740   69.   Guven, M., Brem, R., Macpherson, P., Peacock, M. & Karran, P. Oxidative damage to RPA limits
1741   the nucleotide excision repair capacity of human cells. *J Invest Dermatol* **135**, 2834 (2015).

1742   70.   Graziano, S. *et al.* Lamin A/C recruits ssDNA protective proteins RPA and RAD51 to stalled
1743   replication forks to maintain fork stability. *Journal of Biological Chemistry* **297**, 101301 (2021).

1744   71.   Perrault, R., Cheong, N., Wang, H., Wang, H. & Iliakis, G. RPA facilitates rejoining of DNA double-
1745   strand breaks in an in vitro assay utilizing genomic DNA as substrate. *int. j. radiat. biol* **77**, 593–
1746   607 (2001).

1747   72.   Krasner, D. S., Daley, J. M., Sung, P. & Niu, H. Interplay between Ku and Replication Protein A in
1748   the Restriction of Exo1-mediated DNA Break End Resection. *J Biol Chem* **290**, 18806–18816
1749   (2015).

1750   73.   Chen, H., Lisby, M. & Symington, L. S. RPA coordinates DNA end resection and prevents
1751   formation of DNA hairpins. *Mol Cell* **50**, 589–600 (2013).

1752   74.   Shuck, S. C. & Turchi, J. J. Targeted inhibition of RPA reveals cytotoxic activity, synergy with
1753   chemotherapeutic DNA damaging agents and insight into cellular function. *Cancer Res* **70**, 3189
1754   (2010).

1755   75.   Reber, S. *et al.* The phase separation-dependent FUS interactome reveals nuclear and
1756   cytoplasmic function of liquid-liquid phase separation. *Nucleic Acids Res* **49**, 7713–7731 (2021).

1757   76.   Pessina, F. *et al.* Functional transcription promoters at DNA double-strand breaks mediate RNA-
1758   driven phase separation of damage-response factors. *Nat Cell Biol* **21**, 1286–1299 (2019).

1759   77.   Lenard, A. J. *et al.* Phosphorylation Regulates CIRBP Arginine Methylation, Transportin-1 Binding
1760   and Liquid-Liquid Phase Separation. *Front Mol Biosci* **8**, (2021).

1761   78.   Huelsmann, M. *et al.* Genes lost during the transition from land to water in cetaceans highlight
1762   genomic changes associated with aquatic adaptations. *Sci Adv* **5**, 6671 (2019).

1763   79.   Pryor, J. M. *et al.* Essential role for polymerase specialization in cellular nonhomologous end
1764   joining. *Proc Natl Acad Sci U S A* **112**, E4537–E4545 (2015).

80. Moon, A. F., Gosavi, R. A., Kunkel, T. A., Pedersen, L. C. & Bebenek, K. Creative template-dependent synthesis by human polymerase mu. *Proc Natl Acad Sci U S A* **112**, E4530–E4536 (2015).

81. Soniat, M. M., Myler, L. R., Kuo, H. C., Paull, T. T. & Finkelstein, I. J. RPA Phosphorylation Inhibits DNA Resection. *Mol Cell* **75**, 145-153.e5 (2019).

82. Vassin, V. M., Wold, M. S. & Borowiec, J. A. Replication Protein A (RPA) Phosphorylation Prevents RPA Association with Replication Centers. *Mol Cell Biol* **24**, 1930 (2004).

83. Shi, W. *et al.* The role of RPA2 phosphorylation in homologous recombination in response to replication arrest. *Carcinogenesis* **31**, 994–1002 (2010).

84. Nam, K. *et al.* Analysis of the FGF gene family provides insights into aquatic adaptation in cetaceans. *Scientific Reports 2017 7:1* **7**, 1–13 (2017).

85. Tejada-Martinez, D., De Magalhães, J. P. & Opazo, J. C. Positive selection and gene duplications in tumour suppressor genes reveal clues about how cetaceans resist cancer. *Proceedings of the Royal Society B* **288**, (2021).

86. Doherty, A. & de Magalhães, J. P. Has gene duplication impacted the evolution of Eutherian longevity? *Aging Cell* **15**, 978 (2016).

87. Vazquez, J. M., Kraft, M. & Lynch, V. J. A CDKN2C retroduplication in Bowhead whales is associated with the evolution of extremely long lifespans and alerted cell cycle dynamics. *bioRxiv* 2022.09.07.506958 (2022) doi:10.1101/2022.09.07.506958.

88. Keil, G., Cummings, E. & de Magalhães, J. P. Being cool: how body temperature influences ageing and longevity. *Biogerontology* **16**, 383 (2015).

89. Lu, J. Y. *et al.* Comparative transcriptomics reveals circadian and pluripotency networks as two pillars of longevity regulation. *Cell Metab* **34**, 836-856.e5 (2022).

90. Evdokimov, A. *et al.* Naked mole rat cells display more efficient excision repair than mouse cells. *Aging* **10**, 1454–1473 (2018).

91. Tian, X. *et al.* SIRT6 Is Responsible for More Efficient DNA Double-Strand Break Repair in Long-Lived Species. *Cell* **177**, 622-638.e22 (2019).

92. Kolora, S. R. R. *et al.* Origins and evolution of extreme life span in Pacific Ocean rockfishes. *Science* **374**, 842–847 (2021).

93. Suárez-Menéndez, M. *et al.* Wild pedigrees inform mutation rates and historic abundance in baleen whales. *Science* **381**, 990–995 (2023).

94. Yoshioka, K. I., Kusumoto-Matsuo, R., Matsuno, Y. & Ishiai, M. Genomic Instability and Cancer Risk Associated with Erroneous DNA Repair. *International Journal of Molecular Sciences 2021, Vol. 22, Page 12254* **22**, 12254 (2021).

95. Yokoyama, A. *et al.* Age-related remodelling of oesophageal epithelia by mutated cancer drivers. *Nature* **565**, 312–317 (2019).

96. Martincorena, I. *et al.* Somatic mutant clones colonize the human esophagus with age. *Science*
362, 911–917 (2018).

97. Jaiswal, S. *et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J*
Med **371**, 2488–2498 (2014).

98. Martincorena, I. *et al.* High burden and pervasive positive selection of somatic mutations in
normal human skin. *Science (1979)* **348**, 880–886 (2015).

99. Mitchell, E. *et al.* Clonal dynamics of haematopoiesis across the human lifespan. *Nature 2022*
606:7913 **606**, 343–350 (2022).

100. Chen, Y. *et al.* Fight to the bitter end: DNA repair and aging. *Ageing Res Rev* **64**, 101154 (2020).

101. Seluanov, A., Vaidya, A. & Gorbunova, V. Establishing Primary Adult Fibroblast Cultures From
Rodents. *J Vis Exp* **44**, (2010).

102. Elsner, R., Meiselman, H. J. & Baskurt, O. K. TEMPERATURE-VISCOSITY RELATIONS OF BOWHEAD
WHALE BLOOD: A POSSIBLE MECHANISM FOR MAINTAINING COLD BLOOD FLOW. *Mar Mamm Sci*
20, 339–344 (2004).

103. George, J. C. Growth, morphology and energetics of bowhead whales (Balaena mysticetus).
(University of Alaska Fairbanks, 2009).

104. Maslov, A. Y. *et al.* Single-molecule, quantitative detection of low-abundance somatic mutations
by high-throughput sequencing. *Sci Adv* **8**, (2022).

105. Quail, M. A. *et al.* Optimal enzymes for amplifying sequencing libraries. *Nat Methods* **9**, 10–11
(2012).

106. Blackburn, M. C. Development of new tools and applications for high-throughput sequencing of
microbiomes in environmental or clinical samples. (2010).

107. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence analysis.
Nature Biotechnology 2019 37:3 **37**, 224–226 (2019).

108. Lee, H. J., Chang, H. Y., Cho, S. W. & Ji, H. P. CRISPRpic: fast and precise analysis for CRISPR-
induced mutations via prefixed index counting. *NAR Genom Bioinform* **2**, (2020).

109. GitHub - jstjohn/SeqPrep: Tool for stripping adaptors and/or merging paired reads with overlap
into single reads. https://github.com/jstjohn/SeqPrep.

110. Franken, N. A. P., Rodermond, H. M., Stap, J., Haveman, J. & van Bree, C. Clonogenic assay of cells
in vitro. *Nat Protoc* **1**, 2315–2319 (2006).

111. Rose, J. C. *et al.* Rapidly inducible Cas9 and DSB-ddPCR to probe editing kinetics. *Nat Methods* **14**,
891 (2017).

112. Hudon, S. F. *et al.* Primers to highly conserved elements optimized for qPCR-based telomere
length measurement in vertebrates. *Mol Ecol Resour* **21**, 59–67 (2021).

113. Dimri, G. P. *et al.* A biomarker that identifies senescent human cells in culture and in aging skin in
vivo. *Proc Natl Acad Sci U S A* **92**, 9363 (1995).

114. Debacq-Chainiaux, F., Erusalimsky, J. D., Campisi, J. & Toussaint, O. Protocols to detect
senescence-associated beta-galactosidase (SA-βgal) activity, a biomarker of senescent cells in
culture and in vivo. *Nature Protocols 2009 4:12* **4**, 1798–1806 (2009).

115. Mao, Z. *et al.* SIRT6 promotes DNA repair under stress by activating PARP1. *Science (1979)* **332**,
1443–1446 (2011).

116. Vu, D.-D. *et al.* Multivalent interactions of the disordered regions of XLF and XRCC4 foster robust
cellular NHEJ and drive the formation of ligation-boosting condensates in vitro. *bioRxiv* (2023)
doi:10.1101/2023.07.12.548668.

117. Roy, S. *et al.* XRCC4/XLF Interaction Is Variably Required for DNA Repair and Is Not Required for
Ligase IV Stimulation. *Mol Cell Biol* **35**, 3017–3028 (2015).

118. Dasovich, M. *et al.* Identifying Poly(ADP-ribose)-Binding Proteins with Photoaffinity-Based
Proteomics. *J Am Chem Soc* **143**, 3037–3042 (2021).

119. Tan, E. S., Krukenberg, K. A. & Mitchison, T. J. Large-scale preparation and characterization of
poly(ADP-ribose) and defined length polymers. *Anal Biochem* **428**, 126–136 (2012).

120. Tian, X. *et al.* SIRT6 Is Responsible for More Efficient DNA Double-Strand Break Repair in Long-
Lived Species. *Cell* **177**, 622-638.e22 (2019).

121. Benchling. https://www.benchling.com/.

122. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

123. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

124. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol* **11**,
(2010).

125. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, (2014).

126. Lange, S. S., Takata, K. I. & Wood, R. D. DNA polymerases and cancer. *Nat Rev Cancer* **11**, 96–110
(2011).

127. Wood, R. D., Mitchell, M. & Lindahl, T. Human DNA repair genes, 2005. *Mutat Res* **577**, 275–283
(2005).

128. Perelli, L. *et al.* Evolutionary fingerprints of EMT in pancreatic cancers. *bioRxiv* (2023)
doi:10.1101/2023.09.18.558231.

129. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data.
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

130.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data.
1871   *Bioinformatics* **30**, 2114–2120 (2014).

1872   131.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
1873   *Bioinformatics* **25**, 1754–1760 (2009).

1874   132.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078 (2009).

1875   133.   Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls: the Genome
1876   Analysis Toolkit best practices pipeline. *Curr Protoc Bioinformatics* **43**, (2013).

1877   134.   Chen, X. *et al.* Manta: rapid detection of structural variants and indels for germline and cancer
1878   sequencing applications. *Bioinformatics* **32**, 1220–1222 (2016).

1879   135.   Nickson, C. M. & Parsons, J. L. Monitoring regulation of DNA repair activities of cultured cells in-
1880   gel using the comet assay. *Front Genet* **5**, (2014).

1881   136.   Gyori, B. M., Venkatachalam, G., Thiagarajan, P. S., Hsu, D. & Clement, M. V. OpenComet: An
1882   automated tool for comet assay image analysis. *Redox Biol* **2**, 457–465 (2014).

1883   137.   Shechter, D., Dormann, H. L., Allis, C. D. & Hake, S. B. Extraction, purification and analysis of
1884   histones. *Nature Protocols 2007 2:6* **2**, 1445–1457 (2007).

1885   138.   da Veiga Leprevost, F. *et al.* Philosopher: a versatile toolkit for shotgun proteomics data analysis.
1886   *Nature Methods 2020 17:9* **17**, 869–870 (2020).

1887   139.   Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I. MSFragger:
1888   ultrafast and comprehensive peptide identification in mass spectrometry–based proteomics.
1889   *Nature Methods 2017 14:5* **14**, 513–520 (2017).

1890   140.   Ma, K., Vitek, O. & Nesvizhskii, A. I. A statistical model-building perspective to identification of
1891   MS/MS spectra with PeptideProphet. *BMC Bioinformatics* **13 Suppl 16**, 1–17 (2012).

1892   141.   Yu, F., Haynes, S. E. & Nesvizhskii, A. I. IonQuant Enables Accurate and Sensitive Label-Free
1893   Quantification With FDR-Controlled Match-Between-Runs. *Mol Cell Proteomics* **20**, (2021).

1894   142.   Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity
1895   platform for reference generation and analysis. *Nature Protocols 2013 8:8* **8**, 1494–1512 (2013).

1896   143.   Bateman, A. *et al.* UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res* **49**,
1897   D480–D489 (2021).

1898   144.   Adusumilli, R. & Mallick, P. Data Conversion with ProteoWizard msConvert. *Methods Mol Biol*
1899   **1550**, 339–368 (2017).

1900   145.   Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nature
1901   Biotechnology 2012 30:10* **30**, 918–920 (2012).

1902   146.   Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: neural networks
1903   and interference correction enable deep proteome coverage in high throughput. *Nature Methods
1904   2019 17:1* **17**, 41–44 (2019).

1905   147.   Lazar, C., Gatto, L., Ferro, M., Bruley, C. & Burger, T. Accounting for the Multiple Natures of
1906   Missing Values in Label-Free Quantitative Proteomics Data Sets to Compare Imputation
1907   Strategies. *J Proteome Res* **15**, 1116–1125 (2016).

1908   148.   Gardner, M. L. & Freitas, M. A. Multiple imputation approaches applied to the missing value
1909   problem in bottom-up proteomics. *Int J Mol Sci* **22**, (2021).

1910   149.   Liu, M. *et al.* Genomic discovery of potent chromatin insulators for human gene therapy. *Nature*
1911   *Biotechnology 2014 33:2* **33**, 198–203 (2015).

1912

1913

1914

1915

1916

1917

1918

1919

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

1932

1933

1934

 Figure Legends

Figure 1. Bowhead whale fibroblasts exhibit senescence with reduced SASP and
low basal p53 activity. a, Growth curves of primary and hTERT-immortalized skin
fibroblasts (n=2 for each cell line). **b**, Telomerase activity and telomere length in skin
fibroblasts. **c**, Quantification of β-gal–positive human and bowhead skin fibroblasts in
response to γ-irradiation (12 days) and replicative senescence (n=3 for each
species). **d**, Representative images of SA-β-gal staining of human and bowhead skin
fibroblasts in response to γ-irradiation and replicative senescence. The bar is 100 µm. **e**,
Apoptosis of human and bowhead whale fibroblasts in response to γ-irradiation. Three
days after γ-irradiation, cells were harvested and subjected to an Annexin V apoptosis
assay using flow cytometry (n=3 for each species). **f**, Log fold change of SASP mRNA
expression in human and bowhead whale skin fibroblasts 12 days after γ-irradiation. **g**,
Relative luciferase expression in mouse, cow, human and bowhead whale fibroblasts
transfected with the p53 reporter vector. Data are shown as ratios of firefly/renilla
luciferase (to normalize for transfection efficiency) expression 24 h after transfections
(n=3 for mouse, human, BW; n=2 for cow). **h**, Apoptosis of mouse, cow, human and
bowhead whale fibroblasts in response to UVC. Two days after UVC, cells were
harvested and subjected to an Annexin V apoptosis assay using flow cytometry. Error
bars represent mean ± SD. * p<0.05, *** p<0.001. Welch's t-test was used to quantify
the significance. Whale, bowhead whale; NMR, naked mole-rat. RS, replicative
senescence.

Figure 2. Fewer tumor suppressor hits are required for oncogenic transformation
of bowhead fibroblasts than for human fibroblasts. a, Images of representative
fibroblast colonies for tested cell lines after 4 weeks of growth in soft agar. The top
panel indicates whether the cell lines in the column below have the indicated protein
overexpressed (+), inactivated (-), or expressed in the active endogenous form (WT).
Text above individual images indicate for that cell line whether tumor suppressors are
inactivated through genetic knockout or SV40 Large T (or LT mutants) or Small T (ST)
antigen. Icons in corners of images indicate species. Scale bar represents 250 µm. **b,**
Volumetric growth curves for the indicated bowhead whale fibroblast cell lines in mouse
xenograft assays. All cell lines shown stably express H-Ras^{G12V} and *hTERT* in addition
to the genotype indicated in the figure legend. Data points represent averages from 3
immunodeficient nude mice injected bilaterally (6 injections) for each cell line, except for
TP53$^{-/-}$RB1$^{-/-}$ double knockouts, for which 2 independent cell lines were tested, for a
total of 6 mice/12 injections. Experiments were terminated based on predetermined
thresholds for maximum tumor length or duration of experiment as described in
Methods. Images in the legend show a representative mouse for the indicated cell line
at the final measured time point. Error bars show SEM. **c,** Western blot for p53 protein
in clonally isolated fibroblast colonies following CRISPR targeting of *TP53*. Underlined
lanes indicate colonies selected for further validation and experiments. **d,** Western blot

for Rb protein in clonally isolated fibroblast colonies following CRISPR targeting of *RB1*
on an existing p53 knockout background.

Figure 3. The bowhead whale repairs DSBs with higher accuracy and efficiency
than other mammals. a,b, NHEJ and HR efficiency were measured using fluorescent
reporter constructs. Successful NHEJ and HR event leads to reactivation of the GFP
gene. NHEJ and HR reporter constructs were integrated into primary, low passage skin
fibroblasts. NHEJ and HR repair efficiency were assayed by transfecting cells with I-
SceI expression vector and a DsRed plasmid as a transfection control. The repair
efficiency was calculated as the ratio of GFP+/DsRed+ cells. Experiments were
repeated at least 3 times for each cell line. Error bars represent SD. * $p < 0.05$ (Welch's t-
test). Whale, bowhead whale. **c**, Percent of binucleated cells containing micronuclei in
human and bowhead whale fibroblasts after 2Gy γ-irradiation (n=4). Error bars
represent SD. ** $p < 0.005$ (Welch's t-test). **d,** Endogenous γ-H2AX/53BP1 foci in human
and whale cells. Results are combined from two independent experiments. 200 nuclei
were analyzed. Error bars represent SEM. *** $p < 0.001$ (two-tailed t-test). **e,**
Representative confocal images of human and bowhead whale cells stained for γ-
H2AX and 53BP1 at no treatment (control) and 1h-24h after bleomycin treatment at
concertation 5μg/mL. Scale bar indicates 10 μm. **f,** Quantification of γH2AX/53BP1 foci
with and without DSB induction. Exponentially growing cells were treated for 1h with
bleomycin (BLM) at concertation 5μg/mL, washed twice with PBS and fresh media was
added. At indicated time-points cells were fixed and processed for
immunocytochemistry. Foci were counted by eye in green and red channels. 150-170
nuclei were analyzed. Error bars represent SEM. * $p < 0.05$, ** $p < 0.01$ (two-tailed t-test).
g, Histograms of CRISPR indel size distribution by species. Data for biological
replicates are superimposed and partially transparent with lines connecting data points
for each sample. Unmodified alleles and alleles with substitutions only are excluded
from this analysis. **h**, Distribution of sequenced PTEN allele variants by species after
CRISPR DSB induction at a conserved region of the endogenous PTEN gene. Data are
averages from multiple primary dermal fibroblast lines isolated from different individual
animals for bowhead whale (n=3), human (n=3), cow (n=2), and mouse (n=3). Error
bars represent SEM. **i**, Allele plots showing 15 most frequent allele types after CRISPR
for one representative cell line per species. Sequences are displayed within a window
centered on the cleavage site and extending 20 bp in each direction. Data bars and
values indicate proportion of total alleles represented by each row. For the purposes of
this display and quantification, all individual alleles with identical sequences in the 40-bp
window have been pooled, so rows represent composites of alleles that may differ
outside the display window.

Figure 4. CIRBP is highly expressed in bowhead whale fibroblasts and promotes
DNA DSB repair and genome stability. a, Western blots of DNA repair proteins in
primary fibroblasts from different species. **b-c**, bwCIRBP promotes NHEJ and HR in
human cells as measured by flow cytometric GFP-reporter assays (see Methods). In
these assays DSBs are induced within inactive NHEJ or HR reporter cassettes by

expressing I-SceI endonuclease. Successful NHEJ or HR events lead to reactivation of
the fluorescent GFP reporters that are scored by flow cytometry. All experiments in
these figures were repeated at least 3 times. **d,e**, Knockdown of CIRBP in bowhead
whale fibroblasts decreases NHEJ and HR efficiency. siNT = non-targeting siRNA. **f**,
Western blot of human fibroblasts overexpressing human CIRBP, whale CIRBP or 9R/A
mutated whale CIRBP; **g,** Western blot of bowhead whale fibroblasts with knockdown of
CIRBP. **h**, Overexpression of CIRBP decreases the percentage of binucleated cells
containing micronuclei in human cells 3d after 2Gy γ-irradiation (n=3) (left panel);
Western blot of human fibroblasts overexpressing human CIRBP, human CIRBP with
optimized codons and whale CIRBP (right panel). Error bars represent mean ± SD. *
$p<0.05$, ** $p<0.01$, *** $p<0.001$ (Welch's t-test). siNT - negative control siRNAs that do
not target any gene product. **i,** Number of endogenous γH2AX/53BP1 foci in human
cells with bwCIRBP overexpression. Error bars represent SEM. *** $p<0.001$ (two-tailed
t-test). **j,** CIRBP stimulates NHEJ-mediated ligation *in vitro in*. Linearized pUC19
plasmid with cohesive ends was mixed with human recombinant proteins XRCC4/
Ligase IV complex, and 0 to 1 μM CIRBP. Where indicated, reaction mixtures contained
Ku70/80 heterodimer, PAXX dimer or XLF dimer. The reaction mixtures were incubated
for 1 hr at 30°C, proteins were denatured with SDS at 65°C and loaded onto agarose
gel. Each sample were loaded onto 0.7% agarose gel, followed by gel electrophoresis.
The gel was stained with ethidium bromide.

Figure 5. RPA and CIRBP contribute to increased DNA repair fidelity. a,
Distribution of sequenced PTEN allele variants in human primary fibroblasts treated with
TDRL-505 or rhRPA after CRISPR DSB induction at a conserved region of the
endogenous PTEN gene. Data are averages from experiments performed in triplicate.
Error bars represent SEM. **b,** Distribution of sequenced PTEN allele variants by species
in bowhead whale primary fibroblasts treated with TDRL-505 after CRISPR DSB
induction at a conserved region of the endogenous PTEN gene. Data are averages from
experiments performed in triplicate. Error bars represent SEM. **c,** Distribution of
sequenced PTEN allele variants by species in human fibroblasts with lentiviral
overexpression of luciferase or bwCIRBP after CRISPR DSB induction at a conserved
region of the endogenous PTEN gene. Data are averages from experiments performed
in triplicate. Error bars represent SEM. * $p<0.05$, **** $p<0.0001$. All charts analyzed by
two-way ANOVA with Fisher's LSD. p-values should be considered nominal indices of
significance. **d,** Graphical summary. The bowhead whale has evolved efficient and
accurate DSB repair mediated by high levels of CIRBP and RPA2. This enhanced DNA
repair may help the bowhead whale resist cancer despite its cells requiring fewer
mutational hits for malignant transformation than human cells. Improved DNA repair
rather than enhanced elimination of damaged cells through apoptosis or senescence
may promote longevity in the bowhead whale.

Extended Data Figure 1. Mutation rates in bowhead whale cells during tumor
progression. a, Western blot for p53 protein in clonally isolated fibroblast colonies
following CRISPR targeting of *TP53*. Underlined lanes indicate colonies selected for

further validation and experiments. **b,** Western blot for Rb protein in clonally isolated
fibroblast colonies following CRISPR targeting of *RB1* on an existing p53 knockout
background. **c,** Ratio of firefly:renilla luciferase luminescence in fibroblasts transfected
with firefly luciferase reporter of p53 transcriptional activity and renilla luciferase control.
Cells were treated with etoposide to induce p53 activity. **d,** Ratio of firefly:renilla
luciferase luminescence in fibroblasts transfected with firefly luciferase reporter of E2F
transcriptional activity and renilla luciferase control. Transfected cells were serum
starved for 24h and returned to complete medium for 24h before luminescence
measurement. Higher E2F activity results from reduced Rb activity. Error bars represent
SD. p<0.001 (two-tailed t test), n=3. **e,** Schematic showing experimental design and
samples processed for WGS (whale N = 9 tumors; human N = 2 tumors; mouse N = 1
tumor). **f,** Bar plot displaying percentages of SNV types across species with similarities
of mutational processes across species. **g-l,** Bar plot showing quantifications of
numbers of SNVs and small indels (size 1-10bp) across species. **j-l**, Bar plot showing
quantification of number of large SVs (size > 6000bp) across species. **m,** Histograms
and trend curves showing distribution of SVs size across species. **n,** Bar plot showing
distribution of small, medium and large (6-50Kb, 50-500Kb, >500Kb respectively) SVs
and deletions across species. Error bars represent SD. P values are a result of ordinary
One-Way Anova with Tukey's multiple comparison test (**g-l**) and chi-square test (**n**). * p
< 0.05; ** p < 0.01; *** p < 0.0001; ns = not significant.

Extended Data Figure 2. Mismatch repair, excision repair, and mutagenesis in
bowhead whale cells. a, MMR reactivation of a heteroduplex eGFP plasmid containing
a G/T mismatch. Growing fibroblasts were transfected with the heteroduplex plasmid
and a DsRed plasmid as a transfection control. The repair efficiency was calculated as
the ratio of GFP+/DsRed+ cells. Each dot represents cell line isolated from different
individual (n=3). **b**, NER efficiency was measured by host cell reactivation assay where
a plasmid containing luciferase reporter is UV-irradiated *in vitro* to induce DNA damage,
transfected into cells, and reactivation of the reporter is measured (n=3 for each cell
line). **c**, Kinetics of cyclobutane pyrimidine dimer repair after 30 J/m^2 UVC. Confluent
human and whale skin fibroblasts were subjected to UVC, harvested at different time-
points, genomic DNA was isolated and analyzed for cyclobutene dimers as described in
Methods (n=2 for each cell line). **d**, BER efficiency was measured by host cell
reactivation where luciferase reporter plasmid is treated with methylene blue and light to
induce oxidative DNA damage, transfected into cells, and luciferase activity measured
as described in Methods. **e**, ENU-induced mutational load by SMM-seq in fibroblasts of
the indicated species. Delta SNV frequency was calculated for each cell line (n=6-8
fibroblasts/species; Kruskal-Wallis test). **f**, Analysis of mutational spectra showing a
pattern typical for ENU. An increase in A>T transversions (orange bars) can be found in
ENU-treated mammalian cells. **g**, HPRT mutagenesis assay in ENU-treated cells,
adjusted by plating efficiency measured for each cell line (n=3 cell lines/species) **h**,
Colony forming efficiency for HPRT mutagenesis assay. Error bars represent mean ±
SD. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001 ns=not significant (heteroscedastic
two-tailed t test). **i,** Apoptosis/necrosis of human and bowhead whale fibroblasts in

response to ENU. Cells at growing stage were treated for 3h with ENU at indicated dosages. After treatment cells were washed in PBS and incubated for 3 days. For measuring Apoptosis/Necrosis cells were stained with AnnexinV/PI and analyzed by flow cytometry. **j**, HPRT mutagenesis assay in cells treated with 2 Gy γ-irradiation, adjusted by plating efficiency measured for each cell line (n=2 cell lines/species) **k**, Colony forming efficiency for HPRT mutagenesis assay.

Extended Data Figure 3. Poly-ADP-ribosylation and DNA repair of oxidative damage in bowhead whale cells. a, Bowhead whale cells show greater poly-ADP-ribosylation in response to hydrogen peroxide treatment. **b,** Bowhead whale cells show greater poly-ADP-ribosylation after γ-irradiation. Cells were harvested immediately or at indicated time-points after radiation for Western blot analysis (top panel). Representative images of comet tails under neutral conditions (bottom panel). Cells were processed immediately after radiation. **c**, Nuclear extracts of bowhead whale fibroblasts exhibit higher endogenous PARP activity (n=3). Error bars represent mean ± SD. * $p < 0.05$ (Welch's t-test). Whale=bowhead whale. **d,** Apoptosis/Necrosis of human and bowhead whale fibroblasts in response to hydrogen peroxide at concentration 700μM. Two days after hydrogen peroxide, cells were harvested and subjected to an Annexin V apoptosis assay using flow cytometry. Error bars represent mean ± SD. *** $p < 0.001$. Welch's t-test was used to quantify the significance (n=12). **e,** Percent tail DNA by alkaline comet assay at various time points after 700 μM H_2O_2 treatment in 2 cell lines each of human and bowhead whale fibroblasts. Points represent individual cells. Representative comet images shown below. Bars indicate mean +- SEM.

Extended Data Figure 4. DSB repair efficiency in bowhead whale cells. a, NHEJ efficiency in extrachromosomal assay. NHEJ reporter construct was pre-digested with I-SceI, purified and co-transfected with DsRed into human and bowhead skin fibroblasts. Three days after transfection cells were harvested and subjected to flow cytometry to calculate NHEJ efficiency (n=3). Error bars represent mean ± SD. *** $p < 0.001$ (two-tailed t-test) **b,** Representative images of human and bowhead whale binucleated cells containing micronuclei after 2 Gy of γ-irradiation. Scale bar indicates 20 μm. **c,** Frequency of micronuclei after DSB induction with I-SceI in primary fibroblasts carrying a chromosomally integrated NHEJ reporter cassette. Each cell line was transiently transfected with a BFP-expressing control plasmid or an I-SceI expression plasmid and micronuclei were quantified after 5d in media containing cytochalasin B to prevent cytokinesis. Micronucleus frequencies for each cell line are shown normalized to BFP control (paired t-test, n=3 cell lines/species). **d,** Pulse-field gel stained with ethidium bromide, showing chromosomal DNA fragmentation in human and bowhead confluent skin fibroblasts immediately after different doses of γ-irradiation 0.7, 1.5, 3 and 6h after 40 Gy of γ-irradiation. **e,** Kinetics of DSB repair measured by PFGE in confluent human and bowhead fibroblasts after 40 Gy of γ-irradiation. n=2 for each species.

Extended Data Figure 5. Sequencing of DNA DSB repair products in bowhead whale cells. a, Possible repair outcomes after induction of DSBs with incompatible

ends by I-SceI in NHEJ reporter construct. **b,** Allele plot of Sanger sequencing products resulting from repair of integrated NHEJ reporter cassette after I-SceI cleavage. **c,** NHEJ fidelity in extrachromosomal assay. NHEJ reporter construct was pre-digested with I-SceI, purified and co-transfected with DsRed into human and bowhead skin fibroblasts. Three days after transfection genomic DNA was isolated, subjected to PCR, cloned and analyzed by Sanger sequencing. At least 100 clones were analyzed for each species. Correct – annealing on 2 of the 4 protruding nucleotides **d,** Time course of CRISPR cleavage measured by digital droplet PCR (ddPCR). PTEN copy number at varying time points after CRISPR RNP transfection was measured with ddPCR using primers flanking the predicted cleavage site and normalized within each sample to a single-copy genomic ultraconserved element as described in Methods. Error bars show confidence intervals of Poisson distribution calculated in QuantaSoft. **e,** Pearson correlation between 5^{th} percentile indel size and species lifespan (r=0.8508, 95% CI = 0.5125 to 0.9605, p=0.0009, n=11). **f,** Absolute frequencies of alleles by base pairs of microhomology across species in CRISPR-targeted PTEN repair products. **g,** Relative proportions of deletion alleles by base pairs of microhomology across species in CRISPR-targeted PTEN repair products.

Extended Data Figure 6. Proteomic quantification of DNA repair proteins. a, Western blot for CIRBP on bowhead whale and mouse organs **b,** Abundance of CIRBP protein by LC-MS in liver tissue of mammal species (n=12 per species; 3 biological x 4 technical replicates; N.D.=not detected) **c,** Abundance of RPA2 protein by LC-MS in liver tissue of mammal species (n=12 per species; 3 biological x 4 technical replicates; N.D.=not detected) **d,** Abundance of CIRBP protein by LC-MS in nuclear extracts of liver tissue of mammal species (n=3 biological replicates per species) **e,** Abundance of RPA2 protein by LC-MS in nuclear extracts of liver tissue of mammal species (n=3 biological replicates per species; N.D.=not detected) **f,** Heatmap of LC-MS protein abundance for primary fibroblasts of the indicated species and proteins. Color intensity scale corresponds to $\log_{10}$ ion intensity. **g,** Per-protein normalized abundance by LC-MS of proteins identified in pulldowns of His-tagged Cas9/dCas9 bound to a plasmid containing the genomic *PTEN* target sequence after incubation in extracts of soluble nuclear proteins from human and bowhead whale.

Extended Data Figure 7. Transcriptome, Western blot, and STED quantification of DNA repair proteins. a, Relative expression level of genes in 6 DNA repair pathways among species. Z-scores are scaled by row. Genes in each pathway are ordered decreasingly based on the expression level in bowhead whale. Genes with higher expression in bowhead whale compared to all 3 other species are highlighted in red text to the right of the heatmap. Genes of each gene set were compiled from 3 resources: MsigDB database, GO ontology, and a curated gene list (www.mdanderson.org/documents/Labs/Wood-Laboratory/human-dna-repair-genes.html) **b,** Western blot abundance of RPA2 in cultured skin fibroblasts, using 2 different monoclonal primary antibodies targeting conserved epitopes and normalized to histone H3. A third polyclonal antibody produced the same results but had higher

background reactivity and is not shown. Each lane is a primary fibroblast line from a different adult individual. Fluorescent secondary antibodies were used to increase linear dynamic range for higher quantitative accuracy. **c,** Western blot for CIRBP with 3 different antibodies in 3 fibroblast lines per species. mAb=monoclonal antibody, pAb=polyclonal antibody. **d,** Stimulated emission depletion (STED) images of RPA2 and CIRBP localization in human and bowhead whale fibroblasts. Target protein in red, nuclear DAPI stain in blue. **e,** Western blot for CIRBP in fibroblasts isolated from various mammalian species.

Extended Data Figure 8. Analysis of CIRBP's role in DNA DSB repair. a, CIRBP localization in whale cells. Before formaldehyde fixation, cells were pre-extracted with CSK buffer +/- RNAseA for 3min. After standard immunocytochemistry procedure images were collected using confocal microscope. **b,** Western blot of bowhead whale fibroblasts with knockdown of CIRBP (left panel) and band intensity quantification from 3 independent experiments (right panel) suggesting partial dependence of RPA2 protein abundance on CIRBP expression. **c,** Ion intensity by LC-MS of RPA2 in human fibroblasts with and without lentiviral overexpression of bwCIRBP (n=3 human cell lines). Error bars show mean +-SEM. **d,** DSBs induce CIRBP enrichment in chromatin. Exponentially growing cells were treated with neocarzinostatin (NCS) for the indicated period of time and lysed in CSK buffer to enrich chromatin-bound fraction. α-Tubulin staining was used to verify the absence of cytoplasmic contamination in chromatin-bound fraction. **e,** DSBs induced by γ-irradiation lead to CIRBP enrichment in chromatin. This enrichment is promoted by RNA. Exponentially growing cells were treated with γ-irradiation and at the indicated period of time were lysed in CSK buffer with/without RNAse A to enrich proteins in chromatin-bound fraction. **f,** Overexpression of CIRBP decreases the percentage of binucleated cells containing micronuclei in human cells after I-Sce1-induced DSBs. Each bar indicates an experimental replicate. At least 150 binucleated cells were scored per condition. **g,** Frequency of chromosomal aberrations in human fibroblasts with and without CIRBP overexpression after 2Gy γ-irradiation. 100 metaphases were analyzed per sample. C=control untreated cells. **h,** Frequency of insertions and deletions >20 bp in NHEJ reporter constructs PCR-amplified from human fibroblasts with and without bwCIRBP overexpression after I-SceI expression. Insertion/deletion frequencies were determined from Nanopore sequencing data of PCR products and normalized within each sample to total frequency of all insertions or deletions. **i,** Frequency of insertions and deletions as shown in (**h**) but for bowhead whale fibroblasts with negative control or CIRBP-targeting siRNAs. **j,** Calculated dissociation constants (K_D) and fluorescence polarization (FP) measurements for CIRBP proteins titrated into solutions containing a fixed concentration (3 nM) of fluorescently labeled PAR of various polymer lengths. **k,** EMSA of increasing amounts of recombinant human CIRBP incubated in vitro with 300 ng sheared chromatin from fibroblasts exposed to UVC and oxidative DNA damage as described in Methods. Chromatin was treated with Proteinase K but not RNAse. Nucleic acids are stained with SYBR Gold. Red overlay indicates saturated pixels. **l,** EMSA of

300 ng sheared purified genomic DNA, purified cellular RNA, or chromatin as described
in (**k**) incubated in vitro with 5 µg rhCIRBP. **m,** Hypothermia promotes NHEJ efficiency
in primary human fibroblasts (left panel). Cells were pre-incubated at 33°C for 2 days,
co-transfected with I-SceI-digested NHEJ reporter and DsRed, and returned to the 33°C
incubator. NHEJ efficiency was measured by flow cytometry 3 days following
transfection (n=3). Western blot showing CIRBP upregulation in human cells exposed to
33°C for 2 days (right panel). Western blot images were analyzed in ImageLab software
(Bio-Rad). Error bars represent mean ± SD. ** $p < 0.01$ (Welch's t-test).

Extended Data Figure 9. Analysis of bwCIRBP coding sequence mutations and
protein expression levels. a, Comparison of amino acid sequences between human
and bowhead whale CIRBP through BLAST analysis. **b,** Phylogenetic tree illustrating
the relationships among CIRBP coding sequences from representative species with
genome sequence information available. The asterisk indicates the presence of BHW-
specific variants in the species. The colors indicate the position of variants shown in (**a**).
c, SwissModel/AlphaFold models of human (left, pink) and bowhead whale (right, blue).
Side chains of whale residues that diverge from human are shown, and their ribbon is
colored yellow in the model. The key takeaway is that all the residues that differ
between whale and human are in the C-terminal disordered region, whereas the N-
terminal RNA recognition motif (RRM) is structured and conserved. **d,** Western blot
abundance of bwCIRBP, hCIRBP, and reciprocal amino acid mutants overexpressed in
human cells. **e,** Calculated codon adaptation index (CAI) for CIRBP coding sequence
variants.

Extended Data Figure 10. Analysis of bowhead whale RPA2 sequence. a,
Comparison of amino acid sequences between human and bowhead whale RPA2
through BLAST analysis. **b,** Phylogenetic tree illustrating the relationships among RPA2
coding sequence from different representative species. The asterisk indicates the
presence of BHW-specific variants in the species. The colors indicate the position of
variants shown in (**a**). **c,** AlphaFold protein structures of human and bowhead whale
RPA2 showing the position of the variants.

Extended Data Figure 11. Bowhead whale CIRBP reduces anchorage-independent
cell growth. a, Images of representative human transformed fibroblast colonies with
and without bwCIRBP overexpression after 23 days of growth in soft agar. 20x
magnification. Bar 100µm. **b,** Quantification of colonies after staining with nitro blue
tetrazolium chloride. Colonies were counted using ImageJ software as described in
Methods. Error bars represent SD. *$p < 0.05$ (Welch's t-test). **c,** Cell proliferation MTT
assay. **d,** Trypan Blue exclusion test of cell viability. **e,** Western blot showing expression
of LT, Ras, p16 and p21 after overexpression of bwCIRBP. **f,** Frequency of
chromosomal aberrations in human transformed cells after bwCIRBP overexpression.
100 metaphases were analyzed per sample.

2267

Figure 1

Figure 1. Bowhead whale fibroblasts exhibit senescence with reduced SASP and low basal p53 activity. a, Growth curves of primary and hTERT-immortalized skin fibroblasts (n=2 for each cell line). **b**, Telomerase activity and telomere length in skin fibroblasts. **c**, Quantification of β-gal–positive human and bowhead skin fibroblasts in response to γ-irradiation (12 days) and replicative senescence (n=3 for each species). **d**, Representative images of SA-β-gal staining of human and bowhead skin fibroblasts in response to γ-irradiation and replicative senescence. The bar is 100 μm. **e**, Apoptosis of human and bowhead whale fibroblasts in response to γ-irradiation. Three days after γ-irradiation, cells were harvested and subjected to an Annexin V apoptosis assay using flow cytometry (n=3 for each species). **f**, Log fold change of SASP mRNA expression in human and bowhead whale skin fibroblasts 12 days after γ-irradiation. **g**, Relative luciferase expression in mouse, cow, human and bowhead whale fibroblasts transfected with the p53 reporter vector. Data are shown as ratios of firefly/renilla luciferase (to normalize for transfection efficiency) expression 24 h after transfections (n=3 for mouse, human, BW; n=2 for cow). **h**, Apoptosis of mouse, cow, human and bowhead whale fibroblasts in response to UVC. Two days after UVC, cells were harvested and subjected to an Annexin V apoptosis assay using flow cytometry. Error bars represent mean ± SD. * $p<0.05$, *** $p<0.001$. Welch's t-test was used to quantify the significance. Whale, bowhead whale; NMR, naked mole-rat. RS, replicative senescence.

Figure 2

Figure 2. Fewer tumor suppressor hits are required for oncogenic transformation of bowhead fibroblasts than for human fibroblasts. a, Images of representative fibroblast colonies for tested cell lines after 4 weeks of growth in soft agar. The top panel indicates whether the cell lines in the column below have the indicated protein overexpressed (+), inactivated (-), or expressed in the active endogenous form (WT). Text above individual images indicate for that cell line whether tumor suppressors are inactivated through genetic knockout or SV40 Large T (or LT mutants) or Small T (ST) antigen. Icons in corners of images indicate species. Scale bar represents 250 μm. **b,** Volumetric growth curves for the indicated bowhead whale fibroblast cell lines in mouse xenograft assays. All cell lines shown stably express H-Ras^{G12V} and *hTERT* in addition to the genotype indicated in the figure legend. Data points represent averages from 3 immunodeficient nude mice injected bilaterally (6 injections) for each cell line, except for *TP53*$^{-/-}$*RB1*$^{-/-}$ double knockouts, for which 2 independent cell lines were tested, for a total of 6 mice/12 injections. Experiments were terminated based on predetermined thresholds for maximum tumor length or duration of experiment as described in Methods. Images in the legend show a representative mouse for the indicated cell line at the final measured time point. Error bars show SEM. **c,** Western blot for p53 protein in clonally isolated fibroblast colonies following CRISPR targeting of *TP53*. Underlined lanes indicate colonies selected for further validation and experiments. **d,** Western blot for Rb protein in clonally isolated fibroblast colonies following CRISPR targeting of *RB1* on an existing p53 knockout background.

Figure 3

Figure 3. The bowhead whale repairs DSBs with higher accuracy and efficiency than other mammals. a,b, NHEJ and HR efficiency were measured using fluorescent reporter constructs. Successful NHEJ and HR event leads to reactivation of the GFP gene. NHEJ and HR reporter constructs were integrated into primary, low passage skin fibroblasts. NHEJ and HR repair efficiency were assayed by transfecting cells with I-SceI expression vector and a DsRed plasmid as a transfection control. The repair efficiency was calculated as the ratio of GFP+/DsRed+ cells. Experiments were repeated at least 3 times for each cell line. Error bars represent SD. * $p<0.05$ (Welch's t-test). Whale, bowhead whale. **c**, Percent of binucleated cells containing micronuclei in human and bowhead whale fibroblasts after 2Gy γ-irradiation (n=4). Error bars represent SD. ** $p<0.005$ (Welch's t-test). **d,** Endogenous γ-H2AX/53BP1 foci in human and whale cells. Results are combined from two independent experiments. 200 nuclei were analyzed. Error bars represent SEM. *** $p<0.001$ (two-tailed t-test). **e,** Representative confocal images of human and bowhead whale cells stained for γ-H2AX and 53BP1 at no treatment (control) and 1h-24h after bleomycin treatment at concertation 5μg/mL. Scale bar indicates 10 μm. **f,** Quantification of γH2AX/53BP1 foci with and without DSB induction. Exponentially growing cells were treated for 1h with bleomycin (BLM) at concertation 5μg/mL, washed twice with PBS and fresh media was added. At indicated time-points cells were fixed and processed for immunocytochemistry. Foci were counted by eye in green and red channels. 150-170 nuclei were analyzed. Error bars represent SEM. * $p<0.05$, ** $p<0.01$ (two-tailed t-test). **g**, Histograms of CRISPR indel size distribution by species. Data for biological replicates are superimposed and partially transparent with lines connecting data points for each sample. Unmodified alleles and alleles with substitutions only are excluded from this analysis. **h**, Distribution of sequenced PTEN allele variants by species after CRISPR DSB induction at a conserved region of the endogenous PTEN gene. Data are averages from multiple primary dermal fibroblast lines isolated from different individual animals for bowhead whale (n=3), human (n=3), cow (n=2), and mouse (n=3). Error bars represent SEM. **i**, Allele plots showing 15 most frequent allele types after CRISPR for one representative cell line per species. Sequences are displayed within a window centered on the cleavage site and extending 20 bp in each direction. Data bars and values indicate proportion of total alleles represented by each row. For the purposes of this display and quantification, all individual alleles with identical sequences in the 40-bp window have been pooled, so rows represent composites of alleles that may differ outside the display window.

Figure 4

Figure 4. CIRBP is highly expressed in bowhead whale fibroblasts and promotes DNA DSB repair and genome stability. a, Western blots of DNA repair proteins in primary fibroblasts from different species. **b-c**, bwCIRBP promotes NHEJ and HR in human cells as measured by flow cytometric GFP-reporter assays (see Methods). In these assays DSBs are induced within inactive NHEJ or HR reporter cassettes by expressing I-SceI endonuclease. Successful NHEJ or HR events lead to reactivation of the fluorescent GFP reporters that are scored by flow cytometry. All experiments in these figures were repeated at least 3 times. **d,e**, Knockdown of CIRBP in bowhead whale fibroblasts decreases NHEJ and HR efficiency. siNT = non-targeting siRNA. **f**, Western blot of human fibroblasts overexpressing human CIRBP, whale CIRBP or 9R/A mutated whale CIRBP; **g,** Western blot of bowhead whale fibroblasts with knockdown of CIRBP. **h**, Overexpression of CIRBP decreases the percentage of binucleated cells containing micronuclei in human cells 3d after 2Gy γ-irradiation (n=3) (left panel); Western blot of human fibroblasts overexpressing human CIRBP, human CIRBP with optimized codons and whale CIRBP (right panel). Error bars represent mean ± SD. * $p<0.05$, ** $p<0.01$, *** $p<0.001$ (Welch's t-test). siNT - negative control siRNAs that do not target any gene product. **i,** Number of endogenous γH2AX/53BP1 foci in human cells with bwCIRBP overexpression. Error bars represent SEM. *** $p<0.001$ (two-tailed t-test). **j,** CIRBP stimulates NHEJ-mediated ligation *in vitro in*. Linearized pUC19 plasmid with cohesive ends was mixed with human recombinant proteins XRCC4/Ligase IV complex, and 0 to 1 µM CIRBP. Where indicated, reaction mixtures contained Ku70/80 heterodimer, PAXX dimer or XLF dimer. The reaction mixtures were incubated for 1 hr at 30°C, proteins were denatured with SDS at 65°C and loaded onto agarose gel. Each sample were loaded onto 0.7% agarose gel, followed by gel electrophoresis. The gel was stained with ethidium bromide.

Figure 5

Figure 5. RPA and CIRBP contribute to increased DNA repair fidelity. a, Distribution of sequenced PTEN allele variants in human primary fibroblasts treated with TDRL-505 or rhRPA after CRISPR DSB induction at a conserved region of the endogenous PTEN gene. Data are averages from experiments performed in triplicate. Error bars represent SEM. **b,** Distribution of sequenced PTEN allele variants by species in bowhead whale primary fibroblasts treated with TDRL-505 after CRISPR DSB induction at a conserved region of the endogenous PTEN gene. Data are averages from experiments performed in triplicate. Error bars represent SEM. **c,** Distribution of sequenced PTEN allele variants by species in human fibroblasts with lentiviral overexpression of luciferase or bwCIRBP after CRISPR DSB induction at a conserved region of the endogenous PTEN gene. Data are averages from experiments performed in triplicate. Error bars represent SEM. * $p<0.05$, **** $p<0.0001$. All charts analyzed by two-way ANOVA with Fisher's LSD. p-values should be considered nominal indices of significance. **d,** Graphical summary. The bowhead whale has evolved efficient and accurate DSB repair mediated by high levels of CIRBP and RPA2. This enhanced DNA repair may help the bowhead whale resist cancer despite its cells requiring fewer mutational hits for malignant transformation than human cells. Improved DNA repair rather than enhanced elimination of damaged cells through apoptosis or senescence may promote longevity in the bowhead whale.

Extended Data Figure 1

Extended Data Figure 1. Mutation rates in bowhead whale cells during tumor progression. a, Western blot for p53 protein in clonally isolated fibroblast colonies following CRISPR targeting of *TP53*. Underlined lanes indicate colonies selected for further validation and experiments. **b,** Western blot for Rb protein in clonally isolated fibroblast colonies following CRISPR targeting of *RB1* on an existing p53 knockout background. **c,** Ratio of firefly:renilla luciferase luminescence in fibroblasts transfected with firefly luciferase reporter of p53 transcriptional activity and renilla luciferase control. Cells were treated with etoposide to induce p53 activity. **d,** Ratio of firefly:renilla luciferase luminescence in fibroblasts transfected with firefly luciferase reporter of E2F transcriptional activity and renilla luciferase control. Transfected cells were serum starved for 24h and returned to complete medium for 24h before luminescence measurement. Higher E2F activity results from reduced Rb activity. Error bars represent SD. $p < 0.001$ (two-tailed t test), n=3. **e,** Schematic showing experimental design and samples processed for WGS (whale N = 9 tumors; human N = 2 tumors; mouse N = 1 tumor). **f,** Bar plot displaying percentages of SNV types across species with similarities of mutational processes across species. **g-l,** Bar plot showing quantifications of numbers of SNVs and small indels (size 1-10bp) across species. **j-l,** Bar plot showing quantification of number of large SVs (size > 6000bp) across species. **m,** Histograms and trend curves showing distribution of SVs size across species. **n,** Bar plot showing distribution of small, medium and large (6-50Kb, 50-500Kb, >500Kb respectively) SVs and deletions across species. Error bars represent SD. P values are a result of ordinary One-Way Anova with Tukey's multiple comparison test (**g-l**) and chi-square test (**n**). * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$; ns = not significant.

Extended Data Figure 2

Extended Data Figure 2. Mismatch repair, excision repair, and mutagenesis in bowhead whale cells. a, MMR reactivation of a heteroduplex eGFP plasmid containing a G/T mismatch. Growing fibroblasts were transfected with the heteroduplex plasmid and a DsRed plasmid as a transfection control. The repair efficiency was calculated as the ratio of GFP+/DsRed+ cells. Each dot represents cell line isolated from different individual (n=3). **b**, NER efficiency was measured by host cell reactivation assay where a plasmid containing luciferase reporter is UV-irradiated *in vitro* to induce DNA damage, transfected into cells, and reactivation of the reporter is measured (n=3 for each cell line). **c**, Kinetics of cyclobutane pyrimidine dimer repair after 30 J/m^2 UVC. Confluent human and whale skin fibroblasts were subjected to UVC, harvested at different time-points, genomic DNA was isolated and analyzed for cyclobutene dimers as described in Methods (n=2 for each cell line). **d**, BER efficiency was measured by host cell reactivation where luciferase reporter plasmid is treated with methylene blue and light to induce oxidative DNA damage, transfected into cells, and luciferase activity measured as described in Methods. **e**, ENU-induced mutational load by SMM-seq in fibroblasts of the indicated species. Delta SNV frequency was calculated for each cell line (n=6-8 fibroblasts/species; Kruskal-Wallis test). **f**, Analysis of mutational spectra showing a pattern typical for ENU. An increase in A>T transversions (orange bars) can be found in ENU-treated mammalian cells. **g**, HPRT mutagenesis assay in ENU-treated cells, adjusted by plating efficiency measured for each cell line (n=3 cell lines/species) **h**, Colony forming efficiency for HPRT mutagenesis assay. Error bars represent mean ± SD. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001 ns=not significant (heteroscedastic two-tailed t test). **i,** Apoptosis/necrosis of human and bowhead whale fibroblasts in response to ENU. Cells at growing stage were treated for 3h with ENU at indicated dosages. After treatment cells were washed in PBS and incubated for 3 days. For measuring Apoptosis/Necrosis cells were stained with AnnexinV/PI and analyzed by flow cytometry. **j**, HPRT mutagenesis assay in cells treated with 2 Gy γ-irradiation, adjusted by plating efficiency measured for each cell line (n=2 cell lines/species) **k**, Colony forming efficiency for HPRT mutagenesis assay.

Extended Data Figure 3

Extended Data Figure 3. Poly-ADP-ribosylation and DNA repair of oxidative damage in bowhead whale cells. a, Bowhead whale cells show greater poly-ADP-ribosylation in response to hydrogen peroxide treatment. **b,** Bowhead whale cells show greater poly-ADP-ribosylation after γ-irradiation. Cells were harvested immediately or at indicated time-points after radiation for Western blot analysis (top panel). Representative images of comet tails under neutral conditions (bottom panel). Cells were processed immediately after radiation. **c**, Nuclear extracts of bowhead whale fibroblasts exhibit higher endogenous PARP activity (n=3). Error bars represent mean ± SD. * $p<0.05$ (Welch's t-test). Whale=bowhead whale. **d,** Apoptosis/Necrosis of human and bowhead whale fibroblasts in response to hydrogen peroxide at concentration 700µM. Two days after hydrogen peroxide, cells were harvested and subjected to an Annexin V apoptosis assay using flow cytometry. Error bars represent mean ± SD. *** $p<0.001$. Welch's t-test was used to quantify the significance (n=12). **e,** Percent tail DNA by alkaline comet assay at various time points after 700 µM H_2O_2 treatment in 2 cell lines each of human and bowhead whale fibroblasts. Points represent individual cells. Representative comet images shown below. Bars indicate mean +- SEM.

Extended Data Figure 4

Extended Data Figure 4. DSB repair efficiency in bowhead whale cells. a, NHEJ efficiency in extrachromosomal assay. NHEJ reporter construct was pre-digested with I-SceI, purified and co-transfected with DsRed into human and bowhead skin fibroblasts. Three days after transfection cells were harvested and subjected to flow cytometry to calculate NHEJ efficiency (n=3). Error bars represent mean ± SD. *** p<0.001 (two-tailed t-test) **b,** Representative images of human and bowhead whale binucleated cells containing micronuclei after 2 Gy of γ-irradiation. Scale bar indicates 20 μm. **c,** Frequency of micronuclei after DSB induction with I-SceI in primary fibroblasts carrying a chromosomally integrated NHEJ reporter cassette. Each cell line was transiently transfected with a BFP-expressing control plasmid or an I-SceI expression plasmid and micronuclei were quantified after 5d in media containing cytochalasin B to prevent cytokinesis. Micronucleus frequencies for each cell line are shown normalized to BFP control (paired t-test, n=3 cell lines/species). **d,** Pulse-field gel stained with ethidium bromide, showing chromosomal DNA fragmentation in human and bowhead confluent skin fibroblasts immediately after different doses of γ-irradiation 0.7, 1.5, 3 and 6h after 40 Gy of γ-irradiation. **e,** Kinetics of DSB repair measured by PFGE in confluent human and bowhead fibroblasts after 40 Gy of γ-irradiation. n=2 for each species.

Extended Data Figure 5

Extended Data Figure 5. Sequencing of DNA DSB repair products in bowhead whale cells. a, Possible repair outcomes after induction of DSBs with incompatible ends by I-SceI in NHEJ reporter construct. **b,** Allele plot of Sanger sequencing products resulting from repair of integrated NHEJ reporter cassette after I-SceI cleavage. **c,** NHEJ fidelity in extrachromosomal assay. NHEJ reporter construct was pre-digested with I-SceI, purified and co-transfected with DsRed into human and bowhead skin fibroblasts. Three days after transfection genomic DNA was isolated, subjected to PCR, cloned and analyzed by Sanger sequencing. At least 100 clones were analyzed for each species. Correct – annealing on 2 of the 4 protruding nucleotides **d,** Time course of CRISPR cleavage measured by digital droplet PCR (ddPCR). PTEN copy number at varying time points after CRISPR RNP transfection was measured with ddPCR using primers flanking the predicted cleavage site and normalized within each sample to a single-copy genomic ultraconserved element as described in Methods. Error bars show confidence intervals of Poisson distribution calculated in QuantaSoft. **e,** Pearson correlation between 5th percentile indel size and species lifespan (r=0.8508, 95% CI = 0.5125 to 0.9605, p=0.0009, n=11). **f,** Absolute frequencies of alleles by base pairs of microhomology across species in CRISPR-targeted PTEN repair products. **g,** Relative proportions of deletion alleles by base pairs of microhomology across species in CRISPR-targeted PTEN repair products.

Extended Data Figure 6

Extended Data Figure 6. Proteomic quantification of DNA repair proteins. a, Western blot for CIRBP on bowhead whale and mouse organs **b,** Abundance of CIRBP protein by LC-MS in liver tissue of mammal species (n=12 per species; 3 biological x 4 technical replicates; N.D.=not detected) **c,** Abundance of RPA2 protein by LC-MS in liver tissue of mammal species (n=12 per species; 3 biological x 4 technical replicates; N.D.=not detected) **d,** Abundance of CIRBP protein by LC-MS in nuclear extracts of liver tissue of mammal species (n=3 biological replicates per species) **e,** Abundance of RPA2 protein by LC-MS in nuclear extracts of liver tissue of mammal species (n=3 biological replicates per species; N.D.=not detected) **f,** Heatmap of LC-MS protein abundance for primary fibroblasts of the indicated species and proteins. Color intensity scale corresponds to $\log_{10}$ ion intensity. **g,** Per-protein normalized abundance by LC-MS of proteins identified in pulldowns of His-tagged Cas9/dCas9 bound to a plasmid containing the genomic *PTEN* target sequence after incubation in extracts of soluble nuclear proteins from human and bowhead whale.

Extended Data Figure 7

Extended Data Figure 7. Transcriptome, Western blot, and STED quantification of DNA repair proteins. a, Relative expression level of genes in 6 DNA repair pathways among species. Z-scores are scaled by row. Genes in each pathway are ordered decreasingly based on the expression level in bowhead whale. Genes with higher expression in bowhead whale compared to all 3 other species are highlighted in red text to the right of the heatmap. Genes of each gene set were compiled from 3 resources: MsigDB database, GO ontology, and a curated gene list (www.mdanderson.org/documents/Labs/Wood-Laboratory/human-dna-repair-genes.html) **b,** Western blot abundance of RPA2 in cultured skin fibroblasts, using 2 different monoclonal primary antibodies targeting conserved epitopes and normalized to histone H3. A third polyclonal antibody produced the same results but had higher background reactivity and is not shown. Each lane is a primary fibroblast line from a different adult individual. Fluorescent secondary antibodies were used to increase linear dynamic range for higher quantitative accuracy. **c,** Western blot for CIRBP with 3 different antibodies in 3 fibroblast lines per species. mAb=monoclonal antibody, pAb=polyclonal antibody. **d,** Stimulated emission depletion (STED) images of RPA2 and CIRBP localization in human and bowhead whale fibroblasts. Target protein in red, nuclear DAPI stain in blue. **e,** Western blot for CIRBP in fibroblasts isolated from various mammalian species.

Extended Data Figure 8

Extended Data Figure 8. Analysis of CIRBP's role in DNA DSB repair. a, CIRBP localization in whale cells. Before formaldehyde fixation, cells were pre-extracted with CSK buffer +/- RNAseA for 3min. After standard immunocytochemistry procedure images were collected using confocal microscope. **b,** Western blot of bowhead whale fibroblasts with knockdown of CIRBP (left panel) and band intensity quantification from 3 independent experiments (right panel) suggesting partial dependence of RPA2 protein abundance on CIRBP expression. **c,** Ion intensity by LC-MS of RPA2 in human fibroblasts with and without lentiviral overexpression of bwCIRBP (n=3 human cell lines). Error bars show mean +-SEM. **d,** DSBs induce CIRBP enrichment in chromatin. Exponentially growing cells were treated with neocarzinostatin (NCS) for the indicated period of time and lysed in CSK buffer to enrich chromatin-bound fraction. α-Tubulin staining was used to verify the absence of cytoplasmic contamination in chromatin-bound fraction. **e,** DSBs induced by γ-irradiation lead to CIRBP enrichment in chromatin. This enrichment is promoted by RNA. Exponentially growing cells were treated with γ-irradiation and at the indicated period of time were lysed in CSK buffer with/without RNAse A to enrich proteins in chromatin-bound fraction. **f,** Overexpression of CIRBP decreases the percentage of binucleated cells containing micronuclei in human cells after I-Sce1-induced DSBs. Each bar indicates an experimental replicate. At least 150 binucleated cells were scored per condition. **g,** Frequency of chromosomal aberrations in human fibroblasts with and without CIRBP overexpression after 2Gy γ-irradiation. 100 metaphases were analyzed per sample. C=control untreated cells. **h,** Frequency of insertions and deletions >20 bp in NHEJ reporter constructs PCR-amplified from human fibroblasts with and without bwCIRBP overexpression after I-SceI expression. Insertion/deletion frequencies were determined from Nanopore sequencing data of PCR products and normalized within each sample to total frequency of all insertions or deletions. **i,** Frequency of insertions and deletions as shown in (**h**) but for bowhead whale fibroblasts with negative control or CIRBP-targeting siRNAs. **j,** Calculated dissociation constants (K_D) and fluorescence polarization (FP) measurements for CIRBP proteins titrated into solutions containing a fixed concentration (3 nM) of fluorescently labeled PAR of various polymer lengths. **k,** EMSA of increasing amounts of recombinant human CIRBP incubated in vitro with 300 ng sheared chromatin from fibroblasts exposed to UVC and oxidative DNA damage as described in Methods. Chromatin was treated with Proteinase K but not RNAse. Nucleic acids are stained with SYBR Gold. Red overlay indicates saturated pixels. **l,** EMSA of 300 ng sheared purified genomic DNA, purified cellular RNA, or chromatin as described in (**k**) incubated in vitro with 5 µg rhCIRBP. **m,** Hypothermia promotes NHEJ efficiency in primary human fibroblasts (left panel). Cells were pre-incubated at 33°C for 2 days, co-transfected with I-SceI-digested NHEJ reporter and DsRed, and returned to the 33°C incubator. NHEJ efficiency was measured by flow cytometry 3 days following transfection (n=3). Western blot showing CIRBP upregulation in human cells exposed to 33°C for 2 days (right panel). Western blot images were analyzed in ImageLab software (Bio-Rad). Error bars represent mean ± SD. ** $p < 0.01$ (Welch's t-test).

a

Score	Expect	Method	Identities	Positives	Gaps
315 bits(806)	1e-116	Compositional matrix adjust.	167/172(97%)	168/172(97%)	0/172(0%)

BHW 1   MASDEGKLFVGGLSFDTNEQSLEQVFSKYGQISEVVVVKDRETQRSRGFGFVTFENIDDA   60

Human 1   MASDEGKLFVGGLSFDTNEQSLEQVFSKYGQISEVVVVKDRETQRSRGFGFVTFENIDDA   60

61   KDAMMAMNGKSVDGRQIRVDQAGKSSDNRSRGYRGGSAGGRGFFRGGRGRGRGFSRGGGD   120

     KDAMMAMNGKSVDGRQIRVDQAGKSSDNRSRGYRGGSAGGRGFFRGGRGRGRGFSRGGGD

61   KDAMMAMNGKSVDGRQIRVDQAGKSSDNRSRGYRGGSAGGRGFFRGGRGRGRGFSRGGGD   120

121   RGYGGSRFESRSGGYGGSRDYYSSRSQGGSCGDRSSGGSYRDSYDSYATHNE   172

     RGYGG+RFESRSGGYGGSRDYYSSRSQ G   DRSSGGSYRDSYDSYATHNE

121   RGYGGNRFESRSGGYGGSRDYYSSRSQSGGYSDRSSGGSYRDSYDSYATHNE   172

BHW variant   BHW variant

b

   0.020   0.040   0.060   0.080   0.10   0.12   0.14

LAGENORHYNCHUS_OBLIQUIDENS_CIRBP *

LIPOTES_VEXILLIFER_CIRBP *

ORCINUS_ORCA_CIRBP *

MONODON_MONOCEROS_CIRBP *

DELPHINAPTERUS_LEUCAS_CIRBP *

BALAENA_MYSTICETUS_CIRBP * *

BALAENOPTERA_ACUTOROSTRATA_CIRBP * *

BALAENOPTERA_MUSCULUS_CIRBP * *

BOS_MUTUS_CIRBP *

BOS_INDICUS_CIRBP *

BOS_TAURUS_CIRBP *

MYOTIS_MYOTIS_CIRBP *

MYOTIS_DAVIDII_CIRBP *

MYOTIS_BRANDTII_CIRBP

MYOTIS_LUCIFUGUS_CIRBP *

HYLOBATES_MOLOCH_CIRBP

NOMASCUS_LEUCOGENYS_CIRBP

PONGO_PYGMAEUS_CIRBP

PAN_PANISCUS_CIRBP

HOMO_SAPIENS_CIRBP

GORILLA_GORILLA_GORILLA_CIRBP

HETEROCEPHALUS_GLABER_CIRBP

PHODOPUS_ROBOROVSKII_CIRBP

RATTUS_NORVEGICUS_CIRBP

MASTOMYS_COUCHA_CIRBP

MUS_MUSCULUS_CIRBP

MUS_PAHARI_CIRBP

* * BHW variant

c

N126   S152   Y151   G150   S148

S126   G152   C151   S150   G148

Human CIRBP   Bowhead whale CIRBP

d

Control   hCIRBP   H5BW   bwCIRBP   BW5H

CIRBP

β-Actin

e

CDS variant	CAI
hCIRBP	0.811
bwCIRBP	0.845
H5BW	0.809
BW5H	0.848

Extended Data Figure 9

Extended Data Figure 9. Analysis of bwCIRBP coding sequence mutations and protein expression levels. a, Comparison of amino acid sequences between human and bowhead whale CIRBP through BLAST analysis. **b,** Phylogenetic tree illustrating the relationships among CIRBP coding sequences from representative species with genome sequence information available. The asterisk indicates the presence of BHW-specific variants in the species. The colors indicate the position of variants shown in (**a**). **c,** SwissModel/AlphaFold models of human (left, pink) and bowhead whale (right, blue). Side chains of whale residues that diverge from human are shown, and their ribbon is colored yellow in the model. The key takeaway is that all the residues that differ between whale and human are in the C-terminal disordered region, whereas the N-terminal RNA recognition motif (RRM) is structured and conserved. **d,** Western blot abundance of bwCIRBP, hCIRBP, and reciprocal amino acid mutants overexpressed in human cells. **e,** Calculated codon adaptation index (CAI) for CIRBP coding sequence variants.

a

b

c

Bowhead whale RPA2

Human RPA2

Extended Data Figure 10

Extended Data Figure 10. Analysis of bowhead whale RPA2 sequence. a,
Comparison of amino acid sequences between human and bowhead whale RPA2
through BLAST analysis. **b,** Phylogenetic tree illustrating the relationships among RPA2
coding sequence from different representative species. The asterisk indicates the
presence of BHW-specific variants in the species. The colors indicate the position of
variants shown in (**a**). **c,** AlphaFold protein structures of human and bowhead whale
RPA2 showing the position of the variants.

Extended Data Figure 11

Extended Data Figure 11. Bowhead whale CIRBP reduces anchorage-independent cell growth. a, Images of representative human transformed fibroblast colonies with and without bwCIRBP overexpression after 23 days of growth in soft agar. 20x magnification. Bar 100μm. **b,** Quantification of colonies after staining with nitro blue tetrazolium chloride. Colonies were counted using ImageJ software as described in Methods. Error bars represent SD. *p<0.05 (Welch's t-test). **c,** Cell proliferation MTT assay. **d,** Trypan Blue exclusion test of cell viability. **e,** Western blot showing expression of LT, Ras, p16 and p21 after overexpression of bwCIRBP. **f,** Frequency of chromosomal aberrations in human transformed cells after bwCIRBP overexpression. 100 metaphases were analyzed per sample.