HISTONE DEACETYLASE 19 REGULATES *SHOOT MERISTEMLESS* EXPRESSION IN THE CARPEL MARGIN MERISTEM CONTRIBUTING TO OVULE NUMBER DETERMINATION AND TRANSMITTING TRACT DIFFERENTIATION

Manrique S[1,8], Cavalleri A[1,8], Guazzotti A[1,8], Villarino GH[2], Simonini S[3], Bombarely A[1], Higashiyama T[4], Grossniklaus U[3], Mizzotti C[1], Pereira AM[5,6], Coimbra S[5,6], Sankaranarayanan S[7], Onelli E[1], Masiero S[1], Franks RG[2], Colombo L[1‡]

[1] Dipartimento di Bioscienze, Università degli Studi di Milano, Via Giovanni Celoria 26, 20133, Milan, Italy

[2] North Carolina State University, Department of Plant and Microbial Biology, Raleigh, NC, 27606, USA

[3] Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, CH-8008, Zurich, Switzerland

[4] Institute of Transformative Bio-Molecules (ITbM), Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8601, Japan

[5] Faculdade de Ciências da Universidade do Porto, Departamento de Biologia, Universidade do Porto, rua do Campo Alegre, 4169-007 Porto, Portugal

[6] LAQV Requimte, Sustainable Chemistry, Universidade do Porto, 4169-007 Porto, Portugal

[7] Department of Biological Sciences and Engineering, Indian Institute of Technology Gandhinagar, Palaj, Gujarat 382355, India.

[8] Co-first author

‡Correspondence to: lucia.colombo@unimi.it

Running title

HDA19 controls CMM activity by repressing *STM*

Abbreviations

CMM-Carpel margin meristem

TT – transmitting tract

OP – ovule primordia

ReT – reproductive tract

ISH – *in situ* hybridization

ChIP - chromatin immunoprecipitation

Abstract

The gynoecium is critical for the reproduction of flowering species as it contains

the ovules and the tissues required for pollen germination and guidance. These

tissues are collectively known as the reproductive tract (ReT) and comprise

stigma, style and transmitting tract (TT). The ovules and the ReT originate from

a meristem within the pistil named carpel margin meristem (CMM).

SHOOT MERISTEMLESS (STM) is a key transcription factor required for

meristem formation and maintenance. In all above-ground meristems, including

the CMM, *STM* has to be locally downregulated to allow proper organ

differentiation. However, how this downregulation is achieved in the CMM is

unknown.

In this work, we have studied HISTONE DEACETYLASE 19 (HDA19) role in

ovule and ReT differentiation, based on the observation that *hda19-3* mutant

displays reduced ovule number and fails to properly differentiate the TT.

Fluorescence activated cell sorting (FACS) coupled with RNA-seq revealed that

in the CMM of *hda19-3* mutant, genes promoting organ development are

downregulated while meristematic markers, including *STM*, are upregulated. We

found that HDA19 is fundamental to downregulate *STM* in the CMM, allowing

ovule formation and TT differentiation. *STM* is ectopically expressed in *hda19-3*

at intermediate stages of pistil development, and its downregulation by RNA

interference alleviated *hda19*-3 phenotypic defects. Furthermore, chromatin

immunoprecipitation assays indicated that *STM* is a direct target of HDA19 during

pistil development and that *SEEDSTICK* (*STK)* is required for the histone

acetylation-mediated regulation of *STM*.

Our results have led to the identification of the factors required for *STM* silencing

in the gynoecium allowing organogenesis and tissue differentiation from the

CMM.

Introduction

The number of seeds produced by a plant is important for the reproductive success of species and for food security. To increase seed number, plants can produce more flowers and/or more seeds per flower. As seeds are formed upon ovule fertilization, final seed number depends partially on ovule number.

The ovules and the reproductive tract (ReT) emerge from the carpel margin meristem (CMM) (Reyes-Olalde et al., 2013a; Chávez Montes et al., 2015). The CMM is a determinate meristem that appears as a ridge in the medial part of the gynoecium at stage 7 of floral development (Figure 1A and Smyth et al., 1990). At stage 8, it produces the placenta, from which ovule primordia (OP) arise at stage 9. Contemporarily to OP formation, the medial ridges fuse forming the septum (Figure 1A). From stages 10 to 12, ovules, stigma, style and transmitting tract (TT) complete their developmental program, and the gynoecium becomes competent for fertilization (Bowman et al., 1999; Roeder and Yanofsky, 2006; Reyes-Olalde et al., 2013b; Chávez Montes et al., 2015; Herrera-Ubaldo et al., 2019).

Ovule formation encompasses two steps: (1) boundary formation and primordium outgrowth and (2) identity determination. The first step is controlled by a network of genes that modulate lateral organ formation such as CUP-SHAPED COTYLEDON 1/2/3 (CUC1/2/3), BLADE-ON-PETIOLE (BOP), PIN-FORMED 1 (PIN1), and AINTEGUMENTA (ANT) (Ishida et al., 2000; Mizukami and Fischer, 2000; Vernoux et al., 2000; Norberg et al., 2005). By contrast, the second step is controlled by organ-specific gene networks. Ovule identity is redundantly determined by three MADS-box transcription factors: SEEDSTICK (STK) and SHATTERPROOF1/2 (SHP1/2). In the triple mutant *stk shp1 shp2*, ovules are replaced by carpelloid structures (Favaro et al., 2003; Pinyopich et al., 2003).

As for ReT tissues, the TT supports the growth of pollen tubes to reach the ovules. It forms in the center of the septum and runs from the top to the base of the pistil (Figure 1A) (Crawford and Yanofsky, 2008). In *Arabidopsis*, pollen tubes grow through the intercellular spaces between TT cells, and their growth is aided by the production of a facilitating extra-cellular matrix, and by the death of the TT cells. This process is developmentally regulated and starts before anthesis (Crawford and Yanofsky, 2008; Pereira et al., 2021). The formation of the TT initially depends on transcription factors controlling the formation of the septum

like ETTIN (ETT), SPATULA (SPT), SEUSS (SEU), ANT or INDEHISCENT (IND)

(Alvarez & Smyth, 2002; Azhakanandam et al., 2008; Kay et al., 2013; Pereira

et al., 2021; Sessions & Zambryski, 1995). Later, genes like *NO TRANSMITTING*

TRACT (*NTT*) and *HECATE1/2/3* (*HEC1/2/3*) control the maturation of TT cells

(Gremski et al., 2007, Herrera-Ubaldo et al., 2019; Marsch-Martínez et al., 2014).

Interestingly, STK and SHP1/2 are also required for TT formation in cooperation

with NTT (Marsch-Martínez et al., 2014; Herrera-Ubaldo et al., 2019) and

CESTA/HALF-FILLED (CES) (Di Marzo et al., 2020).

The CMM is a determinate meristem, so that stem cells are maintained

transiently, and consumed once ovule and tissue differentiation are completed

(Alvarez & Smyth, 2002; Reyes-Olalde & de Folter, 2019). Nevertheless, genetic

and transcriptomic analyses have shown its similarity with undetermined

meristems like the shoot apical meristem (SAM) (Wynn et al., 2011; Heidstra and

Sabatini, 2014; Gaillochet and Lohmann, 2015; Villarino et al., 2016; Reyes-

Olalde and de Folter, 2019; Zúñiga-Mayo et al., 2019).

HISTONE DEACETYLASE 19 (HDA19) is a major regulator of the maintenance

of the SAM (Long et al., 2006; Gorham et al., 2018; Chung et al., 2019a), the

flower meristem (FM) (Krogan et al., 2012a; Bollier et al., 2018) and the root

apical meristem (RAM) (Pi et al., 2015a). HDA19 is an epigenetic modifier that

removes acetyl groups from histones, repressing the transcription of the genes

associated with such histones (Kumar et al., 2021). HDA19 target specificity is

provided by the interaction with transcription factors and other cofactors (Kumar

et al., 2021). For instance, in the inflorescence meristem (IM), a complex between

HDA19 and the auxin response factors ETT, ARF4 and the YABBY

FILAMENTOUS FLOWERS (FIL) downregulates the *KNOTTED-1-like*

homeobox (*KNOX*) transcription factor *SHOOT MERISTEMLESS (STM*),

allowing flower primordia specification (Chung et al., 2019). Interestingly, it has

been previously suggested that STM role is similar in both the SAM and the CMM

(Scofield et al., 2007; Durbak and Tax, 2011; Poulios and Vlachonasios, 2018)

as it is required to maintain meristematic identity in both contexts. *STM* has to be

locally downregulated to allow the formation of organ primordia from the SAM

(Long et al., 1996; Kanrar et al., 2006; Landrein et al., 2015). Likewise, in pistils,

its overexpression interferes with ovule identity determination and septum

formation (Scofield et al., 2007).

Here, we show that mutation of *HDA19* leads to ovule number reduction and TT

defects. We identified targets of HDA19 during TT differentiation and ovule

formation, through transcriptomic analysis of cells isolated by fluorescence

activated cell sorting (FACS), using a *STK* marker line active in the TT, placenta,

and ovules. We observed by RNA-seq and in situ hybridization (ISH) that *STM*

was over- and ectopically expressed in TT and ovules in *hda19-3* in respect to

wild-type. Furthermore, chromatin immunoprecipitation (ChIP) experiments

showed that *STM* locus was more acetylated in *hda19-3* flowers. Our work shows

also that *stk* mutant has similar defects to *hda19-3* in ovule number and TT

differentiation and that STK binds *STM* locus and regulates its acetylation state.

Overall, our results indicate that correct development of ovules and TT requires

the repression of *STM* by HDA19 and that STK is also necessary for histone

acetylation-mediated downregulation of *STM* in the CMM.

RESULTS

Development of the reproductive tract requires HDA19

It was previously observed that *hda19* mutants have a reduced seed set (Zhou

et al., 2013), a phenotype that can be caused by pre- (Kuhn et al., 2020) and/or

post-fertilization defects (Zhou et al., 2013). Despite these descriptions, the

molecular basis of *hda19* reduced seed production has not been investigated.

Therefore, in this work we have focused on the involvement of HDA19 on pre-

fertilization stages and specially in its role during ovule number determination and

TT differentiation.

We performed a phenotypic characterization on *hda19-3* mutant pistils

considering traits as ovule density and reproductive tract morphology. At first

instance, we analysed ovule number in *wild-type* and *had19-3*. We observed that,

even though *hda19-3* pistils have similar size respect to *wild-type* (Figure 1B, C),

they contain significantly fewer ovules (51±3) compared to *wild-type* (67±4

ovules) (Figure 1D), suggesting that ovule density is lower. Next, we analysed

pistils radial organization observing that, in *hda19-3,* the medial ridges remain

unfused at stage 12 (Figure 1E-H). Consequently, the TT is not properly formed

and the degeneration of the subepidermal cells required for TT maturation is also

drastically reduced in *hda19-3* (Figure 1F, H) respect to the *wild-type* (Figure 1E,

G). These defects impair pollen tube growth along the pistil (Figure 1I,
Supplemental Figure 1), and reduce fertility predominantly at the bottom-half of
the silique (Figure 1J, K). In hand-pollination experiments using *wild-type* pollen,
the percentage of unfertilized ovules in *wild-type* siliques at 16 hours after
pollination was low and similar between the top- and bottom -half of the siliques
(Figure 1J). In *hda19-3*, fertility is overall reduced in respect to *wild-type* (Figure
1J, K). In addition, the percentage of unfertilized ovules at the bottom-half of the
siliques was significatively higher respect to the top-half (Figure 1K). Altogether,
we concluded that HDA19 is required for proper ovule primordia specification and
TT development.

Figure 1. Ovule density and transmitting tract are abnormal in *hda19-3*. A) Scheme of the development of the reproductive tract from the CMM from stage 8 to 12 of flower development according to Smyth et al., 1990. Abbreviations: CMM=carpel margin meristem; OP=ovule primordia; TT= transmitting tract. **B-C)** Cleared pistils of *wild-type* (**B**) and *hda19-3* (**C**) at stage 9, showing differences

in ovule density. Scale bar = 50 μm. **D)** Ovule number of *wild-type* (n = 70 pistils)

and *hda19-3* (n= 85 pistils) plants. Bars represent the mean ± s.e. value. *** =

Student's T-test with p-value < 0.001. **E-H)** Transversal section of *wild-type* (**E**,

G) and *hda19-3* (**F**, **H**) pistils at stage 12. *hda19-3* shows unfused septum and

defective transmitting tract due to the lack of degeneration of septum cells.

Images **G** and **H** are close up images of a portion of images **E** and **F** respectively.

Abbreviations: TT = transmitting tract; OV = ovule; mr = medial ridge. Scale bar

= 50 μm. **I)** Ratio of pollen tube length/pistil length at 12 and 24 HAP in *wild-type*

(n= 8), heterozygous (n= 8) and homozygous (n= 12) *hda19-3* pistils, hand-

pollinated with wild-type pollen. Bars represent the mean ± s.e. value. *** =

Student's T-test with p-value < 0.001. **J-K)** Seed set at the top-half and bottom-

half of *wild-type* (n = 10) (**J**) and *hda19-3* (n = 10) (**K**) siliques, hand-pollinated

with *wild-type* pollen, showing that the bottom part of *hda19-3* siliques shows

higher levels of unfertilized ovules than the top. Bars represent the mean ± s.e.

value.

Transcriptomic analysis of *hda19-3* sorted cells supports a role of HDA19

in reproductive tract development and ovule number determination

As *HDA19* is highly expressed in most tissues of the inflorescence (Zhou et al.,

2005) and we wanted to focus on its role during ovule number determination and

TT differentiation, we designed a strategy based on the isolation of these specific

cell types to perform transcriptomic analysis.

We crossed *hda19-3* with a line containing the genomic region of *STK* fused to

GFP (*pSTK::STK-GFP,* Mizzotti et al., 2014), as *STK* is expressed in the placenta,

ovules and TT from stage 8 to 12 (Mizzotti et al., 2014; Herrera-Ubaldo et al., 2019).

After verifying that the spatial expression pattern of *pSTK::STK-GFP* was

equivalent in both genetic backgrounds (Supplemental Figure 2), fluorescent

protoplasts were isolated by FACS and RNA was extracted for RNA-sequencing

to identify differentially expressed genes.

Before proceeding with library preparation, enrichment of *STK-GFP* in sorted

protoplasts was validated by qRT-PCR (Supplemental Figure 3). After the

validation, three samples of RNA from *wild-type* GFP-positive cells, and four

samples from *hda19-3* GFP-positive cells (80-100.000 cells/replicate) were used
for library preparation and RNA Illumina sequencing (HiSeq2500).
Since HDA19 is an epigenetic regulator, we used a stringent approach to narrow
down the number of differentially expressed genes (DEGs). We used Cufflinks
(Trapnell et al., 2012), DeSeq2 (Love et al., 2014) and EdgeR (Robinson et al.,
2009) to analyze the RNA-seq data (Supplemental File 1-3). Then, we compared
the DEGs obtained by the three software and selected only those in common
among the three programs (Supplemental File 4). This approach retrieved 554
upregulated genes and 408 downregulated genes (Figure 2A).
To survey the roles of the DEGs, we performed an enrichment analysis of
'Biological Process' GO terms associated with the upregulated and
downregulated genes (Figure 2 B, C and Supplemental File 5, 6) using Panther
(Mi et al., 2018, 2019). Among the upregulated genes we found GO terms as
'response to salicylic acid', 'defense response' or 'aging' that directly reflect
previously described roles of HDA19 (Zhou et al., 2005; Kim et al., 2008; Tanaka
et al., 2008; Zhu et al., 2010; Choi et al., 2012) (Figure 2B, Supplemental File 5).
Likewise, 'NAD(P)H dehydrogenase complex assembly' relates to the role of
HDA19 in light response regulation (Benhamed et al., 2006; Guo et al., 2008; Jan
et al., 2011).
Among the downregulated genes, the most enriched GO terms (Figure 2C) were
'transmitting tissue development' and 'carpel formation', both with a fold
enrichment of 51.28 (Figure 2C, Supplemental File 6).
Overall, GO terms enrichment analysis showed that genes associated with
known roles of HDA19 were present among the DEGs, supporting the validity of
our dataset and of the sampling strategy, and revealed that genes necessary for
the normal progression of carpel and TT development were downregulated in
hda19-3 sorted cells.
242

Figure 2. Reproductive tract transcriptomic comparison of *wild-type* and *hda19-3*. A) Venn diagrams showing the overlap of up- and down-regulated genes (FDR<0.001 and logFC>|1.5|) detected by Cufflinks, EdgeR and DeSeq2.

B) Enrichment analysis of 'Biological Process' GO terms associated with up-
regulated genes (logFC>|1.5|) detected by all three programs. **C)** Enrichment
analysis of 'Biological Process' GO terms associated with down-regulated genes
(logFC>|1.5|) detected by all three programs (fold enrichment > 4). Graphs show
overrepresented child-most (tip) GO terms. Parent GO terms (root terms) have
not been plotted for simplicity. Enrichment analysis was performed using Panther
(Mi et al., 2018, 2019) with the default parameters set by the program.
254
255

Meristematic markers and genes involved in ovule and transmitting tract
development are deregulated in *hda19-3* mutant
HDA19 regulates the maintenance of the stem cell population in several
meristems (Long et al., 2006; Krogan et al., 2012b; Pi et al., 2015b; Bollier et al.,
2018). Therefore, we checked for deregulated meristematic markers in our
dataset, and we found that *CLAVATA1* (*CLV1*) as well as the KNOX genes *STM*,
KNAT1/BREVIPEDICELLUS (*BP*) and *KNAT 6* showed higher expression in
hda19-3 in respect to *wild-type* (Figure 3A).
STM, *BP* and *KNAT6* are partially redundant for SAM maintenance (Byrne et al.,
2002), and their local downregulation at the boundaries of the meristem is
necessary for organ initiation (Belles-Boix et al., 2006; Ragni et al., 2008; Zhao
et al., 2015). Therefore, their upregulation in *hda19-3* could be hindering the
formation of CMM-derived structures.
Furthermore, to identify CMM-specific genes associated to ReT differentiation
and ovule initiation, we cross-referenced our datasets with data from two articles
that described sets of CMM-specific genes (Wynn et al. 2011; Villarino et al.,
2016). Regarding Villarino´s set (stages 5-8), we found that several genes
involved in TT and style development like *HEC1*, *HEC2*, *SPT* (Gremski et al.,
2007; Nahar et al., 2012) and *STYLISH* (*STY1*) (Sohlberg et al., 2006; Kuhn et
al., 2020) were downregulated in *hda19-3* (Figure 3B). This might explain the
defects in the TT (Figure 1E-H) and the style (Kuhn et al., 2020) of *hda19-3.*
Additionally, several well-known players of ovule initiation from Wynn's dataset
(stages 8-10) (Wynn et al., 2011a), like *PHABULOSA* (*PHB*), *ANT*, *PERIANTHIA*
(*PAN*) and *PIN1* (Azhakanandam et al., 2008; Cucinotta et al., 2014; Galbiati et
al., 2013; Lee & Clark, 2015; Nole-Wilson, 2006) were downregulated in *hda19-*

3 (Figure 3C). This observation is consistent with the reduced number of ovules

observed in *hda19-3* (Figure 1D)*.*

Only four CMM-specific genes were upregulated in *hda19-3*: *NGATHA4* (*NGA4*)

and *MYB55* (Villarino et al., 2016) (Figure 3B); and *YABBY3* (*YAB3*) and

INDOLE-3-ACETIC ACID INDUCIBLE 18 (IAA18) (Wynn et al., 2011) (Figure

3C). *NGA4* and *YAB3* are negative regulators of lateral organ formation (Lee et

al., 2015, Alvarez et al., 2009, González-Reig et al., 2012), agreeing with *hda19-*

3's ReT phenotypes. The role of *IAA18* and *MYB55* in the pistil is not known, but

iaa18 mutants have reduced fertility (Uehara et al., 2008), and *MYB55* is a direct

target of BRASSINAZOLE-RESISTANT 1 (BZR1) (He, 2005), which participates

in organ boundary establishment in the SAM (Bell et al., 2012; Gendron et al.,

2012). Therefore, upregulation of these genes might also interfere with ReT

development.

Finally, we checked the expression of some additional regulators of TT and ovule

number determination that were not included in Villarino's or Wynn's datasets

due to the experimental conditions chosen in their work. For example, *HEC3* and

STK itself were downregulated (Figure 3D). It is worth noting that, although *STK*

shows a logFC close to -4 in *hda19-3*, STK spatial pattern in *hda19-3* was

unchanged in respect to *wild-type* (Supplemental Figure 2).

Overall, analysis of CMM-specific genes revealed that several *KNOX* genes

involved in meristem maintenance (*STM*, *BP*, *KNAT6*) and negative regulators of

lateral organ development (*NGA4* and *YAB3*) were upregulated in *hda19-3* pistils,

while positive regulators of TT differentiation and ovule initiation were

downregulated (e.g., *PIN 1* and *ANT*). As HDA19 is a transcriptional repressor,

this suggests that HDA19 might directly regulate meristematic genes and/or

genes involved in repressing the proper differentiation and formation of CMM-

derived structures.

Figure 3. CMM-specific genes are deregulated in *hda19-3* mutant. A) Meristem marker genes deregulated in *hda19-3* mutant. **B)** CMM-specific genes from stages 6-8 of pistil development according to Villarino *et al.* (2016) that are deregulated in *hda19-3*. **C)** CMM-specific genes from stages 8-10 of pistil development according to Wynn *et al.* (2011) that are deregulated in *hda19-3*. **D)** Notable ovule number and TT regulators are deregulated in *hda19-3*. Genes plotted correspond to genes with an FDR < 0.001 and logFC > |1.2| in the RNA-seq data according to Cufflinks (C), EdgeR (E) and DeSeq2 (D). Genes were hierarchically clustered by average clustering using Euclidean distances. Heatmaps were obtained using Heatmapper (http://heatmapper.ca/). The color scale represents the logFC (*wild-type* vs *hda19-3*) as detected by each program and the color is relative to the maximum and minimum values of each table.

STM is ectopically expressed and over-acetylated in the reproductive tract of *hda19-3* mutant

Based on its role in other meristems and on our data, we hypothesized that *STM* could be one of the main end-goal of HDA19's repressing activity in the CMM. To gain insight on the spatial expression pattern of *STM* in *wild-type* and *hda19-3* pistils, we performed *in-situ* hybridization (ISH) with a *STM*-specific antisense probe (Figure 4A-H). At stage 7, the expression pattern of *STM* was similar in both genotypes (Figure 4A, E). However, from stages 8 to 10, the expression of *STM* in the medial ridge increased in *hda19-3* and expanded to the placenta, OP and even to the lateral domain (Figure 4B-C, F-G). After stage 10, *STM* expression became restricted to the adaxial replum zone in both genotypes (Figure 4D, H). As STM promotes meristematic fate, its ectopic expression in the placenta, OP and TT from stage 8-10 might interfere with ovule initiation and TT development, causing the phenotypes observed in *hda19-3* mutant.

To analyze whether the changes in *STM* expression pattern could be caused by variations in its histone acetylation levels, resulting from the absence of HDA19, we collected flowers at stages 1-12 from *wild-type* and *hda19-3* plants and performed ChIP followed by quantitative Real-Time PCR (ChIP-qPCR). We used an antibody against HISTONE 3 (H3) acetylated in Lysine 9 and 14 (K9/14) (AcH3K9/14) (Figure 4I-J). Since no other histone de/acetylases are significantly

deregulated in *hda19-3* sorted cells (Supplemental Figure 4), the changes in *STM*

histones acetylation observed with this antibody should be caused mostly by the

direct action of HDA19.

We tested the acetylation levels in several regions of *STM* locus: (1) the distal

promoter (-2 kb) (Figure 4I), because it was reported that HDA19 can deacetylate

STM in these regions (Chung et al., 2019a); (2) the transcription start-site (TSS),

as it generally correlates with active gene expression (Wang et al., 2017; Kim et

al., 2021; Kumar et al., 2021); and (3) the first intron. We observed a significant

increase in acetylation at the *STM* TSS in *hda19-3*, and a moderate increase at

the promoter region, while no significative increment in acetylation was found in

the first intron (Figure 4J). In conclusion, our results suggest that at the stages

tested, HDA19 is required for the deacetylation of *STM*.

Figure 4. *STM* is ectopically expressed and over-acetylated in *hda19-3*

pistils. A-H) Detection of *STM* mRNA by *in-situ* hybridization in *wild-type* **(A-D)**

and *hda19-3* **(E-H)** pistils. At stage 6, *STM* mRNA is specifically expressed in the

CMM of *wild-type* pistils (**A-B**). Then expression decreases along following
stages until getting restricted to the adaxial replum zone (**C-D**). Instead, in *hda19-*
3 during stages 7-10, when ovule primordia arise, the expression of *STM*
increases and expands outside of the CMM (**F-G**), to finally become also
restricted to the replum later on (**H**). Abbreviations: CMM = carpel margin
meristem; LD= lateral domain; OP = ovule primordia; AdR = adaxial replum.
Scale bar = 50 μm. **I)** Regions of *STM* locus tested for changes in AcH3K9/14. **J)**
AcH3K9 ChIP-qPCR of *STM* locus in *wild-type* and *hda19-3*. Graph shows the
mean ± sd result of *STM* regions enrichment in *hda19-3* background in respect
to the *wild-type*, among three independent biological replicates of the AcH3K9
ChIP-qPCR experiment performed on *wild-type*, *hda19-3, stk* and *stkshp1shp2*
flowers. ** = Student's T-test with p-value < 0. 01.

Downregulation of *STM* alleviates reproductive tract defects of *hda19-3*
mutant in an expression-level dependent manner
To verify if the ectopic expression of *STM* was involved in the reduction of ovule
number and in the TT defects of *hda19-3*, we used an RNA interference (*RNAi*)
construct to reduce *STM* levels in *hda19-3* mutant background. (Fornara et al.,
2004).
We analyzed eight *hda19-3* independent lines transformed with *35S::STM-*
RNAi::pFRH (Figure 5, Supplemental Figure 5). We measured *STM* mRNA
levels in the inflorescences of *wild-type*, *hda19-3* and *35S::STM-RNAi::pFRH*
(*hda19-3 STMRNAi* hereafter) transformed plants by qRT-PCR. The levels of
STM transcript in *hda19-3* mutant were 2-fold those of *wild-type* (Figure 5A). All
the analyzed *hda19-3 STMRNAi* transformants showed levels of *STM* mRNA in
between those of *hda19-3* and the *wild-type* (Figure 5A).
Ovule number and density increased in the *hda19-3 STMRNAi* plants according
to the level of *STM* downregulation (Figure 5B), suggesting that *STM* expression
level in the gynoecium was related to proper ovule number determination. Lines
4 and 9 showed the mildest reduction of *STM* levels (ca. 75% of *hda19-3* levels)
(Figure 5B), and in accordance did not show significant changes in ovule number.
Line 5 was the only exception, as it showed a reduction in *STM* levels similar to
line 1 (ca. 35% reduction) (Figure 5A), but while Line 1 showed an increase in

ovule number, Line 5 did not (Figure 5B). As expected, placenta length remained unchanged in all inspected lines (Figure 5C). Therefore, the increase in ovule number represents an increase in ovule density (Figure 5B-C).

Regarding the TT, we selected lines 1 and 10 for the inspection of TT phenotypes as they were comparable in terms of *STM* expression levels and ovule density. Although the septum of Line 1 and 10 was still partially unfused (Supplemental Figure 5A-C), the ridges were contacting more closely (Supplemental Figure 5B, C) and the subepidermal cells exhibited a *wild-type* like phenotype (Supplemental Figure 5A, C). In accordance, pollen tube growth-rate improved in the transgenic lines respect to *hda19-3* (Figure 5D, Supplemental Figure 5D-F)*.* We hand-pollinated *wild-type*, *hda19-3* and *hda19-3 STMRNAi* pistils (lines 1 and 10) with *wild-type* pollen and 14 hours after pollination (14 HAP) we stained them with aniline blue (Figure 5D, Supplemental Figure 5D-F). We determined in a preliminary experiment that in *wild-type* pistils pollen tubes reach the bottom of the pistil in 16 hours. Therefore, at 14 HAP we would be able to precisely quantify the differences between each genotype. At 14 HAP, pollen tubes grew through 70% of *wild-type* pistils and 60% of *hda19-3 STMRNAi* pistils, a significant increase in respect to *hda19-3*, where pollen tubes barely grew 40% of the length of the pistil (Figure 5D, Supplemental Figure 5D-F). Overall, even if the anatomical changes in the TT are mild, they were sufficient to improve pollen tube growth in respect to *hda19-3* mutant.

In conclusion, our results strongly suggest that *STM* ectopic expression (Figure 4A-H) is mostly responsible for the ReT phenotypes present in *hda19*-3 (Figure 1) as reducing *STM* expression in *hda19-3* to *wild-type* levels leads to a recovery of ovule number and density (Figure 5B, C) and of TT function (Figure 5D, Supplemental Figure 5D-F).

Figure 5. Downregulation of *STM* alleviates *hda19-3* reproductive tract phenotypes. A) Level of *STM* mRNA in inflorescences of *wild-type*, *hda19-3* and 8 independent *hda19-3 STMRNAi* lines. **B)** Ovule number per pistil of *wild-type* (n=25), *hda19-3* (n=27) and *hda19-3 STMRNAi* lines (in order from line 1 to line 10: n=8, 4, 10, 4, 6, 7, 6, 6). The statistical significance of ovule number was determined by ANOVA and a Bonferroni *post-hoc* test for multiple comparisons. Letters over boxes denote significantly different averages and asterisks the level

of signification (* = p-value < 0.05; ** = p-value < 0.01; *** = p-value < 0.001). **C)** Placenta length of *wild-type*, *hda19-3* and the *hda19-3 STMRNAi* lines that showed significant changes in ovule number (Lines 1, 3,6, 8, 10). **D)** Pollen tube growth measured as pollen tube length /total length of the pistil. Pollen tubes were allowed to grow for 14h after manual pollination (14 HAP) of emasculated flowers. Growth was visualized by aniline blue staining and length of pollen tubes and pistils were measured using ImageJ. *Wild-type* (n=17 pistils); *hda19-3* (n = 17 pistils); *hda19-3 STMRNAi* (pool of lines 1 and 10) (n=18 pistils). Letters over box-plots indicate statistical differences as determined by a one-way ANOVA test (p<0.05) and a Bonferroni *post hoc* test for multiple comparisons.

stk shows defects in the reproductive tract and increased histone acetylation at *STM* locus

HDA19 does not bind DNA directly, therefore, it has to be recruited to the DNA by transcription factors and other cofactors to regulate its targets. In the IM, HDA19 interacts with ETT, ARF4 and FIL (Chung et al., 2019b) to downregulate *STM*. However, in the CMM, the action of HDA19 is probably mediated by a different complex as at stages 8-9, ETT, ARF4 and FIL are mainly expressed in the lateral domains of the gynoecium (Sessions et al., 1997; Pekker et al., 2005; González-Reig et al., 2012).

Among the transcriptional factors expressed contextually to *HDA19* in the CMM, we highlighted STK as a possible candidate to partner with HDA19 to regulate *STM* expression. STK acts in redundancy with SHP1/2 to determine the ovule identity. Beyond this, STK is also required for septum and TT development (Colombo et al., 2010; Herrera-Ubaldo, et al., 2019; DiMarzo, et al., 2020), and displays repressor activity associated with histone deacetylation of its targets (Mizzotti et al 2014). To verify the role of STK, and potentially of SHP1/2, in the regulation of the expression of *STM* in the CMM, we analysed *STM* expression between *wild-type* and *stk shp1 shp2* mutant pistils by ISH (Figure 6A-H). *STM* resulted ectopically and overexpressed in *stk shp1 shp2* (Figure 6E-H), similarly to what we previously observed in *hda19-3* (Figure 4A-H). This observation suggests that STK and/or SHP1/2 could be involved in *STM* regulation during ReT development. Subsequentially, we measured ovule number and placenta length. We observed that *stk* mutant showed lower ovule number (Figure 6I)

without changes in placenta length (Figure 6J), similarly to *hda19-3* (Fig. 1).

Nevertheless, no statistically significative additive effects were noted in the *stk*

shp1 shp2 triple mutant (Fig. 6I, J), suggesting that SHP1 and SHP2 are not likely

be involved in ovule number determination.

Finally, to corroborate our hypothesis that STK and HDA19 collaborate to

downregulated *STM* to drive ReT differentiation, we evaluated the TT morphology

in *stk* pistils. As expected, we found that subepidermal cells of *stk* 's medial ridges

appear like those of *hda19-3* (Figure 6K-N).

Figure 6. STK regulates *STM* expression in the CMM. A-H) Detection of *STM*
mRNA in *wild-type* (**A-D**) and *stk shp1 shp2* mutant (**E-H**) by *in situ* hybridization.
STM is over and ectopically expressed in *stk shp1 shp2* triple mutant at stages
7-10 of pistil development (**F-G**) in respect to the *wild-type* situation (**B-C**).
Abbreviations: CMM = carpel margin meristem; LD= lateral domain; OP = ovule
primordia; AdR = adaxial replum. Scale bar = 50 µm. **I-J)** Ovule number (**I**) and
placenta length (**J**) in *wild-type, stk* and *stk shp1 shp2* mutants. Letters over
violin-plots indicate statistical differences as determined by a one-way ANOVA
test (p<0.05) and a Bonferroni *post-hoc* test. **K-N)** Transversal sections of stage
10-11 pistils of *wild-type* (**K**), *hda19-3* (**M**) and *stk* (**N**). *hda19-3* and *stk* show
defective subepidermal cells in the TT. Abbreviations: TT = transmitting tract; OV
= ovule; mr = medial ridge. Scale bar = 50 µm.

STK binds directly *STM* locus and is involved in its deacetylation
To explore whether STK activity was required to regulate *STM* acetylation state,
we performed ChIP-qPCR using an anti AcH3K9/14 antibody on flowers from
wild-type, *hda19-3, stk* and *stk shp1 shp2* (Figure 7A, B), testing the acetylation
pattern of the regions described previously (Figure 4I). Similar to *hda19-3, stk*
and *stk shp1 shp2*, showed an increase in H3K9/14 acetylation at the level of the
TSS, and a moderate increase at the promoter (Figure 7A, B), suggesting a
similar function of HDA19 and STK in regulating the acetylation state of the
regions tested.
Finally, we verified whether STK could directly bind to the *STM* genomic region
by performing a ChIP-qPCR using an anti-GFP antibody on flowers of *wild-type*
and *pSTK::STK-GFP* plants. MADS-box proteins bind DNA motifs called CarG
boxes. We found, and tested by ChIP-qPCR, CArG boxes in the proximal
promoter area, in the first intron and second intron of *STM* (Figure 7C). In
addition, we also tested the region near the TSS of *STM* as it was the most
enriched in the AcH3K9/14 ChIP-qPCR (Figure 7A). As a positive control we used
VERDANDI (*VDD*), a direct target of STK (Matias-Hernandez et al., 2010) (Figure
7D). The first *STM* intron showed the strongest enrichment, followed by the TSS.
The second intron and the promoter showed no enrichment (Figure 7D). This

result indicates that STK can directly bind the *STM* locus, and that its action is
required to regulate its expression level (Figure 6 A-H).
Based on our phenotypic analysis, expression studies and ChIP-qPCR
experiments, we show that HDA19 and STK have a similar effect over *STM* in
the regulation of its levels and domain of expression. Such activity is required to
regulate the CMM functionality and allow ReT development and ovule number
determination, both traits essential for plant fitness (Figure 7E).
513

Figure 7. STK binds to *STM* locus. A-B) AcH3K9 ChIP-qPCR of *STM* TSS (**A**)
and of a distal promoter region (-2kb) (**B**) in *wild-type, hda19-3, stk* and *stk shp1*
shp2 mutants. The pattern of acetylation in the *STM* locus is similar among the
different mutant genotypes. Graphs show the mean ± sd value of enrichment
among three independent biological replicates of the AcH3K9 ChIP-qPCR
experiment performed on *wild-type, hda19-3, stk and stkshp1sp2* flowers.
Student's T-test, ** = p-value < 0. 01, *** = p-value < 0. 001. **C)** Regions of *STM*
locus tested by anti-GFP ChIP-qPCR on *wild-type* and *pSTK::STK-GFP* flowers.
Triangles mark CaRG boxes detected by AthaMap (Steffens et al., 2004). The

differentially acetylated region at the TSS of *STM* is marked with an orange line.

D) ChIP-qPCR to test binding of STK (*pSTK::STK-GFP*) to the promoter, TSS,

first intron and second intron of *STM*. *VDD* is a known direct target of STK, so it

was used as positive control. Graph shows the mean ± sd enrichment value

among four independent biological replicates of the anti-GFP ChIP-qPCR

experiment. **E)** Model of the regulation of *STM* expression in the CMM involving

HDA19 and STK. Until stage 7, the expression of *STM* in the CMM would be high

thanks to the presence of acetylated histones in association with *STM* locus. At

stages 8-9, when the CMM has to start producing the reproductive tract organs,

HDA19 would deacetylate the histones associated with *STM* locus, to

downregulate it and allow the formation of ovule primordia and TT. Direct binding

of STK to *STM* locus would facilitate the action of HDA19, potentially by

generating multimeric complexes involved in HDA19 binding and stabilization to

STM.

DISCUSSION

In this work, we observed that HDA19 is necessary for ovule initiation and TT

differentiation (Figure 1, Supplemental Figure 1). To study the cause of these

defects, we performed an RNA-seq on ReT cells sorted by FACS with the aid of

the *pSTK::STK-GFP* line. The transcriptomic analysis has given an overview of

gene expression in tissues expressing STK-GFP fusion protein that include the

CMM, TT, placenta and late sporophytic ovule tissues. Comparing cells

expressing STK-GFP in *wild-type* and *hda19-3* allowed us to focus on the role of

HDA19 in these tissues. Consistent with *hda19-3* phenotype, several genes

involved in meristem maintenance, including *STM,* were upregulated in *hda19-3*

ReT. Instead, genes involved in organogenesis like *CUC1/2*, *ANT*, *PAN*, *PIN1,*

or *HEC1* and *NTT* were downregulated in *hda19-3* sorted cells (Figure 3). Our

results show that HDA19 is required to repress *STM* in the CMM to allow organ

formation and tissue differentiation, and that STK is also involved in this process.

In plants in which either *HDA19* or *STK* are mutated, the *STM* locus is more

acetylated respect to the *wild-type* situation. This leads to *STM* upregulation and

ectopic expression in the CMM leading to a reduced ovule number and defects

in TT differentiation.

To assess whether *STM* de-regulation was the main responsible for the ReT
phenotypes of *hda19-3*, we reduced *STM* levels in *hda19-3* plants. *stm* loss of
function mutants lack a functional SAM and consequently fail to produce flowers.
Therefore, we used a weak constitutive promoter, showing that the
downregulation of *STM* in *hda19-3* improved ovule number and density and
growth rate of pollen tubes in the TT (Figure 5, Supplemental Figure 5). None of
the transgenic lines obtained rescue completely the *hda19-3* phenotype. This is
not surprising as HDA19 regulates many processes, so it is unlikely that ectopic
expression of *STM* is the only cause of all *hda19-3* defects. In addition, *KNAT6*
and *BP*, which are partially redundant with STM in the SAM (Byrne et al., 2002)
were also upregulated in *hda19-3*'s CMM, therefore, their downregulation might
be necessary for a better complementation.
Previously, Scofield and collaborators observed that the overexpression of *STM*
using an inducible overexpression system led to the transformation of ovules into
carpelloid structures (Scofield et al., 2007). This aligns with our observations but
represents a more extreme phenotype probably because the levels of *STM*
expressed in the flower of the transgenic plants were much higher than those
observed in *hda19-3.* If STM negatively regulates *STK, SHP1* and *SHP2*, its
ectopic expression in the ovules could lead to a downregulation of STK and
SHP1/2 strong enough to cause the transformation of the ovule integuments into
carpelloid structures as described by Scofield et al., 2007. Our transcriptome
analysis clearly showed the downregulation of genes necessary for organ
primordia formation as well as ovule identity in *hda19-3* (Figure 3), which might
be due to the action of STM. This is similar to what happens in other meristems
like the SAM or the IM (Long et al., 2006; Chung et al., 2019) providing functional
evidence to support the idea that the CMM functions in a similar way as its
indeterminate counterparts. However, how its activity is terminated requires
further research, as ISH showed that after stage 10, *STM* expression in *hda19-3*
is similar than in *wild-type* (Figure 4A-I), suggesting that besides HDA19, other
factors contribute to *STM* silencing in the CMM, since lack of HDA19 delays *STM*
silencing but does not avoid it completely.
STK was first described as a member of the ovule identity complex redundant
with *SHP1/2* (Pinyopich et al., 2003; Favaro et al., 2003). However, STK and
SHP1/2 have also non-redundant functions along reproductive development

(Liljegren et al., 2000; Pinyopich et al., 2003; Mizzotti et al., 2012, 2014, Ezquer

et al., 2016; DiMarzo et al.,2020). Our results support that STK is the main player

required for HDA19's action over *STM* in the CMM as the pattern of histone

acetylation of the *STM* locus is similar in both mutants (Figure 7A-B). However,

while *stk* shows a reduction of ovule number and density similar to *hda19-3*, the

phenotype regarding the TT in *stk* is milder than *hda19-3*´s. The role of STK in

the development of the TT was already described (Marsch-Martínez et al., 2014;

Herrera-Ubaldo et al., 2019, Di Marzo et al., 2020), but single *stk* mutants showed

very mild defects that became more severe when *stk* was combined with *ntt* and

ces mutants. Overall, this suggests that in the TT, HDA19 might be partnering

with other proteins besides STK to regulate *STM*.

Finally, we found that STK binds *STM* locus at the level of the TSS and the first

intron (Figure 7C-D). MADS-box proteins form tetramers able to bind

simultaneously to two DNA binding sites, creating DNA loops (Smaczniak et al.,

2012). As the TSS is the region that is over-acetylated in *hda19-3* and *stk*

mutants, and STK binds at the level of the TSS and the first intron, a plausible

model could be that a MADS-box complex containing STK and HDA19, binds to

STM locus forming a loop between the TSS and the first intron. Then, HDA19

would deacetylate the TSS region of *STM* reducing its expression, allowing organ

initiation from the CMM (Figure 7E). It has been previously shown that STK

displays both activator and repressor properties (Mizzotti et al., 2014), suggesting

that only a fraction of its direct targets are regulated through histone de-

acetylation (Mizzotti et al., 2014). This supports STK ability to work in different

multimeric complexes involving different partners. By Yeast Two Hybrid assay we

have not been able to highlight a direct interaction between HDA19 and STK. For

this reason, it is likely the STK and HDA19 interaction is indirect and/or regulated

by post-translational modifications. Exploring this is outside the scope of the

present work, but it will be extremely interesting to investigate the mechanism of

their interaction in the future.

Another interesting observation is that *STK* expression starts at stage 8, in the

flat placenta, well before OP start to form. In light of our results, the onset of *STK*

expression at stage 8, would lead to the reduction of the levels of *STM,* allowing

ovule initiation and septum fusion (Pinyopich et al., 2003). This idea is supported

by the observation that the levels of *STM* are normal in *hda19-3* before stage 8

(Figure 4 A-H) suggesting that HDA19 exerts its action over *STM* during stages

8-10, when STK starts to accumulate.

To our knowledge, the only work on transcriptomics of sorted gynoecium cells

was done with the aid of a dexamethasone-inducible *pAP1::AP1-GR ap1cal*

system (Villarino et al., 2016). This system allows to obtain many synchronized

flowers, but dexamethasone can introduce variability and/or artifacts. Indeed, the

authors observed occasional ectopic expression of SHP2-YFP outside the

normal expression domain of *SHP2* (Villarino et al., 2016). Our work

demonstrates that it is possible to perform consistent FACS-coupled

transcriptomics of gynoecium cells using markers of moderate expression without

the aid of inducible systems. Moreover, as our dataset contains TT cells and

placenta and sporophytic ovule cells, it will be highly useful for future studies of

CMM and ovule development. Indeed, we already found hints of the possible role

of several genes (Wynn et al., 2011; Villarino et al., 2016). For example, *YAB3*

was proposed to have a non-cell autonomous role in CMM development (Wynn

et al., 2011), but our data indicates otherwise (Figure 3). *YAB3* represses medial

factors (González-Reig et al., 2012) and its mutation affects CMM development

(Nole-Wilson, 2006; Wynn et al., 2011b). Wynn and collaborators did not detect

YAB3 in the medial domain by ISH, so its effect was proposed to be non-cell

autonomous (Wynn et al., 2011). As *pSTK::STK-GFP* is not expressed in the

lateral domain, our data indicates that *YAB3* is indeed expressed in the middle

domain at low levels in the wild-type that might be undetectable by ISH.

Our results show that the tight control of *STM* levels achieved through the action

of HDA19 and STK are essential to maximize reproductive fitness of the plant, as

even moderate changes in its expression lead to defects in the ReT that reduce

seed set.

EXPERIMENTAL PROCEDURS

Plant materials and growth conditions

Plants for phenotypical analysis were grown in a greenhouse under long day

conditions (16 hours light, 8 hours dark) at 22-24 °C.

For sorting experiments, plants were grown in a walk-in growth chamber under

continuous light at 22 °C.

Wild-type Col-0 seeds were already available in the laboratory. *hda19-3*
(SALK_139445) was provided by the Nottingham Arabidopsis Stock Center
(NASC).
pSTK::STK-GFP line was available in the laboratory and first described in Mizzotti
et al., 2014.
Stk and *STK +/- shp1 shp2* seeds (used to obtain the *stk shp1 shp2* triple mutant
from segregation) were already available in the laboratory and previously
described by (Pinyopich et al., 2003; Favaro et al., 2003).
666

Ovule number and placenta length measurement
Inflorescences were fixed overnight in ethanol:acetic acid 9:1 and rehydrated in
an ethanol series. Pistils (stages 6-10) were dissected and mounted on chloral
hydrate:glycerol:water (8:1:3, w/v/v) and immediately observed under a Zeiss
Axiophot D1 microscope (Carl Zeiss MicroImaging) equipped with DIC optics and
an Axiocam MRc5 camera (Zeiss) with Axiovision software (version 4.1). Ovules
were counted manually. Placenta length was measured using Axiovision
software.
675

Pistil histology and imaging
Individual flowers from *wild-type*, *hda19-3, 35sRNAi:STM* and *stk* were fixed in
2% (w/v) paraformaldehyde and 2,5% (w/v) glutaraldehyde in PIPES buffer
[0.025 M, pH 7, 0,001 % (v/v) Tween-80], placed under vacuum for 1h and then
at 4ºC overnight. Material was dehydrated in an ethanol series and embedded in
LR White resin. Thick sections (500 µm) were obtained with a Leica EM UC7
Ultramicrotome stained with 1 % (w/v) toluidine blue (Sigma-Aldrich, St Louis,
MO, US) and mounted with DPX (Sigma). Slides were observed under a Zeiss
AxioImager AZ microscope equipped with a Zeiss Axiocam MRc3 camera using
Zen Imaging software (Zen 2011 SP1).
686

Aniline blue staining for pollen tube growth analysis
Flowers were emasculated and pollinated with *wild-type* pollen. Pollen tubes
were allowed to grow for different times depending on the experiment (12, 14 or
24h). Then, pistils were fixed overnight in 9:1 ethanol:acetic acid, washed three

times with water, incubated in 1M NaOH overnight, washed three times in water

and incubated in 0,1% aniline blue (w/v). Pistils were imaged using a Zeiss

AxioPhot D1 microscope equipped with fluorescence filters. Images were

recorded with an Axiocam MRc5 camera (Zeiss) with Axiovision software (version

4.1). Pollen tubes length was measured using ImageJ (Schneider et al., 2012).

Protoplast recovery and Fluorescence Activated Cell Sorting (FACS)

pSTK::STK-GFP construct was introgressed into *hda19-3* background and then

used for cell sorting, along with the parental *pSTK::STK-GFP*. Starting material

consisted in flowers from stages 1-12. Protoplasting and sorting was performed

as described by Villarino et al., 2016, with some modifications. Cells were sorted

into RLT buffer from the RNAeasy Microkit (Qiagen) supplemented with 1% β-

mercaptoethanol (v/v), keeping a ratio of 3 volumes of RLT buffer per 1 volume

of sorted cells (in our sorting conditions 50,000 cells occupied 200 µL). Cell

sorting was performed using a Moflo XDP (Beckman Coulter Inc.). Each

experiment yielded between 80,000 and 100,000 GFP-positive cells.

RNA extraction and expression analysis

RNA was extracted immediately after sorting using an RNAeasy Microkit (Qiagen)

with a few modifications. We used one column for every 25,000 cells. Columns

were eluted twice, using the first eluate for the second elution. Finally, all eluates

from the same sample were pooled, yielding 70-100 ng of total RNA. For qRT-

PCR validation, the columns were eluted a third time, using 14 µL of nuclease-

free water that was sequentially used to elute all the columns corresponding to

the same sample. This yielded around 5 ng of total RNA that was retro-

transcribed using SuperScript III First- Strand Synthesis System (Invitrogen/Life

Technologies). Then, expression of *STK-GFP* was measured by qRT-PCR using

a Thermal Cycler from Applied Biosystems and a QuantiTect SYBR Green PCR

Kit (Qiagen). *MON1* (*AT2G28390*) was used as a reference gene for

normalization based on Czechowski et al., 2005.

For evaluation of *STM* levels in RNAi lines, RNA was extracted using Trizol

(Invitrogen) and 500 ng of RNA were retrotranscribed using iScript cDNA

synthesis kit (BioRad). qRT-PCRs were performed using iTaq Universal SYBR

Green Supermix (BioRad) in a Bio-Rad iCycler iQ thermal cycler. *ACT8* gene
(*AT1G49240*) was used as reference gene for normalization because, according
to our RNA-seq data, its RNA levels do not change in *hda19-3* plants.
All primers used in qRT-PCR experiments showed an amplification efficiency
close to 100%, so the $2^{-\Delta\Delta CT}$ method was used for calculation of expression
changes.
All primers used are listed in Supplemental Table 1.
731

RNA sequencing and analysis
Three validated samples of RNA from *wild-type* GFP-positive cells and 4 samples
from *hda19-3* GFP-positive cells were used for library preparation. Libraries were
sequenced in one lane of the HiSeq2500 Illumina platform with a yield of 250
million reads and an average of 35 million reads per sample. Around 10 million
reads were filtered out and of the 240 million reads remaining, 230 million
mapped uniquely (96% of the reads after filtering) and were used for subsequent
analysis.
All steps of library preparation, RNA sequencing and bioinformatics analysis were
essentially performed as in Villarino et al., 2016. Briefly, we used Cufflinks
(Trapnell et al., 2012), DeSeq2 (Love et al., 2014) and EdgeR (Robinson et al.,
2009) to analyze the RNA-seq data (Supplemental File 1-3). Then, we crossed
the data obtained by the 3 software and selected the DEGs detected by the three
programs, with an FDR < 0.001 and with a logFC > |1.2| (Supplemental File 4).
Genes whose expression value was zero in either *wild-type* or *hda19-3* were
filtered out.
748

Chromatin immunoprecipitation and qPCR analysis
Chromatin immunoprecipitation was performed as described in Mizzotti et al.,
2014, using a rabbit anti-H3 acetyl K9 (Upstate 07-352, Sigma-Aldrich) and a
mouse monoclonal anti-GFP antibody from Roche (11814460001).
qPCR was performed using iTaq Universal SYBR Green Supermix (BioRad) in a
Bio-Rad iCycler iQ thermal cycler.

Enrichment of tested areas was calculated as fold change against a control non-
enriched region (*GAPDH* locus) as previously described by Matias-Hernandez et
al., 2010. Primers used for ChIP-qPCR are listed in Supplemental Table 1.
758

759 ***In-situ* hybridization**
760 Sectioning and *in situ* hybridization were performed as described by (Dreni et al.,
761 2007). *STM* probe was obtained as described in Simonini & Kater, 2014.
762

Generation of RNAi lines
For *35s::STM-RNAi* construct, a fragment of 352 bp of *STM* coding sequence
was amplified (see primer list in Supplemental Table 3), cloned into pDONR207
and then subcloned into pFRH (derived from pFGC5941; NCBI accession
number AY310901) using Gateway technology (Invitrogen).
The construct was transformed into *Agrobacterium tumefaciens* strain EHA105
and *hda19-3* plants were transformed by floral dip (Clough and Bent, 1998).
Transformants were selected in MS media (Murashige and Skoog, 1962)
supplemented with 20 mg/L of hygromycin. Presence of the construct was
confirmed by genotyping (see primers in Supplemental Table 1).
773

774 **Clustering analysis**
775 Heatmaps were obtained using Heatmapper (Babicki et al., 2016). Genes were
776 hierarchically clustered by Euclidean distances using average linkage method.
777

778 **Data availability**
779 RNA-seq data are available at the accession number (XXXXX)
780

Funding
Manrique S, Cavalleri A, Colombo L, were founded by MIUR PRIN2017.
Guazzotti A, Colombo L were founded by H2020-EU.1.3. (SEXSEED ID: 690946).
Manrique S was founded by EMBO SHORT TERM (2015). Manrique S, Masiero
S, Mizzotti C, were founded by MC-IRSES - International research staff exchange
scheme (IRSES) (FRUITLOOK ID: 612640). Coimbra S was founded by
SeedWheels FCT Project - POCI-01-0145-FEDER-027839.

Author contribution

Conceived and design the research: MS, CL, CA, GA, FR. Performed the experiments: MS, CA, GA, VG, OE, PAM, CS, SS. Contributed reagents/materials/analysis tools: CL, FR, HT, GU. Analysed the data: MS, AC, VG, BA. Wrote the manuscript: MS, CA, CL, GA

Acknowledgments
Thanks to Sarah Schuett (NCSU) and Claudia Bazzini (UniMi) for technical assistance with FACS and the Genomic Sciences Laboratory Research Facility (NCSU) for library preparation and Illumina sequencing.

Competing interests
The Authors declare no competing interests.

Literature cited
Alvarez, J., and Smyth, D. R. (2002). CRABS CLAW and SPATULA genes regulate growth and pattern formation during gynoecium development in Arabidopsis thaliana. *Int J Plant Sci* 163, 17–41. doi: 10.1086/324178.

Azhakanandam, S., Nole-Wilson, S., Bao, F., and Franks, R. G. (2008). SEUSS and AINTEGUMENTA Mediate Patterning and Ovule Initiation during Gynoecium Medial Domain Development 1[W][OA]. doi: 10.1104/pp.107.114751.

Babicki, S., Arndt, D., Marcu, A., Liang, Y., Grant, J. R., Maciejewski, A., et al. (2016). Heatmapper: web-enabled heat mapping for all. *Nucleic Acids Res* 44, W147–W153. doi: 10.1093/nar/gkw419.

Bell, E. M., Lin, W. C., Husbands, A. Y., Yu, L., Jaganatha, V., Jablonska, B., et al. (2012). Arabidopsis lateral organ boundaries negatively regulates brassinosteroid accumulation to limit growth in organ boundaries. *Proc Natl Acad Sci U S A* 109, 21146–21151. doi: 10.1073/pnas.1210789109.

Belles-Boix, E., Hamant, O., Witiak, S. M., Morin, H., Traas, J., and Pautot, V. (2006). *KNAT6*: An *Arabidopsis* Homeobox Gene Involved in Meristem Activity and Organ Separation. *Plant Cell* 18, 1900–1907. doi: 10.1105/tpc.106.041988.

Benhamed, M., Bertrand, C., Servet, C., and Zhou, D. X. (2006). Arabidopsis
GCN5, HD1, and TAF1/HAF2 interact to regulate histone acetylation
required for liqht-responsive gene expression. *Plant Cell* 18, 2893–2903. doi:
10.1105/tpc.106.043489.

Bollier, N., Sicard, A., Leblond, J., Latrasse, D., Gonzalez, N., Gévaudant, F., et
al. (2018). At-MINI ZINC FINGER2 and SL-INHIBITOR of MERISTEM
ACTIVITY, a conserved missing link in the regulation of floral meristem
termination in arabidopsis and tomato. *Plant Cell* 30, 83–100. doi:
10.1105/tpc.17.00653.

Bowman, J. L., Baum, S. F., Eshed, Y., Putterill, J., and Alvarez, J. (1999).
Molecular genetics of gynoecium development in Arabidopsis. *Curr Top Dev*
Biol 45, 155–161. doi: 10.1016/s0070-2153(08)60316-6.

Byrne, M. E., Simorowski, J., and Martienssen, R. A. (2002). ASYMMETRIC
LEAVES1 reveals know gene redundancy in Arabidopsis. *Development* 129,
1957–1965. doi: 10.1242/dev.129.8.1957.

Chávez Montes, R. A., Herrera-Ubaldo, H., Serwatowska, J., and de Folter, S.
(2015). Towards a comprehensive and dynamic gynoecium gene regulatory
network. *Curr Plant Biol* 3–4, 3–12. doi: 10.1016/j.cpb.2015.08.002.

Choi, S.-M., Song, H.-R., Han, S.-K., Han, M., Kim, C.-Y., Park, J., et al. (2012).
HDA19 is required for the repression of salicylic acid biosynthesis and
salicylic acid-mediated defense responses in Arabidopsis. *The Plant Journal*
71, 135–146. doi: 10.1111/j.1365-313X.2012.04977.x.

Chung, Y., Zhu, Y., Wu, M. F., Simonini, S., Kuhn, A., Armenta-Medina, A., et al.
(2019a). Auxin Response Factors promote organogenesis by chromatin-
mediated repression of the pluripotency gene SHOOTMERISTEMLESS. *Nat*
Commun 10, 1–11. doi: 10.1038/s41467-019-08861-3.

Chung, Y., Zhu, Y., Wu, M.-F., Simonini, S., Kuhn, A., Armenta-Medina, A., et al.
(2019b). Auxin Response Factors promote organogenesis by chromatin-
mediated repression of the pluripotency gene SHOOTMERISTEMLESS. *Nat*
Commun 10, 886. doi: 10.1038/s41467-019-08861-3.

Clough, S. J., and Bent, A. F. (1998). Floral dip: A simplified method for
Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant*
Journal 16, 735–743. doi: 10.1046/j.1365-313X.1998.00343.x.

Colombo, M., Brambilla, V., Marcheselli, R., Caporali, E., Kater, M. M., and
Colombo, L. (2010). A new role for the SHATTERPROOF genes during
Arabidopsis gynoecium development. *Dev Biol* 337, 294–302. doi:
10.1016/j.ydbio.2009.10.043.

Crawford, B. C. W., and Yanofsky, M. F. (2008). The Formation and Function of
the Female Reproductive Tract in Flowering Plants. *Current Biology* 18,
R972–R978. doi: 10.1016/j.cub.2008.08.010.

Cucinotta, M., Colombo, L., and Roig-Villanova, I. (2014). Ovule development, a
new model for lateral organ formation. *Front Plant Sci* 5, 117. doi:
10.3389/fpls.2014.00117.

Czechowski, T., Stitt, M., Altmann, T., Udvardi, M. K., and Scheible, W.-R. (2005).
Genome-Wide Identification and Testing of Superior Reference Genes for
Transcript Normalization in Arabidopsis. *Plant Physiol* 139, 5–17. doi:
10.1104/pp.105.063743.

Di Marzo, M., Herrera-Ubaldo, H., Caporali, E., Novák, O., Strnad, M., Balanzà,
V., et al. (2020a). SEEDSTICK Controls Arabidopsis Fruit Size by Regulating
Cytokinin Levels and FRUITFULL. *Cell Rep* 30, 2846-2857.e3. doi:
10.1016/j.celrep.2020.01.101.

Di Marzo, M., Roig-Villanova, I., Zanchetti, E., Caselli, F., Gregis, V., Bardetti, P.,
et al. (2020b). MADS-Box and bHLH Transcription Factors Coordinate
Transmitting Tract Development in Arabidopsis thaliana. *Front Plant Sci* 11,
526. doi: 10.3389/fpls.2020.00526.

Dreni, L., Jacchia, S., Fornara, F., Fornari, M., Ouwerkerk, P. B. F., An, G., et al.
(2007). The D-lineage MADS-box gene OsMADS13 controls ovule identity
in rice. *The Plant Journal* 52, 690–699. doi: 10.1111/j.1365-
313X.2007.03272.x.

Durbak, A. R., and Tax, F. E. (2011). CLAVATA signaling pathway receptors of
arabidopsis regulate cell proliferation in fruit organ formation as well as in
meristems. *Genetics* 189, 177–194. doi: 10.1534/genetics.111.130930.

Favaro R, Pinyopich A, Battaglia R, Kooiker M, Borghi L, Ditta G,Yanofsky MF,
Kater MM, C. L. (2003). MADS-Box Protein Complexes Control Carpel and
Ovule Development in Arabidopsis. *Plant Cell* 15, 2603–2611. Available at:
http://www.plantcell.org/content/plantcell/15/11/2603.full.pdf   [Accessed
November 3, 2018].

Fornara, F., Pařenicová, L., Falasca, G., Pelucchi, N., Masiero, S., Ciannamea,
S., et al. (2004). Functional Characterization of *OsMADS18* , a Member of
the *AP1/SQUA* Subfamily of MADS Box Genes. *Plant Physiol* 135, 2207–
2219. doi: 10.1104/pp.104.045039.

Gaillochet, C., and Lohmann, J. U. (2015). The never-ending story: from
pluripotency to plant developmental plasticity. *Development* 142, 2237–
2249. doi: 10.1242/dev.117614.

Galbiati, F., Sinha Roy, D., Simonini, S., Cucinotta, M., Ceccato, L., Cuesta, C.,
et al. (2013). An integrative model of the control of ovule primordia formation.
Plant J 76, 446–55. doi: 10.1111/tpj.12309.

Gendron, J. M., Liu, J. S., Fan, M., Bai, M. Y., Wenkel, S., Springer, P. S., et al.
(2012). Brassinosteroids regulate organ boundary formation in the shoot
apical meristem of Arabidopsis. *Proc Natl Acad Sci U S A* 109, 21152–
21157. doi: 10.1073/pnas.1210799110.

González-Reig, S., Ripoll, J. J., Vera, A., Yanofsky, M. F., and Martínez-Laborda,
A. (2012). Antagonistic Gene Activities Determine the Formation of Pattern
Elements along the Mediolateral Axis of the Arabidopsis Fruit. *PLoS Genet*
8, e1003020. doi: 10.1371/journal.pgen.1003020.

Gorham, S. R., Weiner, A. I., Yamadi, M., and Krogan, N. T. (2018). HISTONE
DEACETYLASE 19 and the flowering time gene FD maintain reproductive
meristem identity in an age-dependent manner. *J Exp Bot* 69, 4757–4771.
doi: 10.1093/jxb/ery239.

Gremski, K., Ditta, G., and Yanofsky, M. F. (2007). The HECATE genes regulate
female reproductive tract development in Arabidopsis thaliana. *Development*
134, 3593–3601. doi: 10.1242/dev.011510.

Guo, L., Zhou, J., Elling, A. A., Charron, J. B. F., and Xing, W. D. (2008). Histone
modifications and expression of light-regulated genes in arabidopsis are
cooperatively influenced by changing light conditions. *Plant Physiol* 147,
2070–2083. doi: 10.1104/pp.108.122929.

He, J.-X. (2005). BZR1 Is a Transcriptional Repressor with Dual Roles in
Brassinosteroid Homeostasis and Growth Responses. *Science (1979)* 307,
1634–1638. doi: 10.1126/science.1107580.

Heidstra, R., and Sabatini, S. (2014). Plant and animal stem cells: Similar yet
different. *Nat Rev Mol Cell Biol* 15, 301–312. doi: 10.1038/nrm3790.

Herrera-Ubaldo, H., Lozano-Sotomayor, P., Ezquer, I., Di Marzo, M., Montes, R.
A. C., Goḿez-Felipe, A., et al. (2019). New roles of NO TRANSMITTING
TRACT and SEEDSTICK during medial domain development in arabidopsis
fruits. *Development (Cambridge)* 146. doi: 10.1242/dev.172395.

Ishida, T., Aida, M., Takada, S., and Tasaka, M. (2000). Involvement of CUP-
SHAPED COTYLEDON Genes in Gynoecium and Ovule Development in
Arabidopsis thaliana. *Plant Cell Physiol* 41, 60–67. doi: 10.1093/pcp/41.1.60.

Jan, I. C., Chun, P. J., Hemmes, H., Jung, C., and Chua, N. H. (2011). Rapid and
reversible light-mediated chromatin modifications of arabidopsis
Phytochrome A Locus. *Plant Cell* 23, 459–470. doi:
10.1105/tpc.110.080481.

Kanrar, S., Onguka, O., and Smith, H. M. S. (2006). Arabidopsis inflorescence
architecture requires the activities of KNOX-BELL homeodomain
heterodimers. *Planta* 224, 1163–1173. doi: 10.1007/s00425-006-0298-9.

Kay, P., Groszmann, M., Ross, J. J., Parish, R. W., and Swain, S. M. (2013).
Modifications of a conserved regulatory network involving
<scp>INDEHISCENT</scp> controls multiple aspects of reproductive tissue
development in Arabidopsis. *New Phytologist* 197, 73–87. doi:
10.1111/j.1469-8137.2012.04373.x.

Kim, K. C., Lai, Z., Fan, B., and Chen, Z. (2008). Arabidopsis WRKY38 and
WRKY62 transcription factors interact with histone deacetylase 19 in basal
defense. *Plant Cell* 20, 2357–2371. doi: 10.1105/tpc.107.055566.

Kim, S., Piquerez, S. J. M., Ramirez-Prado, J. S., Mastorakis, E., Veluchamy, A.,
Latrasse, D., et al. (2021). GCN5 modulates salicylic acid homeostasis by
regulating H3K14ac levels at the 5' and 3' ends of its target genes. *Nucleic*
Acids Res 48, 5953–5966. doi: 10.1093/NAR/GKAA369.

Krogan, N. T., Hogan, K., and Long, J. A. (2012a). APETALA2 negatively
regulates multiple floral organ identity genes in Arabidopsis by recruiting the
co-repressor TOPLESS and the histone deacetylase HDA19. *Development*
139, 4180–90. doi: 10.1242/dev.085407.

Krogan, N. T., Hogan, K., and Long, J. A. (2012b). APETALA2 negatively
regulates multiple floral organ identity genes in Arabidopsis by recruiting the
co-repressor TOPLESS and the histone deacetylase HDA19. *Development*
139, 4180–90. doi: 10.1242/dev.085407.

Kuhn, A., Harborough, S. R., McLaughlin, H. M., Natarajan, B., Verstraeten, I., Friml, J., et al. (2020). Direct ETTIN-auxin interaction controls chromatin states in gynoecium development. *Elife* 9. doi: 10.7554/eLife.51787.

Kumar, V., Thakur, J. K., and Prasad, M. (2021). Histone acetylation dynamics regulating plant development and stress responses. *Cellular and Molecular Life Sciences* 78, 4467–4486. doi: 10.1007/s00018-021-03794-x.

Landrein, B., Kiss, A., Sassi, M., Chauvet, A., Das, P., Cortizo, M., et al. (2015). Mechanical stress contributes to the expression of the STM homeobox gene in Arabidopsis shoot meristems. *Elife* 4. doi: 10.7554/eLife.07811.001.

Lee, B. H., Kwon, S. H., Lee, S. J., Park, S. K., Song, J. T., Lee, S., et al. (2015). The Arabidopsis thaliana NGATHA transcription factors negatively regulate cell proliferation of lateral organs. *Plant Mol Biol* 89, 529–538. doi: 10.1007/s11103-015-0386-y.

Lee, C., and Clark, S. E. (2015). A WUSCHEL-Independent Stem Cell Specification Pathway Is Repressed by PHB, PHV and CNA in Arabidopsis. *PLoS One* 10, e0126006. doi: 10.1371/journal.pone.0126006.

Liljegren, S. J., Ditta, G. S., Eshed, Y., Savidge, B., Bowmant, J. L., and Yanofsky, M. F. (2000). SHATTERPROOF MADS-box genes control dispersal in Arabidopsis. *Nature* 404, 766–770. doi: 10.1038/35008089.

Long, J. A., Moan, E. I., Medford, J. I., and Barton, M. K. (1996). A member of the KNOTTED class of homeodomain proteins encoded by the STM gene of Arabidopsis. *Nature* 379, 66–69. doi: 10.1038/379066a0.

Long, J. A., Ohno, C., Smith, Z. R., and Meyerowitz, E. M. (2006). TOPLESS regulates apical embryonic fate in Arabidopsis. *Science* 312, 1520–3. doi: 10.1126/science.1123841.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15, 550. doi: 10.1186/s13059-014-0550-8.

Marsch-Martínez, N., Zúñiga-Mayo, V. M., Herrera-Ubaldo, H., Ouwerkerk, P. B. F., Pablo-Villa, J., Lozano-Sotomayor, P., et al. (2014). The NTT transcription factor promotes replum development in Arabidopsis fruits. *The Plant Journal* 80, 69–81. doi: 10.1111/tpj.12617.

Matias-Hernandez, L., Battaglia, R., Galbiati, F., Rubes, M., Eichenberger, C., Grossniklaus, U., et al. (2010). VERDANDI is a direct target of the MADS

domain ovule identity complex and affects embryo sac differentiation in
Arabidopsis. *Plant Cell* 22, 1702–15. doi: 10.1105/tpc.109.068627.

Mi, H., Muruganujan, A., Ebert, D., Huang, X., and Thomas, P. D. (2018).
PANTHER version 14: more genomes, a new PANTHER GO-slim and
improvements in enrichment analysis tools. *Nucleic Acids Res* 47, D419–
D426. doi: 10.1093/nar/gky1038.

Mi, H., Muruganujan, A., Huang, X., Ebert, D., Mills, C., Guo, X., et al. (2019).
Protocol Update for large-scale genome and gene function analysis with the
PANTHER classification system (v.14.0). *Nat Protoc* 14, 703–721. doi:
10.1038/s41596-019-0128-8.

Mizukami, Y., and Fischer, R. L. (2000). Plant organ size control:
AINTEGUMENTA regulates growth and cell numbers during organogenesis.
Proc Natl Acad Sci U S A 97, 942–947. doi: 10.1073/pnas.97.2.942.

Mizzotti, C., Ezquer, I., Paolo, D., Rueda-Romero, P., Guerra, R. F., Battaglia,
R., et al. (2014). SEEDSTICK is a master regulator of development and
metabolism in the Arabidopsis seed coat. *PLoS Genet* 10, e1004856. doi:
10.1371/journal.pgen.1004856.

Mizzotti, C., Mendes, M. A., Caporali, E., Schnittger, A., Kater, M. M., Battaglia,
R., et al. (2012). The MADS box genes SEEDSTICK and ARABIDOPSIS
Bsister play a maternal role in fertilization and seed development. *Plant J* 70,
409–20. doi: 10.1111/j.1365-313X.2011.04878.x.

Murashige, T., and Skoog, F. (1962). A Revised Medium for Rapid Growth and
Bio Assays with Tobacco Tissue Cultures. *Physiol Plant* 15, 473–497. doi:
10.1111/j.1399-3054.1962.tb08052.x.

Nahar, Most. A.-U., Ishida, T., Smyth, D. R., Tasaka, M., and Aida, M. (2012).
Interactions of CUP-SHAPED COTYLEDON and SPATULA Genes Control
Carpel Margin Development in Arabidopsis thaliana. *Plant Cell Physiol* 53,
1134. doi: 10.1093/PCP/PCS057.

Norberg, M., Holmlund, M., and Nilsson, O. (2005). The BLADE ON PETIOLE
genes act redundantly to control the growth and development of lateral
organs. *Development* 132, 2203–2213. doi: 10.1242/dev.01815.

Pekker, I., Alvarez, J. P., and Eshed, Y. (2005). Auxin Response Factors Mediate
Arabidopsis Organ Asymmetry via Modulation of KANADI Activity. *Plant Cell*
17, 2899–2910. doi: 10.1105/tpc.105.034876.

Pereira, A. M., Moreira, D., Coimbra, S., and Masiero, S. (2021). Paving the Way for Fertilization: The Role of the Transmitting Tract. *Int J Mol Sci* 22, 2603. doi: 10.3390/ijms22052603.

Pi, L., Aichinger, E., van der Graaff, E., Llavata-Peris, C. I., Weijers, D., Hennig, L., et al. (2015a). Organizer-Derived WOX5 Signal Maintains Root Columella Stem Cells through Chromatin-Mediated Repression of CDF4 Expression. *Dev Cell* 33, 576–88. doi: 10.1016/j.devcel.2015.04.024.

Pi, L., Aichinger, E., van der Graaff, E., Llavata-Peris, C. I., Weijers, D., Hennig, L., et al. (2015b). Organizer-Derived WOX5 Signal Maintains Root Columella Stem Cells through Chromatin-Mediated Repression of CDF4 Expression. *Dev Cell* 33, 576–588. doi: 10.1016/j.devcel.2015.04.024.

Pinyopich, A., Ditta, G. S., Savidge, B., Liljegren, S. J., Baumann, E., Wisman, E., et al. (2003). Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature* 424, 85–88. doi: 10.1038/nature01741.

Poulios, S., and Vlachonasios, K. E. (2018). Synergistic action of *GCN5* and *CLAVATA1* in the regulation of gynoecium development in *Arabidopsis thaliana*. *New Phytologist* 220, 593–608. doi: 10.1111/nph.15303.

Ragni, L., Belles-Boix, E., Günl, M., and Pautot, V. (2008). Interaction of KNAT6 and KNAT2 with Brevipedicellus and Pennywise in Arabidopsis inflorescences. *Plant Cell* 20, 888–900. doi: 10.1105/tpc.108.058230.

Reyes-Olalde, J. I., and de Folter, S. (2019). Control of stem cell activity in the carpel margin meristem (CMM) in Arabidopsis. *Plant Reprod* 32, 123–136. doi: 10.1007/s00497-018-00359-0.

Reyes-Olalde, J. I., Zuñiga-Mayo, V. M., Chávez Montes, R. A., Marsch-Martínez, N., and de Folter, S. (2013a). Inside the gynoecium: At the carpel margin. *Trends Plant Sci* 18, 644–655. doi: 10.1016/j.tplants.2013.08.002.

Reyes-Olalde, J. I., Zuñiga-Mayo, V. M., Chávez Montes, R. A., Marsch-Martínez, N., and de Folter, S. (2013b). Inside the gynoecium: At the carpel margin. *Trends Plant Sci* 18, 644–655. doi: 10.1016/j.tplants.2013.08.002.

Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2009). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140. doi: 10.1093/bioinformatics/btp616.

Roeder, A. H. K., and Yanofsky, M. F. (2006). Fruit Development in Arabidopsis. *Arabidopsis Book* 4, e0075. doi: 10.1199/tab.0075.

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9, 671–675. doi: 10.1038/nmeth.2089.

Scofield, S., Dewitte, W., and Murray, J. A. H. (2007). The KNOX gene SHOOT MERISTEMLESS is required for the development of reproductive meristematic tissues in Arabidopsis. *The Plant Journal* 50, 767–781. doi: 10.1111/j.1365-313X.2007.03095.x.

Sessions, A., Nemhauser, J. L., McColl, A., Roe, J. L., Feldmann, K. A., and Zambryski, P. C. (1997). ETTIN patterns the Arabidopsis floral meristem and reproductive organs. *Development* 124, 4481–4491. doi: 10.1242/dev.124.22.4481.

Sessions, R. A., and Zambryski, P. C. (1995). Arabidopsis gynoecium structure in the wild type and in ettin mutants. *Development* 121, 1519–1532. doi: 10.1242/dev.121.5.1519.

Simonini, S., and Kater, M. M. (2014). Class I BASIC PENTACYSTEINE factors regulate HOMEOBOX genes involved in meristem size maintenance. *J Exp Bot* 65, 1455–1465. doi: 10.1093/jxb/eru003.

Smaczniak, C., Immink, R. G. H., Angenent, G. C., and Kaufmann, K. (2012). Developmental and evolutionary diversity of plant MADS-domain factors: insights from recent studies. *Development* 139, 3081–98. doi: 10.1242/dev.074674.

Smyth, D. R., Bowman, J. L., and Meyerowitz, E. M. (1990). Early flower development in Arabidopsis. *Plant Cell* 2, 755–767. doi: 10.1105/tpc.2.8.755.

Sohlberg, J. J., Myrenås, M., Kuusk, S., Lagercrantz, U., Kowalczyk, M., Sandberg, G., et al. (2006). STY1 regulates auxin homeostasis and affects apical-basal patterning of the Arabidopsis gynoecium. *Plant J* 47, 112–23. doi: 10.1111/j.1365-313X.2006.02775.x.

Staci Nole-Wilson, B. A. K. (2006). AINTEGUMENTA Contributes to Organ Polarity and Regulates Growth of Lateral Organs in Combination with YABBY Genes. *Plant Physiol* 141, 977–987. doi: 10.1104/pp.104.040394.

Tanaka, M., Kikuchi, A., and Kamada, H. (2008). The arabidopsis histone
deacetylases HDA6 and HDA19 contribute to the repression of embryonic
properties after germination. *Plant Physiol* 146, 149–161. doi:
10.1104/pp.107.111674.

Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D. R., et al. (2012).
Differential gene and transcript expression analysis of RNA-seq experiments
with TopHat and Cufflinks. *Nat Protoc* 7, 562–578. doi:
10.1038/nprot.2012.016.

Uehara, T., Okushima, Y., Mimura, T., Tasaka, M., and Fukaki, H. (2008).
Domain II Mutations in CRANE/IAA18 Suppress Lateral Root Formation and
Affect Shoot Development in Arabidopsis thaliana. *Plant Cell Physiol* 49,
1025–1038. doi: 10.1093/pcp/pcn079.

Vernoux, T., Kronenberger, J., Grandjean, O., Laufs, P., and Traas, J. (2000).
PIN-FORMED 1 regulates cell fate at the periphery of the shoot apical
meristem. *Development* 127, 5157–5165. doi: 10.1242/dev.127.23.5157.

Wang, L., Zhang, F., Rode, S., Chin, K. K., Ko, E. E., Kim, J., et al. (2017).
Ethylene induces combinatorial effects of histone H3 acetylation in gene
expression in Arabidopsis. *BMC Genomics* 18, 538. doi: 10.1186/s12864-
017-3929-6.

Wynn, A. N., Rueschhoff, E. E., and Franks, R. G. (2011a). Transcriptomic
Characterization of a Synergistic Genetic Interaction during Carpel Margin
Meristem Development in Arabidopsis thaliana. *PLoS One* 6, e26231. doi:
10.1371/journal.pone.0026231.

Wynn, A. N., Rueschhoff, E. E., and Franks, R. G. (2011b). Transcriptomic
characterization of a synergistic genetic interaction during carpel margin
meristem development in arabidopsis thaliana. *PLoS One* 6, e26231. doi:
10.1371/journal.pone.0026231.

Zhao, M., Yang, S., Chen, C.-Y., Li, C., Shan, W., Lu, W., et al. (2015).
Arabidopsis BREVIPEDICELLUS Interacts with the SWI2/SNF2 Chromatin
Remodeling ATPase BRAHMA to Regulate KNAT2 and KNAT6 Expression
in Control of Inflorescence Architecture. *PLoS Genet* 11, e1005125. doi:
10.1371/journal.pgen.1005125.

Zhou, C., Zhang, L., Duan, J., Miki, B., and Wu, K. (2005). Histone Deacetylase19 is involved in jasmonic acid and ethylene signaling of pathogen response in Arabidopsis. *Plant Cell* 17, 1196–1204. doi: 10.1105/tpc.104.028514.

Zhou, Y., Tan, B., Luo, M., Li, Y., Liu, C., Chen, C., et al. (2013). HISTONE DEACETYLASE19 interacts with HSL1 and participates in the repression of seed maturation genes in Arabidopsis seedlings. *Plant Cell* 25, 134–48. doi: 10.1105/tpc.112.096313.

Zhu, Z., Xu, F., Zhang, Y., Cheng, Y. T., Wiermer, M., Li, X., et al. (2010). Arabidopsis resistance protein SNC1 activates immune responses through association with a transcriptional corepressor. *Proc Natl Acad Sci U S A* 107, 13960–13965. doi: 10.1073/pnas.1002828107.

Zúñiga-Mayo, V. M., Gómez-Felipe, A., Herrera-Ubaldo, H., and de Folter, S. (2019). Gynoecium development: networks in Arabidopsis and beyond. *J Exp Bot* 70, 1447–1460. doi: 10.1093/jxb/erz026.