Title:

Pseudoirreversible inhibition elicits persistent efficacy of a sphigosine-1-phosphate receptor-1

antagonist

Authors:

Yuya Maruyama[1, 2, 3], Yusuke Ohsawa[3], Takayuki Suzuki[3], Yuko Yamauchi[3], Kohsuke Ohno[3],

Hitoshi Inoue[3], Akitoshi Yamamoto[3], Morimichi Hayashi[3], Yuji Okuhara[3], Wataru Muramatsu[1, 2],

Kano Namiki[1, 2], Maki Miyauchi[1, 2], Takahisa Miyao[1], Naho Hagiwara[1], Tatsuya Ishikawa[1, 2],

Kenta Horie[1], Mio Hayama[1], Nobuko Akiyama[1,2], Takatsugu Hirokawa[4,5] and Taishin Akiyama[1,

2, *]

Affiliations:

[1]Laboratory for Immune Homeostasis, RIKEN Center for Integrative Medical Sciences,

Yokohama 230-0045, Japan

[2]Immunobiology, Graduate School of Medical Life Science, Yokohama City University,

Yokohama 230-0045, Japan

[3]Central Research Laboratory, Kissei Pharmaceutical Co., Ltd., 4365-1 Hotaka-Kashiwabara,

Azumino, Nagano 399-8304, Japan

[4]Division of Biomedical Science, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai,

Tsukuba, Ibaraki 305-8575, Japan

[5]Transborder Medical Research Center, University of Tsukuba, 1-1-1 Tennodai, Tsukuba,

Ibaraki 305-8575, Japan

Contact Information: Taishin Akiyama, Ph.D.

Postal address: 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama 230-0045, Japan

Email: taishin.akiyama@riken.jp

Abstract

Sphingosine 1-phosphate receptor 1 (S1PR1), a G protein-coupled receptor, is required for lymphocyte trafficking, and is a promising therapeutic target in inflammatory diseases. To find potent S1PR1 antagonists, identification of the structural basis for drug efficacy is important. Here, we synthesized a novel antagonist, KSI-6666, that persistently inhibits S1PR1 activity and effectively suppresses pathogenic inflammation. Metadynamics simulation suggested that the interaction of a benzene ring moiety in KSI-6666 with a methionine residue in the ligand-binding pocket of S1PR1 inhibits the dissociation of KSI-6666 from S1PR1, generating a metastable binding state. Consistently, *in vitro* functional and mutational analyses revealed that KSI-6666 causes pseudoirreversible inhibition of S1PR1, dependent on the methionine residue of the protein and substituents on the distal benzene ring of KSI-6666. Moreover, *in vivo* study suggested that this pseudoirreversible inhibition is responsible for the persistent activity of KSI-6666. These findings will contribute to the rational design of potent S1PR1 antagonists for the treatment of inflammatory disorders.

Introduction

Sphingosine 1-phosphate (S1P), which is a metabolite of membrane sphingolipids, regulates cell migration, inflammatory response, angiogenesis and neurogenesis[1]. Physiological functions of S1P are mediated by its binding to five subtypes of the sphingosine 1-phosphate receptor, which belongs to the G protein-coupled receptor (GPCR) family[1]. Among S1P receptors, S1P receptor 1 (S1PR1) is critical for lymphocyte trafficking and progression of immune and inflammatory responses. Binding of S1P to S1PR1 causes activation of Gi, a subfamily of guanine nucleotide-binding proteins, leading to the inhibition of adenylate cyclase-induced cAMP synthesis and subsequent activation of several intracellular signaling pathways. The binding also induces the phosphorylation-dependent recruitment of β-arrestin and subsequent cellular internalization of the ligand–receptor complex.

Given its roles in lymphocyte trafficking, S1PR1 is a target for the treatment of autoimmune and inflammatory diseases. FTY720, also named fingolimod, is a first-generation S1PR1 modulator that was approved as a drug for treatment of multiple sclerosis (MS)[2,3]. Mechanistically, FTY720 is phosphorylated *in vivo* into FTY720-phosphate (FTY720-P), which serves as a functional antagonist of S1P receptors including S1PR1.[2] Thus, binding of FTY720-P activates S1PR1 signaling, leading to the internalization and subsequent degradation of S1PR1. As a result, administration of FTY720 causes long-lasting inhibition of S1PR1 signaling through the depletion of cell surface S1PR1. However, in addition to lymphocyte trafficking, S1PR1 controls the function of cardiac cells[4]. Therefore, agonistic activity of FTY720 provokes bradycardia[4-6], a temporal reduction of the heart rate, as an undesirable side-effect.

Competitive antagonists of S1PR1 would not cause side effects ascribed to the agonistic activity. W146, which has a similar structure to FTY720-P, was the first competitive antagonist of S1PR1[7,8]. However, the *in vivo* efficacy of W146 is poor and transient[8]. In addition to FTY720-based S1PR1 modulators such as W146, structurally distinct S1PR1 modulators have been developed[9]. Among them, NIBR-0213 is a competitive antagonist of S1PR1 that displayed persistent efficacy *in vivo*[10]. Further development of potent S1PR1 antagonists may offer more effective methods of treatment for autoimmune diseases such as MS.

The efficacy of a drug is affected by its residence time in the target protein in addition to its equilibrium binding affinity and pharmacokinetics[11-13]. Therefore, the determination of drug–protein binding kinetics is useful for predicting drug efficacy[14]. Some studies have shown that antagonists with slow dissociation rates cause pseudoirreversible inhibition of targets with insurmountable antagonism[15-17], which is defined as the inability of the pre-bound antagonist to be completely replaced by a competing ligand in nonequilibrium conditions[18]. Drugs with pseudoirreversible inhibition properties show persistent *in vivo* efficacies because of their long residence time[19].

Computational approaches can be used to investigate the binding behavior of drugs to target proteins[20-26]. Molecular dynamics simulations can be used to evaluate binding and release pathways, kinetics, and thermodynamics[27]. Metadynamics (MetaD), which is a sampling method enhanced by escaping local free energy minima[24,28], and other approaches[24] are available for detailed description of binding and unbinding kinetics[20,22,23,25]. In addition to these force field-based calculations, quantum mechanical methods such as the fragment molecular orbital (FMO) method have been applied for *ab initio* characterization of drug interactions with proteins[29,30]. However, the linking of these computational estimations for binding and dissociation kinetics to practical *in vitro* and *in vivo* efficacies of a drug remains a challenge.

In this study, we aimed to find a potent S1PR1 antagonist with persistent efficacy and to elucidate the structural basis underlying its persistency. We discovered a S1PR1 antagonist, KSI-6666, that showed potent and persistent efficacy *in vivo*. A combination of MetaD simulation, FMO calculation, and experimental verification suggests that KSI-6666 is a pseudoirreversible inhibitor because of its hindered dissociation from S1PR1. The pseudoirreversible inhibition requires the interaction between a methionine residue in S1PR1 and an aromatic ring moiety of KSI-6666.

Results

KSI-6666 is a potent and selective antagonist of S1PR1 and suppresses intestinal inflammation

We started developing new competitive antagonists of S1PR1 from the lead compound **1** (Fig. 1a), which is similar to a compound previously discovered by Novartis company. For the optimization of compound **1**, we performed *in vitro* screening to evaluate the antagonistic effects of compounds on GTP binding to Gi in S1PR1-expressing human embryonic kidney (HEK293) cells induced by an S1PR1 agonist made by Merck (Merck S1PR1 agonist)[31] (Fig. 1a). In addition, the reduction in blood leukocyte count induced by inhibition of S1PR1 after the oral administration of compounds was evaluated in *in vivo* screening in rats (Fig. 1b). The selection identified a novel S1PR1 antagonist *(RS)*-KSI-6666 (hereafter referred to as KSI-6666 for simplicity), a racemic form with a half-maximal inhibitory concentration (IC$_{50}$) of 6.4 nM in GTP binding assay. Reduction in blood leukocyte count caused by KSI-6666 was observed at 48 h after administration (Fig 1b), and thus it may be more persistent than NIBR-0213 since its response duration was reported as 24 h at 30 mg/kg administration[10]. Consequently, we concluded that KSI-6666 is a potent antagonist with persistent efficacy.

We further confirmed that KSI-6666 had no agonistic activity on S1PR1 signaling. Binding of an agonist to S1PR1 results in the activation of Gi and receptor internalization[1]. Consistently, FTY720-P and the Merck S1PR1 agonist led to an increment in the GTP binding to Gi and a decrease of surface S1PR1 on HEK293 cells (Fig. 1c and d). In contrast, treatment with KSI-6666 caused a decrease in the GTP binding activity and an increment in the surface amount of S1PR1 even in the absence of agonists (Fig. 1c and d). These data indicated that KSI-6666 has no agonistic activity and indeed has inverse agonistic activity, which is defined as inhibition of spontaneous receptor activation[32], thereby suppressing the constitutive activation of S1PR1. In the case of W146, inverse agonistic activity was detected at only high concentrations (Fig. 1d).

We next tested the antagonistic activity of KSI-6666 toward ligand binding to S1PR1 in other assays. S1PR1 activation increases Ca^{2+} mobilization in cells over-expressing Gα15, a subunit

of G protein alpha. KSI-6666 competitively inhibited the agonist-dependent Ca^{2+} mobilization

of S1PR1, but not for other S1PRs (Fig. 1e and Extended Data Fig. 1). KSI-6666 inhibited

mouse and rat S1PR1 with similar inhibitory activity as human S1PR1, demonstrating its cross-

species efficacy (Fig. 1e and Extended Data Fig. 1). Furthermore, the agonist-dependent

internalization of S1PR1 on HEK293 cells was antagonized by KSI-6666 in a dose-dependent

manner (Fig. 1f). *In vivo*, in guinea pig, pretreatment with KSI-6666 completely blocked the

reduction of heart rate induced by administration of FTY720 (Fig. 1g). These data suggest that

KSI-6666 is a selective and competitive antagonist of S1PR1 in mammals, and therefore would

not cause bradycardia as a side effect.

S1PR1 signaling is critical for lymphocyte trafficking, and therefore the inhibition of S1PR1

impairs lymphocyte migration from lymphoid organs to the bloodstream, thereby causing

lymphopenia[1]. In Transwell assays *in vitro*, the addition of KSI-6666 inhibited the motility of

murine splenocytes induced by S1P (Fig. 1h). Consistently, flow cytometric analysis showed

that the numbers of CD4-, CD8-, and CD19-positive cells in mouse blood were significantly

reduced 4 h after intravenous injection of KSI-6666 (Fig. 1i). In contrast, injection of W146

caused a slight reduction of B cell number in the blood at 1 h, but not at 4 h (Fig. 1i). In addition

to rodents, the administration of KSI-6666 effectively diminished blood lymphocyte numbers in

cynomolgus monkeys, confirming efficacy in primates. The data suggest that, compared with

W146, KSI-6666 has a potent and persistent effect in suppressing lymphocyte trafficking in

rodents and primates.

In addition to the lymphocyte trafficking, S1PR1 modulators have a common effect on the

endothelial barrier[7]. Consistently, we observed that the repetitive oral administration of KSI-

6666 caused vascular hyperpermeability-related effects such as edema in the lungs (Extended

Data Fig. 2 and Supplementary Table 1), further supporting that KSI-6666 can inhibit the

functions of S1PR1 *in vivo*.

We further considered whether KSI-6666 suppresses inflammation in pathological rodent

models. Studies in a T-cell transfer colitis model (Fig. 2a) and an oxazolone-induced colitis

model (Fig. 2b) revealed that administration of KSI-6666 significantly suppressed the induced

intestinal inflammation, comparable with the effect of KRP-203, a functional agonistic inhibitor

of S1PR1[33]. These data suggest that KSI-6666 is a potent and persistent S1PR1 antagonist with

anti-inflammatory activity.

Molecular dynamics simulation reveals a slow dissociation of KSI-6666 from S1PR1

The structure of KSI-6666 is largely divisible into a zwitterionic part consisting of a carboxylic

acid and an aliphatic tertiary amine, and a hydrophobic moiety consisting of two aromatic rings

with bulky substituents; one chiral carbon atom in the linker is also between the two aromatic

rings (marked with an asterisk in Fig. 1a). Among the enantiomers, the S1PR1 inhibitory

activity of the *(R)-* KSI-6666 was more potent than that of the *(S)-* KSI-6666 (Fig 3a and b).

This suggests that the spatial arrangement of the distal benzene ring may be important for the

potent binding and inhibitory activity toward S1PR1.

To gain more insight into the structural basis of the activity of KSI-6666, we first performed

molecular docking analysis for the binding of *(R)-*KSI-6666 with S1PR1. Co-crystallographic

data for the complex of W146 and S1PR1 (PDB: 3v2y)[34] were used for the docking study of

*(R)-*KSI-6666. The pose of *(R)-*KSI-6666 with the lowest binding free energy was selected from

among the docking complexes, and compared with the crystallographic structure of W146 (Fig.

3c). Overlay of the pose of *(R)-*KSI-6666 with the W146 structure suggested that the two

compounds would adopt a similar pose in the ligand-binding pocket. Thus, a polar group (a

carboxylic acid in *(R)-*KSI-6666 and a phosphate in W146) and a benzene ring (the internal

benzene ring of *(R)-*KSI-6666 and the benzene ring of W146) overlapped, and the end of the

hydrophobic moiety (CF_3– in *(R)-*KSI-6666 and CH_3– in W146) was in a similar position for the

two inhibitors. The minimum binding free energy to S1PR1 was estimated to be −11.79

kcal/mol for *(R)-*KSI-6666 and −8.83 kcal/mol for W146, supporting stable binding of *(R)-*KSI-

6666.

To detail the interaction between *(R)-*KSI-6666 and S1PR1, the FMO method was performed for

the docking complexes, and the interfragment interaction energies (IFIEs) were estimated. Thus,

S1PR1 (amino acids 15–314) was fragmented into units containing a single amino acid, and the

interaction energies between each unit and *(R)*-KSI-6666 were evaluated by molecular orbital

calculations. Moreover, pair interaction energy decomposition analysis (PIEDA) in FMO

calculation divides the IFIE into four energy components: electrostatic (ES), exchange repulsion

(EX), charge-transfer and mixing term (CT+mix), and dispersion (DI) energies that in part

contribute to hydrophobic interactions. The FMO method suggested strong ES interactions of

the zwitterionic moiety of *(R)*-KSI-6666 with residues Arg120, Tyr29, Glu121, and Lys34 of

S1PR1 (Fig. 3d). Residues Phe125, Asn101, Leu297, Trp269, Phe210, Leu272, Phe273, and

Tyr98 contribute to the interaction with *(R)*-KSI-6666 through DI energy (Fig. 3d). Thus, FMO

calculation supported that hydrophobic interaction due to DI energy, along with ES energy, is

critical for the binding of *(R)*-KSI-6666 to S1PR1.

The dissociation kinetics of an antagonist from a target molecule are one of the critical

determinants for its therapeutic effect[11-13]. Because KSI-6666 showed persistent efficacy

compared with W146, we next performed MetaD simulation, which is used for biasing toward

dissociation by applying external forces, to explore and compare the dissociation process from

S1PR1 between *(R)*-KSI-6666 and W146. In total, 10 simulation runs for 100 ns each were

performed for each compound. A plot of binding free energy profiles (kcal/mol) against the

distance of *(R)*-KSI-6666 (or W146) from the binding site in enhanced sampling using MetaD

simulation suggested that a stable binding state of *(R)*-KSI-6666, but not of W146, may exist

around 2 □ away from the docking position (Fig. 3e). Moreover, to evaluate the dissociation

process, the root-mean-square distance (RMSD) of *(R)*-KSI-6666 and W146 from the docking

position was plotted against the simulation time. Assuming that the ligand was dissociated from

S1PR1 when the RMSD was >30 □, dissociation of W146 occurred in 7 of 10 runs by 100 ns

from the start of the simulation (Fig. 3f), but dissociation of *(R)*-KSI-6666 occurred in only 4 of

10 runs. Furthermore, in four runs for *(R)*-KSI-6666, the RMSD converged at around 2 □ after

100 ns, which is consistent with the binding free energy profiles (Fig. 3e). Overall, MetaD

simulation suggested that *(R)*-KSI-6666 is more resistant than W146 to dissociation from

S1PR1.

To verify the findings from MetaD simulation, binding and dissociation properties of KSI-6666

were determined experimentally by the functional reversibility assay[35] (Fig. 3g). After S1PR1-

expressing cells were pretreated with KSI-6666 at various concentrations, serially diluted

amounts of the agonistic FTY720-P were added (Fig. 3g). The replacement of KSI-6666 with

agonist FTY720-P on S1PR1 was monitored via the level of cellular surface S1PR1 (Fig. 3g).

As described above, in the absence of the agonist, the addition of KSI-6666 caused inverse

agonistic activity (labeled "w/o" in Fig. 3g). As expected, the addition of FTY720-P or S1P

ligand caused the replacement of KSI-6666 in a dose-dependent manner. Of note, however, the

maximum agonist effect (E_{max}) did not reach 100% when the pretreatment was with a high

concentration of KSI-6666 (Fig. 3g). Furthermore, E_{max} decreased with increasing

concentrations of pretreated KSI-6666 (Fig. 3g right). This phenomenon occurred for both

agonists, FTY720-P and the natural ligand S1P. Moreover, the increment of GTP binding

activity on agonist addition did not reach the maximum in the presence of a high concentration

of KSI-6666 (Fig. 3g). This indicates that the replacement of KSI-6666 by the agonists was not

complete in the assay. In contrast, when W146 was used in the pretreatment, the agonistic

responses reached the maximum level, suggesting the complete replacement of W146 by these

agonists. Consequently, these data suggest that KSI-6666 is an insurmountable antagonist with

pseudoirreversible inhibitory activity, which is consistent with the hindered dissociation of KSI-

6666 predicted by MetaD simulation.

242   Residue Met124 of S1PR1 is critical for the pseudoirreversible inhibition by KSI-6666

We next sought to determine the interaction responsible for the pseudoirreversible inhibition of

S1PR1 by KSI-6666. When the MetaD trajectories of *(R)*-KSI-6666 were visually inspected,

(R)-KSI-6666 appeared to intimately interact with hydrophobic residue Met124 of S1PR1,

whereas W146 did not (Fig. 4a and Supplementary Videos 1 and 2). To confirm this observation,

100 snapshots from each MetaD trajectory were selected at equal intervals and analyzed by

using the protein–ligand interaction fingerprint (PLIF) tool implemented in Molecular

Operating Environment (MOE) software, in which the interaction between a compound and

each amino acid residue in a protein is represented as a fingerprint bit according to its

interaction type. The frequency of bit entry for *(R)*-KSI-6666 relative to W146 was estimated.

Data suggested that, among the amino acids that *(R)*-KSI-6666 contacts during the dissociation

process, Met124 was the amino acid with the largest difference in contact frequency between

(R)-KSI-6666 and W146, with *(R)*-KSI-6666 making more frequent surface contact with

Met124 than doing W146 (Fig. 4b). The PLIF analysis also showed that arene interaction and

hydrogen bonding of the side chain, but neither ionic nor other types of hydrogen bonding,

contribute to the interaction of *(R)*-KSI-6666 with Met124, suggesting a major involvement of

hydrophobic interaction in the contact of *(R)*-KSI-6666 with Met124. Overall, the MetaD

simulation suggested that the slow dissociation of *(R)*-KSI-6666 from S1PR1 depends on its

contact with Met124, which may result in the pseudoirreversible inhibition of this inhibitor.

We experimentally verified the significance of Met124 for the interaction predicted by the

MetaD analysis. HEK293 cells expressing mutant S1PR1 where Met124 was replaced by Val

(Val124) or Leu (Leu124) were established (Extended Data Fig. 3). Functional reversibility

assays by measuring the internalization of cell surface S1PR1 showed that, in the Val124 mutant,

pre-bound KSI-6666 was almost completely replaced by the added agonist (Fig. 4c), suggesting

that the replacement of Met124 with Val abolished the pseudoirreversible inhibition by KSI-

6666. The insurmountable inhibitory activity of KSI-6666 in agonist-induced cAMP production

was also significantly impaired in the Val124 mutant, although the effect of the mutation was

partial (Fig. 4d) as compared with assay monitoring of the S1PR1 amount (Fig. 4c). This may

be attributable to the shorter incubation time in the cAMP production assay. Compared with the

mutation to Val, the influence of the replacement of Met124 with Leu was mild (Fig. 4c), which

may be consistent with the prediction from MetaD that hydrophobic interaction with the side

chain of Met124 contributes to the pseudoirreversible inhibition by KSI-6666.

Met124 is critical for the pseudoirreversible inhibition, and therefore we next focused on the

structural basis of the interaction of Met124 with *(R)*-KSI-6666 in MetaD simulation. Further

visible inspection of the MetaD trajectories suggested that Met124 may interact with the distal

benzene ring of *(R)*-KSI-6666, which has trifluorocarbon (CF_3-) and chloride ($Cl-$) substituents,

rather than with the zwitterionic end of the ligand (Fig. 4a and Supplementary Video 1). To

clarify this observation, the distance from Met124 (the γ-carbon atom) to the carbon atom of the

CF_3- group was compared with that from Met124 to a hydroxyl oxygen atom of the carboxylic

acid in the zwitterionic moiety (hereafter referred to as O1) for the all selected poses in the

MetaD trajectories (Fig. 5a). In the original structure determined by the docking study (Fig. 3c),

the distances from Met124 were 10.1 Å for CF_3– and 9.3 Å for O1. A correlation plot of the two

distances in the simulation suggested that two major poses were present (named "gate A" and

"gate B," respectively, see Fig. 5a) besides scattered unbound poses. Gate A has a median

Met124–CF_3 distance of 9.42 Å and a median Met124–O1 distance of 9.97 Å, which is close to

the values in the docking structure (yellow asterisk in gate A of Fig. 5a, right). Gate B has a

median Met124–CF_3 distance of 6.27 Å and a median Met124–O1 distance of 12.95 Å. This

suggests that, in the dissociation process of *(R)*-KSI-6666, there is a metastable binding state in

which the distal benzene ring in *(R)*-KSI-6666 possessing CF_3– and Cl– substituents becomes

close to Met124.

To address if this metastable binding state requires the interaction of Met124 with *(R)*-KSI-6666,

we performed the equivalent MetaD simulation for the Val124 mutant. Notably, the frequency

of the poses in gate B decreased significantly in the MetaD simulation for the Val124 mutant

(Fig. 5a). Overall, a metastable binding state (hereafter referred to as the "Met pocket") is

generated by the interaction of *(R)*-KSI-6666 with Met124, and its formation is correlated with

the slow dissociation of *(R)*-KSI-6666 from S1PR1.

We next evaluated the time-dependent transition of *(R)*-KSI-6666 binding poses by plotting the

distance between the CF_3– group and Met124 in each run versus time. After starting the MetaD

simulation from the docking position, *(R)*-KSI-6666 quickly moved to the Met pocket in seven

runs (Runs 1, 3, 4, 7, 8, 9, and 10; Fig. 5b). Subsequently, *(R)*-KSI-6666 dissociated from the

Met pocket in three runs (Run 3, 8, and 9); moved to another binding pocket close to the starting

position (hereafter referred as to the "deep pocket") in two runs (Runs 7 and 10); and stayed in

the Met pocket in one run (Run 4). In Run 1, although the distal benzene ring moved out of the

pocket, *(R)*-KSI-6666 was still bound to S1PR1. In the other three runs (Runs 2, 5 and 6), *(R)*-

KSI-6666 was located in the deep pocket soon after starting the simulation. Thereafter, *(R)*-KSI-

6666 stayed in the deep pocket in two of the runs (Runs 5 and 6). Finally, in Run 2, *(R)*-KSI-

6666 dissociated from the deep pocket. As expected, for the Val124 mutant, *(R)*-KSI-6666

stayed in the deep pocket after starting the simulation and dissociated without entering the Met

pocket, except in Run 8, in which it may have been temporarily located in the Met pocket from

around 30 to 50 ns (Supplementary Video 3). Overall, these data suggest that Met124 is

involved in the formation of the Met pocket, a metastable binding pocket for *(R)*-KSI-6666 that

acts as a "gatekeeper" to limit the dissociation of *(R)*-KSI-6666 from S1PR1.

FMO calculation suggests a contribution of dispersion energy to the interaction between
KSI-6666 and Met124 during the dissociation process

To further characterize the Met pocket in S1PR1, the FMO method was applied to estimate the

IFIEs of the complexes composed of S1PR1 and *(R)*-KSI-6666 predicted in the MetaD

simulations. Because of computational limitations, we selected eight complexes in which *(R)*-

KSI-6666 adopted some typical poses interacting with Met124: seven poses from Run 7

(snapshot indexes 601, 618, 628, 633, 646, 661, and 696), which would allow monitoring of the

time-dependent change in the interaction energies during the dissociation, and one pose from

Run 3 (index 201) (Supplementary Table 2). As shown in the distance plot (Fig. 6a), poses in

indexes 601, 661, and 696 were positioned in the deep pocket (gate A in Fig 6a), and those in

indexes 201, 618, 628, 633, and 646 were in the Met pocket (gate B in Fig. 6a). The FMO

method was applied to these eight complexes to determine IFIEs between *(R)*-KSI-6666 and

amino acid residues in S1PR1. On the basis of IFIE values, we performed estimation of Pearson

correlation coefficients, clustering and principal component analyses among these eight

complexes (Fig. 6b). Interestingly, the poses in the Met pocket (indexes 201, 618, 628, 633, and

646) appeared to be further separated into two groups depending on the IFIEs: the first group

included poses in indexes 201 and 618 (early stage in the dissociation), and the second group

included poses in indexes 623, 628, and 646 (late stage in the dissociation). In addition, the

poses in the deep pocket were also separated into two: a pose in the early stage of the

dissociation (index 601), and poses in the late stage of the dissociation (indexes 661 and 696).

The total IFIE between *(R)*-KSI-6666 and Met124 was lowest in the group of poses in the Met

pocket, especially in the early stage (indexes 201 and 618), suggesting that the interaction

between Met124 and *(R)*-KSI-6666 would be enhanced in the Met pocket. Moreover, PIEDA

showed that, among DI, CT, and ES energies, the DI energy contributed most to the interaction

between *(R)*-KSI-6666 and Met124 (Fig. 6d), which may explain the involvement of

hydrophobic interaction in the binding of *(R)*-KSI-6666 and Met124.

The pseudoirreversible inhibition contributes to the persistent efficacy of KSI-6666

The distal benzene ring of *(R)*-KSI-6666 may interact with Met124 in S1PR1 during the dissociation process, and therefore we speculated that a change in the electronic state of the benzene ring may influence the dissociation of the ligand from S1PR1. We then tested the contribution of the substituents on the benzene ring (i.e., CF_3 and Cl, highlighted in Fig. 7a) to the IFIEs between *(R)*-KSI-6666 and Met124. Thus, FMO calculation was re-run for the original complexes with a modification in which the two substituents (CF_3– and Cl–) on the distal benzene ring in *(R)*-KSI-6666 were replaced by hydrogen atoms while the poses of *(R)*-KSI-6666 in S1PR1 were maintained. The calculated IFIE values suggested that the interaction in the Met pocket was weakened (Fig. 7b), whereas that in the deep pocket was increased, in the modified complex (Fig. 7b). In addition, PIEDA calculation suggested that the destabilizing effect in the modified complex was mainly ascribable to DI energy, which reflects the hydrophobic interaction. Consequently, this finding may be consistent with the prediction from the PLIF analysis of MetaD for the involvement of hydrophobic interaction between Met124 and *(R)*-KSI-6666. These data suggest that the CF_3– and Cl– substituents of the distal benzene ring are involved in the binding of *(R)*-KSI-6666 in the Met pocket of S1PR1 by stabilizing the interaction between *(R)*-KSI-6666 and Met124.

We experimentally tested if replacing substituents on the benzene ring with hydrogen atoms influenced the dissociation properties of KSI-6666. Notably, derivative compounds of KSI-6666 lacking the CF_3– substituent (compounds **4** and **5** in Fig. 7a) were almost completely replaced with increased addition of FTY720-P in the functional reversibility assay monitoring the amount of surface S1PR1, suggesting no pseudoirreversible inhibitory activity for these compounds (Fig. 7c), which is consistent with predictions from the FMO calculation. We further tested whether the *in vivo* efficacy of orally administered compounds **4** and **5** was altered compared with that of KSI-6666 by evaluation of their effects on blood leukocyte number. Notably, in contrast to KSI-6666, the reduction of blood leukocytes recovered to the basal level by 32 h after the administration of compound **4** or **5** (Fig. 7c), even though the plasma concentrations of KSI-6666 and compound **4** were comparable (Extended Data Fig. 4), indicating that replacement of the CF_3– group with a hydrogen atom impaired the persistency of the efficacy of KSI-6666. Overall, the data suggest that pseudoirreversible inhibition elicits the persistent efficacy of KSI-

379   6666 as an S1PR1 antagonist.

Discussion

Slower dissociation of antagonists from receptors elicits longer and more potent inhibition of target molecules *in vivo*[36-39]. For instance, Aprepitant[19], a pseudoirreversible NK1 receptor antagonist used as an antiemetic, potently and persistently inhibited the NK1 receptor *in vivo*. In contrast, the inhibitory effect of a surmountable antagonist CP-99,994 was weaker and lasted for a short period. Our data suggest that KSI-6666 is a new pseudoirreversible inhibitor of S1PR1. Consistently, KSI-6666 showed persistent efficacy *in vivo*. In contrast, W146 appeared not to be a pseudoirreversible inhibitor. In addition to its poor pharmacokinetics, the transient effect of W146 *in vivo* may be in part ascribed to this dissociation property of W146. Our data are consistent with the concept that the slow dissociation of drugs benefits their efficacy[36-39].

The two-step induced fit model may explain the slow dissociation of KSI-6666 from S1PR1. In this model, stable binding of a drug with its target occurs via a two-step mechanism: in the first step, binding to the target forms an encounter complex; then, in the second step, a conformational change of the complex occurs and strengthens the drug binding to the target. In binding via the two-step induced fit mechanism, the residence time of a drug–target complex may be considerably augmented. MetaD simulation of KSI-6666 in the dissociation process from S1PR1 suggested the presence of another stable binding pose requiring residue Met124 (i.e., the "Met pocket") in addition to the pose close to the original binding pose (i.e., the "deep pocket"). Of note, the replacement of Met124 with Val abolished not only the slow dissociation in the simulation but also the pseudoirreversible inhibition by KSI-6666 *in vitro*. Therefore, KSI-6666 possibly binds to S1PR1 via a two-step induced fit mechanism; it binds to the Met pocket in the first step, and then forms another stable binding structure in the deep pocket, resulting in pseudoirreversible inhibition of S1PR1.

A previous study showed that replacement of Met124 with Lys caused a severe reduction of S1PR1 activity in cell culture. This finding suggests that the hydrophobicity of Met124 is critical for the formation of the binding pocket for the S1PR1 ligand. Our study showed that the replacement of Met124 with Val or Leu did not practically affect the activity of FTY720-P, but severely impaired the slow dissociation of KSI-6666. Thus, Met124 may play roles not only in

the formation of a hydrophobic "wall" for the binding pocket but also as a "gatekeeper" for the

binding and unbinding processes of S1PR1 modulators.

The benzene ring moiety in KSI-6666 appears to be critical for the interaction with Met124.

S1PR1 modulators can be divided into two types: FTY720-based S1PR1 modulators, which

have a flexible lipophilic chain, and structurally divergent ones, most of which have benzene

rings[9]. The S1PR1 modulators with benzene rings may use interaction with Met124 to form

stable interactions, which results in slow dissociation. Future testing of this hypothesis may help

in the design of new S1PR1 antagonists.

Functional antagonists of S1PR1, such as FTY720, have been used for the treatment of MS for

more than a decade. In addition to MS, new applications, such as treatment of ulcerative colitis,

have been proposed for the S1PR1 agonists, ozanimod and etrasimod[40,41]. In the present study,

KSI-6666, a competitive antagonist, showed an efficacy comparable to functional antagonists in

rodent colitis models. Accordingly, competitive S1PR1 antagonists with pseudoirreversible

inhibition, such as KSI-6666, could potentially be applicable for the treatment of these

inflammatory diseases in future.

Computational prediction of drug binding behavior and free energy may be used widely in the

development of new drugs in the future. One important issue is to address if computational

simulation can predict the actual behavior and efficacy of drugs. In this study, we used

molecular dynamics and quantum mechanics methods to explain the structural basis of the

pseudoirreversible inhibition of S1PR1 by KSI-6666, and validated the computational

predictions in experiments using cell culture and animals. This knowledge regarding the

correlation between drug efficiency and the structural basis of the drug dissociation kinetics

may help in the design of effective S1PR1 antagonists to suppress inflammatory diseases. In

such cases, computer simulation would be helpful for predicting the structural basis of drug

binding and unbinding kinetics in addition to equilibrium-binding structures.

Materials and Methods

Chemicals

KSI-6666, *(R)*-KSI-6666, *(S)*-KSI-6666, compounds **1**, **2**, **3**, **4**, and **5**, the Merck S1PR1 agonist, W146, and KRP-203 were synthesized by Kissei Pharmaceutical Co., Ltd. Detailed synthetic schemes and synthetic procedures are presented in the Supplemental Experimental Procedures.

Animals

BALB/cA mice (CLEA Japan and Charles River Laboratories Japan), BALB/cCr mice (Japan SLC), C.B-17 SCID mice (Charles River Laboratories Japan), Sprague-Dawley (SD) rats (Japan SLC), and Hartley guinea pigs (Japan SLC) were maintained in standard controlled conditions with a 12-h lighting cycle and access to chow and water *ad libitum*. Experiments were performed in accordance with the protocols approved by the Laboratory Animal Committee of Kissei Pharmaceutical Co., Ltd. or the Guidelines of the Institutional Animal Care and Use Committee of RIKEN, Yokohama Branch (2018-075). Cynomolgus monkeys (Del Mundo Trading and SICONBREC Inc.) were maintained in standard controlled conditions with a 12-h lighting cycle and access to chow from 9:00 to 13:00 and water *ad libitum*, and experiments were performed by Ina Research Inc. according to protocols approved by the Institutional Animal Care and Use Committee. In total, seven male monkeys were used for the study.

Compound-induced reduction in blood leukocyte count in rats

Compounds dissolved in the vehicle, which consisted of 5%–10% dimethyl sulfoxide (Wako) and 5%–10% Wellsolve (Celeste) mixed in distilled water (Otsuka), were orally administered to 6–16-week-old male SD rats. Blood samples were chronologically collected in EDTA-2K-containing blood collection tubes (BD) under anesthesia before and after compound administration. Leukocytes in blood were counted by an automated hematology analyzer KX-21NV (Sysmex). In some experiments, plasma fractions were isolated and concentrations of compounds were measured by liquid chromatography-mass spectrometry (Q Exactive, Thermo Fisher Scientific).

Induction of lymphopenia in cynomolgus monkeys

Three-to-five-year-old male cynomolgus monkeys were used. The test compound dissolved in

vehicle consisting of 0.5% methyl cellulose 400 solution (Wako) and 0.01 M hydrochloric acid

(Wako) mixed in distilled water (Otsuka) was orally administered to monkeys. Blood samples

were chronologically collected in EDTA-2K-containing blood collection tubes while the

monkey was awake, before and after compound administration. Lymphocytes in blood were

counted by a using hematology analyzer ADVIA120 (Siemens).

GTP binding assay

Membranes were prepared from HEK293 cells stably transfected with human S1PR1. GTP

binding assay was performed using a GTP Gi Binding Assay Kit (Cisbio) according to the

manufacturer's protocol. In the antagonist assay, serial dilutions of the test compound and 10

nM Merck S1PR1 selective agonist were coincubated with 1.25 μg/well of cell membranes

suspended in the stimulation buffer supplied by the manufacturer containing 50 mM $MgCl_2$ and

2 μM GDP, followed by the addition of GTP Eu Cryptate (donor) and d2 antibody (acceptor).

Membranes were incubated overnight at room temperature and the fluorescence signals from the

donor (620 nm) and the acceptor (665 nm) were read using a PHERAstar FSX microplate reader

(BMG Labtech). The fluorescence ratio (signal 665 nm/signal 629 nm$\times 10^4$) was calculated. The

half-maximal inhibitory concentration (IC_{50}) values were calculated by nonlinear regression

analysis of the concentration–response curve in GraphPad Prism 6 software. In assays to

evaluate the effect of the test compound alone, membranes were treated with serial dilutions of

the compound and were subjected to the same operations as noted above. In functional

reversibility assays, serial dilutions of the test compound and serial dilutions of FTY720-P were

co-incubated with membranes followed by the same operations as noted above. The

fluorescence ratio obtained from the response to 10 μM FTY720-P alone was calculated as

100% response ($E_{max} = 100\%$), and that in the condition without FTY720-P or the test

compound was defined as 0%. The maximum value of the response to FTY720-P in cotreatment

with the test compound (calculated by nonlinear regression analysis of the concentration–

response curve in GraphPad Prism 6) was regarded as the E_{max} value.

Receptor internalization assay

HEK293 cells transiently transfected with pBiT3.1-N in which the coding sequence of human

S1PR1 was inserted (pBiT3.1-N-WT-S1PR1), resulting in the expression of S1PR1 with an

HiBiT-tag at the *N*-terminus, were prepared using Lipofectamine 2000 (Thermo Fisher

Scientific). In the functional reversibility assay, serial dilutions of the test compound were

preincubated with cells in culture medium consisting of Dulbecco's Modified Eagle's Medium

(Wako) supplemented with 10% fetal bovine serum (Biological Industries) for 10 min at 37°C,

followed by the induction of receptor internalization by the addition of serial dilutions of

FTY720-P (Cayman) or S1P (Cayman). Following 60-min incubation at 37°C, cell surface

levels of HiBiT-tagged human S1PR1 were monitored using a Nano Glo HiBiT Extracellular

Detection System (Promega) according to the manufacturer's protocol. The surface level of

receptors, represented by the luminescence, was read by using a PHERAstar FSX microplate

reader. The luminesce induced by 10 µM FTY720-P or 100 µM S1P alone was calculated as

100% response ($E_{max} = 100\%$), and that in the condition without agonist or the test compound

was defined as 0%. The maximum value of the response to FTY720-P or S1P following

pretreatment of cells with the test compound (calculated by nonlinear regression analysis of the

concentration–response curve in GraphPad Prism 6) was regarded as the E_{max} value. In

functional reversibility assays with cells expressing mutant S1PR1, pBiT3.1-N with insertion of

the mutated human S1PR1 sequence where the original codon for Met124 (ATG) was replaced

by GTG or CTG, leading to the expression of M124V (pBiT3.1-N-M124V-S1PR1) or M124V

(pBiT3.1-N-M124L-S1PR1), respectively, was transfected into cells. In assays to evaluate the

effect of the test compound alone, cells in culture medium were treated with the compound for

60 min at 37°C and the surface S1PR1 levels were evaluated as noted above. In assays to

evaluate the antagonistic effect of the test compound, cells in the culture medium were treated

with serial dilutions of the compound for 10 min at 37°C, followed by the addition of 10 nM

FTY720-P.

Ca^{2+} mobilization assay

CHO-K1 cells transiently transfected with human S1PR2 or S1PR3, or cells transiently co-

transfected with human S1PR1, S1PR4, S1PR5, mouse S1PR1, or rat S1PR1 and human Gα15,

were prepared using FuGENE HD (Promega). Ca^{2+} mobilization assay using a Fluo-4 Direct

Calcium Assay Kit (Thermo Fisher Scientific) was carried out according to the manufacturer's

protocol. Following the pretreatment of Fluo4-loaded cells with serial dilutions of the test

compound in an assay buffer consisting of Hanks' Balanced Salt Solution (Thermo Fisher

Scientific) and 20 mM HEPES (Thermo Fisher Scientific) supplemented with 0.1% bovine

serum albumin (Sigma-Aldrich) and 2.5 mM probenecid (Thermo Fisher Scientific) for 15 min,

Ca^{2+} mobilization was induced by 10 nM Merck S1PR1 selective agonist for cells expressing

S1PR1, or by 100 nM S1P (Cayman) for human S1PR2-, S1PR3-, S1PR4-, or S1PR5-

transfected cells. The increase in cellular calcium level was evaluated using an FDSS 7000 plate

reader (Hamamatsu Photonics). IC_{50} values were calculated by nonlinear regression analysis of

the concentration–response curve in GraphPad Prism 6 software.

cAMP assay

HEK293 cells transiently transfected with pBiT3.1-N-WT-S1PR1 or pBiT3.1-N-M124V-S1PR1

were prepared using Lipofectamine 2000. Cells were pretreated with serial dilutions of the test

compound in assay buffer consisting of Hanks' Balanced Salt Solution, 20 mM HEPES

(Thermo Fisher Scientific), and 0.5 mM 3-isobutyl-1-methylxanthine (Wako) for 10 min,

followed by the addition of serial dilutions of FTY720-P (Cayman). Following incubation for 10

min, 5 µM forskolin (Wako) was added for 20 min. Cellular cAMP levels were examined using

a LANCE Ultra cAMP Kit (PerkinElmer) according to the manufacturer's protocol, and the

time-resolved fluorescence resonance energy transfer (TR-FRET) emission (665 nm) was read

by using a PHERAstar FSX microplate reader. All procedures were performed at room

temperature. The TR-FRET fluorescence obtained from the response to 1 µM FTY720-P alone

was calculated as 100% response ($E_{max} = 100\%$) and that in the condition without FTY720-P or

the test compound was defined as 0%. The highest value of the response to FTY720-P in cells

co-treated with the test compound (calculated by nonlinear regression analysis of the

concentration–response curve in GraphPad Prism 6) was regarded as the E_{max} value.

Effect on heart rate in guinea pigs

Six-to-seven-week-old male Hartley guinea pigs anesthetized by intraperitoneal injection of 35

mg/kg pentobarbital sodium (Kyoritsu Seiyaku) and 2.5 mg/kg of butorphanol tartrate (Meiji

Seika Pharma) were subjected to tracheotomy after tracheary incision. The thorax was incised,

and the carotid artery was catheterized with a PE90 catheter (BD) connected to a blood pressure

amplifier (Nihon Kohden) and heart rate counter (Nihon Kohden) via a DTXPlus Transducer

(Merit Medical). Signals were monitored and recorded by using a PowerLab 8/35 instrument

(AD Instruments) and LabChart software (AD Instruments). KSI-6666 (30 mg/kg) dissolved in

dimethyl sulfoxide was intravenously administered to the animals, followed by intravenous

injection of 0.3 mg/kg fingolimod hydrochloride (LC Laboratories) dissolved in dimethyl

sulfoxide 15 min after the administration of KSI-6666. The average heart rate counted for 1 min

at each time point was used for data analysis.

Chemotaxis assay

Twelve-week-old female BALB/cA mice were sacrificed, and spleens were collected.

Splenocytes were prepared after the removal of red blood cells using RBC Lysis Buffer

(BioLegend). Cells (2×10^7 cells/mL) in assay medium consisting of phenol red-free RPMI-1640

(Thermo Fisher Scientific) supplemented with 0.1% fatty acid-free bovine serum albumin

(Sigma) were incubated with serial dilutions of the test compound for 5 min, and 100 µL cell

suspensions were transferred to Transwell inserts (Corning) loaded in 24-well culture plates

(BD) filled with 600 µL assay medium per well and 20 nM S1P. Cells were incubated for 3 h at

37°C in a 5% CO_2 incubator, followed by removal of the inserts. Migrated cells in plates were

stained with CyQuant (Thermo Fisher Scientific) for 1 h at 37°C in the CO_2 incubator and

fluorescence intensity, correlated with cell number, was measured by using a PHERAstar FSX

microplate reader.

Flow cytometric analysis of lymphocytes

Compounds were dissolved in 10 mM Na_2CO_3 and 40% (2-hydroxypropyl)-β-cyclodextrin and

intraperitoneally administrated to 8-week-old female mice (10 mg/kg). At 1, 2 and 4 h after the

administration, whole blood was collected into tubes pretreated with EDTA. After the lysis of

erythrocytes by using RBC Lysis Buffer, cell suspensions were stained with anti-mouse CD19

(BD 557399), CD4-PECy7 (BioLegend 100528), and CD8-APCCy7 (BioLegend 100714)

antibodies after blocking by anti-CD16/32 antibodies (BioLegend 101302). Dead cells were

excluded via 7-aminoactinomycin D staining.

Naïve CD4+CD45Rbhi T cell adoptive transfer colitis model

Eight-to-nine-week-old female BALB/cA mice were sacrificed, and spleens were collected.

Splenocytes were prepared after the removal of red blood cells using RBC Lysis Buffer. Naïve T

cells were isolated by negative selection using a Mouse Naive T Cell $CD4^+/CD62L^+/CD44^{low}$

Column Kit (R&D Systems). Naïve CD4 T cells were treated with fluorescein isothiocyanate-

labeled (FITC) anti-mouse CD45RB antibody (BioLegend), followed by washing. Cells treated

with anti-FITC MicroBeads (Miltenyi Biotec) were magnetically separated via a MACS column

(Miltenyi Biotec) and $CD4^+CD45RB^{high}$ T cells were thus purified. Cell suspension (6×10^5

cell/mL) was intraperitoneally transferred into 8–10-week-old female C.B-17 SCID mice (500

µL/mouse). Two weeks after the transfer, oral administration of compounds twice daily was

started and lasted for 14–16 days. The day after the last administration, the DAI score,

consisting of the sum (0–12 points) of the grades for three observations [weight loss (%; none:

0; 0–5: 1; 5–10: 2; 10–20: 3; >20: 4); stool consistency (normal: 0; loose stool: 2; diarrhea: 4);

and hematochezia (absence: 0; presence: 4)] was evaluated. Mice were sacrificed and colons

were collected, weighed and further evaluations were performed. The histological score,

consisting of the sum (0–9 points) of the grades for three observations [decrease in goblet cells

(0–3); inflammatory cell infiltration (0–3); and mucosal thickening (0–3)], was determined. For

mRNA expression analysis, total RNA was extracted from colons by using an RNeasy Mini Kit

(QIAGEN). cDNA was reverse transcribed by using a PrimeScript RT Reagent Kit Perfect Real

Time (TaKaRa Bio Inc.). Real-time PCR was performed using a 7500 Fast Real-Time PCR

System (Applied Biosystems). SYBR Premix Ex *Taq* Perfect Real Time (TaKaRa Bio Inc.) for

S100a9 with a primer set (MA058882) obtained from TaKaRa Bio Inc. and Premix Ex *Taq*

Perfect Real Time (TaKaRa Bio Inc.) for 18S rRNA with TaqMan Ribosomal RNA Control

Reagents (Applied Biosystems) were used for the reaction. The ΔΔCt method was applied to

calculate the relative expression level of target mRNA using the 18S rRNA gene as the

reference.

Oxazolone-induced colitis model

A total of 150 µL of 3% 4-ethoxymethylene-2-phenyl-2-oxazolin-5-one (oxazolone; Sigma-

Aldrich) dissolved in 100% ethanol (Wako) was applied on shaved back skins of 9–10-week-old

male BALB/cCr mice for sensitization. One week after sensitization, 100 µL of 1% oxazolone

dissolved in 50% ethanol oxazolone was intrarectally administered to the mice. On the same day,

test compounds were orally administered to mice twice (before and after sensitization). Mice

were sacrificed 24 h after sensitization, and colons were collected and weighed. Homogenates

of colons were prepared using TisseLyser (QIAGEN) and the amount of interleukin (IL)-4

protein was evaluated using a Mouse IL-4 Quantikine ELISA Kit (R&D Systems).

Molecular docking analysis

Our approach to docking the *(R)*-KSI-6666 molecule to the potential binding sites in S1PR1

used two steps: (1) structure and ligand preparation for docking, and (2) docking of the *(R)*-KSI-

6666 molecule. An initial structure of human S1PR1 was modeled from the X-ray structure of

S1PR1 with the W146 molecule excluding the *Escherichia* virus T4 structure (PDB: 3v2y)[34] and

refined for docking simulations using the Protein Preparation Wizard Script within Maestro[42]

(Schrödinger LLC, New York, NY, USA). For the *(R)*-KSI-6666 molecule, ionization and

energy minimization were performed by the OPLS3 force field in the LigPrep Script in Maestro.

A minimized structure of *(R)*-KSI-6666 was employed as the input structure for docking

simulation. To account for both ligand and receptor flexibility in the first step, the Glide

'Induced Fit Docking (IFD)' protocol[43] (Schrödinger LLC) was used, followed by iteratively

combining rigid receptor docking using Glide and protein remodeling by side chain searching

and minimization using Prime (Schrödinger LLC) techniques. We generated 20 initial

orientations of *(R)*-KSI-6666 in a grid box defined by the center of the co-crystalized ligand

(W146) using Glide docking [standard precision (SP) mode]. Next, we applied the soften-

potential docking options, which involved scaling the van der Waals radii by 0.5 for receptor

and *(R)*-KSI-6666 atoms. In the protein remodeling stage, all residues within a 5.0-Å radius of

each initial docked *(R)*-KSI-6666 were refined using Prime. *(R)*-KSI-6666 was then redocked

into the refined receptor structure using Glide in SP mode. Finally, the best pose for *(R)*-KSI-

6666 was rescored according to GlideScore (SP mode). The Docking score of co-crystalized

ligand W146 was calculated using the score-in-place procedure of Glide SP mode.

Molecular dynamics and metadynamics simulation

Binding models of *(R)*-KSI-6666 and W146 to the structure of human S1PR1 were examined in

100-ns MetaD simulations using Desmond version 2.3 (Schrödinger LLC). MetaD simulation is

a widely used enhanced sampling method that allows the sampling of free energy landscapes. In

this simulation, we defined the biasing collective variables as the distance between the center of

mass of the ligands and the ligand-binding residues (Tyr29, Lys34, Tyr98, Asn101, Ser105,

Gly106, Thr109, Trp117, Arg120, Glu121, Met124, Phe125, Leu128, Ser192, Val194, Leu195,

Pro196, Tyr198, Ile203, Cys206, Thr207, Phe210, Trp269, Leu272, Phe273, Leu276, Glu294,

Leu297, Val298, Ala300 and Val301 for *(R)*-KSI-6666; Tyr29, Lys34, Tyr98, Asn101, Ser105,

Gly106, Thr109, Trp117, Arg120, Glu121, Met124, Phe125, Leu128, Thr193, Val194, Leu195,

Pro196, Ile203, Cys206, Thr207, Phe210, Trp269, Leu272, Phe273, Leu276, Glu294, Leu297,

Ala300 and Val301 for W146). The initial Gaussian hill height was set at 0.3. The OPLS3e force

field[44,45] was used for the simulations. The human S1PR1 structure was placed in a large

palmitoyl-oleoyl-phosphatidylcholine (POPC) bilayer, and transferable intermolecular potential

3 point (TIP3P) water molecules were solvated with 0.15 M NaCl. After minimization and

relaxation of the model, the production molecular dynamics phase was performed for 10

independent 100-ns simulations in the isothermal–isobaric (NPT) ensemble at 300 K and

1 bar using a Langevin thermostat. The long-range electrostatic interactions were computed

using the Smooth Particle Mesh Ewald method. All system setups were performed using

Maestro. The MetaD procedure for *(R)*-KSI-6666 in the M124V variant was identical to that for

wild-type S1PR1. The model of M124V S1PR1 was constructed using Maestro.

Protein–ligand interaction fingerprint analysis

PLIF analyses from MetaD trajectories were performed using the PLIF modules of MOE

2020.0901 (Chemical Computing Group Inc., Montreal, Quebec, Canada). Energy-based

contacts, including side chain hydrogen bonds (donor or acceptor), backbone hydrogen bonds

(donor or acceptor), solvent hydrogen bonds (donor or acceptor), ionic interactions, metal

binding interactions, and π interactions, and surface contact fingerprints, were employed. After

the PLIFs had been generated based on 1000 snapshot structures (every 1 ns in 10 independent

100-ns simulations) for each ligand, the probability of each bit occurring for each residue for

(R)-KSI-6666 compared with W146 was calculated as Qb.

$$Q_b = \frac{C_b}{C_0} \cdot 2\left(P_b - \frac{1}{2}\right)$$

where Cb is the number of snapshots within the set that have the fingerprint bit set, and C0 is

the total number of snapshots. Pb is the relative probability of the fingerprint bit entry over the

snapshots for *(R)*-KSI-6666 compared with that for W146 if it contains the bit. Qb takes values

from −1 to 1. A higher positive value indicates a higher probability of the bit occurring for *(R)*-

KSI-6666.

Fragment molecular orbital method

The initial complex structure of the *(R)*-KSI-6666-bound S1PR1 and selected structures from

trajectories in MetaD simulation for *(R)*-KSI-6666 dissociation were minimized using the

Impref module (Schrödinger LLC) with the OPLS3e force field. To evaluate the contribution of

the two substituents (CF_3 and Cl) on the distal benzene ring of *(R)*-KSI-6666, an additional

complex in which the original *(R)*-KSI-6666 was modified by replacing both substituents with

hydrogen atoms was also generated. FMO calculation[46] was then performed using the ABINIT-

MP program[47] at the FMO2-MP2/6-31G(d) level and IFIE with PIEDA was calculated using

MIZUHO/BioStation viewer.

Conflict of Interest

YM, YOh, TS, YY, KO, HI, AY, MH, and YOk are employees of Kissei Pharmaceutical Co., Ltd.

Acknowledgement

This work was supported by the Research Support Project for Life Science and Drug Discovery

(Basis for Supporting Innovative Drug Discovery and Life Science Research [BINDS]) from

AMED (grant number JP22ama121029j0001 to T.H.), Grants-in-Aid for Scientific Research

from JSPS (17H04038, 17K08622, 20K07332, and 20H03441 to T.A. and N.A.), Grants-in-Aid

for Scientific Research in Innovative Areas from MEXT (18H04989 and 19H04821 to T.A. and

N.A.) and by CREST from the Japan Science and Technology Agency (JPMJCR2011 to T.A.).

The authors declare no competing financial interests. We thank Edanz (https://jp.edanz.com/ac)

for editing a draft of this manuscript.

Author contributions

YM, YOh, TS, MM, TM, and NA performed the experiments. KO and HI synthesized the

chemicals. AY performed LC-MS analysis. MH and YOk performed histological analysis. YY

and TH performed *in silico* calculations. YM, YY, TI, KH, MH, TH, and TA analyzed the

MetaD and FMO data. YM and TA designed the study and wrote the paper.

	Figure legends

Figure 1. KSI-6666 persistently antagonizes S1PR1.

a. Lead optimization of S1PR1 antagonists with long-lasting efficacy. Structures of compounds

synthesized for sequential optimization are shown. The IC_{50} value in GTP binding assay using

the membrane fraction of HEK293 cells stably expressing human S1PR1 was calculated. S1PR1

was activated by 10 nM Merck S1PR1 agonist. Data were obtained from three independent

experiments (mean ± s.e.m.). The chiral carbon atom in KSI-6666 is indicated by an asterisk.

b. Effect of KSI-6666 and compounds **2** and **3** on leukocyte number in blood of rats. Each

compound was orally administered to Sprague Dawley (SD) rats at 3 mg/kg. Leukocytes in the

blood were counted at the indicated times. Leukocyte number was normalized as a percentage

from the baseline count before administration (mean ± s.e.m., $n = 4$–6 /group). Statistical

analyses compared with KSI-6666 were performed by two-way analysis of variance (ANOVA).

c. Effect of KSI-6666, FTY720-phosphate (FTY720-P), and Merck S1PR1 agonist on GTP

binding activity in HEK293 cells expressing S1PR1. Data were obtained from three independent

experiments (mean ± s.e.m.).

d. Effect of KSI-6666, FTY720-P, and W146 on the level of S1PR1 on the surface of HEK293

cells. Data were obtained from three independent experiments (mean ± s.e.m.).

e. Effect of KSI-6666 on Ca^{2+} mobilization induced by the activation of S1P receptor subtypes.

Ca^{2+} mobilization was induced by 10 nM S1PR1 agonist (for S1PR1 of human, mouse, or rat),

or by 100 nM S1P (for human S1PR2, S1PR3, S1PR4, or S1PR5) in CHO-K1 cells expressing

S1P receptors. Gα15 was co-expressed with S1PR1, S1PR4, or S1PR5 to enable signaling from

these receptors to phospholipase C. The IC_{50} value of the test compound was calculated. Data

are the mean ± s.e.m. of three independent experiments.

f. Effect of KSI-6666 pretreatment on downregulation of surface S1PR1 induced by FTY720-P.

HEK293 cells expressing S1PR1 were treated with KSI-6666 at the indicated concentrations.

w/o indicates no treatment with KSI-6666. Data were obtained from three independent

experiments (mean ± s.e.m.).

g. Effect of KSI-6666 pretreatment on the FTY720-P-induced reduction in the heart rate of

guinea pigs. Heart rate over time was measured for 45 min in total (mean ± s.e.m., $n = 3$ /group).

Statistical analysis was performed by two-way ANOVA.

h. Effect of KSI-6666 pretreatment on the FTY720-P-induced chemotactic activity of

splenocytes. Transwell assay was performed using murine splenocytes. Data are the mean ±

s.e.m. of the results from three mice. Statistical analysis *vs*. treatment with S1P alone was

performed by one-way ANOVA followed by Dunnett's test.

i. Effect of KSI-6666 and W146 on blood lymphocytes in mice. After each drug was

intravenously administrated (10 mg/kg), frequency and cell numbers of $CD4^+$, $CD8^+$, and

$CD19^+$ cells were analyzed by flow cytometry. $n = 6$ each for 1 and 4 h, $n = 9$ for the control (0

h). P values (two-tailed Student's t test) were indicated in the figure.

j. Effect of KSI-6666 on lymphocyte number in blood of cynomolgus monkeys. KSI-6666 was

orally administered to monkeys at 3 mg/kg. Lymphocytes in the blood were counted at the

indicated times. Lymphocyte number was normalized as a percentage from the baseline count

before administration (mean ± s.e.m., $n = 3$ /group). Statistical analysis *vs*. vehicle was

performed by two-way ANOVA.

Figure 2. KSI-6666 is a potent anti-inflammatory agent.

a. Therapeutic efficacies of KSI-6666 (15 mg/kg b.i.d.) and KRP-203 (1.5 mg/kg b.i.d.) in T-cell

transfer colitis model. Isolated naive $CD4^+CD45RB^{high}$ T cells from BALB/cA mice were

injected intraperitoneally into SCID mice and colitis was elicited. Two weeks after the transfer,

twice daily oral administration of each compound was started and lasted for 14–16 days.

Animals were sacrificed on the day after the final administration. Colon weight, disease activity

index (DAI) score, histological score, and *S100a9* mRNA expression level relative to ribosomal

RNA expression were evaluated in each group. Statistical analysis was performed by either

Student's *t*-test or Welch's *t*-test according to F-test results (mean ± s.e.m., $n = 15–35$ /group).

b. A representative example of the histology.

c. Efficacies of KSI-6666 (15 mg/kg b.i.d.) and KRP-203 (1.5 mg/kg b.i.d.) in oxazolone-

induced colitis model. Oxazolone (3% solution) was applied on back skins of BALB/cA mice

on day 1 (sensitization), followed by intrarectal administration of 1% oxazolone on day 8

(challenge), and colitis was elicited. Each test compound was orally administered twice on day 8

and animals were sacrificed on day 9. Colon weight and interleukin (IL)-4 protein amount in the

colon in each group were evaluated and the percentage of controls (colitis-induced mice to

which vehicle was administered) was calculated. Statistical analysis was performed by either

Student's *t*-test or Welch's *t*-test according to F-test results (mean ± s.e.m., $n = 9–19$).

Figure 3. KSI-6666 is a pseudoirreversible inhibitor.

a. Difference between KSI-6666 enantiomers in inhibitory activities against GTP binding to

S1PR1 induced by 10 nM Merck S1PR1 agonist. Data were obtained from three independent

experiments. Results are expressed as the mean ± s.e.m. Magenta line, *(RS)*-KSI-6666; cyan line,

(R)-KSI-6666; green line, *(S)*-KSI-6666.

b. Effect of *(RS)*-KSI-6666, *(R)*-KSI-6666, and *(S)*-KSI-6666 on leukocyte number in blood of

rats. Each compound was orally administered to SD rats at 3 mg/kg. Leukocytes in the blood

were counted at the indicated times. Leukocyte number was normalized as a percentage from

the baseline count before the administration (mean ± s.e.m., $n = 3$ /group).

c. Docking pose of *(R)*-KSI-6666 (left) and crystallographic pose of W146 (middle) with S1PR1.

In the merge figure, *(R)*-KSI-6666 and W146 are depicted in green and gray, respectively.

d. Interfragment interaction energies (IFIEs) between *(R)*-KSI-6666 and amino acids in S1PR1.

IFIEs below −2 kcal/mol are shown in the figure. The results of pair interaction energy

decomposition analysis are shown as green bars for electrostatic energies (ES), red bars for

dispersion energies (DI), orange bars for charge transfer and mix energies (CT+ mix), and blue

bars for exclusion energies (EX).

e. Plots of binding free energy profiles (kcal/mol) against the distance between the center of

mass of the ligand-binding residues and the ligand in metadynamics (MetaD) trajectories for

(R)-KSI-6666 (left) and W146 (right) in the dissociation pathway from S1PR1. The results from

all 10 simulation runs are plotted for each compound.

f. Plots of ligand root-mean-square deviation against simulation time in the MetaD trajectories

for *(R)*-KSI-6666 (left) and W146 (right) in the dissociation pathways from S1PR1. The results

from all 10 runs are plotted for each compound.

g. Functional reversibility of KSI-6666 and W146 examined by agonist-induced S1PR1

internalization assay. HEK293 cells expressing HiBiT-tagged human S1PR1 were treated with

each compound and concentration-dependent S1PR1 internalization induced by FTY720-P

(upper left) or S1P (lower left), represented by luciferase activity, was measured. Calculated

E_{max} values for the FTY720-P (upper right) and S1P (lower right)-induced response on treatment

with each compound at different concentrations. Data for the upper and lower graphs were

obtained from five and three independent experiments, respectively. Statistical analysis was

performed by one-way ANOVA, followed by Tukey's test for KSI-6666. Results are expressed

812 as the mean ± s.e.m.

813 **h**. Functional reversibility of KSI-6666 and W146 examined by agonist-induced GTP binding

814 assay (upper). Calculated E_{max} values for the FTY720-P-induced response on treatment with

815 each compound at different concentrations (lower). Data were obtained from eight independent

816 experiments. Statistical analysis was performed by one-way ANOVA, followed by Tukey's test

817 for KSI-6666 ($n = 4$–8 /condition of antagonist treatment, mean ± s.e.m.).

819 **Figure 4**. Met124 of S1PR1 is critical for pseudoirreversible inhibition by KSI-6666.

820 **a**. Typical poses of *(R)*-KSI-6666 in the ligand binding pocket of S1PR1 in MetaD simulation.

821 Residues Thr207, Phe210, and Phe273, representing the inner pocket, are shown in line style,

822 and Met124 is shown in CPK style.

823 **b**. Significant protein–ligand interactions of *(R)*-KSI-6666 relative to W146 in protein–ligand

824 interaction fingerprint (PLIF) analysis. The interaction between the compound and S1PR1

825 during MetaD simulation was represented in a fingerprint scheme by the PLIF tool. The

826 interaction between the compound and each amino acid residue of S1PR1 was evaluated in two

827 main categories: potential-energy based contacts and surface contacts. When an interaction

828 above a threshold was observed for the residue, a fingerprint bit representing the interaction was

829 assigned. Hydrogen bonding, ionic interaction, and π interaction (arene attraction) were

830 identified as potential contacts with energies >0.5 kcal/mol and assigned as fingerprint bits in

831 this analysis. For hydrogen bonding, each amino acid residue was broken down into side chain

832 (SC) and backbone (BB) categories, and each was classified as either electron donor or acceptor.

833 When a surface contact with area >15 $\square^2$ was observed, a fingerprint bit was assigned. The

834 probability of each bit occurring for each residue for *(R)*-KSI-6666 compared with W146 was

835 calculated as Qb, where Qb takes values from −1 to 1. A higher positive value indicates a higher

836 probability of the bit occurring for *(R)*-KSI-6666. The columns show the value of Qb for each

837 residue. The fingerprint bits shown are total surface contacts (left); and arene attraction

838 (representing π interaction), ionic attraction, and BB hydrogen acceptor, BB hydrogen donor,

839 SC hydrogen acceptor, and SC hydrogen donor interactions with *(R)*-KSI-6666 (right, different

840 interactions shown in different colors).

841 **c**. Functional reversibility of KSI-6666 with Val124 and Leu124 mutants of S1PR1 examined by

842 the S1PR1 internalization assay. HEK293 cells expressing HiBiT-tagged human S1PR1 Val124

mutant (M124V, left) or Leu124 mutant (M125L, middle) were treated with KSI-6666 and

concentration-dependent receptor internalization stimulated by FTY720-P was measured.

Calculated E_{max} values for the FTY720-P-induced response on treatment with KSI-6666 at

different concentrations are plotted for wild-type S1PR1 and the two mutants (right). The plot

for the wild-type shown in Fig. 3g is reiterated here as a control. Data for the Val124 and

Leu124 mutants were obtained from four and three independent experiments, respectively.

Statistical analyses *vs*. wild-type were performed by two-way ANOVA for the Val124 and

Leu124 mutants. Results are expressed as the mean ± s.e.m.

d. cAMP assay evaluating the functional reversibility of KSI-6666 in wild-type S1PR1 and the

Val124 mutant. HEK293 cells expressing HiBiT-tagged wild-type S1PR1 (left) or the Val124

mutant (middle) were treated with KSI-6666 and concentration-dependent inhibition by

FTY720-P of forskolin-induced cAMP production was measured. Calculated E_{max} values for the

FTY720-P-induced response on treatment with KSI-6666 at different concentrations are plotted

for the wild-type and the mutant (right). Data were obtained from three independent

experiments, respectively. Statistical analysis was performed by two-way ANOVA. Results are

expressed as the mean ± s.e.m.

Figure 5. Metadynamics simulation suggests the presence of a metastable binding pocket for KSI-6666 in S1PR1.

a. Plot of distance from Met124 (γ-carbon atom) to the carbon atom of the CF_3– group on the distal benzene ring of *(R)*-KSI-6666 against the distance from Met124 to the carboxylic acid (O1 atom) of *(R)*-KSI-6666 in MetaD snapshots of wild-type S1PR1 (left) and the Val124 mutant (middle). Snapshots ($n = 1000$) were selected every 1 ns from 10 independent 100-ns MetaD simulations for each form of S1PR1. The area where both distances were <25 □ is enlarged in the right-hand figure. The frequency of poses assigned to gate A (distance between Met124 or Val124 and the carbon atom of the CF_3– group 2.5–7.8 □, distance between Met124 or Val124 and the oxygen atom of the carboxylic acid 10–15 □) and gate B (distance between Met124 or Val124 and the carbon atom of the CF_3– group 7.8–12.5 □, distance between Met124 or Val124 and the oxygen atom of the carboxylic acid, 7.5–12.5 □) is shown for the wild-type (blue) and Val124 mutant (red) proteins.

b. Time-dependent changes in the distance between the carbon atom of the CF_3– group of *(R)*-KSI-6666 and the γ-carbon atom of Met124 (left) or Val124 (right) of S1PR1. The results of all runs are shown in the upper figures. The results of runs in which the binding of *(R)*-KSI-6666 lasted for at least 100 ns are extracted in the lower figures.

Figure 6. Fragment molecular orbital (FMO) calculation suggested the interaction of *(R)*-KSI-6666 and Met124 is strengthened in a metastable binding pocket.

a. Plot of distance from Met124 (γ-carbon atom) of S1PR1 to the carbon atom of the CF_3– group on the distal benzene ring of *(R)*-KSI-6666 against distance from Met124 to the carboxylic acid (O1 atom) of *(R)*-KSI-6666 corresponding to the poses selected for FMO calculation. Snapshot indexes selected are shown. The plot corresponding to the docked structure is indicated by an orange circle.

b. Pearson correlation and clustering of poses based on total interaction energy calculated by the FMO method using complexes of *(R)*-KSI-6666 and S1PR1.

c. Principal component analysis of poses based on total interaction energy calculated by the FMO method using complexes of *(R)*-KSI-6666 and S1PR1.

d. IFIEs of *(R)*-KSI-6666 with Met124 in each group of poses. The snapshot index to which each pose belongs is shown above the bar chart. The average IFIE of poses in the group is

shown.

Figure 7. Derivative compounds of KSI-6666 that lack pseudoirreversible inhibition have poor

persistent efficacy.

a. Structures of KSI-6666 and related compounds **4** and **5**.

b. Differential IFIEs of modified and original complexes. The complex of *(R)*-KSI-6666 and

Met124 was modified by replacing the two substituents on the distal benzene ring of *(R)*-KSI-

6666 (i.e., CF_3 and Cl) with hydrogen atoms. The two complexes were subjected to FMO

calculation and the differences between their IFIEs were calculated. The pose index for each

group is shown above the bar chart. The average IFIE of poses was employed.

c. Functional reversibility of compounds **4** and **5** examined by S1PR1 internalization assay.

HEK293 cells expressing HiBiT-tagged human S1PR1 were treated with compound **4** (left) or **5**

(middle) and concentration-dependent receptor internalization stimulated by FTY720-P was

measured. Calculated E_{max} values for the FTY720-P-induced response on treatment with these

compounds at different concentrations are plotted (right). The plot of KSI-6666 data shown in

Fig. 3g (upper right) is reiterated as a control. Data for compounds **4** and **5** were obtained from

three independent experiments for each. Statistical analyses *vs*. KSI-6666 were performed by

two-way ANOVA for compounds **4** and **5**. Results are expressed as the mean ± s.e.m.

d. *In vivo* efficacy of KSI-6666, and compounds **4** and **5**. Each compound was orally

administered to SD rats at the dose indicated and the efficacy, represented as the magnitude of

reduction in blood leukocyte count, was evaluated at the indicated times after administration

(left). Leukocyte number was normalized as a percentage from the baseline count before

administration. Data for a single individual belonging to the control shown in Fig. 1b are

reiterated for comparison. Statistical analyses *vs*. KSI-6666 were performed by two-way

ANOVA for compounds **4** and **5**. Results are expressed as the mean ± s.e.m. ($n = 4–8$ /group).

Corresponding plasma concentrations of KSI-6666 and compound **4** at two time points are

shown in the right-hand graph (mean ± s.e.m.).

References

1.   Cartier, A. & Hla, T. Sphingosine 1-phosphate: Lipid signaling in pathology and therapy.
Science **366**(2019).

2.   Chun, J., Kihara, Y., Jonnalagadda, D. & Blaho, V.A. Fingolimod: Lessons Learned and
New Opportunities for Treating Multiple Sclerosis and Other Disorders. *Annu Rev*
Pharmacol Toxicol **59**, 149-170 (2019).

3.   Brinkmann, V. et al. Fingolimod (FTY720): discovery and development of an oral drug
to treat multiple sclerosis. *Nat Rev Drug Discov* **9**, 883-97 (2010).

4.   Jozefczuk, E., Guzik, T.J. & Siedlinski, M. Significance of sphingosine-1-phosphate in
cardiovascular physiology and pathology. *Pharmacol Res* **156**, 104793 (2020).

5.   Camm, J., Hla, T., Bakshi, R. & Brinkmann, V. Cardiac and vascular effects of
fingolimod: mechanistic basis and clinical implications. *Am Heart J* **168**, 632-44 (2014).

6.   Fryer, R.M. et al. The clinically-tested S1P receptor agonists, FTY720 and BAF312,
demonstrate subtype-specific bradycardia (S1P(1)) and hypertension (S1P(3)) in rat.
PLoS One **7**, e52985 (2012).

7.   Sanna, M.G. et al. Enhancement of capillary leakage and restoration of lymphocyte
egress by a chiral S1P1 antagonist in vivo. *Nat Chem Biol* **2**, 434-41 (2006).

8.   Tarrason, G. et al. The sphingosine-1-phosphate receptor-1 antagonist, W146, causes
early and short-lasting peripheral blood lymphopenia in mice. *Int Immunopharmacol* **11**,
1773-9 (2011).

9.   Marciniak, A., Camp, S.M., Garcia, J.G.N. & Polt, R. An update on sphingosine-1-
phosphate receptor 1 modulators. *Bioorg Med Chem Lett* **28**, 3585-3591 (2018).

10.   Quancard, J. et al. A potent and selective S1P(1) antagonist with efficacy in
experimental autoimmune encephalomyelitis. *Chem Biol* **19**, 1142-51 (2012).

11.   Copeland, R.A., Pompliano, D.L. & Meek, T.D. Drug-target residence time and its
implications for lead optimization. *Nat Rev Drug Discov* **5**, 730-9 (2006).

12.   Copeland, R.A. The drug-target residence time model: a 10-year retrospective. *Nat Rev*
Drug Discov **15**, 87-95 (2016).

13.   Tautermann, C.S. Impact, determination and prediction of drug-receptor residence times
for GPCRs. *Curr Opin Pharmacol* **30**, 22-26 (2016).

14.   Bernetti, M., Cavalli, A. & Mollica, L. Protein-ligand (un)binding kinetics as a new

paradigm for drug discovery at the crossroad between experiments and modelling.

Medchemcomm **8**, 534-550 (2017).

15.   Sykes, D.A. et al. Extrapyramidal side effects of antipsychotics are linked to their

association kinetics at dopamine D2 receptors. *Nat Commun* **8**, 763 (2017).

16.   Sykes, D.A. et al. Fevipiprant (QAW039), a Slowly Dissociating CRTh2 Antagonist

with the Potential for Improved Clinical Efficacy. *Mol Pharmacol* **89**, 593-605 (2016).

17.   Agren, R. et al. Ligand with Two Modes of Interaction with the Dopamine D2

Receptor-An Induced-Fit Mechanism of Insurmountable Antagonism. *ACS Chem*

Neurosci **11**, 3130-3143 (2020).

18.   Vauquelin, G., Van Liefde, I., Birzbier, B.B. & Vanderheyden, P.M. New insights in

insurmountable antagonism. *Fundam Clin Pharmacol* **16**, 263-72 (2002).

19.   Hale, J.J. et al. Structural optimization affording 2-(R)-(1-(R)-3, 5-

bis(trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4- (3-oxo-1,2,4-triazol-5-

yl)methylmorpholine, a potent, orally active, long-acting morpholine acetal human NK-

1 receptor antagonist. *J Med Chem* **41**, 4607-14 (1998).

20.   Tiwary, P., Limongelli, V., Salvalaglio, M. & Parrinello, M. Kinetics of protein-ligand

unbinding: Predicting pathways, rates, and rate-limiting steps. *Proc Natl Acad Sci U S A*

112, E386-91 (2015).

21.   Buch, I., Giorgino, T. & De Fabritiis, G. Complete reconstruction of an enzyme-

inhibitor binding process by molecular dynamics simulations. *Proc Natl Acad Sci U S A*

108, 10184-9 (2011).

22.   Tiwary, P., Mondal, J. & Berne, B.J. How and when does an anticancer drug leave its

binding site? *Sci Adv* **3**, e1700014 (2017).

23.   Kokh, D.B. et al. Estimation of Drug-Target Residence Times by tau-Random

Acceleration Molecular Dynamics Simulations. *J Chem Theory Comput* **14**, 3859-3869

(2018).

24.   Cavalli, A., Spitaleri, A., Saladino, G. & Gervasio, F.L. Investigating drug-target

association and dissociation mechanisms using metadynamics-based algorithms. *Acc*

Chem Res **48**, 277-85 (2015).

25.   Callegari, D. et al. Metadynamics Simulations Distinguish Short- and Long-Residence-

Time Inhibitors of Cyclin-Dependent Kinase 8. *J Chem Inf Model* **57**, 159-169 (2017).

26.   Decherchi, S. & Cavalli, A. Thermodynamics and Kinetics of Drug-Target Binding by

981   Molecular Simulation. *Chem Rev* **120**, 12788-12833 (2020).

982   27.   Sledz, P. & Caflisch, A. Protein structure-based drug design: from docking to molecular

983   dynamics. *Curr Opin Struct Biol* **48**, 93-102 (2018).

984   28.   Laio, A. & Parrinello, M. Escaping free-energy minima. *Proc Natl Acad Sci U S A* **99**,

985   12562-6 (2002).

986   29.   Heifetz, A. et al. Analyzing GPCR-Ligand Interactions with the Fragment Molecular

987   Orbital (FMO) Method. *Methods Mol Biol* **2114**, 163-175 (2020).

988   30.   Heifetz, A., James, T., Southey, M., Bodkin, M.J. & Bromidge, S. Guiding Medicinal

989   Chemistry with Fragment Molecular Orbital (FMO) Method. *Methods Mol Biol* **2114**,

990   37-48 (2020).

991   31.   Ibrahim, M.A. et al. Discovery of a novel class of potent and orally bioavailable

992   sphingosine 1-phosphate receptor 1 antagonists. *J Med Chem* **55**, 1368-81 (2012).

993   32.   Stewart, G.D., Sexton, P.M. & Christopoulos, A. Inverse Agonists. in *Encyclopedia of*

994   *Psychopharmacology* (ed. Stolerman, I.P.) 661-667 (Springer Berlin Heidelberg, Berlin,

995   Heidelberg, 2010).

996   33.   Shimizu, H. et al. KRP-203, a novel synthetic immunosuppressant, prolongs graft

997   survival and attenuates chronic rejection in rat skin and heart allografts. *Circulation* **111**,

998   222-9 (2005).

999   34.   Hanson, M.A. et al. Crystal structure of a lipid G protein-coupled receptor. *Science* **335**,

1000   851-5 (2012).

1001   35.   Guo, D., Hillger, J.M., AP, I.J. & Heitman, L.H. Drug-target residence time--a case for

1002   G protein-coupled receptors. *Med Res Rev* **34**, 856-92 (2014).

1003   36.   Dahl, G. & Akerud, T. Pharmacokinetics and the drug-target residence time concept.

1004   *Drug Discov Today* **18**, 697-707 (2013).

1005   37.   Lu, H. & Tonge, P.J. Drug-target residence time: critical information for lead

1006   optimization. *Curr Opin Chem Biol* **14**, 467-74 (2010).

1007   38.   Guo, D., Mulder-Krieger, T., AP, I.J. & Heitman, L.H. Functional efficacy of adenosine

1008   A(2)A receptor agonists is positively correlated to their receptor residence time. *Br J*

1009   *Pharmacol* **166**, 1846-59 (2012).

1010   39.   Schneider, E.V., Bottcher, J., Huber, R., Maskos, K. & Neumann, L. Structure-kinetic

1011   relationship study of CDK8/CycC specific compounds. *Proc Natl Acad Sci U S A* **110**,

1012   8081-6 (2013).

1013   40.   Sandborn, W.J. et al. Ozanimod as Induction and Maintenance Therapy for Ulcerative

1014   Colitis. *N Engl J Med* **385**, 1280-1291 (2021).

1015   41.   Sandborn, W.J. et al. Efficacy and Safety of Etrasimod in a Phase 2 Randomized Trial

1016   of Patients With Ulcerative Colitis. *Gastroenterology* **158**, 550-561 (2020).

1017   42.   Sastry, G.M., Adzhigirey, M., Day, T., Annabhimoju, R. & Sherman, W. Protein and

1018   ligand preparation: parameters, protocols, and influence on virtual screening

1019   enrichments. *J Comput Aided Mol Des* **27**, 221-34 (2013).

1020   43.   Sherman, W., Day, T., Jacobson, M.P., Friesner, R.A. & Farid, R. Novel procedure for

1021   modeling ligand/receptor induced fit effects. *J Med Chem* **49**, 534-53 (2006).

1022   44.   Roos, K. et al. OPLS3e: Extending Force Field Coverage for Drug-Like Small

1023   Molecules. *J Chem Theory Comput* **15**, 1863-1874 (2019).

1024   45.   Kurki, M., Poso, A., Bartos, P. & Miettinen, M.S. Structure of POPC Lipid Bilayers in

1025   OPLS3e Force Field. *J Chem Inf Model* (2022).

1026   46.   Kitaura, K., Ikeo, E., Asada, T., Nakano, T. & Uebayasi, M. Fragment molecular orbital

1027   method: an approximate computational method for large molecules. *Chemical Physics*

1028   *Letters* **313**, 701-706 (1999).

1029   47.   Starikov, E.B. et al. *Modern methods for theoretical physical chemistry of biopolymers*,

1030   (Elsevier, Amsterdam ; Boston, 2006).

1031

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a

Wild-type

Val mutant

B
13.3%
1.8%

A

16.7%
28.5%

Distance of Met/Val124-O1

Distance of Met/Val124-CF3

Distance of Met124-O1

Distance of Met124-CF3

Distance of Val124-O1

Distance of Val124-CF3

Met124 (Val)

b

Wild-type

Val124 mutatnt

Distance of Met124-CF3

Time (nsec)

Distance of Val124-CF3

Time (nsec)

Figure 6

Figure 7