Title: A *de novo* long-read genome assembly is contiguous and complete enough to study paleo-
polyploidy and herbivore-induced transcriptional responses of the sacred datura plant (*Datura wrightii*)

Authors: Jay K Goldberg[1]*, Aaron Olcerst[2], Michael McKibben[1], J Daniel Hare[2], and Judith Bronstein[1]

Author for correspondence: jaykgold@arizona.edu

Running Title: Datura genome paper

Author affiliations: [1] Department of Ecology and Evolutionary Biology, University of Arizona, Tucson,
Arizona, USA; [2] Department of Entomology, University of California Riverside, Riverside, CA, USA

Abstract: The sacred datura plant (*Datura wrightii*) has been used to study plant-herbivore interactions
for decades and, due to this wealth of literature regarding it's ecological interactions, has potential as a
model system for future research on the ecological and evolutionary genomics of plant/insect
interactions. In this paper, we present a *de novo Datura wrightii* genome that we assembled using
PacBio HiFi long-reads. Our assembly is highly complete and contiguous (N50 = 179Mb, BUSCO
Complete = 97.7%). To demonstrate the utility of our assembly to future studies, we successfully used
multiple programs to test for the presence of an ancient whole genome duplication and classify the
duplicated genes identified within our assembly. We also used this assembly as the basis for a genome-
guided differential expression analysis to identify the induced responses of this plant to one of its
specialist herbivores (Coleoptera: Chrysomelidae: *Lema daturaphila*). We find over 6000 significantly
differentially expressed genes between control and herbivore-induced plants, and further that these
changes appear to be sustained and last multiple days. Overall, our results demonstrate that HiFi-only
draft plant genomes are of sufficient quality to serve as the basis for varied biological insights and
expand our knowledge of herbivore-induced plant transcriptional responses.

Introduction

Datura wrightii is a fascinating plant species with great potential as a model system for research
into the ecology and evolution of various plant traits including both physical and chemical defenses
(Hare and Walling 2006; Goldberg et al 2020), tolerance to herbivory (Bronstein et al 2009), floral
phenotypes (Elle and Hare 2002; Riffell et al 2008), perennial life history (Dupin and Smith 2018; 2019)
and ontogenetic changes in defense production throughout an individual plant's lifetime (van Dam et al
1999; Kariñho-Betancourt et al 2015). This is due to the wealth of ecological knowledge already
gathered regarding this common plant and some of its specialist herbivores (Goldberg et al. 2021; Zhang
et al 2022). Furthermore, field-based methodologies have already provided some insights into the
evolutionary processes maintaining a trichome dimorphism in naturally occurring populations (Hare et al
2007; Goldberg et al 2020). As such, the publication of a reference genome for this species would
accelerate research into both the molecular mechanisms governing the expression of plant traits and
their evolutionary trajectories in a rapidly changing environment.

In this paper, we present a *de novo* genome assembly for the sacred datura plant (*Datura
wrightii*) generated using highly accurate long-reads (PacBio HiFi; lang et al 2020). As a result, our
assembly is highly complete and contiguous when compared to recent assemblies of closely related
species (Rajewski et al 2021; De-la-Cruz et al 2021; Xu et al 2017) To demonstrate the utility of our
genome to future research, we conducted two downstream analyses with it: 1) an assessment of the
duplication history of coding regions with our assembly and 2) a differential expression study examining
herbivore-induced transcriptional changes. Together, they show that our *de novo* assembly will have
prolonged value to studies of molecular mechanisms of plant responses to stress and comparative
analyses of flowering plant evolution.

Gene and whole genome duplications (WGDs) have broad implications for evolutionary
processes including the generation and adaptation of novel traits (Schranz et al 2012; Selmecki et al
2015). WGDs have occurred numerous times throughout the diversification of angiosperms (Landis et al
2018); and these 'ancient' WGDs have been shown to underlie the arms-race dynamic occurring
between plants and herbivores (Edger et al 2015). There is widespread evidence of an ancient hexploidy
that occurred early in the evolution of the nightshade family (Solanaceae; Xu et al 2017; Landis et al
2018). As such we used two different analyses – one of which uses only contiguous collinear gene blocks
– to test for evidence of this ancient WGD in our genome.

We further use our assembly to conduct a differential expression study into the induced
responses of *D. wrightii* to one of its specialist herbivores, the three-lined potato beetle (*Lema
daturaphila*; Goldberg et al. 2021; Zhang et al 2022). This analysis utilized older data, generated roughly
a decade ago (Olcerst 2017). Using both pairwise and time course analyses, we identified 1000s of genes
that are significantly differentially expressed when plants are under attack by one of their closest natural
enemies and further show that many of them remain upregulated for multiple days.

Methods

Initial sample collection and preparation, and sequencing

Datura wrightii seeds were originally collected in June 2016 from a wild plant growing at the
intersection of Portal Rd. and Foothills Rd. in Portal, Arizona (see

https://www.inaturalist.org/observations/156947070 for more details). A cohort of seeds from this
collection were germinated in May 2021. Leaf tissue collected for genome sequencing originated from a
single individual, whereas the bud and root tissue for annotation were collected from other individuals
in this cohort. All samples were flash frozen and ground under liquid nitrogen immediately before
storage at -80C.

Nucleic acid extractions and sequencing strategies

DNA extraction and SMRT bell library preparation, and sequencing (PacBio HiFi, Pacific
Bioscience, San Francisco, CA, USA) were performed by the Arizona Genomics Institute (University of
Arizona, Tucson, AZ, USA). High molecular weight DNA was extracted from young leaves using the
protocol of Doyle and Doyle (1987) with minor modifications. Flash-frozen young leaves were ground to
a fine powder in a frozen mortar with liquid nitrogen followed by very gentle extraction in 2% CTAB
buffer (that included proteinase K, PVP-40 and beta-mercaptoethanol) for 30min to 1h at 50 °C. After
centrifugation, the supernatant was gently extracted twice with 24:1 chloroform:isoamyl alcohol. Take
out the upper phase and add $1/10^{th}$ volume 3 M NaAc, gently mixed, and DNA precipitated with iso-
propanol. DNA was collected by centrifugation, washed with 70% ethanol, air dried for 20 minutes and
dissolved thoroughly in EB at room temperature followed by RNAse treatment. DNA purity was
measured with Nanodrop, DNA concentration measured with Qubit HS kit (Invitrogen) and DNA size was
validated by Femto Pulse System (Agilent).

DNA was sheared to an appropriate size range (10-20 kb) using Megaruptor 3 (Diagenode)
followed by Ampure bead purification. The sequencing library was constructed following manufacturers
protocols using SMRTbell Prep kit 3.0. The final library was size selected on a Pippin HT (Sage Science)
using S1 marker with a 10-25 kb size selection. The recovered final library was quantified with Qubit HS
kit (Invitrogen) and size checked on Femto Pulse System (Agilent). The final library was prepared for
sequencing with PacBio Sequel II Sequencing kit 2.0 for HiFi library, loaded on 8M SMRT cells, and
sequenced in CCS mode for 30 hours.

RNA was extracted from root and bud tissue using a ZYMO (Irvine, CA, USA) direct-zol miniprep
kit (Cat. # R2050) and sequenced using NovaSeq (Illumina, San Diego, CA, USA) paired-end (150bp)
sequencing performed by Novogene (Sacramento, CA, USA). RNA for differential expression was
extracted and sequenced as described in Olcerst (2017).

Genome assembly/annotation and gene duplication analyses

CCS output (ie: HiFi reads; 3952061 reads; 65.26Gb total; mean length = 16524) were assembled
using hifiasm-0.16.0 (Cheng et al. 2021) with default settings. Repetitive elements were identified using
RepeatModeler v2.0.1 and softmasked using RepeatMasker v4.1.0 (Tarailo-Graovac and Chen 2009).
Structural annotation was carried out using the funannotate pipeline v1.8.14 (Palmer and Stajich 2019)
using all mRNA-seq libraries (N=2, 150bp paired-end; N=36 50bp single-end) as evidence during the
'funannotate train' step. Protein sequences from a publicly available *D. wrightii* transcriptome
(downloaded from medplantrnaseq.com) were used as evidence in the 'funanotate predict' step.
Default settings were used, except for max intron length which was set to 10000bp. Functional
annotation was performed in InterProScan v5.57-90.0 (Jones et al 2014). Genome quality was examined
using Bandage v0.8.1 (Wick et al 2015), BUSCO v5.1.3 (Figure 1; Seppey et al 2019), and the blobtools
v1.1 pipeline (Figure 2; Laetsch and Blaxter 2017) employing minimap v2-2.24 (Li 2018) for alignment

and the nt sequence database for taxonomic identification (BLAST 2.13.0). We used two programs,
DupPipe (Barker et al 2010) and McScanX (Wang et al 2012), to investigate the history of gene
duplication events that generated our observed *D. wrightii* gene content and test for signatures of
ancient WGDs. We used our *D. wrightii* transcriptome in addition to transcriptomes from *Solanum*
lycopersicum (Hosmani et al 2019), and *S. tuberosum* (The Potato Genome Consortium 2011) as a
reference protein database for DupPipe. MCscanX was run on default settings following the removal of
isoforms and Ka/Ks values were calculated using the add_kaks_to_synteny.pl script (Wang et al 2012).
Synonymous (Ks) divergence calculations for the DupPipe and MCscanX were then visualized using
matplotlib (Hunter 2007).

Differential expression analysis

Samples for differential expression analysis were collected from two experiments conducted at
UC Riverside in 2012/13 (Olcerst 2017). Both studies were conducted with the MVV6 line, a backcrossed
line originating from seeds originally collected in southern California and which induces production of
defenses in response to herbivore attack (Hare, 2007). The first consisted of three replicates of
control/*Lema*-induced leaf samples (N = 6 total RNAseq libraries) collected from greenhouse grown
plants in May 2012. The second was a time course experiment in which samples were taken at 5
timepoints following herbivory (0h [before treatment], 12h, 24h, 48h, 96h; 6 per timepoint, except 24h
which was excluded due to lack of replication). Additional details regarding growing conditions and
sample collection can be found in Olcerst (2017).

Raw RNAseq reads were aligned to the annotated reference genome and counted using STAR
(Dobin et al 2013) with the default parameters. STAR was selected for its flexible alignment parameters
(up to 10 mismatches per read, so long as total mismatches do not exceed 30% of their length), and
automatic read counting functionality when given an annotation (a .gff3 file in our case). Reads aligned
to our reference genome at an average rate of 80% per library, with a multi-mapping rate of less than
10% for all libraries. An average of 0.3% of reads from each library was removed for mapping to over 10
loci. Alignment details can be found in Table S3. Untransformed read counts were passed to DESeq2
(Love et al 2014) in R (R Core Team 2022) for statistical analysis of differential gene expression. Pairwise
analysis was conducted using only the samples collected in May 2012 (N = 6; Figure 3). Time course
dataset (N = 24) was also analyzed in DESeq2 using a likelihood ratio test (LRT) approach. Our full model
consisted of an ANCOVA design, with treatment, timepoint, and their interaction term as variables. Our
reduced model lacked the interaction term; thus, this analysis tested the significance of the interaction
term specifically (i.e. genes for which expression over time differed by treatment). Due to a lack of
replication for the 24h timepoint, it was excluded from the time course analysis. Data visualization was
performed in the DEseq2 (PCA plots), EnhancedVolcano, and pheatmap packages in R (R Core Team
2022).

Results and Discussion

Genome assembly and annotation

Our *de novo* assembly is highly complete and contiguous (N50 = 179Mb; Longest contig =
202Mb; 1144 contigs; BUSCO Complete=97.7% [Single-copy=92.9%, Duplicated=4.8%],
Fragmented=0.2%, Missing=2.1%; Figure 1). The total length of our assembly is 2.2Gb, comparable to
other *Datura* assemblies (Rajewski et al 2021; De-la-Cruz et al 2021) and making our final coverage

27.7X across the whole genome (18.1X when counting only reads that mapped back to the assembly;
Figure 1). When compared to previous *Datura* assemblies we find that our assembly contains far fewer
contigs that are orders of magnitude longer. This improvement is due entirely to our use of PacBio HiFi
reads, rather than previous generation technologies. Blobtools analysis determined that no
contamination was present in our assembly and that all contigs/reads were identified as belonging to
the Solanaceae family (Figure 1). 86.11% of the assembly is repetitive elements, primarily retroelements
such as long terminal repeats (Table S1).

Our annotation, which incorporated both mRNA and protein evidence, identified 49439 genes,
but BUSCO analysis showed that it was less complete than our assembly (Complete=85.7%, [Single-
copy=81.1%, Duplicated=4.6%], Fragmented=5.5%, Missing=8.8%; Figure 1). This is likely due to the
shortcomings of our annotation pipeline choice (funannotate) as it is known to be conservative and
fragment genes with incomplete evidence. Even so, with over 85% of BUSCOs present, our annotation
proved suitable for downstream applications.

Gene Duplication Analyses

We used two separate tools to test for the presence of ancient whole genome duplications
(WGD) within our genome. Both use a similar Ks approach, but one looks at all genes in the genome with
greater than 40% sequence similarity (DupPipe; Barker et al 2010) whereas the other looks specifically at
gene pairs within collinear blocks of genes (MCSCanX; Wang et al 2012). Both analyses provided similar
results (Figure 2) that confirm the presence of the ancient WGD that has been previously documented in
Solanceaous plant genomes (Xu et al 2017; Landis et al 2018). We further used MCScanX to classify all
the gene duplications within the largest 50 contigs of our genome (Table 1, S2). We found that dispersed
duplications are most common (N = 17951; Table 1), although it is important to note that McScanX
tends to inflate this number with improperly classified duplications. Ancient whole genome duplications
are likely to have generated a significant amount of the gene content observed in the *D. wrightii*
genome as well (N = 7714 genes; Table 1) and this number is likely underestimated.

Differential Expression analysis

Our pairwise analysis of *Lema*-induced transcriptional changes showed that control and *Lema*-
induced samples did not have separate profiles, but that induction increased the variance in plant gene
expression profiles (Figure 3A). Our analyses identified 6358 significantly differentially expressed genes
between control and *Lema*-induced plants. (Figure 3; Table S4). Roughly half of these (N = 3114) were up
regulated in response to *L. daturaphila* feeding (Figure 3B). We did not conduct any dedicated GO term
enrichment analysis, but our functional annotation identified some of these genes as playing a role in
responses to herbivory by other plant species (Table S4, S5).

Our time course analysis generated a shorter list of differentially expressed genes (N = 368;
Figure 4; Table S5), but similar results to our pairwise analysis, in terms of most genes being upregulated
in response to herbivory (Figure 4B). We did notice a strong differentiation between 0h (pre-herbivory)
and 12h post-herbivory treatments (Figure 4A), regardless of treatment type (control or herbivory),
which may be a sign of background circadian rhythms in gene expression (Xu et al 2022). Our results
show clear signs of sustained long-term herbivore-induced changes to gene expression (shown in our
48h and 96h timepoints; Figure 4B).

Concluding Remarks

In sum, we present a high quality long-read genome assembly for the sacred datura plant (*Datura wrightii*). We then analyze its gene duplication history and use it as the basis for a genome-guided analysis of herbivore-induced gene expression changes. Multiple tools supported the presence of a well-documented ancient whole-genome duplication event in *Datura wrightii*; and our analysis identified 1000s of differentially expressed genes, some of which have known functional annotations based on comparison to existing reference genomes. Together these data provide a valuable resource for plant scientists more broadly and will contribute to future studies of angiosperm evolution and plant-insect interactions.

Data Availability

Assembly and raw HiFi reads can be found on NCBI (BioProject: PRJNA966699; BioSample: SAMN34546691). Supporting datasets, which includes annotation, can be found at https://github.com/caterpillar-coevolution/Datura-wrightii-genome-project alongside scripts used for computational analyses. Raw transcriptome data will be available on NCBI GenBank (BioProject: PRJNA966699)

Acknowledgments & Funding

We thank Professor Luciano Matzkin and Mike Barker for their guidance throughout the process of working on the project. We would also like to thank the 2016 staff and volunteers of the Southwestern Research Station. Funding was provided by an NSF postdoctoral research fellowship in biology to JKG (#2010772).

Conflict of Interest

The authors declare no conflict of interest.

Literature Cited

Barker, M.S., Dlugosch, K.M., Dinh, L., Challa, R.S., Kane, N.C., King, M.G., Rieseberg, L.H., 2010. EvoPipes.net: Bioinformatic Tools for Ecological and Evolutionary Genomics. Evol Bioinform Online 6, EBO.S5861. https://doi.org/10.4137/EBO.S5861

Bronstein, J.L., Huxman, T., Horvath, B., Farabee, M., Davidowitz, G., 2009. Reproductive biology of Datura wrightii: the benefits of a herbivorous pollinator. Annals of Botany 103, 1435–1443. https://doi.org/10.1093/aob/mcp053

Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., Li, H., 2021. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. Nat Methods 18, 170–175. https://doi.org/10.1038/s41592-020-01056-5

De-la-Cruz, I.M., Hallab, A., Olivares-Pinto, U., Tapia-López, R., Velázquez-Márquez, S., Piñero, D., Oyama, K., Usadel, B., Núñez-Farfán, J., 2021. Genomic signatures of the evolution of defence against its natural enemies in the poisonous and medicinal plant Datura stramonium (Solanaceae). Sci Rep 11, 882. https://doi.org/10.1038/s41598-020-79194-1

Dupin, J., Smith, S.D., 2019. Corrigendum to: "Phylogenetics of Datureae (Solanaceae), including
description of the new genus Trompettia and re-circumscription of the tribe" [in Taxon 67: 359–375.
2018]. TAXON 68, 419–419. https://doi.org/10.1002/tax.12056

Dupin, J., Smith, S.D., 2018. Phylogenetics of Datureae (Solanaceae), including description of the new
genus Trompettia and re–circumscription of the tribe. TAXON 67, 359–375.
https://doi.org/10.12705/672.6

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras,
T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21.
https://doi.org/10.1093/bioinformatics/bts635

Doyle, J.J. and J.L. Doyle. 1987. A rapid DNA isolation procedure from small quantities of fresh leaf
tissues. Phytochem Bull. 19:11–15.

Edger, P.P., Heidel-Fischer, H.M., Bekaert, M., Rota, J., Glöckner, G., Platts, A.E., Heckel, D.G., Der, J.P.,
Wafula, E.K., Tang, M., Hofberger, J.A., Smithson, A., Hall, J.C., Blanchette, M., Bureau, T.E., Wright, S.I.,
dePamphilis, C.W., Eric Schranz, M., Barker, M.S., Conant, G.C., Wahlberg, N., Vogel, H., Pires, J.C.,
Wheat, C.W., 2015. The butterfly plant arms-race escalated by gene and genome duplications.
Proceedings of the National Academy of Sciences 112, 8362–8366.
https://doi.org/10.1073/pnas.1503926112

Elle, E., Hare, J.D., 2002. Environmentally Induced Variation in Floral Traits Affects the Mating System in
Datura wrightii. Functional Ecology 16, 79–88.

Eric Schranz, M., Mohammadin, S., Edger, P.P., 2012. Ancient whole genome duplications, novelty and
diversification: the WGD Radiation Lag-Time Model. Current Opinion in Plant Biology, Genome studies
molecular genetics 15, 147–153. https://doi.org/10.1016/j.pbi.2012.03.011

Goldberg, J.K., Lively, C.M., Sternlieb, S.R., Pintel, G., Hare, J.D., Morrissey, M.B., Delph, L.F., 2020.
Herbivore-mediated negative frequency-dependent selection underlies a trichome dimorphism in
nature. Evolution Letters 4, 83–90. https://doi.org/10.1002/evl3.157

Hare, J.D., 2007. Variation in Herbivore and Methyl Jasmonate-Induced Volatiles Among Genetic Lines of
Datura wrightii. J Chem Ecol 33, 2028–2043. https://doi.org/10.1007/s10886-007-9375-1

Hare, J.D., Elle, E., van Dam, N.M., 2003. Costs of Glandular Trichomes in Datura Wrightii: A Three-Year
Study. Evolution 57, 793–805. https://doi.org/10.1111/j.0014-3820.2003.tb00291.x

Hare, J.D., Walling, L.L., 2006. Constitutive and Jasmonate-Inducible Traits of Datura wrightii. J Chem
Ecol 32, 29–47. https://doi.org/10.1007/s10886-006-9349-8

Hosmani, P. S., Flores-Gonzalez, M., van de Geest, H., Maumus, F., Bakker, L. V., Schijlen, E., … Saha, S.,
2019. An improved de novo assembly and annotation of the tomato reference genome using single-
molecule sequencing, Hi-C proximity ligation and optical maps. Cold Spring Harbor Laboratory.
https://doi.org/10.1101/767764

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A.,
Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., Hunter,

S., 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics 30, 1236–1240.
https://doi.org/10.1093/bioinformatics/btu031

Kariñho-Betancourt, E., Agrawal, A.A., Halitschke, R., Núñez-Farfán, J., 2015. Phylogenetic correlations
among chemical and physical plant defenses change with ontogeny. New Phytologist 206, 796–806.
https://doi.org/10.1111/nph.13300

Laetsch, D.R., Blaxter, M.L., 2017. BlobTools: Interrogation of genome assemblies.
https://doi.org/10.12688/f1000research.12232.1

Landis, J.B., Soltis, D.E., Li, Z., Marx, H.E., Barker, M.S., Tank, D.C., Soltis, P.S., 2018. Impact of whole-
genome duplication events on diversification rates in angiosperms. American Journal of Botany 105,
348–363. https://doi.org/10.1002/ajb2.1060

Lang, D., Zhang, S., Ren, P., Liang, F., Sun, Z., Meng, G., Tan, Y., Li, X., Lai, Q., Han, L., Wang, D., Hu, F.,
Wang, W., Liu, S., 2020. Comparison of the two up-to-date sequencing technologies for genome
assembly: HiFi reads of Pacific Biosciences Sequel II system and ultralong reads of Oxford Nanopore.
GigaScience 9, giaa123. https://doi.org/10.1093/gigascience/giaa123

Li, H., 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34, 3094–3100.
https://doi.org/10.1093/bioinformatics/bty191

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol 15, 1–21. https://doi.org/10.1186/s13059-014-0550-8

Olcerst, A. 2017. Variation in Induced Responses of Datura wrightii to Herbivore Attack: Plasticity of
Volatile Organic Compound Emissions and Gene Expression Across Genotypes, Ontogeny, and a Single
Attack – Doctoral Disseration accessed via ProQuest:
https://www.proquest.com/openview/920bf36262446dfa983f0a67e1755bff/1?pq-
origsite=gscholar&cbl=18750

Palmer, J., Stajich, J., 2019. nextgenusfs/funannotate: funannotate v1.5.3.
https://doi.org/10.5281/zenodo.2604804

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rajewski, A., Carter-House, D., Stajich, J., Litt, A., 2021. Datura genome reveals duplications of
psychoactive alkaloid biosynthetic genes and high mutation rate following tissue culture. BMC Genomics
22, 201. https://doi.org/10.1186/s12864-021-07489-2

Riffell, J.A., Alarcón, R., Abrell, L., 2008. Floral trait associations in hawkmoth-specialized and mixed
pollination systems. Communicative & Integrative Biology 1, 6–8. https://doi.org/10.4161/cib.1.1.6350

Selmecki, A.M., Maruvka, Y.E., Richmond, P.A., Guillet, M., Shoresh, N., Sorenson, A.L., De, S., Kishony,
R., Michor, F., Dowell, R., Pellman, D., 2015. Polyploidy can drive rapid adaptation in yeast. Nature 519,
349–352. https://doi.org/10.1038/nature14187

Seppey, M., Manni, M., Zdobnov, E.M., 2019. BUSCO: Assessing Genome Assembly and Annotation
Completeness, in: Kollmar, M. (Ed.), Gene Prediction: Methods and Protocols, Methods in Molecular
Biology. Springer, New York, NY, pp. 227–245. https://doi.org/10.1007/978-1-4939-9173-0_14

Tarailo-Graovac, M., Chen, N., 2009. Using RepeatMasker to Identify Repetitive Elements in Genomic
Sequences. Current Protocols in Bioinformatics 25, 4.10.1-4.10.14.
https://doi.org/10.1002/0471250953.bi0410s25

The Potato Genome Sequencing Consortium, 2011. Genome sequence and analysis of the tuber crop
potato. *Nature* 475, 189–195. https://doi.org/10.1038/nature10158

van Dam, N., Hare, J., Elle, E., 1999. Inheritance and distribution of trichome phenotypes in Datura
wrightii. Journal of Heredity 90, 220–227. https://doi.org/10.1093/jhered/90.1.220

Wang, Y., Tang, H., DeBarry, J.D., Tan, X., Li, J., Wang, X., Lee, T., Jin, H., Marler, B., Guo, H., Kissinger,
J.C., Paterson, A.H., 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny
and collinearity. Nucleic Acids Research 40, e49. https://doi.org/10.1093/nar/gkr1293

Xu, S., Brockmöller, T., Navarro-Quezada, A., Kuhl, H., Gase, K., Ling, Z., Zhou, W., Kreitzer, C., Stanke,
M., Tang, H., Lyons, E., Pandey, P., Pandey, S.P., Timmermann, B., Gaquerel, E., Baldwin, I.T., 2017. Wild
tobacco genomes reveal the evolution of nicotine biosynthesis. Proceedings of the National Academy of
Sciences 114, 6133–6138. https://doi.org/10.1073/pnas.1700073114

Xu, X., Yuan, L., Xie, Q., 2022. The circadian clock ticks in plant stress responses. Stress Biology 2, 15.
https://doi.org/10.1007/s44154-022-00040-7

Zhang, J., Komail Raza, S.A., Wei, Z., Keesey, I.W., Parker, A.L., Feistel, F., Chen, J., Cassau, S., Fandino,
R.A., Grosse-Wilde, E., Dong, S., Kingsolver, J., Gershenzon, J., Knaden, M., Hansson, B.S., 2022.
Competing beetles attract egg laying in a hawkmoth. Current Biology 32, 861-869.e8.
https://doi.org/10.1016/j.cub.2021.12.021

Figure Legends & Table

Figure 1. (left) Bar charts showing the gene content represented in both our assembly (top bar) and our
annotation (bottom bar). (right) Blobtools coverage plot showing that 64.79% of raw reads mapped to
our final assembly, and that all mapped reads were taxonomically identified as Solanaceae, indicating
that our assembly is contaminant free. The 35.21% of raw reads that did not map to our assembly is
likely the result of the haplotig purging step of the hifiasm process, which produces a haplotype-specific
assembly.

Figure 2. Results of DupPipe (left) and MCScanX (right) showing the frequency of non-synonymous
substitions throughout the genome. DupPipe analyzes global gene content, whereas MCScanX only
examines genes within syntenic blocks. Both analyses show a large peak at Ks = 0.5, indicative of ancient
whole-genome duplication.

Figure 3. Results of pairwise differential expression analysis using only the samples collected in May
2012. (A) Results of exploratory PCA to qualitatively screen for differences between treatment groups.
There was a clear qualitative difference between the control and *Lema*-induced samples. (B) Volcano
plot showing the relationship between significance and fold-change for each gene in our *D. wrightii*
genome annotation. A change of 2/-2 is marked for reference but was not used as a cutoff for any
analysis.

Figure 4. (A) Exploratory PCA of time course data. Although our samples did not form clear clusters,
there are definitive patterns. The 48h and 96h *Lema*-induced samples cluster in the top right, whereas
the controls for these time points cluster with the 0h samples in the top left. Ranges for 0h and 12h
samples do not overlap at all, suggesting that there may also be circadian cycles occurring. (B) Heatmap
showing the fold-change of top 50 most significant genes from our time course analysis (N = 368 total
significant genes).

Table 1. Results of McScanX gene duplication classifier function. Most of the gene content is the results
of dispersed duplications, but ancient whole-genome-duplications (WGD) have also generated
thousands of genes within the *D. wrightii* genome. IDs refer to duplication identifiers in the full gene list
(Table S2).

Duplication Type	ID	Gene count
Singleton	0	8301
Dispersed	1	17951
Proximal	2	2649
Tandem	3	3556
WGD/Segmental	4	7714

Supplemental Materials

Table S1. Detailed output table from RepeatMasker.

Table S2. Detailed results of MCScanX gene duplication classifier function

Table S3. Detailed output of STAR alignments

Table S4. Complete list of significant genes from pairwise differential expression analysis

Table S5. Complete list of significant genes from time course expression analysis

Total Gene Content
(DupPipe)

Frequency

Mean Block Ks

Syntenic Blocks Only
(McScanX)

Frequency

Collinear Homologous Gene Pair Ks