Curbing zoonotic disease spread in multi-host-species systems will require integrating novel data streams and analytical approaches: evidence from a scoping review of bovine tuberculosis

Kimberly Conteddu[1*], Holly M. English[1], Andrew W. Byrne[2], Bawan Amin[1], Laura L. Griffin[1], Prabhleen Kaur[3], Virginia Morera-Pujol[1], Kilian J. Murphy[1], Michael Salter-Townshend[3], Adam F. Smith[4,5,6], Simone Ciuti[1*]

1. Laboratory of Wildlife Ecology and Behaviour, School of Biology and Environmental Science, University College Dublin, Dublin, Ireland
2. One Health Scientific Support Unit, Department of Agriculture, Food and the Marine, Dublin, Ireland
3. School of Mathematics and Statistics, University College Dublin, Dublin, Ireland
4. Department of Wildlife Ecology and Management, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany
5. The Frankfurt Zoological Society, Frankfurt, Germany
6. Department of Visitor Management and National Park Monitoring, Bavarian Forest National Park, Grafenau, Germany

*For correspondence: kimberly.conteddu@ucdconnect.ie; simone.ciuti@ucd.ie

Present address: School of Biology and Environmental Science, University College Dublin, Ireland

Table of Contents

Abstract

Background: Zoonotic diseases represent a significant societal challenge in terms of their health and economic impacts. One Health approaches to managing zoonotic diseases are becoming more prevalent, but require novel thinking, tools and cross-disciplinary collaboration. Bovine tuberculosis (bTB) is one example of a costly One Health challenge with a complex epidemiology involving human, domestic animal, wildlife and environmental factors, which require sophisticated collaborative approaches.

Objective: We undertook a scoping review of multi-host bTB epidemiology to identify recent trends in species publication focus, methodologies, scales and One Health approaches. We aimed to identify research gaps where novel research could provide insights to inform control policy, for bTB and other zoonoses.

Results: The review included 167 articles. We found different levels of research attention across episystems, with a significant proportion of the literature focusing on the badger-cattle-TB episystem, with far less attention given to the multi-host episystems of southern Africa. We found a limited number of studies focusing on management solutions and their efficacy, with very few studies looking at modelling exit strategies. Surprisingly, only a small number of studies looked at the effect of human disturbances on the spread of bTB involving wildlife hosts. Most of the studies we reviewed focused on the effect of badger vaccination and culling on bTB dynamics with few looking at how roads, human perturbations and habitat change may affect wildlife movement and disease spread. Finally, we observed a lack of studies considering the effect of weather variables on bTB spread, which is particularly relevant when studying zoonoses under climate change scenarios.

Conclusions: Significant technological and methodological advances have been applied to bTB episystems, providing explicit insights into its spread and maintenance across populations. We identified a prominent bias towards certain species and locations. Generating more high-quality

empirical data on wildlife host distribution and abundance, high-resolution individual behaviours and

greater use of mathematical models and simulations are key areas for future research. Integrating

data sources across disciplines, and a "virtuous cycle" of well-designed empirical data collection linked

with mathematical and simulation modelling could provide additional gains for policy-makers and

managers, enabling optimised bTB management with broader insights for other zoonoses.

Key words: Zoonosis, bovine tuberculosis, multi-host episystem, methodologies, management,

mathematical modelling, simulations, disease transmission

Introduction

Emerging infectious diseases represent a significant public health concern as they become more prevalent worldwide [1-3]. It is estimated that about 60% of emerging infectious diseases are zoonotic, 72% of which have been estimated to originate from wildlife [2, 3]. In 2019, thirteen different zoonoses had confirmed cases in humans within the European Union [4]. This has been promoted by human technological advances that have caused an exponential growth in global population size and mobility, leading to increased probability of human-wildlife interactions and, therefore, exposure to zoonotic diseases [5, 6]. Zoonotic disease exposure is particularly worrying when combined with the current human population trend and the expected increase in contacts among humans, livestock and other captive animals, as well as wildlife species [6-8].

A multidisciplinary approach and coordinated collaborations between the public health sector, veterinarians, ecologists and wildlife managers have become key to managing existing and to prevent emerging zoonotic diseases [1]. The importance of interdisciplinary approaches is highlighted by the interlinked nature of human, animal and ecosystem health, which led to the new concept of "One World One Health™" [9, 10]. Due to the complexity of zoonotic disease dynamics, the One Health approach offers the most effective means of studying zoonosis thanks to the collaboration of experts in a variety of research fields. Despite such multidisciplinary efforts, the effect of stressors – i.e., direct and/or indirect disturbances such as hunting, habitat loss, and more broadly habitat and climate change – on animal ecology within human dominated landscapes and the potential emergence of zoonotic disease is still understudied [1]. Overall, there is a lack of empirical evidence on how human perturbations can alter disease transmission and the emergence of zoonotic diseases.

Transmission dynamics of zoonoses involve multiple agents including humans and a diverse range of wild and domestic animals. In order to understand the processes behind their transmission, it is essential to clearly disentangle the role of each agent involved [11]. Due to the complexity of disease transmission and the maintenance of infection within multiple wildlife hosts, the individual

components of the transmission chain are often studied separately, which can limit our understanding

of the subtle underlying effects explaining disease emergence and transmission. Therefore, a holistic

approach is essential to develop a complete picture of the transmission dynamics of zoonotic diseases

[11], with research on rabies being an outstanding example of how empirical data can be used to

elucidate epidemiological dynamics [12].

Evidence from empirical data can be boosted by mathematical simulations, which are

powerful tools for predicting disease transmission trajectories [13]. Simulations of disease

transmission through compartmental models (e.g., the Susceptible, Infectious, and/or Removed (SIR)

model and its variations) have been used in a variety of disease systems, including the recent COVID-

19 pandemic. COVID-19, however, is exceptional in the level of global concern garnered and related

significant investment in funding, resulting in the generation of large and accessible empirical datasets

[14]. Zoonoses are typically more difficult to model due to the lack of empirical data on disease

transmission and associated hosts [15]. Mathematical simulations, therefore, offer opportunities to

model zoonoses where empirical data are limited. In addition, such simulations allow us to undertake

experiments that are currently logistically unfeasible, too costly, too complex or on "unobservable"

phenomena [13, 16].

Studying interactions between and within host species as well as the role played by each host

in the transmission chain is a powerful tool that can be used to better understand zoonotic disease

dynamics. Wild animal interactions are heterogeneous by nature and vary significantly between

different populations as well as individuals. Therefore, it is important to account for this variability to

understand the mechanisms behind transmission and subsequently be able to predict and control

disease spread [7]. This can be achieved by using network modelling, where heterogeneous contacts

between animals can be used to simulate disease transmission [7, 15]. Contact networks can also be

used to understand and describe the contact structure among animals using social network analysis

(SNA) [7]. For example, SNA can be beneficial for disease management since it is able to identify

'super-spreaders' (i.e., highly connected individuals) which can then be targeted for vaccination,

allowing for a dramatic reduction in transmission [1, 7]. In addition, new research is looking at

integrating SNA with molecular epidemiology (phylodynamics) to better estimate transmission

pathways and direction of transmission between individuals [17].

Another important tool in epidemiology is the use of statistical modelling to explain and

predict disease risk and distribution over space and time [18, 19], identify disease clusters [5] and

model host abundance [20]. Spatial analyses make it possible to closely monitor and manage zoonotic

diseases [21]. Additionally, using a diverse range of temporal scales can be useful in examining long-

term disease dynamics, as well as the short-term effects of environmental variables [22]. As ecological

processes occur at different scales (from single study sites to macroecological scales), the spatial scale

used for disease distribution modelling is crucial in understanding how these processes exacerbate

the spread of zoonotic diseases [23, 24]. Large spatial scales (i.e., global, continental) can examine the

broader picture and disentangle how host abundance and abiotic factors influence disease prevalence

[11]. Smaller spatial scales (i.e., country, region) can be used to examine population dynamics and

pathogen genetic diversity at the local level [11]. Therefore, spatially-explicit analyses can be used to

inform One Health approaches, advancing our knowledge when high-frequency data on animal

movement are collected. For zoonotic pathogens, consideration of animal movement in the wild (e.g.,

dispersal and ranging dynamics) or domestic hosts (e.g., intraherd movements across farm habitats or

interherd trade movements) at an individual level can provide additional insights into spatial

processes. Temporal patterns are important to consider as many zoonotic diseases show seasonal

variations (e.g., Zoonotic enteric diseases such as *Salmonella* spp, *Escherichia coli*, *Giardia* spp) as well

as possible daily variations (i.e., due to the circadian rhythm of microbes and pathogens as well as

chronobiology of wildlife hosts) in their infection patterns [25-27].

Human alterations to the environment can also shape and modify zoonotic transmission [11,

21, 28, 29]. However, we currently lack a framework to fully understand their impact on host

transmission dynamics [11]. Current research has shown different ways in which pathogen

transmission can be affected. For example, human-driven changes in the environment can modify

interactions between hosts, change host and vector densities, and alter host longevity and movement

[11, 29]. A study by Castillo-Neyra *et al*. showed that rabies transmission was spatially linked to water

channels, which act as ecological corridors connecting multiple susceptible populations and facilitating

pathogen spread and persistence [29]. With cities expanding and providing urban corridors to wildlife,

pathogen persistence could become even more of an issue [29], confirming the importance of

studying the effect of human perturbations on animal ecology and related implications in disease

ecology.

Understanding the epidemiological dynamics of zoonosis is critical from both a health and

economic perspective. A critical example is Zoonotic Tuberculosis (zoonotic TB), which was estimated

in 2016 to be linked to 147,000 human cases and 12,500 deaths worldwide [30]. Zoonotic TB is driven

mainly by *Mycobacterium bovis* (i.e., the causative agent of Bovine Tuberculosis – known also as

bovine TB or bTB), which is transmitted by several wildlife hosts and livestock. Britain and Ireland, as

well as many other countries worldwide [30], have been increasingly impacted by bTB, resulting in

significant economic loss. In Ireland, for instance, 4.2% of cattle herds tested positive for bTB in 2020,

leading to the humane killing of 21,467 cattle [31]. This is in addition to the economic costs associated

with the national bTB eradication program with €92 million spent in 2018 alone [32]. Similar trends

can be observed in the UK, with £70 million spent annually for bTB prevention and control [33]. This

disease also raises welfare concerns for wildlife hosts, especially considering its high prevalence in the

wild. Badgers *Meles meles*, for example, have been shown to have a bTB prevalence exceeding 40% in

hotspot areas in Ireland [34], and red deer in Spain have been estimated to have a prevalence of up

to 50% [35].

Bovine TB eradication is prioritised by governments and researchers due to the significant

health concerns and economic (trade) impacts. Despite decades of control efforts in several countries,

the pathogen has successfully avoided eradication. There are complex reasons as to why this is the

case [36], but a primary factor relates to its complex dynamics of transmission and maintenance across

differing hosts and the environment. Therefore, new thinking may be required to further investigate

if disease control can be driven toward eradication. Detecting gaps in the current bTB literature is an

essential step required to identify target areas for future research and to further hone government

eradication strategies. Here, we aimed to uncover empirical and methodological gaps in the peer-

reviewed literature on bTB. Our intention is to use bTB as an example of a complex multihost zoonotic

disease for which recent developments with sampling design, animal monitoring tools and technology,

and mathematical modelling has helped to fill the gaps in knowledge and improve our understanding

and ability to combat zoonotic diseases more generally.

To achieve our goal, we developed a scoping review of bTB multihost epidemiology focusing

on 18 research questions (reported in **Table 1** and conceptually summarised in **Fig. 1**) regarding the

type of study, whether, which and how wildlife species have been monitored, what kind of sampling

designs and methodological approaches have been used, and whether epidemiological empirical data

have been collected. We then gathered data from the peer-reviewed literature on the mechanisms

driving inter- and intraspecies bTB transmission, looking in particular at novel and multi-disciplinary

approaches, therefore focusing on the scientific literature published in the last six years (January 2017

- July 2022). Our goal is that our work will spark renewed discussion on how to monitor and deal with

zoonotic diseases, direct future research, and stimulate focused funding efforts.

Review

1. Methods

We conducted a scoping review (as per PRISMA guidelines) [37] by sourcing peer-reviewed papers using Web of Science (Clarivate, 2021 Online Version) focusing on bovine tuberculosis, and more specifically its most common cause, *Mycobacterium bovis*, in cattle and several key wildlife hosts. The search terms and list of articles have been summarised in **Additional file 1**. We identified 3,531 potentially relevant papers (i.e., the search included all years of publication) which were uploaded and screened for duplicates using EndNote (Clarivate, Version 20.1.0.15341). Relevant articles were then selected using a PEO (Population, Exposure, Outcomes) eligibility criterium structure [38]. The aim of the PEO is to identify articles of interest by selecting the "Population" (i.e., the subject being affected by the disease/health condition) for a particular "Exposure" (i.e., a disease/health condition) and either a particular "Outcome" or "Themes'' to examine [38, 39]. The PEO eligibility criterium was chosen since it was in line with the recommendations given for scoping reviews that target literature on etiology and risk factors, such as a particular disease. We decided to use a modified version of the PEO framework structure which also includes themes of interest as potential "Outcomes" [39], as summarised in **Table 2** to guarantee reproducibility. All papers that did not meet the eligibility criteria listed in **Table 2** were removed. bTB has received significant attention in the last few years due to its financial impact on rural economies and the risk it poses to human health. We selected the most recent papers published in the period January 2017 - July 2022 (see **Fig.2** for a summary of the step-by-step process), making up 30% of all relevant papers identified, to capture the most recent methodological approaches used to monitor and combat bTB. The papers were screened to answer the questions of interest summarised in **Table 1,** stored in an excel spreadsheet. The results were then imported and plotted using *ggplot2* in R version 4.1.1 [40].

2. Results

Our results are based on 167 peer-reviewed papers published between January 2017 and July 2022. The study location of the papers was representative of 6 continents and 37 different countries (**Fig. 3**). The continent with the highest number of studies on bTB is Europe (n = 103, 52 of which were from the UK), significantly higher than those carried out in much larger continents such as Africa, Asia, and both Americas (**Fig. 3**). We screened all papers for 18 different variables (addressing our 18 questions, see **Table 1**) which we summarised in the following section under the heading: 2.1 **general characteristics** (Sub-headings: "Study species and wildlife species"; "Management and data type"), 2.2 **data analysis** (Sub-headings: "Spatial analysis, spatial scale and temporal scale"; "Farm environment and human perturbations"), and 2.3 **epidemiological analysis** (Sub-headings: "Intra- and interspecies interactions", "Direction of transmission and compartmental models"). Note that most plots presented below have a sample size of n = 167, corresponding to the number of papers screened, with a few exceptions where this sample size is higher (for example, in relation to species included in the study, because a paper reported data about both wildlife and cattle, therefore contributing to multiple levels of a category) or lower (for example, in relation to epidemiology, where variables of interest were analysed only in the subset of papers describing studies that included epidemiological interactions).

2.1. General Characteristics

2.1.1. Study species and wildlife species

We found that 55% of bTB papers focused on cattle only, whereas 26% of them included both cattle and wildlife species and 19% targeted only wildlife species (**Fig. 4a**). Among those papers reporting wildlife data, we found that the European badger attracted most research effort (60% of wildlife studies), followed by wild boar (29%) and cervid species (24%: 17% red deer, 7% fallow deer, 5% white-tailed deer, 1% roe deer 1%, and 1% wapiti elk, from hereinafter referred to as simply elk) (**Fig. 4b**). Other species were included in 12% of wildlife papers (brush-tailed possum: 8%, wild buffalo: 4%).

 2.1.2. Management and data type

 Our results highlighted that only 13% of the studies dealt with management solutions (**Fig. 4c**).

 Management strategies mainly included vaccination (4%) or culling (2%), with 1% looking at both

 vaccination and culling. We also found that most papers gathered original empirical data (85%), and

 papers using simulations were very limited (4%), with a remaining 11% of papers combining empirical

 data and simulations (**Fig. 4d**).

 ## 2.2. Data Analysis

 2.2.1. Spatial analysis, spatial scale and temporal scale

 We found that the majority of papers did not include any spatial analysis. Those that did largely

 focused on spatial patterns in cattle (31%, **Fig. 5a**) slightly more than wildlife (25%, **Fig. 5b**). Among the

 51 papers that investigated spatial analysis in cattle, 47% looked at bTB risk and probability of infection;

 20% investigated cattle movement outside the farm, 12% looked at cattle interactions with wildlife, 10%

 analysed the distribution of bTB positive biological samples, and 8% focused on detection rate by spatial

 unit (e.g., 1 km^2 grid). Movement inside the farm and interactions of farm animals with both wildlife and

 cattle were included in 2% of papers. Among the 41 studies which investigated spatial behaviour in

 wildlife (Fig 5b), analysis was undertaken using a variety of methodologies; direct observations (34%),

 satellite GPS telemetry (32%), camera traps (7%), genetic samples (12%) and proximity loggers (5%), or

 spatial patterns predicted by future scenarios modelling or mathematical simulations (12%).

 We also found that most papers included spatial scales at the regional level or smaller (74%),

 with less than 4% papers looking at national and/or international spatial scales (**Fig. 5c**). In regard to

 temporal scales, 35% of the studies considered interannual variability, whereas 20% tackled intra-

 annual variability, with a small fraction (9%) of studies including both intra- and interannual variability.

 Thirty percent of the studies did not analyse any intra- or interannual temporal variability (**Fig. 5d**). Only

 5% of the studies looked at fine-scale variability (e.g., days), whereas in a few instances the year of study

270 was not reported at all (5%). Finally, only 6% of papers included predictions for temporal patterns into

271 future scenarios.

2.2.2. Farm environment and human perturbations

When looking at farm characteristics, 44% of the studies included some type of herd data (e.g.,

herd size, bTB history), with 40% not including any type of in-farm environmental variables (**Fig. 6a**) and

15% of papers incorporating other types of farm characteristics. These included environmental

conditions on the farm (e.g., natural habitats, land-fragmentation; 7%), farm location in respect to other

farms (6%) and farm location in respect to wildlife (5%).

Environmental conditions outside the farm were included in 24% of the papers' data analysis

(**Fig. 6b**). These studies mainly looked at habitat characteristics around the farm (e.g., wildlife presence,

natural habitats), with one paper also including variables focusing on habitat variation (e.g., forest

clearfell, new artificial plantations). We also looked at weather variables (e.g., temperature, rainfall) and

observed that 5% of papers included these as part of their analysis (**Fig. 6d**). Finally, 10% of the papers

screened included human perturbation variables with the vast majority looking at the effect of

vaccination and culling on transmission dynamics (**Fig. 6c**).

2.3. Epidemiological analysis

2.3.1. Intra- and interspecies interactions

We found that most papers (74%) did not include an analysis on interactions, with 17% of papers

looking at intraspecies transmission and 9% at interspecies transmission (**Fig. 7a**). Among the interaction

studies, 75% included direct interactions and 25% included indirect interactions (**Fig. 7b**). In addition,

interaction data were mostly collected using technological tools (43%; e.g., GPS, proximity loggers,

camera traps), followed by direct observations (39%) and a minority using simulations (11%) (**Fig. 7c**).

The methodology used to analyse interaction data also varied between papers with most papers using

social network analysis (57%), and the others (48%) using a variety of statistical techniques (e.g., linear

models, logistic models, differential equations) (**Fig. 7d**).

2.3.2. Direction of transmission and compartmental models

We found that a limited proportion of the papers (12%) included direction of transmission in

their analysis (**Fig. 8a**). We also found that epidemiological modelling techniques (e.g., compartmental

models and transmission rates) were adopted in 11% of the studies (**Fig. 8b**).

3. Discussion

In this review we examined bTB papers published between January 2017 to July 2022

corresponding to a third of all bTB papers ever published meeting our selection criteria, highlighting the

high level of interest on this zoonosis by researchers aimed at curbing its spread and economic impacts.

We found that there has been a great deal of research effort focusing on the badger-cattle bTB

episystem; we acknowledge, however, that we found a very limited number of studies on other

episystems [41-43]. Our spatially-explicit overview of bTB research efforts (**Fig. 3**) highlights how the

badger-cattle episystem has been the focus of research across Europe and particularly in Britain and

Ireland. Our review also shows that there has been a different level of attention across episystems - e.g.,

there has been huge investment of effort and money, resulting in huge output on the badger-cattle-TB

episystem, but far less on the multi-host episystems of southern Africa, and we believe we have more

to learn from these other systems which are chronically understudied.

Our scoping review found a limited number of studies focusing on management solutions and

their efficacy, with very few studies looking at modelling exit strategies [44, 45]. This is not surprising

considering the scarcity of studies using mathematical simulations, not only to better understand and

predict possible outputs of management solutions, but also to explore long-term bTB dynamics under

different scenarios, e.g. [46, 47]. Surprisingly, only a small number of studies have looked at the effect

of human disturbances on the spread of bTB involving wildlife host species, and this knowledge gap

indeed needs to be tackled in light of research showing the significant role of human perturbations on zoonotic outbreaks and spread [48-50]. Most of the studies we reviewed have focused on the effect of badger vaccination and culling on bTB dynamics with only three studies looking at how roads or human perturbations may affect these dynamics [51-53], and one focusing on the effect of habitat change (clearfell forest operations) on bTB breakdowns [54]. The latter, nevertheless, lacked empirical data on wildlife, which has been indicated as one of the drivers of bTB outbreaks in cattle farms [54]. Finally, we observed that there is a lack of studies looking at the effect of weather variables (i.e., rainfall, soil humidity, temperature etc.) on bTB spread [55]. This is especially important when thinking about wildlife-cattle transmission since it is now thought to occur also through environmental sources [56].

We have carefully evaluated the outcome of our scoping review, and in the following sections we have summarised data types and methodological approaches which, we believe, could contribute to gaining further insights into bTB epidemiology. Based on our review, we have identified a significant gap when it comes to prediction and simulation models, which would be a useful tool for managers to assess disease risk under different land use and climate change scenarios. One major gap that we have to stress is the lack of integration between empirically informed tactical (short-term decision support) and strategic (larger spatial scales and longer term) models blending multiple approaches (though note exceptions before the screening period of our scoping review, see for instance Brooks-Pollock *et al.* [57]). Future research should include compartmental models fitted across space, linked via meta-populations and/or real-landscape multi-host episystems; or agent-based models (ABMs) with empirical data feedback loops. We describe such modelling approaches and their prerequisites in the following sections, beginning with data and monitoring programs, and we continue with recent advances in technology, mathematical tools and analytical solutions.

3.1. Empirical data and long-term monitoring programs: involving stakeholders and setting up fixed long-term monitoring stations across large spatial scales

In order to better understand zoonotic disease dynamics in wild animals we need reliable data sources to model spatial distribution and abundance of the host species involved in transmission, as good quality data is the cornerstone to building informed strategies that involve wildlife interventions. In reference to the badger-cattle bTB episystem in western Europe, both badgers (with several examples among the literature: [58, 59]), and cattle [60] have been extensively monitored. However, in some populations it is possible that deer and wild boar may also play a role in the local spread and maintenance of infection [61]. In Britain and Ireland, in the past decades, the significance of deer as a wildlife host impacting bTB epidemiology has been uncertain [62]. However, recent research is starting to uncover the role deer may play at local scales where conditions favour the transmission between badgers-deer-cattle [63]. There could be opportunities to gather data in collaboration with hunters (as has been used in France [64] and Spain [65], for example) to have access to an incredibly high number of deer samples, within and across countries (large spatial scales) and across years (long-term temporal scales). Involving stakeholders like hunters may provide the unique opportunity to use field tests for bTB [66, 67] to be submitted via smartphone applications, see [68], and to monitor the dynamic of the disease across multiple spatio-temporal scales and in relation to bTB occurrence in the two other hosts in the system (badgers and cattle). The ability to intercept the collaboration of stakeholders widely spread across large spatial scales (e.g., hunters, farmers, foresters) may help to establish systematic, relatively inexpensive, and long-term monitoring programmes able to provide data for Bayesian species and disease distribution models (described in section 3.3), allowing managers to access up to date risk scenarios. This approach can also highlight hotspots of disease outbreak that would drive very focused longitudinal studies involving satellite telemetry on multiple species simultaneously to better disentangle species overlaps and contact rates [69, 70]. The role of stakeholders/citizen scientists in this bTB example would be confirming infection, which is almost never economical, although there is the

hope that cheaper field tests will be released in the next decade – or at least, we hope there will be

investment in that direction. For the time being, a well distributed number of samples could be collected

from hunters to cover large areas systematically and limit the costs required for testing.

When it comes to establishing long-term monitoring programs, fixed long-term sampling

stations across large-spatial scales can capture wildlife population spatio-temporal dynamics. This can,

on one hand, provide data on occurrence, relative density, and spatio-temporal overlaps of the host

species and, on the other hand, gather key empirical data required to parameterise mathematical

simulations. Camera traps are a popular and effective tool for estimating state variables of wildlife

populations [71]. For ungulates, they have successfully been used to understand temporal behaviour

(e.g., diel activity patterns, [72]), spatial behaviour (e.g., occupancy, [73]), and abundance (e.g., density,

[74]). Camera traps have been used for quantifying temporal and spatial overlap of wild ungulates with

domestic animals in open systems [75, 76] with varying results [77]. Kukielka *et al*. demonstrated their

use in identifying hotspots of indirect wildlife–livestock overlap for the prevention of TB crossover [75].

For wildlife, especially ungulates, camera traps offer powerful monitoring solutions not only to measure

abundance and spatial overlap, but also to understand behavioural dynamics that may align closely with

disease risk. An example is the use of camera traps to individually recognise animals, which has been

shown to be possible in a recent study by Hinojo *et al*. [78]. They demonstrated how roe deer (*Capreolus*

capreolus) antler shapes could be used to identify distinct individuals. This data could be used to obtain

better estimates of abundance as well as to build wildlife social networks (which will be discussed in

more detail in section 3.3) to collect data on contact rates between and within species. The parameters

from these analyses would be useful as an input for mathematical simulations to help better understand

disease transmission dynamics in wildlife populations.

3.2. Recent advances with technology can help to gather data for mathematical simulations: interindividual variability within animal populations matters and should be taken into account

Detailed questions concerning the animal behaviour and movement patterns promoting disease transmission can be posed and answered using animal-attached sensors, i.e., biologging [79, 80]. GPS units are the most widely used of these sensors, providing data on animal locations and space use. Proximity sensors can detect when two or more sensor-equipped animals interact and can be used to detect direct encounters which may result in disease transmission. Collars with both GPS units and proximity sensors have been used concurrently on badgers and cattle, finding that while badgers show a habitat preference for cattle pastures, there were no direct contacts between the two species, suggesting that environmental transmission may play an important role in the case of bTB [81]. As such, proximity sensors allow insights which are not obtainable through investigating shared space use alone. Proximity sensors can also provide information on whether and how the duration of exposure to an infected individual affects transmission rate, where disease state of individuals is known [82]. Other biologging sensors, including accelerometers, magnetometers and gyroscopes, are used to classify distinct behaviours from logged datasets [80]. Behaviour classification allows activity budgets to be built, so that behaviours which increase the likelihood of acquiring or transmitting pathogens can be detected and mapped in the landscape. Accelerometers have also been used to compare micro-movements in diseased and healthy animals, with diseased animals exhibiting differences in posture, gait dynamism (e.g., the "bounce" in subsequent walking steps) and energy levels [83]. Monitoring such micro-movements in cattle could act as a warning sign to test herds for bTB when signs of illness are detected, by adapting existing systems in place to monitor lameness through accelerometry [84]. Further, these effects of disease on the internal state of animals yield important insights into how disease status impacts animal movement patterns and therefore disease spread.

Biologging and satellite telemetry monitoring can, on one hand, provide answers aimed at

disentangling the transmission dynamics within multi-host disease systems [81, 85, 86] and, on the other

hand, provide highly valuable empirical data that are strongly needed by parameter hungry

mathematical simulations [82]. When tracking animals, however, special care should be spent to

understand which animals we are monitoring, and whether we are following a special subset of

individuals that are easier to trap. This applies also to where we study animals which will provide

empirical data for mathematical simulations, because the behaviour and movement ecology may vary

significantly depending on the level of human disturbance.

The concept of One Health has indeed highlighted the role of human activities in the spread of

zoonotic diseases [87]. For example, urbanisation, improper waste disposal, and the intentional feeding

of wildlife have been shown to result in wildlife movement into human-dominated areas [88], which

may facilitate disease transfer to humans and other animal communities [89]. However, evidence has

shown that only a select proportion of individuals within wildlife populations will engage in interactions

with humans [90] or utilise these human-dominated areas [88, 91]. Individual variation in movement

patterns [92], sociability [93] and immunological defence [94] among others, impacts disease

transmission and spread [95]. There is supportive evidence that certain behavioural types have higher

infection rates than others (e.g. [96, 97]), although the causal direction may be difficult to determine

since infections also alter host behaviour [96, 98]. Regardless, to gain a more complete understanding

of disease spread, future studies should incorporate this individual variation. These studies often utilise

direct behavioural observations, since these are an invaluable data source that can be used to determine

which individuals in a known population are more likely to engage in close-contact interactions with

humans [90] or access human areas (e.g., farmland) [99]. This can provide us with information on which

individuals in a population may be at "higher risk" of transferring disease to human or other animal

populations.

3.3. Modelling and mathematical simulations: social network analysis, Bayesian species distribution models, and Agent Based Models

Social network analysis is a powerful tool for understanding the causes and consequences of disease transmission within the animal community [100, 101]. In the past decade social network analysis has mainly focused on understanding contact and transport networks of cattle and livestock movements, as well as wildlife movements [102-104]. Nonetheless, the area of network analysis can be expanded to better understand the dynamics of disease transmission between wildlife populations and livestock [101]. Unlike cattle monitoring the movements and interactions of wildlife can be challenging, where typically, a small proportion of individuals are monitored using recent advances with biologging and satellite telemetry, as discussed earlier. Recent advances in statistical analysis of social networks have paved the way to obtain better inferences from limited data [105-107]. Once the network metrics affecting disease transmission dynamics are identified (e.g., transitivity, betweenness centrality) [105, 106], those can be tested via pre-network permutations of available observations to ensure that the available data sufficiently captures non-random interactions among the animals. This can help estimate the scale of disease transmission in wild habitats. Global metrics of a social network help estimate the changes in the overall structure. For example, the network metric transitivity represents the tendency of a population to cluster together and is considered to be negatively correlated with disease transmission rates [104]. However, recent research [106] shows that it is noisily estimated at low sample sizes. Therefore, stable metrics with respect to low sample sizes should be identified before making inferences. Local network metrics such as betweenness centrality represent the tendency of an individual to serve as a bridge between one part of the community and another, helping the selection of individuals to be vaccinated/removed from the population. Research on data collected for wild ungulates [106] shows that the betweenness centrality values of smaller samples remain well correlated with those in larger samples. Similar correlation analysis can be done for other network metrics, mainly in cases of limited data availability for disease transmission. Whenever limited animals from a population

are monitored, confidence intervals around the network metrics should also be obtained to make

informed decisions using statistical evidence. Using the methodologies discussed by Silk *et al*. and Kaur

et al. [104, 106] we now have the possibility of analysing all telemetry data collected thus far on species

involved in bTB transmission (e.g., badgers, wild boar; but also applicable to species from other disease

systems) to test hypotheses on disease transmission dynamics. In addition, it will help in collecting future

data aimed at answering very precise questions such as understanding the role of deer species in bTB

transmission by collecting simultaneous telemetry data of badgers and deer species in Ireland.

In reducing zoonotic risk, knowing the distribution and abundance of wildlife vectors is also

essential [28, 108]. To that aim, Species Distribution Models (SDMs) can be used to produce models of

the distribution and abundance of species based on occurrence data [109]. In recent years spatial

modelling has undergone a conceptual and technical revolution. New modelling techniques within

Bayesian [110] and Machine Learning frameworks [111] allow us to develop spatially explicit models of

animal abundance and distributions with unprecedented accuracy, and the improvement of

computational power allows computers to rise to the challenge and cope with the high computational

demands of these models. The flexibility of the new techniques allows us to use different types of data

(e.g., individual tracking data, survey data, and even citizen science data) and combine them in what is

called Integrated Species Distribution Models (ISDMs), while still taking into account the different

observational processes of each type of data, to produce accurate models even in data scarce systems

[112]. In addition, these new techniques also allow for the calculation of uncertainty in a spatially explicit

manner, which will help us evaluate the quality of the models and better interpret the results. Bayesian

ISDMs using INLA (i.e., Integrated Nested Laplace Approximation) [113] were used to model the

distribution of red, sika and fallow deer in Ireland, which are vectors of bTB [68]. The models produced,

for the first time, relative abundance and distribution maps for each species, which will be an essential

tool for deer population management and thus towards bTB eradication. They are already being used

to determine high sika-density areas for a pilot study on the effect of deer on biodiversity, which will

provide further management tools for the overabundant deer populations in Ireland. In addition,

hierarchical Bayesian models are also the basis of a new project aimed at modelling European badger

sett distribution, badger density, and their body condition. These three models will be linked to bTB

infection in badgers and outbreaks in cattle, in an attempt for the first time to link badger spatial ecology

to bTB management and eradication in Ireland (V. Morera-Pujol 2023, personal communication).

Agent-based simulations are a powerful tool for understanding the transient effects of wildlife

disease in human-dominated landscapes, both as a complementary tool to traditional methods and as

a surrogate for scenarios where data is limited or not available [16]. These models serve as a

computational laboratory that allow researchers to plug-in available real-world data and parameterise

both agents (for instance, a badger) and the environment (for instance, a mosaic of natural habitats and

farms) to empirically test if animal behaviour in response to landscape change or management

interventions modulates disease risk dynamics over time and space [114]. Recent technological

advancements have bolstered agent-based simulations allowing for high-resolution spatio-temporal

models that incorporate geographic information systems (GIS) data to create hyper realistic

environments, and machine learning algorithms to introduce cognition and applied decision making for

agents. Furthermore, process-driven agent-based models (e.g., disease transmission) can be integrated

into larger mechanistic agent-based models (e.g., ecosystem scale epi-dynamics) for increasingly higher-

resolution models that reduce uncertainty and overly-theoretical parameterisation of model entities

[16]. The development of hyper-realistic agent-based simulations, parameterised with high-resolution

data, for the management of bovine tuberculosis in multi-host systems can contribute to answering

important policy questions and how best to select management directions. In practice, this allows for

the totality of data collected in complex multi-host systems to be incorporated into a single environment

where they may be measured against one another in the simulation to deduce the possible effects of

each predictor. Take for example the European badger as the primary wildlife host in Ireland as a case

study. Badgers are prevalent in the agroecological mosaic of natural habitats and farms in Ireland. Agent-

based simulations can utilise data from badger tracking studies [53, 115, 116], habitat suitability [117],

culling and vaccination programmes [118] and disturbance regimes [54, 119] to simulate badger

movement and behaviour realistically. GIS data for farm location and characteristics [120], as well as

ecological and environmental data streams, can then place the badger agent into a highly realistic

environment to examine how these factors affect badger movement, behaviour, and other parameters,

for instance, contact rates with domestic animals. Interactions between agents and the environment

can be modulated by sub-models to further increase the strength of the model, for instance, weather

sub-models (e.g., rainfall) may influence agricultural practice and thus contact rates, alternatively,

disease transmission could also be sub-modelled so that contact rates may/may not result in infection

[16]. Finally, management decisions can be trialled within the simulation to see how likely decisions

change the status of disease within the study system, allowing for "What if?" scenarios to play out

without risk to animal or human welfare or livelihoods.

Our review is a starting point to better understand the current gaps in the zoonotic disease

literature. We are aware that this review has some limitations such as the use of only one search engine,

as well as focusing on one zoonotic disease system and date range. However, we hope that it will incite

new and exciting possibilities for the future which could expand on this current work, by for example

looking at multiple disease systems as well as other date ranges to try and uncover more information

on possible way forwards to combat zoonotic disease transmission.

Conclusion

Our exploration of the recent literature on multi-host bTB episystems, as an example of zoonotic One

Health challenges, has revealed a significant body of work utilising a diversity of methodologies at

different spatio-temporal scales and subjects (individual vs. group) levels. There was a significant bias in

the literature towards one particular episystem, the badger-cattle system that predominates in north-

western Europe, reflecting large financial burdens (for both governments as well as the agricultural

sector) and research funding investments. Alternatively, there were comparatively less publications

from the global south, especially in complex muti-host episystems in southern Africa. In such episystems,

the cost-effective and efficient collection, collation, and use of data are essential to drive greatest added

value to inform on policy options. A theme from the present work is the essential need to use new and

emerging technologies to derive new insights from data that may have been collected for other

purposes.

Given the results from our scoping review, we reflect on several areas where progress could be

made, including the need for high-quality data on wildlife hosts, even in episystems where significant

research investments have already been made. Such careful collection and utilisation of empirical data

could then feed the development of social network analyses, Bayesian distribution models and

eventually Agent Based Models. Mathematical simulations such as ABMs trained on synthetic data and

parameterised by real empirical data can answer questions that would otherwise be too costly,

unethical, or both. Such models can also be used to explore different scenarios in an increasingly human-

dominated world, under different levels of land-use and climate change, or with the appearance of

invasive species in already complex multi-host epidemiological systems. In addition, it can help build

cross-disciplinary bridges with other areas, deriving significant insights into interspecific transmission

like phylodynamic modelling.

Taking Ireland as an example, with the intention not to stay local but rather inspire researchers

from across the globe; Ireland invests considerably in surveying, culling, and vaccinating badgers [121].

However, the question remains - which applies to other countries and zoonotic episystems - should we

be doing more, or being smarter with the data we already have? We suggest the latter. Yes, we need to

be smarter, arranging *ad hoc* data collections using the latest technological tools to estimate unknown

parameters. We need to focus our efforts on mathematical modelling (ABMs, INLA-Bayesian) to optimise

our information gain from the large, high-quality datasets collected over the last few decades (and for

sparser datasets, taking advantage of recently developed statistical tools for enhanced inferences, see

[54, 68]). We have (almost) all the data required to parameterise simulations with significant utility: this

should be one main focus in future research. We believe that, ideally, the feedback of simulation and

mathematical tools to inform data collection, and the "virtuous cycle" of feeding this new data to

improve the next generation of model is a priority for decision making tools for policy makers and

programme managers.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are available in the '*Curbing zoonotic*

disease spread in multi-host-species systems will require integrating novel data streams and analytical

approaches: evidence from a scoping review of bovine tuberculosis' repository,

https://doi.org/10.5281/zenodo.7886368

Competing interests

The authors declare that they have no competing interests.

Funding

This publication has emanated from research conducted with the financial support of Science

Foundation Ireland under Grant number 18/CRT/6049. For the purpose of Open Access, the author has

applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this

submission.

In addition, HME is funded by an Irish Research Council Government of Ireland postgraduate

scholarship.

Authors' contributions

KC - conceived and designed the review, acquired, analysed and interpreted the data, wrote the first

draft of the manuscript.

HME - edited and revised the ms and contributed to the interpretation of the data.

AWB - edited and revised the ms and contributed to the interpretation of the data, drafted the policy

and research implications.

BA, LLG, PK, VMP, KJM, AFS, MST - edited specific sections of the discussion and revised the whole ms.

SC - conceived and designed the review, edited the ms and supervised the process.

All Authors approved the final version of this review.

Acknowledgements

We would like to acknowledge Sylvia Power for helping in reviewing the final version of the

manuscript.

References

1. Destoumieux-Garzón D, Mavingui P, Boetsch G, Boissier J, Darriet F, Duboz P, Fritsch C, Giraudoux P, Le Roux F, Morand S et al (2018) The One Health concept: 10 years old and a long road ahead. Frontiers in Veterinary Science 5: 14.

2. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, Daszak P (2008) Global trends in emerging infectious diseases. Nature 451(7181): 990-993.

3. Karesh WB, Dobson A, Lloyd-Smith JO, Lubroth J, Dixon MA, Bennett M, Aldrich S, Harrington T, Formenty P, Loh EH et al (2012) Ecology of zoonoses: natural and unnatural histories. The Lancet 380(9857): 1936-1945.

4. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control) (2021) The European Union One Health 2019 Zoonoses Report. EFSA Journal 19(2).

5. Beard R, Wentz E, Scotch M (2018) A systematic review of spatial decision support systems in public health informatics supporting the identification of high risk areas for zoonotic disease outbreaks. International Journal of Health Geographics 17(1).

6. Jones BA, Grace D, Kock R, Alonso S, Rushton J, Said MY, Mckeever D, Mutua F, Young J, Mcdermott J et al (2013) Zoonosis emergence linked to agricultural intensification and environmental change. Proceedings of the National Academy of Sciences 110(21): 8399-8404.

7. Craft ME (2015) Infectious disease transmission and contact networks in wildlife and livestock. Philosophical Transactions of the Royal Society B: Biological Sciences 370(1669): 20140107.

8. Johnson CK, Hitchens PL, Pandit PS, Rushmore J, Evans TS, Young CCW, Doyle MM (2020) Global shifts in mammalian population trends reveal key predictors of virus spillover risk. Proceedings of the Royal Society B: Biological Sciences 287(1924): 20192736.

9. One Health Commission (2022) Twelve Manhattan principles formulated. Available at: https://www.onehealthcommission.org/index.cfm/37526/75802/twelve_manhattan_principles_formulated (Accessed: 25/02/2022)

10. Zinsstag J, Schelling E, Waltner-Toews D, Tanner M (2011) From "one medicine" to "one health" and systemic approaches to health and well-being. Preventive Veterinary Medicine 101(3): 148-156.

11. Estrada-Peña A, Ostfeld RS, Peterson AT, Poulin R, De La Fuente J (2014) Effects of environmental change on zoonotic disease risk: an ecological primer. Trends in Parasitology 30(4): 205-214.

12. Fisher CR, Streicker DG, Schnell MJ (2018) The spread and evolution of rabies virus: conquering new frontiers. Nature Reviews Microbiology 16(4): 241-255.

13. Siettos CI, Russo L (2013) Mathematical modeling of infectious disease dynamics. Virulence 4(4): 295-306.

14. Adam D (2020) Special report: The simulations driving the world's response to COVID-19. Nature 580(7803): 316-318.

15. Craft ME, Caillaud D (2011) Network Models: An underutilized tool in wildlife epidemiology? Interdisciplinary Perspectives on Infectious Diseases 2011: 1-12.

16. Murphy KJ, Ciuti S, Kane A (2020) An introduction to agent-based models as an accessible surrogate to field-based research and teaching. Ecology and Evolution 10(22): 12482-12498.

17. Vasylyeva TI, Friedman SR, Paraskevis D, Magiorkinis G (2016) Integrating molecular epidemiology and social network analysis to study infectious diseases: Towards a socio-molecular era for public health. Infection, Genetics and Evolution 46: 248-255.

18. Schrödle B, Held L (2011) Spatio-temporal disease mapping using INLA. Environmetrics 22(6): 725-734.

19. Talisuna AO, Okiro EA, Yahaya AA, Stephen M, Bonkoungou B, Musa EO, Minkoulou EM, Okeibunor J, Impouma B, Djingarey HM et al (2020) Spatial and temporal distribution of infectious disease epidemics, disasters and other potential public health emergencies in the World Health Organisation Africa region, 2016–2018. Globalization and Health 16(1): 9.

643   20.   Charron MV, Kluiters G, Langlais M, Seegers H, Baylis M, Ezanno P (2013) Seasonal and spatial
644   heterogeneities in host and vector abundances impact the spatiotemporal spread of
645   bluetongue. Veterinary Research 44(1): 44.
646   21.   Brownstein JS, Rosen H, Purdy D, Miller JR, Merlino M, Mostashari F, Fish D (2002) Spatial
647   analysis of West Nile Virus: Rapid risk assessment of an introduced vector-borne zoonosis.
648   Vector-Borne and Zoonotic Diseases 2(3): 157-164.
649   22.   Tao Y, Hite JL, Lafferty KD, Earn DJD, Bharti N (2021) Transient disease dynamics across ecological
650   scales. Theoretical Ecology 14(4): 625-640.
651   23.   Brock PM, Fornace KM, Grigg MJ, Anstey NM, William T, Cox J, Drakeley CJ, Ferguson HM, Kao
652   RR (2019) Predictive analysis across spatial scales links zoonotic malaria to deforestation.
653   Proceedings of the Royal Society B: Biological Sciences 286(1894): 20182351.
654   24.   Cohen JM, Civitello DJ, Brace AJ, Feichtinger EM, Ortega CN, Richardson JC, Sauer EL, Liu X, Rohr
655   JR (2016) Spatial scale modulates the strength of ecological processes driving disease
656   distributions. Proceedings of the National Academy of Sciences 113(24): E3359-E3364.
657   25.   Diallo AB, Coiffard B, Leone M, Mezouar S, Mege J-L (2020) For whom the clock ticks: Clinical
658   chronobiology for infectious diseases. Frontiers in Immunology 11: 1457.
659   26.   Kronfeld-Schor N, Bloch G, Schwartz WJ (2013) Animal clocks: when science meets nature.
660   Proceedings of the Royal Society B: Biological Sciences 280(1765): 20131354.
661   27.   Lal A, Hales S, French N, Baker MG (2012) Seasonality in human zoonotic enteric diseases: A
662   systematic review. PLoS ONE 7(4): e31883.
663   28.   Allen T, Murray KA, Zambrana-Torrelio C, Morse SS, Rondinini C, Di Marco M, Breit N, Olival KJ,
664   Daszak P (2017) Global hotspots and correlates of emerging zoonotic diseases. Nature
665   Communications 8(1): 1124.
666   29.   Castillo-Neyra R, Zegarra E, Monroy Y, Bernedo R, Cornejo-Rosello I, Paz-Soldan V, Levy M (2017)
667   Spatial association of canine rabies outbreak and ecological urban corridors, Arequipa, Peru.
668   Tropical Medicine and Infectious Disease 2(3): 38.
669   30.   World Health Organization (WHO), Food and Agriculture Organization of the United Nations
670   (FAO), World Organisation for Animal Health (OIE) (2017) Roadmap for zoonotic tuberculosis.
671   World Health Organization: Switzerland
672   31.   Department of Agriculture, Food and the Marine (2021) National bovine TB statistics. Available
673   at: https://www.gov.ie/en/publication/5986c-national-bovine-tb-statistics-2020/ Accessed 8
674   Nov 21
675   32.   Department of Agriculture, Food and the Marine (DAFM) (2019) Spending review 2019. Animal
676   health: TB Eradication. Department of Agriculture, Food and the Marine: Dublin.
677   33.   Department of Environment, Food and Rural Affairs (DEFRA) (2018) Bovine TB strategy review.
678   Department of Environment, Food and Rural Affairs: England.
679   34.   Corner LAL, O'Meara D, Costello E, Lesellier S, Gormley E (2012) The distribution of
680   *Mycobacterium bovis* infection in naturally infected badgers. The Veterinary Journal 194(2): 166-
681   172.
682   35.   Vicente J, Höfle U, Garrido JM, Fernández-De-Mera IG, Juste R, Barral M, Gortazar C (2006) Wild
683   boar and red deer display high prevalences of tuberculosis-like lesions in Spain. Veterinary
684   Research 37(1): 107-119.
685   36.   Allen AR, Skuce RA, Byrne AW (2018) Bovine Tuberculosis in Britain and Ireland – A perfect
686   storm? The confluence of potential ecological and epidemiological impediments to controlling
687   a chronic infectious disease. Frontiers in Veterinary Science 5: 109.
688   37.   Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MDJ, Horsley T,
689   Weeks L et al (2018) PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and
690   explanation. Annals of Internal Medicine 169(7): 467-473.
691   38.   Munn Z, Stern C, Aromataris E, Lockwood C, Jordan Z (2018) What kind of systematic review
692   should I conduct? A proposed typology and guidance for systematic reviewers in the medical
693   and health sciences. BMC Medical Research Methodology 18(1): 5.

39. City University of London (2021) Determine the key concepts of your topic. Available at:
https://libguides.city.ac.uk/SHS-Litsearchguide/frameworks. Accessed 05 Mar 2021
40. R Core Team (2020) R: a language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
41. Katale BZ, Fyumagwa RD, Mjingo EE, Sayalel K, Batamuzi EK, Matee MI, Keyyu JD, Muumba J,
Mdaki M, Mbugi EV et al (2017) Screening for bovine tuberculosis in african buffalo (*Syncerus*
caffer) in Ngorongoro conservation area, Northern Tanzania: implications for public health.
Journal of Wildlife Diseases 53(4): 711-717.
42. Meunier NV, Sebulime P, White RG, Kock R (2017) Wildlife-livestock interactions and risk areas
for cross-species spread of bovine tuberculosis. Onderstepoort Journal of Veterinary Research
84(1): 1221.
43. Wielgus E, Caron A, Bennitt E, De Garine-Wichatitsky M, Cain B, Fritz H, Miguel E, Cornélis D,
Chamaillé-Jammes S (2021) Inter-group social behavior, contact patterns and risk for pathogen
transmission in Cape Buffalo populations. The Journal of Wildlife Management 85(8): 1574-
1590.
44. Gormley AM, Anderson DP, Nugent G (2018) Cost-based optimization of the stopping threshold
for local disease surveillance during progressive eradication of tuberculosis from New Zealand
wildlife. Transboundary and Emerging Diseases 65(1): 186-196.
45. Smith GC, Budgey R (2021) Simulating the next steps in badger control for bovine tuberculosis
in England. PLOS ONE 16(3): e0248426.
46. Birch CPD, Goddard A, Tearne O (2018) A new bovine tuberculosis model for England and Wales
(BoTMEW) to simulate epidemiology, surveillance and control. BMC Veterinary Research 14(1):
273.
47. Macdonald DW, Riordan P, Mathews F (2006) Biological hurdles to the control of TB in cattle: A
test of two hypotheses concerning wildlife to explain the failure of control. Biological
Conservation 131(2): 268-286.
48. Ellwanger JH, Fearnside PM, Ziliotto M, Valverde-Villegas JM, Veiga ABGD, Vieira GF, Bach E,
Cardoso JC, Müller NFD, Lopes G et al (2022) Synthesizing the connections between
environmental disturbances and zoonotic spillover. Anais da Academia Brasileira de Ciências
94(suppl 3): e20211530.
49. Gibb R, Redding DW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, Jones KE (2020) Zoonotic
host diversity increases in human-dominated ecosystems. Nature 584(7821): 398-402.
50. Keesing F, Ostfeld RS (2021) Impacts of biodiversity and biodiversity loss on zoonotic diseases.
Proceedings of the National Academy of Sciences 118(17): e2023540118.
51. Bastida AZ, Tellez MH, Bustamante Montes LP, Jaramillo Paniagua JN, Jaimes Beni Tes MA,
Mendoza Barrera GN, Rami Rez-Dura NN (2017) Spatial analysis of bovine tuberculosis in the
State of Mexico, Mexico. Veterinaria Italiana 53(1): 39-46.
52. Gaughran A, Mullen E, Macwhite T, Maher P, Kelly DJ, Kelly R, Good M, Marples NM (2021)
Badger territoriality maintained despite disturbance of major road construction. PLOS ONE
16(9): e0242586.
53. O'Hagan MJH, McCormick CM, Collins SF, McBride KR, Menzies FD (2021) Are major roads
effective barriers for badger (*Meles meles*) movements? Research in Veterinary Science 138: 49-
52.
54. Murphy KJ, Morera-Pujol V, Ryan E, Byrne AW, Breslin P, Ciuti S (2022) Habitat availability alters
the relative risk of a bovine tuberculosis breakdown in the aftermath of a commercial forest
clearfell disturbance. Journal of Applied Ecology 59(9): 2333-2345.
55. Brunton LA, Alexander N, Wint W, Ashton A, Broughan JM (2017) Using geographically weighted
regression to explore the spatially heterogeneous spread of bovine tuberculosis in England and
Wales. Stochastic Environmental Research and Risk Assessment 31(2): 339-352.

56.   Allen AR, Ford T, Skuce RA (2021) Does *Mycobacterium tuberculosis var. bovis* survival in the
environment confound bovine tuberculosis control and eradication? A Literature Review.
Veterinary Medicine International 2021: 1-19.

57.   Brooks-Pollock E, Roberts GO, Keeling MJ (2014) A dynamic model of bovine tuberculosis spread
and control in Great Britain. Nature 511(7508): 228-231.

58.   Redpath SHA, Marks NJ, Menzies FD, O'Hagan MJH, Wilson RP, Smith S, Magowan EA, Mcclune
DW, Collins SF, Mccormick CM et al (2023) Impact of test, vaccinate and remove protocol on
home ranges and nightly movements of badgers in a medium density population. Scientific
Reports 13(1): 2592.

59.   Gaughran A, Macwhite T, Mullen E, Maher P, Kelly DJ, Good M, Marples NM (2019) Dispersal
patterns in a medium-density Irish badger population: Implications for understanding the
dynamics of tuberculosis transmission. Ecology and Evolution 9(23): 13142-13152.

60.   Palisson A, Courcoul A, Durand B (2016) Role of cattle movements in bovine tuberculosis spread
in france between 2005 and 2014. PLoS One 11(3): e0152578.

61.   Gortazar C, Diez-Delgado I, Barasona JA, Vicente J, De La Fuente J, Boadella M (2015) The wild
side of disease control at the wildlife-livestock-human interface: A review. Frontiers in
Veterinary Science 1: 27.

62.   Delahay CRJ, Smith GC, Barlow AM, Walker N, Harris A, Clifton-Hadley RS, C.L. (2007) Bovine
tuberculosis infection in wild mammals in the South-West region of England: A survey of
prevalence and a semi-quantitative assessment of the relative risks to cattle. The Veterinary
Journal 173(2): 287-301.

63.   Crispell J, Cassidy S, Kenny K, Mcgrath G, Warde S, Cameron H, Rossi G, Macwhite T, White PCL,
Lycett S et al (2020) *Mycobacterium bovis* genomics reveals transmission of infection between
cattle and deer in Ireland. Microbial Genomics 6(8).

64.   Réveillaud É, Desvaux S, Boschiroli M-L, Hars J, Faure É, Fediaevsky A, Cavalerie L, Chevalier F,
Jabert P, Poliak S et al (2018) Infection of wildlife by *Mycobacterium bovis* in france assessment
through a national surveillance system, Sylvatub. Frontiers in Veterinary Science 5: 262.

65.   Castillo L, Fernández-Llario P, Mateos C, Carranza J, Benítez-Medina JM, García-Jiménez W, and
FB-M (2011) Management practices and their association with *Mycobacterium tuberculosis*
complex prevalence in red deer populations in Southwestern Spain. Preventive Veterinary
Medicine 98(1): 58-63.

66.   Ashford RT, Anderson P, Waring L, Davé D, Smith F, Delahay RJ, Gormley E, Chambers MA,
Sawyer J, Lesellier S (2020) Evaluation of the Dual Path Platform (DPP) VetTB assay for the
detection of *Mycobacterium bovis* infection in badgers. Preventive Veterinary Medicine 180:
105005.

67.   Menzies FD, Mccormick CM, O'Hagan MJH, Collins SF, Mcewan J, Mcgeown CF, Mchugh GE, Hart
CD, Stringer LA, Molloy C et al (2021) Test and vaccinate or remove: Methodology and
preliminary results from a badger intervention research project. Veterinary Record 189(5): e248.

68.   Morera-Pujol V, Mostert PS, Murphy KJ, Burkitt T, Coad B, Mcmahon BJ, Nieuwenhuis M,
Morelle K, Ward AI, Ciuti S (2023) Bayesian species distribution models integrate presence-only
and presence–absence data to predict deer distribution and relative abundance. Ecography
2023(2): e06451.

69.   Gaidet N, Cappelle J, Takekawa JY, Prosser DJ, Iverson SA, Douglas DC, Perry WM, Mundkur T,
Newman SH (2010) Potential spread of highly pathogenic avian influenza H5N1 by wildfowl:
dispersal ranges and rates determined from large-scale satellite telemetry. Journal of Applied
Ecology 47(5): 1147-1157.

70.   Raizman EA, Rasmussen HB, King LE, Ihwagi FW, Douglas-Hamilton I (2013) Feasibility study on
the spatial and temporal movement of Samburu's cattle and wildlife in Kenya using GPS radio-
tracking, remote sensing and GIS. Preventive Veterinary Medicine 111(1): 76-80.

71.   Wearn O, Glover-Kapfer P (2017) Camera-trapping for conservation: a guide to best-practices.
United Kingdom: WWF-UK.

72. Vazquez C, Rowcliffe JM, Spoelstra K, Jansen PA (2019) Comparing diel activity patterns of
wildlife across latitudes and seasons: Time transformations using day length. Methods in
Ecology and Evolution 10(12): 2057-2066.

73. Wevers J, Beenaerts N, Casaer J, Zimmermann F, Artois T, Fattebert J (2021) Modelling species
distribution from camera trap by-catch using a scale-optimized occupancy approach. Remote
Sensing in Ecology and Conservation 7(3): 534-549.

74. Henrich M, Hartig F, Dormann CF, Kühl HS, Peters W, Franke F, Peterka T, Šustr P, Heurich M
(2022) Deer behavior affects density estimates with camera traps, but is outweighed by spatial
variability. Frontiers in Ecology and Evolution 10: 881502.

75. Kukielka VE, Barasona JA, Cowie CE, Drewe JA, Gortazar C, Cotarelo I, Vicente J (2013) Spatial
and temporal interactions between livestock and wildlife in South Central Spain assessed by
camera traps. Preventive Veterinary Medicine 112(3): 213-221.

76. Gaudiano L, Pucciarelli L, Mori E (2021) Livestock grazing affects movements and activity pattern
of Italian roe deer in Southern Italy. European Journal of Wildlife Research 67(4): 66.

77. Triguero-Ocaña R, Vicente J, Palencia P, Laguna E, Acevedo P (2020) Quantifying wildlife-
livestock interactions and their spatio-temporal patterns: Is regular grid camera trapping a
suitable approach? Ecological Indicators 117: 106565.

78. Hinojo A, Christe P, Moreno I, Hofmeister RJ, Dandliker G, Zimmermann F (2022) Estimating roe
deer density using motion-sensitive cameras in Switzerland. The Journal of Wildlife
Management 86(8): e22307.

79. Wilmers CC, Nickel B, Bryce CM, Smith JA, Wheat RE, Yovovich V (2015) The golden age of bio-
logging: how animal-borne sensors are advancing the frontiers of ecology. Ecology 96(7): 1741-
1753.

80. Williams HJ, Taylor LA, Benhamou S, Bijleveld AI, Clay TA, Grissac S, Demšar U, English HM,
Franconi N, Gómez-Laich A et al (2020) Optimizing the use of biologgers for movement ecology
research. Journal of Animal Ecology 89(1): 186-206.

81. Woodroffe R, Donnelly CA, Ham C, Jackson SYB, Moyes K, Chapman K, Stratton NG, Cartwright
SJ (2016) Badgers prefer cattle pasture but avoid cattle: implications for bovine tuberculosis
control. Ecology Letters 19(10): 1201-1208.

82. Dougherty ER, Seidel DP, Carlson CJ, Spiegel O, Getz WM (2018) Going through the motions:
incorporating movement analyses into disease research. Ecology Letters 21(4): 588-604.

83. Wilson RP, Grundy E, Massy R, Soltis J, Tysse B, Holton M, Cai Y, Parrott A, Downey LA, Qasem L
et al (2014) Wild state secrets: ultra-sensitive measurement of micro-movement can reveal
internal processes in animals. Frontiers in Ecology and the Environment 12(10): 582-587.

84. Vázquez Diosdado JA, Barker ZE, Hodges HR, Amory JR, Croft DP, Bell NJ, Codling EA (2015)
Classification of behaviour in housed dairy cows using an accelerometer-based activity
monitoring system. Animal Biotelemetry 3(1): 1-14.

85. Barasona JA, Latham MC, Acevedo P, Armenteros JA, Latham ADM, Gortazar C, Carro F, Soriguer
RC, Vicente J (2014) Spatiotemporal interactions between wild boar and cattle: implications for
cross-species disease transmission. Veterinary Research 45(1): 122.

86. Woodroffe R, Donnelly CA, Chapman K, Ham C, Moyes K, Stratton NG, Cartwright SJ (2021)
Successive use of shared space by badgers and cattle: implications for *Mycobacterium bovis*
transmission. Journal of Zoology 314(2): 132-142.

87. Shaheen MNF (2022) The concept of one health applied to the problem of zoonotic diseases.
Reviews in Medical Virology 32(4): e2326.

88. Lowry H, Lill A, Wong BBM (2013) Behavioural responses of wildlife to urban environments.
Biological Reviews 88(3): 537-549.

89. Soulsbury CD, White PCL (2015) Human–wildlife interactions in urban areas: a review of
conflicts, benefits and opportunities. Wildlife Research 42(7): 541-553.

90. Griffin LL, Haigh A, Amin B, Faull J, Norman A, Ciuti S (2022) Artificial selection in human-wildlife
feeding interactions. Journal of Animal Ecology 91(9): 1892-1905.

91.   Carrete M, Tella JL (2017) Behavioral correlations associated with fear of humans differ between
846   rural and urban burrowing owls. Frontiers in Ecology and Evolution 5: 54.
847   92.   Spiegel O, Leu ST, Bull CM, Sih A (2017) What's your move? Movement as a link between
848   personality and spatial dynamics in animal populations. Ecology Letters 20(1): 3-18.
849   93.   Gartland LA, Firth JA, Laskowski KL, Jeanson R, Ioannou CC (2022) Sociability as a personality
850   trait in animals: methods, causes and consequences. Biological Reviews 97(2): 802-816.
851   94.   Oster M, Scheel M, Muráni E, Ponsuksili S, Zebunke M, Puppe B, Wimmers K (2015) The fight-
852   or-flight response is associated with PBMC expression profiles related to immune defence and
853   recovery in swine. PLoS One 10(3): e0120153.
854   95.   Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM (2005) Superspreading and the effect of
855   individual variation on disease emergence. Nature 438(7066): 355-359.
856   96.   Barber I, Dingemanse NJ (2010) Parasitism and the evolutionary ecology of animal personality.
857   Philosophical Transactions of the Royal Society B: Biological Sciences 365(1560): 4077-4088.
858   97.   Santicchia F, Romeo C, Ferrari N, Matthysen E, Vanlauwe L, Wauters LA, Martinoli A (2019) The
859   price of being bold? Relationship between personality and endoparasitic infection in a tree
860   squirrel. Mammalian Biology 97: 1-8.
861   98.   Poulin R (2013) Parasite manipulation of host personality and behavioural syndromes. Journal
862   of Experimental Biology 216(1): 18-26.
863   99.   Ward AI, Judge J, Delahay RJ (2010) Farm husbandry and badger behaviour: Opportunities to
864   manage badger to cattle transmission of *Mycobacterium bovis*? Preventive Veterinary Medicine
865   93(1): 2-10.
866   100.   Triguero-Ocaña R, Martínez-López B, Vicente J, Barasona JA, Martínez-Guijosa J, Acevedo P
867   (2020) Dynamic network of interactions in the wildlife-livestock interface in mediterranean
868   Spain: An epidemiological point of view. Pathogens 9(2): 120.
869   101.   Wilber MQ, Pepin KM, Campa H, Hygnstrom SE, Lavelle MJ, Xifara T, Vercauteren KC, Webb CT
870   (2019) Modelling multi-species and multi-mode contact networks: Implications for persistence
871   of bovine tuberculosis at the wildlife–livestock interface. Journal of Applied Ecology 56(6): 1471-
872   1481.
873   102.   Fielding HR, Silk MJ, McKinley TJ, Delahay RJ, Wilson-Aggarwal JK, Gauvin L, Ozella L, Cattuto C,
874   McDonald RA (2021) Spatial and temporal variation in proximity networks of commercial dairy
875   cattle in Great Britain. Preventive Veterinary Medicine 194: 105443.
876   103.   Tratalos JA, Madden JM, Mcgrath G, Graham DA, Collins ÁB, More SJ (2020) Spatial and network
877   characteristics of Irish cattle movements. Preventive Veterinary Medicine 183: 105095.
878   104.   Silk MJ, Croft DP, Delahay RJ, Hodgson DJ, Boots M, Weber N, Mcdonald RA (2017) Using social
879   network measures in wildlife disease ecology, epidemiology, and management. BioScience
880   67(3): 245-257.
881   105.   Farine DR, Whitehead H (2015) Constructing, conducting and interpreting animal social network
882   analysis. Journal of Animal Ecology 84(5): 1144-1163.
883   106.   Kaur P, Ciuti S, Ossi F, Cagnacci F, Morellet N, Loison A, Atmeh K, McLoughlin P, Reinking AK,
884   Beck JL et al (2023) Assessing bias and robustness of social network metrics using GPS based
885   radio-telemetry data. Preprint at bioRxiv:2023.2003.2030.534779.
886   107.   Sosa S, Sueur C, Puga-Gonzalez I (2021) Network measures in animal social network analysis:
887   Their strengths, limits, interpretations and uses. Methods in Ecology and Evolution 12(1): 10-21.
888   108.   Newton-Cross G, White PCL, Harris S (2007) Modelling the distribution of badgers *Meles meles*:
889   comparing predictions from field-based and remotely derived habitat data. Mammal Review
890   37(1): 54-70.
891   109.   Elith J, Leathwick JR (2009) Species Distribution Models: Ecological explanation and prediction
892   across space and time. Annual Review of Ecology, Evolution, and Systematics 40(1): 677-697.
893   110.   Lindgren F, Rue H (2015) Bayesian spatial modelling with R-INLA. Journal of Statistical Software
894   63(19): 1-25.

895   111.   Elith J (2019) Machine learning, random forests, and boosted regression trees. In: Brennan LA,
896   Tri AN, and Marcot BG (Eds) Quantitative analyses in wildlife science. Johns Hopkins University
897   Press, Baltimore, Maryland, USA: 281-287.
898   112.   Isaac NJB, Jarzyna MA, Keil P, Dambly LI, Boersch-Supan PH, Browning E, Freeman SN, Golding
899   N, Guillera-Arroita G, Henrys PA et al (2020) Data integration for large-scale models of species
900   distributions. Trends in Ecology & Evolution 35(1): 56-67.
901   113.   Bakka H, Rue H, Fuglstad GA, Riebler A, Bolin D, Illian J, Krainski E, Simpson D, Lindgren F (2018)
902   Spatial modeling with R-INLA: A review. WIREs Computational Statistics 10(6): e1443.
903   114.   Tracey JA, Bevins SN, Vandewoude S, Crooks KR (2014) An agent-based movement model to
904   assess the impact of landscape fragmentation on disease transmission. Ecosphere 5(9): 119.
905   115.   Gaughran A, Kelly DJ, Macwhite T, Mullen E, Maher P, Good M, Marples NM (2018) Super-
906   ranging. A new ranging strategy in European badgers. PLoS One 13(2): e0191818.
907   116.   O'Hagan MJH, Gordon AW, Mccormick CM, Collins SF, Trimble NA, Mcgeown CF, Mchugh GE,
908   Mcbride KR, Menzies FD (2021) Effect of selective removal of badgers (*Meles meles*) on ranging
909   behaviour during a 'Test and Vaccinate or Remove' intervention in Northern Ireland.
910   Epidemiology and Infection 149: 1-31.
911   117.   Byrne AW, Acevedo P, Green S, O'Keeffe J (2014) Estimating badger social-group abundance in
912   the Republic of Ireland using cross-validated species distribution modelling. Ecological Indicators
913   43: 94-102.
914   118.   Abdou M, Frankena K, O'Keeffe J, Byrne AW (2016) Effect of culling and vaccination on bovine
915   tuberculosis infection in a European badger (*Meles meles*) population by spatial simulation
916   modelling. Preventive Veterinary Medicine 125: 19-30.
917   119.   Barroso P, Breslin P, McGrath G, Jamie MM, Tratalos JA, More SJ, Ryan E, Byrne AW, Barrett D
918   (2022) Is there an association between road building and bovine tuberculosis herd risk? A three
919   time-point study in Ireland, 2011–2019. Preventive Veterinary Medicine 198: 105542.
920   120.   Zimmerman J, Fealy RM, Lydon K, Mockler EM, O'Brien P, Packham I, Smith G, Green S (2016)
921   The Irish Land-Parcels Identification System (LPIS)–Experiences in ongoing and recent
922   environmental research and land cover mapping. Biology and Environment: Proceedings of the
923   Royal Irish Academy 116B(1): 53-62.
924   121.   Martin SW, O'Keeffe J, Byrne AW, Rosen LE, White PW, McGrath G (2020) Is moving from
925   targeted culling to BCG-vaccination of badgers (*Meles meles*) associated with an unacceptable
926   increased incidence of cattle herd tuberculosis in the Republic of Ireland? A practical non-
927   inferiority wildlife intervention study in the Republic of Ireland (2011-2017). Preventive
928   Veterinary Medicine 179: 105004.

Table 1: Description of the 18 questions (and related variables) used to screen bTB papers published in the period between 2017-2022. The variables have been divided into three groups depending on the question type. Variables looking at general aspects of the paper (e.g., type of host species included in the study) were grouped as *general characteristics*. Variables concerning the type of data analysis conducted in the studies were grouped as *data analysis*. Finally, variables focused on the type of epidemiological analysis conducted in the papers of interest were grouped as *epidemiology*.

Variables		
Variable Types	**Variable Names**	**18 questions**
General Characteristics	Study Type	Is the study on cattle and/or wildlife?
	Wildlife Species	Which wildlife species are included in the study: i.e., cervid, badger, wild boar or other (brush-tailed possum, wild buffalo)?
	Management	Does the study look at possible management solutions for controlling the spread of bTB from wildlife to livestock and their efficacy?
	Data Type	Is the study based on empirical data and/or simulations?
Data Analysis	Spatial Analysis – Cattle	Does the study conduct spatial analysis in cattle (e.g., looking at risk and probability of infection)?
	Spatial Analysis – Wildlife	Does the study conduct spatial analysis in wildlife (e.g., using GPS collars)? (Which does not imply testing for disease but rather whether the spatial behaviour of the potential wildlife host has been studied)
	Spatial Scale	Does the study include a spatial scale and if so which type: local, regional, national and international?
	Temporal Scale	What type of temporal scale does the study include, if any? Temporal scale type: daily variability, snapshot (1 point in time), intra-annual variability, interannual variability, future scenario predictions.
	Farm Environment	Does the data analysis include variables that explain the environment characteristics inside the farm (e.g., herd data, habitat within the farm)?
	Outside Farm Environment	Does the data analysis include variables that explain the environmental characteristics around the farm (e.g., habitat characteristics)?
	Weather	Does the data analysis include climate variables (e.g., rainfall, soil humidity)?
	Human Perturbation	Does the data analysis include human perturbation variables (e.g., culling, road construction, forest clearfell)?
Epidemiology	Intra and Interspecies Interactions	Does the study look at interactions between animals and, if so, does it look at intraspecies or interspecies interactions?
	Interaction Type	If the study looks at interactions, what type of interactions does it look at? Type of interactions: indirect or direct.
	Interaction Equipment	If the study looks at interactions, how were the data collected? Collection type: direct observations, technological tools (GPS, proximity loggers, camera traps), simulation.
	Interaction Methodology	If the study looks at interactions, what methodology was used for the data analysis? Methodology type: social network analysis or other (e.g., linear regression, logistic regression).
	Direction	Does the study include a direction of transmission (e.g., using whole genome sequencing of genetic samples) and/or a clear and direct path of contacts between animals (e.g., using a social network)?
	Epidemiological Modelling	Does the study look at compartmental models (i.e., SIR) and/or transmission rates?

Table 2: Eligibility criteria used in the selection of papers for the systematic scoping review. The criteria follow a PEO (Population, Exposure, Outcomes/Themes) structure.

Eligibility Criteria			
PEO Criteria	**Explanation**	**Scoping Review Criteria**	**Eligible Paper**
Population	Who is the focus of the scoping review?	Cattle, badger, deer, elk, wild boar, wild buffalo, brushed-tailed possum (i.e., known bTB hosts)	Contains at least one of the species of interest as a study species.
Exposure	What issue is the focal point of the scoping review?	Bovine tuberculosis (i.e., *Mycobacterium bovis*)	Focal point of the paper must be bovine tuberculosis, more specifically the most common cause: *Mycobacterium bovis*.
Outcomes/ Themes	What themes, in relation to the issue, will the scoping review focus on?	● **Risk**: Does the study assess or evaluate potential risks of spreading bTB and its drivers (biotic and/or abiotic factors)? ● **Management**: Does the study look at potential management solutions for controlling the spread of bTB from wildlife to livestock and their efficacy? ● **bTB occurrence**: Does the study look at the presence of bTB in cattle and/or reservoir hosts at a spatial and/or temporal level? ● **bTB spread**: Does the study look at the mechanisms behind the active spread of bTB between individuals of the same species and/or interspecies transmission?	Focuses on at least one of the themes of interest.

Figure legends

Fig. 1: Key host species and topics of interest we screened for in the bovine tuberculosis scientific literature published in the period between 2017-2022. bTB host species include cattle as well as a range of wild species: badger, wild boar, cervid species (with the following species identified in the literature screened: white-tailed deer, red deer, fallow deer, roe deer and wapiti elk), brush-tailed possum and wild buffalo. The circles on the outside illustrate the key information sought in peer-reviewed papers dealing with bTB, which has been expanded and clarified in **Table 1**: type of data collected by researchers; whether spatial analyses were carried out (i.e. in cattle and or wildlife); what type of spatial and temporal scales were considered; whether environmental variables were taken into account (i.e. environment in the farm, environment around the farm and/or climate variables); whether the methodological approach captured the direction of disease transmission; whether the study used common epidemiological modelling techniques (i.e. compartmental models, transmission rates), or whether the study included intra/interspecies interactions in their methodology (i.e. what type of interactions did they look at – e.g. direct and/or indirect, what type of equipment was used to get interactions data and what methodology was used to analyse the data); finally, if human perturbations

(i.e. forest felling, culling, vaccination) were taken into account when looking at variables affecting bTB spread, and management solutions to offset the spread of bTB, if any. Animal silhouettes were downloaded from PhyloPic (https://www.phylopic.org/). Cattle, cervid, brushed-tailed possum and wild boar silhouettes are under: CC0 1.0 Universal (CC0 1.0) Public Domain Dedication. Buffalo silhouette is by Jan A. Venter, Herbert H. T. Prins, David A. Balfour & Rob Slotow (vectorized by T. Michael Keesey) under: Attribution 3.0 Unported (CC BY 3.0), link: https://creativecommons.org/licenses/by/3.0/. Badger silhouette is by Anthony Caravaggi under: Attribution-NonCommercial-ShareAlike 3.0 Unported (CC BY-NC-SA 3.0), link: https://creativecommons.org/licenses/by-nc-sa/3.0/.

Fig. 2: Cascade diagram of the process used in the selection of relevant papers.

Fig. 3: World map showing number of papers screened per country. Number of papers per continent: Europe (103), Africa (26), Asia (19), South America (11), North America (8), Oceania (7).

Fig. 4: Number of papers screened and reporting data on **a)** study species type (whether the study was on cattle and/or wildlife), **b)** wildlife species, **c)** management (whether a paper investigated potential management solutions and their efficacy), **d)** and data type. Animal silhouettes were downloaded from PhyloPic (https://www.phylopic.org/). Cattle, cervid, brushed-tailed possum and wild boar silhouettes are under: CC0 1.0 Universal (CC0 1.0) Public Domain Dedication. Buffalo silhouette is by Jan A. Venter, Herbert H. T. Prins, David A. Balfour & Rob Slotow (vectorized by T. Michael Keesey) under: Attribution 3.0 Unported (CC BY 3.0), link: https://creativecommons.org/licenses/by/3.0/. Badger silhouette is by Anthony Caravaggi under: Attribution-NonCommercial-ShareAlike 3.0 Unported (CC BY-NC-SA 3.0), link: https://creativecommons.org/licenses/by-nc-sa/3.0/.

Fig. 5: Number of papers screened and reporting data on **a)** spatial analysis of cattle (whether the study included any type of spatial analysis), **b)** spatial analysis of wildlife, **c)** spatial scale, and **d)** temporal scale.

Fig. 6: Number of papers screened and reporting data on **a)** in-farm environment (whether the paper analysis included variables explaining environmental characteristics inside the farm), **b)** outside farm environment, **c)** human perturbations (whether the paper analysed the effect of human disturbances on bTB transmission dynamics), and **d)** weather variables.

Fig. 7: Number of papers screened and reporting data on **a)** interactions (inclusion of interaction analysis i.e., intra- and/or interspecies interactions), **b)** interaction type, **c)** the way interactions were monitored, and **d)** interaction data analysis statistical approach.

Fig. 8: Number of papers screened and reporting data on **a)** direction of transmission (whether this was analysed in the paper, e.g., transmission across species), **b)** epidemiological modelling (i.e., papers included compartmental models and/or transmission rates in the analysis).