Dynamic PRC1-CBX8 stabilizes a porous structure of chromatin condensates

Michael Uckelmann[1], Vita Levina[1], Cyntia Taveneau[1,2], Xiao Han Ng[1], Varun Pandey[3],

Jasmine Martinez[4], Shweta Mendiratta[4], Justin Houx[5], Marion Boudes[1], Hari Venugopal[6],

Sylvain Trépout[6], Qi Zhang[1], Sarena Flanigan[1], Minrui Li[1,7], Emma Sierecki[5], Yann Gambin[5],

Partha Pratim Das[3], Oliver Bell[4], Alex de Marco[1,2,8], Chen Davidovich[1,9*]

[1] Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute,
Faculty of Medicine, Nursing and Health Sciences, Monash University; Clayton, Victoria,
3800, Australia.

[2] ARC Centre of Excellence in Advanced Molecular Imaging, Monash University; Clayton,
Victoria, 3800, Australia.

[3] Department of Anatomy and Developmental Biology, Development and Stem Cells
Program, Monash Biomedicine Discovery Institute, Monash University; Clayton, Victoria,
3800, Australia.

[4] Departments of Biochemistry and Molecular Medicine, and Stem Cell and Regenerative
Medicine, Norris Comprehensive Cancer Center, Keck School of Medicine, University of
Southern California, Los Angeles, CA 90033, USA.

[5] EMBL Australia Node for Single Molecule Science and School of Biomedical Sciences,
Faculty of Medicine, The University of New South Wales, Sydney, NSW, 2052 Australia.

[6] Ramaciotti Centre for Cryo-Electron Microscopy, Monash University, Monash, Victoria,
Australia.

[7] Faculty of Information Technology, Monash University; Clayton, Victoria, 3800, Australia.

[8] Simons Electron Microscopy Center, New York Structural Biology Center, New York 10027
NY, United States of America

[9] EMBL-Australia; Clayton, Victoria, 3800, Australia.

* Correspondence: Chen.Davidovich@monash.edu

Abstract

The compaction of chromatin is a prevalent paradigm in gene repression. Chromatin
compaction is commonly thought to repress transcription by restricting chromatin
accessibility. However, the spatial organisation and dynamics of chromatin compacted by
gene-repressing factors are unknown. Using cryo-electron tomography, we solved the three-
dimensional structure of chromatin condensed by the Polycomb Repressive Complex 1
(PRC1) in a complex with CBX8. PRC1-condensed chromatin is porous and stabilised through
multivalent dynamic interactions of PRC1 with chromatin. Mechanistically, positively
charged residues on the internally disordered regions (IDRs) of CBX8 mask negative charges

on the DNA to stabilize the condensed state of chromatin. Within condensates, PRC1

remains dynamic while maintaining a static chromatin structure. In differentiated mouse

embryonic stem cells, CBX8-bound chromatin remains accessible. These findings challenge

the idea of rigidly compacted polycomb domains and instead provides a mechanistic

framework for dynamic and accessible PRC1-chromatin condensates.

Main

Chromatin structure is intricately linked to transcriptional activity[1]. Compacted or "closed"

chromatin is generally associated with inhibition of transcription while "open", more

accessible chromatin is more prone to being transcribed[1]. Polycomb Repressive Complex 1

(PRC1) is a repressive chromatin modifier critical for organismal development[2,3]. PRC1 has

been proposed to inhibit gene expression by tightly compacting chromatin in a process that

often is considered to restrict chromatin accessibility[2–11]. However, direct evidence for

PRC1-compacted chromatin being inaccessible is sparse and mechanistic explanations

remain unsatisfactory (reviewed in[11]). Furthermore, recent studies show that changes in

chromatin accessibility are more gradual than the simple binary classification into "open"

and "closed" chromatin suggests[12–15]. A challenge in consolidating these seemingly

contradictory findings is limited information into how PRC1 influences the three dimensional

structure of chromatin.

PRC1 complexes can include one of five different chromobox proteins (CBX), all homologous

to the fly Polycomb (Pc)[16]. The CBX protein CBX2 forms condensates through liquid-liquid

phase separation, providing a potential mechanism for the compartmentalization of

facultative heterochromatin[9,10]. Phase separation is emerging as a mechanism for chromatin

organisation through the association of self-similar domains[17]. Chromatin can form

condensates in the presence of divalent cations and histone tails[17–20]. Within these

condensates, chromatin has variably been described as liquid-like, formed through liquid-

liquid phase separation[17,19], or as solid[20]. A recent structure of liquid-liquid phase-separated

chromatin, condensed by magnesium cations without protein binding partners, revealed

that nucleosomes organise into irregular assemblies[19]. The lack of apparent periodicity in

chromatin geometry has also been noted in computational simulations and in first attempts

to image chromatin in cells by cryo-electron tomography (cryo-ET)[21,22]. However, the

structural arrangement of chromatin condensed by a repressive factor remained unknown.

Herein we describe the three dimensional cryo-ET structure of chromatin condensed by a

polycomb-repressive complex. We focus on a PRC1 complex that includes CBX8 (PRC1[C8]), a

chromobox protein that is upregulated during cell differentiation[23] and has oncogenic

potential[24,25]. We show that dynamic interactions between PRC1[C8] and chromatin promote

condensates through phase separation. Mechanistically, positive charges on the internally

disordered regions (IDRs) of CBX8 are required for DNA binding and chromatin

condensation. Contrary to expectations, PRC1-condensed chromatin is not tightly

compacted but stabilises a porous chromatin structure that allows largely unhindered
diffusion of PRC1^{C8}.

Results

PRC1-chromatin condensates are porous and accessible

To determine the structure of polycomb-compacted chromatin and the mechanisms of
polycomb-driven chromatin compaction, we reconstituted the system *in vitro*. The
reconstitution included a chromatinized polycomb target gene (3,631 bp DNA) with a
sequence from the human ATOH1 locus, which can harbour roughly up to 20 nucleosomes,
assuming 150-200 bp per nucleosome. This construct is referred to as chromatin hereafter.
We used a native DNA sequence for chromatin reconstitution, as regular spacing using
artificial nucleosome stabilising sequences were previously reported to drive the liquid-
liquid phase separation of chromatin[17]. The nucleosomes on the chromatin that we
reconstituted are not evenly phased (Extended Data Fig. 1a). The purified recombinant PRC1
complex is composed of RING1B, BMI1 and CBX8 (PRC1^{C8}) (Fig. 1a). The PRC1^{C8} complex is
pure (Fig. 1b), monodispersed (Fig. 1c) and retains H2A ubiquitylation activity comparable to
the RING1b-BMI1 heterodimer (Fig. 1d). This also applies to all other protein complexes
used in this study (Extended Data Fig. 2 and 1b,c).

When combined, chromatin and PRC1^{C8} were sufficient to form spherical phase-separated
condensates, apparent in differential interference contrast (DIC) and fluorescence imaging
(Fig. 1e). Using two different fluorescence labels, we confirmed the presence of both
chromatin and PRC1^{C8} within the same condensates (Fig. 1e). Importantly, both PRC1^{C8} and
chromatin are necessary for chromatin condensation, while the individual components do
not phase-separate (Fig. 1e). PRC1^{C8}-chromatin condensates are preserved on an EM grid
after vitrification (Fig. 1f). To study these structures by cryo-electron tomography, we
reduced the salt concentration and increased chromatin concentration (see Methods
section). This was done to decrease condensate size and abundance, which was necessary
for high-quality data collection (Fig. 1g). We used a PRC1 complex with MBP-tagged CBX8
for cryo-tomography. The MBP tag does not affect condensation, as condensates still form
after tag cleavage (Extended Data Fig. 1d). We collected tomograms near the borders of
condensates, to observe the boundary conditions (Extended Data Fig. 3a for an example of a
condensate on the grid). Subtomogram averaging allowed us to identify the orientation and
position of individual nucleosomes in the tomographic volume (Extended Data Fig. 3b-d).
The reconstruction reveals a dense network of hundreds of nucleosomes with a distinct
condensate boundary (Fig. 1g and Movie S1). We could not unambiguously assign density to
PRC1^{C8}, possibly because it adapts various conformations while simultaneously using
multiple surfaces to interact with chromatin (more below). The final structure reflects the
arrangement of nucleosomes in PRC1^{C8}-chromatin condensate (Fig. 1g, second panel).
Unexpectedly, the structure shows that PRC1^{C8} does not compact nucleosomes into an

impassable barrier. Instead, PRC1[C8] rather stabilises chromatin in a porous mesh-like structure (Fig. 1g). Analysing the orientation of individual nucleosomes towards their neighbouring nucleosomes shows no obvious orientation bias (Extended Data Fig. 3c,d). We conclude that PRC1[C8] does not induce a substantial inter-nucleosome orientation bias, but rather supports forming a porous chromatin structure.

We next wished to determine the size of macromolecules that could diffuse into PRC1-chromatin condensates. We used the condensate structure to calculate solvent-excluded volumes[26] with variable probe radii ranging from 0.2 nm to 20 nm (Fig. 1h,i). Interestingly, the analysis shows that the condensate is accessible for macromolecules of a considerable size of up to 8 nm in radius (equivalent to approximately 600 kDa). Small macromolecules (<10 kDa), with radii below 2 nm, would have enough room to access every single nucleosome. Conversely, access is increasingly restricted for molecules with a radius above 8 nm (approximately 600 kDa). This suggests that PRC1-chromatin condensates are surprisingly accessible and that PRC1[C8] itself would be able to move within these condensates largely unhindered.

To compare the structure of PRC1[C8]-chromatin condensates to PRC1[C8]-free chromatin, we generated cryo-tomograms of chromatin without PRC1[C8] (Fig. 1j, movie S2). Low magnification cryo-EM images show that condensation happens only in the presence of PRC1[C8] (Extended Data Fig. 4a,b) and condensates are preserved on the EM grid. The structure of chromatin in the absence of PRC1[C8] is less dense than in the presence of PRC1[C8], with distances between neighbouring nucleosomes that are on average significantly longer (Fig. 1j,k). These results confirm that PRC1[C8] facilitates large-scale chromatin restructuring.

We observe some sporadic areas of high nucleosome density, even without PRC1 (Extended Data Fig. 4c,d). The median distance to the next neighbouring nucleosomes in these sporadic dense PRC1[C8]-free condensates are very similar to distances measured in the PRC1[C8]-chromatin condensates (9.8 nm and 10.9 nm, respectively, Extended Data Fig. 4d). This matches closely to distances reported for chromatin condensed by MgCl2, where the radial distribution function of nucleosomes peaked at 10.6 nm[19]. At the nuclear periphery in cells, the median distance between neighbouring nucleosomes is about 12 nm, which is again remarkably similar[27]. Overall, this raises the possibility that PRC1 thermodynamically stabilises a naturally-occurring condensed chromatin state, rather than actively compacts chromatin. By doing so, PRC1 may cause multiple compacted arrays to cluster together more often.

PRC1[C8] is mobile and chromatin is static within PRC1-chromatin condensates

To gain further insights into PRC1[C8]-chromatin condensation, we tested the dependency of condensation on the concentration of PRC1 and chromatin. Condensates form under close to physiological monovalent salt concentrations of 122.5 mM (90 mM KCl and 32.5 mM

NaCl), at PRC1^{C8} concentrations as low as 250 nM and are dependent on PRC1^{C8} (Fig. 2a). At

a high concentration of PRC1^{C8} (2,000 nM), most efficient condensation occurs at high

chromatin concentration (850 nM nucleosome concentration). Yet, at lower concentration

of PRC1^{C8} (250 nM), ideal condensation appears at lower chromatin concentration and the

condensation efficiency is then reduces when the chromatin concentration increases (Fig

2a). This is possibly because at high chromatin concentration the large amount of potential

binding sites for PRC1^{C8} reduces the average per-site-occupancy of PRC1^{C8}. This may lead to

less efficient phase separation. A similar observation was recently made for a PRC1 complex

with CBX2 and PHC2[28]. Altering the salt concentrations confirms that PRC1^{C8}-chromatin

condensates most readily form close to physiological salt concentration (Extended Data Fig.

5). We conclude that PRC1^{C8} is sufficient to drive the formation of the chromatin

condensates under physiologically relevant conditions. Furthermore, the efficiency of phase

separation depends on the ratio of PRC1 to chromatin. We next asked if chromatin and

PRC1 show different dynamics within the condensates.

Fluorescence recovery after photobleaching (FRAP) of PRC1^{C8}-chromatin condensates shows

fast recovery kinetics for GFP-labelled PRC1 (Fig. 2b, in green; $T_{1/2}$ = 71 ± 8 s). Conversely, we

observed a very slow recovery for Cy5-labelled chromatin (Fig. 2b, in red). The GFP signal

does not recover up to 100 % but rather plateaus at 72 ± 2%. This is possibly because a

substantial percentage of the condensate has been bleached, while redistribution of GFP-

labelled PRC1^{C8} within the same condensate likely drives fluorescence recovery during the

monitored timescale. However, we cannot exclude the possibility that incomplete recovery

is due to an immobile fraction of CBX8. To exclude the possibility that a large protein tag on

CBX8 prevents it from remaining static on chromatin we used a synthetic fluorescence dye

to sparsely label random lysine residues of PRC1^{C8}. We then compared FRAP recovery for

two different PRC1 complexes: one complex included CBX8 with an N-terminal MBP tag and

the other included an untagged CBX8 (Extended Data Fig. 6a,b). The PRC1^{C8} with untagged

CBX8 is still dynamic (Extended Data Fig. 6a, $T_{1/2}$ = ~1300 s), albeit recovery is slower when

compared to MBP-tagged PRC1^{C8} (Extended Data Fig. 6b, $T_{1/2}$ = 362 ± 29 s) and the GFP-

tagged protein (Fig. 2b; $T_{1/2}$ = 71 ± 8 s). Chromatin remains static in all samples. These

results indicate that although the tagging and labelling strategies affect the dynamics of the

system quantitatively, its overall behaviour remains qualitatively the same: within

condensates, PRC1 is mobile while chromatin itself is static (Fig. 2c and Extended Data Fig.

6). This confirms that PRC1^{C8} can diffuse within PRC1^{C8}-chromatin condensates, in line with

our structural analysis (Fig. 1h,i). Recent publications have been at odds as to the solid or

liquid state of chromatin[17,20]. Our data suggests that chromatin behaves as a solid-like

material when condensed by PRC1^{C8}.

We next wished to gain insights into PRC1^{C8} -chromatin condensate formations at low

protein concentrations that better resemble physiological concentrations. To this end, we

employed a single-molecule confocal microscope that allows tracking of individual

condensates through a confocal volume[29] (Extended Data Fig. 7a,b). We used GFP-labelled

PRC1[C8], where GFP peaks indicate the formation of bright protein assemblies. Importantly,
this system allows the detection of protein assemblies smaller than what can be identified
by standard fluorescence microscopes[30]. In the presence of chromatin, assemblies are
observed at PRC1[C8] concentrations as low as 62.5 nM (Extended data Fig. 7c). This indicates
that condensates can form at physiologically relevant PRC1 concentrations, previously
estimated as 130 nM in polycomb bodies in cells[31]. In the absence of chromatin, GFP peaks
are only detected sporadically, even at the highest PRC1[C8] concentration (Extended Data Fig.
7 b,c). Overall, this data indicates that PRC1[C8]-chromatin condensates form under
physiologically relevant PRC1[C8] concentration, but do not form without chromatin.

Multivalent interactions between PRC1[C8] and chromatin induce phase separation

A scaffold-client based phase separation model has recently been proposed for PRC1-CBX2
complexes[28,35], where CBX2[35] or chromatin[28] act as a scaffold that induces phase separation
of PRC1 proteins. Since PRC1[C8] is insufficient to phase-separate without chromatin (Fig 1e),
we wished to test if chromatin might act as a scaffold that concentrates PRC1[C8] and induces
phase separation. To test this model, we probed the different interaction sites between
PRC1 and chromatin. The whole PRC1[C8] complex (RING1b, BMI1 and CBX8) is necessary and
sufficient to condense chromatin, while the PRC1 core or CBX8 alone do not condense
chromatin (Fig. 3a). This suggests multivalent interactions between the PRC1[C8] complex and
chromatin, involving different chromatin interacting surfaces in both PRC1 and CBX8. To
identify the different interaction sites, we first used crosslinking mass spectrometry (XL-MS)
to probe for protein-protein interactions within PRC1[C8]–chromatin condensates (Fig. 3b and
Supplementary Data 4; PRC1 with an MBP-tagged CBX8 was used). As expected, we
identified extensive crosslinks between the RING domains of RING1B and BMI1. RING1B and
BMI1 did not crosslink to histones. This is likely because these proteins bind to the acidic
patch of the nucleosome[36], which is unlikely to be crosslinked by the BS3 crosslinker that
reacts preferentially with lysine residues. The results show multiple crosslinks from the CBX8
chromodomain to the H3 histone tail (Fig. 3b), indicative of binding. Interactions between
CBX-proteins and H3K27me3-modified H3-histone tails have been proposed to recruit PRC1
to chromatin modified by PRC2[2,3]. However, CBX8 did crosslink to unmethylated H3 tails
(Fig. 3b) and a trimethyl-lysine analogue (MLA) at H3K27 did not improve the chromatin-
condensation activity of PRC1[C8] (Fig. 3d). We conclude that H3K27me3 is not necessary for
the chromatin condensation activity of PRC1[C8] and that the H3 histone tail, even if
unmodified, provides an interaction site for PRC1 on chromatin.

We also observe extensive self-crosslinks in the IDRs of CBX8 (Fig. 3b). This may indicate
inter- or intramolecular interaction within this flexible lysine-rich region. However, these
self-crosslinks within the IDRs appear even in the absence of chromatin, suggesting that
they are not related to condensation (Extended Data Fig. 8 and Supplementary Data 3).

DNA has previously been shown to bind CBX8[37] and could provide another interaction site
for PRC1[C8] on chromatin. We tested DNA binding in solution using a 24bp double stranded
DNA probe and found that CBX8 is necessary for the DNA-binding activity of PRC1[C8] (Fig. 3c).
Hence, CBX8 binding to DNA provides a second interaction site of PRC1 with chromatin.

We conclude that PRC1 interacts with chromatin via at least three distinct sites: PRC1[C8]
binds to DNA and the H3 tail via CBX8, as shown herein (Fig. 3), and binds the nucleosome
acidic patch via RING1b-BMI1 as shown elsewhere[36]. These multivalent interactions would
have to change dynamically while PRC1 maintains the condensed state of chromatin and
diffuses through it at the same time (Fig. 2b). Collectively, we propose that PRC1 induces
chromatin condensation via phase separation, through dynamic multivalent interactions
between PRC1 and chromatin.

DNA binding by the CBX8 IDRs is required for efficient phase separation

To test how different PRC1[C8]-chromatin interaction sites affect phase separation, we
generated several different chromatin and CBX8 mutants (Fig. 4a and Extended Data Fig. 9).
Removing the CBX8 chromodomain (PRC1[C8ΔChromo]), which interacts with the H3 tail[37,38] (Fig.
3b), does not have a significant effect on phase separation (Fig. 4b,c). We then mutated 21
positively charged residues in the CBX8 IDRs to alanine (PRC1[C8KR21A]). PRC1[C8KR21A] is
analogous to a phase separation-deficient CBX2 mutant that was previously studied[10].
Accordingly, PRC1[C8KR21A] shows a clear defect in phase separation activity (Fig. 4b,c).

To further dissect the mechanism, we tested the DNA binding activity of PRC1[C8KR21A] and
PRC1[C8ΔChromo] against a 24 bp double-strand DNA probe with a sequence from a polycomb-
target gene. Both mutants are defective in DNA binding (Fig 4d). Hence, while DNA binding
of the chromodomain was reported previously[37], our data now suggests that the IDRs of
CBX8 are also contributing to DNA binding. We hypothesise that electrostatic interactions
between the negatively charged DNA to positive charges in the IDR lead to charge screening,
which promotes phase separation of chromatin (Fig. 4e).

We next set to determine if the H2A acidic patch of the nucleosome affects PRC1[C8]-
chromatin condensation. For that, we mutated residues in the H2A acidic patch that were
previously shown to interfere with the interaction between PRC1 and the nucleosome[36]
(Extended Data Fig. 9). We observed a change in the condensate morphology (Extended
Data Fig. 9a): while PRC1[C8]-chromatin condensates appear spherical, the condensates with
mutated acidic patch chromatin adapt elongated and branched structures (Extended Data
Fig. 9a; most apparent at lower PRC1[C8] concentrations). The effects of the PRC1[C8KR21A]
mutant and the acidic patch chromatin mutant are additive (Extended Data Fig. 9b),
suggesting that both interaction sites contribute to phase separation independently. The
PRC1[C8ΔChromo] mutant does not substantially affect phase separation, regardless of the
chromatin used (Extended Data Fig. 9c). Overall, our data supports a model where PRC1[C8]

uses its chromodomain and IDRs to bind DNA. Then, PRC1^{C8} condenses chromatin through

interactions between the IDRs and DNA and, independently, between PRC1 and the acidic

patch on the nucleosomes (Fig 4e).

CBX8 binding sites on chromatin in mouse embryonic stem cells are accessible

Given the porous structure of PRC1^{C8}-condensed chromatin in vitro (Fig. 1) and the dynamic

diffusion of PRC1^{C8} within condensates (Fig. 2b), we next wished to probe for the

accessibility of PRC1^{C8}-bound chromatin in cells. We carried out the Assay for Transposase

Accessible Chromatin (ATAC-seq) in differentiated mESC, combined with ChIP-seq for CBX8

and H3K27me3. We used differentiated mESC, because CBX8 is expressed at very low levels

in pluripotent mESC and is upregulated during retinoic acid-induced cell differentiation ([23]

and Fig. 5a). The DNA-loaded Tn5 used in ATAC-seq experiments forms a dimeric complex of

approximately 130 kDa with an estimated hydrodynamic radius[39] of 4.6 nm (based on PDB

code 1MUH[40]). In agreement with the accessibility analysis *in vitro* (Fig. 1h), the majority of

CBX8 ChIP-seq peaks in cells overlapped with ATAC-seq peaks (Figure 5b), indicating they

are accessible to Tn5. This observation was persistent across the genome, where ATAC-seq

peaks are co-localised with CBX8 and H3K27me3 peaks (Fig. 5c), indicating that CBX8-target

genes are largely accessible. The insufficiency of CBX8 to restrict chromatin accessibility is

further supported by the similar ATAC-seq profiles of wildtype and *Cbx8* knockout mESCs

(Fig. 5c,f, compare blue to orange). Hence, although the overall chromatin accessibility is

reduced during mESC differentiation (Fig. 5d,e), in agreement with previous works,[41] this

process is not dependent on CBX8. Collectively, we show that CBX8-bound polycomb-

repressed chromatin is largely accessible in differentiated mESCs (Fig. 5b-f).

Data thus far suggest that the deletion of CBX8 does not change chromatin accessibility in

mESCs (Fig.5c,f). Yet, other CBX proteins could potentially compensate for the loss of CBX8

in endogenous polycomb target genes. Therefore, we wished to test CBX8 in a system

where its recruitment is sufficient to trigger gene repression. To this end, we used a mESC

line that expresses a TetR-CBX8 fusion and includes a GFP reporter cassette downstream of

a TetO DNA binding array, stably integrated at chromosome 15 (coordinates: mm10 chr15:

79,013,675; Fig. 5g) [42]. In the absence of doxycycline (Dox), the TetR-CBX8 fusion is recruited

to the TetO-GFP reporter together with RING1B and initiates transcriptional repression (Fig.

5g and Extended Data Fig. 10a). However, GFP repression is reversible upon Dox addition,

which causes release of both TetR-CBX8 and RING1B from the TetO DNA binding array (Fig.

5g and Extended Data Fig. 10a).

Remarkably, ATAC-seq after prolonged Dox treatment did not reveal a substantial increase

in chromatin accessibility over the promoter region (Fig 5h). Small changes in the ATAC-seq

signal can be measured across samples and replicates, but these follow the sample-to-

sample variations globally (see Extended Data Fig. 10b for ATAC-seq coverage over the HoxA

genes cluster) and are therefore unlikely to be indicative of increased local accessibility.

The only increment in accessibility upon Dox treatment occurred downstream of the GFP

cassette, at a considerable distance from the TetO recruitment site and its adjacent PGK

reporter (Fig. 5h). That distal site includes another PGK promoter that is placed over 5 kbp

downstream to the TetO array. That distal PGK promoter originated from the construct that

was used to generate the reporter ([42] and references therein) and does not express a

functional protein-coding mRNA. Hence, we cannot exclude the possibility that CBX8 affects

accessibility at sites distant from the main recruitment hub, possibly through indirect

effects. Nevertheless, the data indicate that CBX8 recruitment and subsequent gene

repression are insufficient to restrict chromatin accessibility at the recruitment site.

Discussion

In conclusion, we have shown that PRC1-CBX8 binds chromatin via multivalent interactions

and induces chromatin condensation using both the nucleosome interacting surface of PRC1

and the IDRs of CBX8. PRC1^{C8} is dynamic within condensates while keeping chromatin in a

static, solid-like state (Fig 3). This is in contrast to the liquid-like state of chromatin

condensates that were observed in vitro, in the absence of PRC1[17]. We established that

PRC1^{C8} is sufficient to induce the solid condensed state of chromatin.

How can PRC1 condense chromatin but yet maintain a highly dynamic behaviour in the

nucleus? PRC1 is characterised with a short residence time on chromatin[31]. Solid-like

chromatin that is condensed by a mobile chromatin binder has been shown *in vitro* for a

truncation of the SAM-domain protein Polyhomeotic (Ph)[43] and in cells for HP1a at

chromocenters[20]. Yet, the mechanism allowing PRC1 and other gene-repressing factors to

condense chromatin while constantly diffusing in it remained largely unknown.

Our cryo-EM structure of PRC1-condensed chromatin explains how PRC1 can move in

condensed chromatin (Fig. 1), owing to the large pores that are formed between condensed

nucleosomes. The multivalent interactions between PRC1-CBX8 to chromatin provide PRC1

with multiple docking sites on chromatin: unmodified and modified H3 tail (Fig 3b-d), DNA

(Fig 3c) and the acidic patch of the nucleosome[36]. Hence, it is possible that PRC1-CBX8 can

constantly change its interactions with chromatin to maintain its condensed structure while

utilising its different chromatin-interacting surfaces to dynamically move around. While

doing so, the positively charged IDRs of CBX8 mask the negative charge of the DNA to bring

together chromatin segments and stabilize the condensed state of chromatin (Fig. 5i).

Neither the DNA binding activity of the chromo domain (Fig. 4b) nor H3K27me3 (Fig. 3d)

seem to be necessary for efficient PRC1^{C8}-chromatin phase separation. A potential limitation

in the usage of a methyl-lysine analogue (Fig 3d) is that it may not always serve as a perfect

histone mimic (discussed in [37]). However, since the chromodomain of CBX8 is dispensable

for chromatin condensation (Fig. 4), it is plausible that histone tail binding may be

dispensable too. Given that the chromodomain is required for DNA binding (Fig. 4d) and

implicated in binding to H3K27me3[37,38], it may play a role predominantly in recruitment.

In cells, canonical PRC1 includes an additional PHC protein, which was previously implicated

in chromatin compaction and condensation [8,28,43]. We reasoned that an in vitro study of a

simplified three-subunit complex, devoid of a PHC subunit, would allow us to characterize

the chromatin condensation activity of the CBX subunit. This minimal complex also allowed

us to overcome difficulties in purifying a PHC-bound PRC1 complex in sufficient quantities

and purity for structural studies. It is plausible that the chromatin compaction activities of

CBX8 and the PHC subunit cooperate in vivo and the absence of a PHC subunit in our

experiments presents a limitation when trying to extrapolate from our in vitro results.

While our experiments were not designed to extensively characterise the contribution of

protein-protein interactions towards phase separation or chromatin condensation, we

cannot exclude their involvement. Indeed, oligomerization of BMI1 was reported[44] and PHC

polymerization influences PRC1-chromatin condensate properties[28]. However, PRC1^{C8} does

not form condensates in the absence of chromatin in vitro (Fig 1e). Furthermore, there are

only about 10 PRC1 molecules per polycomb domain in cells[31], which is a factor that needs

to be considered when attempting to link protein oligomerization to chromatin compaction.

More studies are needed to determine how PRC1 molecules are distributed and work

together within polycomb domains in vivo and how protein-protein and protein-DNA

interactions contribute to this process.

Our data indicate that chromatin condensation together with dynamic behaviour within

chromatin condensates is an intrinsic biophysical property of PRC1-CBX8. It is plausible that

the dynamic behaviour of PRC1 within chromatin condensates is required in order to allow

PRC1 to modify nucleosomes by the H2AK119ub mark while holding them together. This

phenomenon might represent a broad paradigm of repressive chromatin. The internal

structure of PRC1-chromatin condensates is a porous network of nucleosomes (Fig. 1). Such

a structure could present a size-selective diffusion barrier, in agreement with its

permeability to PRC1 diffusion in vitro (Fig. 2) and Tn5 accessibility in cells (Fig. 5). The

existence of such a size-selective diffusion barrier remains to be identified in vivo, where it

may contribute to gene repression by selectively excluding transcriptional coactivators,

which are commonly large protein complexes- (>1MDa[12,45–47]). ATAC-seq experiments reach

their limitations in this context, because of the small size of the Tn5 used in these assays.

Hence, from our experiments in cells, we can only conclude that CBX8-bound chromatin is

not entirely inaccessible. Future studies may develop size-selective probes to directly

address questions of size-selective chromatin accessibility genome-wide. The hypothesis

that polycomb-mediated repression antagonises Pol II transcription without blocking all

proteins has been made nearly three decades ago[48]. This idea was conceived based on the

observation that T7 polymerase (~100 kDa) can initiate transcription from a polycomb-

repressed locus but GAL4-dependent transcriptional activation does not take place there.
This is in agreement with the inverse correlation between the density of chromatin domains
and the molecular weight of the chromatin modifiers present in them[12]. Transcription factor
size has also been suggested to determine access to different chromatin domains based on
simulations[49]. Combining our results with earlier findings[12,19,20,49], we propose that size-
selective exclusion may be part of a broader mechanism by which chromatin-interacting
proteins regulate the accessibility of repressive chromatin.

Acknowledgements: We would like to thank the Ramaciotti centre for electron microscopy
at Monash university for providing instrumentation, technical support and collecting cryo-ET
data, the Monash Proteomics facility for providing instrumentation and technical support
and the Monash Micro Imaging facility for providing instrumentation and technical support.
We would also like to acknowledge the MASSIVE HPC platform for providing high-
performance computing resources
388
Funding: This work was supported by an Australian research council (ARC) DECRA fellowship
DE210101669 (M.U.), NIH-NIMH R01MH122565 (O.B), start-up funding from the Norris
Comprehensive Cancer Center at Keck School of Medicine of USC (O.B), NIH-T32 training
grant T32HD060549 (J.M.), Sylvia and Charles Viertel Senior Medical Research Fellowship
(C.D.) and the National Health and Medical Research Council (NHMRC) grant numbers
APP1162921, APP1184637 and APP2011767 (C.D.). C.D. is an EMBL-Australia Group Leader.
395
Authors contributions: C.D. and M.U. conceptualised the project and acquired funding,
M.U., V.L., C.T., X.H.N, J.M., J.H., S.M., H.V., S.T., M.B. and M.L. carried out experiments and
investigated, P.P.D. and V.P. generated cell lines, S.F. and Q.Z. cloned and purified histone
mutants, M.L. and M.U. developed software, C.D., A.d.M., O.B., P.P.D., Y.G. and E.G.
supervised, M.U. and C.D. wrote the original draft and all authors reviewed and edited the
manuscript.
402
403 **Declaration of interests:** The authors declare no conflict of interest.
404
Data and materials availability: Next generation sequencing data (ATAC-seq and ChIP-seq)
are available under GEO accession number GSE220140. The maps for tomograms in Fig. 1 g,
and j have been deposited to the EMD (EMD-29022 and EMD-43554, respectively). All cryo-
ET raw data has been deposited to EMPIAR under accession numbers EMPIAR-11344 and
EMPIAR-11883, of chromatin in the presence and absence of PRC1-CBX8, respectively. XL-
MS data has been deposited to Pride (PXD039589 and PXD049094).

411
412

Fig. 1. The molecular structure of PRC1-chromatin condensates is porous and accessible to macromolecules. a, Schematics introducing the workflow. **b,** SDS-PAGE of purified PRC1^{C8} complex that includes RING1b, BMI1 and MBP-tagged CBX8. **c,** Size exclusion chromatography of the purified PRC1^{C8} using a HiLoad Sephacryl 300 16/60 column. **d,** In vitro ubiquitylation assay comparing PRC1^{C8} to a RING1b-BMI1 heterodimer. Samples in lane 2 and 3 included E1, E2, ubiquitin and ATP. Ubiquitylation is detected by western blot using an anti-H2A antibody. **e,** Chromatin condensates induced by the PRC1^{C8} complex and the individual proteins, visualised by confocal (left and centre) and phase contrast (right, from independent experiments) microscopy. CBX8 is GFP-labelled and chromatin is Cy5 labelled. Protein and chromatin concentrations are 1 μM and 20 nM (estimated nucleosome concentration), respectively. **f,** Cryo-confocal microscopy of vitrified PRC1^{C8}-chromatin condensates. **g,** Cryo-electron tomography of a PRC1^{C8}-chromatin condensate. Shown is a central slice through the reconstruction (left image). Nucleosome subtomogram averages (centre, bottom) are then placed in a volume the size of the tomographic slice, at the position and orientation determined by template matching and subtomogram averaging (right image). 1330 nM PRC1^{C8} and 3500 nM chromatin (estimated nucleosome concentration) were assayed in 3.5 mM HEPES-KOH pH 7.5, 6.8 mM TRIS-HCl PH 7.5, 21 mM NaCl, 7 mM KCl, 0.8 mM DTT. See Supplementary Data 1 for a list of cross correlation peaks. **h,** Surface representation of the volume of a subset of the PRC1^{C8}-chromatin condensate structure that is inaccessible to probes of given radii. **i,** Inaccessible volumes for a given probe radii are plotted, with exemplary molecules indicated (in grey) and selected probes coloured as in **h**. For the indicated complexes, the hydrodynamic radius was estimated[32] using resolved domains from published structures (see Methods section for PDB accessions) as a minimum size estimate. **j,** As in g but without PRC1^{C8}. See Supplementary Data 2 for a list of cross correlation peaks. **k,** Pairwise distances of each individual nucleosome to its nearest three neighbouring nucleosomes in 3D space in tomograms with and without PRC1^{C8}. Only unique pairs are plotted from two tomograms with (+PRC1) and three tomograms without (-PRC1). Whiskers extend from the 5th to the 95th percentiles. Significance was tested using a Brown-Forsythe ANOVA with a Games-Howell post hoc test. **** = p-value < 0.0001.

Fig. 2. PRC1^{C8} is mobile while chromatin is static within PRC1-chromatin condensates.

a, Titration of Chromatin against PRC1^{C8}. Condensates were assessed with a fluorescence widefield microscope imaging a Cy5 label on the chromatin. Presented are representative micrographs of three replicates, including two with MBP-tagged PRC1^{C8} (presented) and one with GFP-tagged PRC1^{C8}. **b,** Representative micrographs of FRAP recorded in PRC1^{C8}-chromatin condensates. CBX8 is GFP labelled and chromatin is Cy5 labelled. Mean fluorescence intensity of the bleached area, normalised to pre-bleach mean signal, is plotted for every time point. Error bars show standard deviation from n=7 (GFP) and n=8 (Cy5) measurements recorded from two independent experiments. The GFP signal recovery was fit with an exponential association model, best fit values for Plateau and fluorescence recovery half time (T$_{1/2}$) are shown with standard error. **c,** Schematic representation of the FRAP experiment.

Fig. 3. Multivalent interactions between PRC1^{C8} and chromatin. a, Chromatin condensation
in response to the whole PRC1^{C8} complex (RING1b, BMI1 and CBX8) or the individual
components CBX8 and the RING1b–BMI1 heterodimer. Representative images from two
replicates. **b,** Intramolecular (purple lines) and intermolecular (green lines) protein-protein
interactions mapped within PRC1-chromatin condensates using crosslinking mass
spectrometry (XL-MS). Data is from three independent replicates. **c,** PRC1^{C8} or PRC1 core
binding to a fluorescein labelled 24bp DNA probe measured by fluorescence polarisation.
Data points show the mean (baseline subtracted) of three independent replicates and the
error bars indicate the standard error. The continuous line represents the fit to a Hill binding
model, when applicable. **d,** Titration of PRC1^{C8} to unmodified chromatin (top) and
H3K27me3-MLA chromatin (bottom) at an identical chromatin concentration (50 ng/µl DNA)
and 150 mM KCl. Micrographs are representative of two independent replicates.

Fig. 4. Positive charges in the CBX8 IDRs are required for DNA binding and phase separation. a, Schematics depicting the different CBX8 mutants used, drawn to scale. **b,** PRC1^{C8}-chromatin condensates in the context of different CBX8 mutants. Varying concentrations of PRC1^{C8} were titrated to a constant concentration of C5-labelled reconstituted chromatin (50 ng/μl). Widefield fluorescence and differential interference contrast (DIC) micrographs are representative of three replicates. **c** Quantification of the total area covered by condensates per micrograph for different PRC1^{C8} mutants and concentrations. Bars represent the means from three independent replicates and error bars represent the standard deviation. **d,** Fluorescence polarisation assay measured the affinity of different PRC1^{C8} mutants for a Fluorescein-labelled 24 bp DNA probe. Data points are the mean (baseline subtracted) of three independent replicates and error bars indicate the

484   standard error. The continuous line represents the fit to a Hill binding model, when

485   applicable. **e,** Model for chromatin condensation by PRC1[C8]: Electrostatic interaction

486   between the CBX8 IDR with DNA provides charge screening and promotes phase separation.

Fig. 5. CBX8 binding sites on chromatin in mouse embryonic stem cells are accessible.
a, Schematics of the experimental setup (left) and anti-CBX8 western blot (right) of wildtype
and *Cbx8* knockout mESC after 48 hours of retinoic acid (RA) treatment. **b,** Overlap of ATAC-
seq peaks and CBX8 and H3K27me3 ChIP-seq peaks. ATAC-seq peaks are defined from two
biological replicates. **c,** ChIP-seq traces for H3K27me3 and CBX8 in wildtype mESC and
representative ATAC-seq traces at four genes in wildtype and CBX8 knockout mESC after 48
hours of RA treatment. **d,** Accessibility changes at all ATAC-seq peaks in wildtype (WT)
mESC, in response to retinoic acid (RA) treatment. **e,** Accessibility changes at all CBX8-target
sites in wildtype mESC, in response to RA treatment. **f,** Comparison of accessibility at CBX8-
target sites between wildtype and *Cbx8* knockout cells after RA treatment. **g,** Top: schematic
representation of the chromosome-integrated reporter. Left panel: ChIP-qPCR using FLAG
antibody (CBX8 is FLAG tagged) at indicated distances from the TetO array, in the presence
and absence of doxycycline (Dox) treatment for six hours. Bars represent the mean bound
over input (Bd/In) normalised to the IAP gene and points represent two replicates. See
Extended Data Fig 10a for ChIP-qPCR using additional antibodies. Right panel: Brightfield
and GFP-fluorescence images of the mECS cells before and after Dox treatment. **h,** ATAC-seq
signal reporting accessibility of the integrated locus before and after Dox treatment for six
days. From left to right: annotated are the TetO array and its proximal PGK promoter that
controls the Puromycin-GFP reporter gene, and the distal PGK promoter. **i,** Model: PRC1
forms multivalent interactions with chromatin, thereby stabilizing chromatin condensates
potentially through charge screening of negatively charged DNA by positive charges in the
CBX8 IDR. These interactions dynamically change as PRC1 diffuses through condensates.

Extended Data Fig. 1. Quality control of chromatin and protein constructs. a, MNase

digestion of reconstituted Chromatin and a naked DNA (same sequence as used for

chromatin reconstitution). DNA fragments post digestion are resolved on a 1.2 % Agarose

gel. Protected bands indicating mono- and di-nucleosome core particles (NCP and diNCP)

are indicated by the arrows. **b,** 3 ug of each protein complex used in this study resolved on a

4-12% SDS-PAGE gel stained with Coomassie. **c,** Ubiquitylation activity of each protein

complex used in this study visualized on a western blot. All samples include UBA1, UBCH5C,

Ubiquitin, ATP and 1 uM chromatin (nucleosome concentration). **d,** Phase separation

experiment comparing chromatin condensation activity of PRC1^{C8} with MBP-tagged CBX8 to

PRC1^{C8} with the tag cleaved.

524
525 **Extended Data Fig. 2. PRC1 complexes used in this study are monodispersed.** HPLC elution
526 profiles from a Shim-pack Bio Diol 200 HPLC column for each of the purified protein
527 complexes, as indicated.
528

Extended Data Fig. 3. Nucleosomes in PRC1[C8]-chromatin condensates show no orientation bias towards neighbouring nucleosomes. a, Low-magnification micrograph showing examples of dense regions suspected to be condensates (dashed circles). Tomograms were collected at the borders of regions such as these. Micrographs are from the same gird from which tomograms were collected. **b,** Structures of the template used for template matching (left), the averaged structure after template matching (middle) and the final structure after the subtomogram averaging routine (right) with the related Fourier shell correlation curve (bottom). **c,** Orientations of nucleosomes towards neighbouring nucleosomes within a cut-off of 20 nm. Individual points represent nucleosomes and lines between points are coloured according to the relative orientation of neighbouring nucleosomes as indicated in the colour key (left). **d,** Distribution of nucleosome-nucleosome orientation for the three nearest neighbours and the 150th neighbour of each nucleosome in tomogram #1. Colours correspond to the same respective orientations as in **c.**

543
Extended Data Fig. 4. a, Representative low-magnification cryo-EM micrograph of a grid
square with vitrified chromatin in absence of PRC1. **b,** Same as (**a**) but with PRC1. Black
arrows indicate presumed condensates. Red crosses relate to stage movement and do not
indicate features in the context of this figure. **c,** The same cryo-tomogram as in Fig. 1j, of
chromatin in the absence of PRC1, with dense regions highlighted. **d,** Distances to the three
nearest neighbouring nucleosomes, where +PRC1 as in Fig. 1k and -PRC1 shows distances
for the highlighted dense region in **c**.
551

553 **Extended Data Fig. 5.** Chromatin condensation in response to variations in salt and PRC1

554 concentration measured by confocal microscopy using Cy5-labelled chromatin at a constant

555 concentration of 50 ng/µl DNA (approximately 400 nM nucleosomes).

Extended Data Fig. 6. Untagged PRC1^{C8} is mobile while chromatin is static within PRC1-chromatin condensates. a, Representative micrographs of FRAP recorded in PRC1^{C8}-chromatin condensates, where within this complex CBX8 is untagged. PRC1^{C8} is labelled with ATTO488 and chromatin is Cy5 labelled. The mean fluorescence intensity of the bleached area, normalised to the pre-bleach mean signal, is plotted for every time point. Error bars show standard deviation from n=6 (GFP) and n=7 (Cy5) measurements that were recorded from four independent experiments that were carried out on different days. The GFP signal recovery was fit with an exponential association model, best fit values for Plateau and fluorescence recovery half time (T$_{1/2}$) are shown. The lower limits of the 95% confidence interval are presented in parentheses (the upper boundaries could not be determined confidently). **b,** Representative micrographs of FRAP recorded in PRC1^{C8}-chromatin condensates, where within this complex CBX8 includes an N-terminal MBP-tag. PRC1^{C8} is labelled with ATTO488 and chromatin is Cy5 labelled. The mean fluorescence intensity of the bleached area, normalised to the pre-bleach mean signal, is plotted for every time point. Error bars show standard deviation from n=7 (GFP) and n=6 (Cy5) measurements recorded from four independent experiments that were carried out on different days. The GFP signal recovery was fit with an exponential association model, best fit values for Plateau and fluorescence recovery half time (T$_{1/2}$) are shown with SEM.

Extended Data Fig. 7. Condensates can be detected at physiologically relevant PRC1^{C8}

concentrations. a, An illustration of single molecule confocal microscopy. Individual

condensates (in grey) are detected when diffusing through the confocal volume (in blue)

and emitting GFP fluorescence (in green). **b,** Top: representative trace tracking GFP signal

over time for samples with 500 nM GFP-labelled PRC1^{C8} and chromatin (200 nM nucleosome

concentration). Traces show a 5 minute window from a 20 minute experiment. Blue lines

show the raw GFP signal and orange lines show the GFP signal after low-pass filtering using

a Butterworth filter. Green dots indicate the maxima of the detected peaks. Bottom: same

as the top plot, but in the absence of chromatin. **c,** GFP peak counts at different PRC1^{C8}

concentrations. Data from two replicates are shown. Bars indicate the mean of two

independent replicates that were carried out on different days, the error bars show the

standard error of the mean and individual data points are presented.

Extended Data Fig. 8. Crosslinking mass spectrometry (XL-MS) profile of PRC1^{C8} is similar in
the absence of chromatin. Same experiment as in Fig. 3b, but without chromatin.

Extended Data Fig. 9. The nucleosome acidic patch is required for efficient PRC1-

chromatin phase separation. a Phase separation of *X. laevis* wildtype and acidic patch

mutant chromatin in response to increasing concentrations of PRC1[C8] wildtype. DIC

micrographs are representative of two replicates. **b** Same as (**a),** but with the PRC1[C8KR21A]

mutant. **c** Same as (**a**) but with the PRC1[C8ΔChromo] mutant.

Extended Data Fig. 10. a, ChIP-qPCR in the presence and absence of doxycycline (Dox)

treatment using RING1B antibody at indicated distances from a chromosome-integrated

TetO array (left), at control genes (right) and using FLAG (CBX8) antibodies at control genes

(middle). Bars represented the bound over input (Bd/In) normalised to the IAP gene and the

dots represent individual data points. **b,** ATAC-seq signal at the HoxA locus of the reporter-

integrated mECS cell line before and after dox treatment. Two independent replicates are

shown.

Extended Data Movie S1. Related to Fig.1; Cryo-tomogram of a PRC1^{C8}-chromatin condensate. The movie shows a scan through the z-axis of denoised[50] tomogram of chromatin in presence of PRC1.

Extended Data Movie S2. Related to Fig.1; Cryo-tomogram of chromatin without PRC1. The movie shows a scan through the z-axis of denoised[50] tomogram of chromatin in absence of PRC1.

Supplemental Table 1. Template matching results for PRC1^{C8}-chromatin condensates. Results from the Dynamo template matching process before manually removing false positive hits, in Excel format. Every entry describes a cross correlation peak above the cut-off of 0.17. Each peak indicates the position of a nucleosome at x-y-z coordinates given in columns 24-26 and at rotations applied to the template as defined by Euler angles in columns 7-9. The table is also provided in Dynamo format (.tbl) for direct import into Dynamo (Supplementary Data 1).

Supplemental Table 2. Template matching results for chromatin without PRC1^{C8}. Results from the dynamo template matching process before manually removing false positive hits in Excel format. Every entry describes a cross correlation peak above the cut-off of 0.23. Each peak indicates the position of a nucleosome at x-y-z coordinates given in columns 24-26 and at rotations applied to the template as defined by Euler angles in columns 7-9. The table is also provided in Dynamo format (.tbl) for direct import into Dynamo.

Supplementary Data 1. Template matching results for chromatin without PRC1^{C8}. Same as Supplemental Table 1, provided in Dynamo format (.tbl) for direct import into Dynamo.

Supplementary Data 2. Template matching results for PRC1^{C8}-chromatin condensates. Same as Supplemental Table 3, provided in Dynamo format (.tbl) for direct import into Dynamo.

Supplementary Data 3. XL-MS-identified crosslinks within PRC1^{C8}. Provided is a list of the crosslinks that were detected using XL-MS within PRC1^{C8} in the absence of chromatin.

641 **Supplementary Data 4. XL-MS-identified crosslinks within PRC1^{C8}-chromatin.** Provided is a

642 list of the crosslinks that were detected using XL-MS within PRC1^{C8} in the presence of

643 chromatin.

Materials and Methods

Plasmids and cloning

Human RING1b (Uniprot ID Q99496) and human BMI1 (Uniprot ID P35226) were cloned into a pFBOH-mhl vector (Addgene plasmid # 62304) cleaved with BseRI using Gibson Assembly® Master Mix (NEB #E2611L) using the primers indicated in Table S1.

Human CBX8 wildtype open reading frame (Uniprot ID Q9HC52-1 and NCBI Reference Sequence was NM_020649.2) was obtained from gene synthesis (Gen9). The CBX8[KR21A] mutant open reading frame was codon optimised for expression in *Trichoplusia ni* insect cells and synthesised (Genscript). The CBX8[ΔChromo] truncation was amplified from the wildtype ORF using primers indicated in Table 1 and then subcloned into a vector with the same backbone as used to expresses the wild type protein. All three CBX8 constructs were cloned into a modified pFastBac1 pFB1.HMBP.A3.PrS.ybbR vector digested by XhoI and XmaI sites to include a N-terminal 6xHis-MBP tag Cloning of the polycomb target gene ATOH1 into the pUC18 vector was described previously[51]. Plasmids for expression of human histones (H2A, H2B, H3.1 and H4) in *E.coli* were a kind gift from David Tremethick, Australian National University. UbcH5c WT pET28a was a gift from Rachel Klevit (Addgene plasmid # 12643; http://n2t.net/addgene:12643; RRID:Addgene_12643)[52]. pET3a-hUBA1 was a gift from Titia Sixma (Addgene plasmid # 63571; http://n2t.net/addgene:63571; RRID:Addgene_63571)[53].

To generate a baculovirus expression vector of a monomeric EGFP-CBX8 (mEGFP-CBX8) construct, first GFP was amplified from a pSpCas9(BB)-2A-GFP vector and CBX8 was amplified from a pFB1.HMBP.A3.PrS.ybbR vector containing CBX8 as an insert. Subsequently, EGFP-CBX8 was cloned into the pFBOH-mhl vector cleaved with BseRI via Gibson assembly, with a Serine-Glycine-Serine linker between EGFP and CBX8. Finally, to generate monomeric mEGFP-CBX8, alanine residue 207 in EGFP was mutated to Lysine using a site directed mutagenesis kit (Takeda) and the mutagenesis primers listed in Table 1.

Table 1: Cloning and mutagenesis primers (5'-3') and sequences

mEGFP mutagenesis A207K fw	CTGAGCACCCAGTCCAAGCTGAGCAAAGACCC	Used for mutagenesis of CBX8-EGFP
mEGFP mutagenesis A207K rv	GGGGTCTTTGCTCAGCTTGGACTGGGTGCTCA	Used for mutagenesis of CBX8-EGFP
EGFP_fw_PFBOH_MHL	TTGTATTTCCAGGGCATGGTGAGCAAGGGCGAG	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector
EGFP rv with SGS linker	TGAAAGCTCACTTCCACTCTTGTACAGCTCGTCCATGC	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector

CBX8 5' with SGS linker	TGTACAAGAGTGGAAGTGAGCTTTCAGCGGTGGGG	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector
CBX8 rv pFBOH_MHL	CAAGCTTCGTCATCATCATCTTTTCTCTTTAAAAAAGCCT	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector
ATOH1 fw	GCAGAGCCCAAACATTCACACA	Used to amplify ATOH1 from pUC18 plasmid
ATOH1 rv	GCGGAGTTTCCTAAAAGACGCC	Used to amplify ATOH1 from pUC18 plasmid
H2BT120C fw	GTGACCTGCTATACCAGCAGCAAATAA	Used for mutagenesis of H2B
H2BT120C rv	GGTATAGCAGGTCACGGCTTTGGTGCC	Used for mutagenesis of H2B
RING1b pFBOH-mhl fw	ttgtatttccagggcTCTCAGGCTGTGCAGACAAAC	Used for Gibson assembly into pFBOH-mhl vector
RING1b pFBOH-mhl rv	caagcttcgtcatcaTTTGTGCTCCTTTGTAGGTGC	Used for Gibson assembly into pFBOH-mhl vector
BMI1 pFBOH-mhl fw	ttgtatttccagggcCATCGAACAACGAGAATCAAG	Used for Gibson assembly into pFBOH-mhl vector
BMI1 pFBOH-mhl rv	caagcttcgtcatcaACCAGAAGAAGTTGCTGATGA	Used for Gibson assembly into pFBOH-mhl vector
24bp DNA duplex Fluorescein labeled	GGCGCCCTGCCCCGCCTCGCTCTG	Fluorescein labeled DNA duplex used for binding assays. Only the top strand is shown. The dye was attached to the 3' end of the top strand.
CBX8 KR21A	ATGGAATTATCAGCAGTAGGTGAACGCGTGTTTGCAGCGGAAGCGCTGCTCAAACGTAGGATTCGGAAGGGTCGCATGGAGTACCTAGTTAAGTGGAAAGGATGGTCACAGAAGTATAGCACATGGGAACCGGAGGAGAACATACTGGATGCTCGCTTGCTAGCAGCGTTTGAGGAGCGGGAACGAGAAATGGAGCTGTACGGACCTAAAAAACGTGGACCCAAGCCAAAGACCTTCCTTTTGAAGGCTCAAGCGGCTGCCAAGGCTGCCACGTATGAATTCAGATCGGATTCTGCCGCAGGTATAAGAATTCCCT	CBX8 KR21A mutant open reading frame (obtained by gene synthesis from GeneScript).

	ACCCAGGGGCCAGTCCACAGGA CTTGGCCTCCACGTCTAGGGCA GCAGAGGGCTTACGGAACATGG GCCTCTCTCCTCCGGCTTCATCTA CTAGTACGAGTAGTACATGTGCC GCGGAAGCGCCAAGGGACGCA GACCGAGACGCAGATCGGGATG CCGAGAGAGATGCGGAACGAGA GGCGGAAAGAGAAGCCGAGCGC GAGGCTGAGCGTGAAGCGGAAC GTGGCACAAGCGCCGTTGATGAC AAACCATCGAGCCCTGGTGATTC CAGCAAAAAACGGGGACCCAAG CCTAGGAAGGAGTTGCCGGACCC ATCCCAAGCCCCGCTTGGTGAAC CATCGGCGGGCCTCGGGGAATAT CTTAAAGGCCGAGCTTTGGACGA TACCCCTAGTGGTGCAGGAAAA TTTCCTGCGGGACATTCGGTTATC CAACTTGCTGCACGACAAGACTC AGATTTAGTACAGTGCGGGGTGA CATCCCCCAGCTCTGCAGAG GCGACCGGGGCCCTAGCTGTCGA CACCTTCCCAGCACGCGTGATAG CGCACCGAGCCGCATTTTTAGAA GCTGCTGGGCAGGGCGCGCTA GATCCCAACGGCACTAGGGTGCG GCACGGTTCAGGACCTCCCTCGT CCGGGGGGGGGCCTATATAGAGA CATGGGAGCTCAGGGGGGGTAGA CCGTCGCTTATCGCTCGTATCCCG GTAGCCCGTATTCTGGGTGACCC GGAGGAAGAATCCTGGTCTCCCA GTTTAACGAATTTAGAGAAA GTCGTAGTCACTGACGTTACGTC AAATTTTCTGACAGTCACCATCAA AGAGAGCAATACTGATCAAGGAT TCTTCAAGGAAAAGCGCTAA	
C8_PFB1_68_fw	TTCTGTTCCAGGGGCCCGGCCCC AAAAAGCGTGGA	Primer used to generate the CBX8$^{\Delta Chromo}$ truncation.
C8_PFB1_68_rv	TCGAGACTGCAGGCTTCATCTTTT CTCTTTAAAAAAGCCT	Primer used to generate the CBX8$^{\Delta Chromo}$ truncation.

Protein expression and purification

PRC1^{C8}, PRC1 core and CBX8 were co-expressed in *Trichoplusia ni* insect cells using the Baculovirus system. CBX8 variably carried an N-terminal 6xHis-mEGFP or a N-terminal 6xHis-MBP tag. The purification protocols were identical regardless of the tag. Baculoviruses were generated as per manufacturers instructions (Thermo Fisher). The viral titre was determined using the MTT assay (Promega #G3580). *Trichoplusia ni* insect cells were infected at a density of 1.5 - 2 x 10^6 cells and incubated for 60 hours at 27 °C in a shaker. The cells were spun down at 1500 relative centrifugal force (RCF). The pellet was resuspended in 100 ml of lysis buffer per litre of cell culture (20 mM HEPES-KOH pH 7.5, 400 mM NaCl, 25 mM Imidazol, 10 % Glycerol, 0.2 mM TCEP, 1 mM PMSF and EDTA-free Complete protease inhibitor (Thermo Fisher)). Lysis proceeded for 30-45 minutes at 4 °C while rotating. The lysate was then centrifuged at 29000 RCF for 20 minutes at 4 °C. The supernatant was transferred to a fresh tube and 5 ml of Ni-NTA resin (Qiagen) was added. The samples were then incubated for 60 minutes at 4 °C while rotating and subsequently centrifuged at 500 RCF for 5 minutes at 4 °C to settle the beads. About 90 % of the supernatant was removed. The beads were resuspended in the remaining 10 % of supernatant and transferred to 25 mm diameter gravity flow columns (Biorad). The beads were allowed to settle before the remaining buffer was drained and the beads were then washed with 12 CV of Buffer B (20 mM HEPES-KOH pH 7.5 at 20 °C, 500 mM NaCl, 25 mM Imidazole, 10 % Glycerol, 0.2mM TCEP), followed by 30 CV of Buffer A (20 mM HEPES pH 7.5 at 20 °C, 100 mM NaCl, 25 mM Imidazole, 10 % Glycerol, 0.5 mM DTT or 0.2 mM TCEP). The protein was then eluted in 6 CV Elution Buffer (20 mM HEPES-KOH pH 7.5 at 20 °C, 100 mM NaCl, 400 mM Imidazole, 10 % Glycerol, 0.2 mM TCEP). The eluted protein was loaded onto a Hitrap 5 ml Heparin column (Cytvia) equilibrated in IX Buffer A (20 mM HEPES pH 7.5 at 20 °C, 100 mM NaCl, 0.5 mM DDT or 0.2 mM TCEP) and the column was washed with 5 CV of IX Buffer A. The proteins were resolved over a 20 CV gradient ranging from 0 % to 100 % IX Buffer B (20 mM HEPES pH7.5 at 20 °C, 1000 mM NaCl, 0.5 mM DTT or 0.2 mM TCEP). The fractions were analysed on SDS-PAGE and fractions containing the protein complex of interest with the expected subunits stoichiometry were pooled. The pooled fractions were concentrated using a Amicon ultra 30K centrifugal filter (Merck, cat UFC903024) and purified by size exclusion chromatography using a HiLoad Sephacryl 300 16/60 column (Cytiva) equilibrated in GF Buffer (20 mM HEPES-KOH pH 7.5 at 20 °C, 150 mM NaCl, 0.5 mM DTT). The collected fractions were analysed on SDS-PAGE and fractions containing the protein complex of interest at the expected stoichiometry were pooled and concentrated to a concentration of 1-2 mg/ml using an Amicon ultra 30K centrifugal filter (Merck, cat UFC903024). The purified protein was then aliquoted and frozen in liquid nitrogen. The purified proteins were stored at -80 °C until use.

For the production of Atto-488 labelled PRC1^{C8} complexes, purification was carried out as above, with some modifications. PRC1^{C8} was purified as described above, without cleaving

the N-terminal MBP tag from CBX8, up until the end of the Ion Exchange Chromatography
(HiTrap 5 ml Heparin column, as above). Then, fractions containing PRC1[C8] were pooled
together and concentrated to 2.4 mg/ml. To generate untagged PRC1[C8], PreScission
protease was added to 1:50 protease:PRC1[C8] mass ratio in a 3.5 ml reaction volume,
incubated overnight at 4-8 °C and then the MBP tag was removed using 0.8 ml amylose
beads through a batch removal, before proceeding to the subsequent labelling reaction. For
the fluorescence labelling of MBP-tagged PRC1[C8], the subsequent labelling reaction was
carried out without tag cleavage. For fluorescence labelling, of either MBP tagged or
untagged PRC1[C8], 2.4 mg/ml protein was supplemented with Atto-488 NHS ester (Merck
41698; the Atto-488 NHS ester was dissolved in DMSO to 4.3 mM before used) to a final
molar stoichiometry of 1.4:1.0 dye:protein and allowed to incubate 1 hour at room
temperature in the dark, and then the protein was loaded on a light-protected HiLoad
Sephacryl 300 16/60 column equilibrated with 25 mM HEPES-KOH pH 7.5, 0.5 mM DTT, and
either 150 mM NaCl or 500 mM NaCl for the MBP-tagged or untagged PRC1[C8], respectively.
The process was completed as described above, with the exception that the fluorescently
labelled proteins were protected from light until experimentation.
737
738 Human UBA1, UBCH5C and Ubiquitin were purified as described previously[54]. Human
739 histone proteins H2A, H2B, H2BT120C, H3 and H4 were purified as described previously[55],
740 except that the gel filtration step was omitted. Purified *Xenopus laevis* histone proteins H2A,
741 H2B, H3, H4 and H2A acidic patch mutant (E56T, E61T, E64T, D90S, E91T, E92T) were bought
742 form from the Histone Source of the Colorado State University, Fort Collins.
743
744
745 **Production and purification of DNA for chromatin reconstitution**
746
747 The ATOH1 polycomb target gene was amplified in a large scale 10 ml PCR reaction including
748 500 nM ATOH1 fwd and reverse primers (see Table 1), 4 μg of ATOH1-pUC18 template[51],
749 200 μM dNTP mix (Invitrogen), 3 % DMSO, 50 mM KCl, 10 mM Tris-HCl, pH 8.8, 1.5 mM
750 MgCl2, 0.01% sterile gelatin. The reaction mixture was divided into 96-well plates to include
751 50 μl per well and the following PCR program was run:
752
753

Temperature (°C)	Time (minutes)	Number of Cycles
95	3:00	1
95	0:15	
63	0:20	25

72	8:00	
72	3:00	1

754
755
The PCR product was purified via ion exchange chromatography using a 5 ml HiTrap Q
column (GE Healthcare). The column was equilibrated in Buffer A (25 mM HEPES 7.5, 250
mM NaCl) and the sample was resolved using a linear gradient ranging from 0 % to 100 %
Buffer B (25 mM HEPES pH 7.5, 2 M NaCl). The fractions containing pure DNA were pooled,
subjected to ethanol-precipitation and resuspended in 10 mM Tris-HCl pH 7.5 at 20 °C, 0.1
mM EDTA.
762
763 **Chromatin reconstitution**
764
Histone Octamers were refolded as described previously[55]. All steps were done at 4 °C or on
ice. Chromatin was reconstituted following the salt-gradient dialysis protocol described
previously[55]. For large scale reconstitution, DNA and octamer were combined at an optimal
ratio that was determined at trial experiments for each batch of octamers and DNA. To
determine the optimal ratio of histone octamer to DNA, titration was carried out using
increasing amounts of octamer to a constant amount of DNA (molar ratios of 16:1, 18:1,
20:1, 22:1 of Octamer:DNA) following by salt gradient dialysis. For the salt gradient dialysis,
samples were initially dialysed in a buffer containing 10 mM Tris-HCl pH 7.5 at 20 °C, 2 M
KCl, 1 mM EDTA, 1 mM DTT. The initial buffer was then gradually exchanged for a low salt
buffer containing 10 mM Tris-HCl pH 7.5 at 20 °C, 250 mM KCl, 1 mM EDTA, 1 mM DTT over
the course of 18 hours, after which the salt exchange was complete. Samples were then
centrifuged at 21000 RCF for one minute, separated on a 1% agarose gel in TAE buffer and
stained with SYBR Safe (Sigma). The highest Octamer:DNA ratio at which chromatin
remained soluble after finishing salt dialysis was finally used for large scale reconstitution.
For large scale reconstitutions, the salt gradient dialysis was performed as above while
scaling up the reaction accordingly. Additionally, after conclusion of the salt gradient, the
samples were transferred to 400 ml of low salt buffer (10 mM TrisRIS-HCl pH 7.5 at 20 °C,
250 mM KCl, 1 mM EDTA, 1 mM DTT) and dialysed for another 2 hours. The samples were
finally dialysed against 1 litre of Chromatin Storage Buffer (10 mM TRIS-HCl pH 7.5, 10 mM
KCl) overnight.
785
786 **Reconstitution of Fluorescently labelled chromatin**
787
To allow site-specific labelling of histone H2B, a cysteine was introduced via site-directed
mutagenesis (H2BK120C, as previously described[56]) using mutagenesis primers indicated in
Table S1. H2BK120C was labelled with Cyanine5-maleimide (Lumiprobe cat #13080) as
described previously[56]. Labelled octamers were refolded as described above. Before

chromatin reconstitution, the unlabelled and labelled octamers were combined at a molar
ratio of 7:1 (unlabeled:labelled). Chromatin was then reconstituted as described above.
794
795   **Chromatin condensation assays**
796
Fluorescently labelled proteins and chromatin were protected from light whenever possible.
Chromatin condensation assays were done as described previously[17] with some
modifications. Assays were done in 384-well plates with #1.5 glass bottoms (MatTek
PBK384G-1.5-C). The wells were treated with 1 M NaOH for 1 hour at room temperature,
NaOH was removed and wells were washed with copious amounts of MilliQ water (MQ).
MQ was removed and 70 ul of 5k mPEG-silane (Sigma #JKA3037-1G, dissolved in 95% EtOH
to a final concentration of 25 mg/ml) was added to each well. The plates were sealed and
incubated overnight at room temperature. The mPEG-silane was removed, the wells were
washed once with 95 % EtOH, then rinsed with copious amounts of MQ and dried
completely in the fume hood.
The wells were then passivated by adding 40 ul of 20 mg/ml BSA (NEB) and incubated for at
least one hour at room temperature. The BSA was removed and the wells were washed
three times with Condensation Buffer (20 mM HEPES pH 7.5 at 20 °C, 0.2 mg/ml BSA, 10 %
Glycerol, 5 mM DTT and 150 mM KCl). The chromatin stock was adjusted to a DNA
concentration of 100 ng/ul (unless otherwise stated) and 150 mM KCl. The PRC1 complex
was diluted in Condensation buffer to a protein concentration equal to twice the final PRC1
assay concentration as stated. To induce condensation, 16 ul of the diluted PRC1 were
combined with 16 ul of the salt-adjusted chromatin dilution in PCR tubes. The samples were
incubated for 30 minutes at room temperature before being transferred to the 384-well
plate and incubated for a further 60 minutes at room temperature, so that the first images
were recorded 90 minutes after induction of condensation.
Images were recorded with a Nikon C1 scanning confocal microscope. GFP was excited with
a 488 nm laser, Cy5 was excited with a 561 nm laser. Linear contrast adjustments were
made with ImageJ. Where several micrographs are compared to each other, the same
contrast settings were used for all micrographs.
For condensation assays in Fig. 2a, Fig. 4, Extended Data Fig. 1d and Extended Data Fig. 9,
chromatin and PRC1 dilutions were prepared in Chromatin Buffer (10 mM TRIS-HCl pH 7.5 at
20 °C, 10 mM KCl, 1 mM DTT) and PRC1 buffer (20 mM HEPES-KOH pH 7.5 at 20 °C, 150 mM
NaCl, 1 mM DTT), respectively. After adjusting BSA and salt concentration and combining
PRC1 and chromatin, the final reaction contained the PRC1 and chromatin concentration
stated in the figures and the following components: 20 mM HEPES-KOH, pH 7.5, 5 mM TRIS-
HCl pH 7.5 at 20 °C, 90 mM KCl, 32.5 mM NaCl, 0.2 mg/ml BSA (NEB), 1mM DTT.
DIC images and fluorescence wide field images were recorded with a Leica DMI8 imaging
system equipped with a 4.2 MP (2k x 2k) sCMOS monochromatic K8 camera and a 63x oil
immersion objective with numerical aperture of 1.3. DIC images in Extended Data Fig. 9
were recorded with a Leica DMI8 imaging system equipped with a 4.2 MP (2k x 2k) sCMOS

monochromatic DFC 9000GT camera and a 40x dry objective with a numerical aperture of
0.8.
Condensate quantification was done using ImageJ. A threshold was set for each image to
segment condensates in the frame using the Cy5 signal. Condensates overlapping the frame
edges were excluded. Then the total area of all condensates in the frame was calculated.
838
839   **Chromatin ubiquitylation assay**
840
The salt concentration of the chromatin stock was adjusted to 100 mM KCl. The nucleosome
equivalent concentration of chromatin arrays was calculated by measuring the molar DNA
concentration and assuming that one DNA molecule is populated by 20 nucleosomes. The
reaction mixture included 750 nM (Fig. 1d) or 1000 nM (Extended Data Fig.1d) of chromatin
(nucleosome equivalent concentration), 500 nM PRC1^{C8} or RING1b-BMI1 dimer, 100 nM
hUBA1, 500 nM UBCH5C and 50 μM ubiquitin in Ub-Buffer (25 mM HEPES-KOH pH 7.5 at
20 °C, 100 mM KCl, 3 mM MgCl and 2 mM DTT) and started by adding ATP to a final
concentration of 3 mM. 15 ul reactions were incubated at 30 °C for 45 minutes or the
indicated time points (Extended Data Fig 1d). The reaction was stopped by adding 5 ul of 4X
NuPage LDS-loading dye (Thermo Scientific cat #NP0008) supplemented with 5 % 2-
mercaptoethanol. The samples were separated on a 4-12% NuPage gel (Thermo Scientific
cat #NP0321BOX) using MES buffer (Thermo Scientific cat #NP0002) in the tank. The gels
were then blotted onto a nitrocellulose membrane (Amersham, cat #GE10600002) in Tris-
Glycine transfer buffer + 20 % Ethanol (v/v) for 90 minutes in the cold room at 310 mAmp in
a blotting tank (BioRad). H2A was detected using anti-H2A primary antibodies (Merck
Millipore Cat. # 07-146, 1:1000 titer) and HRP-conjugated secondary antibodies (Santa Cruz,
cat #sc-2357, titer 1:5000).
858
859   **Sample preparation for cryo-electron tomography**
860
The PRC1^{C8} complex (MBP-tagged CBX8) was combined with chromatinized ATOH1 at a final
concentration of 1.6 μM PRC1^{C8} and 500 ng/ul DNA at a final salt concentration of 25 mM
NaCl and 8.3 mM KCl. Samples were incubated at room temperature for 30 minutes. Just
before freezing, 5 nm gold nanoparticles were added at a 1:6 ratio (v/v). 3.5 ul of sample
was applied to a Quantifoil grid (R1.2/1.3 on 200 copper mesh, Quantifoil cat #N1-
C14nCu20-01) and vitrified in liquid ethane using the Vitrobot plunge freezer (Thermo
Scientific) with the following settings: Temperature 4°C, blot force -3, blot time 4 seconds,
humidity 100%. The final sample composition after addition of gold was 1330 nM PRC1^{C8}
and 3500 nM chromatin (estimated nucleosome concentration) in 3.5 mM HEPES-KOH pH
7.5, 6.8 mM TRIS-HCl PH 7.5, 21 mM NaCl, 7 mM KCl, 0.8 mM DTT. The sample of chromatin
in absence of PRC1 was prepared the same but instead of adding PRC1 an equivalent
volume of the PRC1 buffer was added.
873

Low magnification cryo-EM image collection

The images in Extended Data Fig. 3a were collected just before cryo-ET data collection using the Titan Krios (Thermo Fisher) at an acceleration voltage of 300 keV and are of the same grid from which tomograms were collected. The images in Extended Data Fig. 4a were recorded using a Talos Arctica TEM (Thermo Fisher) at an acceleration Voltage of 200 keV.

Cryo-electron tomography data collection and processing

The data was collected with a Titan Krios electron microscope (Thermo Fisher) at 300 keV acceleration voltage using a Gatan K2 (+PRC1 sample) or K3 (-PRC1 sample) Summit camera (Gatan). A tilt series was acquired ranging from -60 to 60 degrees with 3 degree increments and a nominal defocus of -2.5 μm. A dose symmetric collection scheme was followed as described previously[57]. Chromatin in the presence of PRC1 was images at pixel size of 1.32 Angstrom. The chromatin samples in the absence of PRC1 were collected in super resolution mode with a nominal pixel size of 1.632 Angstrom (0.815 Angstrom super resolution pixel size). The total dose per tomogram was 144.32 e/A^2 (+PRC1) and 160.72 e/A^2 (-PRC1). Four frames were collected per tilt (dose per frame 0.88 e/A^2 for the +PRC1 sample and 0.98 e/A^2 for the -PRC1 sample). The movies were motion corrected using motioncor2[58]. The motion corrected images were combined into stacks and further processed with Imod[59] version 4.9.9. Imaging artefacts were identified and removed with Imod's Ccderaser function. Tilts were aligned using the gold fiducial markers and the final aligned stack was binned by a factor of 4. The defocus was estimated using emClarity[60] and the estimated defocus was used in Imod for CTF correction. The final tomogram was calculated using Imod's implementation of weighted back projection. For visualisation, the tomogram was denoised using Topaz[50]. The denoised tomogram was only used for visualisation (Figure 1 and Movie S1). The original non-denoised tomograms were used for all further processing, including template matching and subtomogram averaging.

Template matching, subtomogram averaging and modelling the chromatin structure of condensates from cryo-electron tomograms

Two tomograms were selected for further processing (Tomogram #1 and Tomogram #2 in the following). To avoid user bias, we used a template matching algorithm[61] with the structure of a single nucleosome as template (EMD-8140[61]) to identify initial positions and orientations for each nucleosome. The pixel size of the template was adjusted using emClarity (version 1.0.0) to match the unbinned pixel size of the tomogram[60]. The template was then subjected to a low-pass filter of 30 Angstrom and binned by a factor of four to match the binned tomogram. Template matching was done with the Matlab implementation of Dynamo (version v-1.1.514)[61] using the "dynamo_match" function. The template was masked with a tight-fitting mask with smooth edges. The scanning range was

set to 360 degrees with 10 degree steps. In-plane rotation was also scanned over 360
degrees with 10 degrees steps. Particles that passed a cross-correlation cut-off of 0.17
(Tomogram #1), 0.19 (Tomogram #2), 0.24 (Tomogram #27), 0.23 (Tomogram #41) and 0.25
(Tomogram #49) were selected for further analysis. Obvious false positives were removed
manually.

The particles were cropped from the tomogram with a box size of 36 pixels. Nucleosome
position and orientation was refined over three rounds of subtomogram averaging[61]. As an
initial template for subtomogram averaging, we used an average from all cropped particles
after template matching, masked with a tight-fitting mask with smooth edges generated in
the Dynamo mask editor. Specific settings for the different rounds of subtomogram
averaging are detailed in Table 2. The resulting average structure of a nucleosome from the
subtomogram averaging was then used to populate a volume the size of the tomogram with
nucleosomes at the determined positions and orientations. The graphic depiction of the
final model (Figure 1G) was generated using the Dynamo Matlab implementation[61].
928
929   Table 2: Settings for subtomogram averaging in Dynamo

	Round 1	Round 2	Round 3
Iterations	3	3	3
References	1	1	1
Cone Aperture	100	16	8
Cone Sampling	5	2	1
Azymuth Rotation Range	100	16	8
Azymuth Rotation Sampling	5	2	1
Refine	0	0	4
Refine Factor	0	0	2
High Pass	2	2	2
Low Pass	7	8	8

Symmetry	c1	c1	c1
Particle Dimension	36	36	36
Shift Limits	5	4	2
Shift Limiting Ways	1	1	1
Seperation in Tomogram	0	0	0
Basic MRA	0	0	0
Threshold Parameter	0.5	0.6	0.6
Threshold Modus	2	2	2

930
931
932 **Analysis of exclusion volume**
933
Exclusion volumes for spherical molecules of various radii were calculated using 3V[26] as
follows. First, the positional and rotational coordinates of the nucleosomes within
Tomogram #1 were determined by template matching followed by subtomogram averaging,
as described above. Next, the positional and rotational coordinates of the nucleosomes
were tabulated from within a section of 300 pixels x 300 pixels x 150 pixels (corresponding
to 158.4 nm x 158.4 nm x 87.0 nm at the x, y and z axes, respectively) centred on the pixel at
location (650,650,80), which was large enough to include hundreds of nucleosomes but yet
sufficiently small to carry out the subsequent computational analysis. The dimensions of the
resulting table were converted to Angstrom by multiplying with a tomogram pixel size of
5.28 Angstrom/pixel. Next, the table including nucleosome positions was used by the
Dynamo function 'dtchimera' to generate a chimera cmd script that placed a pdb structure
of a single nucleosome (PDB 1KX4[63]) at the position and orientation defined for each
nucleosome in the table. At the end of this process, each of the nucleosomes within the
tomogram section is represented by the pdb coordinates of the nucleosome template. The
resulting model was saved as a pdb file and used as input for 3V[26]. Varying probe radii
ranging from 2 to 20 nm were used.
950
951
952
953

Hydrodynamic radius calculation using protein structures

The hydrodynamic radii of various proteins (Figure 1I) were calculated using the HullRad[39] web server (http://52.14.70.9/). The PDB codes for the structures used are: 6LTJ (BAF[45]), 1MUH (Tn5[40]), 7O4J (PolII-PIC[47]), 6C24 (PRC2.2[64]), 8GXS (PolII-Mediator[65]). 6LTJ, 6C24, 8GXS and 7O4J are only partial models as not all residues were assigned, therefore the calculated hydrodynamic radius represents an estimate of the minimal complex size, while the actual size of these complexes could be larger.

Analysis of nucleosome-nucleosome orientation and distances

Nucleosome-nucleosome orientation was classified into face-face, face-side and side-side, as described in[22]. The orientations were calculated from the output table of the template matching and subtomogram averaging process in Dynamo[61] using the Matlab script 'calculate_orientation.m'. The nucleosome-nucleosome orientation plot (Fig. S1A) was generated using the 'plot_edges' function from the python notebook NCP_orientation_analysis.ipyn. Both scripts are available on Github (https://github.com/MichaUckelmann/Chromatin-Structure-Analysis). Centre-centre distances between neighbouring nucleosomes were calculated using the Matlab function 'knnsearch'. Double-counted distances were removed before further analysis.

Cryo-light microscopy

The samples were prepared as described for cryo-electron tomography, without gold nanoparticles and in a buffer containing 100 mM KCl. The vitrified grids were imaged using a ZEISS LSM900 Airyscan2 with a Linkam CMS196V Cryo stage.

Fluorescence recovery after photobleaching (FRAP)

For FRAP of PRC1[C8] that was fluorescently labelled using a GFP-CBX8, samples and plates were prepared and images were recorded as described above for chromatin condensation assays. 1 µM PRC1[C8] with an N-terminal GFP tag was used. The FRAP experiments were set up with the NIS-Elements software (Nikon). Regions of interest (ROIs) were defined and a single image was recorded before bleaching. Then the ROI was bleached using 488 nm (GFP) and 561 (Cy5) lasers. The bleaching time and laser power was set so that approximately 80 % of fluorescence signal within the ROI was quenched. Recovery was measured over 424 seconds, recording a total of 13 images. The data was analysed with ImageJ, the mean pixel intensity of the bleached ROI was quantified for each timepoint.

For FRAP of PRC1[C8] that was fluorescently labelled using spared labelling of lysine residues using Atto-488, samples and plates were prepared and images were recorded as

described above for chromatin condensation assays. To bring both the untagged PRC1^{C8} and

the MBP-tagged PRC1^{C8} (i.e. an N-terminal MBP tag on CBX8) to the same concentration, the

stock solutions of the Atto-488-labelled untagged PRC1^{C8} was adjusted to 8 μM PRC1^{C8}, 150

mM NaCl and 25mM HEPES pH7.5. Both the PRC1^{C8} stocks, with or without the MBP tag,

were then diluted to 4 μM protein in a condensation buffer (25 mM HEPES pH7.5, 150 mM

KCl, 0.2 mg/mL BSA, 2 mM DTT). 8 μl of 4 μM PRC1 solution and 8 μl of 100 ng/μl chromatin

(DNA concentration) were combined in a well to induce condensation, at final

concentrations of 2 μM PRC1^{C8}, with or without the MBP tag, 50 ng/μl chromatin (DNA

concentration) and with a final buffer composition of 25 mM HEPES pH7.5, 37.5 mM NaCl,

112.5 mM KCl, 0.1 mg/mL BSA and 1 mM DTT. Regions of interest (ROI) were defined and a

single image was recorded before bleaching. Then the ROI was bleached using 488 nm

(Atto-488) and 561 nm (Cy5) lasers. The bleaching time and laser power were set so that

approximately 80 % of the fluorescence signal within the ROI was quenched. Recovery was

measured over 24 minutes, with a frequency of ~ 1 image/minute. The data was analysed

with NIS-Elements software.

1010

1011 **Single molecule confocal microscopy**

1012

Cy5 labelled reconstituted chromatin (200 nM nucleosome concentration) and GFP labelled

PRC1^{C8} at the concentration indicated in the figure were combined in assay buffer (25 mM

HEPES-KOH pH 7.5, 100 mM KCl, 15 mM NaCl, 1 mM DTT, 0.1 mg/ml BSA). Samples were

immediately loaded into a custom-made silicone well plate with a 70 x 80 mm glass

coverslip (ProSciTech, Kirwan, QLD, Australia). Plates were analysed at room temperature on

a custom setup based on a Zeiss Axio Observer microscope (Zeiss, Oberkochen, Germany).

Illumination is provided by a 488 nm and a 561 nm laser beams, cofocussed in the sample

volume using a ×40 magnification, 1.2 Numerical Aperture water immersion objective (Zeiss,

Oberkochen, Germany). This creates a very small observation volume in solution (~1 fL),

through which fluorescent proteins diffuse and emitting light in specific wavelengths as their

fluorescent tags are excited by the laser beams. Light emitted by the fluorophores is split

into GFP and mCherry channels by a 560 nm dichroic mirror. The fluorescence of GFP is

measured through a 525/50 nm band-pass filter and the fluorescence of mCherry is

measured through a 580 nm long-pass filter. Fluorescence is detected by two photon

counting detectors (Micro Photon Devices, Bolzano, Italy). Photons of the two channels are

recorded simultaneously in 1 ms time bins by a custom Lab-VIEW 2018 program (National

Instruments)[66]. The data were analysed using a custom python script (Spyder version 4.1.5)

that automatically detects peaks, as described in[67,68].

1031

1032 **DNA binding assays**

1033

DNA binding was assayed using a 24 bp DNA (sequence see Table 1) with Fluorescein

attached to the top strand. The probe was protected from light wherever possible. The

probes were synthesised and delivered as duplexes (IDT). Probes were dissolved at a
concentration of 5 mM DNA in milliQ water. The probes were then diluted to 4 µM in
annealing buffer (20 mM HEPES-KOH, 150 mM NaCl) and heated to 95 °C for 5 minutes. The
probe was then left at room temperature for at least 1 hour to anneal before being diluted
to 20 nM in 20 mM HEPES-KOH pH 7.5 at 20 °C, 150 mM NaCl, 1 mg/ml BSA (NEB cat
#B9000S), 0.1% Tween20, 1 mM DTT. Serial protein dilutions of the PRC1[C8] and RING1b-
BMI1 complexes were prepared in Protein Dilution Buffer (20 mM HEPES-KOH, 150 mM
NaCl, 1 mM DTT) ranging from 8000 nM to 3.9 nM (Fig. 3) and from 7200 nM to 7 nM (Fig.
4). 20 ul of probe were mixed with 20 ul of the respective protein dilution and transferred to
a 384-well plate. The samples were incubated for 30 minutes in the dark at room
temperature and then read using a Pherastar plate reader (BMG Labtech). The fluorescence
anisotropy signal was normalised and the curves were fitted with a specific binding model
with Hill slope (GraphPad Prism).
1049
1050 **Crosslinking mass spectrometry (XL-MS)**
1051
Sample preparation, processing and data analysis was identical for samples with and
without chromatin. Reconstituted chromatin (if used) was dialysed overnight against 1 litre
of XL buffer (25 mM HEPES-KOH pH 7.5 at 20 °C, 150 mM NaCl, 1 mM DTT). The chromatin
was combined with PRC1[C8] in a 1:1.5 molar ratio (Nucleosomes:PRC1[C8]). Specifically, the
nucleosome equivalent concentration of the chromatin array was calculated from the
measured DNA concentration assuming 18 nucleosomes per DNA molecule. 1.5 µM PRC1
and 1 µM nucleosome equivalent concentration of chromatin arrays were then combined in
XL buffer and incubated for 30 minutes at room temperature. Crosslinking was done as
described before[69]. The bis(sulfosuccinimidyl)suberate (BS3) crosslinker was added to a final
concentration of 500 µM and crosslinking proceeded for 20 minutes at room temperature at
a reaction volume of 15 µl. The reaction was stopped by the addition of Tris-HCl pH 8 at
20 °C to a final concentration of 125 mM. The samples were then diluted to a volume of
100 µL using a buffer containing 50 mM Tris pH 8 at 20 °C and 150 mM NaCl. TCEP was
added to a final concentration of 10 mM and the samples were incubated at 60 °C for
30 min. Chloroacetamide was added to a final concentration of 40 mM and the samples
were incubated in the dark for 20 min. The samples were then digested using trypsin
(Promega cat #V5280) at 37 °C overnight. The digest was stopped by adding formic acid to a
final concentration of 1% v/v. The digested samples were purified using 100 µl ZipTip pipette
tips (Merck cat #ZTC18S960) according to the manufacturer's instructions. Samples were
then concentrated to ~5 µL using a SpeedVac vacuum centrifuge and diluted with 20 µL
Buffer A (0.1% v/v formic acid).
1073
The peptides were analyzed by online nano-high-pressure liquid chromatography (UHPLC)
electrospray ionization-tandem mass spectrometry (MS/MS) on an Q Exactive Plus
Instrument connected to an Ultimate 3000 UHPLC (Thermo-Fisher Scientific). Peptides

reconstituted in 0.1% formic acid were loaded onto a trap column (Acclaim C18 PepMap 100
nano Trap, 2 cm × 100 µm I.D., 5-µm particle size and 300-Å pore size; Thermo-Fisher
Scientific) at 15 µL/min for 3 min before switching the precolumn in line with the analytical
column (Acclaim C18 PepMap RSLC nanocolumn, 75 µm ID × 50 cm, 3-µm particle size, 100-
Å pore size; Thermo-Fisher Scientific). The separation of peptides was performed at
250 nL/min using a non-linear acetonitrile (ACN) gradient of buffer A (0.1% formic acid) and
buffer B (0.1% formic acid, 80% ACN), starting at 2.5% buffer B to 42.5% over 95 min. Data
were collected in positive mode using a Data Dependent Acquisition m/z of 375–2000 as the
scan range, and higher-energy collisional dissociation (HCD) for MS/MS of the 12 most
intense ions with z 2–5. Other instrument parameters were: MS1 scan at 70,000 resolution,
MS maximum injection time 118 ms, AGC target 3E6, ion intensity threshold of 4.2e4 and
dynamic exclusion set to 15 s. MS/MS resolution of 35000 at Orbitrap with the maximum
injection time of 118 ms, AGC of 5e5 and HCD with collision energy = 27%.
For the data analysis, Thermo raw files were analysed using the pLink 2.3.4 search engine[70]
to identify crosslinked peptides, searching against the sequences of RING1b, BMI1, CBX8,
H2A, H2B, H3 and H4. The default settings for searches were used. N-terminal acetylation
and methionine oxidation were used as variable modifications and carbamidomethyl on
cysteines as a fixed modification. False discovery rates of 1% for peptide spectrum match
level were applied by searching a reverse database. Crosslinked peptides were further
analysed using the crissrosslinkeR package[71]. Specifically, peptides were retained by
crissrosslinkeR only if they passed a p-value cutoff of 0.05 or were present in at least two of
three replicates. Subsequent visualisation of retained peptide was carried out with xiNET[72].
1099
1100 **Generation of Cbx8 KO mESC lines using CRISPR/Cas9**
1101
Paired sgRNAs were designed to delete exons 1–4 of the murine *Cbx8* gene. The Golden
Gate Cloning method was used to clone the sgRNAs (sequence below) into the lentiguide-
mCherry-Cas9 plasmid[73,74]. 2 million mESCs were transfected with 1 µg of each plasmid
carrying sgRNAs-mCherry-Cas9, using electroporation (Neon™ Transfection System
MPK5000). The following day, mCherry-positive mESCs were sorted by FACS and plated on a
10 cm dish at a very low density for single-cell clone picking. After 5-6 days, individual
colonies (derived from single cells) were picked, expanded, and genotyped using genomic
PCR to identify homozygous/biallelic deletions of Cbx8 KO mESC colonies. Selected Cbx8 KO
mESC lines were further confirmed by western blot for CBX8 (Cell Signalling, CBX8 (D2O8C),
cat # 14696S, titer 1:50 (Fig. 4a)) and HRP-conjugated secondary antibodies (Santa Cruz, cat
#sc-2357, titer 1:10000).
1113
1114 **Cbx8 sgRNA sequences (5' and 3' sgRNAs)**
1115 CBX8-5'Fw: CACCTGCGAATGCGCCGCTTCAGG
1116 CBX8-5'Rv: AAACCCTGAAGCGGCGCATTCGCA
1117 CBX8-3'Fw: CACCCTCTATGGCCCCAAAAAGCG

1118   CBX8-3'Rv: AAACCGCTTTTTGGGGCCATAGAG

1119

1120   **Cbx8 genotyping primers:**

1121   Deletion_Fw: GCCTTCTGGTGCAGCTAAGT

1122   Deletion_Rv: GACGTCAGCGGGAGAGTATT

1123   Internal_Fw: CACCAAATGAATGCTCCAAA

1124   Internal_Rv (same as the Deletion_Rv): GACGTCAGCGGGAGAGTATT

1125

1126   **Mouse embryonic stem cell culture**

1127

Wildtype and *Cbx8* knockout mES cells were grown on gelatinized culture dishes in DMEM,

20 % FBS, 1x non-essential amino acids (Gibco #11140-050), 1x Glutamax (Gibco #35050-

061), PenStrep 100 u/ml (Thermo Fischer), 0.5 x EmbryoMax 2-Mercaptoethanol (Merck

Millipore #ES-007-E), 2.5 µg/ml Plasmocin (Invivogen), 1000 U/ml ESGRO Leukemia

Inhibitory Factor (LIF) (Merck Millipore cat #ESG1107). For ATAC-seq experiments with the

reporter-integrated mECS line, the cells were treated for 6 days with 1 µg/ml Doxycycline

(passaged every 48 h) before ATAC-seq was performed as described below.

1135

1136   **Mouse embryonic stem cell differentiation ahead of ChIP-seq and ATAC-seq**

1137

Differentiation was induced by seeding cells at a density of 0.3×10^6 cells per well in 6-well

plates (for ATAC-seq) or at 1.5×10^6 cells per 10-cm culture dish (for ChIP-seq) in media

containing 1 µM all-trans retinoic acid (RA,Sigma-Aldrich R2625-50MG) and no LIF. Cells

were differentiated for 48 hours and the media was changed after 24 hours. Control cells

were treated with a DMSO volume equivalent to the RA volume in the differentiating cells.

After 48 hours, the cells were harvested, washed once with PBS, counted and used

immediately in either ChIP-seq or ATAC-seq experiments.

1145

1146   **Mouse embryonic stem cell culture for ChIP-qPCR**

1147

TetR-CBX8 reporter-integrated mESCs were cultivated without feeders in high-glucose-

DMEM (Corning 10-013-CV) supplemented with 13.5% fetal bovine serum (Corning 35-015-

CV), 10 mM HEPES pH 7.4 (Corning, 25-060-CI), 2 mM GlutaMAX (Gibco, 35050-061), 1 mM

Sodium Pyruvate (Corning 25-000-CI), 1% Penicillin/Streptomycin (Sigma, P0781), 1X non-

essential amino acids (Gibco, 11140-050), 50 mM β-mercaptoethanol (Gibco, 21985-023)

and recombinant LIF. Cells were incubated at 37 °C and 5 % CO2 and were passaged every

48 h by trypsinization in 0.25 % 1x Trypsin-EDTA (Gibco, 25200-056). To reverse TetR-CBX8

binding, 1 µg/ml Doxycycline (Sigma, D9891) was added to cell culture medium for 6 hours.

1156

1157

1158

ChIP qPCR

25 x 10^6 reporter-integrated mESCs were collected, washed once in 1x PBS and cross-linked for 7 min in 1 % formaldehyde. The crosslinking was quenched by addition of 125 mM glycine and incubated on ice. The cross-linked ESCs were pelleted by centrifugation for 5 min at 1200 g at 4 °C. Nuclei were prepared by washes with NP-Rinse buffer 1 (10 mM Tris pH 8.0, 10 mM EDTA pH 8.0, 0.5 mM EGTA, 0.25 % Triton X-100) followed by NP-Rinse buffer 2 (10 mM Tris pH 8.0, 1 mM EDTA, 0.5 mM EGTA, 200 mM NaCl). Afterwards, the nuclei were washed twice with shearing buffer (1 mM EDTA pH 8.0, 10 mM Tris-HCl pH 8.0, 0.1 % SDS) and subsequently resuspended in 900 µL shearing buffer including 1x protease inhibitors Complete Mini cocktail (Roche). Chromatin was sheared by sonication in 15 ml Bioruptor tubes (Diagenode, C01020031) with 437.5 mg sonication beads (Diagenode, C03070001) for 6 cycles (1 min on and 1 min off) on a Bioruptor Pico sonicator (Diagenode). ChIP lysates equivalent to 50 ug DNA were incubated in 1x IP buffer (50 mM HEPES/KOH pH 7.5, 300 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% DOC, 0.1% SDS), and 1.5 ul of FLAG M2 antibody (Sigma Aldrich Sigma F1804) overnight. Antibody-bound chromatin was captured using Dynabeads protein G beads (Thermofisher #10004D) for 4 h at 4 °C. ChIPs were washed 5x with 1x IP buffer (50 mM HEPES/KOH pH 7.5, 300 mM NaCl, 1 mM EDTA, 1% Triton-X100, 0.1 % DOC, 0.1 % SDS), followed by 3x washes with DOC buffer (10 mM Tris pH 8, 0.25 mM LiCl, 1 mM EDTA, 0.5 % NP40, 0.5 % DOC) and 1x with TE/50 mM NaCl. ChIP DNA was eluted twice with elution buffer (1 % SDS, 0.1 M NaHCO₃) at 65 °C for 20 min, and subsequently treated with RNase A (60 ug final, Invitrogen) for 30 min at 37 °C, and Proteinase K (15ug, NEB) for 3 h at 55 °C and crosslinks were reversed overnight at 65 °C. The following day, ChIP samples and corresponding inputs were purified by AMPure XP beads (Beckman Coulter A63880).

ChIP-qPCR primers:

Name	Forward primer sequence	Reverse primer sequence
TetO	AAGATGGGCTGCAGGAATTC	ATACACGCCTACCTCGACATAC
+0.6 kb	GCAGGACGTGACAAATGGAAG	AAAGCGAAGGAGCAAAGCTG
+1.0 kb	GATCCGGACCGCCACATC	ACACCTTGCCGATGTCGAG
+1.9 kb	CAGTGCCACGTTGTGAGTTG	GCCCCTTGTTGAATACGCTTG
IAP	CTCCATGTGCTCTGCCTTCC	CCCCGTCCCTTTTTTAGGAGA

ChIP-seq

ChIP-seq was done as described previously[75]. 1.5 µg of CBX8 antibody (Cell Signalling, CBX8 (D2O8C), cat # 14696S) and 3 µg of H3K27me3 antibody (Tri-Methyl-Histone H3 (Lys27) (C36B11), cat #35861SF) were used. Libraries were prepared using NEBNext ultra II DNA library kit for Illumina (NEB Biolabs) according to the manufacturer instructions. The

1193 resulting libraries were assessed for quality on a High Sensitivity D1000 Screen Tape

1194 (Agilent) and were sequenced using an Illumina Novaseq 6000 sequencer (Genewiz/Azenta).

1195

1196 **Data processing for Chip-seq**

1197

The reads were quality-trimmed and adapters were removed using Trim Galore!, a wrapper

script to cutadapt[76], in paired end mode and using default settings. The reads were then

aligned to the mouse mm10 genome build using bowtie2 (version 2.3.5)[77] with the option

"very-sensitive". The data was reduced to only properly paired reads using "samtools view"

(Samtools version 1.9) with the flag "-f 3". PCR duplicates were removed using the

RemoveDuplicates function from Picard Tools (version 2.19.0). Read mates were fixed using

samtools fixmate. BigWig files were calculated using BamCoverage (Deeptools version 3.5.2)

with CPM normalisation.

Peaks were called with Macs2 (version 2.1.1)[78] callpeak function, using the input sample as

control. For the CBX8 ChIP-seq data set, default settings with a q-value cut-off of 0.05 were

used. For H3K27me3 ChIP-seq data set, broad mode was used with a q-value and broad

cutoff of 0.001. Peaks overlapping with ENCODE blacklisted regions[79] were removed.

1210

1211 **ATAC-seq**

1212

1213 ATAC-seq was done using a commercial kit (Diagenode Cat.# C01080002) according to the

1214 instructions of the manufacturer. The resulting libraries were assessed for quality control on

1215 a High Sensitivity D1000 Screen Tape (Agilent) and libraries were sequenced using an

1216 Illumina Novaseq 6000 sequencer (Genewiz/Azenta).

1217

1218 **Data processing ATAC-seq**

1219

ATAC-seq data was processed as described previously[80]. Specifically, reads were quality-

trimmed and adapters were removed using Trim Galore!, a wrapper script to cutadapt[76], in

paired end mode using default settings. The reads were then aligned to the mouse mm10

genome build using bowtie2 (version 2.3.5)[77] with the option "very-sensitive". Reads were

sorted and indexed using Samtools (version 1.9). Mitochondrial reads were removed using a

python script from Harvard Bioinformatics (available at

https://github.com/harvardinformatics/ATAC-seq). The data was reduced to only properly

paired reads using "samtools view" with the flag "-f 3". The library complexity was

estimated and the data sets were subsampled to reach a similar complexity as described

previously[80]. Data shown in Fig. 5h and Extended Data Fig 8b were not subsampled because

complexity was nearly identical. PCR duplicates were removed using the RemoveDuplicates

function from Picard Tools (version 2.19.0). Read mates were fixed using samtools fixmate.

BigWig files were calculated using BamCoverage (Deeptools version 3.5.2) with CPM

normalisation

For peak calling the bam files were converted to BEDPE files and the Tn5 shift was corrected
by running a bash script provided at https://github.com/reskejak/ATAC-seq
(bedpeTn5Shift.sh). Files were then converted to minimal bed format and peaks were called
using Macs2 (version 2.1.1)[78] callpeak function in broad mode with broad-cutoff set to 0.05.
Peaks overlapping with ENCODE blacklisted regions[79] were removed. Consensus peaks for
each condition were defined as the intersect of peaks from both biological replicates. Venn
diagrams were generated using the ChIPPeakAnno package[81].
1241
1242
1243

1244
1245
1246

References

1. Misteli, T. The Self-Organizing Genome: Principles of Genome Architecture and

Function. *Cell* **183**, 28–45 (2020).

2. Schuettengruber, B., Bourbon, H.-M., Di Croce, L. & Cavalli, G. Genome Regulation by

Polycomb and Trithorax: 70 Years and Counting. *Cell* **171**, 34–57 (2017).

3. Blackledge, N. P., Rose, N. R. & Klose, R. J. Targeting Polycomb systems to regulate

gene expression: modifications to a complex story. *Nat Rev Mol Cell Biol* **16**, 643–649

(2015).

4. Eskeland, R. *et al.* Ring1B Compacts Chromatin Structure and Represses Gene

Expression Independent of Histone Ubiquitination. *Molecular Cell* **38**, 452–464 (2010).

5. Francis, N. J., Kingston, R. E. & Woodcock, C. L. Chromatin Compaction by a Polycomb

Group Protein Complex. *Science* **306**, 1574–1577 (2004).

6. Grau, D. J. *et al.* Compaction of chromatin by diverse Polycomb group proteins requires

localized regions of high charge. *Genes Dev.* **25**, 2210–2221 (2011).

7. Lau, M. S. *et al.* Mutation of a nucleosome compaction region disrupts Polycomb-

mediated axial patterning. *Science* **355**, 1081–1084 (2017).

8. Isono, K. *et al.* SAM Domain Polymerization Links Subnuclear Clustering of PRC1 to

Gene Silencing. *Developmental Cell* **26**, 565–577 (2013).

9. Tatavosian, R. *et al.* Nuclear condensates of the Polycomb protein chromobox 2 (CBX2)

assemble through phase separation. *Journal of Biological Chemistry* **294**, 1451–1463

(2019).

10. Plys, A. J. *et al.* Phase separation of Polycomb-repressive complex 1 is governed by a

charged disordered region of CBX2. *Genes Dev.* **33**, 799–813 (2019).

11. Illingworth, R. S. Chromatin folding and nuclear architecture: PRC1 function in 3D.

Current Opinion in Genetics & Development **55**, 82–90 (2019).

12. Miron, E. *et al.* Chromatin arranges in chains of mesoscale domains with nanoscale

functional topography independent of cohesin. *Science Advances* **6**, eaba8811 (2020).

13. Nozaki, T. *et al.* Dynamic Organization of Chromatin Domains Revealed by Super-

Resolution Live-Cell Imaging. *Molecular Cell* **67**, 282-293.e7 (2017).

14. Murphy, S. & Boettiger, A. N. Polycomb repression of Hox genes involves spatial

feedback but not domain compaction or demixing. 2022.10.14.512199 Preprint at

https://doi.org/10.1101/2022.10.14.512199 (2022).

15. King, H. W., Fursova, N. A., Blackledge, N. P. & Klose, R. J. Polycomb repressive

complex 1 shapes the nucleosome landscape but not accessibility at target genes.

Genome Res. **28**, 1494–1507 (2018).

16. Chittock, E. C., Latwiel, S., Miller, T. C. R. & Müller, C. W. Molecular architecture of

polycomb repressive complexes. *Biochemical Society Transactions* **45**, 193–205 (2017).

17. Gibson, B. A. *et al.* Organization of Chromatin by Intrinsic and Regulated Phase

Separation. *Cell* **179**, 470-484.e21 (2019).

18. Maeshima, K. *et al.* Nucleosomal arrays self-assemble into supramolecular globular

structures lacking 30-nm fibers. *The EMBO Journal* **35**, 1115–1132 (2016).

19. Zhang, M. *et al.* Molecular organization of the early stages of nucleosome phase

separation visualized by cryo-electron tomography. *Molecular Cell* **82**, 3000-3014.e9

(2022).

20. Strickfaden, H. *et al.* Condensed Chromatin Behaves like a Solid on the Mesoscale In

Vitro and in Living Cells. *Cell* **183**, 1772-1784.e13 (2020).

21. Cai, S., Böck, D., Pilhofer, M. & Gan, L. The in situ structures of mono-, di-, and

trinucleosomes in human heterochromatin. *MBoC* **29**, 2450–2457 (2018).

22. Farr, S. E., Woods, E. J., Joseph, J. A., Garaizar, A. & Collepardo-Guevara, R.

Nucleosome plasticity is a critical element of chromatin liquid–liquid phase separation

and multivalent nucleosome interactions. *Nat Commun* **12**, 2883 (2021).

23. Morey, L. *et al.* Nonoverlapping Functions of the Polycomb Group Cbx Family of

Proteins in Embryonic Stem Cells. *Cell Stem Cell* **10**, 47–62 (2012).

24. Chung, C.-Y. *et al.* Cbx8 Acts Non-canonically with Wdr5 to Promote Mammary

Tumorigenesis. *Cell Reports* **16**, 472–486 (2016).

25. Tan, J. *et al.* CBX8, a Polycomb Group Protein, Is Essential for MLL-AF9-Induced

Leukemogenesis. *Cancer Cell* **20**, 563–575 (2011).

26. Voss, N. R. & Gerstein, M. 3V: cavity, channel and cleft volume calculator and extractor.

Nucleic Acids Research **38**, W555–W562 (2010).

27. Hou, Z., Nightingale, F., Zhu, Y., MacGregor-Chatwin, C. & Zhang, P. Structure of native

chromatin fibres revealed by Cryo-ET in situ. *Nature Communications* **14**, 6324 (2023).

28. Stefan Niekamp, Sharon K. Marr, Theresa A. Oei, Radhika Subramanian, & Robert E.

Kingston. Modularity of PRC1 Composition and Chromatin Interaction define

Condensate Properties. *Mol Cell*. **84**, 1651-1666 (2024) doi:10.1101/2023.10.26.564217.

29. Gambin, Y. *et al.* Single-Molecule Fluorescence Reveals the Oligomerization and

Folding Steps Driving the Prion-like Behavior of ASC. *Journal of Molecular Biology* **430**,

491–508 (2018).

30. Sierecki, E. *et al.* Nanomolar oligomerization and selective co-aggregation of α-synuclein

pathogenic mutants revealed by single-molecule fluorescence. *Scientific Reports* **6**,

37630 (2016).

31. Huseyin, M. K. & Klose, R. J. Live-cell single particle tracking of PRC1 reveals a highly

dynamic system with low target site occupancy. *Nat Commun* **12**, 887 (2021).

32. Barbieri, M. *et al.* Complexity of chromatin folding is captured by the strings and binders

switch model. *Proceedings of the National Academy of Sciences* **109**, 16173–16178

(2012).

33. Erdel, F. & Rippe, K. Formation of Chromatin Subcompartments by Phase Separation.

Biophysical Journal **114**, 2262–2270 (2018).

34. Lawrimore, J. *et al.* The rDNA is biomolecular condensate formed by polymer–polymer

phase separation and is sequestered in the nucleolus by transcription and R-loops.

Nucleic Acids Research **49**, 4586–4598 (2021).

35. Brown, K. *et al.* Principles of assembly and regulation of condensates of Polycomb

repressive complex 1 through phase separation. *Cell Reports* **42**, (2023).

36. McGinty, R. K., Henrici, R. C. & Tan, S. Crystal structure of the PRC1 ubiquitylation

module bound to the nucleosome. *Nature* **514**, 591–596 (2014).

37. Connelly, K. E. *et al.* Engagement of DNA and H3K27me3 by the CBX8 chromodomain

drives chromatin association. *Nucleic Acids Research* **47**, 2289–2305 (2019).

38. Bernstein, E. *et al.* Mouse Polycomb Proteins Bind Differentially to Methylated Histone

H3 and RNA and Are Enriched in Facultative Heterochromatin. *Molecular and Cellular*

Biology **26**, 2560–2569 (2006).

39. Fleming, P. J. & Fleming, K. G. HullRad: Fast Calculations of Folded and Disordered

Protein and Nucleic Acid Hydrodynamic Properties. *Biophysical Journal* **114**, 856–869

(2018).

40. Davies, D. R., Goryshin, I. Y., Reznikoff, W. S. & Rayment, I. Three-Dimensional

Structure of the Tn5 Synaptic Complex Transposition Intermediate. *Science* **289**, 77–85

(2000).

41. Meléndez-Ramírez, C. *et al.* Dynamic landscape of chromatin accessibility and

transcriptomic changes during differentiation of human embryonic stem cells into

dopaminergic neurons. *Sci Rep* **11**, 16977 (2021).

42. Suh, J. L. *et al.* Reprogramming CBX8-PRC1 function with a positive allosteric

modulator. *Cell Chemical Biology* **29**, 555-571.e11 (2022).

43. Seif, E. *et al.* Phase separation by the polyhomeotic sterile alpha motif

compartmentalizes Polycomb Group proteins and enhances their activity. *Nat Commun*

11, 5609 (2020).

44. Gray, F. *et al.* BMI1 regulates PRC1 architecture and activity through homo- and hetero-

oligomerization. *Nat Commun.* **7**, 13343 (2016).

45. He, S. *et al.* Structure of nucleosome-bound human BAF complex. *Science* **367**, 875–

881 (2020).

46. Wang, H. *et al.* Structure of the transcription coactivator SAGA. *Nature* **577**, 717–720

(2020).

47. Schilbach, S., Aibara, S., Dienemann, C., Grabbe, F. & Cramer, P. Structure of RNA

polymerase II pre-initiation complex at 2.9 Å defines initial DNA opening. *Cell* **184**, 4064-

4072.e28 (2021).

48. McCall, K. & Bender, W. Probes of chromatin accessibility in the Drosophila bithorax

complex respond differently to Polycomb-mediated repression. *The EMBO Journal* **15**,

569–580 (1996).

49. Maeshima, K. *et al.* The physical size of transcription factors is key to transcriptional

regulation in chromatin domains. *J. Phys.: Condens. Matter* **27**, 064116 (2015).

50. Bepler, T., Kelley, K., Noble, A. J. & Berger, B. Topaz-Denoise: general deep denoising

models for cryoEM and cryoET. *Nat Commun* **11**, 5208 (2020).

51. Zhang, Q. *et al.* PALI1 facilitates DNA and nucleosome binding by PRC2 and triggers an

allosteric activation of catalysis. *Nat Commun* **12**, 4592 (2021).

52. Brzovic, P. S., Lissounov, A., Christensen, D. E., Hoyt, D. W. & Klevit, R. E. A

UbcH5/Ubiquitin Noncovalent Complex Is Required for Processive BRCA1-Directed

Ubiquitination. *Molecular Cell* **21**, 873–880 (2006).

53. Pichler, A. *et al.* SUMO modification of the ubiquitin-conjugating enzyme E2-25K. *Nat*

Struct Mol Biol **12**, 264–269 (2005).

54. Uckelmann, M. *et al.* USP48 restrains resection by site-specific cleavage of the BRCA1

ubiquitin mark from H2A. *Nat Commun* **9**, 229 (2018).

55. Luger, K., Rechsteiner, T. J. & Richmond, T. J. Preparation of nucleosome core particle

from recombinant histones. in *Methods in Enzymology* vol. 304 3–19 (Academic Press,

1999).

56. Sabantsev, A., Levendosky, R. F., Zhuang, X., Bowman, G. D. & Deindl, S. Direct

observation of coordinated DNA movements on the nucleosome during chromatin

remodelling. *Nat Commun* **10**, 1720 (2019).

57. Hagen, W. J. H., Wan, W. & Briggs, J. A. G. Implementation of a cryo-electron

tomography tilt-scheme optimized for high resolution subtomogram averaging. *Journal of*

Structural Biology **197**, 191–198 (2017).

58. Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for

improved cryo-electron microscopy. *Nat Methods* **14**, 331–332 (2017).

59. Mastronarde, D. N. Dual-Axis Tomography: An Approach with Alignment Methods That

Preserve Resolution. *Journal of Structural Biology* **120**, 343–352 (1997).

60. Himes, B. A. & Zhang, P. emClarity: software for high-resolution cryo-electron

tomography and subtomogram averaging. *Nat Methods* **15**, 955–961 (2018).

61. Castaño-Díez, D., Kudryashev, M., Arheit, M. & Stahlberg, H. Dynamo: A flexible, user-

friendly development tool for subtomogram averaging of cryo-EM data in high-

performance computing environments. *Journal of Structural Biology* **178**, 139–151

(2012).

62. Chua, E. Y. D. *et al.* 3.9 Å structure of the nucleosome core particle determined by

phase-plate cryo-EM. *Nucleic Acids Res* **44**, 8013–8019 (2016).

63. Davey, C. A., Sargent, D. F., Luger, K., Maeder, A. W. & Richmond, T. J. Solvent

Mediated Interactions in the Structure of the Nucleosome Core Particle at 1.9Å

Resolution††We dedicate this paper to the memory of Max Perutz who was particularly

inspirational and supportive to T.J.R. in the early stages of this study. *Journal of*

Molecular Biology **319**, 1097–1113 (2002).

64. Kasinath, V. *et al.* Structures of human PRC2 with its cofactors AEBP2 and JARID2.

Science **359**, 940–944 (2018).

65. Chen, X. *et al.* Structures of +1 nucleosome–bound PIC-Mediator complex. *Science* **378**,

62–68 (2022).

66. Leitão, A. D. G. *et al.* Selectivity of Lewy body protein interactions along the aggregation

pathway of α-synuclein. *Communications Biology* **4**, 1124 (2021).

67. Lau, D. *et al.* Single Molecule Fingerprinting Reveals Different Amplification Properties of

α-Synuclein Oligomers and Preformed Fibrils in Seeding Assay. *ACS Chem. Neurosci.*

13, 883–896 (2022).

68. Bhumkar, A. *et al.* Single-Molecule Counting Coupled to Rapid Amplification Enables

Detection of α-Synuclein Aggregates in Cerebrospinal Fluid of Parkinson's Disease

Patients. *Angewandte Chemie International Edition* **60**, 11874–11883 (2021).

69. Horn, V. *et al.* Structural basis of specific H2A K13/K15 ubiquitination by RNF168. *Nat*

Commun **10**, 1751 (2019).

70. Yang, B. *et al.* Identification of cross-linked peptides from complex samples. *Nat*

Methods **9**, 904–906 (2012).

71. Gail, E. H., Shah, A. D., Schittenhelm, R. B. & Davidovich, C. crisscrosslinkeR:

identification and visualization of protein–RNA and protein–protein interactions from

crosslinking mass spectrometry. *Bioinformatics* **36**, 5530–5532 (2020).

72. Combe, C. W., Fischer, L. & Rappsilber, J. xiNET: Cross-link Network Maps With

Residue Resolution*. *Molecular & Cellular Proteomics* **14**, 1137–1147 (2015).

73. Canver, M. C. *et al.* A saturating mutagenesis CRISPR-Cas9–mediated functional

genomic screen identifies cis- and trans-regulatory elements of Oct4 in murine ESCs.

Journal of Biological Chemistry **295**, 15797–15809 (2020).

74. Seruggia, D. *et al.* TAF5L and TAF6L Maintain Self-Renewal of Embryonic Stem Cells

via the MYC Regulatory Network. *Molecular Cell* **74**, 1148-1163.e7 (2019).

75. Healy, E. *et al.* PRC2.1 and PRC2.2 Synergize to Coordinate H3K27 Trimethylation.

Molecular Cell **76**, 437-452.e6 (2019).

76. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17**, 10–12 (2011).

77. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357–359 (2012).

78. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biology* **9**, R137

(2008).

79. Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of

Problematic Regions of the Genome. *Sci Rep* **9**, 9354 (2019).

80. Reske, J. J., Wilson, M. R. & Chandler, R. L. ATAC-seq normalization method can

significantly affect differential accessibility analysis and interpretation. *Epigenetics &*

Chromatin **13**, 22 (2020).

81. Zhu, L. J. *et al.* ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and

ChIP-chip data. *BMC Bioinformatics* **11**, 237 (2010).

1442