Dynamic PRC1-CBX8 stabilizes a porous structure of chromatin condensates

Michael Uckelmann[1], Vita Levina[1], Cyntia Taveneau[1,2], Xiao Han Ng[1], Varun Pandey[3], Qi

Zhang[1], Sarena Flanigan[1], Minrui Li[1,4], Partha Pratim Das[3], Alex de Marco[1,2], Chen

Davidovich[1,5,*]

[1] Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute,
Faculty of Medicine, Nursing and Health Sciences, Monash University; Clayton, Victoria,
3800, Australia.

[2] ARC Centre of Excellence in Advanced Molecular Imaging, Monash University; Clayton,
Victoria, 3800, Australia.

[3] Department of Anatomy and Developmental Biology, Development and Stem Cells
Program, Monash Biomedicine Discovery Institute, Monash University; Clayton, Victoria,
3800, Australia.

[4] Faculty of Information Technology, Monash University; Clayton, Victoria, 3800, Australia.

[5] EMBL-Australia; Clayton, Victoria, 3800, Australia.

* Correspondence: Chen.Davidovich@monash.edu

Abstract

The compaction of chromatin is a prevalent paradigm in gene repression. Chromatin
compaction is commonly thought to repress transcription by restricting chromatin
accessibility. However, the spatial organisation and dynamics of chromatin compacted by
gene-repressing factors are unknown. Using cryo-electron tomography, we solved the three-
dimensional structure of chromatin condensed by the Polycomb Repressive Complex 1
(PRC1) in a complex with CBX8. PRC1-condensed chromatin is porous and stabilised through
multivalent dynamic interactions of PRC1 with chromatin. Within condensates, PRC1
remains dynamic while maintaining a static chromatin structure. In differentiated mouse
embryonic stem cells, CBX8-bound chromatin remains accessible. These findings challenge
the idea of rigidly compacted polycomb domains and instead provides a mechanistic
framework for dynamic and accessible PRC1-chromatin condensates.

Main

Chromatin structure is intricately linked to transcriptional activity[1]. Compacted or "closed"
chromatin is generally associated with inhibition of transcription while "open", more
accessible chromatin is more prone to being transcribed[1]. Polycomb Repressive Complex 1
(PRC1) is a repressive chromatin modifier critical for organismal development[2,3]. PRC1 has
been proposed to inhibit gene expression by tightly compacting chromatin in a process that
often is considered to restrict chromatin accessibility[2–11]. However, direct evidence for

PRC1-compacted chromatin being inaccessible is sparse and mechanistic explanations
remain unsatisfactory (reviewed in[11]). Furthermore, recent studies show that changes in
chromatin accessibility are more gradual than the simple binary classification into "open"
and "closed" chromatin suggests[12–15]. A challenge in consolidating these seemingly
contradictory findings is limited information into how PRC1 influences the three dimensional
structure of chromatin.

PRC1 complexes can include one of five different chromobox proteins (CBX), all homologous
to the fly Polycomb (Pc)[16]. The CBX protein CBX2 forms condensates through liquid-liquid
phase separation, providing a potential mechanism for the compartmentalization of
facultative heterochromatin[9,10]. Phase separation is emerging as a mechanism for chromatin
organisation through the association of self-similar domains[17]. Chromatin can form
condensates in the presence of divalent cations and histone tails[17–20]. Within these
condensates, chromatin has variably been described as liquid-like, formed through liquid-
liquid phase separation[17,19], or as solid[20]. A recent structure of liquid-liquid phase-separated
chromatin, condensed by magnesium cations without protein binding partners, revealed
that nucleosomes organise into irregular assemblies[19]. The lack of apparent periodicity in
chromatin geometry has also been noted in computational simulations and in first attempts
to image chromatin in cells by cryo-electron tomography (cryo-ET)[21,22]. However, the
structural arrangement of chromatin condensed by a repressive factor remained unknown.

Herein we describe the three dimensional cryo-ET structure of chromatin condensed by a
polycomb-repressive complex. We focus on a PRC1 complex that includes CBX8 (PRC1^{C8}), a
chromobox protein that is upregulated during cell differentiation[23] and has oncogenic
potential[24,25]. We show that dynamic interactions between PRC1^{C8} and chromatin promote
condensates through polymer-polymer phase separation. Contrary to expectations, PRC1-
condensed chromatin is not tightly compacted but stabilises a porous chromatin structure
that allows largely unhindered diffusion of PRC1^{C8}.

Results

PRC1-chromatin condensates are porous and accessible

To determine the structure of polycomb-compacted chromatin and the mechanisms of
polycomb-driven chromatin compaction, we reconstituted the system *in vitro*. The
reconstitution included a chromatinized polycomb target gene (referred to as chromatin
hereafter) and a purified recombinant PRC1 complex composed of RING1b, BMI1 and CBX8
(PRC1^{C8}) (Fig. 1a). The PRC1^{C8} complex is pure (Fig. 1b), monodispersed (Fig. 1c) and retains
H2A ubiquitylation activity comparable to the RING1b-BMI1 heterodimer (Fig. 1d).

When combined, chromatin and PRC1^{C8} were sufficient to form spherical phase-separated
condensates, apparent in phase-contrast and fluorescence imaging (Fig. 1e). Using two
different fluorescence labels, we confirmed the presence of both chromatin and PRC1^{C8}

within the same condensates (Fig. 1e). Importantly, both PRC1^{C8} and chromatin are
necessary for chromatin condensation, while the individual components do not phase-
separate (Fig. 1e). PRC1^{C8}-chromatin condensates are preserved on an EM grid after
vitrification (Fig. 1f). This allowed us to study the internal molecular structure of PRC1^{C8}-
chromatin condensates by cryo-electron tomography (Fig. 1g). We collected tomograms
near the condensate border to observe the boundary conditions. The reconstruction allows
individual nucleosomes to be identified, revealing a dense network of hundreds of
nucleosomes with a distinct condensate boundary (Fig. 1G and Movie S1). We could not
unambiguously assign density to PRC1^{C8}, possibly because it adapts various conformations
while simultaneously using multiple surfaces to interact with chromatin (more below). The
final structure reflects the arrangement of nucleosomes in PRC1^{C8}-chromatin condensate
(Fig. 1g, second panel). Unexpectedly, the structure shows that PRC1^{C8} does not compact
nucleosomes into an impassable barrier. Instead, PRC1^{C8} rather stabilises chromatin in a
porous mesh-like structure (Fig. 1g). Analysing the orientation of individual nucleosomes
towards their neighbouring nucleosomes shows no obvious orientation bias (Extended Data
Fig. 1a,b). We conclude that PRC1^{C8} does not induce a substantial inter-nucleosome
orientation bias, but rather supports forming a porous chromatin structure.

We next wished to determine the size of macromolecules that could diffuse into PRC1-
chromatin condensates. We used the condensate structure to calculate solvent-excluded
volumes[26] with variable probe radii ranging from 0.2 nm to 20 nm (Fig. 1h,i). Interestingly,
the analysis shows that the condensate is accessible for macromolecules of a considerable
size of up to 8 nm in radius (equivalent to approximately 600 kDa). Small macromolecules
(<10 kDa), with radii below 2 nm, would have enough room to access every single
nucleosome. Conversely, access is increasingly restricted for molecules with a radius above
8 nm (approximately 600 kDa). This suggests that PRC1-chromatin condensates are
surprisingly accessible and that PRC1^{C8} itself would be able to move within these
condensates largely unhindered. To test this we next ventured to analyse PRC1 dynamics
within condensates experimentally.

PRC1^{C8} is mobile and chromatin is static within PRC1-chromatin condensates

Chromatin condensates form under close to physiological salt concentrations, at PRC1^{C8}
concentrations as low as 500 nM and are dependent on PRC1^{C8} (Fig. 2a) .We conclude that
PRC1^{C8} is sufficient to drive the formation of the chromatin condensates under
physiologically relevant conditions. We next asked if chromatin and PRC1 show different
dynamics within the condensates. Fluorescence recovery after photobleaching (FRAP) of
PRC1^{C8}-chromatin condensates shows fast recovery kinetics for GFP-labelled PRC1 (Fig. 2b,
in green). Conversely, we observed a very slow recovery for Cy5-labelled chromatin (Fig. 2b,
in red). We conclude that, within condensates, PRC1 is mobile while chromatin itself is static
(Fig. 2c). This confirms that PRC1^{C8} can diffuse largely unhindered within PRC1^{C8}-chromatin
condensates, in line with our structural analysis (Fig. 1h,i).

Multivalent interactions between PRC1[C8] and chromatin induce phase separation

Previous studies have concluded that magnesium-driven chromatin condensates form *in*
vitro via liquid-liquid phase separation[17,19]. This view is challenged by the observation of
solid-like chromatin condensates in living cells[20]. Our observation that chromatin is static
within PRC1-chromatin condensates does not agree with liquid-like behaviour, but rather
suggests polymer-polymer phase separation (PPPS) as a mechanism for condensate
formation[27–29]. PPPS depends on one polymer acting as a scaffold that is condensed by the
second molecule binding to it via multivalent interactions, effectively acting like a
crosslinker[27–29].

To test for polymer-polymer phase separation of condensates, we next probed for
multivalent interactions between PRC1 and chromatin. The whole PRC1[C8] complex (RING1b,
BMI1 and CBX8) is necessary and sufficient to condense chromatin, while the PRC1 core or
CBX8 alone do not condense chromatin (Fig. 3a). This suggests multivalent interactions
between the PRC1[C8] complex and chromatin, involving different chromatin interacting
surfaces in both PRC1 and CBX8. To identify the different interaction sites, we first used
crosslinking mass spectrometry (XL-MS) to probe for protein-protein interactions within
PRC1[C8] –chromatin condensates (Fig. 3b). The results show multiple crosslinks from the
CBX8 chromodomain to the H3 histone tail (Fig. 3b), indicative of binding. Interactions
between CBX-proteins and H3K27me3-modified H3-histone tails have been proposed to
recruit PRC1 to chromatin modified by PRC2[2,3]. However, CBX8 did crosslink to
unmethylated H3 tails (Fig. 3b) and a trimethyl-lysine analogue (MLA) at H3K27 did not
improve the chromatin-condensation activity of PRC1[C8] (Fig. 3d). We conclude that
H3K27me3 is not necessary for the chromatin condensation activity of PRC1[C8] and that the
H3 histone tail, even if unmodified, provides an interaction site for PRC1 on chromatin.

DNA has previously been shown to bind CBX8[30] and could provide another interaction site
for PRC1[C8] on chromatin. We tested DNA binding in solution using a 24bp double stranded
DNA probe and found that CBX8 is necessary for the DNA-binding activity of PRC1[C8] (Fig. 3c).
Hence, CBX8 binding to DNA provides a second interaction site of PRC1 with chromatin.

We conclude that PRC1 interacts with chromatin via at least three distinct sites: PRC1[C8]
binds to DNA and the H3 tail via CBX8, as shown herein (Fig. 3), and binds the nucleosome
acidic patch via RING1b-BMI1 as shown elsewhere[31]. These multivalent interactions would
have to change dynamically while PRC1 maintains the condensed state of chromatin and
diffuses through it at the same time (Fig. 2b). Collectively, we propose that PRC1 induces
chromatin condensation via polymer-polymer phase separation, through dynamic
multivalent interactions between PRC1 and chromatin.

CBX8 binding sites on chromatin in mouse embryonic stem cells are accessible

Given the porous structure of PRC1^{C8}-condensed chromatin in vitro (Fig. 1) and the dynamic
diffusion of PRC1^{C8} within condensates (Fig. 2b), we next wished to probe for the
accessibility of PRC1^{C8}-bound chromatin in cells. We carried out the Assay for Transposase
Accessible Chromatin (ATAC-seq) in differentiated mESC, combined with ChIP-seq for CBX8
and H3K27me3. We used differentiated mESC, because CBX8 is expressed at very low levels
in pluripotent mESC and is upregulated during retinoic acid-induced cell differentiation ([23]
and Fig. 4a). The DNA-loaded Tn5 used in ATAC-seq experiments forms a dimeric complex of
approximately 130 kDa with an estimated hydrodynamic radius[32] of 4.6 nm (based on PDB
code 1MUH[33]). In agreement with the accessibility analysis *in vitro* (Fig. 1h), the majority of
CBX8 ChIP-seq peaks in cells overlapped with ATAC-seq peaks (Figure 4B), indicating they
are accessible to Tn5. This observation was persistent across the genome, where ATAC-seq
peaks are co-localised with CBX8 and H3K27me3 peaks (Fig. 4c), indicating that CBX8-target
genes are largely accessible. The insufficiency of CBX8 to restrict chromatin accessibility is
further supported by the similar ATAC-seq profiles of wildtype and *Cbx8* knockout mESCs
(Fig. 4c,f, compare blue to orange). Hence, although the overall chromatin accessibility is
reduced during mESC differentiation (Fig. 4d,e), in agreement with previous works,[34] this
process is not dependent on CBX8. Collectively, we show that CBX8-bound polycomb-
repressed chromatin is largely accessible in differentiated mECSs (Fig. 4b-f).

Discussion

In conclusion, we have shown that PRC1-CBX8 binds chromatin via multivalent interactions
and induces chromatin condensation through polymer-polymer phase separation. PRC1^{C8} is
dynamic within condensates while keeping chromatin in a static, solid-like state. We
established that PRC1^{C8} is sufficient to induce the solid condensed state of chromatin.

How can PRC1 condense chromatin but yet maintain a highly dynamic behaviour in the
nucleus? PRC1 is characterised with a short residence time on chromatin[35]. Solid-like
chromatin that is condensed by a mobile chromatin binder has been shown *in vitro* for a
truncation of the SAM-domain protein Polyhomeotic (Ph)[36] and in cells for HP1a at
chromocenters[20]. Yet, the mechanism allowing PRC1 and other gene-repressing factors to
condense chromatin while constantly diffusing in it remained largely unknown. Our cryo-EM
structure of PRC1-condensed chromatin explains how PRC1 can move in condensed
chromatin (Fig. 1), owing to the large pores that are formed between condensed
nucleosomes. The multivalent interactions between PRC1-CBX8 to chromatin provide PRC1
with multiple docking sites on chromatin: unmodified and modified H3 tail (Fig 3b-d), DNA
(Fig 3c) and the acidic patch of the nucleosome[31]. Hence, it is possible that PRC1-CBX8 can
constantly change its interactions with chromatin to maintain its condensed structure while
utilising its different chromatin-interacting surfaces to dynamically move around. Our data
indicate that chromatin condensation together with dynamic behaviour within chromatin
condensates is an intrinsic biophysical property of PRC1-CBX8. It is plausible that the
dynamic behaviour of PRC1 within chromatin condensates is required in order to allow PRC1

to modify nucleosomes by the H2AK119ub mark while holding them together. This
phenomenon might represent a broad paradigm of repressive chromatin.The internal
structure of PRC1-chromatin condensates is a porous network of nucleosomes (Fig. 1). Such
a structure could present a size-selective diffusion barrier, in agreement with its
permeability to PRC1 diffusion in vitro (Fig. 2) and Tn5 accessibility in cells (Fig. 4). Such a
size-selective diffusion barrier may contribute to gene repression by selectively excluding
transcriptional coactivators, which are commonly large protein complexes (>1MDa[12,37–39]).
This is in agreement with the inverse correlation between the density of chromatin domains
and the molecular weight of the chromatin modifiers present in them[12]. Transcription factor
size has also been suggested to determine access to different chromatin domains based on
simulations[40]. Combining our results with earlier findings[12,19,20,40], we propose that size-
selective exclusion may be part of a broader mechanism by which chromatin-interacting
proteins regulate the accessibility of repressive chromatin.

Acknowledgements: We would like to thank the Ramaciotti centre for electron microscopy
at Monash university for providing instrumentation, technical support and collecting cryo-ET
data, the Monash Proteomics facility for providing instrumentation and technical support
and the Monash Micro Imaging facility for providing instrumentation and technical support.
We would also like to acknowledge the MASSIVE HPC platform for providing high-
performance computing resources
205
206 **Funding:** This work was supported by an Australian research council (ARC) DECRA fellowship
207 DE210101669 (M.U.), Sylvia and Charles Viertel Senior Medical Research Fellowship (C.D.)
208 and the National Health and Medical Research Council (NHMRC) grant numbers
209 APP1162921, APP1184637 and APP2011767 (C.D.). C.D. is an EMBL-Australia Group Leader.

210
211 **Authors contributions:** C.D. and M.U. conceptualised the project and acquired funding,
212 M.U., V.L., C.T., X.H.N and M.L. carried out experiments and investigated, P.P.D. and V.P.
213 generated cell lines, S.F. and Q.Z. cloned and purified histone mutants, M.L. and M.U.
214 developed software, C.D., A.d.M. and P.P.D. supervised, M.U. and C.D. wrote the original
215 draft and all authors reviewed and edited the manuscript.
216
217 **Declaration of interests:** The authors declare no conflict of interest.
218
219 **Data and materials availability:** Next generation sequencing data (ATAC-seq and ChIP-seq)
220 are available under GEO accession number GSE220140. The map for tomogram #1 has been
221 deposited to the EMD (EMD-29022). All cryo-ET raw data has been deposited to EMPIAR
222 under accession number EMPIAR-11344. XL-MS data has been deposited to Pride
223 (PXD039589)
224

Fig. 1. The molecular structure of PRC1-chromatin condensates is porous and accessible to macromolecules. a, Schematics introducing the workflow. **b,** SDS-PAGE of purified PRC1^{C8} complex that includes RING1b, BMI1 and MBP-tagged CBX8. **c,** Size exclusion chromatography of the purified PRC1^{C8} using a HiLoad Sephacryl 300 16/60 column. **d,** In vitro ubiquitylation assay comparing PRC1^{C8} to a RING1b-BMI1 heterodimer. Samples in lane 2 and 3 included E1, E2, ubiquitin and ATP. Ubiquitylation is detected by western blot using an anti-H2A antibody. **e,** Chromatin condensates induced by the PRC1^{C8} complex and the individual proteins, visualised by confocal (left and centre) and phase contrast (right, from independent experiments) microscopy. CBX8 is GFP-labelled and chromatin is Cy5 labelled. Protein and chromatin concentrations are 1 μM and 20 nM, respectively. **f,** Cryo-confocal microscopy of vitrified PRC1^{C8}-chromatin condensates. **g,** Cryo-electron tomography of a PRC1^{C8}-chromatin condensate. Shown is a central slice through the

reconstruction (left image). Nucleosome subtomogram averages (centre, bottom) are then
placed in a volume the size of the tomographic slice, at the position and orientation
determined by template matching and subtomogram averaging (right image). **h,** Surface
representation of the volume of a subset of the PRC1^{C8}-chromatin condensate structure that
is inaccessible to probes of given radii. **i,** Inaccessible volumes for a given probe radii are
plotted, with exemplary molecules indicated (in grey) and selected probes coloured as in **h**.
For the indicated complexes, the hydrodynamic radius was estimated[32] using resolved
domains from published structures (see Methods section for PDB accessions) as a minimum
size estimate. See also Extended Data Figure 1.

Fig. 2. PRC1^{C8} is mobile while chromatin is static within PRC1-chromatin condensates.
a, Chromatin condensation in response to salt and PRC1 concentration measured by
confocal microscopy using Cy5-labelled chromatin at a constant concentration of 50 ng/µl
DNA. **b,** Representative micrographs of FRAP recorded in PRC1^{C8}-chromatin condensates.
CBX8 is GFP labelled and chromatin is Cy5 labelled. Mean fluorescence intensity of the
bleached area, normalised to pre-bleach mean signal, is plotted for every time point. Error
bars show standard deviation from n=7 (GFP) and n=8 (Cy5) measurements recorded from
two independent experiments. **c,** Schematic representation of the FRAP experiment.

Fig. 3. Multivalent interactions between PRC1^{C8} and chromatin. a, Chromatin condensation in response to the whole PRC1^{C8} complex (RING1b, BMI1 and CBX8) or the individual components CBX8 and the RING1b–BMI1 heterodimer. Representative images from two replicates. **b,** Intermolecular (purple lines) and intramolecular (green lines) protein-protein interactions mapped within PRC1-chromatin condensates using crosslinking mass spectrometry (XL-MS). Data is from three independent replicates. **c,** PRC1^{C8} (blue) or PRC1 core (red) binding to a fluorescein labelled 24bp DNA probe measured by fluorescence anisotropy. Error bars show standard deviation based on two independent replicates. **d,** Titration of PRC1^{C8} to unmodified chromatin (top) and H3K27me3-MLA chromatin (bottom) at an identical chromatin concentration (50 ng/μl DNA) and 150 mM KCl. Micrographs are representative of two independent replicates.

Fig. 4. CBX8 binding sites on chromatin in mouse embryonic stem cells are accessible.
a, Schematics of the experimental setup (left) and anti-CBX8 western blot (right) of wildtype and *Cbx8* knockout mESC after 48 hours of retinoic acid (RA) treatment. **b,** Overlap of ATAC-seq peaks and CBX8 and H3K27me3 ChIP-seq peaks. ATAC-seq peaks are defined from two biological replicates. **c,** ChIP-seq traces for H3K27me3 and CBX8 in wildtype mESC and representative ATAC-seq traces at four genes in wildtype and CBX8 knockout mESC after 48 hours of RA treatment. **d,** Accessibility changes at all ATAC-seq peaks in wildtype (WT) mESC, in response to retinoic acid (RA) treatment. **e,** Accessibility changes at all CBX8-target sites in wildtype mESC, in response to RA treatment. **f,** Comparison of accessibility at CBX8-

280 target sites between wildtype and *Cbx8* knockout cells after RA treatment. **g,** Model: PRC1
281 forms multivalent interactions with chromatin, thereby stabilizing chromatin condensates.
282 These interactions dynamically change as PRC1 diffuses through condensates.
283

284   **Extended Data**

a

b

Extended Data Fig. 1. Related to Fig. 1; Nucleosomes in PRC1[C8]-chromatin condensates

show no orientation bias towards neighbouring nucleosomes. a, Orientations of

nucleosomes towards neighbouring nucleosomes within a cut-off of 20 nm. Individual points

represent nucleosomes and lines between points are coloured according to the relative

orientation of neighbouring nucleosomes as indicated in the colour key (left). **b,** Distribution

of nucleosome-nucleosome orientation for the three nearest neighbours and the 150th

neighbour of each nucleosome in tomogram #1. Colours correspond to the same respective

orientations as in **a**.

Extended Data Movie S1. Related to Fig.1; Cryo-tomogram of a PRC1^{C8}-chromatin

condensate. The movie shows a scan through the z-axis of denoised[51] tomogram #1.

Materials and Methods

Plasmids and cloning

Human RING1b (Uniprot ID Q99496) and human BMI1 (Uniprot ID P35226) were cloned into a pFBOH-mhl vector (Addgene plasmid # 62304) cleaved with BseRI using Gibson Assembly® Master Mix (NEB #E2611L) using the primers indicated in Table S1.

Human CBX8 open reading frame (Uniprot ID Q9HC52-1 and NCBI Reference Sequence was NM_020649.2) was obtained from gene synthesis (Gen9) cloned into a modified pFastBac1 pFB1.HMBP.A3.PrS.ybbR vector digested by XhoI and XmaI sites to include a N-terminal 6xHis-MBP tag. Cloning of the polycomb target gene ATOH1 into the pUC18 vector was described previously[41]. Plasmids for expression of human histones (H2A, H2B, H3.1 and H4) in *E.coli* were a kind gift from David Tremethick, Australian National University. UbcH5c WT pET28a was a gift from Rachel Klevit (Addgene plasmid # 12643; http://n2t.net/addgene:12643; RRID:Addgene_12643)[42].

pET3a-hUBA1 was a gift from Titia Sixma (Addgene plasmid # 63571; http://n2t.net/addgene:63571; RRID:Addgene_63571)[43].

To generate a baculovirus expression vector of a monomeric EGFP-CBX8 (mEGFP-CBX8) construct, first GFP was amplified from a pSpCas9(BB)-2A-GFP vector and CBX8 was amplified from a pFB1.HMBP.A3.PrS.ybbR vector containing CBX8 as an insert. Subsequently, EGFP-CBX8 was cloned into the pFBOH-mhl vector cleaved with BseRI via Gibson assembly, with a Serine-Glycine-Serine linker between EGFP and CBX8. Finally, to generate monomeric mEGFP-CBX8, alanine residue 207 in EGFP was mutated to Lysine using a site directed mutagenesis kit (Takeda) and the mutagenesis primers listed in Table 1.

Table 1: Cloning and mutagenesis primers (5'-3')

mEGFP mutagenesis A207K fw	CTGAGCACCCAGTCCAAGCTGAG CAAAGACCC	Used for mutagenesis of CBX8-EGFP
mEGFP mutagenesis A207K rv	GGGGTCTTTGCTCAGCTTGGACT GGGTGCTCA	Used for mutagenesis of CBX8-EGFP
EGFP_fw_PFBOH_MHL	TTGTATTTCCAGGGCATGGTGAG CAAGGGCGAG	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector
EGFP rv with SGS linker	TGAAAGCTCACTTCCACTCTTGTA CAGCTCGTCCATGC	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector
CBX8 5' with SGS linker	TGTACAAGAGTGGAAGTGAGCTT TCAGCGGTGGGG	Used for Gibson assembly of mEGFP-CBX8 into pFBOH-mhl vector
CBX8 rv pFBOH_MHL	CAAGCTTCGTCATCATCATCTTTT	Used for Gibson assembly of

	CTCTTTAAAAAAGCCT	mEGFP-CBX8 into pFBOH-mhl vector
ATOH1 fw	GCAGAGCCCAAACATTCACACA	Used to amplify ATOH1 from pUC18 plasmid
ATOH1 rv	GCGGAGTTTCCTAAAAGACGCC	Used to amplify ATOH1 from pUC18 plasmid
H2BT120C fw	GTGACCTGCTATACCAGCAGCAAATAA	Used for mutagenesis of H2B
H2BT120C rv	GGTATAGCAGGTCACGGCTTTGGTGCC	Used for mutagenesis of H2B
RING1b pFBOH-mhl fw	ttgtatttccagggcTCTCAGGCTGTGCAGACAAAC	Used for Gibson assembly into pFBOH-mhl vector
RING1b pFBOH-mhl rv	caagcttcgtcatcaTTTGTGCTCCTTTGTAGGTGC	Used for Gibson assembly into pFBOH-mhl vector
BMI1 pFBOH-mhl fw	ttgtatttccagggcCATCGAACAACGAGAATCAAG	Used for Gibson assembly into pFBOH-mhl vector
BMI1 pFBOH-mhl rv	caagcttcgtcatcaACCAGAAGAAGTTGCTGATGA	Used for Gibson assembly into pFBOH-mhl vector
24bp DNA duplex Fluorescein labeled	GGCGCCCTGCCCCGCCTCGCTCTG	Fluorescein labeled DNA duplex used for binding assays. Only the top strand is shown. The dye was attached to the 3' end of the top strand.

Protein expression and purification

PRC1[C8], PRC1 core and CBX8 were co-expressed in *Trichoplusia ni* insect cells using the Baculovirus system. CBX8 variably carried an N-terminal 6xHis-mEGFP or a N-terminal 6xHis-MBP tag. The purification protocols were identical regardless of the tag. Baculoviruses were generated as per manufacturers instructions (Thermo Fisher). The viral titre was determined using the MTT assay (Promega #G3580). *Trichoplusia ni* insect cells were infected at a density of 1.5 - 2 x 10^6 cells and incubated for 60 hours at 27 °C in a shaker. The cells were spun down at 1500 relative centrifugal force (RCF). The pellet was resuspended in 100 ml of lysis buffer per litre of cell culture (20 mM HEPES-KOH pH 7.5, 400 mM NaCl, 25 mM Imidazol, 10 % Glycerol, 0.2 mM TCEP, 1 mM PMSF and EDTA-free Complete protease inhibitor (Thermo Fisher)). Lysis proceeded for 30-45 minutes at 4 °C while rotating. The lysate was then centrifuged at 29000 RCF for 20 minutes at 4 °C. The supernatant was transferred to a fresh tube and 5 ml of Ni-NTA resin (Qiagen) was added.

The samples were then incubated for 60 minutes at 4 °C while rotating and subsequently
centrifuged at 500 RCF for 5 minutes at 4 °C to settle the beads. About 90 % of the
supernatant was removed. The beads were resuspended in the remaining 10 % of
supernatant and transferred to 25 mm diameter gravity flow columns (Biorad). The beads
were allowed to settle before the remaining buffer was drained and the beads were then
washed with 12 CV of Buffer B (20 mM HEPES-KOH pH 7.5 at 20 °C, 500 mM NaCl, 25 mM
Imidazole, 10 % Glycerol, 0.2mM TCEP), followed by 30 CV of Buffer A (20 mM HEPES pH 7.5
at 20 °C, 100 mM NaCl, 25 mM Imidazole, 10 % Glycerol, 0.5 mM DTT or 0.2 mM TCEP). The
protein was then eluted in 6 CV Elution Buffer (20 mM HEPES-KOH pH 7.5 at 20 °C, 100 mM
NaCl, 400 mM Imidazole, 10 % Glycerol, 0.2 mM TCEP). The eluted protein was loaded onto
a Hitrap 5 ml Heparin column (Cytvia) equilibrated in IX Buffer A (20 mM HEPES pH 7.5 at
20 °C, 100 mM NaCl, 0.5 mM DDT or 0.2 mM TCEP) and the column was washed with 5 CV of
IX Buffer A. The proteins were resolved over a 20 CV gradient ranging from 0 % to 100 % IX
Buffer B (20 mM HEPES pH7.5 at 20 °C, 1000 mM NaCl, 0.5 mM DTT or 0.2mM TCEP). The
fractions were analysed on SDS-PAGE and fractions containing the protein complex of
interest with the expected subunits stoichiometry were pooled. The pooled fractions were
concentrated using a Amicon ultra 30K centrifugal filter (Merck, cat UFC903024) and
purified by size exclusion chromatography using a HiLoad Sephacryl 300 16/60 column
(Cytiva) equilibrated in GF Buffer (20 mM HEPES-KOH pH 7.5 at 20 °C, 150 mM NaCl, 0.5 mM
DTT). The collected fractions were analysed on SDS-PAGE and fractions containing the
protein complex of interest at the expected stoichiometry were pooled and concentrated to
a concentration of 1-2 mg/ml using an Amicon ultra 30K centrifugal filter (Merck, cat
UFC903024). The purified protein was then aliquoted and frozen in liquid nitrogen. The
purified proteins were stored at -80 °C until use.
364
365 Human UBA1, UBCH5C and Ubiquitin were purified as described previously[44]. Histone
366 proteins H2A, H2B, H2BT120C, H3 and H4 were purified as described previously[45], except
367 that the gel filtration step was omitted.
368
369
370 **Production and purification of DNA for chromatin reconstitution**
371
372 The ATOH1 polycomb target gene was amplified in a large scale 10 ml PCR reaction including
373 500 nM ATOH1 fwd and reverse primers (see Table 1), 4 µg of ATOH1-pUC18 template[41],
374 200 µM dNTP mix (Invitrogen), 3 % DMSO, 50 mM KCl, 10 mM Tris-HCl, pH 8.8, 1.5 mM
375 MgCl2, 0.01% sterile gelatin. The reaction mixture was divided into 96-well plates to include
376 50 µl per well and the following PCR program was run:
377
378

Temperature (°C)	Time	Number of

	(minutes)	Cycles
95	3:00	1
95	0:15	
63	0:20	25
72	8:00	
72	3:00	1

The PCR product was purified via ion exchange chromatography using a 5 ml HiTrap Q
column (GE Healthcare). The column was equilibrated in Buffer A (25 mM HEPES 7.5, 250
mM NaCl) and the sample was resolved using a linear gradient ranging from 0 % to 100 %
Buffer B (25 mM HEPES pH 7.5, 2 M NaCl). The fractions containing pure DNA were pooled,
subjected to ethanol-precipitation and resuspended in 10 mM Tris-HCl pH 7.5 at 20 °C, 0.1
mM EDTA.

Chromatin reconstitution

Histone Octamers were refolded as described previously[45]. All steps were done at 4 °C or on
ice. Chromatin was reconstituted following the salt-gradient dialysis protocol described
previously[45]. For large scale reconstitution, DNA and octamer were combined at an optimal
ratio that was determined at trial experiments for each batch of octamers and DNA. To
determine the optimal ratio of histone octamer to DNA, titration was carried out using
increasing amounts of octamer to a constant amount of DNA (molar ratios of 16:1, 18:1,
20:1, 22:1 of Octamer:DNA) following by salt gradient dialysis. For the salt gradient dialysis,
samples were initially dialysed in a buffer containing 10 mM Tris-HCl pH 7.5 at 20 °C, 2 M
KCl, 1 mM EDTA, 1 mM DTT. The initial buffer was then gradually exchanged for a low salt
buffer containing 10 mM Tris-HCl pH 7.5 at 20 °C, 250 mM KCl, 1 mM EDTA, 1 mM DTT over
the course of 18 hours, after which the salt exchange was complete. Samples were then
centrifuged at 21000 RCF for one minute, separated on a 1% agarose gel in TAE buffer and
stained with SYBR Safe (Sigma). The highest Octamer:DNA ratio at which chromatin
remained soluble after finishing salt dialysis was finally used for large scale reconstitution.
For large scale reconstitutions, the salt gradient dialysis was performed as above while
scaling up the reaction accordingly. Additionally, after conclusion of the salt gradient, the
samples were transferred to 400 ml of low salt buffer (10 mM TrisRIS-HCl pH 7.5 at 20 °C,
250 mM KCl, 1 mM EDTA, 1 mM DTT) and dialysed for another 2 hours. The samples were
finally dialysed against 1 litre of Chromatin Storage Buffer (10 mM TRIS-HCl pH 7.5, 10 mM
KCl) overnight.

Reconstitution of Fluorescently labelled chromatin

To allow site-specific labelling of histone H2B, a cysteine was introduced via site-directed mutagenesis (H2BK120C, as previously described[46]) using mutagenesis primers indicated in Table S1. H2BK120C was labelled with Cyanine5-maleimide (Lumiprobe cat #13080) as described previously[46]. Labelled octamers were refolded as described above. Before chromatin reconstitution, the unlabelled and labelled octamers were combined at a molar ratio of 7:1 (unlabeled:labelled). Chromatin was then reconstituted as described above.

Chromatin condensation assays

Fluorescently labelled proteins and chromatin were protected from light whenever possible. Chromatin condensation assays were done as described previously[17] with some modifications. Assays were done in 384-well plates with #1.5 glass bottoms (MatTek PBK384G-1.5-C). The wells were treated with 1 M NaOH for 1 hour at room temperature, NaOH was removed and wells were washed with copious amounts of MilliQ water (MQ). MQ was removed and 70 ul of 5k mPEG-silane (Sigma #JKA3037-1G, dissolved in 95% EtOH to a final concentration of 25 mg/ml) was added to each well. The plates were sealed and incubated overnight at room temperature. The mPEG-silane was removed, the wells were washed once with 95 % EtOH, then rinsed with copious amounts of MQ and dried completely in the fume hood.

The wells were then passivated by adding 40 ul of 20 mg/ml BSA (NEB) and incubated for at least one hour at room temperature. The BSA was removed and the wells were washed three times with Condensation Buffer (20 mM HEPES pH 7.5 at 20 °C, 0.2 mg/ml BSA, 10 % Glycerol, 5 mM DTT and 150 mM KCl unless stated otherwise). The chromatin stock was adjusted to a DNA concentration of 100 ng/ul and 150 mM KCl. The PRC1 complex was diluted in Condensation buffer to a protein concentration equal to twice the final PRC1 assay concentration as stated. To induce condensation, 16 ul of the diluted PRC1 were combined with 16 ul of the salt-adjusted chromatin dilution in PCR tubes. The samples were incubated for 30 minutes at room temperature before being transferred to the 384-well plate and incubated for a further 60 minutes at room temperature, so that the first images were recorded 90 minutes after induction of condensation.

Images were recorded with a Nikon C1 scanning confocal microscope. GFP was excited with a 488 nm laser, Cy5 was excited with a 561 nm laser. Linear contrast adjustments were made with ImageJ. Where several micrographs are compared to each other, the same contrast settings were used for all micrographs.

Phase contrast images were recorded with a Leica DMI8 imaging system.

Chromatin ubiquitylation assay

The salt concentration of the chromatin stock was adjusted to 100 mM KCl. The nucleosome
equivalent concentration of chromatin arrays was calculated by measuring the molar DNA
concentration and assuming that one DNA molecule is populated by 18 nucleosomes. The
reaction mixture included 750 nM of chromatin (nucleosome equivalent concentration), 500
nM PRC1[C8] or RING1b-BMI1 dimer, 100 nM hUBA1, 500 nM UBCH5C and 50 µM ubiquitin in
Ub-Buffer (25 mM HEPES-KOH pH 7.5 at 20 °C, 100 mM KCl, 3 mM MgCl and 2 mM DTT) and
started by adding ATP to a final concentration of 3 mM. 15 ul reactions were incubated at
30 °C for 45 minutes and the reaction was stopped by adding 5 ul of 4X NuPage LDS-loading
dye (Thermo Scientific cat #NP0008) supplemented with 5 % 2-mercaptoethanol. The
samples were separated on a 4-12% NuPage gel (Thermo Scientific cat #NP0321BOX) using
MES buffer (Thermo Scientific cat #NP0002) in the tank. The gels were then blotted onto a
nitrocellulose membrane (Amersham, cat #GE10600002) in Tris-Glycine transfer buffer +
20 % Ethanol (v/v) for 90 minutes in the cold room at 310 mAmp in a blotting tank (BioRad).
H2A was detected using anti-H2A primary antibodies (Merck Millipore Cat. # 07-146, 1:1000
titer) and HRP-conjugated secondary antibodies (Santa Cruz, cat #sc-2357, titler 1:5000).
466
467   **Sample preparation for cryo-electron tomography**
468
The PRC1[C8] complex was combined with chromatinized ATOH1 at a final concentration of
1.6 µM PRC1[C8] and 500 ng/ul DNA at a final salt concentration of 25 mM NaCl and 8.3 mM
KCl. Samples were incubated at room temperature for 30 minutes. Just before freezing, 5
nm gold nanoparticles were added at a 1:6 ratio (v/v). 3.5 ul of sample was applied to a
Quantifoil grid (R1.2/1.3 on 200 copper mesh, Quantifoil cat #N1-C14nCu20-01) and vitrified
in liquid ethane using the Vitrobot plunge freezer (Thermo Scientific) with the following
settings: Temperature 4°C, blot force -3, blot time 4 seconds, humidity 100%.
476
477   **Cryo-electron tomography data collection and processing**
478
The data was collected with a Titan Krios electron microscope (Thermo Fisher) at 300 keV
acceleration voltage using a Gatan K2 Summit camera (Gatan). A tilt series was acquired
ranging from -60 to 60 degrees with 3 degree increments and a nominal defocus of -2.5 µm.
A dose symmetric collection scheme was followed as described previously[47]. The pixel size
was 1.32 Angstrom, the total dose per tomogram was 144.32 e/A^2. Four frames were
collected per tilt (dose per frame 0.88 e/A^2).
The movies were motion corrected using motioncor2[48]. The motion corrected images were
combined into stacks and further processed with Imod[49] version 4.9.9. Imaging artefacts
were identified and removed with Imod's Ccderaser function. Tilts were aligned using the
gold fiducial markers and the final aligned stack was binned by a factor of 4. The defocus
was estimated using emClarity[50] and the estimated defocus was used in Imod for CTF
correction. The final tomogram was calculated using Imod's implementation of weighted
back projection. For visualisation, the tomogram was denoised using Topaz[51]. The denoised

tomogram was only used for visualisation (Figure 1 and Movie S1). The original non-
denoised tomograms were used for all further processing, including template matching and
subtomogram averaging.
495

Template matching, subtomogram averaging and modelling the chromatin structure of
condensates from cryo-electron tomograms

Two tomograms were selected for further processing (Tomogram #1 and Tomogram #2 in
the following). To avoid user bias, we used a template matching algorithm[52] with the
structure of a single nucleosome as template (EMD-8140[53]) to identify initial positions and
orientations for each nucleosome. The pixel size of the template was adjusted using
emClarity to match the unbinned pixel size of the tomogram[50]. The template was then
subjected to a low-pass filter of 30 Angstrom and binned by a factor of four to match the
binned tomogram. Template matching was done with the Matlab implementation of
Dynamo[52] using the "dynamo_match" function. The template was masked with a tight-
fitting mask with smooth edges. The scanning range was set to 360 degrees with 10 degree
steps. In-plane rotation was also scanned over 360 degrees with 10 degrees steps. Particles
that passed a cross-correlation cut-off of 0.17 (Tomogram #1) or 0.19 (Tomogram #2) were
selected for further analysis. Obvious false positives were removed manually.

The particles were cropped from the tomogram with a box size of 36 pixels. Nucleosome
position and orientation was refined over three rounds of subtomogram averaging[52]. As an
initial template for subtomogram averaging, we used an average from all cropped particles
after template matching, masked with a tight-fitting mask with smooth edges generated in
the Dynamo mask editor. Specific settings for the different rounds of subtomogram
averaging are detailed in Table 2. The resulting average structure of a nucleosome from the
subtomogram averaging was then used to populate a volume the size of the tomogram with
nucleosomes at the determined positions and orientations. The graphic depiction of the
final model (Figure 1G) was generated using the Dynamo Matlab implementation[52].
519

520 Table 2: Settings for subtomogram averaging in Dynamo

	Round 1	Round 2	Round 3
Iterations	3	3	3
References	1	1	1
Cone Aperture	100	16	8
Cone Sampling	5	2	1

Azymuth Rotation Range	100	16	8
Azymuth Rotation Sampling	5	2	1
Refine	0	0	4
Refine Factor	0	0	2
High Pass	2	2	2
Low Pass	7	8	8
Symmetry	c1	c1	c1
Particle Dimension	36	36	36
Shift Limits	5	4	2
Shift Limiting Ways	1	1	1
Seperation in Tomogram	0	0	0
Basic MRA	0	0	0
Threshold Parameter	0.5	0.6	0.6
Threshold Modus	2	2	2

Analysis of exclusion volume

Exclusion volumes for spherical molecules of various radii were calculated using 3V[26] as

follows. First, the positional and rotational coordinates of the nucleosomes within

Tomogram #1 were determined by template matching followed by subtomogram averaging,

as described above. Next, the positional and rotational coordinates of the nucleosomes

were tabulated from within a section of 300 pixels x 300 pixels x 150 pixels (corresponding

to 158.4 nm x 158.4 nm x 87.0 nm at the x, y and z axes, respectively) centred on the pixel at

location (650,650,80), which was large enough to include hundreds of nucleosomes but yet

sufficiently small to carry out the subsequent computational analysis. The dimensions of the
resulting table were converted to Angstrom by multiplying with a tomogram pixel size of
5.28 Angstrom/pixel. Next, the table including nucleosome positions was used by the
Dynamo function 'dtchimera' to generate a chimera cmd script that placed a pdb structure
of a single nucleosome (PDB 1KX4[54]) at the position and orientation defined for each
nucleosome in the table. At the end of this process, each of the nucleosomes within the
tomogram section is represented by the pdb coordinates of the nucleosome template. The
resulting model was saved as a pdb file and used as input for 3V[26]. Varying probe radii
ranging from 2 to 20 nm were used.
541
542 **Hydrodynamic radius calculation using protein structures**
543
The hydrodynamic radii of various proteins (Figure 1I) were calculated using the HullRad[32]
web server (http://52.14.70.9/). The PDB codes for the structures used are: 6LTJ (BAF[37]),
1MUH (Tn5[33]), 7O4J (PolII-PIC[39]), 6C24 (PRC2.2[55]), 8GXS (PolII-Mediator[56]). 6LTJ, 6C24, 8GXS
and 7O4J are only partial models as not all residues were assigned, therefore the calculated
hydrodynamic radius represents an estimate of the minimal complex size, while the actual
size of these complexes could be larger.
550
551 **Analysis of nucleosome-nucleosome orientation**
552
Nucleosome-nucleosome orientation was classified into face-face, face-side and side-side,
as described in[22]. The orientations were calculated from the output table of the template
matching and subtomogram averaging process in Dynamo[52] using the Matlab script
'calculate_orientation.m'. The nucleosome-nucleosome orientation plot (Fig. S1A) was
generated using the 'plot_edges' function from the python notebook
NCP_orientation_analysis.ipyn. Both scripts are available on Github
(https://github.com/MichaUckelmann/Chromatin-Structure-Analysis).
560
561 **Cryo-light microscopy**
562
563 The samples were prepared as described for cryo-electron tomography, without gold
564 nanoparticles and in a buffer containing 100 mM KCl. The vitrified grids were imaged using a
565 ZEISS LSM900 Airyscan2 with a Linkam CMS196V Cryo stage.
566
567 **Fluorescence recovery after photobleaching (FRAP)**
568
Samples and plates were prepared and images were recorded as described above for
chromatin condensation assays. 1 µM PRC1[C8] with an N-terminal GFP tag was used. The
FRAP experiments were set up with the NIS-Elements software (Nikon). Regions of interest
(ROIs) were defined and a single image was recorded before bleaching. Then the ROI was

bleached using 488 nm (GFP) and 561 (Cy5) lasers. The bleaching time and laser power was
set so that approximately 80 % of fluorescence signal within the ROI was quenched.
Recovery was measured over 424 seconds, recording a total of 13 images.
The data was analysed with ImageJ, the mean pixel intensity of the bleached ROI was
quantified for each timepoint.
578
579 **DNA binding assays**
580
DNA binding was assayed using a 24 bp DNA (sequence see Table 1) with Fluorescein
attached to the top strand. The probe was protected from light wherever possible. The
probes were synthesised and delivered as duplexes (IDT). Probes were dissolved at a
concentration of 5 mM DNA in milliQ water. The probes were then diluted to 4 µM in
annealing buffer (20 mM HEPES-KOH, 150 mM NaCl) and heated to 95 °C for 5 minutes. The
probe was then left at room temperature for at least 1 hour to anneal before being diluted
to 20 nM in 20 mM HEPES-KOH pH 7.5 at 20 °C, 150 mM NaCl, 1 mg/ml BSA (NEB cat
#B9000S), 0.1% Tween20, 1 mM DTT. Serial protein dilutions of the PRC1^{C8} and RING1b-
BMI1 complexes were prepared in Protein Dilution Buffer (20 mM HEPES-KOH, 150 mM
NaCl, 1 mM DTT) ranging from 10000 nM to 2.44 nM. 20 ul of probe were mixed with 20 ul
of the respective protein dilution and transferred to a 384-well plate. The samples were
incubated for 30 minutes in the dark at room temperature and then read using a Pherastar
plate reader (BMG Labtech). The fluorescence anisotropy signal was normalised and the
curves were fitted with a specific binding model with Hill slope (GraphPad Prism).
595
596 **Crosslinking mass spectrometry (XL-MS)**
597
Reconstituted chromatin was dialysed overnight against 1 litre of XL buffer (25 mM HEPES-
KOH pH 7.5 at 20 °C, 150 mM NaCl, 1 mM DTT). The chromatin was combined with PRC1^{C8} in
a 1:1.5 molar ratio (Nucleosomes:PRC1^{C8}). Specifically, the nucleosome equivalent
concentration of the chromatin array was calculated from the measured DNA concentration
assuming 18 nucleosomes per DNA molecule. 1.5 µM PRC1 and 1 µM nucleosome
equivalent concentration of chromatin arrays were then combined in XL buffer and
incubated for 30 minutes at room temperature. Crosslinking was done as described
before[57]. The bis(sulfosuccinimidyl)suberate (BS3) crosslinker was added to a final
concentration of 500 µM and crosslinking proceeded for 20 minutes at room temperature at
a reaction volume of 15 µl. The reaction was stopped by the addition of Tris-HCl pH 8 at
20 °C to a final concentration of 125 mM. The samples were then diluted to a volume of
100 µL using a buffer containing 50 mM Tris pH 8 at 20 °C and 150 mM NaCl. TCEP was
added to a final concentration of 10 mM and the samples were incubated at 60 °C for
30 min. Chloroacetamide was added to a final concentration of 40 mM and the samples
were incubated in the dark for 20 min. The samples were then digested using trypsin
(Promega cat #V5280) at 37 °C overnight. The digest was stopped by adding formic acid to a

final concentration of 1% v/v. The digested samples were purified using 100 µl ZipTip pipette
tips (Merck cat #ZTC18S960) according to the manufacturer's instructions. Samples were
then concentrated to ~5 µL using a SpeedVac vacuum centrifuge and diluted with 20 µL
Buffer A (0.1% v/v formic acid).

The peptides were analyzed by online nano-high-pressure liquid chromatography (UHPLC)
electrospray ionization-tandem mass spectrometry (MS/MS) on an Q Exactive Plus
Instrument connected to an Ultimate 3000 UHPLC (Thermo-Fisher Scientific). Peptides
reconstituted in 0.1% formic acid were loaded onto a trap column (Acclaim C18 PepMap 100
nano Trap, 2 cm × 100 µm I.D., 5-µm particle size and 300-Å pore size; Thermo-Fisher
Scientific) at 15 µL/min for 3 min before switching the precolumn in line with the analytical
column (Acclaim C18 PepMap RSLC nanocolumn, 75 µm ID × 50 cm, 3-µm particle size, 100-
Å pore size; Thermo-Fisher Scientific). The separation of peptides was performed at
250 nL/min using a non-linear acetonitrile (ACN) gradient of buffer A (0.1% formic acid) and
buffer B (0.1% formic acid, 80% ACN), starting at 2.5% buffer B to 42.5% over 95 min. Data
were collected in positive mode using a Data Dependent Acquisition m/z of 375–2000 as the
scan range, and higher-energy collisional dissociation (HCD) for MS/MS of the 12 most
intense ions with z 2–5. Other instrument parameters were: MS1 scan at 70,000 resolution,
MS maximum injection time 118 ms, AGC target 3E6, ion intensity threshold of 4.2e4 and
dynamic exclusion set to 15 s. MS/MS resolution of 35000 at Orbitrap with the maximum
injection time of 118 ms, AGC of 5e5 and HCD with collision energy = 27%.
For the data analysis, Thermo raw files were analysed using the pLink 2.3.4 search engine[58]
to identify crosslinked peptides, searching against the sequences of RING1b, BMI1, CBX8,
H2A, H2B, H3 and H4. The default settings for searches were used. N-terminal acetylation
and methionine oxidation were used as variable modifications and carbamidomethyl on
cysteines as a fixed modification. False discovery rates of 1% for peptide spectrum match
level were applied by searching a reverse database. Crosslinked peptides were further
analysed using the crissrosslinkeR package[59]. Specifically, peptides were retained by
crissrosslinkeR only if they passed a p-value cutoff of 0.05 or were present in at least two of
three replicates. Subsequent visualisation of retained peptide was carried out with xiNET[60].

Generation of Cbx8 KO mESC lines using CRISPR/Cas9

Paired sgRNAs were designed to delete exons 1–4 of the murine *Cbx8* gene. The Golden
Gate Cloning method was used to clone the sgRNAs (sequence below) into the lentiguide-
mCherry-Cas9 plasmid[61,62]. 2 million mESCs were transfected with 1 µg of each plasmid
carrying sgRNAs-mCherry-Cas9, using electroporation (Neon™ Transfection System
MPK5000). The following day, mCherry-positive mESCs were sorted by FACS and plated on a
10 cm dish at a very low density for single-cell clone picking. After 5-6 days, individual
colonies (derived from single cells) were picked, expanded, and genotyped using genomic
PCR to identify homozygous/biallelic deletions of Cbx8 KO mESC colonies. Selected Cbx8 KO

mESC lines were further confirmed by western blot for CBX8 (Cell Signalling, CBX8 (D2O8C),

cat # 14696S, titer 1:50 (Fig. 4a)) and HRP-conjugated secondary antibodies (Santa Cruz, cat

#sc-2357, titer 1:10000).

Cbx8 sgRNA sequences (5' and 3' sgRNAs)

CBX8-5'Fw: CACCTGCGAATGCGCCGCTTCAGG

CBX8-5'Rv: AAACCCTGAAGCGGCGCATTCGCA

CBX8-3'Fw: CACCCTCTATGGCCCCAAAAAGCG

CBX8-3'Rv: AAACCGCTTTTTGGGGCCATAGAG

Cbx8 genotyping primers:

Deletion_Fw: GCCTTCTGGTGCAGCTAAGT

Deletion_Rv: GACGTCAGCGGGAGAGTATT

Internal_Fw: CACCAAATGAATGCTCCAAA

Internal_Rv (same as the Deletion_Rv): GACGTCAGCGGGAGAGTATT

Mouse embryonic stem cell culture

Wildtype and *Cbx8* knockout mES cells were grown on gelatinized culture dishes in DMEM,

20 % FBS, 1x non-essential amino acids (Gibco #11140-050), 1x Glutamax (Gibco #35050-

061), PenStrep 100 u/ml (Thermo Fischer), 0.5 x EmbryoMax 2-Mercaptoethanol (Merck

Millipore #ES-007-E), 2.5 µg/ml Plasmocin (Invivogen), 1000 U/ml ESGRO Leukemia

Inhibitory Factor (LIF) (Merck Millipore cat #ESG1107).

Mouse embryonic stem cell differentiation ahead of ChIP-seq and ATAC-seq

Differentiation was induced by seeding cells at a density of $0.3x10^6$ cells per well in 6-well

plates (for ATAC-seq) or at $1.5x10^6$ cells per 10-cm culture dish (for ChIP-seq) in media

containing 1 µM all-trans retinoic acid (RA,Sigma-Aldrich R2625-50MG) and no LIF. Cells

were differentiated for 48 hours and the media was changed after 24 hours. Control cells

were treated with a DMSO volume equivalent to the RA volume in the differentiating cells.

After 48 hours, the cells were harvested, washed once with PBS, counted and used

immediately in either ChIP-seq or ATAC-seq experiments.

ChIP-seq

ChIP-seq was done as described previously[63]. 1.5 µg of CBX8 antibody (Cell Signalling, CBX8

(D2O8C), cat # 14696S) and 3 µg of H3K27me3 antibody (Tri-Methyl-Histone H3 (Lys27)

(C36B11), cat #35861SF) were used. Libraries were prepared using NEBNext ultra II DNA

library kit for Illumina (NEB Biolabs) according to the manufacturer instructions. The

resulting libraries were assessed for quality on a High Sensitivity D1000 Screen Tape
(Agilent) and were sequenced using an Illumina Novaseq 6000 sequencer (Genewiz/Azenta).
696
697 **Data processing for Chip-seq**
698
The reads were quality-trimmed and adapters were removed using Trim Galore!, a wrapper
script to cutadapt[64], in paired end mode and using default settings. The reads were then
aligned to the mouse mm10 genome build using bowtie2[65] with the option "very-sensitive".
The data was reduced to only properly paired reads using "samtools view" with the flag "-f
3". PCR duplicates were removed using the RemoveDuplicates function from Picard Tools.
Read mates were fixed using samtools fixmate. BigWig files were calculated using
BamCoverage with CPM normalisation.
Peaks were called with Macs2[66] callpeak function, using the input sample as control. For the
CBX8 ChIP-seq data set, default settings with a q-value cut-off of 0.05 were used. For
H3K27me3 ChIP-seq data set, broad mode was used with a q-value and broad cutoff of
0.001. Peaks overlapping with ENCODE blacklisted regions[67] were removed.
710
711 **ATAC-seq**
712
713 ATAC-seq was done using a commercial kit (Diagenode Cat.# C01080002) according to the
714 instructions of the manufacturer. The resulting libraries were assessed for quality control on
715 a High Sensitivity D1000 Screen Tape (Agilent) and libraries were sequenced using an
716 Illumina Novaseq 6000 sequencer (Genewiz/Azenta).
717
718 **Data processing ATAC-seq**
719
ATAC-seq data was processed as described previously[68]. Specifically, reads were quality-
trimmed and adapters were removed using Trim Galore!, a wrapper script to cutadapt[64], in
paired end mode using default settings. The reads were then aligned to the mouse mm10
genome build using bowtie2[65] with the option "very-sensitive". Reads were sorted and
indexed using Samtools. Mitochondrial reads were removed using a python script from
Harvard Bioinformatics (available at https://github.com/harvardinformatics/ATAC-seq). The
data was reduced to only properly paired reads using "samtools view" with the flag "-f 3".
The library complexity was estimated and the data sets were subsampled to reach a similar
complexity as described previously[68]. PCR duplicates were removed using the
RemoveDuplicates function from Picard Tools. Read mates were fixed using samtools
fixmate. BigWig files were calculated using BamCoverage with CPM normalisation
For peak calling the bam files were converted to BEDPE files and the Tn5 shift was corrected
by running a bash script provided at https://github.com/reskejak/ATAC-seq
(bedpeTn5Shift.sh). Files were then converted to minimal bed format and peaks were called
using Macs2[66] callpeak function in broad mode with broad-cutoff set to 0.05. Peaks

overlapping with ENCODE blacklisted regions[67] were removed. Consensus peaks for each

condition were defined as the intersect of peaks from both biological replicates. Venn

diagrams were generated using the ChIPPeakAnno package[69].

References

1. Misteli, T. The Self-Organizing Genome: Principles of Genome Architecture and Function. *Cell* **183**, 28–45 (2020).

2. Schuettengruber, B., Bourbon, H.-M., Di Croce, L. & Cavalli, G. Genome Regulation by Polycomb and Trithorax: 70 Years and Counting. *Cell* **171**, 34–57 (2017).

3. Blackledge, N. P., Rose, N. R. & Klose, R. J. Targeting Polycomb systems to regulate gene expression: modifications to a complex story. *Nat Rev Mol Cell Biol* **16**, 643–649 (2015).

4. Eskeland, R. *et al.* Ring1B Compacts Chromatin Structure and Represses Gene Expression Independent of Histone Ubiquitination. *Molecular Cell* **38**, 452–464 (2010).

5. Francis, N. J., Kingston, R. E. & Woodcock, C. L. Chromatin Compaction by a Polycomb Group Protein Complex. *Science* **306**, 1574–1577 (2004).

6. Grau, D. J. *et al.* Compaction of chromatin by diverse Polycomb group proteins requires localized regions of high charge. *Genes Dev.* **25**, 2210–2221 (2011).

7. Lau, M. S. *et al.* Mutation of a nucleosome compaction region disrupts Polycomb-mediated axial patterning. *Science* **355**, 1081–1084 (2017).

8. Isono, K. *et al.* SAM Domain Polymerization Links Subnuclear Clustering of PRC1 to Gene Silencing. *Developmental Cell* **26**, 565–577 (2013).

9. Tatavosian, R. *et al.* Nuclear condensates of the Polycomb protein chromobox 2 (CBX2) assemble through phase separation. *Journal of Biological Chemistry* **294**, 1451–1463 (2019).

10. Plys, A. J. *et al.* Phase separation of Polycomb-repressive complex 1 is governed by a charged disordered region of CBX2. *Genes Dev.* **33**, 799–813 (2019).

11. Illingworth, R. S. Chromatin folding and nuclear architecture: PRC1 function in 3D. *Current Opinion in Genetics & Development* **55**, 82–90 (2019).

12. Miron, E. *et al.* Chromatin arranges in chains of mesoscale domains with nanoscale functional topography independent of cohesin. *Science Advances* **6**, eaba8811 (2020).

13. Nozaki, T. *et al.* Dynamic Organization of Chromatin Domains Revealed by Super-

Resolution Live-Cell Imaging. *Molecular Cell* **67**, 282-293.e7 (2017).

14. Murphy, S. & Boettiger, A. N. Polycomb repression of Hox genes involves spatial

feedback but not domain compaction or demixing. 2022.10.14.512199 Preprint at

https://doi.org/10.1101/2022.10.14.512199 (2022).

15. King, H. W., Fursova, N. A., Blackledge, N. P. & Klose, R. J. Polycomb repressive

complex 1 shapes the nucleosome landscape but not accessibility at target genes.

Genome Res. **28**, 1494–1507 (2018).

16. Chittock, E. C., Latwiel, S., Miller, T. C. R. & Müller, C. W. Molecular architecture of

polycomb repressive complexes. *Biochemical Society Transactions* **45**, 193–205 (2017).

17. Gibson, B. A. *et al.* Organization of Chromatin by Intrinsic and Regulated Phase

Separation. *Cell* **179**, 470-484.e21 (2019).

18. Maeshima, K. *et al.* Nucleosomal arrays self-assemble into supramolecular globular

structures lacking 30-nm fibers. *The EMBO Journal* **35**, 1115–1132 (2016).

19. Zhang, M. *et al.* Molecular organization of the early stages of nucleosome phase

separation visualized by cryo-electron tomography. *Molecular Cell* **82**, 3000-3014.e9

(2022).

20. Strickfaden, H. *et al.* Condensed Chromatin Behaves like a Solid on the Mesoscale In

Vitro and in Living Cells. *Cell* **183**, 1772-1784.e13 (2020).

21. Cai, S., Böck, D., Pilhofer, M. & Gan, L. The in situ structures of mono-, di-, and

trinucleosomes in human heterochromatin. *MBoC* **29**, 2450–2457 (2018).

22. Farr, S. E., Woods, E. J., Joseph, J. A., Garaizar, A. & Collepardo-Guevara, R.

Nucleosome plasticity is a critical element of chromatin liquid–liquid phase separation

and multivalent nucleosome interactions. *Nat Commun* **12**, 2883 (2021).

23. Morey, L. *et al.* Nonoverlapping Functions of the Polycomb Group Cbx Family of

Proteins in Embryonic Stem Cells. *Cell Stem Cell* **10**, 47–62 (2012).

24. Chung, C.-Y. *et al.* Cbx8 Acts Non-canonically with Wdr5 to Promote Mammary

Tumorigenesis. *Cell Reports* **16**, 472–486 (2016).

25. Tan, J. *et al.* CBX8, a Polycomb Group Protein, Is Essential for MLL-AF9-Induced

800   Leukemogenesis. *Cancer Cell* **20**, 563–575 (2011).

801   26. Voss, N. R. & Gerstein, M. 3V: cavity, channel and cleft volume calculator and extractor.

802   *Nucleic Acids Research* **38**, W555–W562 (2010).

803   27. Barbieri, M. *et al.* Complexity of chromatin folding is captured by the strings and binders

804   switch model. *Proceedings of the National Academy of Sciences* **109**, 16173–16178

805   (2012).

806   28. Erdel, F. & Rippe, K. Formation of Chromatin Subcompartments by Phase Separation.

807   *Biophysical Journal* **114**, 2262–2270 (2018).

808   29. Lawrimore, J. *et al.* The rDNA is biomolecular condensate formed by polymer–polymer

809   phase separation and is sequestered in the nucleolus by transcription and R-loops.

810   *Nucleic Acids Research* **49**, 4586–4598 (2021).

811   30. Connelly, K. E. *et al.* Engagement of DNA and H3K27me3 by the CBX8 chromodomain

812   drives chromatin association. *Nucleic Acids Research* **47**, 2289–2305 (2019).

813   31. McGinty, R. K., Henrici, R. C. & Tan, S. Crystal structure of the PRC1 ubiquitylation

814   module bound to the nucleosome. *Nature* **514**, 591–596 (2014).

815   32. Fleming, P. J. & Fleming, K. G. HullRad: Fast Calculations of Folded and Disordered

816   Protein and Nucleic Acid Hydrodynamic Properties. *Biophysical Journal* **114**, 856–869

817   (2018).

818   33. Davies, D. R., Goryshin, I. Y., Reznikoff, W. S. & Rayment, I. Three-Dimensional

819   Structure of the Tn5 Synaptic Complex Transposition Intermediate. *Science* **289**, 77–85

820   (2000).

821   34. Meléndez-Ramírez, C. *et al.* Dynamic landscape of chromatin accessibility and

822   transcriptomic changes during differentiation of human embryonic stem cells into

823   dopaminergic neurons. *Sci Rep* **11**, 16977 (2021).

824   35. Huseyin, M. K. & Klose, R. J. Live-cell single particle tracking of PRC1 reveals a highly

825   dynamic system with low target site occupancy. *Nat Commun* **12**, 887 (2021).

36. Seif, E. *et al.* Phase separation by the polyhomeotic sterile alpha motif

compartmentalizes Polycomb Group proteins and enhances their activity. *Nat Commun*

11, 5609 (2020).

37. He, S. *et al.* Structure of nucleosome-bound human BAF complex. *Science* **367**, 875–

881 (2020).

38. Wang, H. *et al.* Structure of the transcription coactivator SAGA. *Nature* **577**, 717–720

(2020).

39. Schilbach, S., Aibara, S., Dienemann, C., Grabbe, F. & Cramer, P. Structure of RNA

polymerase II pre-initiation complex at 2.9 Å defines initial DNA opening. *Cell* **184**, 4064-

4072.e28 (2021).

40. Maeshima, K. *et al.* The physical size of transcription factors is key to transcriptional

regulation in chromatin domains. *J. Phys.: Condens. Matter* **27**, 064116 (2015).

41. Zhang, Q. *et al.* PALI1 facilitates DNA and nucleosome binding by PRC2 and triggers an

allosteric activation of catalysis. *Nat Commun* **12**, 4592 (2021).

42. Brzovic, P. S., Lissounov, A., Christensen, D. E., Hoyt, D. W. & Klevit, R. E. A

UbcH5/Ubiquitin Noncovalent Complex Is Required for Processive BRCA1-Directed

Ubiquitination. *Molecular Cell* **21**, 873–880 (2006).

43. Pichler, A. *et al.* SUMO modification of the ubiquitin-conjugating enzyme E2-25K. *Nat*

Struct Mol Biol **12**, 264–269 (2005).

44. Uckelmann, M. *et al.* USP48 restrains resection by site-specific cleavage of the BRCA1

ubiquitin mark from H2A. *Nat Commun* **9**, 229 (2018).

45. Luger, K., Rechsteiner, T. J. & Richmond, T. J. Preparation of nucleosome core particle

from recombinant histones. in *Methods in Enzymology* vol. 304 3–19 (Academic Press,

1999).

46. Sabantsev, A., Levendosky, R. F., Zhuang, X., Bowman, G. D. & Deindl, S. Direct

observation of coordinated DNA movements on the nucleosome during chromatin

remodelling. *Nat Commun* **10**, 1720 (2019).

47. Hagen, W. J. H., Wan, W. & Briggs, J. A. G. Implementation of a cryo-electron

tomography tilt-scheme optimized for high resolution subtomogram averaging. *Journal of*

Structural Biology **197**, 191–198 (2017).

48. Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for

improved cryo-electron microscopy. *Nat Methods* **14**, 331–332 (2017).

49. Mastronarde, D. N. Dual-Axis Tomography: An Approach with Alignment Methods That

Preserve Resolution. *Journal of Structural Biology* **120**, 343–352 (1997).

50. Himes, B. A. & Zhang, P. emClarity: software for high-resolution cryo-electron

tomography and subtomogram averaging. *Nat Methods* **15**, 955–961 (2018).

51. Bepler, T., Kelley, K., Noble, A. J. & Berger, B. Topaz-Denoise: general deep denoising

models for cryoEM and cryoET. *Nat Commun* **11**, 5208 (2020).

52. Castaño-Díez, D., Kudryashev, M., Arheit, M. & Stahlberg, H. Dynamo: A flexible, user-

friendly development tool for subtomogram averaging of cryo-EM data in high-

performance computing environments. *Journal of Structural Biology* **178**, 139–151

(2012).

53. Chua, E. Y. D. *et al.* 3.9 Å structure of the nucleosome core particle determined by

phase-plate cryo-EM. *Nucleic Acids Res* **44**, 8013–8019 (2016).

54. Davey, C. A., Sargent, D. F., Luger, K., Maeder, A. W. & Richmond, T. J. Solvent

Mediated Interactions in the Structure of the Nucleosome Core Particle at 1.9Å

Resolution††We dedicate this paper to the memory of Max Perutz who was particularly

inspirational and supportive to T.J.R. in the early stages of this study. *Journal of*

Molecular Biology **319**, 1097–1113 (2002).

55. Kasinath, V. *et al.* Structures of human PRC2 with its cofactors AEBP2 and JARID2.

Science **359**, 940–944 (2018).

56. Chen, X. *et al.* Structures of +1 nucleosome–bound PIC-Mediator complex. *Science* **378**,

62–68 (2022).

57. Horn, V. *et al.* Structural basis of specific H2A K13/K15 ubiquitination by RNF168. *Nat*

Commun **10**, 1751 (2019).

58. Yang, B. *et al.* Identification of cross-linked peptides from complex samples. *Nat*

Methods **9**, 904–906 (2012).

59. Gail, E. H., Shah, A. D., Schittenhelm, R. B. & Davidovich, C. crisscrosslinkeR:

identification and visualization of protein–RNA and protein–protein interactions from

crosslinking mass spectrometry. *Bioinformatics* **36**, 5530–5532 (2020).

60. Combe, C. W., Fischer, L. & Rappsilber, J. xiNET: Cross-link Network Maps With

Residue Resolution*. *Molecular & Cellular Proteomics* **14**, 1137–1147 (2015).

61. Canver, M. C. *et al.* A saturating mutagenesis CRISPR-Cas9–mediated functional

genomic screen identifies cis- and trans-regulatory elements of Oct4 in murine ESCs.

Journal of Biological Chemistry **295**, 15797–15809 (2020).

62. Seruggia, D. *et al.* TAF5L and TAF6L Maintain Self-Renewal of Embryonic Stem Cells

via the MYC Regulatory Network. *Molecular Cell* **74**, 1148-1163.e7 (2019).

63. Healy, E. *et al.* PRC2.1 and PRC2.2 Synergize to Coordinate H3K27 Trimethylation.

Molecular Cell **76**, 437-452.e6 (2019).

64. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17**, 10–12 (2011).

65. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357–359 (2012).

66. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biology* **9**, R137

(2008).

67. Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of

Problematic Regions of the Genome. *Sci Rep* **9**, 9354 (2019).

68. Reske, J. J., Wilson, M. R. & Chandler, R. L. ATAC-seq normalization method can

significantly affect differential accessibility analysis and interpretation. *Epigenetics &*

Chromatin **13**, 22 (2020).

69. Zhu, L. J. *et al.* ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and

ChIP-chip data. *BMC Bioinformatics* **11**, 237 (2010).