Rational engineering of binding pocket's structure and dynamics in penicillin G acylase for selective degradation of bacterial signaling molecules

Michal Grulich[1,‡,*], Bartlomiej Surpeta[2,3,‡], Andrea Palyzová[4], Helena Marešová[4], Jiří Zahradník[5] and Jan Brezovsky[2,3,*]

1. Laboratory of Modulation of Gene Expression, Institute of Microbiology,v.v.i., Academy of Sciences of the Czech Republic, Videnska 1083, 142 20 Prague 4, Czech Republic

2. Laboratory of Biomolecular Interactions and Transport, Department of Gene Expression, Institute of Molecular Biology and Biotechnology, Faculty of Biology, Adam Mickiewicz University, Uniwersytetu Poznanskiego 6, 61-614 Poznan, Poland

3. International Institute of Molecular and Cell Biology in Warsaw, Ks Trojdena 4, 02-109 Warsaw, Poland

4. Laboratory of Molecular Structure Characterization, Institute of Microbiology,v.v.i., Academy of Sciences of the Czech Republic, Videnska 1083, 142 20 Prague 4, Czech Republic

5. Laboratory of Protein Engineering, First Medical Faculty at Biocev, Charles University, Vestec-Prague, Průmyslová 595, 252 50, Czech Republic

‡ These authors contributed equally.

* Email: grulich@biomed.cas.cz (M.G); janbre@amu.edu.pl or jbrezovsky@iimcb.gov.pl (J.B.)

ABSTRACT

The rapid increase in antibiotic-resistant bacteria and the inability to provide new generations of potent antimicrobials necessitates the search for new, unconventional solutions. Methods based on targeting bacterial communication induced by signaling molecules, known as quorum sensing, are gaining increasing interest. Quorum quenching (QQ), as the process of interrupting this communication is called, can be achieved by enzymatic degradation of signaling molecules and represents a promising solution as it limits the expression of genes responsible for virulence, biofilm formation, and drug resistance. It is also believed to circumvent common resistance mechanisms. Therefore, enzymes with QQ activity represent potential next-generation antimicrobial agents for use in medicine, industry, and other areas of life. This work focuses on a biotechnologically optimized penicillin G acylase from *Escherichia coli* (ecPGA), for which primary QQ activity for smaller signaling molecules was recently confirmed. Herein, we introduced triple-point mutations within the binding pocket by an ensemble-based design aimed at modulating the pocket structure and the dynamics of its entrance gates. Next, we proposed a computational workflow to select promising combinations for further modeling. We selected three candidates for experimental evaluation using molecular dynamics simulations of the constructs with six different, biologically relevant signaling molecules. These comprised (i) the VAF variant with enhanced activity towards the medium-sized ligands like the signaling molecule of *Burkholderia cenocepacia*, C08-HSL (N-octanoyl-L-homoserine lactone); (ii) the YAF variant preferring longer substrates like the signaling compound of pathogenic *Vibrio* species, C10-HSL (N-decanoyl-L-homoserine lactone); and finally (iii) the MSF variant with improved efficacy for the longest substrate, C12-3O-HSL (N-3-oxo-dodecanoyl-L-homoserine lactone), the signaling molecule of *Pseudomonas aeruginosa*. In-depth analyses of these engineered variants revealed modulated topology and dynamics of the binding pockets. While we could consistently expand the pockets in these variants, the reactive binding of longer substrates became limited, due to either overly promoted dynamics of the pocket in the VAF variant or an overstabilized pocket in the MSF variant. In summary, we demonstrated the designability of ecPGA for improved QQ and provided insights into the role of adequately modulated pocket dynamics for the activity. Such knowledge, together with the methodology developed for filtering and scoring large datasets of potential variants that reflect the outcomes of our biochemical assays, may provide a suitable toolbox for future exploration and design of tailored QQ acylases toward particular signaling molecules, making them viable antimicrobial agents.

KEYWORDS quorum quenching, protein engineering, rational design, bacterial antibiotics resistance, molecular dynamics,

molecular modeling

INTRODUCTION

Antibiotic-resistant bacteria have become an issue of global priority and will require effective approaches and solutions in the coming years. As estimated in some reports, it may lead to millions of excess deaths per year by 2050.[1] Increasing bacterial resistance derives from several sources. First, it is due to multiple bacterial mechanisms that help them to cope with the antibiotics or drug-like molecules, including inactivation of the compounds, reduced accumulation achieved through efflux pumps or alterations of porins, bacterial target site modification, and finally bacterial persistence due to the formation of biofilms.[2] Those mechanisms are promoted by the direct action of antibiotics on microorganisms, which exerts strong pressure to evolve multiple resistance approaches to survive.[3] It is further promoted by the still widespread misuse of antibiotics in various areas of life to maximize profits, rather than taking care of their sustainable management, keeping them only for critical medical use to protect life.[4] In addition, the widespread unjustified prophylactic prescription of antibiotics without determining the cause of infection constitutes a severe problem, further promoting the progress and spread of bacterial antibiotic resistance.[5] Finally, it was shown that resistance genes could be transferred between the same or even different bacterial species without exposure to the antimicrobial agent, resulting in the widespread of resistant species in all environments, including soil, sea, drinking water, and food.[6–9]

Collaborative efforts on a global and local scale are required to cope with the above-mentioned challenges. While the incorrect prescriptions and misuse of the medication, while challenging, can be gradually reduced by education, appropriate control, and regulations, substituting antibiotics in various fields or overcoming already widespread resistance mechanisms and inefficiency of known antibiotics constitute non-trivial tasks. It is further complicated by the void in discovering new generations of antibiotics, with the last inventions arising nearly 30 years ago.[10,11] Therefore, the constant search for new alternatives is crucial. Such approaches should preferably be able to escape the existing resistance mechanisms and limit the potential development of resistance in the future. Interestingly, one recently studied approach targets a bacterial communication process called quorum sensing (QS). In gram-negative bacteria, this process is primarily induced by N-acyl-homoserine lactones (AHLs) used as signaling molecules for synthase/transcriptional regulator systems. In these systems, the synthase enzyme produces the signaling molecule, while the receptor recognizes the signal and affects the expression of QS-dependent genes, including the synthase.[12] This process enables the collective response of bacteria and controls the expression of genes responsible for virulence, swarming, spreading resistance, or biofilm formation in a cell-density-dependent manner. This communication can be disrupted by the quorum quenching (QQ) process, achieved either by inhibiting the specific synthase involved in the signaling or its receptor, or by enzymatically inactivating the signaling molecule. While inhibiting intracellular proteins may suffer from similar resistance mechanisms as for current antibiotics, signal inactivation constitutes a more promising approach due to its extracellular targeting, markedly reducing the selective pressure.[13] Amongst three enzyme classes that process AHLs, amidases/acylases are of significant interest due to their irreversible hydrolytic action.[14]

Notably, QQ acylases have shown promise in many applications in different fields, including medicine, industry, agriculture, aquaculture, and wastewater treatment among others.[14–19] *Pseudomonas aeruginosa* AHL acylase (paPvdQ), one of the first enzymes described to display QQ activity,[20] was formulated as a dry and stable inhalable powder that was shown to limit the virulence of *P. aeruginosa* infections initially in the *Caenorhabditis elegans* model, followed by the successful trials in the mouse.[21–23] Its mutant was then successfully designed towards shorter AHLs exhibiting operational activity on the *Burkholderia cenocepacia* larvae infection model.[19] Coating medical implants and catheters were another target for QQ enzyme application,[17,24] taking advantage of enzymes' immobilization to improve device stability and reuse potential.[25,26] Furthermore,

bacterial biofilms are the source of biofouling,[27] which causes considerable financial losses in wastewater treatment, industrial processes utilizing membranes, or structures that are in contact with seawater, including ships, boats, fish nets, or pipelines. Prevention of bacterial colonization on surfaces aims at improved and longer permeability of membranes, or avoiding additional and unwanted cleaning of devices and surfaces.[28] To address these drawbacks, immobilized acylase in a wastewater treatment model system incubated with *P. aeruginosa* PAO1 demonstrated the filtration membrane retained its permeability for up to 14 days due to the prevention of biofilm maturation on the membrane, which confirms the high application potential of this biotechnology.[29,30] In terms of the anti-biofouling solutions for the water-contacting equipment, the current solutions are based on special paints and coatings, which frequently contain environmentally unfriendly ingredients such as copper that, although preventing biofouling, are hazardous for the water environments.[16,27] Therefore, QQ enzymes constitute a promising eco-friendly approach without hazardous effects due to their rapid environmental degradation. However, stability might represent a considerable limitation and challenge to overcome.[15,31] Finally, AHL acylases were recently shown to inhibit the spoilage of food caused by bacteria, which represents yet another application area for QQ technologies.[18,32]

Given the above facts, the increased application potential of QQ enzymes is evident in various areas. However, it is worth noting the obstacles that limit their applicability. When considering broad-scale utilization, aspects such as established procedures for mass production, immobilization, high tolerance to industrial processes, and shelf-life are critical. Although numerous enzymes have been shown to exhibit non-negligible QQ activity, not all are suitable for wide-scale adoption. For example, paPvdQ, despite being the most investigated AHL acylase with verified QQ activity, [19,22,23] lacks an optimized procedure for industrial-scale production. Furthermore, PF2571 AHL acylase was shown to possess high activity towards long AHLs, and its variant has been modulated to cleave the shorter substrates efficiently. However, while these enzymes show high potential for preventing food spoilage, the lack of optimized and cost-effective production is currently considered a major application obstacle.[18,32]

Instead of devising processes for improving the applicability of individual QQ enzymes, we propose to focus on engineering the QQ activity of already well-established *Escherichia coli* penicillin G acylase (ecPGA), best known for its role in the synthesis of semi-synthetic β-lactam antibiotics.[33–36] ecPGA represents a perfect template for engineering due to the availability of multiple high-quality crystal structures, broad knowledge of its properties, and multiple protocols for expression and purification, as documented by successful rational engineering cases.[37,38] Previously, we have shown that ecPGA possesses QQ activity towards short- to medium-long AHLs.[39] Moreover, by contrasting its mechanism of action with paPvdQ, we have revealed dynamic determinants that govern the QQ activity and constitute promising targets for engineering the potency of ecPGA towards AHLs. These include (i) residues forming the bottom part of ecPGA's acyl-binding cavity that limit the depth and compromise productive binding of longer substrates, and (ii) the dynamics of molecular gates themselves that control the access to the acyl-binding site.[39] Since bacterial species, including the beneficial microbiota, often utilize distinct AHLs,[13] we believe that relevant QQ solutions for various applications could be devised by engineering these molecular determinants.

In this work, we successfully designed ecPGA variants with improved activity towards varied bacterial signaling molecules using computationally-driven workflow targeting the structure and dynamics of the acyl-binding pocket of this enzyme. This work confirms our ability to customize ecPGA towards tailored QQ specificity. By in-depth analysis of the mutants, we provide molecular details underlying the improved activity and altered specificity. Furthermore, we point out additional limitations in our strategy that can guide subsequent engineering efforts.

MATERIALS AND METHODS

Computational

Preparation of templates for mutagenesis. The three best quality structures of ecPGA were selected from the PDB database (PDB IDs: 1GK9, 1GM7, and 1GM9),[40] and water molecules, and organic and inorganic compounds were removed. Structures were further modified to achieve a wild-type composition; 1GM7 Ala241β was converted to native Asn, and Sme16α was replaced by Met for 1GM7 and 1GM9 structures using PyMOL 2.0.1 (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.) mutagenesis wizard. The *RepairPDB* module of FoldX 4[41] was then applied to improve the quality of the structures, including correcting inappropriate torsion angles and too-close contacts indicated in the structures' validation reports.

***In-silico* site-saturation mutagenesis and filtering.** The *BuildModel* module of FoldX 4 software was used to construct the mutant structures. Each mutagenesis experiment was performed using the module's default settings, namely ionStrength 0.05, pH 7, temperature 298K, vdWDesign 2.0, and was repeated ten times (numberOfRuns 10). The predicted stability of the introduced mutation was averaged from the mutagenesis obtained independently for each of the structures, considering both orders of introducing mutations for two-point mutations. Overall, an ensemble of 60 structures was generated for each mutant (30 x 2) and further evaluated with CAVER 3.02[42] software to calculate the tunnel covering the binding pocket entrance. The starting point for the CAVER calculations was set up as a center of mutated residues, namely Phe138α, Met142α, and Ile177β, with *probe_radius* of 0.9 Å, *shell_radius* of 3 and *shell_depth* set up to 7. The results of the CAVER calculation for each mutant were processed separately by the comparative analysis module of TransportTools 0.9.4 software.[43] Tunnel clusters were joined into superclusters using the average-linkage clustering method with a 4.0 Å cutoff. Filtering was applied to consider only superclusters containing tunnels that occurred in at least two structures per ensemble that were generated for each variant (*min_snapshot_num = 2*). The variants were then evaluated based on the stability estimated by FoldX software and the geometric filtering based on comparative TransportTools analysis. The variables we considered were the number of frames with an open pocket, average length, average and maximal bottleneck radii, and average throughput.

Preparation of protein and ligand structures for molecular docking. At this stage, only the variants we obtained from the best-quality template of the wild-type enzyme (PDB-ID 1GK9) were used for further modeling. Designed and repaired wild-type structures were protonated by H++ web-server[44,45] with default settings, namely pH 7.5, salinity 0.15, and internal and external dielectric constants of 10 and 80, respectively. Protonation of the crucial Ser1β residue was adjusted to resemble the state ready for the first stage of the reaction. Autodock4.2 was used to perform molecular docking experiments to obtain protein-ligand complexes for all designed variants and the wild-type with the following set of AHLs: C06-HSL (N-hexanoyl-L-homoserine lactone), C06-3O-HSL (N-3-oxo-hexanoyl-L-homoserine lactone), C08-HSL, C08-3O-HSL (N-3-oxo-octanoyl-L-homoserine lactone), C10-HSL, and C12-3O-HSL. Structures of ligands were created in Avogadro,[46] and optimized using Gaussian09 software[47] at HF/6-31G* level of theory. Furthermore, charges were derived from the RESP-A1 charge model with a multi-conformational RESP fitting procedure using the RESP ESP web server.[48,49]

Molecular docking calculations. The grid box for docking was defined using the center of mass of Ser1β and Phe138α residues and was adjusted manually using Autodock PyMOL plug-in[50] to cover the entire binding site and catalytic machinery. The box dimension was set up to be 17.5 x 16.25 x 21.25 Å in size. Ligand molecules were treated as flexible during docking, while the receptor was considered rigid except for the mutated residue at position 138α. Receptor and ligand structures were prepared

using *prepare_receptor4.py* and *prepare_ligand4.py* scripts from AutoDockTools4 utilities.[51] Docking procedure was performed using the default settings of the Lamarckian Genetic Algorithm search, which was repeated 250 times for each complex. Representative complex structures were selected based on the favorable binding score from the Autodock4.2 scoring function and were filtered by the criteria defined by proper ligand stabilization for the nucleophile attack reaction.[37,39,52] This criteria included oxyanion hole stabilization interactions (distances between the carbonyl oxygen of the ligand molecule and hydrogens from Ala69β-NH and Asn241β-NδH), nucleophile attack (distance between carbonyl carbon of the ligand and hydroxyl oxygen of Ser1β), and additional stabilization provided by Gln23β (distance between amide hydrogen of the ligand and backbone oxygen Gln23β-O) that is known to play a crucial role in the tetrahedral intermediate stabilization. Complexes with distances satisfying a cutoff of 3.0 Å and favorable binding score were used as inputs for the following molecular dynamics (MD) simulations (Figure 1).

Figure 1. Geometric criteria based on the nucleophilic attack reaction mechanism. Four distances were considered during the selection of the best poses from molecular docking for subsequent simulations. Those included nucleophile attack distance (black dashed line), oxyanion hole stabilization from Ala69β and Asn241β, and additional stabilization from Gln23β (yellow dashed lines). We applied a 3.0 Å cutoff for all listed distances.

Molecular dynamics simulations of protein-ligand complexes. Pre-selected complex structures were merged with the crystallographic waters, and placed in an octahedral box containing TIP3P water molecules with a minimum distance of 10 Å between the solute and the edge of the box. Na+ and Cl- ions were added to the system to neutralize the charge and obtain a salt concentration of approximately 0.1 M. Water molecules that were too close to the solute were removed to avoid potential instabilities in the system during simulation. Parameters and topologies were obtained using the *tleap* module from AmberTools17[53], applying the ff14SB force field.[54] The protonation of the N-terminal serine was designed to resemble the nucleophile-attack ready state (^+H_3N-Ser-O$^-$).[39,55] Charges were obtained analogously to the procedure performed for ligands, and the different atom types were derived analogously to the Cornell et al. force field.[56] Energy minimization and MD simulations were performed using the *pmemd* and *pmemd.CUDA* engines of AMBER 16 package.[53] All complexes were energy minimized in five consecutive stages, each composed of 1000 steps of the steepest descent method followed by 1000 steps of the conjugate gradient method, with gradually decreasing restraints on the protein heavy atoms (initially 500 for heavy atoms, and later restraints applied only on the backbone atoms 500, 125, 25, 0.001 kcal/mol^{-1} Å^{-2}) and ligand (initially 500 for heavy atoms, continued by restraints used only on the amide bond atoms, 50, 25, 5, 0.01 kcal/mol^{-1} Å^{-2}). Equilibration was

performed in 4 stages: (i) 100 ps NVT heating stage from 0 to 100 K using a Langevin thermostat with 5 kcal/mol^{-1}Å^{-2} harmonic restraints on the heavy atoms, followed by three NPT stages using a Langevin thermostat and Monte Carlo barostat: (ii) 300 ps in 100 K with 5 kcal/mol^{-1}Å^{-2} harmonic restraints on the heavy atoms, (iii) 600 ps gradually heating the system to 310 K in the first 200 ps applying harmonic restraints on the backbone heavy atoms using force constant of 5 kcal/mol^{-1}Å^{-2} and distance restraints on the crucial hydrogen bonds stabilizing the ligand (ligand carbonyl carbon to oxyanion hole stabilizing residues Ala69β-NH and Asn241β-NδH) with additional stabilization of the ligand amide hydrogen (by Gln23β-O) using force constant of 25 kcal/mol^{-1} Å^{-2}, and (iv) 1 ns gradually decreasing the distances that restraint was applied in stage (iii) from 25 to 0 kcal/mol^{-1}Å^{-2}. Furthermore, 5 ns NPT production runs were performed using a Langevin thermostat at 310 K, applying a Monte Carlo barostat. All MD simulation stages were run using a 2 fs time-step. The trajectory was generated by saving coordinates every 10 ps, and simulation steps (minimization, equilibration) were performed at least 30 times, with the exception of the production which was continued only for the replicates that maintained the substrate in close proximity to the productive state and properly stabilized by the active site machinery. Otherwise, such replicates were considered non-productive (with zero frames sampling reactive stabilization of the ligand), and terminated after equilibration stage. Root mean square deviation (RMSD) of protein and ligand separately were used to inspect the stability of the system and complex itself during production runs (Supplementary File 2). Crucial distances required for estimating enzymes' capabilities to maintain reactive stabilization of the complex were measured using *cpptraj* from AmberTools17. Heat maps representing the relative change of the potency of each variant were generated using the Seaborn Python library.[57]

Molecular dynamics simulations of free enzymes. Wild-type ecPGA and paPvdQ enzyme structures (PDB ID: 1GK9 and 4M1J[58]), as well as the three best mutants (VAF, YAF, and MSF), were subjected to an additional round of free enzyme MD simulations. Systems were prepared in an analogous way to protein-ligand complexes described in the previous section with three major differences: (i) a lack of ligand in the system, (ii) neutral (resting) N-terminal serine protonation state (H$_2$N-Ser-OH)[39,55] and (iii) hydrogen atom masses were repartitioned[59] to enable 4 fs time-steps during MD simulations. Parameters for this neutral protonation state were obtained from our previous study.[39] The minimization was performed with the *pmemd* module of Amber18,[60] while equilibration and production runs were performed with the *pmemd.CUDA* module of the same package, using analogous settings as for the protein-ligand complexes. All stages were performed in triplicate, and the production simulation was 500 ns long, with the trajectory generated by saving coordinates every 20 ps. The stability of the systems was inspected in terms of RMSD (Figure S1). Crucial distances between catalytically relevant atoms were measured using the *cpptraj* module from AmberTools18. Then, principal component analysis (PCA) was applied to perform dimensionality reduction of this distances network, as implemented in Python scikit-learn library.[61] First, two principal components (PC1 and PC2) were clustered using hdbscan[62] with *min_cluster_size* of 500 and *cluster_selection_epsilon* of 0.001. From these clustering results, members featuring the strongest affiliation to a particular cluster (*probability_* = 1.0) were used to calculate average distances that characterize the cluster representatives, which were further transformed to the original PC space using the PCA model.

Characterization of binding site properties. *Cpptraj* from AmberTools18 was used to generate an ensemble of every tenth snapshot from free enzyme trajectories (2500 snapshots per trajectory) in PDB file format for further geometrical analysis. First, a pocket search was performed using mdpocket[63,64] run (mode 1) with default settings for wild-type and the three shortlisted mutant structures. Then, grid points corresponding to the binding pocket cavity of interest were selected, and the following run (mode 2) was performed to calculate the pocket descriptors. Additionally, mdpocket-generated information

on the most conserved residues that form particular pockets was used to select the starting point for tunnel detection using the CAVER software.[42] Those covered the center of mass of the residues Val138α, Val56β and Trp154β for mutant VAF; Tyr52β, Val56β and Trp154β for mutants YAF and MSF; Met142α, Ser67β and Ile177β for ecPGA wild-type; and Leu146α, Leu53β, and Trp162β for paPvdQ. Tunnel calculations were performed using a probe radius of 0.7 Å. The clustering threshold was 3.5 Å, and one representative was selected for each path. Profiles to visualize binding cavities were generated based on filtered CAVER paths, and we considered only those that corresponded to frames with cavities meeting the following criteria; (i) minimum depth of the cavity greater than 5 Å, (ii) the narrowest point at the entrance to the cavity is greater than 1.4 Å, and (iii) the maximum distance between the desired and actual starting point for tunnel calculations and end point of the tunnel below 2.0 Å.

All plots were generated using matplotlib[65] or seaborn Python libraries. Figures containing structural representations were generated using PyMOL 2.0.1.

Experimental

Chemicals – details on used AHLs. C10-HSL (molecular weight (MW) of 255.4) was obtained from Merck AG, and the remaining AHLs, namely C06-HSL (MW of 199.2), C06-3O-HSL (MW of 213.2), C08-HSL (MW of 227.3), C08-3O-HSL (MW of 241.3), and C12-3O-HSL (MW of 297.4) were purchased from Cayman Chemical, USA and prepared as 10 mM solution dissolved in DMSO and stored in aliquots in -20 °C. For screening biosensor assay, AHLs were dissolved to 300 µM in Luria-Bertani (LB) medium before use.

Bacterial strains and growth conditions. *E. coli* BL21(DE3), *E. coli* TOP10 (for routine cloning) and the control strain RE3(pKA18),[66] were routinely cultivated in LB medium (in g/L: tryptone 10, yeast extract 5, NaCl 10, pH 7.2) supplemented with kanamycin (35 mg/mL) for 24 h at 37°C, 180 rpm. *Chromobacterium violaceum* CV026, the AHL biosensor strain, was used to screen quorum sensing inhibitory activity and was cultured aerobically in LB medium supplemented with kanamycin (25 mg/mL) for 24 h at 28°C, 180 rpm. To produce mutated ecPGAs, recombinant strains were cultivated in 50 mL of mineral MCHGly medium for 24 h at 28°C, 180 rpm, and protein expression was induced with 0.1 mM isopropyl alcohol β-D-thio-galactopyranoside.[67] To prepare solid media for screening or a short-term strain maintenance, LB medium was supplemented with agar (15 g/L) or kanamycin (25 or 35 mg/mL).

Construction of expression system and mutagenesis. The *ecpga* gene was initially cloned in the pET-26b vector using restriction cloning. In brief, the *ecpga* gene (GenBank: X04114.1) was amplified by PCR using Q5 polymerase (NEB, USA) from our previous plasmid pKA18[68] with ecPGA cloning primers bearing *Nde*I and *Eco*RI site overlaps. The purified (NucleoSpin® Gel and PCR Clean-up, Macherey-Nagel, Germany) and digested PCR product by *Nde*I and *Eco*RI restriction enzymes (NEB, USA) was ligated using T4 DNA ligase (NEB, USA) into the pET-26b vector cleaved with the same enzymes. The construct was directly transformed in electrocompetent *E. coli* TOP10 cells. Single colonies were screened by colony PCR and verified by sequencing with sequencing primers. Mutagenesis steps were performed by the implementation of the restriction-free (RF) cloning method.[69] Initially, mega-primers were amplified by PCR reactions (KAPA HiFi DNA Polymerase, Roche, Switzerland) with specific primers bearing mutations and the reverse primer overlapping the integration sites in the destination vector. Gel-purified mega-primers (NucleoSpin®) were incorporated in the destination plasmid pET-26-PGA by second whole-plasmid amplification PCR. The parental vector was digested by the *Dpn*I enzyme (NEB, USA), and the ligation mixture was directly electroporated into *E. coli* TOP10 cells. The successful incorporation of the mega-primers into the destination vector was

verified by colony PCR and sequencing. All primers are listed in Table S1. The constructed plasmids bearing mutated *pga* gene were purified from *E. coli* TOP10 cells and transformed to the *E. coli* BL21(DE3).

Enzyme purification and hydrolytic assay: The three chosen mutants were purified as previously described by Kutzbach and Rauenbusch.[70] The activity of one unit (U) was defined as the amount of ecPGA cleaving 1 μmol of the corresponding AHL per minute in 0.05 M sodium phosphate buffer containing 2 % (w/v) AHL at 35°C and pH 7.0.

Biosensors assay of AHL and ecPGA activity. To screen QQ enzyme activity, cells grown in MCHGly medium with IPTG (8000 g, 5 min, 4°C) were washed by centrifugation with 0.05 M Na phosphate buffer, pH 8.0 and sonicated twice in the same buffer on ice for 30 s. The cell supernatant obtained after centrifugation (13400 rpm, 5 min, 4°C) was used in screening biosensor assays with the *C. violaceum* strains CV026 and 300 μM C06-HSL, C08-HSL and C08-3O-HSL according to Last et al.,[71] i.e., 40 μL of cell supernatant was mixed with 300 μM AHL in LB medium in an Eppendorf tube, and the mixture was cultivated for 1 h at 37°C and 100 rpm. After the biocatalytic reaction, 160 μL of diluted *C. violaceum* CV026 (in the ratio of 1:50) from an overnight culture grown in LB medium was added to the Eppendorf tube. The ecPGA activity assay using the harvested cells was performed according to Sobotková et al.[66]

Effect of temperature on the activity of AHL degradation. The optimal temperature for the activity of individual ecPGA variants was determined by conducting cleavage experiments as described in detail in Surpeta *et al*.[39] for all the AHLs signaling molecules investigated herein.

Enzyme kinetics. Kinetic characterization of AHL degradation by ecPGA and its mutant variants was carried out in 0.05 M phosphate buffer at pH 7.0 at the optimal temperature for ecPGA (35°C), as described previously.[39] Monitoring of reactant concentration was performed at 210 nm using an HPLC apparatus and samples were prepared for HPLC analyses as follows. First, the reaction aliquots were adjusted to pH 2 to terminate the reaction, followed by evaporating to dryness at 35°C. The resulting residues were dissolved in 0.2 mL of acetonitrile. An aliquot of 20 μL of the resulting solution was loaded onto an analytical RP-C18 HPLC column (250 x 4.6, 5 μm particle size, Hypersil ODS). The elution profile proceeded under the following conditions: i) isocratic elution in methanol-water (50:50, v/v) for 10 minutes, ii) linear gradient from 50 to 90 % methanol when in water for over 15 minutes, iii) isocratic elution for 25 minutes. The flow rate for HPLC was set at 0.4 mL/min. The retention times of the substrates of interest are listed in Table S2. The relationship between the initial reaction rate and the substrate concentration (1 - 1000 μM) was determined for each substrate in three independent experiments. The kinetic parameters K_M and V_{max} were calculated using Hans-Volf plots and an ANOVA calculator.

RESULTS AND DISCUSSION

Identifying hot spots in acyl-binding sites of ecPGA to tailor its QQ activities. A recent investigation found insufficient accommodation of moderate-length C08-HSL in the wild-type ecPGA acyl-binding pocket.[39] This, together with arrangements in the paPvdQ enzyme that are selective towards longer AHLs (Figure 2A-B),[20] inspired the exploration of residues whose modification could expand the ecPGA pocket, while improving the stabilization of the bound substrates by altering the structure and dynamics of the entrance to the pocket. Using MD simulations that were generated in our previous work,[39] we identified residues that frequently formed the pocket of wild-type ecPGA (Figure 2C-D, Figure S2) and excluded catalytic residues Ser1β, Gln23β, Ala69β, Asn241β, and gating residues Arg145α, Phe146α and Phe24β (Figure S2). This procedure left us with two hot-spot resides, namely Met142α to deepen the pocket (Figure 2C) and Ile177β to narrow and stabilize the entrance (Figure 2D). Additionally, we considered the third position with the capacity to further elongate the engineered pocket. In this case, we used the binding pose of the transition state analog present in the paPvdQ structure (PDB ID: 4M1J) to estimate the direction of pocket expansion (Figure 2A-C). From the residues forming the target region, Pro49β, Phe57β, Phe138α, and Trp154β, we chose the most mutable residue identified by HotSpot Wizard 3.0 tool,[72,73] namely Phe138α, as the third hot-spot (Figure S2).

Figure 2. Rational design of triple-point ecPGA mutants. A. Side view of the paPvdQ binding pocket profile with transition state analog shown as cyan sticks. B. Entrance to the paPvdQ binding pocket. C. Side view of the ecPGA wild-type binding pocket with paPvdQ transition stage analog shown as cyan sticks superimposed, residues selected for mutagenesis – Phe138α and Met142α – are shown as sticks. D. Entrance to the ecPGA binding pocket, residue Ile177β selected for mutagenesis to narrow the entrance is shown as sticks. The coloring of ecPGA was based on the conservation of contribution to the binding pocket formation from mdpocket analyses (C-D).

Computational saturation mutagenesis and filtering identified six variants with enhanced pocket ensembles. Mutagenesis was performed in a two-step workflow using the FoldX tool (Figure 3A-B). First, we considered mutations of residues Met142α and Ile177β. For Met142α, we allowed 19 substitutions to all standard amino acids. In the case of Ile177β, we limited the mutagenesis to residues bulkier than the native isoleucine without introducing additional hydrogen bonding capacity, reducing the possible substitutions at this position to phenylalanine only. By introducing the mutations in both possible orders, we generated 60 structures for each mutant. Using TransportTools comparative module, we analyzed the pocket conformation in the structural ensembles of mutants and identified a consistently present supercluster that best represented the acyl-binding pocket. By considering the geometric properties of engineered pockets and the predicted effects of mutations on ecPGA stability, we pre-selected only the most promising candidates for the second design step. The criteria for

selection, irrespective of the order of introduced mutations, were as follows: (i) a large-margin destabilization penalty (defined here as average energy change extended by the corresponding standard deviation) per mutation that cannot exceed 2 kcal/mol, (ii) conformational behavior of the engineered pocket from the comparative analysis was improved over the wild-type pocket considering the number of frames with an open pocket, average length, average and maximal bottleneck radii, and average throughput (Table S3). Such a procedure resulted in prioritizing two mutations for the next step – Met142αAla & Ile177βPhe and Met142αSer & Ile177βPhe, which were analyzed together with their reverse combinations (Table S4).

Figure 3. Schematic representation of the *in-silico* mutagenesis procedure. A. ecPGA wild-type pocket with residues selected for mutagenesis shown as blue, purple, and green sticks. B. Step-wise procedure of possible variants generation and filtering based on their predicted stability and geometries of engineered pockets. C. Six best candidates selected for subsequent detailed molecular investigation.

Then, we introduced 19 substitutions instead of the third hot-spot residue, Phe138α, into all generated structures of two double-point mutants and followed an analogous procedure for their evaluation, i.e., stability was estimated with FoldX and pocket ensemble geometry was investigated with TransportTools. The second filtering round consisted of the same maximal destabilization penalty of 2 kcal/mol per mutation, and stricter filters based on TransportTools geometries, requiring significantly improved average length, average and maximal bottleneck radii, and near permanent occurrence (Table S5). Reverse order mutations were already considered when calculating properties from the ensembles, which simplified the filtering process. Such filtering yielded seven three-point variants, including two very similar variants, VAF and VSF, with much higher overall destabilization penalties at over 5 kcal/mol (Table S6). While both displayed geometrical improvements, we decided to retain only the VAF variant with the deepest pocket to avoid the costly processing of two potentially unstable variants and produce a set of the six best candidates for further examination (Table S6, Figure 3C).

Computational evaluation of substrate binding in selected variants predicted three variants with specificity tailored towards AHLs used by different pathogenic bacteria. The six variants selected were used as receptors for molecular docking experiments with a set of signaling molecules of different sizes, being C06-, C06-3O-, C08-, C08-3O-, C10-, and C12-3O-HSL. Following strict filtering based on adherence to the reaction mechanism (Figure 1) and favorable binding score, we selected appropriate and pre-organized complexes for the follow-up MD simulations (Table S7).

These simulations revealed improved binding and stabilization for various AHLs for the designed mutants compared to the wild-type enzyme. Starting from the complexes that exhibited favorable ligand stabilization for the first reaction step, we performed repetitive, 5 ns long MD simulations for each system. Here, we focused on the ability of the variant to maintain reactive stabilization for the nucleophilic attack reaction. Building on top of the known mechanism-based criteria to estimate reactivity,[37,52,74] we predicted potential K_M change by counting the fraction of the simulations that sampled proper AHL stabilization. This was then combined with the information of the average number of frames having properly stabilized AHLs by interactions with the oxyanion hole-stabilizing residues Ala69β and Asn241β, and exhibiting nucleophile attack distance and angle within 3.3 Å and 75-105°, respectively (Table S8, Supplementary File 2). From here on, this metric is referred to as the *'reactive stabilization score'.* By contrasting this score with the stabilization observed for wild-type ecPGA for individual AHLs, we estimated relative activity improvements for each investigated variant (Figure 4). Overall, our analysis indicated that the effects of the mutations that were introduced when contrasted to the wild-type enzyme (i) was primarily neutral or only slightly favorable for all ecPGA variants with shorter substrates (C06-, C06-3O-HSL), (ii) most of the designed mutants presented improved activities for medium-size ligands (C08-, C08-3O-HSL), and (iii) few candidates presented markedly improved activities for the longest C12-3O-HSL, which is also the most problematic one for the wild-type ecPGA with a shallow pocket.[39]

We decided to take three of the most interesting candidates for experimental evaluation. The VAF mutant was chosen due to its unique, narrow specificity towards medium-size AHLs, including C08-HSL, a signaling molecule of pathogenic *Burkholderia cenocepacia*.[19] The YAF mutant was selected for its overall improvement for longer substrates (C08-3O – C12-3O-HSL), in contrast to other mutants, especially the LSF mutant with seemingly similar properties but lacking the improvement for C10-HSL.[75] Finally, the MSF mutant was shortlisted, due to its preference for C12-3O-HSL, the signaling molecule of *P. aeruginosa*, and the preferred substrate of prototypical paPvdQ.[19,20]

Figure 4. Predicted activities based on reactive stabilization score of six pre-selected ecPGA variants with a set of AHLs. Rows correspond to the studied AHLs, and columns to particular ecPGA variants, including the wild-type ecPGA. The coloring scheme was applied according to the reactive stabilization score relative to the ecPGA wild-type (relative reactive stabilization score ranges: decreased: 0 – 0.5, neutral: 0.5 – 15, improved: >15).

Experimental characterization confirmed activity profiles of the three best variants agreed with the predictions. The VAF, YAF and MSF mutants were successfully expressed and subjected to a hydrolytic activity assay. The ability of the ecPGAs variants to degrade AHLs was initially tested by a qualitative bioassay using *C. violaceum* CV026. Different production of violacein by *C. violaceum* CV026 and the loss of ecPGA activity of all mutated constructs compared with the product of the original *pga* gene from *E. coli* RE3(pKA18) cloned into *E. coli* BL21(DE3)pETPGA indicated changes in enzyme activities for various substrates (Table 1 and Supplementary File 3). From the experimental assays, we observed good correspondence between the ranking provided by our computational predictions and experimental K_M values, with a Spearman correlation coefficient of -0.69 (Figure S3), suggesting the developed '*reactive stabilization score*' as a relevant metric to guide rational engineering of PGAs. Our introduced mutations had a primarily neutral effect on enzyme activity with shorter substrates compared to wild-type ecPGA. The only exception was the YAF mutant, which exhibited a significant drop of the catalytic efficiency below 10% of the wild-type, caused both by the decreased k_{cat} and increased K_M. This observation contrasts with the computational prediction of the neutral effect, indicating that the activity drop originates from subsequent reaction stages not investigated in the simulations. Overall, we observed satisfactory agreement between the experimental assay and predictions for medium and long substrates. The YAF mutant performed well with the long substrates, reaching a 3-fold improvement for C08-3O- and C12-3O-HSLs. The VAF mutant exhibited notably improved activities with medium substrates, i.e., a nearly 2- and 5-fold improvement for C08-3O- and C08-HSLs, respectively. Interestingly, the VAF mutant had relatively

low activity for long substrates, similar to the MSF mutant with C10-HSL, which, conversely, performed well for the longer C12-3O-HSL. Intrigued by these tendencies and considering the promising correlation between experiments and predictions for most variants, we investigated the origins of these non-trivial effects in more detail.

Table 1. Experimental results for wild-type and designed ecPGA variants. The data is represented as the mean ± stdev.

AHL/variant	ecPGA	VAF	YAF	MSF
k_{cat} [s^{-1}]				
C06	0.0090 ± 0.0003	0.0096 ± 0.0010	0.0031 ± 0.0002	0.0099 ± 0.0006
C06-3O	0.0100 ± 0.0002	0.0098 ± 0.0006	0.0030 ± 0.0003	0.0090 ± 0.0005
C08	0.0085 ± 0.0003	0.0178 ± 0.0010	0.0147 ± 0.0010	0.0098 ± 0.0009
C08-3O	0.0077 ± 0.0004	0.0091 ± 0.0010	0.0120 ± 0.0010	0.0110 ± 0.0008
C10	0.0045 ± 0.0004	0.0045 ± 0.0004	0.0040 ± 0.0002	0.0044 ± 0.0004
C12-3O	0.0043 ± 0.0002	0.0041 ± 0.0003	0.0055 ± 0.0002	0.0050 ± 0.0006
K$_M$ [mM]				
C06	0.59 ± 0.07	0.52 ± 0.05	2.42 ± 0.15	0.49 ± 0.01
C06-3O	0.51 ± 0.03	0.53 ± 0.03	1.34 ± 0.09	0.62 ± 0.02
C08	0.70 ± 0.04	0.28 ± 0.03	0.38 ± 0.03	0.49 ± 0.01
C08-3O	0.88 ± 0.04	0.57 ± 0.02	0.41 ± 0.04	0.42 ± 0.01
C10	2.98 ± 0.20	2.98 ± 0.20	1.44 ± 0.08	3.01 ± 0.30
C12-3O	2.50 ± 0.10	2.81 ± 0.20	1.06 ± 0.04	1.12 ± 0.20
k_{cat}/K$_M$ [mM^{-1} s^{-1}]				
C06	0.0153 ± 0.0010	0.0185 ± 0.0020	0.0013 ± 0.0006	0.0200 ± 0.0040
C06-3O	0.0196 ± 0.0020	0.0185 ± 0.0040	0.0022 ± 0.0005	0.0146 ± 0.0040
C08	0.0122 ± 0.0020	0.0636 ± 0.0040	0.0387 ± 0.0040	0.0200 ± 0.0030
C08-3O	0.0087 ± 0.0010	0.0160 ± 0.0020	0.0300 ± 0.0010	0.0262 ± 0.0010
C10	0.0015 ± 0.0002	0.0015 ± 0.0003	0.0028 ± 0.0008	0.0015 ± 0.0003
C12-3O	0.0017 ± 0.0003	0.0015 ± 0.0003	0.0052 ± 0.0009	0.0045 ± 0.0005

Engineered variants maintain their ability to open the entrance to the acyl-binding pocket and arrange their catalytic machinery for the productive binding of AHLs. To determine the molecular basis of substrate specificity change upon the introduced mutations, we performed extended free enzyme MD simulations of 0.5 µs in length for three replicates for each variant and contrasted those results with previously produced simulations of ecPGA and paPvdQ.[39] The simulations indicated that engineered variants maintained the overall folds well, confirming the absence of severe destabilizing effects due to the introduced mutations (Figure S1). Furthermore, we inspected the tendency of the catalytic machinery, consisting of Ser1β, Ala69β, Asn241β, and Gln23β, to adopt properly pre-organized conformations for productive stabilization of AHLs. Analogously to our previous work,[39] we measured distances between functional atoms of these residues and performed PCA to characterize conformations adopted by these crucial residues (Figure S4). These residues often adopt the favorable state for productive binding, characterized by the serine hydroxyl oxygen approximately equidistant to the remaining functional (stabilizing) atoms. This state was visited in 59 % and 51 % of YAF and MSF simulations, respectively, even surpassing the 36% occurrence in paPvdQ simulations (Table S9). For mutant VAF, the appropriate state was not dominantly adopted, with approximately 21% occurrence, similar to 24 % occurrence in ecPGA (Table S9). Additionally, we inspected the capabilities of the acyl-binding site opening to track potential undesirable gating events introduced by mutations. From this perspective, all variants exhibited sufficiently frequent openings to enable AHLs binding comparable to ecPGA (Figures S5-S8).

The engineered variants demonstrated uniform expansion of their pockets, although this altered the dynamics of their entrances and the stability of the deepest pocket regions. To evaluate the effect of mutations on the targeted acyl-binding pockets, we compared their behavior between the variants, wild-type ecPGA and paPvdQ. Starting from the average volumes that we found in the simulations, all engineered pockets were successfully expanded over the volume observed for the ecPGA pocket, often reaching or even exceeding the volume of the paPvdQ pocket (Figure 5A). By inspecting the details of the geometrical profiles of the pockets (Figure 5B and Figure S9), we observed increased pocket depth in all variants, almost reaching the depth of pocket in paPvdQ, and approximately twice as deep as the ecPGA wild-type. Interestingly, the VAF mutant pocket presented the least stable profile, particularly at the bottom portion (the depth of 15-20 Å). Here, structural fluctuations resulting in the opening of extra space were regularly observed, while well-defined and stable pocket bottoms were present in the other two mutants (Figure 5C). To understand the potential source of the differential pocket stability, we measured the distances between the three pairs of mutated residues in all variants (Figure S10). This data displays different behavior of mutated sites, i.e., mutants YAF and MSF present analogous behavior, where the residues at the 138α and 142α positions formed the closest pair, followed by the 142α-177β pair, and finally the 138α-177β pair. Conversely, in the mutant VAF, we observed two essential differences, namely (i) separation between the 138α-142α pair was increased, similarly to wild-type, reaching the ranges seen for the 142α-177β pair, and (ii) the distances between both the 138α-142α and 138α-177β pairs were less stable, often adopting much larger distances. This more dynamic cross-talk between substituted residues correlates with the observed increased fluctuation of the binding pocket volume for this mutant (Figure 5A) and more considerable deviation in the bottom part of the pocket, as illustrated by the profile analysis (Figure 5B).

Additionally, the initial part of the pocket profile in the engineered variants provided intriguing observations (Figure 5B). Here, we noticed a constriction located below the 5 Å depth that was not present in wild types. Since the constricted area was within the proximity of the mutated site (177β), we analyzed the effect of this perturbation on the conformational dynamics of gating residues Phe24β and Phe146α. The wild-type ecPGA is known to frequently visit two main states of Phe146α (♥ and ♦) while occasionally also adopting an alternative conformation of Phe24β (♣).[39] Interestingly, we observed markedly restricted dynamics of the gating residues in all three variants (Figure 5D), predominantly displaying the single conformation of Phe146α (♥). While the alternative conformation of Phe24β (♣) was rarely visited in the case of VAF and YAF variants, The MSF variant did not display this conformation. To understand how the Ile177βPhe substitution affects the behavior of the gate, we inspected the distribution of spatial separation between the center of mass of the 177β residue and the two gating residues (Figure S11). The most striking difference was the limited range of the distances between gating Phe146α and the introduced Phe177β observed in all mutants when compared to the wild-type ecPGA, where the separation between these two residues reached over 10 Å. Curiously, the restricted mobility of this pair was less pronounced in the VAF mutant, in which the distribution of the respective distances was broader and exhibited bimodal behavior. The increased mobility of the entrance and the initial part of the pocket in this mutant is in agreement with the increased fluctuations detected for this mutant in the pocket profile, volume and residues forming the bottom part of the cavity, rendering this pocket the most dynamic and unstable among the evaluated variants.

Figure 5. Detailed investigation of structure and dynamics of engineered binding pockets. A. Average binding pocket volumes calculated from free enzyme MD simulations using mdpocket. B. Profiles of the binding pocket in the free enzyme MD simulations, based on CAVER tunnel calculations analyzed from properly pre-organized structures. C. Isomesh represents the open voids that occurred in at least 50 % of analyzed frames, with mutated residues shown as sticks. D. Modulated dynamics of the gating residues of designed mutants compared to wild-type ecPGA enzyme. Graphs show conformations sampled by the gating residues as maps of chi1 angles of Phe24β versus Phe146α. Structures located in frames around the maps show (sticks) representative conformations for each state.

Suboptimal dynamics of the enlarged pockets prevent productive stabilization of longer AHLs in VAF and MSF mutants.

One of the most intriguing discrepancies between the structural analysis of the investigated variants and their preferences for various AHLs is the minimal activity of the VAF mutant that was observed with C10- and C12-3O-HSLs, despite its most extensive and flexible binding pocket. Our previous analyses of the ecPGA-C08-HSL complex identified the acyl-binding pocket as being too shallow, causing suboptimal stabilization of this substrate by the oxyanion hole residues. The increased distance of C08-HSL to Ala69β (from 2.0 to 2.5 Å) was found to be a relevant indicator reflecting the increased tendency of the substrate to leave the binding site.[39] Hence, we analyzed these distances in our simulations to evaluate the proper stabilization of AHLs in all variants (Table S10). In line with our expectation, the VAF mutant exhibited markedly longer distances to the oxyanion-hole stabilizing Ala69β with both of the longer substrates, reaching 2.9 ± 0.1 and 2.6 ± 0.3 Å for C10- and C12-3O-HSL, respectively. These distances were much higher when compared to the remaining active complexes. Unlike in our previous study with ecPGA-C08-HSL, the instability of longer AHLs was not caused by a problematic fit into a too-shallow pocket, but instead by the increased dynamics of the pocket.

Additionally, we observed a similar increase in the Ala69β distance for these AHLs with the MSF mutant (Table S10). The pocket of this mutant remained stable following the introduction of Ser142α, as this residue made frequent hydrogen bonds with Tyr52β, stabilizing the bottom section of the pocket. We envision two possible mechanisms behind the enhanced expulsion of longer AHLs from the binding pocket of MSF mutant, being i) the stability introduced into the pocket could limit the range of adjustments necessary to accommodate such AHLs, as corroborated by the conformational changes observed at the bottom of the paPvdQ pocket upon C12-HSL binding,[20] and ii) such an arrangement results into the exposure of multiple polar groups inside the pocket, decreasing complementarity with the hydrophobic tails of the bound C10- and C12-3O-HSL. Surprisingly, these effects translated to the less frequent formation of a reactive binding pose and lower activity for only the C10-HSL, while the combined inference into C12-3O-HSL showed improved catalytic performance, which is quite controversial considering that the later AHL is considerably longer. Attempting an explanation of this inconsistency, we realized that although both C10- and C12-3O-HSL represent rather long substrates, they also differ by the presence of a 3-oxo substituent. Thus, we analyzed hydrogen bonding interactions formed with this group. Aside from rare interactions with Ser1β and Gln23β, the most frequent interaction of the 3-oxo group was noted with the amide group of Ala69β (Table S11). When inspecting the occurrence of these hydrogen bonding events and the reactive complexes, we noted the recurring recovery of reactive poses after the hydrogen bonds are formed with AHLs carrying the 3-oxo substituent (Supplementary File 4 and Figure S12). This observation indicates that although the binding pocket carrying Ser142α is not optimal for the longer substrates, the 3-oxo substitution in C12-3O-HSL, absent in C10-HSL, partially offsets the repulsion, helping the substrate stay in the vicinity of the catalytic machinery and promoting the restoration of the productive complex. This is further supported by analogous behavior observed in simulations of the LSF mutant carrying the same mutation (Figure 4).

CONCLUSIONS

QQ enzymes have gained significant interest as promising antimicrobial agents in the last decade, due mainly to their environmentally friendly character, potential to escape common resistance mechanisms, and their application potential in medicine, various fields of industry, biotechnology, and agriculture.[14,15] Furthermore, the current antibiotic discovery void accentuates the need to investigate antibiotic alternatives to effectively fight bacterial infections.[1,10] Several studies have previously investigated engineered QQ acylases, examining improvements in QQ potency or modified substrate specificity, includ-

ing PvdQ acylase,[19] MacQ bifunctional acylase,[76] and PF2571 AHL acylase[32]. Furthermore, significant efforts have been invested in developing and improving methods for the efficient engineering of QQ enzymes.[77,78] However, to our knowledge, most of the previous systems are not yet fully developed for efficient and sustainable applications on a broader scale.

To address this issue, and building on top of our previous findings of QQ activity of industrially potent penicillin G acylases towards short- to medium-size AHLs,[39] we engineered ecPGA specificity towards a group of AHLs of different sizes. We developed a workflow for the computational design of pocket ensembles, facilitated by the recently developed TransportTools library[43] that enabled us to compare differences in pocket geometries across the large dataset in a unified and reproducible manner. We used this workflow to introduce triple-point mutations in ecPGA's binding pocket to alter its ability to accommodate AHLs while deprioritizing mutants with increased destabilization. Furthermore, the reactive stabilization score that we developed to select candidates from MD simulations for experimental evaluation displayed an appreciable correlation to the results of biochemical assays, suggesting its ability to screen for analogous QQ enzymes.

Detailed analyses of three experimentally constructed mutants exposed changes in the structure-dynamic-function relationships that are relevant to the specificity of these enzymes towards AHLs of different lengths. The VAF mutant, which lacked enhanced activity with longer AHLs and presented the highest instability and plasticity of the binding site, was also the variant with the highest predicted destabilization energy in the initial design despite its viable expression *in vivo*. This highlighted the multifacet consequences of destabilizing effects during rational engineering campaigns. Conversely, more stable YAF and MSF mutants featured stable binding pockets capable of proper stabilization of longer AHLs, although we also identified drawbacks of potential overstabilization of the pocket in variants carrying Met142αSer mutation. Furthermore, we detected modulated dynamics of the Phe146α and Phe24β gating residues compared to wild-type ecPGA, tracking the source of this perturbation to the introduction of the Phe177β residue, which interacts more closely with the gating residue Phe146α, restricting its conformational freedom. Although limiting the dynamics of the gate and driving it closer to more stable gates observed in the prototypical paPvdQ,[39] the perturbation had not narrowed down the entrance as designed. Instead, the entrance remained as open as that seen in the wild-type, whereas the subsequent part of the pocket became more constricted (Figures 3C and 5B). We hypothesize that this constriction may instead compromise the efficient binding of the AHLs that need to adopt strained conformations to fit such a curved pocket. Although revealing unfavorable effects that were introduced with the Ile177βPhe substitution, this uncovered an additional hot-spot contribution to the dynamic control of gating in ecPGA, opening new opportunities for modulation of QQ activity and specificity. When all our data is taken into account, we have demonstrated the ability to design industrially proficient ecPGA towards enhanced QQ activity, identifying several relevant targets enabling further engineering of ecPGA as a practical antibacterial agent.

AUTHOR INFORMATION

Corresponding Author

* Email: grulich@biomed.cas.cz (M.G); janbre@amu.edu.pl or jbrezovsky@iimcb.gov.pl (J.B.)

Author Contributions

M.G.‡ designed and coordinated the experimental work, performed detailed characterization of all the ecPGA designed mutants (YAF, VAF and MSF), and analyzed and interpreted data. B.S.‡ designed ecPGA variants and performed all computational work, analyzed and interpreted data. H.M. and A.P. executed all of the bacterial flasks and fed-batch cultivation in a bioreactor to produce enzymes. J.Z. designed the set of used primers for the ecPGA mutant variants. J.B. conceived and coordinated the project, designed computational analyses, and analyzed and interpreted data. The manuscript was written through the contributions of all authors. All authors have approved the final version of the manuscript.
‡These authors contributed equally.

Funding Sources

This work was supported by the National Science Centre, Poland (Grant Number 2021/41/N/NZ2/01365), Institutional Research project RVO61388971 from the Institute of Microbiology of the CAS and by the Czech Foundation grant 21-17044S. B.S. is a recipient of a scholarship provided by POWER project POWR.03.02.00-00-I022/16. The computations were performed at the Poznan Supercomputing and Networking Center.

ABBREVIATIONS

QQ, quorum quenching; HSL, homoserine lactone; ecPGA, Escherichia coli penicillin G acylase; QS, quorum sensing; AHLs, N-acyl-homoserine lactones; paPvdQ, Pseudomonas aeruginosa acyl-homoserine lactone acylase; C06-HSL, N-hexanoyl-L-homoserine lactone; C06-3O-HSL, N-3-oxo-hexanoyl-L-homoserine lactone; C08-HSL, N-octanoyl-L-homoserine lactone; C08-3O-HSL, N-3-oxo-octanoyl-L-homoserine lactone; C10-HSL, N-decanoyl-L-homoserine lactone; C12-3O-HSL, N-3-oxo-dodecanoyl-L-homoserine lactone; MD, molecular dynamics; RMSD, root mean square deviation; PCA, principal component analysis; PC1, principal component; PC1, principal component 1; PC2, principal component 2; mw, molecular weight; LB, Luria-Bertani; RF, restriction-free cloning method; LAF, Phe138αLeu & Met142αAla & Ile177βPhe ecPGA variant; MAF, Phe138αMet & Met142αAla & Ile177βPhe ecPGA variant; VAF, Phe138αVal & Met142αAla & Ile177βPhe ecPGA variant; YAF, Phe138αTyr & Met142αAla & Ile177βPhe ecPGA variant; LSF, Phe138αLeu & Met142αSer & Ile177βPhe ecPGA variant; MSF, Phe138αMet & Met142αSer & Ile177βPhe ecPGA variant;

REFERENCES

(1)   Murray, C. J.; Ikuta, K. S.; Sharara, F.; Swetschinski, L.; Aguilar, G. R.; Gray, A.; Han, C.; Bisignano, C.; Rao, P.; Wool, E.; Johnson, S. C.; Browne, A. J.; Chipeta, M. G.; Fell, F.; Hackett, S.; Haines-Woodhouse, G.; Hamadani, B. H. K.; Kumaran, E. A. P.; McManigal, B.; Agarwal, R.; Akech, S.; Albertson, S.; Amuasi, J.; Andrews, J.; Aravkin, A.; Ashley, E.; Bailey, F.; Baker, S.; Basnyat, B.; Bekker, A.; Bender, R.; Bethou, A.; Bielicki, J.; Boonkasidecha, S.; Bukosia, J.; Carvalheiro, C.; Castañeda-Orjuela, C.; Chansamouth, V.; Chaurasia, S.; Chiurchiù, S.; Chowdhury, F.; Cook, A. J.; Cooper, B.; Cressey, T. R.; Criollo-Mora, E.; Cunningham, M.; Darboe, S.; Day, N. P. J.; Luca, M. D.; Dokova, K.; Dramowski, A.; Dunachie, S. J.; Eckmanns, T.; Eibach, D.; Emami, A.; Feasey, N.; Fisher-Pearson, N.; Forrest, K.; Garrett, D.; Gastmeier, P.; Giref, A. Z.; Greer, R. C.; Gupta, V.; Haller, S.; Haselbeck, A.; Hay, S. I.; Holm, M.; Hopkins, S.; Iregbu, K. C.; Jacobs, J.; Jarovsky, D.; Javanmardi, F.; Khorana, M.; Kissoon, N.; Kobeissi, E.; Kostyanev, T.; Krapp, F.; Krumkamp, R.; Kumar, A.; Kyu, H. H.; Lim, C.; Limmathurotsakul, D.; Loftus, M. J.; Lunn, M.; Ma, J.; Mturi, N.; Munera-Huertas, T.; Musicha, P.; Mussi-Pinhata, M. M.; Nakamura, T.; Nanavati, R.; Nangia, S.; Newton, P.; Ngoun, C.; Novotney, A.; Nwakanma, D.; Obiero, C. W.; Olivas-Martinez, A.; Olliaro, P.; Ooko, E.; Ortiz-Brizuela, E.; Peleg, A. Y.; Perrone, C.; Plakkal, N.; Ponce-de-Leon, A.; Raad, M.; Ramdin, T.; Riddell, A.; Roberts, T.; Robotham, J. V.; Roca, A.; Rudd, K. E.; Russell, N.; Schnall, J.; Scott, J. A. G.; Shivamallappa, M.; Sifuentes-Osornio, J.; Steenkeste, N.; Stewardson, A. J.; Stoeva, T.; Tasak, N.; Thaiprakong, A.; Thwaites, G.; Turner, C.; Turner, P.; Doorn, H. R. van; Velaphi, S.; Vongpradith, A.; Vu, H.; Walsh, T.; Waner, S.; Wangrangsimakul, T.; Wozniak, T.; Zheng, P.; Sartorius, B.; Lopez, A. D.; Stergachis, A.; Moore, C.; Dolecek, C.; Naghavi, M. Global Burden of Bacterial Antimicrobial Resistance in 2019: A Systematic Analysis. *The Lancet* **2022**, *399* (10325), 629–655. https://doi.org/10.1016/S0140-6736(21)02724-0.

(2)   De Oliveira, D. M. P.; Forde, B. M.; Kidd, T. J.; Harris, P. N. A.; Schembri, M. A.; Beatson, S. A.; Paterson, D. L.; Walker, M. J. Antimicrobial Resistance in ESKAPE Pathogens. *Clinical Microbiology Reviews* **2020**, *33* (3), e00181-19. https://doi.org/10.1128/CMR.00181-19.

(3)   Defoirdt, T. Quorum-Sensing Systems as Targets for Antivirulence Therapy. *Trends in Microbiology* **2018**, *26* (4), 313–328. https://doi.org/10.1016/j.tim.2017.10.005.

(4)   Berendonk, T. U.; Manaia, C. M.; Merlin, C.; Fatta-Kassinos, D.; Cytryn, E.; Walsh, F.; Bürgmann, H.; Sørum, H.; Norström, M.; Pons, M.-N.; Kreuzinger, N.; Huovinen, P.; Stefani, S.; Schwartz, T.; Kisand, V.; Baquero, F.; Martinez, J. L. Tackling Antibiotic Resistance: The Environmental Framework. *Nature Reviews Microbiology* **2015**, *13* (5), 310–317. https://doi.org/10.1038/nrmicro3439.

(5)   Roope, L. S. J.; Smith, R. D.; Pouwels, K. B.; Buchanan, J.; Abel, L.; Eibich, P.; Butler, C. C.; Tan, P. S.; Walker, A. S.; Robotham, J. V.; Wordsworth, S. The Challenge of Antimicrobial Resistance: What Economics Can Contribute. *Science* **2019**, *364* (6435), eaau4679. https://doi.org/10.1126/science.aau4679.

(6)   Wright, G. D. Antibiotic Resistance in the Environment: A Link to the Clinic? *Current Opinion in Microbiology* **2010**, *13* (5), 589–594. https://doi.org/10.1016/j.mib.2010.08.005.

(7)   Vaz-Moreira, I.; Nunes, O. C.; Manaia, C. M. Bacterial Diversity and Antibiotic Resistance in Water Habitats: Searching the Links with the Human Microbiome. *FEMS Microbiology Reviews* **2014**, *38* (4), 761–778. https://doi.org/10.1111/1574-6976.12062.

(8)   Baquero, F.; Martínez, J.-L.; Cantón, R. Antibiotics and Antibiotic Resistance in Water Environments. *Current Opinion in Biotechnology* **2008**, *19* (3), 260–265. https://doi.org/10.1016/j.copbio.2008.05.006.

(9)   Allen, H. K.; Donato, J.; Wang, H. H.; Cloud-Hansen, K. A.; Davies, J.; Handelsman, J. Call of the Wild: Antibiotic Resistance Genes in Natural Environments. *Nature Reviews Microbiology* **2010**, *8* (4), 251–259. https://doi.org/10.1038/nrmicro2312.

(10)   Silver, L. L. Challenges of Antibacterial Discovery. *Clin Microbiol Rev* **2011**, *24* (1), 71–109. https://doi.org/10.1128/CMR.00030-10.

(11)   Boyd, N. K.; Teng, C.; Frei, C. R. Brief Overview of Approaches and Challenges in New Antibiotic Development: A Focus On Drug Repurposing. *Frontiers in Cellular and Infection Microbiology* **2021**, *11*.

(12)   Utari, P. D.; Vogel, J.; Quax, W. J. Deciphering Physiological Functions of AHL Quorum Quenching Acylases. *Front. Microbiol.* **2017**, *8*, 1123. https://doi.org/10.3389/fmicb.2017.01123.

(13)   Grandclément, C.; Tannières, M.; Moréra, S.; Dessaux, Y.; Faure, D. Quorum Quenching: Role in Nature and Applied Developments. *FEMS Microbiol Rev* **2016**, *40* (1), 86–116. https://doi.org/10.1093/femsre/fuv038.

(14)   Fetzner, S. Quorum Quenching Enzymes. *Journal of Biotechnology* **2015**, *201*, 2–14. https://doi.org/10.1016/j.jbiotec.2014.09.001.

(15)   Bzdrenga, J.; Daudé, D.; Rémy, B.; Jacquet, P.; Plener, L.; Elias, M.; Chabrière, E. Biotechnological Applications of Quorum Quenching Enzymes. *Chemico-Biological Interactions* **2017**, *267*, 104–115. https://doi.org/10.1016/j.cbi.2016.05.028.

(16)   Ciriminna, R.; Bright, F. V.; Pagliaro, M. Ecofriendly Antifouling Marine Coatings. *ACS Sustainable Chem. Eng.* **2015**, *3* (4), 559–565. https://doi.org/10.1021/sc500845n.

(17)   Ivanova, K.; Fernandes, M. M.; Mendoza, E.; Tzanov, T. Enzyme Multilayer Coatings Inhibit Pseudomonas Aeruginosa Biofilm Formation on Urinary Catheters. *Appl Microbiol Biotechnol* **2015**, *99* (10), 4373–4385. https://doi.org/10.1007/s00253-015-6378-7.

(18)   Wang, D.; Chen, H.; Li, J.; Li, T.; Ren, L.; Liu, J.; Shen, Y. Screening and Validation of Quorum Quenching Enzyme PF2571 from Pseudomonas Fluorescens Strain PF08 to Inhibit the Spoilage of Red Sea Bream Filets. *International Journal of Food Microbiology* **2022**, *362*, 109476. https://doi.org/10.1016/j.ijfoodmicro.2021.109476.

(19)   Koch, G.; Nadal-Jimenez, P.; Reis, C. R.; Muntendam, R.; Bokhove, M.; Melillo, E.; Dijkstra, B. W.; Cool, R. H.; Quax, W. J. Reducing Virulence of the Human Pathogen Burkholderia by Altering the Substrate Specificity of the Quorum-Quenching Acylase PvdQ. *Proceedings of the National Academy of Sciences* **2014**, *111* (4), 1568–1573. https://doi.org/10.1073/pnas.1311263111.

(20)   Bokhove, M.; Jimenez, P. N.; Quax, W. J.; Dijkstra, B. W. The Quorum-Quenching N-Acyl Homoserine Lactone Acylase PvdQ Is an Ntn-Hydrolase with an Unusual Substrate-Binding Pocket. *Proceedings of the National Academy of Sciences* **2010**, *107* (2), 686–691. https://doi.org/10.1073/pnas.0911839107.

(21)   Wahjudi, M.; Murugappan, S.; van Merkerk, R.; Eissens, A. C.; Visser, M. R.; Hinrichs, W. L. J.; Quax, W. J. Development of a Dry, Stable and Inhalable Acyl–Homoserine–Lactone–Acylase Powder Formulation for the Treatment of Pulmonary Pseudomonas Aeruginosa Infections. *European Journal of Pharmaceutical Sciences* **2013**, *48* (4–5), 637–643. https://doi.org/10.1016/j.ejps.2012.12.015.

(22)   Papaioannou, E.; Wahjudi, M.; Nadal-Jimenez, P.; Koch, G.; Setroikromo, R.; Quax, W. J. Quorum-Quenching Acylase Reduces the Virulence of Pseudomonas Aeruginosa in a Caenorhabditis Elegans Infection Model. *Antimicrobial Agents and Chemotherapy* **2009**, *53* (11), 4891–4897. https://doi.org/10.1128/AAC.00380-09.

(23)   Utari, P. D.; Setroikromo, R.; Melgert, B. N.; Quax, W. J. PvdQ Quorum Quenching Acylase Attenuates Pseudomonas Aeruginosa Virulence in a Mouse Model of Pulmonary Infection. *Front Cell Infect Microbiol* **2018**, *8*, 119. https://doi.org/10.3389/fcimb.2018.00119.

(24)   Ivanova, K.; Fernandes, M. M.; Francesko, A.; Mendoza, E.; Guezguez, J.; Burnet, M.; Tzanov, T. Quorum-Quenching and Matrix-Degrading Enzymes in Multilayer Coatings Synergistically Prevent Bacterial Biofilm Formation on Urinary Catheters. *ACS Appl. Mater. Interfaces* **2015**, *7* (49), 27066–27077. https://doi.org/10.1021/acsami.5b09489.

(25)   Beladiya, C.; Tripathy, R. K.; Bajaj, P.; Aggarwal, G.; Pande, A. H. Expression, Purification and Immobilization of Recombinant AiiA Enzyme onto Magnetic Nanoparticles. *Protein Expression and Purification* **2015**, *113*, 56–62. https://doi.org/10.1016/j.pep.2015.04.014.

(26)   Lee, B.; Yeon, K.-M.; Shim, J.; Kim, S.-R.; Lee, C.-H.; Lee, J.; Kim, J. Effective Antifouling Using Quorum-Quenching Acylase Stabilized in Magnetically-Separable Mesoporous Silica. *Biomacromolecules* **2014**, *15* (4), 1153–1159. https://doi.org/10.1021/bm401595q.

(27)   Cordeiro, A. L.; Werner, C. Enzymes for Antifouling Strategies. *Journal of Adhesion Science and Technology* **2011**, *25* (17), 2317–2344. https://doi.org/10.1163/016942411X574961.

(28)   Yeon, K.-M.; Cheong, W.-S.; Oh, H.-S.; Lee, W.-N.; Hwang, B.-K.; Lee, C.-H.; Beyenal, H.; Lewandowski, Z. Quorum Sensing: A New Biofouling Control Paradigm in a Membrane Bioreactor for Advanced Wastewater Treatment. *Environ. Sci. Technol.* **2009**, *43* (2), 380–385. https://doi.org/10.1021/es8019275.

(29)   Jiang, W.; Xia, S.; Liang, J.; Zhang, Z.; Hermanowicz, S. W. Effect of Quorum Quenching on the Reactor Performance, Biofouling and Biomass Characteristics in Membrane Bioreactors. *Water Research* **2013**, *47* (1), 187–196. https://doi.org/10.1016/j.watres.2012.09.050.

(30)   Weerasekara, N. A.; Choo, K.-H.; Lee, C.-H. Hybridization of Physical Cleaning and Quorum Quenching to Minimize Membrane Biofouling and Energy Consumption in a Membrane Bioreactor. *Water Research* **2014**, *67*, 1–10. https://doi.org/10.1016/j.watres.2014.08.049.

(31)   Lejars, M.; Margaillan, A.; Bressy, C. Fouling Release Coatings: A Nontoxic Alternative to Biocidal Antifouling Coatings. *Chem. Rev.* **2012**, *112* (8), 4347–4390. https://doi.org/10.1021/cr200350v.

(32)   Wang, D.; Cui, F.; Ren, L.; Tan, X.; Li, Q.; Li, J.; Li, T. Enhancing the Inhibition Potential of AHL Acylase PF2571 against Food Spoilage by Remodeling Its Substrate Scope via a Computationally Driven Protein Design. *J. Agric. Food Chem.* **2022**, *70* (45), 14510–14521. https://doi.org/10.1021/acs.jafc.2c05753.

(33)   Grulich, M.; Štěpánek, V.; Kyslík, P. Perspectives and Industrial Potential of PGA Selectivity and Promiscuity. *Biotechnology Advances* **2013**, *31* (8), 1458–1472. https://doi.org/10.1016/j.biotechadv.2013.07.005.

(34)   Arroyo, M.; de la Mata, I.; Acebal, C.; Castillón, M. P. Biotechnological Applications of Penicillin Acylases: State-of-the-Art. *Appl. Microbiol. Biotechnol.* **2003**, *60* (5), 507–514. https://doi.org/10.1007/s00253-002-1113-6.

(35)   Bruggink, A.; Roos, E. C.; de Vroom, E. Penicillin Acylase in the Industrial Production of β-Lactam Antibiotics. *Org. Process Res. Dev.* **1998**, *2* (2), 128–133. https://doi.org/10.1021/op9700643.

(36)   Avinash, V. S.; Pundle, A. V.; Ramasamy, S.; Suresh, C. G. Penicillin Acylases Revisited: Importance beyond Their Industrial Utility. *Critical Reviews in Biotechnology* **2016**, *36* (2), 303–316. https://doi.org/10.3109/07388551.2014.960359.

(37)   Grulich, M.; Brezovský, J.; ŠtĿpánek, V.; Palyzová, A.; Kyslíková, E.; Kyslík, P. Resolution of α/β-Amino Acids by Enantioselective Penicillin G Acylase from Achromobacter Sp. *Journal of Molecular Catalysis B: Enzymatic* **2015**, *122*, 240–247. https://doi.org/10.1016/j.molcatb.2015.09.008.

(38)   Grulich, M.; Brezovský, J.; Štěpánek, V.; Palyzová, A.; Marešová, H.; Zahradník, J.; Kyslíková, E.; Kyslík, P. In-Silico Driven Engineering of Enantioselectivity of a Penicillin G Acylase towards Active Pharmaceutical Ingredients. *Journal of Molecular Catalysis B: Enzymatic* **2016**, *133*, S53–S59. https://doi.org/10.1016/j.molcatb.2016.11.014.

(39)   Surpeta, B.; Grulich, M.; Palyzová, A.; Marešová, H.; Brezovsky, J. Common Dynamic Determinants Govern Quorum Quenching Activity in N-Terminal Serine Hydrolases. *ACS Catal.* **2022**, *12* (11), 6359–6374. https://doi.org/10.1021/acscatal.2c00569.

(40)   McVey, C. E.; Walsh, M. A.; Dodson, G. G.; Wilson, K. S.; Brannigan, J. A. Crystal Structures of Penicillin Acylase Enzyme-Substrate Complexes: Structural Insights into the Catalytic Mechanism11Edited by K. Nagai. *Journal of Molecular Biology* **2001**, *313* (1), 139–150. https://doi.org/10.1006/jmbi.2001.5043.

(41)   Schymkowitz, J.; Borg, J.; Stricher, F.; Nys, R.; Rousseau, F.; Serrano, L. The FoldX Web Server: An Online Force Field. *Nucleic Acids Res* **2005**, *33* (suppl_2), W382–W388. https://doi.org/10.1093/nar/gki387.

(42)   Chovancova, E.; Pavelka, A.; Benes, P.; Strnad, O.; Brezovsky, J.; Kozlikova, B.; Gora, A.; Sustr, V.; Klvana, M.; Medek, P.; Biedermannova, L.; Sochor, J.; Damborsky, J. CAVER 3.0: A Tool for the Analysis of Transport Pathways in Dynamic Protein Structures. *PLoS Computational Biology* **2012**, *8* (10), e1002708. https://doi.org/10.1371/journal.pcbi.1002708.

(43)   Brezovsky, J.; Thirunavukarasu, A. S.; Surpeta, B.; Sequeiros-Borja, C. E.; Mandal, N.; Sarkar, D. K.; Dongmo Foumthuim, C. J.; Agrawal, N. TransportTools: A Library for High-Throughput Analyses of Internal Voids in Biomolecules and Ligand Transport through Them. *Bioinformatics* **2022**, *38* (6), 1752–1753. https://doi.org/10.1093/bioinformatics/btab872.

(44)   Gordon, J. C.; Myers, J. B.; Folta, T.; Shoja, V.; Heath, L. S.; Onufriev, A. H++: A Server for Estimating p Ka s and Adding Missing Hydrogens to Macromolecules. *Nucleic Acids Res* **2005**, *33* (suppl_2), W368–W371. https://doi.org/10.1093/nar/gki464.

(45)   Anandakrishnan, R.; Aguilar, B.; Onufriev, A. V. H++ 3.0: Automating PK Prediction and the Preparation of Biomolecular Structures for Atomistic Molecular Modeling and Simulations. *Nucleic Acids Research* **2012**, *40* (W1), W537–W541. https://doi.org/10.1093/nar/gks375.

(46)   Hanwell, M. D.; Curtis, D. E.; Lonie, D. C.; Vandermeersch, T.; Zurek, E.; Hutchison, G. R. Avogadro: An Advanced Semantic Chemical Editor, Visualization, and Analysis Platform. *Journal of Cheminformatics* **2012**, *4* (1), 17. https://doi.org/10.1186/1758-2946-4-17.

(47)   M. J. Frisch; G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone; B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng; J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao; H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin; V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts; R. E. Stratmann, O. Yazyev, A.

J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma; V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman; J. V. Ortiz, J. Cioslowski, D. J. Fox. Gaussian 09, Revision D.01, 2013.

(48) Dupradeau, F.-Y.; Pigache, A.; Zaffran, T.; Savineau, C.; Lelong, R.; Grivel, N.; Lelong, D.; Rosanski, W.; Cieplak, P. The R.E.D. Tools: Advances in RESP and ESP Charge Derivation and Force Field Library Building. *Phys. Chem. Chem. Phys.* **2010**, *12* (28), 7821–7839. https://doi.org/10.1039/C0CP00111B.

(49) Cieplak, P.; Cornell, W. D.; Bayly, C.; Kollman, P. A. Application of the multimolecule and multiconformational RESP methodology to biopolymers: Charge derivation for DNA, RNA, and proteins. *Journal of Computational Chemistry* **1995**, *16* (11), 1357–1377. https://doi.org/10.1002/jcc.540161106.

(50) Seeliger, D.; de Groot, B. L. Ligand Docking and Binding Site Analysis with PyMOL and Autodock/Vina. *J Comput Aided Mol Des* **2010**, *24* (5), 417–422. https://doi.org/10.1007/s10822-010-9352-6.

(51) Morris, G. M.; Huey, R.; Lindstrom, W.; Sanner, M. F.; Belew, R. K.; Goodsell, D. S.; Olson, A. J. AutoDock4 and Auto-DockTools4: Automated Docking with Selective Receptor Flexibility. *Journal of Computational Chemistry* **2009**, *30* (16), 2785–2791. https://doi.org/10.1002/jcc.21256.

(52) Daniel, L.; Buryska, T.; Prokop, Z.; Damborsky, J.; Brezovsky, J. Mechanism-Based Discovery of Novel Substrates of Haloalkane Dehalogenases Using in Silico Screening. *J. Chem. Inf. Model.* **2015**, *55* (1), 54–62. https://doi.org/10.1021/ci500486y.

(53) D.A. Case; D.S. Cerutti, T.E. Cheatham, III, V.W.D. Cruzeiro, T.A. Darden,; I.Y. Ben-Shalom; S.R. Brozel; R.E. Duke, D. Ghoreishi, M.K. Gilson, H. Gohlke, A.W. Goetz, D. Greene, R Harris, N. Homeyer, Y. Huang,; S. Izadi, A. Kovalenko, T. Kurtzman, T.S. Lee, S. LeGrand, P. Li, C. Lin, J. Liu, T. Luchko, R. Luo, D.J.; Mermelstein, K.M. Merz, Y. Miao, G. Monard, C. Nguyen, H. Nguyen, I. Omelyan, A. Onufriev, F. Pan, R.; Qi, D.R. Roe, A. Roitberg, C. Sagui, S. Schott-Verdugo, J. Shen, C.L. Simmerling, J. Smith, R. Salomon-; Ferrer, J. Swails, R.C. Walker, J. Wang, H. Wei, R.M. Wolf, X. Wu, L. Xiao, D.M. York and P.A. Kollman. AMBER 2017, 2017.

(54) Maier, J. A.; Martinez, C.; Kasavajhala, K.; Wickstrom, L.; Hauser, K. E.; Simmerling, C. Ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from Ff99SB. *J. Chem. Theory Comput.* **2015**, *11* (8), 3696–3713. https://doi.org/10.1021/acs.jctc.5b00255.

(55) Grigorenko, B. L.; Khrenova, M. G.; Nilov, D. K.; Nemukhin, A. V.; Švedas, V. K. Catalytic Cycle of Penicillin Acylase from *Escherichia Coli* : QM/MM Modeling of Chemical Transformations in the Enzyme Active Site upon Penicillin G Hydrolysis. *ACS Catal.* **2014**, *4* (8), 2521–2529. https://doi.org/10.1021/cs5002898.

(56) Cornell, W. D.; Cieplak, P.; Bayly, C. I.; Gould, I. R.; Merz, K. M.; Ferguson, D. M.; Spellmeyer, D. C.; Fox, T.; Caldwell, J. W.; Kollman, P. A. A Second Generation Force Field for the Simulation of Proteins, Nucleic Acids, and Organic Molecules. *J. Am. Chem. Soc.* **1995**, *117* (19), 5179–5197. https://doi.org/10.1021/ja00124a002.

(57) Waskom, M. L. Seaborn: Statistical Data Visualization. *Journal of Open Source Software* **2021**, *6* (60), 3021. https://doi.org/10.21105/joss.03021.

(58) Clevenger, K. D.; Wu, R.; Er, J. A. V.; Liu, D.; Fast, W. Rational Design of a Transition State Analogue with Picomolar Affinity for Pseudomonas Aeruginosa PvdQ, a Siderophore Biosynthetic Enzyme. *ACS Chem. Biol.* **2013**, *8* (10), 2192–2200. https://doi.org/10.1021/cb400345h.

(59) Hopkins, C. W.; Le Grand, S.; Walker, R. C.; Roitberg, A. E. Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *J. Chem. Theory Comput.* **2015**, *11* (4), 1864–1874. https://doi.org/10.1021/ct5010406.

(60) D.A. Case; I.Y. Ben-Shalom; S.R. Brozel; D.S. Cerutti, T.E. Cheatham, III, V.W.D. Cruzeiro, T.A. Darden,; R.E. Duke, D. Ghoreishi, M.K. Gilson, H. Gohlke, A.W. Goetz, D. Greene, R Harris, N. Homeyer, Y. Huang,; S. Izadi, A. Kovalenko, T. Kurtzman, T.S. Lee, S. LeGrand, P. Li, C. Lin, J. Liu, T. Luchko, R. Luo, D.J.; Mermelstein, K.M. Merz, Y. Miao, G. Monard, C. Nguyen, H. Nguyen, I. Omelyan, A. Onufriev, F. Pan, R.; Qi, D.R. Roe, A. Roitberg, C. Sagui, S. Schott-Verdugo, J. Shen, C.L. Simmerling, J. Smith, R. Salomon-; Ferrer, J. Swails, R.C. Walker, J. Wang, H. Wei, R.M. Wolf, X. Wu, L. Xiao, D.M. York and P.A. Kollman. AMBER 2018, 2018.

(61) Pedregosa, F.; Varoquaux, G.; Gramfort, A.; Michel, V.; Thirion, B.; Grisel, O.; Blondel, M.; Prettenhofer, P.; Weiss, R.; Dubourg, V.; Vanderplas, J.; Passos, A.; Cournapeau, D.; Brucher, M.; Perrot, M.; Duchesnay, É. Scikit-Learn: Machine Learning in Python. *Journal of Machine Learning Research* **2011**, *12* (85), 2825–2830.

(62) McInnes, L.; Healy, J.; Astels, S. Hdbscan: Hierarchical Density Based Clustering. *Journal of Open Source Software* **2017**, *2* (11), 205. https://doi.org/10.21105/joss.00205.

(63) Schmidtke, P.; Bidon-Chanal, A.; Luque, F. J.; Barril, X. MDpocket: Open-Source Cavity Detection and Characterization on Molecular Dynamics Trajectories. *Bioinformatics* **2011**, *27* (23), 3276–3285. https://doi.org/10.1093/bioinformatics/btr550.

(64) Le Guilloux, V.; Schmidtke, P.; Tuffery, P. Fpocket: An Open Source Platform for Ligand Pocket Detection. *BMC Bioinformatics* **2009**, *10* (1), 168. https://doi.org/10.1186/1471-2105-10-168.

(65) Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science Engineering* **2007**, *9* (3), 90–95. https://doi.org/10.1109/MCSE.2007.55.

(66) Sobotková, L.; Štěpánek, V.; Plháčková, K.; Kyslík, P. Development of a High-Expression System for Penicillin G Acylase Based on the Recombinant Escherichia Coli Strain RE3(PKA18). *Enzyme and Microbial Technology* **1996**, *19* (5), 389–397. https://doi.org/10.1016/S0141-0229(96)00052-X.

(67) Marešová, H.; Palyzová, A.; Kyslík, P. The C-Terminal Region Controls Correct Folding of Genus Trametes Pyranose 2-Oxidases. *Journal of Biotechnology* **2007**, *130* (3), 229–235. https://doi.org/10.1016/j.jbiotec.2007.04.023.

(68) Valešová, R.; Hollerová-Sobotková, L.; Štěpánek, V.; Kyslík, P. Optimization of the Host–Plasmid Interaction in the Recombinant Escherichia Coli Strains Overproducing Penicillin G Acylase. *Enzyme and Microbial Technology* **2004**, *35* (1), 74–80. https://doi.org/10.1016/j.enzmictec.2004.03.015.

(69) Peleg, Y.; Unger, T. Application of the Restriction-Free (RF) Cloning for Multicomponents Assembly. In *DNA Cloning and Assembly Methods*; Valla, S., Lale, R., Eds.; Methods in Molecular Biology; Humana Press: Totowa, NJ, 2014; pp 73–87. https://doi.org/10.1007/978-1-62703-764-8_6.

(70) Kutzbach, C.; Rauenbusch, E. Preparation and General Properties of Crystalline Penicillin Acylase from Escherichia Coli ATCC 11 105. *Hoppe-Seyler's Z. Physiol. Chem.* **1974**, *355* (1), 45–53. https://doi.org/10.1515/bchm2.1974.355.1.45.

(71) Last, D.; Krüger, G. H. E.; Dörr, M.; Bornscheuer, U. T. Fast, Continuous, and High-Throughput (Bio)Chemical Activity Assay for N-Acyl-l-Homoserine Lactone Quorum-Quenching Enzymes. *Applied and Environmental Microbiology* **2016**, *82* (14), 4145–4154. https://doi.org/10.1128/AEM.00830-16.

(72) Bendl, J.; Stourac, J.; Sebestova, E.; Vavra, O.; Musil, M.; Brezovsky, J.; Damborsky, J. HotSpot Wizard 2.0: Automated Design of Site-Specific Mutations and Smart Libraries in Protein Engineering. *Nucleic Acids Res* **2016**, *44* (W1), W479–W487. https://doi.org/10.1093/nar/gkw416.

(73) Sumbalova, L.; Stourac, J.; Martinek, T.; Bednar, D.; Damborsky, J. HotSpot Wizard 3.0: Web Server for Automated Design of Mutations and Smart Libraries Based on Sequence Input Information. *Nucleic Acids Res* **2018**, *46* (W1), W356–W362. https://doi.org/10.1093/nar/gky417.

(74) Novikov, F. N.; Stroganov, O. V.; Khaliullin, I. G.; Panin, N. V.; Shapovalova, I. V.; Chilov, G. G.; Švedas, V. K. Molecular Modeling of Different Substrate-Binding Modes and Their Role in Penicillin Acylase Catalysis. *The FEBS Journal* **2013**, *280* (1), 115–126. https://doi.org/10.1111/febs.12054.

(75) Laj, N.; Elayadeth-Meethal, M.; Huxley, V. A. J.; Hussain, R. R.; Saheer Kuruniyan, M.; Poonkuzhi Naseef, P. Quorum-Sensing Molecules: Sampling, Identification and Characterization of N-Acyl-Homoserine Lactone in Vibrio Sp. *Saudi Journal of Biological Sciences* **2022**, *29* (4), 2733–2737. https://doi.org/10.1016/j.sjbs.2021.12.062.

(76) Yasutake, Y.; Kusada, H.; Ebuchi, T.; Hanada, S.; Kamagata, Y.; Tamura, T.; Kimura, N. Bifunctional Quorum-Quenching and Antibiotic-Acylase MacQ Forms a 170-KDa Capsule-Shaped Molecule Containing Spacer Polypeptides. *Scientific Reports* **2017**, *7* (1), 8946. https://doi.org/10.1038/s41598-017-09399-4.

(77) Murugayah, S. A.; Gerth, M. L. Engineering Quorum Quenching Enzymes: Progress and Perspectives. *Biochem. Soc. Trans.* **2019**, *47* (3), 793–800. https://doi.org/10.1042/BST20180165.

(78) Billot, R.; Plener, L.; Jacquet, P.; Elias, M.; Chabrière, E.; Daudé, D. Engineering Acyl-Homoserine Lactone-Interfering Enzymes toward Bacterial Control. *J. Biol. Chem.* **2020**, *295* (37), 12993–13007. https://doi.org/10.1074/jbc.REV120.013531.