Title: Integration of leaf spectral reflectance variability facilitates identification of plant

leaves at different taxonomic levels.

Authors:

Quinteros Casaverde, Natalia L.[1,3],

Serbin, Shawn P.[2]

Daly, Douglas C.[1,3]

Affiliations:

[1] Biology Department, Graduate Center of the City University of New York

[2] Brookhaven National Laboratory

[3] New York Botanical Garden

ORCID:
Quinteros Casaverde, Natalia L. 0000-0003-1926-640X
Serbin, Shawn P. 0000-0003-4136-8971
Daly, Douglas C. 0000-0003-1205-9491

Abstract

Plant identification is crucial to the conservation and management of natural areas. The shortwave

spectral reflectance of leaves is a promising tool for rapidly identifying species at different

taxonomic ranks. However, the spectral reflectance of leaves changes in response to biotic and

abiotic conditions. This investigation asked how this variability affects the accuracy of methods

used to predict plant taxonomies and what factors most influence the spectral signature of leaves.

To answer these questions, we measured the reflectance of leaves of 62 woody species from the

living collection at the NYBG twice in two pairwise samplings. We found that PLS-DA accuracy

improved when we used a larger sample of natural variance in the classification model. Finally, to

evaluate whether there was an influence of the species' relatedness or the growing environment on

structural and biochemical traits predicted from the leaf reflectance, we ran a phylogenetic signal

analysis and a series of mixed effects model analyses that showed no phylogenetic but an

environmental influence. We found that the increase in temperature and relative humidity

variability explained the increment of predicted carotene and the decrease of Nitrogen content for

the first pairwise analysis. For the second pairwise analysis, we found that the reduction of relative

humidity variability explained leaf water and Nitrogen content decrease, and relative humidity

decrease combined with day length decrease explained a decline in LMA.

Key words: spectral reflectance, taxonomic classification, PLS-DA, PROSPECT-D RTM, PLSR,

predicted structural and biochemical traits, phylogenetic signal, environmental influence.

1. INTRODUCTION

Plant identification is key to conservation (Trias-Blasi & Vorontsova, 2015) and management (Noss, 2001), and hence essential for human survival (Stuessy et al., 2014). Recent studies have shown that the shortwave spectral reflectance of leaves (0.3 to 2.5 microns), captured with proximal remote sensing instruments, is a promising tool for rapidly identifying species of plants into their different taxonomic ranks with high accuracy (Asner et al., 2014; Buitrago et al., 2018; Cavender-Bares et al., 2016, 2017; Cochrane, 2000; Durgante et al., 2013; Féret & Asner, 2011; Hesketh & Sánchez-Azofeifa, 2012; Lang et al., 2015, 2017; Meireles et al., 2020; Paiva et al., 2021; Price, 1994; Stasinski et al., 2021). It is expected that with the advent of new hyperspectral satellites (Qian, 2021), form the ground (e.g. in rapid floristic assessments; Higgins & Ruokolainen, 2004; Marcelo-Peña et al., 2015) the ability to identify plant species over large areas and through time with spectral methods will improve.

However, the spectral reflectance of leaves of any given species is excepted to change over time (Hesketh & Sánchez-Azofeifa, 2012). Because of this, before the addition of the spectra to the plethora of traits required for taxonomy or phylogenetics (Vale, 2019), we need to understand the effect of these variability on the spectral signature of leaves (Castro-Esau et al., 2006; Hesketh & Sánchez-Azofeifa, 2012).

The spectral signature or the spectral reflectance pattern of a leaf is the intensity of incident radiation reflected to the environment, measured across the visible (VIS), near infrared (NIR), and shortwave infrared (SWIR) portions of the electromagnetic spectrum, from 400 nm to 2400 nm. This reflectance pattern varies depending on the chemical and structural attributes of the

leaf, (e.g. photosynthetic pigments, water, cellulose, lignin; Brewer, 2012), and secondary

metabolites concentrations (Fine et al., 2021). Because these components absorb, reflect, and

transmit portions of the electromagnetic spectrum differentially (Campbell & Norman, 1998;

Schweiger et al., 2017) the spectral signature is a phenotypic manifestation of leaves. Moreover,

the environment and the evolutionary history (phylogenetic effect) of a plant promote variation

in leaf phenotype (Díaz et al., 2016; Doyle, 2007; Fernando et al., 2007; Little et al., 2010) and

the spectral signature is not an exception (Cavender-Bares et al., 2016; Meireles et al., 2020).

Phylogenetic effects impact wavelengths differentially at higher taxonomic ranks, and differ

among classes (Meireles et al., 2020). The spectral signatures of plant species, from dry and

moist habitats in Panama, have shown to vary with the seasons of the year (Hesketh & Sánchez-

Azofeifa, 2012). Spectral reflectance also differ across individuals of the same species when

found in diffent habitas (Castro-Esau et al., 2006).

Most of the traits affecting the spectral signature of leaves are labile (Ball et al., 2015; Cotrozzi

et al., 2017; De La Riva et al., 2016; Fyfe, 2003; Gamon et al., 2015; González-Rebeles et al.,

2021; Li et al., 2018; Nguyen et al., 2017; Nunes et al., 2019; Paz-Kagan & Asner, 2017; Reich

et al., 1991; Yang et al., 2016; Young, 1991). It is possible to predict many chemical and

structural leaf traits from the spectral signature by using inversion of leaf level radiative transfer

models (Fourty et al., 1996; Jacquemoud & Baret, 1990) regressions methods such as Partial

Least Squares (PLSR; Serbin et al., 2014), and previous studies suggest that reflectance data

closely tracks underlying changes in leaf biochemistry (e.g. Yang et al., 2016). Hence, we can

use these traits to evaluate spectral responses to a fluctuating environment. Moreover, since the

predicted values of these chemical and structural traits are a good approximation of empiric

measurements (Blackburn, 2007; Burnett, Anderson, et al., 2021; Ely et al., 2019; Féret & Asner,

2011; Fourty et al., 1996; Nakaji et al., 2019; Serbin et al., 2014; Shiklomanov et al., 2016), it is

possible to make inference on what trait is contributing the most to the temporal variability of the

spectra.

The present work explores how the spectral signature of leaves and their temporal variability

help us to identify tropical woody species from several regions of the world yet found in an

artificial environment: the Enid A. Haup Conservatory (EAHC), at the New York Botanical

Garden (NYBG). First, we ask how the variation of the spectral signature of leaves impacts the

identification of species into their taxonomic levels (from family to the species) and how to

better integrate the variation of the spectra into the classification models. We expect that the

integration of all the temporal spectral variability will outperform the classification compared to

the classification of species based on just one collecting season. Second, we ask what

environmental factors within the EAHC (mean and standard deviation of temperature and

relative humidity, and mean day length) have an effect over the predicted values of leaf

photosynthetic pigments (total chlorophyl a & b content, carotene content), leaf water content,

leaf nitrogen content, and leaf dry mass per area (LMA), where we expect variation of these

traits as a response to the changing environmental conditions.

2. METHODS:

2.1. Data Collection

We collected leaves from a total of 62 woody species (Table 1S. One individual per species)

belonging to the botanical families Acanthaceae, Ericaceae, Fabaceae, Myrtaceae, Rubiaceae,

and Solanaceae from the Rainforest Pavilion at the EAHC. The initial collection was made

during the days of the year (DOY) 105 and 112 during the spring of 2019 where we collected

five leaves from 51 species (1 individual per species). The second collection was during the

summer of 2020 (DOY 251) where we collected 3 leaves from 24 species previously sampled

and collected 18 new species (1 individual per species). The final collection took place during

the winter of 2021 (DOY 57) where we also collected 3 leaves from 16 species previously

collected and 3 new species (1 individual per species). All collections were made between 10 am

and 3pm, the number of leaves were different during the second and third collection due to the

COVID19 pandemic restrictions at the NYBG.

Two spectroradiometers were used in 2019: A Spectra Vista Corporation (SVC HR 1024i) and

Spectral Evolution (SE PSR+) using the same type of foreoptic, an SVC LC-RP-Pro leaf clip

with an internal, calibrated and full-spectrum light source. All leaf reflectance measurements

were made inside the LC-RP-Pro leaf clips and were calibrated with an Spectralon® white

reference. Despite using two different spectroradiometers, because they both covered the same

spectral range, were radiometrically calibrated, and used the same Spectralon® references, the

data could be combined together without needing any additional inter-calibration (e.g. Serbin et

al., 2019). In 2020 and 2021 we used the SVC HR 1024i with the same leaf clip.

2.2. **Prediction of Leaf Structural and Chemical Traits.**

To estimate pigment (chlorophylls and carotenoids) and leaf water content, we used the Bayesian

inversion of the PROSPECT-D RTM as implemented in the rrtm package

(https://github.com/ashiklom/rrtm). In PROSPECT, the leaf is simulated as an N number of

semi-translucid flat plates, where each plate has a specific wavelength-dependent transmissivity

(Feret et al., 2008). Moreover, the model transmissivity is based on the linear combination of

specific spectral absorption measurements of the total photosynthetic pigment content

(chlorophyll a + b and carotenoids) as well as water and dry matter content, which are multiplied

by their respective quantities (Cab µg/cm^2, Car µg/cm^2, Cw g/cm^2) obtained from known

datasets: LOPEX, ANGERS, HAWAII, and CALMIT. To evaluate the accuracy in the resulting

parameter values, we used Shiklomanov et al., (2016) framework that provides the joint

probability distribution of the PROSPECT parameters, with the residual's standard deviation as

outputs.

To obtain leaf mass per area (LMA) and leaf Nitrogen content per area, we used the PLSR

framework implemented in Burnett et al. (2021), which predicts these traits by using published

coefficients (https://zenodo.org/badge/latestdoi/222699149). The output of PLSR modeling is a

linear algorithm of regression coefficients by waveband that explains the contribution of each

specific band to the prediction of a specific leaf functional trait (Burnett, Serbin, et al., 2021;

Serbin et al., 2014)

2.3. Statistical Analysis

2.3.1. **Effects of temporal variation on the classification of leaf spectra at different**

taxonomic levels. We investigated how temporal changes in leaf properties and

resulting reflectance signatures would impact classification approaches. To do this,

we used the Partial Least Squared Discriminant Analysis (PLS-DA) to classify leaves

into their corresponding families and genera using samples from one season, and then

apply to a different sampling date. At the family level the PLS-DA was ran with 43

species (one individual per species; 5 leaves per ind.) from the six botanical families

mention above. At the genus level, we ran the PLS-DA with 22 species (one

individual per species) belonging to 9 genera. For both analyses, we trained the model

with 70% of the data and tested them with 30% and set the cross-validated by using

bootstrap cross-validation (Rodríguez-Pérez et al., 2018) by using the trainControl

and train functions, respectively. The trainControl and train functions are part of the

caret package (Kuhn & Johnson, 2013). Once we obtained the optimal number of

components from the trained data and used them to run the PLS-DA with the plsda

function also from the caret package (Kuhn & Johnson, 2013) in R. To test for the

effect of spectral variability, we ran the PLS-DA using leaves sampled in 2019 for

training and 2020 – 2021 leaves for testing and classify leaves at the family, genus,

and species level. For the analyses at the species level, we used 43 individuals. After

we ran another PLS-DA for all datasets together (2019, 2020, 2021) by creating a

train/test datasets (70/30) with a stratified random data partition across species and

collection dates with the same parameters stated for the first PLS-DA. To detect

areas of high variability within the spectral signature, we calculated the coefficient of

variation (CV) between the leaves of species sampled in 2019 and resampled in 2020

and 2021. We calculated the CV separately for each resampling season because the

group of species sampled were different each time. To visualize these both the

spectral signatures and the resulting CV differences, we plotted this information into

heatmaps using the base function heatmap from R.

2.3.2. **Phylogenetic signal and principal component analyses:** To determine the

contribution of the phylogeny into the spectral signature variability, we ran a

phylogenetic signal analysis on the structural and chemical traits derived from

195 PROSPECT-D and PLSR models as proxies for directly measured functional traits.

196 For that, we split the species based on date of resampling and calculated the

197 interquartile rage to account for trait variation. We ran a phylogenetic signal analysis

198 using the function physig from the phytools package (Revell, 2012) based on Pagel's

199 lambda (Pagel, 1999) which accounts for polytomies (Münkemüller et al., 2012)

200 within the *Macleania* clade (Hooker, 1837). We extracted the phylogenetic distances

201 from the mega-tree GBOTB.extended.tre provided by the package V.Phylomaker in

202 R (Jin & Qian, 2019) for each analysis. We then ran a principal component analysis

203 (PCA) to determine trait contribution of each trait to the variation of species using the

204 function prcomp from the stats package and the function fviz_pca_var from the

205 package factoextra (Kassambara & Mundt, 2020).

206 2.3.3. **Effects of temperature, relative humidity, and length of the day:** To test for the

207 effect of the environment over the derived traits, we first run a Pearson correlation to

208 build the models, to avoid bias created by the strength of the correlation (Schielzeth

209 et al., 2020). For that, we use the cor.test function from the package stats in R. After

210 the strength of the correlation was established, we built the models to test for the

211 effect of temperature (°C , mean/standard deviation), relative humidity (%,

212 mean/standard deviation), and mean length of the day (hours) over the derived traits.

213 The environmental data was downloaded from sensors inside EAHC, which are

214 connected to a PRIVA® system that collects temperature and relative humidity every

215 5 minutes. To calculate the length of the day we used the function daylength from the

216 insol package (Corripio, 2019). We calculated the mean and standard deviation of all

217 environmental variables for 30 days prior and including the measurement day (Van

Goethem et al., 2013). We build seven mixed-effect models by first establishing the

random effect by nesting species into collection date, and the fixed effects

environmental variables as fixed effects. The first five models included mean

temperature, mean relative humidity, and mean day length and the respective

interactions. The sixth model included the standard deviation of temperature and the

seventh the standard deviation of relative humidity. We use the function lmer from

the lme4 package (Bates et al., 2015) in R and mixed from the afex package which

calculates the p-value using the Satterthwaite's method (Henrik Singmann et al.,

2021). We used the BoxCoxME function from the tramME package to transform

heteroscedastic models with the error distribution set as Standard Gaussian and a

Bernstein basis transformation (Tamási & Hothorn, 2021). Each mixed-effect models

were run separately for individuals resampled in 2020 and 2021.

3. RESULSTS

The leaf classification, when using only samples from 2019, was very successful. The PLS-DA

model was able to classify the leaves with an accuracy of 0.95 (Figure 1A, Table 2). At the genus

level the accuracy level was 1 (Figure 1B). However, the inclusion of temporal variability into

our classification models decreased its accuracy when the PLS-DA data was spited (training data

with leaves from 2019; testing data with leaves from 2020 & 2021), but the accuracy increased

tremendously when the PLS-DA model took the whole sample's variability into account. PLS-

DA models using the 2019 dataset for training and the other two as testing datasets had a poorer

performance (Figure 2: A, C, E) compared to the PLS-DA models containing all three datasets

together and split into train/test datasets by using a stratified partition method (Figure 2: B, D, F).

At the family level the classification accuracy improved from 0.39 to 0.94. At the genus level,

from 0.47 to 0.98. At the species level, from 0.21 to 0.96. All the models had an accuracy

significantly different and substantially higher than the no information rate, and the kappa values

for the stratified models were higher than 0.9 (Table 2).

Table 1. PLS-DA analysis for Split datasets, where 2019 leaves were used as training and 2020 and 2021 and

Stratified datasets where all leaves were classified together and partitioned considering collection date. N of Comp is

the number of components, 95% CI is the 95% confidence interval, and No Inf Rate is the no information rate.

Model	N of Comp	Accuracy	95% CI	No Inf Rate	p-value	Kappa
Family 2019	26	0.93	0.84, 0.97	0.2	< 0.001	0.91
Family Split	59	0.29	0.22, 0.36	0.24	< 0.001	0.14
Family Stratified	35	0.9	0.84, 0.94	0.22	< 0.001	0.9
Genus 2019	36	1	0.9,1	0.23	<0.001	1
Genus Split	50	0.34	0.24, 0.47	0.2	< 0.001	0.2
Genus Stratified	33	0.98	0.91, 0.99	0.22	< 0.001	0.98
Species Split	60	0.07	0.04,0.13	0.04	< 0.001	0.04
Species Stratified	60	0.97	0.91,0.99	0.06	< 0.001	0.97

We observed that the leaf spectral signatures measured at two different times of the year varied

tremendously. The comparison between species resampled in 2020 (Figure 3C) shows that the

areas of the spectrum with highest variability are in the visible (VIS 500-700 nm) and in the red-

near infrared transition zone or red edge (~720 nm). For the same period, most species showed

increased variability at the beginning of the first downward curve of the short-wave infrared

known as SWIR I, as well as at 1900 nm which is where the second downward curve of SWIR

starts (SWIR II).

We did not find a phylogenetic signal for the interquartile range of any of the derived traits. The

phylogenetic signal analysis was not different from zero for any of the traits for both datasets (p-

value > 0.05). The PCA indicates that the traits that contribute most of the variation between the

species resampled in 2020 are LMA (PC1: 24.2%; PC2: 16.4%), total chlorophyll content/area

(PC1: 21%; PC2: 25.6%), and carotene content/area (PC1: 19.9%; PC2: 25.6%). For this PCA,

PC1 accounted for 62.1% of the variation and PC2 for 24.3%. For species resampled in 2021,

the traits that contribute to most of the variation were nitrogen content/area (PC1: 13.6%; PC2:

47.1%), LMA (PC1: 23.8%; PC2: 15.4%), and total chlorophyll content/area (PC1: 27.3%; PC2:

3.8%). For this PCA, PC1 accounted for 61.1% of the variation and PC2 for 21.6% (Figure 4).

For the environmental variables the correlation between mean temperature, mean relative

humidity, and mean daylength was weak (Fig. 2S). The correlation coefficients between the

standard deviation of both temperature and relative humidity were high with all other variables

and between themselves, except with mean relative humidity (Fig. 2S). Hence, we build seven

mixed-effect models according to these results (Table 3).

Model	Model Specification
M₁	$y = \beta \ T_{mean} + u \ (individual/year) + \varepsilon$
M₂	$y = \beta_1 \ T_{mean} + \beta_2 \ RH_{mean} + u \ (individual/year) + \varepsilon$
M₃	$y = \beta_1 \ T_{mean} + \beta_2 \ RH_{mean} + \beta_3 \ DL_{mean} + u \ (individual/year) + \varepsilon$
M₄	$y = \beta_1 \ T_{mean} * RH_{mean} + \beta_3 \ DL_{mean} + u \ (individual/year) + \varepsilon$
M₅	$y = \beta_1 \ T_{mean} * RH_{mean} * \beta_3 \ DL_{mean} + u \ (individual/year) + \varepsilon$
M₆	$y = \beta \ T_{sd} + u \ (individual/year) + \varepsilon$
M₇	$y = \beta \ RH_{sd} + u \ (individual/year) + \varepsilon$

The results for the mixed-model analysis showed significant contribution of these environmental

factors on the variation of the derived structural and chemical traits. The mixed effect models for

species resampled in 2020 (Table 4) showed that the increased of daily temperature variation

(sdT$_{2019}$ < sdT$_{2020}$, Table 2S, Fig 2S), explained the increase of carotene content/area (p-value <

0.01; Fig 7C), and that the increase of daily relative humidity variation (sdRH$_{2019}$ < sdRH$_{2020}$,

Table 2S), explained the slight decrease in leaf nitrogen content/area (p-value < 0.01; Fig 7I). The

mixed effect models for species resampled in 2021 (Table 5) showed that the decrease in the

variation of relative humidity (sdRH$_{2019}$ > sdRH$_{2021}$, Table 2S, Fig 2S) had a downward effect over

nitrogen leaf content/area and the Box-Cox transformation of leaf water content/area, (p-value <

0.01), similar results are shown by untransformed model. Finally, the decrease in the average

relative humidity, combined with a decrease of the mean daylength had a downward effect of LMA

(p-values < 0.05; Fig 7H). All models presented were non heteroscedastic (Fligner test p-value >

0.05).

Table 3. Results for most parsimonious mixed effect models for species resampled in 2020.

Trait	Environmental Variable	Estimated value	Std. Error
Total Chlorophyll	Intercept	0.63356	0.03553
	Stand. Dev. Temperature (C)	0.03459	0.02356
Carotene	(Intercept)	0.164452	0.007807
	Stand. Dev. Temperature (C)	0.013512	0.004162
Leaf Mass per Area (transformed)	Stand. Dev. Relative Humidity (%)	-0.20381	0.15369
Nitrogen	(Intercept)	1.65447	0.10353
	Stand. Dev. Relative Humidity (%)	-0.14313	0.03536

Table 4. Results for most parsimonious mixed effect models for species resampled in 2021.

Trait	Environmental Variable	Estimated value	Std. Error
Total Chlorophyll	(Intercept)	0.67477	0.04248
	Mean Temperature (C)	0.04785	0.04172
Carotene	(Intercept)	0.169376	0.007304
	Mean Temperature (C)	0.008505	0.007184
Water (transformed)	Stand. Dev. Relative Humidity (%)	1.20086	0.20834
Leaf Mass per Area	(Intercept)	35.435	7.883
	Mean Temperature (C)	1.765	8.401
	Mean Relative Humidity (%)	8.208	12.501
	Mean Daylength (Hours)	10.865	8.917
	Mean Temperature (C) * Mean Relative Humidity (%)	-4.534	12.11
	Mean Temperature (C) * Mean Daylength (hours)	9.843	6.66
	Mean Relative Humidity (%) * Mean Daylength (hours)	19.229	7.123
	Mean Temperature (C) * Mean Relative Humidity (%) * Mean Daylength (hours)	-6.656	8.203
Nitrogen	(Intercept)	1.3465	0.10977
	Stand. Dev. Relative Humidity (%)	0.44352	0.08419

4. DISCUSSION

The identification of species at the taxonomic ranks of family, genus, and species were highly

accurate when the variability of the spectral collection was taking into consideration (Figure 2 B,

D, F, & Table 2). Kappa > 90% in all cases, which is indicative of good model performance (Landis

& Koch, 1977; Schreier, 2012). It is therefore important to build spectral libraries for PLS-DA or

other types of classification methods that consider the variability of leaf reflectance within and

across species and though time to improve species identification. This is consistent with other

studies exploring the ability to estimate traits with spectra (Schmitt et al., 2021). The errors made

by the classification models in some cases could be an effect of plant relatedness at higher

taxonomic levels, where spectral reflectance have shown phylogenetic signal (Meireles et al.,

2020). For instance, for the classification at the family level in the 2019 dataset, the

misidentification of leave corresponding to the Rubiaceae family as belonging to the Acanthaceae

family, could be linked to both belonging to from the Lamiids clade (Stevens, 2012). In the case

of leaves misidentified with the stratified method, leaves belonging to the Solanaceae family were

misidentified as Rubiaceae and vice versa, and that could be due to their shared relationship as part

of Lamiid clade (Asterids) as well. At the genus level, Thunbergia was misidentified as

Aphelandra, both genera belong to the Acanthaceae family (Lamiid/Asterids). At the species level,

Pterocarpus officinalis, misclassified as *Tipuana tipu,* they both belong to the Fabaceae (Rosids)

and *Thunbergia grandiflora* misidentified as *Aphelandra flava*, are both Acanthaceaes (Lamiids;

Stevens, 2012).

When analyzing the variability of the spectra (Figure 3) we found high variability in the red edge

zone (~720 nm), supporting Hesketh & Sánchez-Azofeifa's (2012) findings with other tropical

species sampled in Panama at different times of the year. We also found that areas of high

coefficient of variation are linked to other important absorption features for water, proteins, and

cellulose. For instance, the individuals resampled in 2020 showed high variability in the visible

and red – NIR transition zone (red-edge) where we can find absorption features of chlorophyl at

660 nm and at around 400 nm where carotenoids absorb radiant energy. For the same comparison,

variability at the SWIR I range increased. In this spectral region we can find absorption features

of water content, specifically at 1450 nm (Curran, 1989; Jacquemoud et al., 1996; Schweiger et

al., 2018; Ustin et al., 2009).

The lack of phylogenetic signal in the interquartile range for each chemical and structural trait

suggests that the evolutionary history is not impacting the plasticity of these traits. These results

could be indicative that the variability observed is most likely coming from the environment or

other attributes of the leaf such as age (Chavana-Bryant et al., 2019).

LMA, total chlorophyl, and carotene contents/area contributed the most to the PC1, for species

resampled in 2020. Nitrogen, LMA, and total chlorophyll where the major contributors were to

PC1, for species resampled in 2021. Since these derived traits underlie leaf spectra (Meireles et

al., 2020) we can expect that the areas of the spectra connected to these traits are varying the most,

and subsequently that the variation of these traits are causing the variation of the spectra.

The growth environment measured as the standard deviation of temperature, and the standard

deviation of relative humidity had statistically significant effect over carotene content/area and

leaf nitrogen content/area for the species resampled in 2020. But, chlorophyll content/area nor

LMA showed statistically significant results. Increases in carotenes are expected during the

summer months when the incident radiation is higher and the photosystems need to be protected

(Young, 1991). Nitrogen content/area showed a decrease (Fig. 1S) related to the increase in the

variability of the EAHC relative humidity (RH$_{sd}$, Table 4). There is no direct evidence of how an

increase in the variability of relative humidity affects photosynthetic (RuBisCo) and non-

photosynthetic nitrogen content in leaves, but there is evidence that when relative humidity

increases, water flux decreases (Kupper et al., 2011) which could affect nutrient uptake from the

soil. At the same time both nitrogen content and LMA, are positively correlated (Onoda et al.,

2017) and we can observe that LMA decreased as well for these species. For species resampled in

2021, the variation of relative humidity (M7; Table 3) had a statistically significant effect over

water (box-cox transformed) and nitrogen content/area respectively. Additionally, the interaction

of mean humidity and length of daylight (M4; Table 3) had a statistically significant effect over

LMA. The drier conditions within EAHC during the winter months could affect leaf water content,

since watering was kept constant, and the relative amount of water in the air affects stomatal

responses, and photosynthesis (Broughton et al., 2021). This second response may affect leaf

nitrogen content as well. Nitrogen is an important component of photosynthetic pigments, such

as chlorophyll, and proteins (such as RuBisCo), so, it is not surprising that reduced relative

humidity and shorter days are impacting photosynthesis and protein production (Meletiou-

Christou et al., 1994; Muller et al., 2011), as well as LMA.

We can conclude that the areas of the spectrum influenced by photosynthetic pigments, water,

nitrogen, and the constituents of LMA are driving the variability of the spectral signature, through

time. This type of spectral variation affects species identification (Hesketh & Sánchez-Azofeifa,

2012) and trait predictions (Schmitt et al., 2021). The use of remote sensing instruments is gaining

momentum in the study of biodiversity from space is a field that is gaining momentum because of

the capacity instruments to cover large areas of the globe year-round (Cavender-Bares et al., 2022).

Spectral variability affects the identification of species. We now know this variability is partly

linked to the environment through its influence over specific leaf chemical and structural traits.

The accuracy of the identification increases when the variability of the spectral signal is accounted

for, and that should be a lesson for the development of more reliable spectral libraries. This will

not only help the study of biodiversity at larger scales with the use of hyperspectral satellites, but

it could be an excellent resource for rapid biodiversity assessments in the ground that become

decision making tools for natural resources managers.

ACKNOWLEDGEMENTS

We would like to thank Drs Amy Berkov, Ana Carnaval, Kyle Dexter, Fabian Michelangeli for

their valuable input in the development of this manuscript. We would like specially to thank Marc

Hachadourian, Christian Primeau, and all the staff of the Enid A. Haup Conservatory at the New

York Botanical Garden for allowing us to work there before and during the COVID19 pandemic

and for maintaining the living collections and LaGuardia Comminuty College students Ludia

Adhikary and Kenia Martinez for their valuable help during data collection. We also would like to

thank the CUNY EEB graduate students who contributed to this article with their comments and

constructive criticism. Finally, we would like to thank the Graduate Center of CUNY for the

research grants that allowed the completion of this project.

Figures

420
421
422 *Figure 1. Confusion Matrix of the partial least square discriminant analysis at the family level (test dataset N =80 leaves)*

423 *and genus level (test dataset N = 35). (A) The PLS-D analysis had an accuracy of 93% and specific accuracy ranges*

424 *from 91% - 100% at the family level. (B) The PLS-D analysis had an accuracy of 100%. Specific accuracy of 100% at*

425 *the genus level.*

426
427
428   *Figure 2. PLS-D analysis with 2019 as the training data set and 2020 and 2021 as test datasets (A, C, E), and all*

429   *datasets together with stratified partition (B, D, F). At the Family level the accuracy, split pls-da had an accuracy of 59%*

430   *and the stratified analysis had an accuracy of 90%. The specific accuracy varied 50% to 69% for the split pls-da, and*

431   *from 89% to 100% for the stratified pls-da.*

459   *Figure 3. Spectral signature from mean reflectance (%) for species sampled in the spring of 2019 (A & D), resampled in the summer*

461   *of 2020 (B) and the winter of 2021 (E). Coefficient of variation (%) calculated from leaves collected in 2019 and resampled in 2020*

462   *(C), and from leaves collected in 2019 and resampled in 2021(F). Spectral reflectance was measured from 400 nm to 2400 nm and*

464   *ranges from 0.72 – 61% (yellow to dark red) and coefficient of variation 0.01- 0.66% (yellow to dark read).*

465
466 *Figure 4. Contribution of traits to the variance explain by PC1 (or Dim 1) and PC2 (or Dim2) for A) species resampled*

467 *in 2020, and B) species resampled in 2021.*

Bibliography

Asner, G. P., Martin, R. E., Carranza-Jim Enez, L., Sinca, F., Tupayachi, R., Anderson, C. B., &

    Martinez, P. (2014). Functional and biological diversity of foliar spectra in tree canopies

    throughout the Andes to Amazon region. *New Phytologist*, *204*, 127–139.

    https://doi.org/10.1111/nph.12895

Ball, A., Sanchez-Azofeifa, A., Portillo-Quintero, C., Rivard, B., Castro-Contreras, S., &

    Fernandes, G. (2015). Patterns of Leaf Biochemical and Structural Properties of Cerrado

    Life Forms: Implications for Remote Sensing. *PLOS ONE*, *10*(2), e0117659.

    https://doi.org/10.1371/JOURNAL.PONE.0117659

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). *lme4: Linear mixed-effects models*

    using Eigen and S4 (p. 127). https://cran.r-project.org/package=lme4

Blackburn, G. A. (2007). Hyperspectral remote sensing of plant pigments. *Journal of*

    Experimental Botany, *58*(4), 855–867. https://doi.org/10.1093/jxb/erl123

Brewer, T. (2012). Remote Sensing of Vegetation: Principles, Techniques and Applications. *Soil*

    Use and Management, *28*(1), 134–134. https://doi.org/10.1111/j.1475-2743.2011.00383.x

Broughton, K. J., Payton, P., Tan, D. K. Y., Tissue, D. T., & Bange, M. P. (2021). Effect of

    vapour pressure deficit on gas exchange of field-grown cotton. *Journal of Cotton Research*,

    4(1), 30. https://doi.org/10.1186/s42397-021-00105-4

Buitrago, M. F., Skidmore, A. K., Groen, T. A., & Hecker, C. A. (2018). Connecting infrared

    spectra with plant traits to identify species. *ISPRS Journal of Photogrammetry and Remote*

    Sensing, *139*, 183–200. https://doi.org/10.1016/J.ISPRSJPRS.2018.03.013

Burnett, A. C., Anderson, J., Davidson, K. J., Ely, K. S., Lamour, J., Li, Q., Morrison, B. D.,

    Yang, D., Rogers, A., & Serbin, S. P. (2021). A best-practice guide to predicting plant traits

from leaf-level hyperspectral data using partial least squares regression. *Journal of*

Experimental Botany. https://doi.org/10.1093/jxb/erab295

Burnett, A. C., Serbin, S. P., & Rogers, A. (2021). Source:sink imbalance detected with leaf- and

canopy-level spectroscopy in a field-grown crop. *Plant Cell and Environment*, *44*(8), 2466–

2479. https://doi.org/10.1111/pce.14056

Campbell, G. S., & Norman, J. M. (1998). An Introduction to Environmental Biophysics. In *An*

Introduction to Environmental Biophysics. Springer New York. https://doi.org/10.1007/978-

1-4612-1626-1

Castro-Esau, K. L., Sánchez-Azofeifa, G. A., Rivard, B., Wright, S. J., & Quesada, M. (2006).

Variability in leaf optical properties of mesoamerican trees and the potential for species

classification. *American Journal of Botany*, *93*(4), 517–530.

https://doi.org/10.3732/ajb.93.4.517

Cavender-Bares, J., Gamon, J. A., Hobbie, S. E., Madritch, M. D., Meireles, J. E., Schweiger, A.

K., & Townsend, P. A. (2017). Harnessing plant spectra to integrate the biodiversity

sciences across biological and spatial scales. *American Journal of Botany*, *104*(7), 966–969.

https://doi.org/10.3732/ajb.1700061

Cavender-Bares, J., Meireles, J., Couture, J., Kaproth, M., Kingdon, C., Singh, A., Serbin, S. P.,

Center, A., Zuniga, E., Pilz, G., Townsend, P., Cavender-Bares, J., Meireles, J. E., Couture,

J. J., Kaproth, M. A., Kingdon, C. C., Singh, A., Serbin, S. P., Center, A., … Townsend, P.

A. (2016). Associations of Leaf Spectra with Genetic and Phylogenetic Variation in Oaks:

Prospects for Remote Detection of Biodiversity. *Remote Sensing*, *8*(3), 221.

https://doi.org/10.3390/rs8030221

Cavender-Bares, J., Schneider, F. D., Santos, M. J., Armstrong, A., Carnaval, A., Dahlin, K. M.,

Fatoyinbo, L., Hurtt, G. C., Schimel, D., Townsend, P. A., Ustin, S. L., Wang, Z., &

Wilson, A. M. (2022). Integrating remote sensing with ecology and evolution to advance

biodiversity conservation. *Nature Ecology & Evolution 2022*, 1–14.

https://doi.org/10.1038/s41559-022-01702-5

Chavana-Bryant, C., Malhi, Y., Anastasiou, A., Enquist, B. J., Cosio, E. G., Keenan, T. F., &

Gerard, F. F. (2019). Leaf age effects on the spectral predictability of leaf traits in

Amazonian canopy trees. *Science of the Total Environment*, *666*, 1301–1315.

https://doi.org/10.1016/j.scitotenv.2019.01.379

Cochrane, M. A. (2000). Using vegetation reflectance variability for species level classification

of hyperspectral data. *International Journal of Remote Sensing*, *21*(10), 2075–2087.

https://doi.org/10.1080/01431160050021303

Corripio, J. (2019). *Package "insol" Type Package Title Solar Radiation Version 1.2.1* (p. 36).

Cotrozzi, L., Couture, J. J., Cavender-Bares, J., Kingdon, C. C., Fallon, B., Pilz, G., Pellegrini,

E., Nali, C., & Townsend, P. A. (2017). Using foliar spectral properties to assess the effects

of drought on plant water potential. *Tree Physiology*, *37*(11), 1582–1591.

https://doi.org/10.1093/treephys/tpx106

Curran, P. J. (1989). Remote sensing of foliar chemistry. In *Remote Sensing of Environment*

(Vol. 30, Issue 3, pp. 271–278). Elsevier. https://doi.org/10.1016/0034-4257(89)90069-2

De La Riva, E. G., Olmo, M., Poorter, H., Ubera, J. L., & Villar, R. (2016). Leaf mass per area

(LMA) and its relationship with leaf structure and anatomy in 34 mediterranean woody

species along a water availability gradient. *PLoS ONE*, *11*(2).

https://doi.org/10.1371/journal.pone.0148788

Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, ian J., Lavorel, S., Dray, S., Reu, B., Kleyer,

565 M., Wirth, C., Colin Prentice, I., Garnier, E., Bönisch, G., Westoby, M., Poorter, H., Reich,

566 P. B., Moles, A. T., Dickie, J., Gillison, A. N., Zanne, A. E., … Gorné, L. D. (2016). The

567 global spectrum of plant form and function. *Nature*, *529*(7585), 167–171.

568 https://doi.org/10.1038/nature16489

569 Doyle, J. A. (2007). Systematic value and evolution of leaf architecture across the angiosperms

570 in light of molecular phylogenetic analyses. *CFS Courier Forschungsinstitut Senckenberg*,

571 *258*(258), 21–37. https://www.researchgate.net/publication/280903742

572 Durgante, F. M., Higuchi, N., Almeida, A., & Vicentini, A. (2013). Species Spectral Signature:

573 Discriminating closely related plant species in the Amazon with Near-Infrared Leaf-

574 Spectroscopy. *Forest Ecology and Management*, *291*, 240–248.

575 https://doi.org/10.1016/j.foreco.2012.10.045

576 Ely, K. S., Burnett, A. C., Lieberman-Cribbin, W., Serbin, S. P., & Rogers, A. (2019).

577 Spectroscopy can predict key leaf traits associated with source-sink balance and carbon-

578 nitrogen status. *Journal of Experimental Botany*, *70*(6), 1789–1799.

579 https://doi.org/10.1093/jxb/erz061

580 Féret, J.-B., & Asner, G. P. (2011). *Spectroscopic classification of tropical forest species using*

581 *radiative transfer modeling*. https://doi.org/10.1016/j.rse.2011.05.004

582 Feret, J.-B., François, C., Asner, G. P., Gitelson, A. A., Martin, R. E., Bidel, L. P. R., Ustin, S.

583 L., le Maire, G., & Jacquemoud, S. (2008). PROSPECT-4 and 5: Advances in the leaf

584 optical properties model separating photosynthetic pigments. *Remote Sensing of*

585 *Environment*, *112*(6), 3030–3043. https://doi.org/10.1016/j.rse.2008.02.012

586 Fernando, V., Ernesto, G., & José, M. G. (2007). Ecological limits to plant phenotypic plasticity.

587 *New Phytologist*, *176*(4), 749–763. http://dx.doi.org/10.1111/j.1469-8137.2007.02275.x

Fine, P. V. A., Salazar, D., Martin, R. E., Metz, M. R., Misiewicz, T. M., & Asner, G. P. (2021).

Exploring the links between secondary metabolites and leaf spectral reflectance in a diverse

genus of Amazonian trees. *Ecosphere*, *12*(2), e03362. https://doi.org/10.1002/ecs2.3362

Fourty, T., Baret, F., Jacquemoud, S., Schmuck, G., & Verdebout, J. (1996). Leaf optical

properties with explicit description of its biochemical composition: Direct and inverse

problems. *Remote Sensing of Environment*, *56*(2), 104–117. https://doi.org/10.1016/0034-

4257(95)00234-0

Fyfe, S. K. (2003). Spatial and temporal variation in spectral reflectance: Are seagrass species

spectrally distinct? *Limnology and Oceanography*, *48*(1part2), 464–479.

https://doi.org/10.4319/LO.2003.48.1_PART_2.0464

Gamon, J. A., Kovalchuck, O., Wong, C. Y. S., Harris, A., & Garrity, S. R. (2015). Monitoring

seasonal and diurnal changes in photosynthetic pigments with automated PRI and NDVI

sensors. *Biogeosciences*, *12*, 4149–4159. https://doi.org/10.5194/bg-12-4149-2015

González-Rebeles, G., Terrazas, T., Méndez-Alonzo, R., Paz, H., Brodribb, T. J., & Tinoco-

Ojanguren, C. (2021). Leaf water relations reflect canopy phenology rather than leaf life

span in Sonoran Desert trees. *Tree Physiology*, *41*(9), 1627–1640.

https://doi.org/10.1093/treephys/tpab032

Henrik Singmann, Ben Bolker, Jake Westfall, Frederik Aust, & Mattan S. Ben-Shachar. (2021).

afex: Analysis of Factorial Experiments (p. 77). https://cran.r-project.org/package=afex

Hesketh, M., & Sánchez-Azofeifa, G. A. (2012). The effect of seasonal spectral variation on

species classification in the Panamanian tropical forest. *Remote Sensing of Environment*,

118, 73–82. https://doi.org/10.1016/j.rse.2011.11.005

Higgins, M. A., & Ruokolainen, K. (2004). Rapid Tropical Forest Inventory: a Comparison of

Techniques Based on Inventory Data from Western Amazonia. In *Conservation Biology*

(Vol. 18, Issue 3).

Hooker, W. J. (1837). Macleania Hook. *Icones Plantarum*, *2*, 109.

Jacquemoud, S., & Baret, F. (1990). PROSPECT: A model of leaf optical properties spectra.

Remote Sensing of Environment, *34*(2), 75–91. https://doi.org/10.1016/0034-

4257(90)90100-Z

Jacquemoud, S., Ustin, S. L., Verdebout, J., Schmuck, G., Andreoli, G., & Hosgood, B. (1996).

Estimating leaf biochemistry using the PROSPECT leaf optical properties model. *Remote*

Sensing of Environment, *56*(3), 194–202. https://doi.org/10.1016/0034-4257(95)00238-3

Jin, Y., & Qian, H. (2019). V.PhyloMaker: an R package that can generate very large

phylogenies for vascular plants. *Ecography*, *42*(8), 1353–1359.

https://doi.org/10.1111/ecog.04434

Kassambara, A., & Mundt, F. (2020). Package 'factoextra': Extract and visualize the results of

multivariate data analyses. *CRAN- R Package*, 84.

https://github.com/kassambara/factoextra/issues

Kuhn, M., & Johnson, K. (2013). Applied predictive modeling. In *Applied Predictive Modeling*.

Springer New York. https://doi.org/10.1007/978-1-4614-6849-3

Kupper, P., Sõber, J., Sellin, A., Lõhmus, K., Tullus, A., Räim, O., Lubenets, K., Tulva, I., Uri,

V., Zobel, M., Kull, O., & Sõber, A. (2011). An experimental facility for free air humidity

manipulation (FAHM) can alter water flux through deciduous tree canopy. *Environmental*

and Experimental Botany, *72*, 432–438. https://doi.org/10.1016/j.envexpbot.2010.09.003

Landis, J. R., & Koch, G. G. (1977). The Measurement of Observer Agreement for Categorical

Data. *Biometrics*, *33*(1), 159. https://doi.org/10.2307/2529310

Lang, C., Almeida, D. R. A., & Costa, F. R. C. (2017). Discrimination of taxonomic identity at

species, genus and family levels using Fourier Transformed Near-Infrared Spectroscopy

(FT-NIR). *Forest Ecology and Management*, *406*(June), 219–227.

https://doi.org/10.1016/j.foreco.2017.09.003

Lang, C., Costa, F. R. C., Camargo, J. L. C., Durgante, F. M., & Vicentini, A. (2015). Near

Infrared Spectroscopy Facilitates Rapid Identification of Both Young and Mature

Amazonian Tree Species. *PLOS ONE*, *10*(8), e0134521.

https://doi.org/10.1371/journal.pone.0134521

Li, Y., He, N., Hou, J., Xu, L., Liu, C., Zhang, J., Wang, Q., Zhang, X., & Wu, X. (2018).

Factors influencing leaf chlorophyll content in natural forests at the biome scale. *Frontiers*

in Ecology and Evolution, *6*(JUN), 64. https://doi.org/10.3389/fevo.2018.00064

Little, S. A., Kembel, S. W., & Wilf, P. (2010). Paleotemperature Proxies from Leaf Fossils

Reinterpreted in Light of Evolutionary History. *PLoS ONE*, *5*(12), 15161.

https://doi.org/10.1371/journal.pone.0015161

Marcelo-Peña, J. L., Huamantupa, I., Särkinen, T., & Tomazello, M. (2015). Identifying

conservation priority areas in the marañón valley (Peru) based on floristic inventories.

Edinburgh Journal of Botany, *73*(1), 95–123. https://doi.org/10.1017/S0960428615000281

Meireles, J. E., Cavender-Bares, J., Townsend, P. A., Ustin, S., Gamon, J. A., Schweiger, A. K.,

Schaepman, M. E., Asner, G. P., Martin, R. E., Singh, A., Schrodt, F., Chlus, A., &

O'Meara, B. C. (2020). Leaf reflectance spectra capture the evolutionary history of seed

plants. *New Phytologist*, *228*(2), 485–493. https://doi.org/10.1111/nph.16771

Meletiou-Christou, M. S., Rhizopoulou, S., & Diamantoglou, S. (1994). Seasonal changes of

carbohydrates, lipids and nitrogen content in sun and shade leaves from four mediterranean

evergreen sclerophylls. *Environmental and Experimental Botany*, *34*(2), 129–140.

https://doi.org/10.1016/0098-8472(94)90032-9

Muller, O., Hirose, T., Werger, M. J. A., & Hikosaka, K. (2011). Optimal use of leaf nitrogen

explains seasonal changes in leaf nitrogen content of an understorey evergreen shrub.

Annals of Botany, *108*(3), 529–536. https://doi.org/10.1093/aob/mcr167

Münkemüller, T., Lavergne, S., Bzeznik, B., Dray, S., Jombart, T., Schiffers, K., & Thuiller, W.

(2012). How to measure and test phylogenetic signal. *Methods in Ecology and Evolution*,

3(4), 743–756. https://doi.org/10.1111/j.2041-210X.2012.00196.x

Nakaji, T., Oguma, H., Nakamura, M., Kachina, P., Asanok, L., Marod, D., Aiba, M., Kurokawa,

H., Kosugi, Y., Kassim, A. R., & Hiura, T. (2019). Estimation of six leaf traits of East

Asian forest tree species by leaf spectroscopy and partial least square regression. *Remote*

Sensing of Environment, *233*, 111381. https://doi.org/10.1016/j.rse.2019.111381

Nguyen, H. T., Meir, P., Sack, L., Evans, J. R., Oliveira, R. S., & Ball, M. C. (2017). Leaf water

storage increases with salinity and aridity in the mangrove Avicennia marina: integration of

leaf structure, osmotic adjustment and access to multiple water sources. *Plant Cell and*

Environment, *40*(8), 1576–1591. https://doi.org/10.1111/pce.12962

Noss, R. F. (2001). Beyond Kyoto: Forest Management in a Time of Rapid Climate Change.

Conservation Biology, *15*(3), 578–590. https://doi.org/10.1046/J.1523-

1739.2001.015003578.X

Nunes, M. H., Both, S., Bongalov, B., Brelsford, C., Khoury, S., Burslem, D. F. R. P., Philipson,

C., Majalap, N., Riutta, T., Coomes, D. A., & Cutler, M. E. J. (2019). Changes in leaf

functional traits of rainforest canopy trees associated with an El Nino event in Borneo.

Environmental Research Letters, *14*(8). https://doi.org/10.1088/1748-9326/ab2eae

Onoda, Y., Wright, I. J., Evans, J. R., Hikosaka, K., Kitajima, K., Niinemets, Ü., Poorter, H.,

Tosens, T., & Westoby, M. (2017). Physiological and structural tradeoffs underlying the

leaf economics spectrum. *New Phytologist*, *214*(4), 1447–1463.

https://doi.org/10.1111/nph.14496

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, *401*(6756),

877–884.

Paiva, D. N. A., Perdiz, R. de O., & Almeida, T. E. (2021). Using near-infrared spectroscopy to

discriminate closely related species: a case study of neotropical ferns. *Journal of Plant*

Research, *134*(3), 509–520. https://doi.org/10.1007/s10265-021-01265-9

Paz-Kagan, T., & Asner, G. P. (2017). Drivers of woody canopy water content responses to

drought in a Mediterranean-type ecosystem: *Ecological Applications*, *27*(7), 2220–2233.

https://doi.org/10.1002/eap.1603

Price, J. C. (1994). How unique are spectral signatures? *Remote Sensing of Environment*, *49*(3),

181–186. https://doi.org/10.1016/0034-4257(94)90013-2

Qian, S. E. (2021). Hyperspectral Satellites, Evolution, and Development History. In *IEEE*

Journal of Selected Topics in Applied Earth Observations and Remote Sensing (Vol. 14, pp.

7032–7056). https://doi.org/10.1109/JSTARS.2021.3090256

Reich, P. B., Uhl, C., Walters, M. B., & Ellsworth, D. S. (1991). Leaf lifespan as a determinant

of leaf structure and function among 23 amazonian tree species. *Oecologia*, *86*(1), 16–24.

https://doi.org/10.1007/BF00317383

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution*, *3*(2), 217–223. https://doi.org/10.1111/j.2041-

210X.2011.00169.x

Rodríguez-Pérez, R., Fernández, L., & Marco, S. (2018). Overoptimism in cross-validation when

using partial least squares-discriminant analysis for omics data: a systematic study.

Analytical and Bioanalytical Chemistry, *410*(23), 5981–5992.

https://doi.org/10.1007/s00216-018-1217-1

Schielzeth, H., Dingemanse, N. J., Nakagawa, S., Westneat, D. F., Allegue, H., Teplitsky, C.,

Réale, D., Dochtermann, N. A., Garamszegi, L. Z., & Araya-Ajoy, Y. G. (2020).

Robustness of linear mixed-effects models to violations of distributional assumptions.

Methods in Ecology and Evolution, *11*(9), 1141–1152. https://doi.org/10.1111/2041-

210X.13434

Schmitt, S., Tysklind, N., Derroire, G., Heuertz, M., & Hérault, B. (2021). Topography shapes

the local coexistence of tree species within species complexes of Neotropical forests.

Oecologia, 1–10. https://doi.org/10.1007/s00442-021-04939-2

Schreier, M. (2012). Qualitative Content Analysis in Practice. In *SAGE Publications*.

Schweiger, A. K., Cavender-Bares, J., Townsend, P. A., Hobbie, S. E., Madritch, M. D., Wang,

R., Tilman, D., & Gamon, J. A. (2018). Plant spectral diversity integrates functional and

phylogenetic components of biodiversity and predicts ecosystem function. *Nature Ecology*

& Evolution, *2*(6), 976–982. https://doi.org/10.1038/s41559-018-0551-1

Schweiger, A. K., Schütz, M., Risch, A. C., Kneubühler, M., Haller, R., & Schaepman, M. E.

(2017). How to predict plant functional types using imaging spectroscopy: linking

vegetation community traits, plant functional types and spectral response. *Methods in*

Ecology and Evolution, *8*(1), 86–95. https://doi.org/10.1111/2041-210X.12642

Serbin, S. P., Singh, A., Mcneil, B. E., Kingdon, C. C., & Townsend, P. A. (2014). Spectroscopic

determination of leaf morphological and biochemical traits for northern temperate and

boreal tree species. *Ecological Applications*, *24*(7), 1651–1669.

Serbin, S. P., Wu, J., Ely, K. S., Kruger, E. L., Townsend, P. A., Meng, R., Wolfe, B. T., Chlus,

A., Wang, Z., & Rogers, A. (2019). From the Arctic to the tropics: multibiome prediction of

leaf mass per area using leaf reflectance. *New Phytologist*, *224*(4), 1557–1568.

https://doi.org/10.1111/nph.16123

Shiklomanov, A. N., Dietze, M. C., Viskari, T., Townsend, P. A., & Serbin, S. P. (2016).

Quantifying the influences of spectral resolution on uncertainty in leaf trait estimates

through a Bayesian approach to RTM inversion. *Remote Sensing of Environment*, *183*, 226–

238. https://doi.org/10.1016/j.rse.2016.05.023

Stasinski, L., White, D. M., Nelson, P. R., Ree, R. H., & Meireles, J. E. (2021). Reading light:

leaf spectra capture fine-scale diversity of closely related, hybridizing arctic shrubs. *New*

Phytologist, *232*(6), 2283–2294. https://doi.org/10.1111/nph.17731

Stevens, P. F. (2012). Angiosperm Phylogeny Website. *Version 12, July 2012*.

Stuessy, T. F., Crawford, D. J., Soltis, D. E., & Soltis, P. S. (2014). *Plant Systematics; The*

origin, interpretation, and ordering of plant biodiversity.

Tamási, B., & Hothorn, T. (2021). tramME: Mixed-Effects Transformation Models Using

Template Model Builder. *R Journal*, *13*(2), 398–418. https://doi.org/10.32614/RJ-2021-075

Trias-Blasi, A., & Vorontsova, M. (2015). Plant identification is key to conservation. *Nature*

2015 521:7551, *521*(7551), 161–161. https://doi.org/10.1038/521161c

Ustin, S. L., Gitelson, A. A., Jacquemoud, S., Schaepman, M., Asner, G. P., Gamon, J. A., &

Zarco-Tejada, P. (2009). Retrieval of foliar information about plant pigment systems from

high resolution spectroscopy. *Remote Sensing of Environment*, *113*, S67–S77.

https://doi.org/10.1016/j.rse.2008.10.019

Van Goethem, D., Smedt, D., Valcke, S., Potters, R., & Samson, G. (2013). Seasonal, Diurnal

and Vertical Variation of Chlorophyll Fluorescence on Phyllostachys humilis in Ireland.

PLoS ONE, *8*(8), 72145. https://doi.org/10.1371/journal.pone.0072145

Yang, X., Tang, J., Mustard, J. F., Wu, J., Zhao, K., Serbin, S., & Lee, J. E. (2016). Seasonal

variability of multiple leaf traits captured by leaf spectroscopy at two temperate deciduous

forests. *Remote Sensing of Environment*, *179*, 1–12.

https://doi.org/10.1016/j.rse.2016.03.026

Young, A. J. (1991). The photoprotective role of carotenoids in higher plants. *Physiologia*

Plantarum, *83*(4), 702–708. https://doi.org/10.1111/j.1399-3054.1991.tb02490.x