New insights into the role of the microbiota in *Ornithodoros moubata*: Influence of endosymbionts on the reproductive fitness of the tick

Taraveau Florian[1], Pollet Thomas[1], Duhayon Maxime[1], Gardès Laëtitia[1], Jourdan-Pineau Hélène*[1]

[1] UMR ASTRE, CIRAD, INRAE, Campus de Baillarguet, F-34398 Montpellier, France

*Corresponding author
Correspondence: helene.jourdan@cirad.fr
First author correspondence: florian.taraveau@cirad.fr

ABSTRACT

Over the past ten years, many studies demonstrated the crucial role of the tick microbiome in tick biology. The soft tick *Ornithodoros moubata* is a hematophagous ectoparasite of *Suidae* particularly known to transmit the *African swine fever virus*. Its bacterial microbiota is characterized by a high prevalence of *Francisella*-like and *Rickettsia* endosymbionts. The present study aims to better understand the potential influence of the microbiota on the reproductive fitness of *O. moubata*. A total of 132 adult female ticks were treated using gentamycin or rifampicin added to the blood meal. Half of the ticks also received a supplementation with B vitamins to address the nutritional role of endosymbionts. Over two periods of 50 days, several traits related to the reproductive fitness were monitored to investigate the importance of *Francisella* and *Rickettsia* for those traits. It appeared that most of the considered reproductive parameters were not affected. However, antibiotic treatments induced an increase in the tick survival indicating a potential fitness cost of harboring endosymbionts for ticks during the reproduction period. Similarly, 366 first stage nymphs of *Ornithodoros moubata* were exposed to the same treatments for molecular quantification of both endosymbionts. Results from qPCR suggested that treatments produced a bacteriostatic effect on endosymbionts without fully eliminating *Francisella* or *Rickettsia*.

Keywords: Soft tick, *Ornithodoros*, microbiota, antibiotic treatment, *Francisella*-like endosymbiont, *Rickettsia*

Introduction

While ticks are among the most important vectors of pathogens affecting humans and animals worldwide, it is now well established that tick-borne pathogens (TBPs) coexist with many other microorganisms in ticks (e.g., Bonnet *et al*. 2017; Estrada-Peña *et al*. 2018). These other tick-borne microbes (microbiota) are likely to confer multiple detrimental, neutral, or beneficial effects to their tick hosts (Duron *et al*. 2018; Duron, Cremaschi, et McCoy 2016). In addition, many studies identified multiple interactions between some members of the tick microbiota and TBPs (e.g., Lejal *et al*. 2021; Aivelo, Norberg, et Tschirren 2019). For instance, these tick microbial communities can play a role in the transmission or multiplication of TBPs (Vayssier-Taussat *et al*. 2014; Bonnet *et al*. 2017; Budachetri *et al*. 2018; Bonnet et Pollet 2021) which question about the future possibility to manipulate tick microbiomes and use them as a new tool in tick and tick-borne disease control.

If the precise role of all tick-borne microbial species remains unclear, they can be broadly classified as environmental or endosymbiotic. Those later intracellular bacteria are further categorized into primary and secondary endosymbionts according to their importance for the host (Jiménez-Cortés *et al*. 2018), although intermediate forms are acknowledged. Primary symbionts are heritable obligate symbionts and represent crucial members of the microbiota for the survival or the development of the arthropod (Bonnet *et al*. 2017). In contrast, secondary endosymbionts are not required for host survival and are more sporadically associated with host individuals. Accordingly, four main bacterial genera are currently well identified as tick primary endosymbionts: *Coxiella*-like endosymbionts (CLE), *Candidatus Midichloria mitochondrii*, *Francisella*-like endosymbionts (FLE), and, less often, *Rickettsia*. All of these endosymbionts were identified to carry a copy of genes involved in essential nutrients synthesis, including B vitamins which are not naturally brought by the strictly hematophagous alimentation. Depending on the tick species, one or two of these genera will be found predominantly and will assume the nutritional function (Duron et Gottlieb 2020; Bonnet et Pollet 2021). For example, this role is assumed by *Coxiella*-like endosymbionts (CLE) in *Amblyomma americanum* whereas it is assumed by *Rickettsia* in *Ixodes pacificus* (Rio, Attardo, et Weiss 2016). Over time, several evolutionary events of loss and/or new endosymbiosis are suspected, leading to the diversity of endosymbionts currently found (Duron *et al*. 2017; Duron et Gottlieb 2020).

However, the role of primary endosymbionts is not limited to nutrient supply. In ticks, they are also known to have an influence on fitness, reproduction (Guizzo *et al*. 2017; Li, Zhang, et Zhu 2018; Zhang *et al*. 2017; Zhong, Jasinskas, et Barbour 2007; Kurlovs *et al*. 2014), immunity, and vector competence (Narasimhan *et al*. 2014; Bonnet et Pollet 2021; Gall *et al*. 2016). Besides primary endosymbionts, many other members of the microbiota revealed by high-throughput sequencing are still poorly known and their effects on tick biology are not elucidated yet.

The life cycle of soft ticks (several nymphal stages, short blood feedings, adult females feeding and laying eggs several times before dying) makes these ticks an ideal model for studying the role of the tick microbiota on tick biology and physiology. *Ornithodoros moubata* is a soft tick widely distributed in Southern and Eastern Africa (Vial 2009; Leeson 1952). Its main hosts include wild and domestic members of the *Suidae* family: warthog, bushpig and domestic pig (Peirce 1974). This tick species is mainly known as a reservoir host for the *African swine fever virus* (ASFV) and is consequently an important concern for animal health issues (Sánchez-Vizcaíno *et al*. 2015).

High-throughput 16S rDNA sequencing in *O. moubata* showed that the most abundant bacterial species belonged to the genus *Francisella* and *Rickettsia* (Duron *et al*. 2018). Both bacteria are recognized as maternally-inherited endosymbionts in ticks (Duron *et al*. 2017). Furthermore, *Francisella*-like endosymbionts (FLE) have been shown to play a key role in the production of B vitamins, including biotin, folic acid and riboflavin (Duron *et al*. 2018).

Apart from their role in tick nutrition, the influence of *Francisella* and *Rickettsia* in the physiology and life history traits of soft ticks remains poorly known. In comparison, past studies suggest a role of CLE and *Rickettsia* for the reproductive fitness of female hard ticks belonging to the genera *Ixodes* (Kurlovs *et al*. 2014), *Rhipicephalus* (Li, Zhang, et Zhu 2018)*, Haemaphysalis* (Zhang *et al*. 2017)*,* and *Amblyomma* (Zhong, Jasinskas, et Barbour 2007). In those species, the reproductive fitness (number of eggs laid, mass of eggs, duration of oviposition and hatching rate) was negatively impacted by the elimination of parts of the microbiota.

The present study aimed to better characterize the role of the microbiota in reproductive fitness in soft ticks. Accordingly, our first objective was to eliminate the main bacterial species of the microbiota of

88 *Ornithodoros moubata* with antibiotic treatments. The second goal was to evaluate the consequences of such
89 a disruption on the reproductive fitness of ticks, using B vitamin supplementation to identify which part of the
90 observed changes was directly attributable to an action of the endosymbionts on the reproduction and which
91 part was a consequence of their nutritional role.

Material and methods

Ticks

All ticks are issued from a colony maintained at the CIRAD laboratory (Montpellier, France, member of the
Vectopole Sud network) since 2008. This colony originated from the Neuchâtel strain initially collected in
Tanzania and maintained at the University of Neuchâtel (Switzerland).
The female ticks used for the experiments had never been mated before and did not receive any blood
meal since molting from the last nymphal stage to the adult stage.
Adult ticks, nymphs, larvae, and eggs were kept in stable conditions in one incubator, at 25°C, 85% relative
humidity, and 0% luminosity.

Feeding and antibiotic treatment

Adult stages were engorged with a first blood meal at the beginning of the experiment and a second one
52 days later. Blood meals were performed using an artificial feeding system with fresh cow blood, collected
1-3 hours before the experiment in 10mL heparinized tubes. All the blood used for this experiment was issued
from the same cow. Blood meals lasted three hours in complete darkness while blood temperature was
maintained at 41°C and stirred at 250 rpm.
Antibiotics and B vitamins were added in the blood for each group following the protocol of Duron *et al.*
(Duron *et al*. 2018) except that dilutions were performed in Milli-Q water instead of methanol due to the
potential toxicity of methanol for arthropods (Kaviraj, Bhunia, et Saha 2004). Concentrations were chosen
according to recommendations for sensitive bacterial species and previous studies in the literature:
gentamycin 10 μg.mL^{-1}, rifampicin 10 μg.mL^{-1}, biotin 1 μg.mL^{-1}, folic acid 30 μg.mL^{-1}, riboflavin 20 μg.mL^{-1}
(Duron *et al*. 2018; Zhong, Jasinskas, et Barbour 2007; Narasimhan *et al*. 2014). Antibiotics and B vitamins were
replaced by the same volumes of Milli-Q water for negative controls.
Each tick was weighted before and after the blood meal. To ensure efficient blood meal, ticks were fed in
group: four to seven females were fed together with 3 males in one well containing 9 mL of blood. Therefore,
we had to mark ticks individually with different colors using Posca felt pens. If ticks gained less than 50 mg at
the end of the first blood meal, the blood feeding was considered as a failure and ticks were fed again during
a second, and, if needed, a third blood meal. If after three attempts some ticks were still not fed, they were
excluded from the study.
Feeding success was not affected by the antibiotic treatment (first 50 days: p=0.5054 ; second 50 days:
p=0.5343) nor by vitamin treatment (first 50 days: p=0.1553 ; second 50 days: p=0.0892). Similarly, tick weight
after bloodmeal was not affected by the antibiotic treatment (first 50 days: p=0.9325 ; second 50 days:
p=0.9050) nor by vitamin treatment (first 50 days: p=0.5556; second 50 days: p=0.9835).
124
Independently from the feeding of adults, nymphal stages were engorged by groups of thirty individuals.
Treatment groups were the same as the one used for adult ticks (control, rifampicin, gentamycin, with or
without B vitamins) with a total of 60 nymphs for each condition. A total of 20 individuals were sampled for
molecular analyses at day 0 (before blood meal), and 10 individuals at day 1, day 3, day 5, day 10, day 15 and
day 30 after blood meal (**Figure 1**). Nymphs which died before DNA extraction were excluded. For sample size
and group size, see supplementary data (**Table S1**).

Figure 1: Antibiotic treatments performed, followed by the evaluation of traits of interest at different stages of the parasitic cycle of *Ornithodoros moubata*. The traits monitored are indicated near the parasitic stage concerned. In blue, experiment conducted on adult ticks to monitor reproductive parameters. In green, experiment conducted on first stage nymphs to monitor bacterial loads after antibiotic treatment. Tick pictures: Françoise Chirara © Cirad-ASTRE.

Reproduction monitoring

After blood meal, females were kept in the same tubes as males for seven days to allow mating, then each female was isolated in an individual plastic tube. The monitoring of the tick reproduction was blinded regarding applied treatments. A total of 132 females were included in the study and monitored every day during two successive 50-days periods after blood meal. For each tick, the laying date, the number of eggs laid and the egg mass were recorded. The number of eggs laid was standardized with the weight of the female tick after blood meal. For each egg, the date of hatching and of molting into first nymphal stage were recorded (**Figure 1**). At the end of the experiment, the remaining females were stored at -80°C for molecular analyses.

DNA extraction

Nymphs and adults were washed 30 seconds in 1% bleach solution and then rinsed for 1 minute in three successive baths of Milli-Q water to eliminate cuticular bacteria (Binetruy *et al*. 2019). Ticks were then crushed individually using a Qiagen TissueLyser instrument with steel and plastic beads in 300 µL of DMEM medium complemented at 10% with calf serum. DNA was extracted from 75 µL of each crushed tick homogenate, using the genomic DNA extraction kit NucleoSpin® Tissue (Macherey-Nagel, Hoerdt, Germany), and following the standard protocol for human or animal tissue and cultured cells. DNA extracts were finally eluted in 50 µl of elution buffer and stored at – 20 C until further use.

Quantitative PCR

Quantitative PCRs were launched using Roche Lightcycler® 96 instruments (Roche Applied Sciences, Beijing, China) using the taqMan method.

Extracted DNA (2 µL) was added to the amplification mix (18 µL: Mix 10 µL, H2O 6.6 µL, forward Primer
Actin 0.6 µM, reverse Primer Actin 0.6 µM, Probe Actin 0.2 µM, forward Primer target gene 0.6 µM, reverse
Primer target gene 0.6 µM, Probe target gene 0.2 µM). Primers used to amplify *Francisella* targeted the *RpoB*
gene coding for the B subunit of RNA polymerase. Primer sequences were Forward: 5'-
GTGGTGTACCTATTGCTACG-3',   Reverse: 5'-TGATGGTCGTACTGGTAAGC-3'   and   Probe: 5'-
ATAGGTGGCTTTGCAACTAA-3'. Primers used to amplify *Rickettsia* targeted the *GltA* gene coding for the citrate
synthase enzyme. Primer sequences were Forward: 5'-AATGATCCGTTTAGGTTGATG-3', Reverse: 5'-
ACCCACTCCTGCAAATAG-3' and Probe: 5'-GTGCCGCAGTACTTAAAGA-3'. The actin gene from *Ornithodoros*
moubata was also amplified as a control for the assessment of DNA extraction success and to standardize the
results with the tick size. Primer sequences were Forward: 5'-CCGGTATTGCCGACCGTATGC-3', Reverse: 5'-
CTCCCTGTCCACCTTCCAGC-3' and Probe: 5'-CGAGAGGAAGTACTCCGTCTGG-3' (Pereira De Oliveira, Vial, et Le
Potier 2022; Duron *et al.* 2018).
Positive and negative controls were included, using 2 µL of DNA of tick previously tested positive for *FLE*
and *Rickettsia* (Duron *et al.* 2018) and 2 µL of Milli-Q water respectively. The PCR cycling program was 1 min
at 95°C, then 45 cycles of 1 min at 95°C, 30 sec at 56°C. Analyses were performed using both FAM and HEX
fluorescence indicators. Samples with a cycle threshold (Ct) beyond 38 were considered as negative. Cts were
converted into concentrations from the standard curves performed with dilutions of a DNA plasmid containing
the targeted sequences. Results are presented as ratio of the concentration of the gene of interest (GltA or
RpoB) divided by the concentration of the actin gene to standardize for the tick size.

Statistical analysis

Statistical analyses were performed using R version 4.2.3 (2023-03-15) to test the effect of antibiotic and
vitamin treatments on DNA concentrations and traits associated with reproductive fitness (R Core Team 2023).
For the analyses of qPCR results, ratios of DNA concentrations were log-transformed and generalized least-
squares (GLS) models were used with the function **gls** to allow different variances for each level of the
explanatory variables (Zuur *et al.* 2009). The best model including optimal variance structure and fixed effects
(namely, antibiotic and vitamin treatments and their interaction) was determined by AIC comparisons. In
nymphs, due to material constraints, two different experimenters performed the qPCR at two different
moments. This was added as a fixed effect in all the models and had a clear effect on measured ratios (GltA:
$p=1.602e^{-05}$, F-value=19.17 and, RpoB: $p=1.144e^{-65}$, F-value=472.7). DNA concentrations were also log-
transformed and analyzed using **glm** with a gamma distribution when ratios were not relevant for the analyzes.
To analyze reproduction parameters, we used generalized linear models with the function **lm**, **glm** or
generalized mixed model with **glmer** (R package **Lme4** (Bates *et al.* 2015)) according to the distribution of the
data and the absence (**glm**) or presence (**glmer**) of a random effect. The random effect was added for analyses
targeting the offspring of tick females to account for the lack of independence of data points (since measured
individuals may be offspring of the same mother). For egg incubation time and time for molting into first
nymphal stage, the level of variability between mother was not sufficient to warrant incorporating it as a
random effect. Depending on the variable, normal, Poisson or binomial distributions were used (see **Table 1**).
Overdispersion and underdispersion of the data were estimated using a quasi-Poisson or quasi-binomial
distribution. Statistical comparisons were performed using either F-test, chi-squared test or Kruskal-Wallis test.
When the mean was presented for a trait, standard deviation (SD) was indicated in brackets. In addition, data
from the two 50-days periods were statistically compared and analyzed using the models described above.
Interactions with antibiotic treatments and B vitamin supplementation were taken into account in the analysis.

Ethics

The use of cows which provided blood for tick engorgement complied with the ethical standards of
European regulations (directive from the European Council from November 24th, 1986 (86/609 / EEC)) and
French legislation (ethic committee CEEA-LR-36 available on the APAFIS platform) on the care and use of
animals. More precisely, blood sampling has been recorded in 2021 under the agreement number
APAFIS#27628-2020101010475151 for the CIRAD-Montpellier rearing facility G 34-172-15.

Results

Molecular results

A total of 366 first stage nymphs were tested for *Rickettsia* (target gene *GltA*) and *Francisella*-like
endosymbiont (target gene *RpoB*).

Out of 366 samples, 98.6% were tested positive for *Rickettsia* (359/364) while two samples were excluded
due to the abnormal qPCR curves. Two nymphs were tested negative in the control group without vitamins (at
day 3 and day 5) and three others in the gentamycin group (day 30 with and without vitamins and day 10
without vitamins). The concentration of *GltA* DNA increased over time after blood meal in the control
(p=0.003528, χ_1^2=22.50) and gentamycin (p=4.74e^{-11}, χ_1^2=59.70) groups but not in the rifampicin group
(p=0.2616, χ_1^2 =2.821). Similar increase also appeared for the actin gene in all groups (p<2.2e^{-16}, χ_1^2=115.5).
After standardization, the ratio of DNA concentration remained significantly lower in the rifampicin group than
in the two other groups (p=1.624e^{-92}, F-value=429.0), independently from vitamin supplementation (**Figure 2A**
and 2B).

Figure 2: Boxplots of the ratios of DNA concentration obtained by qPCR for *Rickettsia* and *Francisella*-like
endosymbiont (FLE) after antibiotic treatment of individual first stage nymphs. (A) Ratio of the concentration of
GltA DNA ([GltA]), indicator of the presence of *Rickettsia* over the concentration of *O. moubata* actin DNA [Actin].
Nymphs supplemented with B vitamins. **(B)** Ratio of [GltA] over [Actin]. Nymphs not supplemented with B vitamins.
(C) Ratio of the concentration of *RpoB* DNA ([RpoB]), indicator of the presence of FLE over [Actin]. Nymphs
supplemented with B vitamins **(D)** Ratio of [RpoB] over [Actin]. Nymphs not supplemented with B vitamins.

Regarding *Francisella*-like endosymbiont, 98.1% of the qPCR were positive (359/366) with one negative
samples at day 0, two in the rifampicin group (day 1 no vitamin, and day 10 with vitamin) and four in the
control group (day 5 and 10 with vitamin, and day 20 and 30 no vitamin). As for *Rickettsia*, there was an
increase in the quantity of *RpoB* DNA detected over time in the gentamycin group (p=0.003095, χ_1^2=23.84) but
not in the control (p=0.08688, χ_1^2=3.190) and rifampicin (p=0.1337, χ_1^2=5.103) groups. After standardization,

the ratio of DNA concentration remained significantly different between the three treatment groups
(p=1.709e^{-12}, F-value=29.42) (**Figure 2C and 2D**).
231
At the end of the reproduction monitoring, a total of 59 females were still alive and were collected for
quantification of *Rickettsia* and *Francisella*-like endosymbionts. Forty-eight females were tested positive for
Rickettsia (14/14 control, 2/6 gentamycin without vitamin, 11/12 gentamycin with vitamin, 10/14 rifampicin
without vitamin, and 11/13 rifampicin with vitamin), all females were tested positive for *Francisella*-like
endosymbiont, and five adults were excluded due to abnormal qPCR curves. Comparing the ratio of
concentration of symbionts, no significant differences were observed between the groups concerning vitamin
supplementation (p>0.05). However, focusing on the different antibiotic treatment groups, it appeared that
ticks treated with rifampicin or gentamycin hosted lower concentrations of *Rickettsia* (p=1.805e^{-06}, F-
value=19.11) and *Francisella*-like endosymbiont (p=1.123e^{-09}, F-value= 31.72) than ticks from the control
group. This effect was stronger in the rifampicin group than in the gentamycin group (**Figure 3A and 3B**).
242

Figure 3: Boxplots of the DNA concentration obtained by qPCR for *Rickettsia* and *Francisella*-like endosymbiont
(FLE) after antibiotic treatment of female ticks. (A) Ratio of the concentration of *GltA* DNA ([GltA]), indicator of the
presence of *Rickettsia* over the concentration of *O. moubata* actin DNA [Actin]. **(B)** Ratio of the concentration of
RpoB DNA ([RpoB]), indicator of the presence of FLE over [Actin].

Monitoring of the reproduction

Adult ticks were separated in three treatment groups with two subgroups in each (half with B vitamin
supplementation and half without). Effects of antibiotic treatment, vitamin supplementation and the
interaction between the two were evaluated for statistical differences on traits associated with reproductive
fitness over two successive periods of 50 days (**Table 1 and 2**).
252

Table 1: Statistical analyses performed on traits associated with reproductive fitness measured in female ticks during 50 days after the first antibiotic treatment and vitamin supplementation.

Variable	Family	Effect of the treatment	Effect of the vitamin	Effect of Interaction
Survival over 50 days after blood meal	Binomial	Pr: 0.009687** Dev: 9.274	Pr: 0.3768 Dev: 0.7810	Pr: 0.1867 Dev: 3.356
Laying probability	Binomial	Pr: 0.3499 Dev: 2.100	Pr: 0.1073 Dev: 2.594	Pr: 0.4483 Dev: 1.605
Number of days to first oviposition	Quasi-Poisson[1]	Pr: 0.3741 Dev: 3.548	Pr: 0.05820 Dev: 6.472	Pr: 0.1609 Dev: 6.590
Number of eggs standardized by mother weight[3]	Quasi-Poisson[1]	Pr: 0.4116 Dev: 18.70	Pr: 0.7698 Dev: 0.9018	Pr: 0.9174 Dev: 1.816
Average egg mass	Normal	Pr: 0.3529 Dev: 0.002267	Pr: 0.1437 Dev: 0.004680	Pr: 0.8892 Dev: $2.531e^{-04}$
Egg incubation time	Quasi-Poisson[2]	Pr: 0.6958 Dev: 0.06174	Pr: 0.7652 Dev: 0.007590	Pr: 0.05697 Dev: 0.4878
Egg hatching rate	Binomial	Pr: 0.7476 Dev: 0.5817	Pr: 0.9567 Dev: 0.002947	Pr: 0.5500 Dev: 1.196
Time for molting into first nymphal stage	Quasi-Poisson[2]	Pr: 0.7098 Dev: 0.04697	Pr: 0.3924 Dev: 0.05011	Pr: 0.7444 Dev: 0.04045
Molting rate	Binomial	Pr: 0.8713 Dev: 0.2756	Pr: 0.1844 Dev: 1.762	Pr: 0.7626 Dev: 0.5419

Note: Pr: P-value. Dev: Deviance, X^2 for binomial and quasi-Poisson distribution, F-value for normal distribution. Degree of freedom was 2 for the effect of the treatment and the effect of the interaction and 1 for the effect of vitamin. [1] Distribution corrected for overdispersion of the data. [2] Distribution corrected for underdispersion of the data. [3] Variable corrected using an offset of the tick weight. Statistically significant results are highlighted with a grey background. Statistical difference code: 0 '***' 0.001 '**' 0.01 '*' 0.05 ' ' 1

In total, 78.8% (104/132) of female ticks survived throughout the experiment. Survival was significantly different between the three treatment groups but not between the vitamin subgroups (**Figure 4A**, **Table 1**), with the control group presenting a lower survival rate (65.2%) than both the gentamycin (90.9%) and rifampicin (81.0%) groups.

The oviposition probability was not significantly different between treatment groups (**Figure 4B, Table 1**). Out of 132 ticks engorged, 74.2% (98/132) started oviposition during the first period of 50 days. Those ticks laid between 2 and 220 eggs, with an average of 109 eggs, over the first 50 days of monitoring (or, after standardization, 0.361 egg per mg of weight of the tick [SD: 0.116]). No differences between any subgroups were found for this trait (**Figure 4D, Table 1**). The mean time between blood meal and first day of oviposition was not affected by treatments with a mean of 23.9 days [SD: 6.81] (**Figure 4C**, **Table 1**). Similarly, no differences were detected for the mean egg mass (**Figure 4E, Table 1**). On average, an egg weighted 0.538 mg [SD: 0.0463].

No statistical differences were found between the treatments neither for the egg incubation time, nor for the egg hatching rate (**Figure 4F and 4G**, **Table 1**). Mean values for hatching parameters were 11.9 days of incubation [SD: 1.01] and 79.6% of hatching success [SD: 15.1].

Figure 4: Histograms and boxplots of the traits associated with reproductive fitness measured in female ticks during 50 days after the first antibiotic treatment and vitamin supplementation. (A) Tick survival rate. **(B)** Laying probability: number of ticks laying eggs/ population size. **(C)** Time between blood meal and first day of oviposition (in days). **(D)** Number of eggs laid per tick corrected for the mass of the blood fed female tick (number of eggs per mg of tick weight). **(E)** Mean egg weight (in mg). **(F)** Egg incubation time (in days). **(G)** Egg hatching rate. **(H)** Time between hatching and molting into first nymphal stage (in days) **(I)** Molting success rate. In histograms, 95% confidence intervals are presented.

No differences were registered for the time needed to molt into first nymphal stage (**Figure 4H**, **Table 1**). On average, larvae successfully molted in 6.30 days [SD: 0.657]. Finally, no significant differences were observed for molting rates which reached 98.4% on average [SD: 4.77]. (**Figure 4I**, **Table 1**).

After the first period of 50 days, the surviving ticks were fed a second time with the same feeding protocol. Female ticks were monitored during a new period of 50 days and parameters associated with reproduction were measured (**Table 2**).

Table 2: Statistical analyses performed on traits associated with reproductive fitness measured in female ticks during 50 days after the second antibiotic treatment and vitamin supplementation.

Variable	Family	Effect of the treatment	Effect of the vitamin	Effect of Interaction
Survival over full duration of the experiment (102 days)	Binomial	Pr: 0.006548** Dev: 10.06	Pr: 0.9598 Dev: 0.002503	Pr: 0.06723 Dev: 5.399
Survival over 50 days after the second blood meal	Binomial	Pr: 0.1501 Dev: 3.793	Pr: 0.7106 Dev: 0.1376	Pr: 0.04820* Dev: 6.065
Laying probability	Binomial	Pr: 0.08903 Dev: 5.146	Pr: 0.7287 Dev: 0.1280	Pr: 0.2472 Dev: 2.974
Number of days to first oviposition	Quasi-Poisson[1]	Pr: 0.6820 Dev: 4.207	Pr: 0.1323 Dev: 12.46	Pr: 0.2084 Dev: 17.24
Number of eggs standardized by mother weight[3]	Quasi-Poisson[1]	Pr: 0.2587 Dev: 132.4	Pr: 0.3037 Dev: 51.79	Pr: 0.01714* Dev: 398.1
Average egg mass	Normal	Pr: 0.9865 Dev: 0.0000662	Pr: 0.8609 Dev: 0.0001511	Pr: 0.3596 Dev: 0.005115
Egg incubation time	Quasi-Poisson[2]	Pr: 0.3250 Dev: 0.2497	Pr: 0.6804 Dev: 0.01885	Pr: 0.01565* Dev: 0.9236
Egg hatching rate	Binomial	Pr: 0.7355 Dev: 0.6145	Pr: 0.1896 Dev: 1.721	Pr: 0.4796 Dev: 1.470
Time for molting into first nymphal stage	Quasi-Poisson[2]	Pr: 0.0003831*** Dev: 1.264	Pr: 0.6249 Dev: 0.01920	Pr: 0.9009 Dev: 0.01676
Molting rate	Binomial	Pr: 0.2100 Dev: 3.121	Pr: 0.717 Dev: 0.1314	Pr: 1.000 Dev: $6.72e^{-5}$

Note: Pr: P-value. Dev: Deviance, X^2 for binomial and quasi-Poisson distribution, F-value for normal distribution. Degree of freedom was 2 for the effect of the treatment and the effect of the interaction and 1 for the effect of vitamin. [1] Distribution corrected for overdispersion of the data. [2] Distribution corrected for underdispersion of the data. [3] Variable corrected using an offset of the tick weight. Statistically significant results are highlighted with a grey background. Statistical difference code: 0 '***' 0.001 '**' 0.01 '*' 0.05 ' ' 1

B vitamins: ☐ Absence of vitamin supplementation ☐ Presence of vitamin supplementation

Figure 5: Histograms and boxplots of the traits associated with reproductive fitness measured in female ticks during 50 days after the second antibiotic treatment and vitamin supplementation. (A) Tick survival rate. **(B)** Laying probability: number of ticks laying eggs/ population size. **(C)** Time between blood meal and first day of oviposition (in days). **(D)** Number of eggs laid per tick corrected for the mass of the blood fed female adult tick (number of eggs per mg of tick weight). **(E)** Mean egg weight (in mg). **(F)** Egg incubation time (in days). **(G)** Egg hatching rate. **(H)** Time between hatching and molting into first nymphal stage (in days) **(I)** Molting success rate. In histograms, 95% confidence intervals are presented.

Figure 6: Histogram of the survival rate of *Ornithodoros moubata* ticks after antibiotic treatment and vitamin supplementation over the whole experiment (102 days). 95% confidence intervals are presented.

Measurements after the second blood meal yielded largely similar results (**Figure 5**, **Table 2, Table S2**). There was no effect of antibiotic and vitamin treatments, except for the time before molting which was increased for larvae of the rifampicin group (6.59 days against 6.38 for the control group and 6.34 for the gentamycin group). No effect was observed regarding vitamin supplementation (**Figure 5H**, **Table 2**).

Considering the whole experiment (two blood meals), a strong effect of the antibiotic treatment group is revealed on survival. The control groups presented some of the lowest survival rates (33.3% and 30.4% respectively in absence and presence of vitamins). The rifampicin group without B vitamins also presented a low survival rate (30.0%), whereas the three other groups presented higher survival rates: 57.1% for rifampicin with vitamin, 54.5% and 76.2% for the gentamycin group with or without vitamins (**Figure 6**, **Table 2**).

Following the ticks over two consecutive blood meals brings information on how the tick reproduction parameters are affected over time in *Ornithodoros moubata* (**Table 3, Table S2**). Almost all traits associated with reproductive fitness were negatively affected during the second 50-days period after blood meal: survival rates, laying probability, number of eggs laid, egg mass and hatching rate decreased whereas egg incubation time and molting time increased. The number of days to oviposition and the molting rate did not change after the second blood meal. These observations are mostly independent from the treatment received during blood meals.

329 **Table 3: Comparison of the reproductive indicators measured in adult female ticks between the two 50 days**
330 **periods after blood meal with antibiotic treatment (gentamycin, rifampicin, or negative control) and vitamin**
331 **supplementation (presence of vitamins and absence of vitamins).**

Variable	Family	Effect of Aging	Effect of Interaction with Antibiotic	Effect of Interaction with Vitamin
Survival over 50 days after blood meal	Binomial	Pr: 0.001892** Dev: 9.652	Pr: 0.4465 Dev: 1.613	Pr: 0.4004 Dev: 0.7072
Laying probability	Binomial	Pr: $1.208e^{-07}$ *** Dev: 28.01	Pr: 0.8867 Dev: 0.2404	Pr: 0.3864 Dev: 0.7502
Number of days to first oviposition	Quasi-Poisson[1]	Pr: 0.5998 Dev: 0.8048	Pr: 0.8483 Dev: 0.9622	Pr: 0.008667** Dev: 20.15
Number of eggs standardized by mother weight[3]	Quasi-Poisson[1]	Pr: 0.0004472*** Dev: 327.6	Pr: 0.1917 Dev: 87.80	Pr: 0.1817 Dev: 47.39
Average egg mass	Normal	Pr: 0.03459* Dev: 0.01311	Pr: 0.9348 Dev: 0.001939	Pr: 0.6345 Dev: 0.006528
Egg incubation time	Quasi-Poisson[2]	Pr: $2e^{-16}$ *** Dev: 7.352	Pr: 0.5287 Dev: 0.1153	Pr: 0.5771 Dev: 0.02813
Egg hatching rate	Binomial	Pr: $<2e^{-16}$ *** Dev: 187.2	Pr: 0.0229* Dev: 7.553	Pr: 0.3265 Dev: 0.9629
Time for molting into first nymphal stage	Quasi-Poisson[2]	Pr: 0.009203** Dev: 0.5013	Pr: 0.0002558*** Dev: 1.223	Pr: 0.3808 Dev: 0.05678
Molting rate	Binomial	Pr: 0.9751 Dev: 0.09712	Pr: 0.2268 Dev: 2.697	Pr: 0.3196 Dev: 0.9905

332 Note: Pr: P-value. Dev: Deviance, X^2 for binomial and quasi-Poisson distribution, F-value for normal distribution. Degree of
333 freedom was 1 for the effect of aging and 2 for the effect of interactions. [1] Distribution corrected for overdispersion of the data.
334 [2] Distribution corrected for underdispersion of the data. [3] Variable corrected using an offset of the tick weight. Statistically
335 significant results are highlighted with a grey background. Statistical difference code: 0 '***' 0.001 '**' 0.01 '*' 0.05 ' ' 1

Discussion

Many experts on ticks, tick-microbe interactions and tick-borne diseases now consider that the tick microbiota manipulation might become a potential new tool to fight against tick and tick-borne pathogens in the future (e.g., Bonnet *et al*. 2017; Bonnet et Pollet 2021). Before considering such an eventuality, it is essential to identify members of the tick microbiota and their role on both tick biology and physiology. It is also crucial to experimentally estimate consequences of the tick microbiota modifications on ticks. In this study, we focused on the microbiota of the tick *Ornithodoros moubata*. *O. moubata* is known to host two main bacterial endosymbionts, *Francisella* (namely *Francisella*-like endosymbionts (FLE)) and *Rickettsia*, with FLE being the one harboring the nutritional function (Duron *et al*. 2018). Here, using antibiotic treatments, we tried to eliminate members of *O. moubata* microbiota, particularly these two obligate endosymbionts, to evaluate the consequences of this dysbiosis on several parameters of the tick biology.

Impacts of antibiotics in an experimental design

Based on previous studies, we decided to use a protocol with an antibiotic treatment in the blood meal instead of using antibiotic injections or hosts treated with antibiotics (Duron *et al*. 2018; Guizzo *et al*. 2017; Zhang *et al*. 2017; Ninio *et al*. 2015). As a lab species since decades, *O. moubata* presents excellent results with artificial feeding and this method was chosen for easier use and more accurate control of antibiotic concentrations. The protocol used by Duron *et al.* (Duron *et al*. 2018) was chosen as a model and adapted for our study. Another option, which was ruled out here, was the use of microinjections to treat the ticks (Guizzo *et al*. 2017; Li, Zhang, et Zhu 2018; Zhang *et al*. 2017; Zhong, Jasinskas, et Barbour 2007; Kurlovs *et al*. 2014; Ninio *et al*. 2015). While this method is extremely efficient to ensure that the right amount of antibiotics is administered, it remains traumatic for the tick and is not adapted for soft tick species (Bonnet et Liu 2012).

The choice of the antibiotic molecules was based on the target symbionts. Duron *et al.* recently sequenced the lineage of *O. moubata* from the Cirad laboratory (Duron *et al*. 2018). Based on this study, we decided to use rifampicin which can eliminate both Rickettsia and FLE, and gentamycin, for which most *Rickettsia* strains are resistant whereas *Francisella* species are sensitive. Therefore, the elimination of *Francisella* alone under gentamycin treatment would allow to evaluate its role independently from *Rickettsia* (for which insights would be given using rifampicin treatment). However, the role of the rest of the microbiota would still need to be considered and evaluated. Other molecules such as kanamycin and tetracycline were ruled out due to their potential toxicity for arthropods (Li, Zhang, et Zhu 2018; Coon, Brown, et Strand 2016). Finally, the vitamin supplementation was chosen based on the identification of essential B vitamins (riboflavin, folic acid and biotin) (Duron *et al*. 2018).

Another important methodological aspect is the use of vitamin supplementation. Indeed, if an effect observed on reproductive parameters under antibiotic treatment is suppressed when supplementing with vitamins, it suggests that this change was linked to the nutritional role of the microbiota. On the opposite, if an effect is not rescued by supplementing in vitamins, it is more likely to be the consequence of a direct role of the symbiont on the reproduction.

Impacts of antibiotics on the tick microbiota

The experiment performed on first stage nymphs brought interesting results on the effect of the blood meal on the bacterial population size. Indeed, for both *Francisella*-like endosymbionts and *Rickettsia*, a significant increase in the amount of bacterial DNA was observed during 30 days after blood meal. It also appeared that the amount of actin DNA increased over the same period of time, suggesting a strong metabolic activity with multiple cellular divisions. This is most probably linked to the molting process which takes 10.5 days [SD: 1.06] under those conditions (Jourdan-Pineau, unpublished data). As intracellular bacteria, it is possible that *Francisella* and *Rickettsia* took advantage of this metabolic increase for their own growth as observed in other species (Sassera *et al*. 2008). Those endosymbionts might be simply more active during this period of intense metabolic activity which would be in agreement with the alterations of the molting process observed after antibiotic treatments (Duron *et al*. 2018).

Focusing on ticks treated with antibiotics, first stage nymphs showed contrasted responses according to the symbiont considered. Over the 30 days after treatment, no effect of the gentamycin was detected on

Rickettsia populations. This result was expected for the gentamycin treatment group as *Rickettsia* are known
to be resistant to this antibiotic (Vanrompay *et al*. 2018). Conversely, the rifampicin treatment significantly
lowered *Rickettsia* population size over time compared to the control group. Focusing on the bacterial
concentration instead of the ratios, it appeared that the concentration of *Rickettsia* slightly decreased or
stayed at the same level as it was at day 0. This might suggest a strong bacteriostatic effect of the rifampicin
treatment on *Rickettsia* populations.
Regarding *Francisella*-like endosymbiont, significant differences were observed in the ratios of
concentration between the treatment groups with a decrease in the rifampicin group. When looking at the
concentration of FLE, as observed for Rickettsia, the concentration of FLE tended to rise after blood meal in
the gentamycin groups, whereas, it remained close to a basal level in the rifampicin one. More information
was obtained from the qPCR performed on adult females following the two antibiotic treatments. We observed
that the ratios of concentrations of both FLE and *Rickettsia* strongly decreased in the rifampicin groups and
were slightly lowered in the gentamycin ones compared to the control groups 50 days after the second
treatment.
400
Overall, the qPCR results suggest that the rifampicin treatments limited the growth and/or decreased the
population size for *Rickettsia* and FLE. It seemed that the effect was bacteriostatic after one treatment on
nymphs during growth. The diminution of the population of symbiont after two treatments can be the
consequence of this bacteriostatic effect over time or the result of a bactericide effect after several consecutive
treatments. In the end, the treatment performed was sufficient to prevent the growth of FLE and *Rickettsia*
even though insufficient to eliminate completely the microbiota.
A slight effect was reported with the gentamycin treatment but this antibiotic (or, at least, the posology
used for this antibiotic) seemed to be less effective than the rifampicin. Finally, no effect of the vitamin group
was reported on the concentration of endosymbiont in adult ticks. This result was expected as the microbiota
does not depend on B vitamins, it rather produces B vitamins.
Our experiments question the use of antibiotics during artificial blood feeding of ticks. In hard ticks,
antibiotics and antifungal molecules are frequently added to the blood to avoid the development of micro-
organisms during the long periods of times needed for the blood meal (Krull *et al*. 2017; Militzer *et al*. 2021;
González, Bickerton, et Toledo 2021). This use of antibiotics and the molecules chosen should be carefully
considered during experiment design as they will modify the tick's microbiota and, potentially, affects the
ability of the tick to develop normally and to be maintained in a colony. This attention should also be increased
for lab experiments performed to identify the potential influence of members of the microbiota on the
presence of pathogens and *vice versa*.

Consequences of the modification of the microbiota on the tick physio-biology

In female ticks, considering only the antibiotic treatments, no effect was detected on the reproduction
itself after the first treatment. This result would suggest that the FLE and *Rickettsia* are not implicated in the
reproduction of the tick. However, at this point of the experiment it is also possible that the effect of treatment
did not eliminate enough symbiont yet. Furthermore, even if offspring of treated mothers showed no impaired
or delayed development until first nymphal stage, we cannot rule out that their microbiota might be modified.
Further experiments on those offspring would allow to characterize their microbiota and see if they suffer any
developmental deficiency during subsequent nymphal stages.
Besides, we observed that the negative control group presented a higher mortality rate than the antibiotic
groups. This surprising result led us to consider the fitness cost of harboring endosymbionts, as already
referenced for other arthropods (Ankrah, Chouaia, et Douglas 2018). For instance, it is possible to make the
hypothesis that *Francisella* would harbor an essential long-term function for the tick (providing essential
nutrients, avoiding a deficiency) but a short-term cost for the tick (increased mortality during reproduction).
Another possible explanation would be that other bacterias than FLE and *Rickettsia* were eliminated by the
antibiotic treatment and that those bacterias presented a fitness cost for the tick. This hypothesis remains less
probable than a role of FLE and Rickettsia which are the most frequent in *O. moubata* (Duron *et al*. 2018).
Nevertheless, the effect of antibiotic treatments should not be considered alone. Several trends can be
noticed for ticks which received vitamin supplementations. Vitamin supplementation tended to be associated
with a reduction of the time between mating and first oviposition, an increase in the success rate of oviposition

and, a slight increase in the success rate of the first molting. This is consistent with the idea that these vitamins
are essential for the tick and could lead to better reproductive and molting parameters and highlights the
importance of nutritional symbionts such as FLE. However, these trends were not significant in our study and
must be considered carefully. It is possible that the duration of our experiment was not sufficient to fully
observe the effects of a vitamin deficiency. Moreover, FLE were maintained at a low concentration but not
completely eliminated here, suggesting that all ticks were already supplied, at some extent, in B vitamins.

If we consider the results obtained after the second antibiotic treatment, two main elements can be
stressed out. First, is the confirmation of an effect on tick survival. After both treatments, the control group
presented a higher mortality than the gentamycin group. However, after the second treatment, the rifampicin
group was also affected by an increased mortality rate. One hypothesis could be that more FLE have been
eliminated after the second treatment and, as a consequence, the deficiency in B vitamin for the tick is stronger
than in the first treatment, leading to the death of adult females. This hypothesis is comforted by the fact that
the survival rate in the rifampicin group after the second treatment was better when the ticks were
supplemented with B vitamins. From these results, it would seem that there is a cost to maintain high levels
of symbionts (increased mortality in the control group), but the absence of symbionts results in a similar death
rate (mortality in the second rifampicin group), and finally in the groups with moderate amount of symbiont,
the benefits of endosymbiont compensate any associated fitness cost (gentamycin treatment groups).
Second, when regarding the parameters associated with reproduction, no clear-cut effect can be
established after the second treatment. The trends noticed during first treatment for vitamin supplementation
are not confirmed and the few significant results (egg incubation time and time to molting) present variations
of low range. If it is difficult to interpret the variations for the egg incubation time, the variations on the molting
time on the opposite are oriented with a longer time period for the rifampicin group. As suggested in the
literature, endosymbionts are crucial during molting (Duron *et al*. 2018), and their elimination through
rifampicin could indeed lead to a deterioration of the parameters associated with molting.
When looking at past literature on the role of endosymbionts for tick reproduction, several effects were
associated with the elimination of *Coxiella*-like endosymbionts (CLE) in hard ticks, including a decrease in
number of eggs laid (Guizzo *et al*. 2017; Zhang *et al*. 2017), an increase in the egg incubation time (Li, Zhang,
et Zhu 2018), a decrease in the hatching rate (Guizzo *et al*. 2017; Li, Zhang, et Zhu 2018; Zhong, Jasinskas, et
Barbour 2007) and a decrease in the molting success (Zhong, Jasinskas, et Barbour 2007). Interestingly, no
effects were reported for *Rickettsia* (Kurlovs *et al*. 2014) or FLE. This is especially interesting as all these three
endosymbionts have a nutritional role. The results presented here indicate no effect of FLE or *Rickettsia* on the
tick reproduction. It is possible that CLE are the only tick symbionts which improve the reproductive fitness of
their host in addition to B vitamin production, leading them to remain one of the most commonly found
endosymbiont in ticks (Duron *et al*. 2017).
In our study, we presented results issued from two successive treatments. Interestingly, in *O. moubata*,
the vitellogenesis processes start during the last nymphal stages (Sonenshine 1991) suggesting that the role of
the microbiota on reproduction may occur before adult stage. In the future, antibiotic treatments will have to
be performed starting from late nymph stages to evaluate the role of the microbiota during these stages on
the reproduction of the adults obtained.
However, blood meal and mating are also critical steps for vitellogenesis processes (Horigane *et al*. 2010)
and most studies performed on tick reproduction targeted directly adult females (Guizzo *et al*. 2017; Li, Zhang,
et Zhu 2018; Zhang *et al*. 2017; Zhong, Jasinskas, et Barbour 2007; Kurlovs *et al*. 2014). Parameters such as the
number of eggs laid depend directly on the efficiency of the blood meal. Moreover, as shown here and in
previous studies (Sassera *et al*. 2008), endosymbiont load blooms after the blood meal strengthening the
hypothesis that they have probably a role during and/or after the meal. Ideally, using germ-free tick lineages
during blood meal would bring striking elements of response on this point. Such lineages are however difficult
to sustain, and antibiotic treatment remains, for now, the main tool available.

Finally, even though it was not the objective of the study, monitoring adult females over two periods of
egg-laying brought data on the effects of ageing on the reproduction. Similarly to results obtained in model
invertebrates (Tatar 2010), the reproductive indicators in soft ticks were strongly deteriorated between the
first and second period of egg-laying. Our results indicate no link between this phenomenon and the presence
or absence of FLE and *Rickettsia* suggesting the effect is solely due to ageing.

492
To conclude, this study suggests that FLE and *Rickettsia* endosymbionts have no direct effect on the
reproduction of their tick host. Several trends associated with these bacterial species were highlighted
including a potential fitness cost, and strong load increase following blood meal.
However, elucidating the role of the tick microbiota remains an arduous task. Antibiotic treatments
constitute a powerful tool for this aim, but protocols need to be adjusted and modified in each tick lineage to
take into account the microbiota composition and sensitivity.

Author contributions

FT performed the experiments described, analyzed the data, and wrote the manuscript. TP provided the
material, obtained the funding, and conceived and coordinated the study. MD provided the material and
helped perform the experiments. LG helped perform the experiments and analyze the data. HJ provided the
material, obtained the funding, conceived and coordinated the study, performed the experiments and
analyzed the data. All authors read and corrected the final manuscript.

Acknowledgements

The authors would like to thank Olivier Duron, Rémi Pereira, and the members of the UMR ASTRE for the
discussions on molecular biology protocols, as well as Facundo Muñoz for the discussions about statistics.
All experiments on arthropods have been performed on the Baillarguet insectarium platform
(https://doi.org/10.18167/infrastructure/00001),member of the National Infrastructure EMERG'IN and of the
Vectopole Sud network (http://www.vectopole-sud.fr/). The Baillarguet insectarium platform is led by the joint
units Intertryp (IRD, Cirad) and ASTRE (Cirad, INRAE).

Data, scripts and codes availability

Data will be available online after peer-review process: https://doi.org/10.5061/dryad.bg79cnpfm.

Supplementary material

Table S1: Sample size and repartition of individuals into groups for the different experiments performed.

Experiment	Treatment group	Vitamin subgroup
Symbiont loads ; nymphs n=366	No treatment n=20	No treatment N=20
	Control n=116	Vitamin ; n=57
		No vitamin ; n=59
	Rifampicin n=110	Vitamin ; n=50
		No vitamin ; n=60
	Gentamicin n=120	Vitamin ; n=60
		No vitamin ; n=60
Symbiont loads ; females n=59	Control n=14	Vitamin ; n=7
		No vitamin ; n=7
	Rifampicin n=18	Vitamin ; n=12
		No vitamin ; n=6
	Gentamicin n=27	Vitamin ; n=13
		No vitamin ; n=14
Reproduction monitoring ;	Control	Vitamin ; n=23

first 50-days period n=132	n=46	No vitamin ; n=23
	Rifampicin n=42	Vitamin ; n=21
		No vitamin ; n=21
	Gentamicin n=44	Vitamin ; n=22
		No vitamin ; n=22
Reproduction monitoring ; second 50-days period n=99	Control n=27	Vitamin ; n=12
		No vitamin ; n=15
	Rifampicin n=33	Vitamin ; n=17
		No vitamin ; n=16
	Gentamicin n=39	Vitamin ; n=20
		No vitamin ; n=19

516   Note: Sample size variations between groups are the result of tick mortality in the group before the start of monitoring.

Table S2: Mean and variance for the traits associated with reproductive fitness after antibiotic treatment and vitamin supplementation.

Variable	First treatment	Second treatment
Laying probability	74.2% (98/132)	40.4% (40/99)
Survival over 50 days after blood meal	78.8% (104/132)	60.6% (60/99)
Number of days to first oviposition	23.9 days [SD: 6.81]	24.7 days [SD: 12.0]
Number of eggs / Number of eggs standardized by mother weight	109 eggs / 0.361 egg/mg [SD: 0.116]	65.2 eggs / 0.228 egg/mg [SD: 0.176]
Average egg mass	0.538 mg [SD: 0.0463]	0.516 mg [SD: 0.0671]
Egg incubation time	11.9 days [SD: 1.01]	12.4 days [SD: 1.18]
Egg hatching rate	79.6% [SD: 15.1]	55.0% [SD: 28.9]
Time for molting into first nymphal stage	6.30 days [SD: 0.657]	6.39 days [SD: 0.724]
Molting rate	98.4% [SD: 4.77]	98.5% [SD: 4.32]

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article.

Funding

This study was supported and financed by the Ecology and Evolution of Infectious Diseases Program, grant no. 2019-67015-28981 from the USDA National Institute of Food and Agriculture, in a project entitled "unraveling the effect of contact networks & socio-economic factors in the emergence of infectious diseases at the wild-domestic interface" (https://www.asf-nifnaf.org/).

References

Aivelo, Tuomas, Anna Norberg, et Barbara Tschirren. 2019. « Bacterial Microbiota Composition of *Ixodes Ricinus* Ticks: The Role of Environmental Variation, Tick Characteristics and Microbial Interactions ». *PeerJ* 7: e8217. https://doi.org/10.7717/peerj.8217.

Ankrah, Nana Y. D., Bessem Chouaia, et Angela E. Douglas. 2018. « The Cost of Metabolic Interactions in Symbioses between Insects and Bacteria with Reduced Genomes ». *MBio* 9 (5). https://doi.org/10.1128/mBio.01433-18.

Bates, Douglas, Martin Mächler, Ben Bolker, et Steve Walker. 2015. « Fitting Linear Mixed-Effects Models Using Lme4 ». *Journal of Statistical Software* 67 (1): 1-48. https://doi.org/10.18637/jss.v067.i01.

Binetruy, Florian, Marlène Dupraz, Marie Buysse, et Olivier Duron. 2019. « Surface Sterilization Methods Impact Measures of Internal Microbial Diversity in Ticks ». *Parasites & Vectors* 12 (1): 268. https://doi.org/10.1186/s13071-019-3517-5.

Bonnet, Sarah I., Florian Binetruy, Angelica M. Hernández-Jarguín, et Olivier Duron. 2017. « The Tick Microbiome: Why Non-Pathogenic Microorganisms Matter in Tick Biology and Pathogen Transmission ». *Frontiers in Cellular and Infection Microbiology* 7: 236. https://doi.org/10.3389/fcimb.2017.00236.

Bonnet, Sarah Irène, et Thomas Pollet. 2021. « Update on the Intricate Tango between Tick Microbiomes and Tick-Borne Pathogens ». *Parasite Immunology* 43 (5): e12813. https://doi.org/10.1111/pim.12813.

Bonnet, Sarah, et Xiangye Liu. 2012. « Laboratory artificial infection of hard ticks: a tool for the analysis of tick-borne pathogen transmission ». *Acarologia* 52 (4): 453-64. https://doi.org/10.1051/acarologia/20122068.

Budachetri, Khemraj, Deepak Kumar, Gary Crispell, Christine Beck, Gregory Dasch, et Shahid Karim. 2018. « The Tick Endosymbiont *Candidatus Midichloria Mitochondrii* and Selenoproteins Are Essential for the Growth of *Rickettsia Parkeri* in the Gulf Coast Tick Vector ». *Microbiome* 6 (1): 141. https://doi.org/10.1186/s40168-018-0524-2.

Coon, Kerri L., Mark R. Brown, et Michael R. Strand. 2016. « Mosquitoes host communities of bacteria that are essential for development but vary greatly between local habitats ». *Molecular ecology* 25 (22): 5806-26. https://doi.org/10.1111/mec.13877.

Duron, Olivier, Florian Binetruy, Valérie Noël, Julie Cremaschi, Karen D. McCoy, Céline Arnathau, Olivier Plantard, et al. 2017. « Evolutionary Changes in Symbiont Community Structure in Ticks ». *Molecular Ecology* 26 (11): 2905-21. https://doi.org/10.1111/mec.14094.

Duron, Olivier, Julie Cremaschi, et Karen D. McCoy. 2016. « The High Diversity and Global Distribution of the Intracellular Bacterium *Rickettsiella* in the Polar Seabird Tick *Ixodes Uriae* ». *Microbial Ecology* 71 (3): 761-70. https://doi.org/10.1007/s00248-015-0702-8.

Duron, Olivier, et Yuval Gottlieb. 2020. « Convergence of Nutritional Symbioses in Obligate Blood Feeders ». *Trends in Parasitology* 36 (10): 816-25. https://doi.org/10.1016/j.pt.2020.07.007.

Duron, Olivier, Olivier Morel, Valérie Noël, Marie Buysse, Florian Binetruy, Renaud Lancelot, Etienne Loire, et al. 2018. « Tick-Bacteria Mutualism Depends on B Vitamin Synthesis Pathways ». *Current Biology: CB* 28 (12): 1896-1902.e5. https://doi.org/10.1016/j.cub.2018.04.038.

Estrada-Peña, Agustín, Alejandro Cabezas-Cruz, Thomas Pollet, Muriel Vayssier-Taussat, et Jean-François Cosson. 2018. « High Throughput Sequencing and Network Analysis Disentangle the Microbial Communities of Ticks and Hosts Within and Between Ecosystems ». *Frontiers in Cellular and Infection Microbiology* 8: 236. https://doi.org/10.3389/fcimb.2018.00236.

Gall, Cory A, Kathryn E Reif, Glen A Scoles, Kathleen L Mason, Michelle Mousel, Susan M Noh, et Kelly A Brayton. 2016. « The bacterial microbiome of *Dermacentor andersoni* ticks influences pathogen susceptibility ». *The ISME Journal* 10 (8): 1846-55. https://doi.org/10.1038/ismej.2015.266.

González, Julia, Mathew Bickerton, et Alvaro Toledo. 2021. « Applications of Artificial Membrane Feeding for *Ixodid* Ticks ». *Acta Tropica* 215 (mars): 105818. https://doi.org/10.1016/j.actatropica.2020.105818.

Guizzo, Melina Garcia, Luís Fernando Parizi, Rodrigo Dutra Nunes, Renata Schama, Rodolpho M. Albano, Lucas Tirloni, Daiane Patrícia Oldiges, et al. 2017. « A *Coxiella* mutualist symbiont is essential to the development of *Rhipicephalus microplus* ». *Scientific Reports* 7 (décembre). https://doi.org/10.1038/s41598-017-17309-x.

Horigane, M., T. Shinoda, H. Honda, et D. Taylor. 2010. « Characterization of a Vitellogenin Gene Reveals Two Phase Regulation of Vitellogenesis by Engorgement and Mating in the Soft Tick *Ornithodoros Moubata* (*Acari*: *Argasidae*) ». *Insect Molecular Biology* 19 (4): 501-15. https://doi.org/10.1111/j.1365-2583.2010.01007.x.

Jiménez-Cortés, J. Guillermo, Rodolfo García-Contreras, Martha I. Bucio-Torres, Margarita Cabrera-Bravo, Alex Córdoba-Aguilar, Giovanni Benelli, et Paz M. Salazar-Schettino. 2018. « Bacterial Symbionts in Human Blood-Feeding Arthropods: Patterns, General Mechanisms and Effects of Global Ecological Changes ». *Acta Tropica* 186 (octobre): 69-101. https://doi.org/10.1016/j.actatropica.2018.07.005.

Kaviraj, A., F. Bhunia, et N. C. Saha. 2004. « Toxicity of Methanol to Fish, Crustacean, Oligochaete Worm, and Aquatic Ecosystem ». *International Journal of Toxicology* 23 (1): 55-63. https://doi.org/10.1080/10915810490265469.

Krull, Christoph, Bettina Böhme, Peter-Henning Clausen, et Ard M. Nijhof. 2017. « Optimization of an artificial tick feeding assay for *Dermacentor reticulatus* ». *Parasites & Vectors* 10 (février): 60. https://doi.org/10.1186/s13071-017-2000-4.

Kurlovs, Andre H., Jinze Li, Du Cheng, et Jianmin Zhong. 2014. « *Ixodes pacificus* Ticks Maintain Embryogenesis and Egg Hatching after Antibiotic Treatment of *Rickettsia* Endosymbiont ». *PLoS ONE* 9 (8). https://doi.org/10.1371/journal.pone.0104815.

Leeson, H. S. 1952. « The Recorded Distribution of *Ornithodoros Moubata* (*Murray*) (*Acarina*) ». *Bulletin of Entomological Research* 43 (2): 407-11. https://doi.org/10.1017/S000748530004058X.

Lejal, E., J. Chiquet, J. Aubert, S. Robin, A. Estrada-Peña, O. Rue, C. Midoux, et al. 2021. « Temporal Patterns in *Ixodes Ricinus* Microbial Communities: An Insight into Tick-Borne Microbe Interactions ». *Microbiome* 9 (1): 153. https://doi.org/10.1186/s40168-021-01051-8.

Li, Lan-Hua, Yi Zhang, et Dan Zhu. 2018. « Effects of antibiotic treatment on the fecundity of *Rhipicephalus haemaphysaloides* ticks ». *Parasites & Vectors* 11 (avril). https://doi.org/10.1186/s13071-018-2807-7.

Militzer, Nina, Alexander Bartel, Peter-Henning Clausen, Peggy Hoffmann-Köhler, et Ard M. Nijhof. 2021. « Artificial Feeding of All Consecutive Life Stages of *Ixodes Ricinus* ». *Vaccines* 9 (4): 385. https://doi.org/10.3390/vaccines9040385.

Narasimhan, Sukanya, Nallakkandi Rajeevan, Lei Liu, Yang O. Zhao, Julia Heisig, Jingyi Pan, Rebecca Eppler-Epstein, Kathleen DePonte, Durland Fish, et Erol Fikrig. 2014. « Gut Microbiota of the Tick Vector *Ixodes scapularis* Modulate Colonization of the Lyme Disease Spirochete ». *Cell host & microbe* 15 (1): 58-71. https://doi.org/10.1016/j.chom.2013.12.001.

Ninio, Camille, Olivier Plantard, Valentina Serra, Claudia Pollera, Nicola Ferrari, Alessandra Cafiso, Davide Sassera, et Chiara Bazzocchi. 2015. « Antibiotic Treatment of the Hard Tick *Ixodes Ricinus*: Influence on *Midichloria Mitochondrii* Load Following Blood Meal ». *Ticks and Tick-Borne Diseases* 6 (5): 653-57. https://doi.org/10.1016/j.ttbdis.2015.05.011.

Peirce, M. A. 1974. « Distribution and Ecology of *Ornithodoros Moubata Porcinus Walton* (*Acarina*) in Animal Burrows in East Africa* ». *Bulletin of Entomological Research* 64 (4): 605-19. https://doi.org/10.1017/S0007485300035896.

Pereira De Oliveira, Rémi, Laurence Vial, et Marie-Frédérique Le Potier. 2022. « Quantification of ASFV DNA and RNA in *Ornithodoros* Soft Ticks ». *Methods in Molecular Biology (Clifton, N.J.)* 2503: 105-18. https://doi.org/10.1007/978-1-0716-2333-6_7.

R Core Team. 2023. « R: A language and environment for statistical computing. » R Foundation for Statistical Computing, Vienna, Austria. 2023. https://www.R-project.org/.

624   Rio, Rita V. M., Geoffrey M. Attardo, et Brian L. Weiss. 2016. « Grandeur Alliances: Symbiont Metabolic
625   Integration and Obligate Arthropod Hematophagy ». *Trends in Parasitology* 32 (9): 739-49.
626   https://doi.org/10.1016/j.pt.2016.05.002.

627   Sánchez-Vizcaíno, J. M., L. Mur, A. D. S. Bastos, et M. L. Penrith. 2015. « New Insights into the Role of Ticks
628   in African Swine Fever Epidemiology ». *Revue Scientifique Et Technique (International Office of*
629   *Epizootics)* 34 (2): 503-11. https://doi.org/10.20506/rst.34.2.2375.

630   Sassera, Davide, Nathan Lo, Edwin A. P. Bouman, Sara Epis, Michele Mortarino, et Claudio Bandi. 2008.
631   « "*Candidatus Midichloria*" Endosymbionts Bloom after the Blood Meal of the Host, the Hard Tick
632   *Ixodes Ricinus* ». *Applied and Environmental Microbiology* 74 (19): 6138-40.
633   https://doi.org/10.1128/AEM.00248-08.

634   Sonenshine, Daniel E. 1991. « Biology of ticks. » *Oxford University Press* Vol. 1
635   https://doi.org/10.1017/S0007485300042668

636   Tatar, Marc. 2010. « Reproductive Aging in Invertebrate Genetic Models ». *Annals of the New York Academy*
637   *of Sciences* 1204 (août): 149-55. https://doi.org/10.1111/j.1749-6632.2010.05522.x.

638   Vanrompay, Daisy, Thi Loan Anh Nguyen, Sally J. Cutler, et Patrick Butaye. 2018. « Antimicrobial Resistance
639   in *Chlamydiales*, *Rickettsia*, *Coxiella*, and Other Intracellular Pathogens ». *Microbiology Spectrum* 6
640   (2). https://doi.org/10.1128/microbiolspec.ARBA-0003-2017.

641   Vayssier-Taussat, Muriel, Emmanuel Albina, Christine Citti, Jean-François Cosson, Marie-Agnès Jacques,
642   Marc-Henri Lebrun, Yves Le Loir, et al. 2014. « Shifting the Paradigm from Pathogens to Pathobiome:
643   New Concepts in the Light of Meta-Omics ». *Frontiers in Cellular and Infection Microbiology* 4: 29.
644   https://doi.org/10.3389/fcimb.2014.00029.

645   Vial, Laurence. 2009. « Biological and Ecological Characteristics of Soft Ticks (*Ixodida*: *Argasidae*) and Their
646   Impact for Predicting Tick and Associated Disease Distribution ». *Parasite (Paris, France)* 16 (3):
647   191-202. https://doi.org/10.1051/parasite/2009163191.

648   Zhang, Chun-Mian, Ning-Xin Li, Tian-Tian Zhang, Zhao-Xi Qiu, Yuan Li, Li-Wu Li, et Jing-Ze Liu. 2017.
649   « Endosymbiont CLS-HI Plays a Role in Reproduction and Development of *Haemaphysalis*
650   *Longicornis* ». *Experimental & Applied Acarology* 73 (3-4): 429-38. https://doi.org/10.1007/s10493-
651   017-0194-y.

652   Zhong, Jianmin, Algimantas Jasinskas, et Alan G. Barbour. 2007. « Antibiotic Treatment of the Tick Vector
653   *Amblyomma americanum* Reduced Reproductive Fitness ». *PLoS ONE* 2 (5).
654   https://doi.org/10.1371/journal.pone.0000405.

655   Zuur, Alain F., Elena N. Ieno, Neil Walker, Anatoly A. Saveliev, et Graham M. Smith. 2009. *Mixed Effects*
656   *Models and Extensions in Ecology with R*. Statistics for Biology and Health. New York, NY: Springer.
657   https://doi.org/10.1007/978-0-387-87458-6.

659