Confidence-based Prediction of Antibiotic Resistance at the Patient Level

Juan S. Inda-Díaz[1,2,3,*], Anna Johnning[1,2,4], Magnus Hessel[2,5], Anders Sjöberg[4,6], Anna

Lokrantz[4], Lisa Helldal[5], Mats Jirstrand[4,6], Lennart Svensson[6] and Erik Kristiansson[1,2*]

[1]Department of Mathematical Sciences, Chalmers University of Technology and University of

Gothenburg; Gothenburg, SE-41296, Sweden.

[2]Centre for Antibiotic Resistance Research (CARe), University of Gothenburg; Gothenburg,

Sweden.

[3]Centre for Microbiome Research, School of Biomedical Sciences, Queensland University of

Technology, Translational Research Institute, Woolloongabba, Australia.

[4]Department of Systems and Data Analysis, Fraunhofer-Chalmers Centre; Gothenburg, SE-

41288, Sweden.

[5]Department of Clinical Microbiology, Sahlgrenska University Hospital, Region Västra

Götaland, Gothenburg, Sweden.

[6]Department of Electrical Engineering, Chalmers University of Technology; Gothenburg,

Sweden.

*Address correspondence to Juan S. Inda-Díaz, inda@chalmers.se and Erik Kristiansson,

erik.kristiansson@chalmers.se.

ABSTRACT

Rapid and accurate diagnostics of bacterial infections are necessary for efficient treatment of antibiotic-resistant pathogens. Cultivation-based methods, such as antibiotic susceptibility testing (AST), are limited by bacterial growth rates and seldom yield results before treatment needs to start, increasing patient risk and contributing to antibiotic overprescription. Here, we present a deep-learning method that leverages patient data and available AST results to predict antibiotic susceptibilities that have not yet been measured. After training on three million AST results from 30 European countries, the method achieved an average accuracy of 93% across bacterial species and antibiotics. It predicted susceptibility with an average major error rate below 5% for quinolones, cephalosporins, and carbapenems, and below 8% and 14% for aminoglycosides and penicillins, respectively. Furthermore, the model predicted resistance with an average very major rate below 10% for cephalosporins, carbapenems, and aminoglycosides, but with higher very major error rates for penicillins and quinolones. We combined the method with conformal prediction and demonstrated accurate estimation of the predictive uncertainty at the patient level. Our results suggest that AI-based decision support may offer new means to meet the growing burden of antibiotic resistance.

IMPORTANCE

Improved diagnostic tools are vital for maintaining efficient treatment of antibiotic-resistant bacteria and for reducing antibiotic overconsumption. In our research, we introduce a new deep learning-based method capable of predicting untested antibiotic resistance phenotypes. The method uses transformers, a powerful AI technique that efficiently leverages both antibiotic susceptibility tests (AST) and patient data simultaneously. The model produces predictions that can be used as time- and cost-efficient alternatives to results from cultivation-based diagnostic

assays. Significantly, our study highlights the potential of AI technologies to address the

increasing prevalence of antibiotic-resistant bacterial infections.

INTRODUCTION

The global rise of antibiotic-resistant bacterial infections threatens human health globally

(1). In 2019, almost five million yearly deaths were attributed to antibiotic-resistant bacteria *(2)*,

a number that is expected to continue to grow in the coming decades *(3)*. Reduced antibiotic

efficacy in treatment increases the risk of performing vital healthcare procedures – including

surgery, chemotherapy, and organ transplantation *(4)* – and, thereby, jeopardizes modern

medicine as a whole.

Accurate and fast diagnostics are necessary for effective treatment of antibiotic-resistant

bacteria. A central method is antibiotic susceptibility testing (AST), a laboratory test in which a

bacterium isolated from a patient sample is cultivated and its resistance phenotype assessed in

the presence of antibiotics *(5)*. However, AST can be time-consuming due to the often low

bacterial growth rate and the large number of antibiotics that may need to be tested for highly

multidrug-resistant isolates. Tier-based testing strategies and time constraints usually yield

incomplete AST results initially. For life-threatening infections, treatment needs to start as early

as possible, often before AST result are available *(6)*. Under these circumstances, the choice of

treatment is reduced to educated guesses based on limited diagnostic information *(7)*. This form

of "empirical" treatment is associated with increased patient risks and overprescription of

antibiotics (8–10).

Antibiotic resistance is commonly caused by resistance genes encoding various defense

mechanisms. These genes are often co-localized on mobile genetic elements, such as plasmids

and transposons *(11, 12)*. Multiple resistance genes can thus be transferred simultaneously

between bacterial cells, giving rise to strong correlations in susceptibility to different antibiotics.

The infecting bacterium and, therefore, its susceptibility profile are, furthermore, dependent on

patient characteristics, including age, sex, and the geographical region where the infection was

acquired (13–15). Indeed, patient data has previously been shown to contain valuable

information for selecting suitable antibiotic therapy for bacterial urinary tract infections (15–17).

There are, however, no AI-based methods that can combine patient data with the initial, often

incomplete AST results and thus guide treatment choice based on all available diagnostic

information. Indeed, integrating patient data with available AST results could enable more

accurate prediction of susceptibility to antibiotics that have not been tested, thereby providing

physicians with comprehensive diagnostic information at an earlier point in time.

Artificial intelligence (AI) and deep learning have been successfully applied to

diagnostics *(18, 19)*, but the focus has primarily been on image-based data commonly used in

radiology and pathology *(20)*. In contrast, methods for non-image multimodal data, which are

more prevalent in the diagnosis of infectious diseases, have received less attention *(21)*. Several

methods for predicting phenotypes from genotypic data have also been proposed (22), as have

machine-learning methods using electronic health records and patient data (23). There are,

however, few AI-based decision support systems for antibiotic treatment selection in use in

clinical settings *(24)*. A major culprit in the development of such methods is the complexity of

the diagnostic data, which is typically categorical (stratified test results and patient data) and

contains dependencies and redundancies. Also, when applied to antibiotic-resistant bacteria, the

incompleteness of the diagnostic data, particularly AST results, requires approaches that can

efficiently handle missing observations. Furthermore, since model accuracy depends on the

amount of available information, any prediction must be accompanied by estimates of its

uncertainty. Indeed, the possibility of disregarding predictions that are insufficiently confident is

vital for critical decision-making and, thus, essential for the adoption of AI-based methodologies

in healthcare settings *(25)*. However, today, most AI-based diagnostic methods are primarily

evaluated in populations and do not provide uncertainty information for patient-level predictions

(26). Existing methods have, so far, been unable to adequately address these challenges and are

either limited to a few specific antibiotics, cannot incorporate missing AST data, or do not

provide any estimates of the predictive uncertainty *(27, 28)*.

In recent years, transformer-based models, such as BERT (bidirectional encoder

representations from transformers) *(29)* and GPT (generative pre-trained transformer) *(30)*, have

transformed natural language processing. These models operate on categorical input data, often

structured into sentences of words, and subject them to multi-head self-attention *(31)*. This

process enables inference of complex word dependencies directly from data and thereby predicts

missing parts. Therefore, we hypothesized that transformers may be suitable for predicting

antibiotic susceptibility results from a combination of incomplete diagnostic information and

patient data. Multimodal self-attention would enable the identification of complex dependencies

among diagnostic data types and facilitate extrapolation to susceptibilities that have not been

tested. Transformers have previously been shown to be highly useful beyond natural language

processing *(32, 33)*, but are rarely used for diagnosing infectious diseases.

In this study, we present a novel transformer-based method that accurately predicts

antibiotic susceptibilities using patient data and incomplete diagnostic information for three

common bacterial pathogens, *Escherichia coli*, *Klebsiella pneumoniae*, and *Pseudomonas*

aeruginosa. We combine the model with conditional inductive conformal prediction *(34)* to

estimate the uncertainty of each prediction at the patient level, thereby enabling dismissal of

predictions with too low certainty. The model was trained and evaluated on a large,

heterogeneous dataset containing AST results from blood infections collected during routine care

across 30 European countries. Our results showed that the model could accurately predict

susceptibility to a wide range of antibiotics, even when only a few AST results were included in

the input. We also show that the model handles increasing information well, providing more

accurate predictions as additional AST results become available. Finally, we demonstrate that

predictions can be made within predefined confidence levels for most bacterial species and

antibiotics, which allows control of both the major and very major error rates. We conclude that

the combination of transformers and conditional inductive conformal predictions constitutes an

appealing class of models for integrating and predicting diagnostic information.

RESULTS

A transformer-based model for antibiotic susceptibility prediction

We developed a transformer-based model to predict unavailable diagnostic information

using multiple classifications. The input to the model is a set of words containing the available

diagnostic information (results of antibiotic susceptibility testing; AST) and patient data (Figure

1). The AST results are assumed to be incomplete, where only a subset of the possible tests has

available results. The model uses two transformers, which summarize the patient and AST

information in two sentence-specific classification (CLS) vectors, one representing patient-

specific information and one representing bacterial isolate-specific information. These vectors

then serve as input to multiple antibiotic-specific feed-forward neural networks, each predicting

the probability of susceptibility to the corresponding antibiotic. The uncertainty is estimated

using inductive conformal prediction, conditioned on the susceptibility; hence, controlling the

false-positive (major error) and false-negative (very major error) rates for each antibiotic *(34,*

35). The full details of the model architecture and the uncertainty estimation are provided in

Methods.

The model was trained and evaluated on data from The European Surveillance System

(TESSy) (https://www.ecdc.europa.eu/en/publications-data/european-surveillance-system-tessy),

which contains AST results for 1,161,303 *E. coli*, 301,506 *K. pneumoniae*, and 166,448 *P.*

aeruginosa isolates collected from blood infections of patients in 30 European countries. The

training and evaluation were performed on 20 (16, 18, and 11 for *E. coli*, *K. pneumoniae*, and *P.*

aeruginosa, respectively) commonly used antibiotics that belong to five large and clinically

relevant antibiotic classes: aminoglycosides, carbapenems, cephalosporins, penicillins, and

quinolones (Table 1). A bacterial isolate was, on average, tested for susceptibility to 7.3, 8.8, and

7.8 antibiotics, with standard deviations of 1.9, 2.4, and 1.7 for *E. coli*, *K. pneumoniae*, and *P.*

aeruginosa, respectively (Figures S1 and S2). The most commonly tested antibiotics were

ceftazidime (CAZ) and ciprofloxacin (CIP), for which at least 93% of isolates were tested,

regardless of pathogen, country of origin, or patient gender. In contrast, less than 10% of the *E.*

coli and *K. pneumoniae* isolates were tested for piperacillin (PIP), moxifloxacin (MFX), and

nalidixic acid (NAL, Figure S3). The susceptibility rate for *E. coli* isolates was lowest for the

penicillins amoxicillin (AMX), ampicillin (AMP), and PIP (43%–50%), whereas for *K.*

pneumoniae isolates it was lowest for PIP (20%). For other antibiotics and *P. aeruginosa*

isolates, the susceptibility rates were higher, representing a more unbalanced dataset. A slightly

higher susceptibility rate was observed among isolates collected from female patients (Figure

S4).

Predictions of antibiotic susceptibility have high performance

To assess how well the model predicts the susceptibility of antibiotics not included as

inputs, we used 70% of the bacterial isolates for 5-fold cross-validation and model training. We

used 15% of the isolates for testing, and the remaining 15% for calibration of conformal

prediction. During training, calibration, and testing, we selected a random number of antibiotics

as input to the model (distribution available in Figure S5; mean 5.7, 6.3, and 5.7 with standard

deviations of 1.3, 1.5, and 1.1 for *E. coli*, *K. pneumoniae*, and *P. aeruginosa* isolates,

respectively; see Methods for details) together with patient data. The susceptibility of the

remaining tested antibiotics was assumed to be unknown and, therefore, masked from the input

data. The model's predictions were then compared with the antibiotics masked from the input to

evaluate performance. In this setting, the model had an overall high performance that did not

differ substantially between folds in the cross-validation, with a standard error (se) below 0.015

for all antibiotics and bacterial species except the major error (ME) for the antibiotic PIP in *K.*

pneumoniae (se = 0.016 for both training and test datasets). The performance was largely

consistent across the training and test processes (Figures 2 and S6). There were, however,

apparent differences in performance between the three bacterial species and between antibiotics.

The F1 score (the harmonic mean of precision and recall) in the test dataset for *E. coli and K.*

pneumoniae isolates was highest for cephalosporins (91% and 96% on average, respectively),

and quinolones (86% and 93% on average, respectively), though lower for penicillins (74% and

88% on average, respectively, Figure 2A). For *P. aeruginosa* isolates, the F1 score was 84% for

penicillins, and 82% for quinolones and cephalosporines. Carbapenems had an average F1 score

of 90% and 76% for *K. pneumoniae* and *P. aeruginosa* isolates, respectively, while

aminoglycosides had the lowest F1 scores: 65% for *E. coli* and 76% for *K. pneumoniae* and *P.*

aeruginosa isolates.

Next, we evaluated the model based on the ME rate, defined as the proportion of

susceptible bacterial isolates erroneously predicted as resistant, and the very major error (VME)

rate, defined as the proportion of resistant isolates erroneously predicted as susceptible. The ME

and VME rates are standard performance measures in antibiotic susceptibility testing, and are

frequently used to evaluate and compare diagnostic methods *(36)*. Based on the test dataset,

cephalosporins had, on average over all predictions in the test data, an ME rate of 1.4% (1%–

1.8%), 2.3% (1.5%–3.9%), and 5.2% (5.1%–5.2%) for *E. coli*, *K. pneumoniae*, and *P.*

aeruginosa isolates, respectively, while the corresponding VME rate was, on average, 6.7%

(5.6%–9.4%), 2.3% (1.2%–3.2%), and 12.2% (9.7%–12.8%) (Figure 2B). For quinolones, the

average ME rate was 2.2% (2%–5%), 2.9% (1.6%–4.7%), and 3.3% (2.5%–4.2%), while the

VME rate was, on average 19.2% (8.9%–28.6%), 8.3% (4.2%–13.4%), and 21.9% (20.3%–

23.5%) for *E. coli*, *K. pneumoniae*, and *P. aeruginosa* isolates, respectively. For *E. coli* isolates,

the penicillins had a higher overall average ME rate of 12.5% (4.1%–16.6%). Here, PIP had the

lowest VME rate (12.2%), while the other penicillins had rates between 23% and 26%. For *K.*

pneumoniae and *P. aeruginosa* isolates, the average ME for penicillins, excluding PIP, was 7.3%

and 6.3%, respectively, while the VME was 11%. We noted that the model struggled to predict

PIP resistance in *K. pneumoniae* with an ME as high as 51%. This may be a consequence of data

imbalance, where *K. pneumoniae* had a resistance rate of 80% to PIP (compared to 50% and 22%

for *E. coli* and *P. aeruginosa*, respectively) but accounted for only 11% of the data for this

antibiotic. Aminoglycosides had average ME/VME rates of 7.3/16.3, 10.2/7.4, and 5.5/9.5, for *E.*

coli, K. pneumoniae, and *P. aeruginosa* isolates, respectively. Finally, the ME and VME rates for

carbapenems for *K. pneumoniae* isolates were 2.3% (1.7%–3.1%) and 4.4% (2.8%–7.3%),

respectively, and for *P. aeruginosa* isolates, 8.4% (5.3%–11.4%) and 14.6% (10.9%–18.4%),

respectively.

The number of available AST results influenced the model's performance. When the

number of AST results used as input to the model increased from four to eight antibiotics,

significant reductions were seen in the ME rate for penicillins, where AMX and AMP dropped

from 34% and 22% to 3% and 7%, respectively, and the VME of CIP, which fell from 36% to

7% in *E. coli* isolates (Figure 3D). Interestingly, the drop for other antibiotics and pathogens was

more modest or non-existent. A substantial reduction in the VME rate was also observed across

most antibiotics and pathogens.

When eight AST results were included in the input, fourteen of sixteen antibiotics in *E.*

coli isolates, sixteen of twenty antibiotics in *K. pneumoniae*, and all antibiotics in *P. aeruginosa*

had an ME rate below 10%, and below 5% for eleven, thirteen, and ten antibiotics in each

pathogens, respectively. Similarly, the VME rate was below 10% for nine, sixteen, and ten, and

below 5% for six, twelve, and three of the antibiotics corresponding to each pathogen. Going

from four to eight input AST results increased the F1 scores from 75% to 87% for *E. coli*, from

85% to 92% for *K. pneumoniae*, and from 75% to 84% for *P. aeruginosa*, on average. The

model's overall high performance, as indicated by its F1 score, ME, and VME rates, suggests that

it can be used to predict complete susceptibility patterns for bacterial isolates accurately.

Control of the major and very major error rates at the patient level

In clinical practice, decisions are taken at the patient level. It is thus vital that decision

support based on predictions also conveys information about the confidence in the prediction.

Indeed, if the uncertainty is too high, the prediction may need to be considered with care or

completely dismissed until more diagnostic information becomes available. Therefore, we

implemented an algorithm that assigns a quantitative measure of uncertainty to each individual

prediction *(34)*. The algorithm was based on conditional inductive conformal prediction, where

the certainty of each prediction was derived from a conformity measure, defined in our case as

the softmax score from each antibiotic-specific neural network (Figure 1B). The distributions of

softmax scores for both susceptible and resistant bacterial isolates were empirically calculated

for each antibiotic and each of the three pathogens, using a dedicated calibration dataset. These

distributions were used to determine whether a new prediction aligns with the susceptible and/or

resistant isolates in the calibration dataset and, based on a predefined confidence level, is deemed

sufficiently certain. The output of a final prediction for one antibiotic is a prediction set, with

either a) a single label, i.e., either susceptible or resistant, when there is enough conformity to

only one of the softmax distributions; b) multiple labels, i.e. both susceptible and resistant, if

there is enough conformity for both softmax distributions; or c) no label if there is no conformity

to either group. The proportion of predictions that will, on average, have the correct label in their

prediction set is governed by the confidence level $1 - \varepsilon$. Note that in this setting, ε corresponds

to the average ME and the VME rates for susceptible and resistant bacterial isolates for each

pathogen, respectively (see Methods for full details).

The empirically derived ME and VME rates were close to the prespecified values of ε for

all antibiotics (Figures S7-S11). For example, at a confidence level of 90% ($\varepsilon = 0.1$), the

observed ME rates were, on average, 10.1%, 9.9%, and 9.9% (standard deviation 0.4%, 1.3%,

and 0.3%), and the observed VME rates were, on average 9.7%, 9.4%, and 9.3% (standard

deviation 1.43%, 0.95%, and 1.7%) for *E. coli*, *K. pneumoniae*, and *P. aeruginosa* isolates,

respectively. The concordance between pre-specified and observed error rates was sustained at

higher confidence levels where, for a confidence level of 95%, the average ME rates were 5.1%,

5.1%, and 5% (standard deviation 0.5%, 1%, and 0.3%) and the average VME rates were 5%,

4.7%, and 5% (standard deviation 0.8%, 0.6%, and 1.4%) for *E. coli*, *K. pneumoniae*, and *P.*

aeruginosa isolates, respectively. Thus, the results showed that the specified confidence levels

calculated from empirical distributions were sufficiently stable between datasets.

The model could confidently predict the phenotype for a large majority of bacterial isolates

and antibiotics

At a confidence level of 90%, as many as 82%, 89%, and 86% of the predictions were

unambiguous and correct (only the correct label was in the prediction set) for *E. coli*, *K.*

pneumoniae, and *P. aeruginosa* isolates, respectively; however, this number varied between

antibiotics (Figure 4A, 5A, 6A). Cephalosporins had a high proportion of correct unambiguous

predictions (90%) across the three pathogens, and quinolones had 90% for *K. pneumoniae*. For

the remaining antibiotics, between 82% and 89% of the predictions were correct and

unambiguous, except for penicillins for *E. coli* (71%) and quinolones for *P. aeruginosa*, which

showed higher uncertainty. The proportion of unambiguous predictions was, as expected,

reduced when the confidence level was increased; to 78%, 89%, and 77% at a 95% confidence

level and, finally, to 72%, 85%, and 63% at a 97.5% confidence level for *E. coli*, *K. pneumoniae*,

and *P. aeruginosa*, respectively. On the other hand, the number of unambiguous predictions

increased as more diagnostic information was provided to the model. At a 90% confidence level,

the proportion of correct unambiguous predictions increased from, on average, 76%, 84%, and

83% when four AST results were included in the input, to 89%, 91%, and 91% when eight AST

results were included in the input for *E. coli, K. pneumoniae*, and *P. aeruginosa* isolates,

respectively (Figure 4B, 5B, 6B). The increase was substantial for predicting susceptibility to the

penicillins AMP and AMX in *E. coli* isolates, to PIP and OFX in *K. pneumoniae* isolates, and for

predicting susceptibility and resistance to CIP, as well as susceptibility to LVX and IPM in *P.*

aeruginosa isolates. With eight input AST results, the model achieved 90% correct,

unambiguous predictions for most of the remaining masked antibiotics. This was also true for

higher confidence levels, where the proportion of unambiguous and correct predictions with

eight input AST results was 89% and 84% for 95% and 97.5% confidence, respectively.

DISCUSSION

In this study, we present a method that uses a transformer model to predict unavailable

diagnostic information. When combined with conditional inductive conformal prediction to

estimate the predictive uncertainty at the patient level, the method can abstain from making

decisions unless the confidence is deemed sufficiently high. The model was applied to the

diagnostics of infectious bacteria, a field facing rapidly rising societal and economic costs due to

the growing challenges related to antibiotic resistance *(37, 38)*. We showed that only a few

antibiotic susceptibility test (AST) results are sufficient to accurately derive a more complete

resistance profile. This has the potential to guide antibiotic therapy for serious infections, where

treatment must begin before all diagnostic tests are completed.

The method was trained on a large, heterogeneous dataset comprising over 12 million

AST results from more than a million bacterial isolates of three pathogens collected from 30

European countries. This dataset was compiled for surveillance purposes and contains AST

results originally produced in routine diagnostics. Validation, in which AST results were

randomly excluded, showed that the model had generally high accuracy in predicting antibiotic

susceptibility. The performance improved further as more ASTs were included in the input data,

demonstrating that the model could efficiently incorporate diagnostic information as it becomes

available and, thus, over time, produce more certain predictions. Indeed, when AST for eight

antibiotics was used as input, the model could predict most susceptibilities with a VME rate

(false-negative rate) as low as 10%. This indicates that AI prediction can serve as a viable

complement to laboratory diagnostic tests and may be utilized to save time, alleviate suffering,

and reduce economic costs.

Providing information about predictive uncertainty is crucial in diagnostics, where

population-level knowledge is used to predict outcomes for individual patients. We addressed

this challenge by implementing an algorithm based on conditional inductive conformal

prediction *(34, 39, 40)* to provide each prediction with an accompanying confidence score. In our

application, the predefined confidence corresponded to ME and VME rates, yielding predictions

that limit false-positive and false-negative rates to the desired levels. During model testing, we

found consistent error rates across datasets and, for cephalosporins and quinolones, the

prediction sets mainly contained a single, correct label. However, a higher variation was

observed for penicillins, resulting in a larger proportion of ambiguous predictions (multiple

labels). Furthermore, our implementation separates the uncertainty for susceptible and resistant

predictions. This is valuable in the diagnostics of bacterial infections, where a VME, i.e., the

incorrect identification of a resistant isolate, may lead to ineffective antibiotic treatment.

Therefore, VMEs are often considered to be the most serious errors, especially for life-

threatening infections. The ability to set confidence scores for susceptible and resistant

predictions individually enables the model to be adjusted for different clinical scenarios.

The model showed a clear difference in predictive performance between antibiotics,

notoriously lower for penicillins, especially compared to cephalosporins. Historically, penicillins

were among the earliest classes of antibiotics and the first beta-lactam antibiotics introduced.

Resistance to beta-lactam antibiotics is typically caused by enzymes that hydrolyze the

antibiotics *(41)*. Penicillins have been widely used for over 80 years, and bacteria can

consequently acquire a wide diversity of resistance genes *(42)*. In contrast, resistance to

cephalosporins – a later generation of beta-lactam antibiotics – is, to a larger extent, dependent

on additional genetic events, such as mutations in chromosomal genes or the acquisition of

broader-spectrum resistance mechanisms *(41)*. These patterns were reflected in the data, where

resistance to penicillins was common. Indeed, as many as 87.7% of the *E. coli* and 61% of the *K.*

pneumoniae bacterial isolates resistant to a single antibiotic were resistant to a penicillin.

Furthermore, it is plausible that the order of evolution of multidrug-resistant bacteria affects the

performance of our model. We also consider resistance phenotypes that are initially acquired to

be harder to predict than those that commonly appear in later stages of evolution, simply because

there are no other correlating susceptibilities. Results from additional diagnostic assays, such as

targeted molecular tests that detect antibiotic resistance genes and genomic data, could further

improve performance. Indeed, the flexibility of the transformers allows incorporating other types

of diagnostic information, including genotypic information, as new words in the input sentence.

Ultimately, the AI methodology presented here has the potential to enhance the diagnosis

of infections caused by antibiotic-resistant bacteria. We argue that data-driven methods have the

potential to replace selected diagnostic assays, thereby providing physicians with more

comprehensive decision support at an earlier stage. This has the potential to improve the

339 treatment of bacterial infections, thereby decreasing patient morbidity and mortality, reducing

340 costs, and limiting the overprescription of antibiotics.

342 **MATERIALS AND METHODS**

343 **Data description**

344  This study is based on data from the European Surveillance System (TESSy), collected as

345 part of surveillance conducted by the European Centre for Disease Prevention and Control. The

346 dataset contains more than 12 million antibiotic susceptibility testing (AST) results for bacteria

347 isolated from blood and cerebrospinal fluid of hospitalized patients across 30 European

348 countries. For each bacterial isolate, we retrieved the species name, AST results, the patient's

349 gender and age, the date of bacterial isolation, and the country where the test was performed.

350 This metadata will be referred to collectively as the patient data. The analysis was limited to the

351 susceptibility of *Escherichia coli, Klebsiella pneumoniae,* and *Pseudomonas aeruginosa* isolates,

352 which were the most common species, from blood infections collected between 2007 and 2020.

353 Tests that resulted in either a resistant (R) or a susceptible (S) result were included, while tests

354 with an I result were excluded. Only antibiotics with a resistance rate of at least 8% were

355 considered. This resulted in data covering sixteen antibiotics from five clinically relevant classes:

356 aminoglycosides, cephalosporins, penicillins, quinolones, and carbapenems (Table 1).

357 Furthermore, isolates tested for fewer than five antibiotics were removed. If an antibiotic was

358 tested multiple times for the same bacterial isolate, only the most recent test was included. The

359 final dataset contained $n = 1{,}629{,}257$ bacterial isolates with 12,433,856 AST results, resulting

360 in an average of 7.6 tests (standard deviation of 2.1) per isolate. After randomization, the

361 bacterial isolates were divided into three datasets: a training dataset (70%), a calibration dataset

362　(15%), and a test dataset (15%). The train dataset was further divided into five groups for 5-fold

363　cross-validation.

Data expansion

To increase the number of AST result combinations, each dataset was expanded. For each

bacterial isolate j, multiple data points were generated by randomly splitting the susceptibility

test results into two groups. The first group, together with the patent information, x_j, was

considered known and, thus, used as input to the model. The second group, y_j, was hidden from

the model and used to evaluate the predictive performance. The full details of the data expansion

are provided in Supplemental Materials (Table S1). A made-up example of the information

available for a bacterial isolate is presented below:

Available information for bacterial isolate j: "ESCCOL SV 30 M 2013_01 LVX_R AMC_S

AMP_S TZP_R CTX_S GEN_S", represents a. *E. coli* bacterium isolated at a Swedish hospital

(SV) from a 30-year-old (30) male (M) patient in January 2013 (2013_01) where the isolated

bacterium was tested against six antibiotics. The AST results indicated resistance to the

antibiotics levofloxacin (LVX) and piperacillin/tazobactam (TZP), and susceptibility to

amoxicillin/clavulanic acid (AMC), ampicillin (AMP), cefotaxime (CTX), and gentamicin

(GEN). Two example data points, (x_k, y_k) and $(x_{k'}, y_{k'})$, that could, potentially, be created from

this isolate:

$(x_k, y_k) = ($"ESCCOL SV 30 M 2013_01 LVX_R AMC_S AMP_S TZP_R", "CTX_S GEN_S"$)$,

$(x_{k'}, y_{k'}) = ($"ESCCOL SV 30 M 2013_01 LVX_R AMC_S AMP_S CTX_S GEN_S", "TZP_R"$)$.

Model Architecture and Training

Given a data point, (x_k, y_k), the model takes the input x_k and makes predictions $\hat{y}_k$ of the

susceptibilities y_k. The input sentence x_k is split into patient data and AST data, both of which

contain the word for the bacterial species, and both are complemented at the start with the

classification word, *CLS*. The patient data and the AST data are padded to a length $L = 6$ and

$L = 21$ with *PAD* words if needed. Each word is then converted into a linear embedding

representation in the form of a d-dimensional vector that provides semantic meaning to the

model ($d = 128$). These word embeddings are passed through two transformer encoder layers,

each with two attention heads, followed by an add-and-normalize layer, a position-wise feed-

forward layer (using 256 nodes), and, finally, another add-and-normalize layer. The first vector

of the output from each encoder– representing the *CLS* word and containing information at the

sentence level– are concatenated and fed to a linear transformation. The output is used as input to

$M = 20$ independent antibiotic-specific feed-forward networks, each of depth two. The

intermediate layer of the network has 128 nodes with rectified linear unit (ReLU) activation

function, followed by a normalization step, while the final layer outputs a linear transformation

to vectors of length two, which are used for binary classification. The isolate was classified as

resistant or susceptible based on the largest output value. Additionally, models with the

combinations of 1 or 2 encoder heads and layers, with word embedding of size 64 or 128, and a

position-wise feed-forward layer of 128 or 256, with patient data and no patient data, were

trained (Figure S13-S15). The proposed model achieved higher accuracy, lower ME, and lower

VME in 23, 14, and 10 out of 45 pathogen-antibiotic combinations, respectively. The

combination of one head, one layer, with word embedding of size 128 and a position-wise feed-

forward layer of size 256 was the only one to have more pathogen-antibiotic combinations with

lower VME (13).

The model was trained as follows. At each epoch, 248,000 bacterial isolates were

randomly sampled from the train dataset and expanded as described in Methods – Data

expansion. The model was then trained on 512,000 randomly selected data points from the data

expansion step, divided into mini-batches of size 512. The loss was based on the cross-entropy

between the known (y_k) and predicted $(\hat{y}_k)$ labels. The Adam optimizer was used to minimize

the loss over 700 epochs with a fixed learning rate of 1×10^{-6}. The model was implemented and

trained using Pytorch version 1.7.1.

Uncertainty control

An algorithm based on conditional inductive conformal prediction was used to quantify

the uncertainty of the predictions *(34)* with respect to the antibiotic and label (i.e., "susceptible"

and "resistant") (34). Conformal prediction uses empirical data to estimate uncertainty and has

previously been shown to suit complex diagnostic data for which valid distributional

assumptions may be hard to make *(43)*. For a data point (x_k, y_k) belonging to pathogen q, the

algorithm estimates a prediction set $\Gamma_k^{\varepsilon,q}$, containing the predictions that are deemed sufficiently

confident given a predefined confidence level $1 - \varepsilon$. The uncertainty for a prediction was based

on its conformity measure, defined as the softmax transformation of the outputs of a neural

network. The conformity measure and, thus, the uncertainty, was derived individually for each

antibiotic and each label (i.e., resistance or susceptible).

We estimated the empirical distributions for each conformity measure from the

calibration dataset separately for each pathogen, which, for an antibiotic a and pathogen q, were

assumed to contain $l_{q,a} = l_{q,a,S} + l_{q,a,R}$ data points, where $l_{q,a,S}$ and $l_{q,a,R}$ are the number of data

points for susceptible and resistant bacterial isolates of pathogen q, respectively. For a data point

(x_k, y_k) derived from pathogen q, let $\alpha_k^{a,S}$ and $\alpha_k^{a,R}$ denote the softmax score for the prediction

of susceptibility and resistance to antibiotic a, respectively. The prediction sets were decided

based on the empirical p-values $p_k^{q,a,S}$ and $p_k^{q,a,R}$, which were calculated according to

$$p_k^{q,a,S} = \frac{|i = 1, \ldots, l_{q,a,S} : \tilde{\alpha}_i^{q,a,S} \leq \alpha_k^{a,S}| + 1}{l_{q,a,S} + 1},$$

$$p_k^{q,a,R} = \frac{|i = 1, \ldots, l_{q,a,R} : \tilde{\alpha}_i^{q,a,R} \leq \alpha_k^{a,R}| + 1}{l_{q,a,R} + 1},$$

433   where $\tilde{\alpha}_i^{q,a,S}$ and $\tilde{\alpha}_i^{q,a,R}$ denotes the softmax scores calculated using the data points in the

434   calibration dataset for isolates from pathogen q. At a confidence $1 - \varepsilon$, the prediction set was

435   then formed by

$$\Gamma_k^{\varepsilon,a} = \{S \text{ if } p_k^{q,a,S} > \varepsilon\} \bigcup \{R \text{ if } p_k^{q,a,R} > \varepsilon\}.$$

Performance

The model's predictive performance was computed based on 257,784, 242,121, and

114,410 train; 285,897, 223,661, and 97,542 calibration; and 280,428, 223,665, and 100,927 test

individual AST results for *E. coli*, *K. pneumoniae*, and *P. aeruginosa* isolates, respectively. To

evaluate the overall model performance, the F1 score was calculated. In addition, the *major error*

(ME) rate, defined as the proportion of true susceptible bacterial isolates erroneously predicted as

resistant, and the *very major error* (VME) rate, defined as the proportion of true resistant isolates

erroneously predicted as susceptible, were also calculated. To measure the performance of the

uncertainty control, true predictions were defined as prediction regions containing the true label

for each antibiotic, and false predictions were those containing either no labels or only the wrong

one.

For comparison, the following naive classifier was included (Figure S7). For each of the

bacterial species, the frequencies between all pairwise antibiotics and their susceptibility profile

were calculated in the training dataset. Then, the ratios $r_{a_R,b} = \frac{|a_R b_R|}{|a_R b_R| + |a_R b_S|}$ and $r_{a_S,b} =$

$\frac{|a_S b_R|}{|a_S b_R| + |a_S b_S|}$, between antibiotic a, with resistance (R) and susceptible (S) profiles, and the

antibiotic b, with resistance profile, were calculated for all pairwise combinations of antibiotics.

Finally, for a new prediction of antibiotic b in y_k, resistance was deemed if the average A =

$\sum_{a_j \in x_k} \frac{r_{a_j b}}{n_a}$ was larger than 0.5, where a_j represent the antibiotics in x_k with resistance $j \in$

$\{R, S\}$, and n_a is the number of antibiotics in x_k, the isolate was deemed susceptible otherwise.

To investigate the effect of patient data, we compared our results with a model based solely on

AST results (Figure S6). The VMEs for the three bacterial species were found to be significantly

smaller (p-val<0.01) when the original model was compared to the model without patient data

and to the naive classifier, as determined by Wilcoxon signed-rank tests.

ACKNOWLEDGMENTS

Data from The European Surveillance System – TESSy between 2007 to 2020, provided

by Andorra, Albania, Armenia, Austria, Azerbaijan, Bosnia and Herzegovina, Belgium,

Bulgaria, Belarus, Switzerland, Cyprus, Czechia, Germany, Denmark, Estonia, Greece, Spain,

Finland, France, Georgia, Croatia, Hungary, Ireland, Israel, Iceland, Italy, Kyrgyzstan,

Kazakhstan, Liechtenstein, Lithuania, Luxembourg, Latvia, Monaco, Republic of Moldova,

Montenegro, Republic of North Macedonia, Malta, Netherlands, Norway, Poland, Portugal,

Romania, Serbia, Russian Federation, Sweden, Slovenia, Slovakia, San Marino, Tajikistan,

Turkmenistan, Turkey, Ukraine, United Kingdom, Uzbekistan, Kosovo and released by The

European Centre for Disease Prevention and Control (ECDC). The views and opinions of the

authors expressed herein do not necessarily state or reflect those of ECDC. The accuracy of the

authors' statistical analysis and the findings they report are not the responsibility of ECDC.

ECDC is not responsible for conclusions or opinions drawn from the data provided. ECDC is not

responsible for the correctness of the data and data management, data merging, and data collation

after the provision of the data. ECDC shall not be held liable for improper or incorrect use of the

data.

DATA AVAILABILITY STATEMENT

The data used in this study belongs to the European Surveillance System – TESSy. We

acknowledge the European Surveillance System – TESSy for data availability. The model is

available in the GitHub repository https://github.com/indajuan/Confidence-based-Prediction-of-

Antibiotic-Resistance.

FUNDING

Swedish Research Council (VR), grant 2018-02835 (EK)

Swedish Research Council (VR), grant 2019-03482 (EK)

Chalmers AI Research Centre (CHAIR), (EK)

National Health and Medical Research Council of Australia under the framework of JPI

AMR, grant 2031902, SEARCHER

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

SUPPLEMENTAL MATERIALS

Figure S1. The number of susceptible (opaque) and resistant (transparent) bacterial isolates tested for each antibiotic and pathogen in the dataset.

Figure S2. Distribution of the number of antibiotics tested per bacterial isolate for each pathogen in the dataset.

Figure S3. Distribution of the proportion of bacterial isolates tested against each antibiotic for female (opaque) and male (transparent) patients and for each pathogen across countries in the dataset. The center line, box limits, and whiskers represent the median, upper, and lower quartiles, and 1.5 times the interquartile range, respectively.

Figure S4. The proportion of bacterial isolates susceptible to each antibiotic for female (opaque) and male (transparent) patients and for each pathogen across countries in the dataset. The center line, box limits, and whiskers represent the median, upper, and lower quartiles, and 1.5 times the interquartile range, respectively.

Figure S5. Histogram of the number of antibiotic susceptibility testing (AST) results used as input for the model during training at each epoch for each pathogen.

Figure S6. Performance of the cross-validation. The major error (red) and very major error (VME) rates for each fold are represented by dots, the horizontal line, box limits, and whiskers represent the median, upper, and lower quartiles, and 1.5 times the interquartile range, respectively, across folds.

Figure S7. Performance of the model with and without patient data and the naive classifier.

Figure S8-12. Empirical and expected error rates and performance of the model after conditional

inductive conformal prediction. For each pathogen and penicillins (S8), cephalosporins (S9),

quinolones (S10), aminoglycosides (S11), and carbapenems (S12) antibiotics.

Figure S13-S15. Performance of the model with different hyperparameters: 1 or 2 heads (1H,

2H), 1 or 2 layers (1L, 2L), word embedding of size 64 and position-wise feed-forward layer of

128 (small), or word embedding of size 128 and position-wise feed-forward layer of 256 (large),

with patient and AST data, or without patient data (no patient), for (S13) *E. coli*, (S14) *K.*

pneumoniae, and (S15) *P. aeruginosa*.

Table S1. Data expansion. For isolate j with A_j AST results, we created a fixed number of data

points at each epoch of training, calibration, and testing. The number of AST results, a_k,

included in the input sentence x_k of data point k was sampled from a binomial distribution with

parameters $N = A_j - 1$ and $p = \frac{D_j}{A_j - 1} - 1$, listed below. If $a_k < 4$, the data point was dismissed.

The number of antibiotic susceptibilities to predict for data point k was given by $A_j - a_k$, and

the true susceptibility test results are summarized in the sequence y_k. The decision on which

antibiotics were included in x_k and y_k was done randomly for all data points.

Table S2. Performance of the transformer model. Results from the test dataset: F1-score, major

error (ME) rate, and very major error (VME) rate shown for each antibiotic and pathogen,

considering all possible numbers of input AST results. Dash (–) indicates combinations of

bacterial species and antibiotics that the model does not cover.

Table S3. Performance of the transformer model as a function of the number of antibiotic

susceptibility testing (AST) results included in the input. Results from the test dataset: F1-score,

major error (ME) rate, and very major error (VME) rate shown for each antibiotic, pathogen, and

the number of antibiotic susceptibility testing (AST) results included in the input. Dash (–)

indicates combinations of bacterial species and antibiotics that the model does not cover. We

limited the table to 8 AST results in the input due to the low number of isolates tested for 10 or

more antibiotics in the data.

TABLES

Table 1. Bacterial isolates summary. The included antibiotics, the number of isolates from

each species tested for each antibiotic, their resistance rates, and the weights used in the loss

function.

Antibiotic	*Escherichia coli*		*Klebsiella pneumoniae*		*Pseudomonas aeruginosa*		Loss function weights
	Isolates	%R[1]	Isolates	%R[1]	Isolates	%R[1]	(S/R)[2]
AMC – Amoxicillin / clavulanic acid	550 937	33%	129 537	42%	–	–	0.45/0.55
AMP – Ampicillin	800 283	57%	–	–	–	–	0.45/0.55
AMX – Amoxicillin	293 369	55%	–	–	–	–	0.45/0.55
PIP – Piperacillin	93 919	50%	21 028	80%	62 218	22%	0.45/0.55
TZP – Piperacillin/ tazobactam	646 892	8%	164 551	29%	152 233	16%	0.15/85
CAZ – Ceftazidime	1 080 891	10%	280 916	30%	159 118	15%	0.3/0.7
CRO – Ceftriaxone	350 947	11%	90 851	30%	–	–	0.3/0.7
CTX – Cefotaxime	990 234	11%	253 971	30%	–	–	0.3/0.7
FEP – Cefepime	336 526	10%	102 213	33%	65 619	16%	0.3/0.7
CIP – Ciprofloxacin	1 117 333	22%	286 899	31%	160 669	20%	0.3/0.7
LVX – Levofloxacin	291 403	23%	73 574	29%	45 636	24%	0.3/0.7
MFX – Moxifloxacin	105 648	26%	23 231	26%	–	–	0.3/0.7
NAL – Nalidixic acid	63 468	27%	16 532	36%	–	–	0.45/0.55
OFX – Ofloxacin	136 704	19%	33 764	36%	–	–	0.3/0.7
AMK – Amikacin	–	–	202 968	9%	130 264	9%	0.15/85
GEN – Gentamicin	1 073 614	9%	278 400	20%	143 645	14%	0.15/85
TOB – Tobramycin	564 295	10%	145 914	27%	119 954	13%	0.15/85
ETP – Ertapenem	–	–	109 997	14%	–	–	0.15/85
IPM – Imipenem	–	–	189 733	10%	121 852	20%	0.15/85

| MEM – Meropenem | – | – | 242 732 | 9% | 130 352 | 15% | 0.15/85 |

[1]Proportion of resistance bacterial isolates to each of the antibiotics.

[2]Weight on the cross-entropy loss function for the two different labels: susceptible/resistant.

Dash (–) indicates combinations of bacterial species and antibiotics that the model does not

cover.

FIGURES

Figure 1. The proposed model architecture and uncertainty control. (**A**) The architecture of

the proposed model. Both input sequences start with a classification word, *CLS*, followed by the

bacterial species name. One of the input sequences also contains patient information (country,

gender, age, and sampling date), while the other includes available antibiotic susceptibility

testing (AST) data. Both input sequences are fixed to lengths $L = 5$ for the patient information

and $L = 21$ for the AST data, padded with the *PAD* word. We use a linear embedding to

represent the input words numerically, which are fed into the transformer encoders. The first

vectors (*CLS*) from the outputs of each encoder are combined into a linear model and fed to m

independent neural networks, each representing one antibiotic. The neural network outputs are

two-dimensional vectors indicating susceptibility and resistance, respectively. (**B**) Uncertainty

control. The neural network outputs undergo a softmax rescaling. A calibration dataset is used to

build empirical distributions of conformity for resistant and susceptible predictions for each

pathogen and antibiotic. The prediction regions for the data points k and $k + 1$ are built based on

the deviation of the observed softmax score from the empirical distribution and the confidence

interval threshold. See Methods for full details.

Figure 2. Performance of the transformer model. Results from the train and test datasets: (**A**)

F1-score, (**B**) major error (ME) rate, and very major error (VME) rate for the transformer model,

shown for each antibiotic and pathogen. Note that, due to data restrictions, not all antibiotics are

assessed for all three pathogens.

Figure 3. The predictive performance of the model as a function of the number of antibiotic

susceptibility testing (AST) results included in the input. Major error (ME) and very major

error (VME) rates from light to dark: 4 to 8 AST results are shown for each antibiotic and

pathogen. Note that, due to data restrictions, not all antibiotics are assessed for all three

pathogens.

Figure 4. The proportion of correct predictions as a function of the number of labels in the

prediction set for *E. coli* isolates. The proportion of correct predictions with a single label

(light) and multiple labels (dark), major errors (MEs), and very major errors (VMEs) with a

single label (light) and empty set (dark) predictions for resistant (R) and susceptible (S) isolates,

shown for each antibiotic and pathogen using three different confidence levels: 90%, 95%, and

97.5%. (**A**) The proportions are shown for each antibiotic at three confidence levels (90%, 95%,

and 97.5%), considering all possible numbers of input AST results. (**B**) The proportions are

shown as a function of the number of input AST results (90% confidence level).

Figure 5. The proportion of correct predictions as a function of the number of labels in the

prediction set for *K. pneumoniae* isolates. The proportion of correct predictions with a single

label (light) and multiple labels (dark), major errors (MEs), and very major errors (VMEs) with a

single label (light) and empty set (dark) predictions for resistant (R) and susceptible (S) isolates,

shown for each antibiotic and pathogen using three different confidence levels: 90%, 95%, and

97.5%. (**A**) The proportions are shown for each antibiotic at three confidence levels (90%, 95%,

and 97.5%), considering all possible numbers of input AST results. **(B)** The proportions are

shown as a function of the number of input AST results (90% confidence level).

Figure 6. The proportion of correct predictions with respect to the number of labels in the

prediction set for *P. aeruginosa* isolates. The proportion of correct predictions with a single

label (light) and multiple labels (dark), major errors (MEs), and very major errors (VMEs) with a

single label (light) and empty set (dark) predictions for resistant (R) and susceptible (S) isolates,

shown for each antibiotic and pathogen using three different confidence levels: 90%, 95%, and

97.5%. **(A)** The proportions are shown for each antibiotic at three confidence levels (90%, 95%,

and 97.5%), considering all possible numbers of input AST results. **(B)** The proportions are

shown as a function of the number of input AST results (90% confidence level).

REFERENCES

1. World Health Organization. 2015. Global action plan on antimicrobial resistance.

2. Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Aguilar GR, Gray A, Han C, Bisignano C,

Rao P, Wool E, others. 2022. Global burden of bacterial antimicrobial resistance in 2019: a

systematic analysis. The Lancet 399:629–655.

3. O'Neill J. 2016. Tackling drug-resistant infections globally: final report and

recommendations. Government of the United Kingdom.

4. Mercer DK, Torres MDT, Duay SS, Lovie E, Simpson L, von Köckritz-Blickwede M, de la

Fuente-Nunez C, O'Neil DA, Angeles-Boza AM. 2020. Antimicrobial Susceptibility

Testing of Antimicrobial Peptides to Better Predict Efficacy. Frontiers in Cellular and

Infection Microbiology 10:326.

5.   Jorgensen JH, Ferraro MJ. 1998. Antimicrobial susceptibility testing: general principles and

contemporary practices. Clin Infect Dis 26:973–980.

6.   Friedman ND, Temkin E, Carmeli Y. 2016. The negative impact of antibiotic resistance.

Clin Microbiol Infect 22:416–422.

7.   Bassetti M, Rello J, Blasi F, Goossens H, Sotgiu G, Tavoschi L, Zasowski EJ, Arber MR,

McCool R, Patterson JV, others. 2020. Systematic review of the impact of appropriate

versus inappropriate initial antibiotic therapy on outcomes of patients with severe bacterial

infections. International journal of antimicrobial agents 56:106184.

8.   Battle SE, Bookstaver PB, Justo JA, Kohn J, Albrecht H, Al-Hasan MN. 2016. Association

between inappropriate empirical antimicrobial therapy and hospital length of stay in Gram-

negative bloodstream infections: stratification by prognosis. Journal of Antimicrobial

Chemotherapy 72:299–304.

9.   Kumar A, Ellis P, Arabi Y, Roberts D, Light B, Parrillo JE, Dodek P, Wood G, Kumar A,

Simon D, others. 2009. Initiation of inappropriate antimicrobial therapy results in a fivefold

reduction of survival in human septic shock. Chest 136:1237–1248.

10. van den Bosch C, Hulscher ME, Akkermans RP, Wille J, Geerlings SE, Prins JM. 2017.

Appropriate antibiotic use reduces length of hospital stay. Journal of Antimicrobial

Chemotherapy 72:923–932.

11. Dolejska M, Papagiannitsis CC. 2018. Plasmid-mediated resistance is going wild. Plasmid
99:99–111.

12. Johnning A, Karami N, Tång Hallbäck E, Müller V, Nyberg L, Buongermino Pereira M,
Stewart C, Ambjörnsson T, Westerlund F, Adlerberth I, Kristiansson E. 2018. The
resistomes of six carbapenem-resistant pathogens - a critical genotype-phenotype analysis.
Microb Genom 4.

13. Dias SP, Brouwer MC, van de Beek D. 2022. Sex and gender differences in bacterial
infections. Infection and Immunity 90:e00283-22.

14. Murray KA, Preston N, Allen T, Zambrana-Torrelio C, Hosseini PR, Daszak P. 2015.
Global biogeography of human infectious diseases. Proceedings of the National Academy
of Sciences 112:12746–12751.

15. Yelin I, Snitser O, Novich G, Katz R, Tal O, Parizade M, Chodick G, Koren G, Shalev V,
Kishony R. 2019. Personal clinical history predicts antibiotic resistance of urinary tract
infections. Nature medicine 25:1143–1152.

16. Kanjilal S, Oberst M, Boominathan S, Zhou H, Hooper DC, Sontag D. 2020. A decision
algorithm to promote outpatient antimicrobial stewardship for uncomplicated urinary tract
infection. Science Translational Medicine 12:eaay5067.

17. Stracy M, Snitser O, Yelin I, Amer Y, Parizade M, Katz R, Rimler G, Wolf T, Herzel E,
Koren G, others. 2022. Minimizing treatment-induced emergence of antibiotic resistance in
bacterial infections. Science 375:889–894.

18.   Secinaro S, Calandra D, Secinaro A, Muthurangu V, Biancone P. 2021. The role of

artificial intelligence in healthcare: a structured literature review. BMC Medical Informatics

and Decision Making 21:1–23.

19.   Yu K-H, Beam AL, Kohane IS. 2018. Artificial intelligence in healthcare. Nature

biomedical engineering 2:719–731.

20.   Benjamens S, Dhunnoo P, Meskó B. 2020. The state of artificial intelligence-based FDA-

approved medical devices and algorithms: an online database. NPJ digital medicine 3:1–8.

21.   Rajpurkar P, Chen E, Banerjee O, Topol EJ. 2022. AI in health and medicine. Nature

Medicine 28:31–38.

22.   Ali T, Ahmed S, Aslam M. 2023. Artificial Intelligence for Antimicrobial Resistance

Prediction: Challenges and Opportunities towards Practical Implementation. Antibiotics

(Basel) 12:523.

23.   Sakagianni A, Koufopoulou C, Feretzakis G, Kalles D, Verykios VS, Myrianthefs P,

Fildisis G. 2023. Using Machine Learning to Predict Antimicrobial Resistance―A

Literature Review. Antibiotics 12:452.

24.   The Medical Futurist. 2022. FDA-approved A.I.-based algorithms.

https://medicalfuturist.com/fda-approved-ai-based-algorithms/

25.   Begoli E, Bhattacharya T, Kusnezov D. 2019. The need for uncertainty quantification in

machine-assisted medical decision making. Nature Machine Intelligence 1:20–23.

26.   Kompa B, Snoek J, Beam AL. 2021. Second opinion needed: communicating uncertainty in
medical machine learning. NPJ Digital Medicine 4:1–6.

27.   Feretzakis G, Loupelis E, Sakagianni A, Kalles D, Martsoukou M, Lada M, Skarmoutsou
N, Christopoulos C, Valakis K, Velentza A, others. 2020. Using machine learning
techniques to aid empirical antibiotic therapy decisions in the intensive care unit of a
general hospital in Greece. Antibiotics 9:50.

28.   Feretzakis G, Sakagianni A, Loupelis E, Kalles D, Skarmoutsou N, Martsoukou M,
Christopoulos C, Lada M, Petropoulou S, Velentza A, others. 2021. Machine learning for
antibiotic resistance prediction: A prototype using off-the-shelf techniques and entry-level
data to guide empiric antimicrobial therapy. Healthcare informatics research 27:214–221.

29.   Devlin J, Chang M-W, Lee K, Toutanova K. 2018. Bert: Pre-training of deep bidirectional
transformers for language understanding. arXiv preprint arXiv:181004805.

30.   Brown T, Mann B, Ryder N, Subbiah M, Kaplan JD, Dhariwal P, Neelakantan A, Shyam P,
Sastry G, Askell A, others. 2020. Language models are few-shot learners. Advances in
neural information processing systems 33:1877–1901.

31.   Vaswani A, Shazeer N, Parmar N, Uszkoreit J, Jones L, Gomez AN, Kaiser \Lukasz,
Polosukhin I. 2017. Attention is all you need. Advances in neural information processing
systems 30.

32.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A,
Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E,

Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals

O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein

structure prediction with AlphaFold. Nature 596:583–589.

33. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD,

Winther O, Brunak S, von Heijne G, Nielsen H. 2022. SignalP 6.0 predicts all five types of

signal peptides using protein language models. Nature biotechnology 1–3.

34. Vovk V. 2012. Conditional validity of inductive conformal predictors, p. 475–490. *In* Asian

conference on machine learning. Proceedings of Machine Learning Research.

35. Vazquez J, Facelli JC. 2022. Conformal Prediction in Clinical Medical Sciences. Journal of

Healthcare Informatics Research 1–12.

36. Reller LB, Weinstein M, Jorgensen JH, Ferraro MJ. 2009. Antimicrobial susceptibility

testing: a review of general principles and contemporary practices. Clinical infectious

diseases 49:1749–1755.

37. European Centre for Disease Prevention and Control, European Medicines Agency. 2009.

The bacterial challenge : time to react : a call to narrow the gap between multidrug-resistant

bacteria in the EU and the development of new antibacterial agents. Publications Office.

38. Ahmad M, Khan AU. 2019. Global economic impact of antibiotic resistance: A review.

Journal of global antimicrobial resistance 19:313–316.

39. Vovk V, Gammerman A, Shafer G. 2005. Algorithmic Learning in a Random World.

Springer-Verlag, Berlin, Heidelberg.

40.   Papadopoulos H. 2008. Inductive Conformal Prediction: Theory and Application to Neural

707   NetworksTools in Artificial Intelligence. IntechOpen, Rijeka.

708   41.   Bush K, Bradford PA. 2020. Epidemiology of β-lactamase-producing pathogens. Clinical

709   microbiology reviews 33:e00047-19.

710   42.   Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A, Allesoe

711   RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B, Werner G,

712   Bender JK, Stingl K, Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar S, Westh H,

713   Pinholt M, Anjum MF, Duggett NA, Kempf I, Nykäsenoja S, Olkkola S, Wieczorek K,

714   Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C, Lund O, Aarestrup FM. 2020.

715   ResFinder 4.0 for predictions of phenotypes from genotypes. Journal of Antimicrobial

716   Chemotherapy 75:3491–3500.

717   43.   Olsson H, Kartasalo K, Mulliqi N, Capuccini M, Ruusuvuori P, Samaratunga H, Delahunt

718   B, Lindskog C, Janssen EA, Blilie A, others. 2022. Estimating diagnostic uncertainty in

719   artificial intelligence assisted pathology using conformal prediction. Nature

720   communications 13:7761.

Figure 1. The proposed model architecture and uncertainty control. (A) The architecture of the proposed model. Both input sequences start with a classification word, *CLS*, followed by the bacterial species name. One of the input sequences also contains patient information (country, gender, age, and sampling date), while the other includes available antibiotic susceptibility testing (AST) data. Both input sequences are fixed to lengths L=5 for the patient information and L=21 for the AST data, padded with the *PAD* word. We use a linear embedding to represent the input words numerically, which are fed into the transformer encoders. The first vectors (*CLS*) from the outputs of each encoder are combined into a linear model and fed to *m* independent neural networks, each representing one antibiotic. The neural network outputs are two-dimensional vectors indicating susceptibility and resistance, respectively. **(B)** Uncertainty control. The neural network outputs undergo a softmax rescaling. A calibration dataset is used to build empirical distributions of conformity for resistant and susceptible predictions for each pathogen and antibiotic. The prediction regions for the data points k and $k + 1$ are built based on the deviation of the observed softmax score from the empirical distribution and the confidence interval threshold. See Methods for full details.

Fig. 2. Performance of the transformer model. Results from the train and testing datasets: **(A)** F1-score, **(B)** major error (ME) rate, and very major error (VME) rate for the transformer model shown for each antibiotic and pathogen. Note that, due to data restrictions, not all antibiotics are assessed for all three pathogens.

746
747

Fig. 3. The predictive performance of the model as a function of the number of antibiotic susceptibility testing (AST) results included in the input. Major error (ME) and very major error (VME) rates from light to dark: 4 to 8 AST results are shown for each antibiotic and pathogen. Note that, due to data restrictions, not all antibiotics are assessed for all three pathogens.

Fig. 4. The proportion of correct predictions with respect to the number of labels in the prediction set for *E. coli* isolates. The proportion of correct predictions with a single label (light) and multiple labels (dark), major errors (MEs), and very major errors (VMEs) with a single label (light) and empty set (dark) predictions for resistant (R) and susceptible (S) isolates, shown for each antibiotic and pathogen using three different confidence levels: 90%, 95%, and 97.5%. **(A)** The proportions are shown for each antibiotic at three confidence levels (90%, 95%, and 97.5%), considering all possible numbers of input AST results. **(B)** The proportions are shown as a function of the number of input AST results (90% confidence level).

767
768

Fig. 5. The proportion of correct predictions with respect to the number of labels in the prediction set for *K. pneumoniae* isolates. The proportion of correct predictions with a single label (light) and multiple labels (dark), major errors (MEs), and very major errors (VMEs) with a single label (light) and empty set (dark) predictions for resistant (R) and susceptible (S) isolates, shown for each antibiotic and pathogen using three different confidence levels: 90%, 95%, and 97.5%. **(A)** The proportions are shown for each antibiotic at three confidence levels (90%, 95%, and 97.5%), considering all possible numbers of input AST results. **(B)** The proportions are shown as a function of the number of input AST results (90% confidence level).

Fig. 6. The proportion of correct predictions with respect to the number of labels in the prediction set for *P. aeruginosa* isolates. The proportion of correct predictions with a single label (light) and multiple labels (dark), major errors (MEs), and very major errors (VMEs) with a single label (light) and empty set (dark) predictions for resistant (R) and susceptible (S) isolates, shown for each antibiotic and pathogen using three different confidence levels: 90%, 95%, and 97.5%. **(A)** The proportions are shown for each antibiotic at three confidence levels (90%, 95%, and 97.5%), considering all possible numbers of input AST results. **(B)** The proportions are shown as a function of the number of input AST results (90% confidence level).