Characterizing fibrinolysis under flow conditions: a simplified mesoscale 3D model

Remy Petkantchin[1*]**, Alexandre Rousseau**[2]**, Omer Eker**[3,4]**, Karim Zouaoui Boudjeltia**[2]**, Franck Raynaud**[1†]**, Bastien Chopard**[1*†]**, and the INSIST investigators**[5]

[1]Scientific and Parallel Computing Group, Computer Science Department, University of Geneva, Switzerland; [2]Laboratory of Experimental Medicine (ULB222), Faculty of Medicine, Université libre de Bruxelles, CHU de Charleroi, Belgium; [3]Department of Neuroradiology, Hôpital Pierre Wertheimer, Hospices Civils de Lyon, Lyon, France; [4]CREATIS Laboratory, UMR 5220, U1206, Université Lyon, INSA-Lyon, Université Claude Bernard Lyon 1, UJM-Saint Etienne, CNRS, Inserm, Lyon, France; [5]see end of this article

*For correspondence:
remy.petkantchin@unige.ch (RP);
bastien.chopard@unige.ch (BC)

[†]These authors share last authorship

Abstract One of the routine clinical treatments to eliminate ischemic stroke thrombi is injecting a biochemical product into the patient's bloodstream, which breaks down the thrombi's fibrin fibers: intravenous or intravascular thrombolysis. Unfortunately, however, the successful outcome of this procedure is still more of an exception than the rule. Understanding why from clinical evidence is close to impossible, given that no existing tool can track the lysis at the thrombus scale and that time is limited due to the emergency. For that reason, we conducted *in vitro* flow-driven fibrinolysis experiments to study pure fibrin thrombi breakdown in controlled conditions and observed that the lysis front evolved non-linearly in time. To understand these results, we developed an analytical 1D lysis model in which the thrombus is considered a porous medium. The lytic cascade is reduced to a second-order reaction involving fibrin and a surrogate pro-fibrinolytic agent. The model was able to reproduce the observed lysis evolution under the assumptions of constant fluid velocity and lysis occurring only at the front. For adding complexity, such as clot heterogeneity or complex flow conditions, we propose a 3-dimensional mesoscopic numerical model of blood flow and fibrinolysis, which validates the analytical model's results. Such a numerical model could help us better understand the spatial evolution of the thrombi breakdown, extract the most relevant physiological parameters to lysis efficiency, and possibly explain the failure of the clinical treatment. These findings suggest that even though real-world fibrinolysis is a complex biological process, a simplified model can recover the main features of lysis evolution.

Introduction

Ischemic stroke, already reported by Hippocrates in 400 B.C. *Nilsen* (*2010*), is still among the first causes of disability and death in developed countries *INSIST* (*2017*), causing around 4.5 million deaths globally per year *Donkor* (*2018*). It occurs when an aggregate of blood components clogs a brain artery, thus impeding sufficient irrigation. These thrombi, or blood clots, are mainly composed of red blood cells (RBCs) and platelets bound together by fibrin fibers, although they may contain several other biological entities. The current treatment options are either thrombolysis alone, a biochemical breakdown of the thrombus, or used with a mechanical removal of the clot

(such as thrombectomy, or thrombus aspiration). The latter, whose efficacy has been demonstrated *Fransen et al.* (*2014*) requires endovascular treatment and is indicated only in large, easily accessible arteries. The former, less invasive and applicable in first aid within a few hours after stroke symptoms onset, is done by injecting a pro-fibrinolytic macromolecule into the bloodstream. This agent, typically tissue-Plasminogen-Activator (tPA), binds to the fibrin fibers to facilitate the conversion of plasminogen to plasmin, which will then cut the fibrin strands and result in thrombus breakdown. Although the biochemistry of fibrinolysis is well understood *Weisel* (*2005*), our knowledge of many aspects of the complete process *in vivo* and *in vitro* remains sparse *Petkantchin et al.* (*2022*); *Bannish and Hudson* (*2021*). For example, the permeability of the thrombus, which is central to the transport of active macromolecules in the bloodstream, is only estimated *via* indirect measurements like perviousness *Arrarte Terreros et al.* (*2020*); the exact amount of drug that reaches the thrombus is unknown; even the dynamics of fibrin strands degradation is still actively investigated *Raynaud et al.* (*2021*); *Lynch et al.* (*2022*); *Collet J. P. et al.* (*2000*); *Longstaff and Kolev* (*2015*); *Bucay et al.* (*2015*); *Feller et al.* (*2018*). Even though patients who receive early intervention show a better three-month prognosis, about fifty percent fail to respond, and the reasons for this are unknown. Hence, theoretical, experimental, and clinical research are essential to understand the underlying mechanisms of the variability of treatment outcomes and eventually improve further drug treatment development.

We present in this study a combination of experiments and modeling to investigate the dynamics of flow-driven lysis profiles of fibrin-pure clots. Experimentally, we monitor the progress of the lysis and estimate the time evolution of the lysis profile and the dynamics of the flow seeping through the clot. The experiments show that the lysis accelerates in time. However, the flow remains constant or slightly increases during the lysis, thus suggesting that the recanalization would occur late. We translate these observations into a simple one-dimensional lysis model with constant flow and all pro-fibrinolytic agents aggregated into a single species, called anti-Fibrin Agent (anti-FA), that is blocked at the front of the clot and interacts through a second-order reaction. This model can be solved analytically and accurately predicts lysis profiles given the initial conditions.

Several numerical models have already been developed *Diamond and Anand* (*1993*); *Zidansek et al.* (*1995*); *Anand et al.* (*2005*); *Bajd and Serša* (*2013*); *Bannish et al.* (*2014*); *Miyazaki and Yamaguchi* (*2003*); *Piebalgs and Xu* (*2015*); *Piebalgs et al.* (*2018*); *Gu et al.* (*2019*). Most of these models couple pharmacokinetics with the transport of proteins to achieve the lysis of millimetric clots. However, their main drawbacks are relying on non-physiological protein concentrations or flow conditions. Furthermore, numerical models often rely on indirect measurements from previously published data, such as total lysis time, to validate simulation results. Moreover, by design, they do not seem well-suited to tackle more complex situations, such as clots with a heterogeneous composition. Finally, we gather our experimental and theoretical results to adopt a different approach. Instead of solving the kinetics of several species and the Navier-Stokes equations, we rely on a mesoscopic approach in which the flow is computed with the open-source lattice-Boltzmann (LB) library Palabos *Latt et al.* (*2020*). The fibrin clot is described as a porous material, and its porosity changes locally with the progress of the lysis. Our framework can incorporate heterogeneity directly into clot porosity and composition to assess to what extent the thrombus heterogeneity can impact stroke treatment and determine whether it is possible to design novel protocols when considering thrombolysis in acute ischemic stroke.

Material and Methods

Experimental setup

Pasteur pipettes, with the tips cut off, were used (VWR). To form the fibrin clot at a height suitable for the experiments, we placed μl of glycerol (Merck) in the bottom of the pipettes Then we place the tubes in a support. The glycerol level rises in the tube to the desired level. For fibrin clot formation, we prepared solutions containing different concentrations of fibrinogen: 2.4 or $3.2\,\mathrm{mg \cdot ml^{-1}}$ (Merck).

Plasminogen (Merck) is at a final concentration of $13\,\mu\text{g}\cdot\text{ml}^{-1}$, Owren (Stago) as a buffer. We add thrombin (Stago) at a final concentration of $2\,\text{U}\cdot\text{ml}^{-1}$ to form the clots (final volume $800\,\mu$l). Then, we incubated for 15 minutes at 37°C. The Pasteur pipettes are then placed on a stand and a solution is added to a height of 2 cm. This consists of Owren buffer, tPA (Merck) at a final concentration of $10\,\text{ng}\cdot\text{ml}^{-1}$, FITC-labelled BSA (Abcam) at a final concentration of $0.1\,\text{mg}\cdot\text{ml}^{-1}$, plasminogen (Merck) at a final concentration of $13\,\mu\text{g}\cdot\text{ml}^{-1}$. Lysis is induced by adding medium every 10 minutes to the mark which is 2 cm from the level of the previously formed clot. We record a photo in daylight and a photo under a UV lamp to visualise the migration of the labelled BSA.

A variant of the experiment without plasminogen was performed to measure the migration of tPA. The tPA is added only in the supernatant. The clot is collected after a migration time of 30, 60, or 90 minutes. . The clot is then cut into four equal parts. Plasminogen is added and the whole is incubated for 16 hours. The lysate is then collected for tPA assay by Elisa (Invitrogen). In another experiment, the clot is allowed to lysate over the first quarter of its length. The clot is then removed and cut into four equal parts, left to incubate for 16 hours and the lysate is recovered for tPA determination by Elisa (Invitrogen).

Minimal one-dimensional model of fibrinolysis

Typical models of fibrinolysis consist of ten or more equations to account for the different reactions between species, free or bound, involved in the fibrinolytic cascade *Piebalgs et al.* (*2018*); *Diamond and Anand* (*1993*); *Bannish et al.* (*2014*); *Shibeko et al.* (*2020*). In *Shibeko et al.* (*2020*), the authors have shown that the number of equations necessary to describe the lysis of fibrin pure clot can be reduced to only 6. Hence, we wonder whether it is possible to lower this number even more, while preserving the main features of the lysis. To do so, we propose a one-dimensional minimal model of lysis that relies on few assumptions and that can be solved analytically. This model aims not to replace previous detailed models of fibrinolysis but rather to identify a set of minimal rules sufficient to reproduce the time evolution of the experimentally observed lysis profiles. We consider that the system is made of only two species. The first one is the fibrin (F), and the second, the anti-FA ($\overline{F}$), a fictive aggregate that accounts for the activation, inhibition, and binding of the different components. We further assume that the fluid velocity u_f is constant, and the anti-FA, which does not diffuse, remains blocked at the front of the clot to mimic the binding of profibrinolytic species, until the fibrin concentration decreases below a certain threshold F^*. We finally assume that all the profibrinolytic processes, such as activation of plasminogen into plasmin, can be included into a single reaction rate k_1. The equations for the reaction and the transport of our two species finally read as:

$$\partial_t F = -k_1 F \overline{F}, \tag{1}$$

$$\partial_t \overline{F} = -u_\text{f} \partial_x \overline{F}\, \Theta\left(F^* - F\right), \tag{2}$$

where $\Theta(x)$ is the Heaviside step function, with $\Theta(x) = 1$ if $x > 0$ and $\Theta(x) = 0$ otherwise. We write F and $\overline{F}$ for $F(x,t)$ and $\overline{F}(x,t)$ respectively for readability, and ∂_i denotes $\frac{\partial}{\partial i}$. To build the model upon the observation that tPA does not penetrate deep into the bulk of the clot, we limit the penetration of anti-FA to a thickness Δ. Thus, we partition the thrombus of initial length L and fibrin concentration F_0, into n slices $x_0^\Delta, x_1^\Delta, ..., x_{n-1}^\Delta$ of thickness Δ. We choose the flow to be directed from the first slice x_0^Δ to the last x_{n-1}^Δ and impose a constant concentration $\overline{F_0}$ at the inlet. Under these considerations, the concentration $\overline{F}(x_0^\Delta, t)$ of anti-FA that accumulates at the first slice x_0^Δ evolves as:

$$\overline{F}(x_0^\Delta, t) = \overline{F_0}\frac{u_\text{f}}{\Delta}t \text{ , if } t \leq t_0. \tag{3}$$

After replacing this term in eq (1) and integrating, we get:

$$\int_{F_0}^{F^*} \frac{dF(x_0^\Delta, t)}{F(x_0^\Delta, t)} = -k_1 \overline{F}_0 \frac{u_{\rm f}}{\Delta} \int_0^{t_0} t \, dt$$

$$\Rightarrow \log\left(\frac{F^*}{F_0}\right) = -k_1 \overline{F}_0 \frac{u_{\rm f}}{2\Delta} t_0^2,$$

132   where t_0 is the time to lyse the first slice, and satisfies $F(x_0^\Delta, t_0) = F^*$. This yields:

$$t_0 = \sqrt{\frac{2\Delta \log\left(\frac{F_0}{F^*}\right)}{k_1 u_{\rm f} \overline{F}_0}}. \tag{4}$$

133   Recursively (see section S1), we obtain the time t_n to lyse the full clot of length $L = n\Delta$:

$$t_n = \frac{\Delta}{u_{\rm f}} + \sqrt{t_{n-1}^2 + \frac{2\Delta}{k_1 u_{\rm f} \overline{F}_0} \log\left(\frac{F_0}{F^*}\right)}. \tag{5}$$

134   If Δ is small enough compared to the length of the clot L, this result is independent of Δ (see
135   Appendix and Figure S2).

Mesoscopic three-dimensional model of fibrinolysis

137   We have demonstrated that considering only one profibrinolytic species blocked at the front of
138   the clot is sufficient to recover non-trivial lysis profiles. We now incorporate those findings in a
139   more elaborated three-dimensional model that accounts for the fluid simulation, the thrombus as
140   a porous medium, and the lysis. These elements are summarized in Figure 1.

Fluid computation

In order to simulate the fluid flow, we rely on the so-called lattice-Boltzmann (LB) method, which al-
lows the simulation of fluids without explicitly solving the Navier-Stokes equations. The LB method
can be seen as a latticed discretization of the kinetic theory, where the probability density function
is replaced by fictitious particles colliding and streaming on the lattice. Inspired by Lattice Gas Au-
tomata *Frisch et al.* (*1986*), the LB method is an efficient alternative to the NS equations *He and*
Luo (*1997a*,b); *Chopard et al.* (*2002*); *Krüger et al.* (*2017*); *Succi* (*2018*), that has been widely used
to address biomedical problems *Malaspinas et al.* (*2015*); *Li et al.* (*2019*); *Kotsalos et al.* (*2019*).
Briefly, it solves the Boltzmann equation, which describes the evolution of so-called "fluid" parti-
cles in a regularly discretized lattice. The spatial discretization enforces a set of q discrete velocities
c_i, $i = 0, 1, ..., q-1$ for the fluid particle densities f_i, also called populations. For example, f_1 is the
density of fluid particles moving with velocity c_1. Figure 1A depicts a 2D domain denoted D2Q9
(2 spatial dimensions and nine discrete velocities associated with nine directions). Macroscopic
variables such as density ρ and momentum $\rho \mathbf{u}$ are recovered by computing the moments of f_i:

$$\rho(\mathbf{x}, t) = \sum_i f_i(\mathbf{x}, t), \tag{6}$$

$$\rho(\mathbf{x}, t)\mathbf{u}(\mathbf{x}, t) = \sum_i \mathbf{c_i} f_i(\mathbf{x}, t). \tag{7}$$

155   The temporal evolution of the populations $f_i(\mathbf{x}, t)$ is a two steps process. First, the collision
156   step computes at each lattice position $\mathbf{x}$ the resulting scattered populations $f_i^*(\mathbf{x}, t)$. The streaming
157   step propagates the populations $f_i^*(\mathbf{x}, t)$ to the neighboring nodes. More formally, the scattered
158   populations are given by:

$$f_i^*(\mathbf{x}, t) = f_i(\mathbf{x}, t) + \Omega_i(f). \tag{8}$$

159   where $\Omega_i(f)$ is the collision operator. In this paper, we consider a relaxation-type collision process
160   between populations, the Bhatnagar-Gross-Krook (BGK) collision operator *Bhatnagar et al.* (*1954*):

$$\Omega_i(f) = -\frac{f_i(\mathbf{x}, t) - f_i^{eq}(\mathbf{x}, t)}{\tau} \Delta t, \tag{9}$$

where Δt is the temporal discretization, and τ is the model relaxation time associated with the fluid kinematic viscosity. The equilibrium populations $f_i^{eq}(\mathbf{x}, t)$ are expressed as follows (see for instance *Shan et al.* (*2006*)):

$$f_i^{eq}(\mathbf{x}, t) = w_i \, \rho \left(1 + \frac{\mathbf{u} \cdot \mathbf{c_i}}{c_s^2} + \frac{(\mathbf{u} \cdot \mathbf{c_i})^2}{2c_s^4} - \frac{\mathbf{u} \cdot \mathbf{u}}{2c_s^2} \right) \tag{10}$$

where w_i are weighting factors that depend on the dimension and the number of discretized velocities. c_s is the speed of sound in the lattice, defined by the lattice topology (see for instance *Krüger et al.* (*2017*)). Finally, the streaming step propagates the scattered populations:

$$f_i(\mathbf{x} + \mathbf{c_i}\Delta t, t + \Delta t) = f_i^*(\mathbf{x}, t). \tag{11}$$

Clot description

Fibrinogen is a soluble blood protein made up of three pairs of polypeptide chain, and converted into fibrin by the cleavage of the fibrinopeptide by thrombin. Due to a change in solubility, fibrin monomers polymerize into protofibrils that aggregate into fibers that merge to form a branched network of fibers. The resulting fibrin fibers' diameter range from 10^{-8} to 10^{-7} m, with pores' diameter from 10^{-7} to 10^{-5} m *Diamond and Anand* (*1993*), and the thrombi are typically millimetric to centimetric *Riedel et al.* (*2011*). Simulating the lysis of such an intermingled mesh at the fiber scale seems unrealistic and computationally expensive, considering the size and time scales involved. We aim to simulate the lysis of centimetric clots over hundreds of minutes while state-of-the-art detailed molecular models of fibers operate at nanometer and microsecond *Yesudasan et al.* (*2018*). Instead, with a coarser description, one can treat clots as fibrous porous media *Zidansek et al.* (*1995*); *Piebalgs et al.* (*2018*); *Wufsus et al.* (*2013*), which can be done in the context of LB *Li et al.* (*2014*) with the so-called partial bounce-back (PBB) method.

Partial bounce-back

When the populations f_i hit an obstacle during the collision step, their velocities get reversed after the streaming. This procedure leads to a zero fluid velocity relative to the boundary of the obstacle and is called solid bounce-back boundary:

$$f_i(\mathbf{x_b}, t + \Delta t) = f_{\hat{i}}^*(\mathbf{x_b}, t) \tag{12}$$

where the subscript b denotes a point at the boundary and $\hat{i}$ is the opposite lattice direction of i. The PBB method is an extension of the solid bounce-back boundary. The underlying idea is to assign a continuous fraction of bounce-back to each volume element, whether none, part, or all of the fluid is reflected at the boundary (Figure 1B). In LB terms, the outgoing populations in direction i are calculated as the weighted sum of the post-collision populations $f_i^*(\mathbf{x}, t)$ and the bounced populations $f_{\hat{i}}(x, t)$ *Walsh et al.* (*2009*):

$$f_i(\mathbf{x} + \mathbf{c_i}\Delta t, t + \Delta t) = (1 - \gamma)f_i^*(\mathbf{x}, t) + \gamma f_{\hat{i}}(\mathbf{x}, t) \tag{13}$$

where γ is the fraction of bounce-back and ranges between 0 (completely fluid voxel) and 1 (completely solid voxel). Note that for $\gamma = 0$, we recover the usual streaming of a fluid node (Eq. (11)), while for $\gamma = 1$, we have the equation of the solid bounce-back boundary (Eq. (12)). Consistently, the macroscopic velocity u_{f} is computed as *Walsh et al.* (*2009*):

$$\mathbf{u_f}(\mathbf{x}, t) = \frac{(1 - \gamma)}{\rho(\mathbf{x}, t)} \sum_i \mathbf{c_i} f_i(\mathbf{x}, t) \tag{14}$$

194 Permeability model

The permeability is a macroscopic quantity that measures the ability of a porous medium to let a
fluid flow through it. It is associated with the material's porosity, the shape of the pores, and their
local arrangements. Consequently, the permeability law of a given material is generally not known
a priori. Instead, it has to be measured experimentally or simulated, with Darcy's law, for instance.
In **Walsh et al.** (**2009**), the authors derived analytically the relationship between the permeability k
and the bounce-back fraction:

$$\gamma = \frac{1}{1 + \frac{2k}{\nu \Delta t}}, \tag{15}$$

where ν is the kinematic fluid viscosity. However, Eq. (15) only relates the fraction of bounce-back
γ to the permeability k. It implies that in order to simulate a porous material with the characteristic
permeability of a thrombus, we have to incorporate in our model a permeability law that provides a
value of k, knowing the radius of the fibers (R_f) and the relative volume occupied by the fibers (solid
fraction n_s^*). Several studies established permeability laws, either experimentally **Davies** (**1952**)
or numerically using randomly organized fibers **Clague et al.** (**2000**) or two-dimensional periodic
square arrays of cylinders **Jackson and James** (**1986**). These different laws yield to similar results,
but **Wufsus et al.** (**2013**) show that the permeability of *in vitro* fibrin clots is best described by Davies'
permeability law **Davies** (**1952**):

$$k_{\mathrm{D}}(R_f, n_s^*) = R_f^2 \left(16 n_s^{*1.5}(1 + 56 n_s^{*3}) \right)^{-1}. \tag{16}$$

210 Finally, we obtain the relationship $\gamma \left(n_s^*, R_f \right)$ by replacing in Eq. (15) the value of k by Davies' formula
211 (16). We show (see Appendix and Figure S 6) that the PBB method can be extended to accurately
212 simulate different types of fibrous porous media over a wide range of solid fractions and reproduce
213 the permeability law of fibrin-pure clots.

214 3D Lysis model

We previously described the clot as a collection of PBB voxels repelling the fluid with an intensity
that depends on the local concentration of fibrin. To model the lysis of the clot, we use a two-
species model as already proposed. Unlike the one-dimensional model, the fluid velocity is not
constant but computed with the LB method that considers the clot's permeability and the change
of the solid fraction due to the decrease of fibrin concentration. The governing equations for the
concentrations of the two species become:

$$\partial_t F = -k_1 F \overline{F}, \tag{17}$$

$$\partial_t \overline{F} = \left(D_{\overline{F}} \nabla^2 \overline{F} - \nabla \cdot \left(\mathbf{u}_{\mathrm{f}} \, \overline{F} \right) \right) \Theta(F^* - F), \tag{18}$$

here again, we write F and $\overline{F}$ for $F(\mathbf{x}, t)$ and $\overline{F}(\mathbf{x}, t)$ respectively for readability. For numerical rea-
sons, we did not consider the anti-FA $\overline{F}$ as a concentration field but treated it as an ensemble of
meso-particles, each carrying a certain amount of anti-FA. The lysis reaction occurs when meso-
particles encounter fibrin voxels. The diffusion coefficient of anti-FA $D_{\overline{F}}$ is taken as the diffusion
coefficient of tPA, $D_{\overline{F}} = 3.3 \cdot 10^{-11}$ m^2·s^{-1} **Shibeko et al.** (**2020**). The numerical resolution of equation
(17) is done by finite forward differences:

$$F(\mathbf{x}, t') = F(\mathbf{x}, t) - k_1 F(\mathbf{x}, t) \overline{F}(\mathbf{x}, t) \Delta t \tag{19}$$

227 where $t' = t + \Delta t$, and the transport of anti-FA (18) is a modified version of the built-in solver of
228 the Palabos library, see section S1. We consider that anti-FA cleaves the strands radially and uni-
229 formly **Diamond and Anand** (**1993**); **Blinc et al.** (**2000**); **Gabriel et al.** (**1992**). As the concentration of
230 fibrin decreases, the radius of the fibers R_f shrinks accordingly **Diamond and Anand** (**1993**):

$$R_f\left(F(\mathbf{x},t')\right) = \sqrt{R_f(F_0)^2 - \frac{F_0 - F(\mathbf{x},t')}{C\frac{L_{\Delta V}}{\Delta V}}} \tag{20}$$

where $L_{\Delta V}$ is the total length of fibrin strands in volume ΔV, computed based on the fibrin concentration and fibers density, and C is a series of constants *Diamond and Anand* (*1993*). Then, upon reaction (19), the solid fraction n_s^* of each voxel needs to be recomputed as:

$$n_s^*\left(F(\mathbf{x},t')\right) = \pi R_f^2\left(F(\mathbf{x},t')\right)\frac{L_{\Delta V}}{\Delta V}. \tag{21}$$

The changes in R_f and n_s^* during lysis thus induce a local change in the permeability k, through Davies' equation (16).

Results

We first present the experimental results, and compare them with the simplified analytical model. This also allows us to estimate a value for the model reaction parameter k_1. In a second step, we show the numerical model results when varying the fibrin and anti-FA concentrations. Finally, we briefly explore the lysis of spatially heterogeneous thrombi with the numerical model.

Penetration of tPA through fibrin pure clots

We first test the hypothesis that anti-FA remains blocked at the fibrin sites. Previous detailed molecular models already reported that due to the strong affinity of tPA with fibrin, most of the tPA is bound at the front of the clot *Shibeko et al.* (*2020*); *Diamond and Anand* (*1993*). However, it has not been assessed experimentally to what extent tPA remains blocked or perfuses through the clot under permeation conditions. Figure 2A schematizes the experiments where we measure the concentration of bound tPA in different regions of the clot (top most part HH, middle top part H, middle bottom part B, and bottom most part BB) after thirty minutes, sixty minutes and two hours of exposure at the clot-tPA interface. Overall, we find that tPA concentration in the HH part is significantly greater than in other parts of the clot and ten to twenty times greater than in the supernatant (Figures 2B-D). As the exposure to tPA increased, the concentration in the regions H, B, and BB increased slightly. However, it remained significantly lower than the concentration in the supernatant (Figure 2C,D). The parts of the clot in which we measure the concentration of bound tPA are considerably bigger than the regions where we assume the anti-FA to be blocked. However, we have demonstrated formally that the lysis profiles do not depend on the size of the blocking area, if it is chosen small enough (Appendix and Figure S2). For the thinner clots, the pore size can slightly hinder the transport properties of the larger molecules. We collected the solution in the vicinity of the clot front and did not measure any excess of tPA in this region (data not shown), indicating that tPA can enter the clot but is rapidly bound to fibrin. From a modeling perspective, these results support our hypothesis that anti-FA can be blocked at fibrin sites.

Time evolution of lysis front position and permeation

As described in the Methods, we measure the flow and the position of the clot front as the lysis proceeds. The flow is driven by pressure gradient $\frac{\Delta P}{L}$, where ΔP is the pressure difference due to the load of the liquid column and L is the length of the clot. With the progress of the lysis, the clot length shortens, and liquid height increases, which should accelerate the fluid velocity, according to Darcy's law. However, for clots prepared with 2.4 and $3.2\,\text{mg·ml}^{-1}$ of fibrinogen, the flow increases moderately (Figure 3A) or stays steady during a significant part of the lysis (Figure 3C). These results are essential regarding thrombolytic treatment and recanalization. They indicate that flow is barely restored until at least half of the clot is removed. These results contrast with *Shibeko et al.* (*2020*), where simulated recanalization occurs while the average fibrin concentration in the occluded vessel is still high.

²⁷² Together with the flow, we also track the position of the front h during the lysis. Our results show
²⁷³ that $h(t)$ varies non-linearly with time; consequently, the lysis velocity increases as the clot is lysed.
²⁷⁴ Previous numerical models found different lysis profiles, typically linear or sub-linear **Piebalgs and**
²⁷⁵ **Xu** (**2015**); **Shibeko et al.** (**2020**). This difference can be explained by the fact that simulated bound
²⁷⁶ tPA at the front of the clot tends to decrease with time, while our experimental data indicate a
²⁷⁷ constant accumulation at the front (Figure 2). Despite a change in the pressure gradient, hydro-
²⁷⁸ dynamics effects cannot be invoked as a primary factor for lysis acceleration, considering that the
²⁷⁹ flow does not change significantly during the lysis. Instead, the spatial distribution of anti-FA may
²⁸⁰ be sufficient to explain the experimental lysis profiles. We measure the averaged fluid velocity
²⁸¹ for each experiment and use this value to solve iteratively equation (5). Next, we fit this solution
²⁸² with the experimental curves, using the corresponding values of fibrinogen and tPA concentrations
²⁸³ (Figures 3B, D solid lines), and estimate the corresponding value of the parameter k_1 (Figure 3E).
²⁸⁴ For 2.4 and $3.2 \, \mathrm{mg \cdot ml^{-1}}$ of fibrinogen, we measure respectively $k_1 = 18.67 \pm 4.33 \, \mathrm{(s \cdot mg \cdot ml^{-1})^{-1}}$ and
²⁸⁵ $k_1 = 7.89 \pm 1.90 \, \mathrm{(s \cdot mg \cdot ml^{-1})^{-1}}$. The theoretical predictions agree with the experimental results, even
²⁸⁶ though the model is oversimplified and has only one free parameter. A limitation of our model,
²⁸⁷ but also present in other computational approaches, is the intrinsic variability between different
²⁸⁸ experiments. Experiments performed with similar conditions result in variable outcomes for both
²⁸⁹ the flow and lysis profiles, and lysis time can be up to twice longer (Figure 3). Despite identical pro-
²⁹⁰ tocol for clot preparation, the variability remains important and difficult to assess. Likely, the clot
²⁹¹ structure is involved, either local changes of the porosity or fibers radius, that may affect the flow
²⁹² of the degradation of the fibers. For instance, Figure S1 shows the dispersion of the permeabil-
²⁹³ ity among clots prepared with identical fibrin concentration. To date, analytical predictions and
²⁹⁴ experimental results have not been directly compared in similar conditions, making our experi-
²⁹⁵ mental data essential for future theoretical or numerical developments. We provide experimental
²⁹⁶ lysis profiles under various experimental conditions that could be used to test and validate new
²⁹⁷ models of fibrinolysis, see Figure S3.

²⁹⁸ **Mesoscopic model of fibrinolysis**
²⁹⁹ We will now show the results of the numerical model varying the fibrin and anti-FA concentrations,
³⁰⁰ and briefly explore the lysis of spatially heterogeneous thrombi. Unless said otherwise, the param-
³⁰¹ eters used for the simulations presented thereafter are gathered in Table 1.

³⁰² Homogeneous thrombi
³⁰³ Our LB numerical model can be used in different settings, for instance, inside patient-specific
³⁰⁴ geometries (Figure 4A), or experimental tubes (Figure 4B). Our computational framework is com-
³⁰⁵ posed of three main building blocks. The clot is described as a porous medium, the fluid computed
³⁰⁶ with the LB method, and flow characteristics depend on the presence of the clot and its gradual
³⁰⁷ lysis (see Methods). More importantly, our model accounts for the interplay between those blocks:
³⁰⁸ with the progression of the lysis, the local concentration of fibrin decreases, which reduces the
³⁰⁹ solid fraction and the permeability at a voxel level and eventually modifies globally the fluid flow
³¹⁰ as well as transport properties. Our implementation of the partial bounce-back method can repro-
³¹¹ duce predicted fluid velocity profiles (Figure 6A) and simulate fibrous porous at any solid fraction
³¹² (Figure 6B-D).

³¹³ Furthermore, we show that the minimal two-species lysis model leads to a non-linear front
³¹⁴ evolution under the assumption that one species is blocked at the clot interface. However, we
³¹⁵ must test whether such assumptions remain valid once implemented in a more complex numeri-
³¹⁶ cal setup. For different initial concentrations of fibrinogen F_0, anti-FA $\overline{F_0}$ and reaction constant k_1
³¹⁷ the time to lyse the first slice of clot Δ is in agreement with the time predicted by Equation (4). The
³¹⁸ layer is considered lysed for the one-dimensional model when the fibrin concentration decreases
³¹⁹ below a threshold value F^*. In three dimensions, not all the voxels are lysed simultaneously due to
³²⁰ the Poiseuille flow profile and the diffusion of the anti-FA. The layer is lysed when the average fibrin

concentration in a slice goes below F^*, which results in small deviations from the theoretical prediction (Figure 4A). Then we simulate the complete lysis of a two-dimensional clot of width 5.8 mm and length 5 mm, with different initial concentrations of fibrin (Figure 4B) and anti-FA (Figure 4C). As observed experimentally, the lysis front velocity increases in time, and the simulated lysis profiles match the theoretical ones (see Figure S7). Interestingly, we observe a systematic deviation from the predicted profiles in the last part of the lysis. In the last third of the simulation, the lysis is faster than expected by Equation (5). This discrepancy shows the effect of considering a constant fluid velocity. Indeed, the flow rates computed from the simulations increase initially very slowly, restoring only around 15% of the final throughput, followed by an abrupt change (Figure 4B, C dashed lines), which manifests itself by a deviation from the theoretical lysis curve. It was already reported that enhanced permeation results in faster lysis *Diamond and Anand* (*1993*). In the experiments we observe a late increase of the flow but less pronounced than in the simulations. It is worth noticing that experimental lysis profiles were not acquired until the end of the lysis. Clot front positions and flow were monitored at constant frame rate (see Methods). Whenever the measurements stopped before the lysis of the full clot, it implies that the clot was removed between two consecutive images, and likely the flow increased abruptly, as in the simulations, but could not be measured.

Heterogeneous thrombi

Thrombi in ischemic strokes are not homogeneous *Staessens and De Meyer* (*2020*); *Dumitriu La-Grange et al.* (*2023*); *Meglio et al.* (*2019*); *Desilles et al.* (*2022*). We take advantage of the high adaptability of the numerical mesoscopic model to explore the lysis under flow of heterogeneous thrombi. We study two types of numerical thrombi. Type 1 thrombi are created by drawing for each site a random fibrin concentration within an uniform distribution, centered around the target concentration of $2\,\mathrm{mg\cdot ml^{-1}}$. The standard deviation of the law comes in four variants: 25, 50, 75 and 100% of the target concentration. For instance, a thrombus generated at the target fibrin concentration of $2\,\mathrm{mg\cdot ml^{-1}}$ and 100% dispersion, can have the concentration of each site vary between 0 and $4\,\mathrm{mg\cdot ml^{-1}}$. Type 2 thrombi are created by randomly placing 25 medium sized-disks with a high fibrin concentration, and by filling the remaining space with a lower concentration, chosen so that the average concentration in the whole thrombus is equal to the target concentration of $2\,\mathrm{mg\cdot ml^{-1}}$. The disks can overlap each other or can be cut if they exceed the domain's boundary. This means that there are thrombi that have disks occupying more space than others, which implies that they have a lower outer-disk fibrin concentration to keep the same average target concentration. We imposed three intra-disk concentrations : 2.5, 3 and $3.5\,\mathrm{mg\cdot ml^{-1}}$, corresponding to an increase of respectively 25, 50 and 75% of the target concentration.

For type 1 clots, we generated and lysed 10 thrombi for each fibrin dispersion percentage. Since the lysis front is no longer a good metric for heterogeneous thrombi, due to the heterogeneity of each slice of thrombus, we measure lysis progression by computing the thrombus remaining mass percentage and recanalization, see Figure 5A and B. We can see that the more heterogeneous the thrombi, the earlier the recanalization starts ($\sim$ 10% faster in average) and the more variability can be observed in the recanalization profile. One of the simulation at 100% standard deviation recanalizes only one tenth of the final flow of the homogeneous simulation after 500 s. Lyses that have a lower final outflow typically have clot remnants on the walls or in initially high concentration spots. The times to lyse 25, 50 and 75% of the clot, $t_{\frac{1}{4}}$, $t_{\frac{1}{2}}$ and $t_{\frac{3}{4}}$ respectively, are reported in Figure 5C. Thrombi that are more heterogeneous lyse on average faster, and show more variability in lysis times. Finally, the lysis velocities standard deviation tends to increase in time. This can be seen from the boxplot Figure 5C, where we can see the quartiles of the boxes vary in time.

For type 2 thrombi, we lysed three sets of 10 thrombi for each of the three intra-disk fibrin concentrations: 2.5, 3, and $3.5\,\mathrm{mg\cdot ml^{-1}}$. As for the previous type of heterogeneity, we see that the more heterogeneous the thrombus, the more variability there is in the lysis curves, see Figure 5D and F, and the less abrupt their recanalization occurs, see Figure 5E. These variations are more

371 pronounced than for type 1 clots, with fluctuation of $\pm 35\%$ on half-lysis time for the most hetero-
372 geneous case. Also, the lower the outer-disk concentration, the higher the average flow inside the
373 thrombus before the lysis, but the harder it will be to accumulate anti-FA at the thrombus disks to
374 lyse them. This is because the higher permeability outer-disks regions act as preferred lesser resis-
375 tance channels for the flow. This can produce lysis curves that reach a plateau, with a significant
376 fraction of clot mass remaining. More variability on the lysis times and fluxes can be observed with
377 type 2 clots than type 1 clots, see Figure 5F.

Discussion

379 We presented original *in vitro* fibrinolysis experiments of homogeneous fibrin-pure thrombi per-
380 fused continuously with tPA, and an analytical formula that could predict the lysis front evolution,
381 provided the effective reaction rate k_1 is well calibrated. These revealed three key points: (1) tPA
382 does not penetrate in the bulk of the fibrin-pure thrombi, (2) the lysis accelerates, and (3) flow is
383 restored suddenly only at the end of the lysis. We also presented an original numerical framework
384 that uses the LB and the PBB methods to predict more specifically the evolution of clot length, per-
385 meability, and hydrodynamic properties during the fibrinolysis. The lysis front evolution matched
386 the analytical model while the flow throughout lysis stayed of the same order of magnitude as the
387 initial flow. Our results reveal that the fibrin concentration and permeability of the thrombus, dic-
388 tating the transport of pro-fibrinolytic macromolecules to the fibers, is of utmost importance for
389 fibrinolysis success.

390 The spatial thrombus heterogeneity alone showed a significant influence on lysis time and re-
391 canalization, with up to $\sim 60\%$ variation in half-lysis time for different instances of type 2 clots,
392 containing high concentration disks of variable size. We could imagine that other internal arrange-
393 ments of thrombi would make an even greater difference, and it would be interesting to study
394 which topology lyses the fastest. To study more realistic thrombi, it is possible with the present
395 model to generate and lyse clots reconstructed from real clot images. Indeed, it is well known that
396 thrombi analyzed after a thrombectomy show a very large heterogeneity in the spatial distribution
397 of the blood's figurative elements but also in the structure of the fibrin network *Staessens et al.*
398 (*2020*). All these elements are not easy to take into account in the design of an experiment. In
399 addition to this spatial heterogeneity, there is the influence of cell surfaces in the control of fibri-
400 nolysis. Whether it is the receptors specific to the factors involved in fibrinolysis *Cesarman-Maus*
401 *and Hajjar* (*2005*) or annexin A2 *Lim and Hajjar* (*2021*). What further complicates the modeling is
402 that one cell type can have completely opposite effects on fibrinolysis. We have recently shown
403 that platelets can have a profibrinolytic or antifibrinolytic action depending on the clinical situation
404 of the patients *Petkantchin et al.* (*2022*). Numerical experimentation will be a complementary tool
405 to address these issues. A showcase example can be found in the Appendix Figure S8, where we
406 lysed 2D slices of patient retrieved thrombi (sliced and analyzed by Staessens *et al.* *Staessens et al.*
407 (*2020*)). 3D reconstructed thrombi could also be lysed, provided 3D data is available. We chose
408 in the present work to treat heterogeneity as a change in fibrin concentration, but other options
409 could be to change locally the reaction rate k_1, or the permeability (fiber radius or permeability
410 law), which is feasible with our numerical model.

411 The numerical model is purposely streamlined, keeping only the essential features needed to
412 replicate the experimental observations. In that regard, it suffers in appearance from not seeming
413 clinically relevant. For instance, we assume steady and laminar flow, whereas it is not the case *in*
414 *vivo*. Recent work from INSIST partners *Arrarte Terreros et al.* (*2021*) identified up to 4 different
415 patterns of flow in clotted regions of ischemic stroke patients: *slow anterograde*, *fast anterograde*,
416 *retrograde* and mixed *anterograde-retrograde* flows. Anterograde flow designates the expected di-
417 rection of the flow, *i.e.* from proximal to distal, while retrograde means the opposite direction.
418 These different flow patterns undoubtedly affect the transport of lytic drugs to the occlusion site
419 and, eventually, the evolution of the lysis. Also, the artery is here modeled by a rigid tube, and clot
420 fragments stay in place until totally lysed, even when disconnected from the wall or the main clot

chunk. Fortunately, thanks to its design, the model could be easily extended to include conditions closer to *in vivo* thrombolysis. We could instead reconstruct *Decroocq et al.* (*2022*) patient-specific geometries with collateral vessels and try to reproduce the complex flow conditions observed in AIS patients *Arrarte Terreros et al.* (*2021*). In doing so, we could hypothesize the effect of artery geometry, thrombus location, and drug injection patterns on lysis efficacy. Seners et al. have reported that distal thrombi lyse faster than proximal thrombi *Seners et al.* (*2016*). One could use the numerical model to assess whether this discrepancy comes from the shape and composition of distal clots, local flow conditions, or both. Fragmentation of the clot could be implemented by adding fluid to clot interaction, and the PBB method is a good technique to simulate moving objects *Rettinger and Rüde* (*2017*); *Tsigginos et al.* (*2022*). It would be interesting to measure the impact of each of these extensions and their necessity in future works.

Acknowledgements

This research is part of the INSIST project *INSIST* (*2017*), which has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No. 777072. This work was also supported by grants from the CHU Charleroi and the Fonds de la Chirurgie Cardiaque.

Conflict of interests

None declared.

INSIST Investigators

Charles Majoie[1], Ed van Bavel[2], Henk Marquering[1,2], Nerea Arrarte-Terreros[1,2], Praneeta Konduri[1,2], Sissy Georgakopoulou[2], Yvo Roos[3], Alfons Hoekstra[4], Raymond Padmos[4], Victor Azizi[4], Claire Miller[4], Max van der Kolk[4], Aad van der Lugt[5], Diederik W.J. Dippel[6], Hester L. Lingsma[7], Nikki Boodt[5,6,7], Noor Samuels[5,6,7], Stephen Payne[8], Tamas Jozsa[8], Wahbi K. El-Bouri[8], Michael Gilvarry[9], Ray McCarthy[9], Sharon Duffy[9], Anushree Dwivedi[9], Behrooz Fereidoonnezhad[10], Kevin Moerman[10], Patrick Mc Garry[10], Senna Staessens[11], Simon de Meyer[11], Sarah Vandelanotte[11], Francesco Migliavacca[12], Gabriele Dubini[12], Giulia Luraghi[12], Jose Felix Rodriguez Matas[12], Sara Bridio[12], Bastien Chopard[13], Franck Raynaud[13], Rémy Petkantchin[13], Vanessa Blanc-Guillemaud[14], Mikhail Panteleev[15,16], Alexey Shibeko[15], Karim Zouaoui Boudjeltia[17].

[1]Department of Radiology and Nuclear Medicine, Amsterdam UMC, location AMC, Amsterdam, the Netherlands; [2]Biomedical Engineering and Physics, Amsterdam UMC, location AMC, Amsterdam, the Netherlands; [3]Department of Neurology, Amsterdam UMC, location AMC, Amsterdam, the Netherlands; [4]Computational Science Lab, Faculty of Science, Institute for Informatics, University of Amsterdam, Amsterdam, the Netherlands; [5]Department of Radiology and Nuclear Medicine, Erasmus MC University Medical Center, PO Box 2040, 3000 CA Rotterdam, the Netherlands; [6]Department of Neurology, Erasmus MC University Medical Center, PO Box 2040, 3000 CA Rotterdam, the Netherlands; [7]Department of Public Health, Erasmus MC University Medical Center, PO Box 2040, 3000 CA Rotterdam, the Netherlands; [8]Institute of Biomedical Engineering, Department of Engineering Science, University of Oxford, Parks Road, Oxford OX1 3PJ, UK; [9]Cerenovus, Galway Neuro Technology Center, Galway, Ireland; [10]College of Engineering and Informatics, National University of Ireland Galway, Ireland; National Center for Biomedical Engineering Science, National University of Ireland Galway, Ireland; [11]Laboratory for Thrombosis Research, KU Leuven Campus Kulak Kortrijk, Kortrijk, Belgium; [12]Laboratory of Biological Structure Mechanics, Department of Chemistry, Materials and Chemical Engineering "Giulio Natta", Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133 Milano, Italy; [13]Computer Science Department, University of Geneva, CUI, 7 route de Drize, 1227 Carouge, Switzerland; [14]Institut de Recherches Internationales Servier, Coubevoie Cedex, France; [15]Center for Theoretical Problems of Physicochemical Pharmacology RAS, Moscow, Russia; [16]Dmitry Rogachev National Research Center of Pediatric Hematology, Oncology and Immunology, Moscow,

468 Russia; Faculty of Physics, Lomonosov Moscow State University, Moscow, Russia; [17]Laboratory of
469 Experimental Medicine (ULB 222 Unit), Université Libre de Bruxelles (ULB), CHU de Charleroi, Bel-
470 gium.

References

Anand M, Rajagopal K, Rajagopal KR. A Model for the Formation and Lysis of Blood Clots. Pathophysiology of
Haemostasis and Thrombosis. 2005; 34(2-3):109–120. https://www.karger.com/Article/FullText/89931, doi:
10.1159/000089931.

Arrarte Terreros N, Tolhuisen ML, Bennink E, de Jong HWAM, Beenen LFM, Majoie CBLM, van Bavel E, Mar-
quering HA. From perviousness to permeability, modelling and measuring intra-thrombus flow in acute
ischemic stroke. Journal of Biomechanics. 2020 Oct; 111:110001. https://linkinghub.elsevier.com/retrieve/pii/
S0021929020304243, doi: 10.1016/j.jbiomech.2020.110001.

Arrarte Terreros N, van Willigen BG, Niekolaas WS, Tolhuisen ML, Brouwer J, Coutinho JM, Beenen LF, Majoie
CB, van Bavel E, Marquering HA. Occult blood flow patterns distal to an occluded artery in acute ischemic
stroke. Journal of Cerebral Blood Flow & Metabolism. 2021 Sep; p. 0271678X211044941. https://doi.org/10.
1177/0271678X211044941, doi: 10.1177/0271678X211044941.

Bajd F, Serša I. Mathematical Modeling of Blood Clot Fragmentation During Flow-Mediated Thrombolysis.
Biophysical Journal. 2013 Mar; 104(5):1181–1190. https://www.cell.com/biophysj/abstract/S0006-3495(13)
00134-3, doi: 10.1016/j.bpj.2013.01.029.

Bannish BE, Keener JP, Fogelson AL. Modelling fibrinolysis: a 3D stochastic multiscale model. Mathematical
Medicine and Biology. 2014 Mar; 31(1):17–44. https://academic.oup.com/imammb/article-lookup/doi/10.1093/
imammb/dqs029, doi: 10.1093/imammb/dqs029.

Bannish BE, Hudson NE. The utility and potential of mathematical models in predicting fibrinolytic outcomes.
Current Opinion in Biomedical Engineering. 2021 Dec; 20:100337. https://www.sciencedirect.com/science/
article/pii/S2468451121000775, doi: 10.1016/j.cobme.2021.100337.

Bhatnagar PL, Gross EP, Krook M. A Model for Collision Processes in Gases. I. Small Amplitude Processes in
Charged and Neutral One-Component Systems. Physical Review. 1954 May; 94(3):511–525. https://link.aps.
org/doi/10.1103/PhysRev.94.511, doi: 10.1103/PhysRev.94.511, publisher: American Physical Society.

Blinc A, Magdic J, Fric J. Atomic force microscopy of fibrin networks and plasma clots during fibrinolysis. . 2000;
p. 12.

Bucay I, O'Brien ET, Wulfe SD, Superfine R, Wolberg AS, Falvo MR, Hudson NE. Physical determinants of fibri-
nolysis in single fibrin fibers. PloS One. 2015; 10(2):e0116350. doi: 10.1371/journal.pone.0116350.

Cesarman-Maus G, Hajjar KA. Molecular mechanisms of fibrinolysis. British Journal of Haematology. 2005
May; 129(3):307–321. doi: 10.1111/j.1365-2141.2005.05444.x.

Chopard B, Dupuis A, Masselot A, Luthi P. Cellular automata and lattice Boltzmann techniques: an approach to
model and simulate complex systems. Advances in Complex Systems. 2002 Jun; 05(02n03):103–246. https:
//www.worldscientific.com/doi/abs/10.1142/S0219525902000602, doi: 10.1142/S0219525902000602.

Clague DS, Kandhai BD, Zhang R, Sloot PMA. Hydraulic permeability of (un)bounded fibrous media using the
lattice Boltzmann method. Physical Review E. 2000 Jan; 61(1):616–625. https://link.aps.org/doi/10.1103/
PhysRevE.61.616, doi: 10.1103/PhysRevE.61.616.

Collet J P, Park D, Lesty C, Soria J, Soria C, Montalescot G, Weisel J W. Influence of Fibrin Network
Conformation and Fibrin Fiber Diameter on Fibrinolysis Speed. Arteriosclerosis, Thrombosis, and Vas-
cular Biology. 2000 May; 20(5):1354–1361. https://www.ahajournals.org/doi/10.1161/01.atv.20.5.1354, doi:
10.1161/01.ATV.20.5.1354.

Darcy H. Les fontaines publiques de la ville de Dijon; 1856. https://gallica.bnf.fr/ark:/12148/bpt6k624312.

Davies CN. The Separation of Airborne Dust and Particles. Inst Mech Eng. 1952; (B1):185–213.

Decroocq M, Lavoue G, Ohta M, Frindel C. A Software to Visualize, Edit, Model and Mesh Vascular Net-
works. Annual International Conference of the IEEE Engineering in Medicine and Biology Society IEEE En-
gineering in Medicine and Biology Society Annual International Conference. 2022 Jul; 2022:2208–2214. doi:
10.1109/EMBC48229.2022.9871365.

Desilles JP, Di Meglio L, Delvoye F, Maïer B, Piotin M, Ho-Tin-Noé B, Mazighi M. Composition and Organization of Acute Ischemic Stroke Thrombus: A Wealth of Information for Future Thrombolytic Strategies. Frontiers in Neurology. 2022; 13. https://www.frontiersin.org/articles/10.3389/fneur.2022.870331.

Diamond SL, Anand S. Inner clot diffusion and permeation during fibrinolysis. Biophysical Journal. 1993 Dec; 65(6):2622–2643. https://linkinghub.elsevier.com/retrieve/pii/S0006349593813146, doi: 10.1016/S0006-3495(93)81314-6.

Donkor ES. Stroke in the 21st Century: A Snapshot of the Burden, Epidemiology, and Quality of Life. Stroke Research and Treatment. 2018 Nov; 2018:3238165. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6288566/, doi: 10.1155/2018/3238165.

Dumitriu LaGrange D, Reymond P, Brina O, Zboray R, Neels A, Wanke I, Lövblad KO. Spatial heterogeneity of occlusive thrombus in acute ischemic stroke: A systematic review. Journal of Neuroradiology. 2023 Jan; https://www.sciencedirect.com/science/article/pii/S0150986123000056, doi: 10.1016/j.neurad.2023.01.004.

Feller T, Hársfalvi J, Csányi C, Kiss B, Kellermayer M. Plasmin-driven fibrinolysis in a quasi-two-dimensional nanoscale fibrin matrix. Journal of Structural Biology. 2018 Sep; 203(3):273–280. doi: 10.1016/j.jsb.2018.05.010.

Fransen PSS, Beumer D, Berkhemer OA, van den Berg LA, Lingsma H, van der Lugt A, van Zwam WH, van Oostenbrugge RJ, Roos YBWEM, Majoie CB, Dippel DWJ, MR CLEAN Investigators. MR CLEAN, a multicenter randomized clinical trial of endovascular treatment for acute ischemic stroke in the Netherlands: study protocol for a randomized controlled trial. Trials. 2014 Sep; 15:343. doi: 10.1186/1745-6215-15-343.

Frisch U, Hasslacher B, Pomeau Y. Lattice-Gas Automata for the Navier-Stokes Equation. Physical Review Letters. 1986 Apr; 56(14):1505–1508. https://link.aps.org/doi/10.1103/PhysRevLett.56.1505, doi: 10.1103/PhysRevLett.56.1505, publisher: American Physical Society.

Gabriel DA, Muga K, Boothroyd EM. The effect of fibrin structure on fibrinolysis. Journal of Biological Chemistry. 1992 Dec; 267(34):24259–24263. https://www.sciencedirect.com/science/article/pii/S0021925818357594, doi: 10.1016/S0021-9258(18)35759-4.

Gu B, Piebalgs A, Huang Y, Roi D, Lobotesis K, Longstaff C, Hughes AD, Chen R, Thom SA, Xu XY. Computational simulations of thrombolysis in acute stroke: Effect of clot size and location on recanalisation. Medical Engineering & Physics. 2019 Nov; 73:9–17. https://linkinghub.elsevier.com/retrieve/pii/S1350453319301493, doi: 10.1016/j.medengphy.2019.07.014.

He X, Luo LS. Lattice Boltzmann Model for the Incompressible Navier–Stokes Equation. Journal of Statistical Physics. 1997 Aug; 88(3/4):927–944. http://link.springer.com/10.1023/B:JOSS.0000015179.12689.e4, doi: 10.1023/B:JOSS.0000015179.12689.e4.

He X, Luo LS. Theory of the lattice Boltzmann method: From the Boltzmann equation to the lattice Boltzmann equation. Physical Review E. 1997 Dec; 56(6):6811–6817. https://link.aps.org/doi/10.1103/PhysRevE.56.6811, doi: 10.1103/PhysRevE.56.6811.

INSIST, In SIlico Clinical Trials for treatment of Acute ischemic stroke (INSIST) H2020 project. https://insist-h2020.eu/; 2017. https://insist-h2020.eu/.

Jackson GW, James DF. The permeability of fibrous porous media. The Canadian Journal of Chemical Engineering. 1986 Jun; 64(3):364–374. http://doi.wiley.com/10.1002/cjce.5450640302, doi: 10.1002/cjce.5450640302.

Jolugbo P, Ariëns RAS. Thrombus Composition and Efficacy of Thrombolysis and Thrombectomy in Acute Ischemic Stroke. Stroke. 2021 Mar; 52(3):1131–1142. doi: 10.1161/STROKEAHA.120.032810.

Kotsalos C, Latt J, Chopard B. Bridging the computational gap between mesoscopic and continuum modeling of red blood cells for fully resolved blood flow. Journal of Computational Physics. 2019 Dec; 398:108905. http://adsabs.harvard.edu/abs/2019JCoPh.39808905K, doi: 10.1016/j.jcp.2019.108905.

Krüger T, Kusumaatmaja H, Kuzmin A, Shardt O, Silva G, Viggen EM. The Lattice Boltzmann Method: Principles and Practice. Graduate Texts in Physics, Cham: Springer International Publishing; 2017. http://link.springer.com/10.1007/978-3-319-44649-3, doi: 10.1007/978-3-319-44649-3.

Kutay ME, Aydilek AH, Masad E. Laboratory validation of lattice Boltzmann method for modeling pore-scale flow in granular materials. Computers and Geotechnics. 2006 Dec; 33(8):381–395. https://linkinghub.elsevier.com/retrieve/pii/S0266352X06000826, doi: 10.1016/j.compgeo.2006.08.002.

Latt J, Malaspinas O, Kontaxakis D, Parmigiani A, Lagrava D, Brogi F, Belgacem MB, Thorimbert Y, Leclaire S, Li S, Marson F, Lemus J, Kotsalos C, Conradin R, Coreixas C, Petkantchin R, Raynaud F, Beny J, Chopard B. Palabos: Parallel Lattice Boltzmann Solver. Computers & Mathematics with Applications. 2020 Apr; https://linkinghub.elsevier.com/retrieve/pii/S0898122120301267, doi: 10.1016/j.camwa.2020.03.022.

Li R, Yang YS, Pan J, Pereira GG, Taylor JA, Clennell B, Zou C. Lattice Boltzmann modeling of permeability in porous materials with partially percolating voxels. Physical Review E. 2014 Sep; 90(3). https://link.aps.org/doi/10.1103/PhysRevE.90.033301, doi: 10.1103/PhysRevE.90.033301.

Li S, Chopard B, Latt J. Continuum model for flow diverting stents in 3D patient-specific simulation of intracranial aneurysms. Journal of Computational Science. 2019 Nov; 38:101045. http://www.sciencedirect.com/science/article/pii/S1877750319310269, doi: 10.1016/j.jocs.2019.101045.

Lim HI, Hajjar KA. Annexin A2 in Fibrinolysis, Inflammation and Fibrosis. International Journal of Molecular Sciences. 2021 Jan; 22(13):6836. https://www.mdpi.com/1422-0067/22/13/6836, doi: 10.3390/ijms22136836, number: 13 Publisher: Multidisciplinary Digital Publishing Institute.

Longstaff C, Kolev K. Basic mechanisms and regulation of fibrinolysis. Journal of Thrombosis and Haemostasis. 2015; 13(S1):S98–S105. doi: 10.1111/jth.12935.

Lynch SR, Laverty SM, Bannish BE, Hudson NE. Microscale structural changes of individual fibrin fibers during fibrinolysis. Acta Biomaterialia. 2022 Mar; 141:114–122. https://www.sciencedirect.com/science/article/pii/S174270612200006X, doi: 10.1016/j.actbio.2022.01.006.

Malaspinas O, Turjman A, Ribeiro de Sousa D, Garcia-Cardena G, Raes M, Nguyen PTT, Zhang Y, Courbebaisse G, Lelubre C, Zouaoui Boudjeltia K, Chopard B. A spatio-temporal model for spontaneous thrombus formation in cerebral aneurysms. bioRxiv. 2015 Jul; http://biorxiv.org/lookup/doi/10.1101/023226, doi: 10.1101/023226.

Meglio LD, Desilles JP, Ollivier V, Nomenjanahary MS, Meglio SD, Deschildre C, Loyau S, Olivot JM, Blanc R, Piotin M, Bouton MC, Michel JB, Jandrot-Perrus M, Ho-Tin-Noé B, Mazighi M. Acute ischemic stroke thrombi have an outer shell that impairs fibrinolysis. Neurology. 2019 Oct; 93(18):e1686–e1698. https://n.neurology.org/content/93/18/e1686, doi: 10.1212/WNL.0000000000008395, publisher: Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology Section: Article.

Miyazaki H, Yamaguchi T. Formation and destruction of primary thrombi under the influence of blood flow and von Willebrand factor analyzed by a discrete element method. Biorheology. 2003 Jan; 40(1,2,3):265–272. https://content.iospress.com/articles/biorheology/bir188.

Nilsen ML. A historical account of stroke and the evolution of nursing care for stroke patients. The Journal of Neuroscience Nursing: Journal of the American Association of Neuroscience Nurses. 2010 Feb; 42(1):19–27. doi: 10.1097/jnn.0b013e3181c1fdad.

Petkantchin R, Padmos R, Boudjeltia KZ, Raynaud F, Chopard B. Thrombolysis: Observations and numerical models. Journal of Biomechanics. 2022 Feb; 132:110902. https://linkinghub.elsevier.com/retrieve/pii/S0021929021006552, doi: 10.1016/j.jbiomech.2021.110902.

Piebalgs A, Gu B, Roi D, Lobotesis K, Thom S, Xu XY. Computational Simulations of Thrombolytic Therapy in Acute Ischaemic Stroke. Scientific Reports. 2018 Dec; 8(1). http://www.nature.com/articles/s41598-018-34082-7, doi: 10.1038/s41598-018-34082-7.

Piebalgs A, Xu XY. Towards a multi-physics modelling framework for thrombolysis under the influence of blood flow. Journal of The Royal Society Interface. 2015 Dec; 12(113):20150949. https://royalsocietypublishing.org/doi/10.1098/rsif.2015.0949, doi: 10.1098/rsif.2015.0949.

Raynaud F, Rousseau A, Monteyne D, Perez-Morga D, Zouaoui Boudjeltia K, Chopard B. Investigating the two regimes of fibrin clot lysis: an experimental and computational approach. Biophysical Journal. 2021 Sep; 120(18):4091–4106. doi: 10.1016/j.bpj.2021.08.005.

Rettinger C, Rüde U. A comparative study of fluid-particle coupling methods for fully resolved lattice Boltzmann simulations. Computers & Fluids. 2017 Sep; 154:74–89. http://arxiv.org/abs/1702.04910, doi: 10.1016/j.compfluid.2017.05.033, arXiv:1702.04910 [physics].

Riedel CH, Zimmermann P, Jensen-Kondering U, Stingele R, Deuschl G, Jansen O. The Importance of Size: Successful Recanalization by Intravenous Thrombolysis in Acute Anterior Stroke Depends on Thrombus Length. Stroke. 2011 Jun; 42(6):1775–1777. https://www.ahajournals.org/doi/10.1161/STROKEAHA.110.609693, doi: 10.1161/STROKEAHA.110.609693.

Seners P, Turc G, Maïer B, Mas JL, Oppenheim C, Baron JC. Incidence and Predictors of Early Recanalization After Intravenous Thrombolysis: A Systematic Review and Meta-Analysis. Stroke. 2016 Sep; 47(9):2409–2412. doi: 10.1161/STROKEAHA.116.014181.

Shan X, Yuan XF, Chen H. Kinetic theory representation of hydrodynamics: a way beyond the Navier–Stokes equation. Journal of Fluid Mechanics. 2006 Mar; 550:413–441. https://www.cambridge.org/core/journals/journal-of-fluid-mechanics/article/abs/kinetic-theory-representation-of-hydrodynamics-a-way-beyond-the-navierstokes-equation/57B640BEE52716398F6C015BC66D189C, doi: 10.1017/S0022112005008153, publisher: Cambridge University Press.

Shibeko AM, Chopard B, Hoekstra AG, Panteleev MA. Redistribution of TPA Fluxes in the Presence of PAI-1 Regulates Spatial Thrombolysis. Biophysical Journal. 2020 Aug; 119(3):638–651. https://linkinghub.elsevier.com/retrieve/pii/S0006349520304963, doi: 10.1016/j.bpj.2020.06.020.

Shin JW, Jeong HS, Kwon HJ, Song KS, Kim J. High red blood cell composition in clots is associated with successful recanalization during intra-arterial thrombectomy. PLoS ONE. 2018 May; 13(5):e0197492. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5962078/, doi: 10.1371/journal.pone.0197492.

Staessens S, De Meyer SF. Thrombus heterogeneity in ischemic stroke. Platelets. 2020 Apr; p. 1–9. https://www.tandfonline.com/doi/full/10.1080/09537104.2020.1748586, doi: 10.1080/09537104.2020.1748586.

Staessens S, Denorme F, Francois O, Desender L, Dewaele T, Vanacker P, Deckmyn H, Vanhoorelbeke K, Andersson T, De Meyer SF. Structural analysis of ischemic stroke thrombi: histological indications for therapy resistance. Haematologica. 2020 Feb; 105(2):498–507. http://www.haematologica.org/lookup/doi/10.3324/haematol.2019.219881, doi: 10.3324/haematol.2019.219881.

Succi S. The Lattice Boltzmann Equation: For Complex States of Flowing Matter. Oxford University Press; 2018. https://doi.org/10.1093/oso/9780199592357.001.0001, doi: 10.1093/oso/9780199592357.001.0001.

Tsigginos C, Meng J, Gu XJ, Emerson DR. Coupled LBM-DEM simulations using the partially saturated method: Theoretical and computational aspects. Powder Technology. 2022 Jun; 405:117556. https://www.sciencedirect.com/science/article/pii/S0032591022004508, doi: 10.1016/j.powtec.2022.117556.

Walsh SDC, Burwinkle H, Saar MO. A new partial-bounceback lattice-Boltzmann method for fluid flow through heterogeneous media. Computers & Geosciences. 2009 Jun; 35(6):1186–1193. https://linkinghub.elsevier.com/retrieve/pii/S009830040800201X, doi: 10.1016/j.cageo.2008.05.004.

Weisel JW. Fibrinogen and fibrin. Advances in Protein Chemistry. 2005; 70:247–299. doi: 10.1016/S0065-3233(05)70008-5.

Wufsus AR, Macera NE, Neeves KB. The Hydraulic Permeability of Blood Clots as a Function of Fibrin and Platelet Density. Biophysical Journal. 2013 Apr; 104(8):1812–1823. https://linkinghub.elsevier.com/retrieve/pii/S0006349513003299, doi: 10.1016/j.bpj.2013.02.055.

Yesudasan S, Wang X, Averett RD. Fibrin polymerization simulation using a reactive dissipative particle dynamics method. Biomechanics and Modeling in Mechanobiology. 2018 Oct; 17(5):1389–1403. https://doi.org/10.1007/s10237-018-1033-8, doi: 10.1007/s10237-018-1033-8.

Zidansek A, Blinc A, Lahajnar G, Keber D, Blinc R. Finger-like lysing patterns of blood clots. Biophysical Journal. 1995 Sep; 69(3):803–809. https://linkinghub.elsevier.com/retrieve/pii/S0006349595799544, doi: 10.1016/S0006-3495(95)79954-4.

Quantity	Definition	Value
v [m^2/s]	Solution kinematic viscosity	$1 \cdot 10^{-6}$
∇P [Pa/m]	Pressure gradient across clot	$7.5 \cdot 10^3$
D [mm]	Tube diameter	5.8
F [mg/ml]	Clot fibrin concentration	$2-4$
F^* [mg/ml]	Fibrin concentration threshold	0.2
$\overline{F}$ [mg/ml]	Anti-FA concentration	$3.3-20 \cdot 10^{-6}$
k [m^2]	Clot permeability	10^{-13}
k_1 [(s·mg·ml^{-1})$^{-1}$]	Estimated fibrin reaction rate	12.5
L [mm]	Clot length	5, specified else
R_f [nm]	Fibrin fiber radius	140
u_{f} [μm·s^{-1}]	Mean fluid velocity (Figure 4)	$4-10$
Δx [m]	Space discretization	$1 \cdot 10^{-4}$
Δt [s]	Time discretization	$2.5 \cdot 10^{-5}$

Table 1. Typical initial conditions for the numerical and analytical models. It is specified when other values are used in the manuscript.

Figure 1. Elements of the fibrinolysis model. **A**: Fluid flow is computed in a simulated tube or vessel using the LB method. The flow is driven by a local pressure drop imposed at the left and right boundaries of the system. A clot is placed, simulated as a porous medium, using Walsh's PBB method *Walsh et al.* (*2009*). **B**: The blocking fraction of each clot voxel can be specified depending on the fiber volume fraction. Meso-particles of pro-fibrinolytic agents, aggregated into anti-FA, are continuously injected and will shrink the fibers' radius upon interaction with clot voxels *Diamond and Anand* (*1993*). **C**: The lysis reaction is of second-order and relates the mean fiber radius, as well as the solid fraction in a voxel, to the fibrin concentration inside it.

Figure 2. tPA does not go through the clot but remains blocked near the interface with the supernatant. **A**: Graphical sketch of the experiment. First, the supernatant containing tPA flows through the clot, then the clot is removed and cut into four pieces corresponding to the top part (HH), the top intermediate part (H), the bottom intermediate part (B), and the most bottom part (BB). Each part is lysed for one night after the addition of plasminogen, and finally, tPA concentrations are measured after centrifuging the fibrin degradation products. Concentration of tPA in different parts of the clot (HH,H,B and BB) after **B**: 30 minutes, **C**: 60 minutes and **D**: 120 minutes of perfusion. The boxplots represent the median as well as 1^{st} and 3^{rd} quartiles. The outliers are labeled as statistically non-significant, being outside of 2.689 standard deviations. The number of samples n varies between figures B,C and D because the corresponding experiments were independent.

Figure 3. A,C: Flow measured in the fibrinolysis experiments of 2.4 (A) and (C) $3.2 \, \mathrm{mg \cdot ml^{-1}}$ fibrin. Experiments were repeated and numbered from 1 to 4. In most of the experiments, the flow was constant during most of the lysis and increased towards the end of lysis. **B, D**: Lysis fronts positions measured for the same lysis experiments, with 2.4 (B) and (D) $3.2 \, \mathrm{mg \cdot ml^{-1}}$ fibrin. The analytical model was fitted to the results by choosing the best value of k_1. These values are reported in fig E. **E**: Best values for the analytical model's reaction parameter k_1 fitted on lysis experiments B,D. The mean value, excluding one non-statistically significant value, is $k_1^{\mathrm{exp}} = 12.5$ $(\mathrm{s \cdot mg \cdot ml^{-1}})^{-1}$. The boxplots show the 1^{st} and 3^{rd} quartiles, and the median.

Figure 4. A: Snapshots of a fibrinolysis simulation with the 3D model in a patient-specific artery. The fibrin thrombus (red) initially blocks part of the flow (blue), which makes it hard for anti-fibrin particles (green) to reach it. When the anti-fibrin finally reaches the thrombus and does the lysis reaction, the artery can be recanalized. **B**: Snapshots of fibrinolysis with the 3D model in a cylinder in which a pressure gradient is applied. The axis shows time evolution at regular time intervals. The fibrin thrombus (red) is lysed by the anti-fibrin particles (green), following the lysis equation (17). At each time step, the view angle changes by approximately 50°. **C**: Comparison between simulated time (3D) and expected time (analytical) to lyse the first slice of the clot, obtained for different reaction parameters k_1 (blue), initial concentrations of anti-FA $\overline{F}_0$ (green) and fibrin F_0 (red). Here $\Delta = \Delta x = 114$ µm. **D, E**: Lysis front positions in time for different concentrations of fibrin and anti-FA, respectively, with the numerical 2D (dots) and the analytical (line) models. Baseline parameters are $F_0 = 3\,\mathrm{mg\cdot ml^{-1}}$, $\overline{F}_0 = 10\,\mathrm{ng\cdot ml^{-1}}$, $k_1 = 280\,(\mathrm{s\cdot mg\cdot ml^{-1}})^{-1}$, $\nabla P_0 = 8250\,\mathrm{Pa\cdot m^{-1}}$, $\Delta x = 1.14\cdot 10^{-4}$ m, $\Delta t = 1.83\cdot 10^{-4}$ s.

Figure 5. 10 type 1 (A-C) uniformly heterogeneous 2D thrombi for each fibrin standard deviation, and 10 type 2 (D-F) heterogeneous 2D thrombi, for each intra-disk concentrations of $2.5, 3$ and $3.5 \, \mathrm{mg \cdot ml^{-1}}$, of length 5 mm and width 5.8 mm were lysed, with $\nabla P = 8000 \, \mathrm{Pa \cdot m^{-1}}$ and $k_1 = 1400 \, (\mathrm{s \cdot mg \cdot ml^{-1}})^{-1}$. The mean fibrin concentration for each clot is $F_M = 2 \, \mathrm{mg \cdot ml^{-1}}$, and the initial anti-FA $\overline{F}_0 = 20 \, \mathrm{ng \cdot ml^{-1}}$. **A,D**: The remaining clot mass is plotted in time. **B,E**: The average throughput inside the clot. **C,F**: The times to reach 25, 50 and 75% of the lysis are plotted. The more heterogeneous the clots are, the higher is the variablity on the lysis times. One simulation with $F_{Disk} = 3.5 \, \mathrm{mg \cdot ml^{-1}}$ did not reach the 75% lysis threshold during the simulation time, and was excluded from this boxplot.

 Appendix 1

Permeability measurement of fibrin clots

We measured the outflow for two concentrations of fibrinogen, $2.4\,\mathrm{mg \cdot ml^{-1}}$ (Figure S1A) and $3.2\,\mathrm{mg \cdot ml^{-1}}$

(Figure S1B). Clots were formed in $800\,\mu\mathrm{l}$ total volume with $38.4 - 51.2\,\mu\mathrm{l}$ of fibrinogen at $50\,\mathrm{mg \cdot ml^{-1}}$,

Owren. $160\,\mu\mathrm{l}$ of thrombin was added to trigger clot formation during 15 minutes at $37°C$. Clots

have typical length of 2.6 cm, diameter 5.8 mm and final concentrations of fibrinogen of $2.4\,\mathrm{mg \cdot ml^{-1}}$

and $3.2\,\mathrm{mg \cdot ml^{-1}}$. Outflow was measured every ten minutes. As expected, outflows were constant

over time, allowing us to measure clot permeability reliably. However, our results show high vari-

ability within the same experimental conditions (Figure S1C), highlighting how internal clot prop-

erties, in particular fiber radius, impact the permeability and, ultimately, the transport of proteins

through the clot. Using Darcy's law, we estimated the permeability from the measurement of the

outflow:

$$k = \frac{Q\mu}{\mathcal{A}} \frac{l_0}{\rho g \left(L - l_0 \right)} \tag{22}$$

670 where $\mathcal{A}$ is the section of the tube, k the permeability, μ the dynamic viscosity, L the length of the

671 tube and l_0 the length of the clot.

Appendix 1—figure 1. Permeability decreases with fibrinogen concentration. Outflow as function of time for fibrinogen concentration of **A**: $2.4\,\mathrm{mg \cdot ml^{-1}}$ and **B**: $3.2\,\mathrm{mg \cdot ml^{-1}}$. Each curve corresponds to a separate experiment, with the corresponding permeability indicated in the legend. **C**: Permeability as function of the concentration of fibrinogen. The boxplots represent the median as well as the first and third quartiles.

672 **Lysis of the n-th slice with the analytical model**

673 Before initiating the lysis of the second slice x_1^Δ, the quantity $\overline{F}_0 \frac{u_f}{\Delta} t$ has to be advected over a dis-

674 tance Δ. Hence, the anti-FA that accumulates in x_1^Δ is given by:

$$\overline{F}(x_1^\Delta, t) = \overline{F}_0 \frac{u_f}{\Delta} \left(t - \frac{\Delta}{u_f} \right) \cdot \Theta \left(t - \left(t_0 + \frac{\Delta}{u_f} \right) \right). \tag{23}$$

675 The Heaviside step function Θ reflects that anti-FA does not reach x_1^Δ until x_0^Δ has been lysed, and

676 the anti-FA has been transported over the distance Δ. Thus, at the time $t = t_0 + \frac{\Delta}{u_f}$, the concentration

677 of anti-FA at x_1^Δ is $\overline{F}_0 \frac{u_f t_0}{\Delta}$.

After injecting (23) into (1) and integrating, we have :

$$\int_{F_0}^{F^*} \frac{dF(x_1^\Delta, t)}{F(x_1^\Delta, t)} = -k_1 \overline{F}_0 \frac{u_f}{\Delta} \int_0^{t_1} \left(t - \frac{\Delta}{u_f}\right) \Theta\left(t - \left(t_0 + \frac{\Delta}{u_f}\right)\right) dt$$

$$= -k_1 \overline{F}_0 \frac{u_f}{\Delta} \int_{t_0 + \frac{\Delta}{u_f}}^{t_1} \left(t - \frac{\Delta}{u_f}\right)$$

$$\Rightarrow \log\left(\frac{F^*}{F_0}\right) = -k_1 \overline{F}_0 \frac{u_f}{2\Delta} \left[t^2 - \frac{2\Delta}{u_f} t\right]_{t_0 + \frac{\Delta}{u_f}}^{t_1}$$

$$= -k_1 \overline{F}_0 \frac{u_f}{2\Delta} \left(t_1^2 - \frac{2\Delta}{u_f} t_1 - \left(t_0 + \frac{\Delta}{u_f}\right)^2 + \frac{2\Delta}{u_f}\left(t_0 + \frac{\Delta}{u_f}\right)\right),$$

678 which after rearrangement gives:

$$\left(t_1 - \frac{\Delta}{u_f}\right)^2 - t_0^2 + \frac{2\Delta \log\left(\frac{F^*}{F_0}\right)}{k_1 \overline{F}_0 u_f} = 0. \tag{24}$$

679 Keeping the positive solution of the resulting quadratic equation, we get the time t_1 at which
680 the second slice x_1^Δ is lysed:

$$t_1 = \frac{\Delta}{u_f} + \sqrt{t_0^2 + \frac{2\Delta}{k_1 u_f \overline{F}_0} \log\left(\frac{F_0}{F^*}\right)}. \tag{25}$$

By applying the same reasoning recursively, we get the formula for the time to lyse n slices :

$$t_n = \frac{\Delta}{u_f} + \sqrt{t_{n-1}^2 + \frac{2\Delta}{k_1 u_f \overline{F}_0} \log\left(\frac{F_0}{F^*}\right)},$$

681 **Analytical model independent of Δ**
682 We show in Figure S2 that the analytical formula for the lysis time of the thrombus does not depend
683 on the choice of the spatial slicing Δ, provided that $\Delta \ll L$.

We demonstrate it here analytically, starting with equation (5) and rearranging terms:

$$t_n = \frac{\Delta}{u_f} + \sqrt{t_{n-1}^2 + \frac{2\Delta}{k_1 u_f \overline{F}_0} \log\left(\frac{F_0}{F^*}\right)}$$

$$= \frac{\Delta}{u_f} + \sqrt{t_{n-1}^2 + \beta\Delta}$$

$$\Rightarrow \left(t_n - \frac{\Delta}{u_f}\right)^2 = t_{n-1}^2 + \beta\Delta$$

$$\Rightarrow t_n^2 - 2\frac{\Delta}{u_f} t_n + \frac{\Delta^2}{u_f^2} = t_{n-1}^2 + \beta\Delta.$$

684 By taking the first order approximation of Δ, we neglect the second order term $\frac{\Delta^2}{u_f^2}$, which yields:

$$t_n^2 - t_{n-1}^2 = 2\frac{\Delta}{u_f} t_n + \beta\Delta. \tag{26}$$

685 On the other hand, we can write :

$$t_n^2 - t_{n-1}^2 = (t_n - t_{n-1})(t_n + t_{n-1}) \approx 2t\, dt, \tag{27}$$

where dt is the variation of lysis time at time t for the slice Δ. At first order in Δ, $t_n + t_{n-1} = 2t$ is a

Appendix 1—figure 2. Lysis profiles of a 1 cm thrombus predicted with the analytical formula. $F_0 = 3.2\,\text{mg·ml}^{-1}$, $\overline{F}_0 = 10\,\text{ng·ml}^{-1}$, $\nabla P = 10^6\,\text{Pa·m}^{-1}$, $k_1 = 12.5(\text{mg·ml}^{-1})^{-1}$. The lysis time predicted with the analytical formula is independent of the slicing of the thrombus Δ, is Δ is small compared to L.

sufficient approximation. Combining (26) and (27), and writing $\Delta = dh$ we get:

$$2t\,dt = \left(\frac{2}{u_\text{f}}t + \beta\right)dh \tag{28}$$

$$\Rightarrow \frac{2t}{\frac{2}{u_\text{f}}t + \beta}dt = dh \tag{29}$$

$$\Rightarrow \left(u_\text{f} - \frac{\beta}{\frac{2}{u_\text{f}^2}t + \frac{\beta}{u_\text{f}}}\right)dt = dh. \tag{30}$$

686   By integrating, and setting $h = 0$ at $t = 0$ we finally get:

$$h(t) = u_\text{f}t - \frac{1}{2}\beta u_\text{f}^2 \log\left(2t/u_\text{f}^2 + \beta/u_\text{f}\right) + \frac{1}{2}\beta u_\text{f}^2 \log\left(\beta/u_\text{f}\right), \tag{31}$$

687 which, using the properties of the log, can be rewritten as:

$$h(t) = u_\text{f}\left(t - \frac{1}{2}\beta u_\text{f} \log\left(\frac{2}{\beta u_\text{f}}t + 1\right)\right), \tag{32}$$

688 or:

$$h(t) = u_\text{f}t - \frac{u_\text{f}\log\left(\frac{F_0}{F^*}\right)}{k_1\overline{F}_0}\log\left(\frac{k_1\overline{F}_0}{\log\left(\frac{F_0}{F^*}\right)}t + 1\right), \tag{33}$$

and which is independent of the slicing.

Experimental lysis fronts with various concentrations.

We put at disposition here lysis front measurements for fibrinolysis experiments that we con-
ducted, while varying initial fibrin concentration (Figure 3A), and the tPA solution concentration
(Figure 3B). The dots represent the experimental measurements made at 15 minutes interval, and
the solid lines the analytical model, assuming constant flow throughout the lysis, estimated with
Darcy's law and Davies' equation for the thrombi permeability.

Appendix 1—figure 3. A: Comparison of lysis fronts obtained with the second set of experiments where fibrin concentration is varied, and the analytical model uses optimized reaction parameter obtained by minimizing the RMSE. tPA concentration is $10\,\mathrm{ng\cdot ml^{-1}}$. **B**: Comparison of lysis fronts obtained with the second set of experiments where tPA concentration is varied, and the analytical model uses optimized reaction parameter. We differentiate here two batches of experiments (dots, triangles) that might have suffered slightly different environmental conditions. Fibrin concentration is $3.2\,\mathrm{mg\cdot ml^{-1}}$.

Pressure boundary conditions

Pressure boundary conditions (BCs) were imposed on both ends of the tube by imposing densities, as is usually done in LB. The 3D Zou-He pressure BC *Kutay et al.* (*2006*) and a novel pressure BC were tested. The novel BC showed better simulation stability when increasing the pressure difference; that is why it was chosen for the numerical model. This novel BC can also simulate a pulsatile pressure profile by applying a time modulation to the inlet density. Figure S4 presents an explanation of how the populations at the boundaries are computed.

Let us show how the density is imposed with this pressure BC. We consider the situation displayed in Figure S4, where we need to impose the density on the outlet boundary node N. After having copied the unknown populations f_i from node $N-1$, the density at node N can be computed as usually in LB:

$$\rho_{\mathrm{tmp}} = \sum_{j} f_j(\mathbf{x_N}, t) + \sum_{i} f_i(\mathbf{x_N}, t) = \sum_{k} f_k(\mathbf{x_N}, t) \tag{34}$$

where the f_j designates the known populations from the streaming step, at node N. The equilibrium function f_k^{eq} at node N, used in the standard BGK collision, reads *Krüger et al.* (*2017*):

$$f_k^{eq}(\mathbf{x_N}, t) = w_k\, \rho_{\mathrm{tmp}} \left(1 + \frac{\mathbf{u_N} \cdot \mathbf{c_k}}{c_s^2} + \frac{(\mathbf{u_N} \cdot \mathbf{c_k})^2}{2c_s^4} - \frac{\mathbf{u_N} \cdot \mathbf{u_N}}{2c_s^2}\right) \tag{35}$$

where w_k are the weighting factors, $\mathbf{c_k}$ the lattice discrete velocities, and c_s the speed of sound magnitude in the lattice. By using the property that $\sum_k f_k^{eq}(\mathbf{x_N}, t) = \sum_k f_k(\mathbf{x_N}, t) = \rho_{\mathrm{tmp}}$ at node N, we

subtract and add the following equilibrium functions to the populations f_k at node N:

$$\rho' = \sum_k f_k(\mathbf{x_N}, t) - f_k^{eq}(\rho_{\text{tmp}}, \mathbf{u_N})$$

$$+ f_k^{eq}(\rho_{\text{target}}, \mathbf{u_N})$$

$$= \sum_k f_k(\mathbf{x_N}, t) - \sum_l f_l^{eq}(\rho_{\text{tmp}}, \mathbf{u_N})$$

$$+ \sum_m f_m^{eq}(\rho_{\text{target}}, \mathbf{u_N})$$

$$= \rho_{\text{tmp}} - \rho_{\text{tmp}} + \rho_{\text{target}}$$

$$= \rho_{\text{target}}$$

709 By juggling with these equilibrium functions, we can thus impose the density ρ_{target} at the boundary.
710 It is worth noting that the velocities $\mathbf{u_N}$ remain unchanged.

Appendix 1—figure 4. Pressure boundary condition used for inlet and outlet, with ρ_N the density and u_N the velocity at node N. The computation steps are the following (here showing outlet): **1**. Pre-computation state: At node N, populations f_j are known from the streaming of neighbor nodes at previous iterations, and populations f_i, density, and velocities are unknown. At node $N - 1$, f_k, density and velocities are known. It is chosen as the previous node in the direction normal to the boundary. **2**. Unknown populations at node N are copied from populations at node $N - 1$. From populations f_j and f_i, a temporary density ρ_{tmp} and the velocities are computed. Two equilibrium functions are then computed, one with ρ_{tmp}, and the other with target density ρ_{target}. We subtract the former and add the latter to the populations at node N. **3**. Node N now has target density ρ_{target} and populations f_k.

Particles transport

712 The particles in the numerical model embody the *anti-FA* entity, which accounts for the lysis of
713 fibrin. They are subject to *advection*, the transport driven by the fluid (lytic solution), and *diffusion*,
714 which is the movement due to thermal agitation and molecular collisions.

Advection

The Palabos library offers a built-in algorithm that automatically transports particles advected by
the fluid. However, it is a one-size-fits-all solution and is not necessarily the most efficient way to
advect the particle for a specific model. Looping over all particles at each iteration step is costly. In
our thrombolysis model, the flow varies very slowly during most of the lysis. It is thus interesting
in our case, to increase the advection time scale according to the flow velocity. Practically, instead
of moving on a distance dx the particles at each lattice-Boltzmann iteration, we move them of $T dx$
after T iterations, where T is computed such that the particles move at most of half a lattice site:

$$T = \frac{0.5}{\text{maximum velocity inside domain}(t)}. \tag{36}$$

723 This computational trick allows a speedup of around a factor 10 in a typical lysis simulation, without
724 any visible difference on the lysis curves. The advection velocity of each particle is computed as
725 with the Palabos solver, using weighted averaging of the velocity at the 4 closest lattice nodes.

Diffusion

727 The Péclet number:

$$\text{Pe} = \frac{uL}{D} \tag{37}$$

728 where u is the mean velocity of the fluid, L is a characteristic dimension, and D the diffusion co-
729 efficient of the diffusing mass in the medium, gives an approximation of the ratio of advection
730 against diffusion. For high Péclet (Pe $\gg$ 1), diffusion can be neglected but should otherwise be
731 taken into account. In the clotted tubes of the experiments presented in the manuscript, and for
732 tPA molecules, the initial Péclet is:

$$\text{Pe}_i = \frac{u_{\text{solution}} d_{\text{tube}}}{D_{\text{tPA}}} \sim \frac{3 \cdot 10^{-7} \cdot 10^{-2}}{3.3 \cdot 10^{-11}} \sim 100 \tag{38}$$

733 which shows that the diffusion of tPA is non-negligible to initiate the lysis of the clot.
734   Since the anti-FA is modeled by particles, the most straightforward way to implement their
735 diffusion is by individually giving a random velocity at each simulation time step dt, of norm :

$$|v_{\text{Diff}}| = \frac{\sqrt{2nD_{\text{tPA}}dt}}{dt} \tag{39}$$

736 where $n = 3$ is the spatial dimension of the model.
737   It can be verified that this method yields the expected collective behavior, by computing the
738 mean squared displacement (MSD) of the N particles:

$$\text{MSD}(t) = \frac{1}{N} \sum_{i=1}^{N} |\mathbf{x_i}(t) - \mathbf{x_i}(0)|^2 \tag{40}$$

739 where $\mathbf{x_i}(t)$ is the 3D position of particle i at time t.
740   It can be shown that the MSD approximates the diffusion coefficient:

$$\text{MSD}(t) = 2nD_{\text{tPA}}t. \tag{41}$$

741 By simulating $N = 100$ particles in a system with no advection, already good agreement can be
742 observed with eq. (41), as shown in fig. 5.

Porosity and permeability validation

744 Let us here validate our implementation of the PBB model for computing flow in simulated fibrin
745 clots.
746   Let us recall Darcy's law for laminar flow **Darcy** (**1856**) :

$$u_{\text{f}} = -\frac{k}{\rho\nu} \frac{\Delta P}{L}. \tag{42}$$

747 with ν the kinematic viscosity, ρ the fluid density, ΔP the pressure drop and k the *permeability*.
748   Walsh et al. show that the permeability k_{PBB} in m^2 of PBB voxels is related to γ and the simulation
749 time step Δt as follows:

$$k_{\text{PBB}} = \frac{(1-\gamma)\nu \Delta t}{2\gamma}. \tag{43}$$

750 Combining (14), (42) and (43), we can express Darcy's law in terms of γ:

$$u_{\text{f}} = -\frac{\Delta t}{2\gamma} \frac{\Delta P}{L\rho}. \tag{44}$$

751   We measured the values of the fluid velocity u_{f} in numerical simulations, and compared with
752 the analytical solution of Darcy and Walsh equation (44) in Figure S6A. Flow was simulated in a
753 3D rectangular duct filled with a porous medium with bounce-back fractions of $\gamma = 0.2, 0.4, 0.8$

Appendix 1—figure 5. Diffusion of $N = 100$ anti-FA particles in a simulation without flow. A linear fit is shown between the MSD and the quantity $2nD_{tPA}t$, which slope is 1.007, as expected from equation (41).

respectively. The velocity profile was measured along the y-axis (transversal) in the center of the duct. In the analytical solution, we imposed no-slip conditions, i.e. $u_f = 0$ at the walls.

Next, we investigate the validity of the PBB method for a typical range of fibrin clot permeabilities. Other commonly used laws to describe fibrous porous media permeabilities are Clague's equation *Clague et al.* (*2000*), a numerical solution of permeability of randomly organized fibrous media:

$$k_C(R_f, n_s^*) = 0.50941 R_f^2 \left(\left(\frac{\pi}{4n_s^*} \right)^{0.5} - 1 \right)^2 e^{-1.8042n_s^*}, \tag{45}$$

and Jackson-James' equation *Jackson and James* (*1986*), the weighted average of the solution to the Stokes equation for flow parallel to or normal to 2D periodic square arrays of cylinders:

$$k_{JJ}(R_f, n_s^*) = R_f^2 \frac{3}{20n_s^*}(-\ln(n_s^*) - 0.931). \tag{46}$$

Permeabilities measured with the model as a function of input solid fractions are shown in Figure S6B-D, and compared with their empirical counterparts equations (16), (45) and (46). The permeability values for the physiological range of low solid fractions ($0.003 - 0.04$, i.e. $1 - 4\,\mathrm{mg \cdot ml^{-1}}$ fibrinogen concentration) do not vary substantially from one model to another; thus, if the clots studied are highly porous, the choice of model has little impact. Clague's law predicts an increase of permeability with the increase of solid fraction in the $0.8 - 1$ range, and Jackson-James' law asymptotically approaches zero as n_s^* approaches 0.4. Thus, in order to be able to simulate the whole range of solid fractions, Davies' permeability law was chosen for the model.

Deviation of numerical from analytical model

We show here the deviation of the mesoscopic numerical lysis front, from the front given by the analytical formula, in the conditions of Figure 4. The relative absolute difference ϵ on the front position in time is shown on Figure S7. We see that the deviation stays below 5% during most of the lysis, and increases up to 20% in the last third of the total lysis time. This increase in the difference is due to the assumption of constant flow made in the analytical model.

Lysis of realistic heterogeneous clots

We generated numerical thrombi directly from ischemic stroke patient thrombi retrieved by thrombectomy and sliced by Staessens et al. *Staessens and De Meyer* (*2020*). Platelet stained slices are

Appendix 1—figure 6. A: Comparison of fluid velocity of a transverse slice of a clot in a rectangular duct, between Darcy's law and our implementation of the PBB method, for three bounce-back fractions $\gamma = 0.2, 0.4, 0.8$. Velocity and position are given in lattice units (LU). Space and time steps were $\Delta x = 1.13 \cdot 10^{-4}$ m and $\Delta t = 1.83 \cdot 10^{-5}$ s, $\nu = 3 \cdot 10^{-6}$ m²·s⁻¹, and the pressure gradient $\nabla P = 7700$ Pa·m⁻¹. The domain size was $N_x = N_y = N_z = 48$ lattice nodes. **B,C,D**: Fibrous thrombus intrinsic permeability k as a function of fiber solid fraction n_s^*. Wufsus et al. **Wufsus et al. (2013)** *in vitro* fibrin gels measurements (black +), Davies' equation (B) **Davies (1952)**, Clague (C) **Clague et al. (2000)** and Jackson and James (D) **Jackson and James (1986)**, according to equations (16), (45) and (46) respectively. Simulations were made in a tube of radius 4.5 mm (23 nodes), length 1 cm (46 nodes), with a fully occluding homogeneous thrombus (i.e., same dimensions as the tube). The space-time discretization was $\Delta x = 2.17 \cdot 10^{-4}$ m and $\Delta t = 1.58 \cdot 10^{-3}$ s, and the pressure gradient $\nabla P = 100$ Pa·m⁻¹ The hydrated fiber radius ranges from $R_f = 51$ to 60 nm for formulae and simulations, as in **Wufsus et al. (2013)**.

converted to grayscale using the 'L' algorithm of the Python Imaging Library. We then assign a fib-
rin concentration in a linear scale ranging from $1\,\text{mg}\cdot\text{ml}^{-1}$ (white, no platelets) to $8\,\text{mg}\cdot\text{ml}^{-1}$ (black,
platelet-rich), Figure S8A. Platelet-rich zones are supposed to be more tPA-resistant and to have
lower permeability **Jolugbo and Ariëns (2021)**, **Meglio et al. (2019)**, **Shin et al. (2018)**, which we
translate in the present model with higher fibrin concentration regions. We lysed a RBC-rich clot
and a PLT-rich clot with the numerical model, as well as homogeneous clots with the same average
concentration of 2.45 and $3.92\,\text{mg}\cdot\text{ml}^{-1}$ respectively, Figure S8B. The lysis of the heterogeneous clots
was slightly faster, and the recanalization occurred slightly before, Figure S8C.

Appendix 1—figure 7. Lysis front position deviation in time, between the numerical 2D and analytical models, for different concentrations of fibrin (**A**) and anti-FA (**B**). Baseline parameters are $F_0 = 3 \, \text{mg·ml}^{-1}$, $\overline{F}_0 = 10 \, \text{ng·ml}^{-1}$, $k_1 = 280 \, (\text{s·mg·ml}^{-1})^{-1}$, $\nabla P_0 = 8250 \, \text{Pa·m}^{-1}$, $\Delta x = 1.14 \cdot 10^{-4}$ m, $\Delta t = 1.83 \cdot 10^{-4}$ s.

Appendix 1—figure 8. A: Platelet stained patient clot slices from *Staessens and De Meyer* (*2020*) are digitalized by assigning a fibrin concentration in a linear scale, based on platelet staining. **B**: These clots are lysed with the numerical model, with an imposed pressure gradient $\nabla P = 8000 \, \text{Pa·m}^{-1}$ and a reaction constant $k_1 = 1400 \, (\text{s·mg·ml}^{-1})^{-1}$, and compared against homogeneous clots with the same concentration. **C**: The mean throughput is monitored during the lysis. RBC-rich clots are more permeable since before the start of the lysis, the average fibrin concentration being lower than the PLT-rich's one.