Birds escape the oxidative stress costs of high blood glucose levels

Csongor I. Vágási[1,*], Orsolya Vincze[1,2], Marie Adámková[3,4], Tereza Kauzálová[3], Ádám Z. Lendvai[6], Laura Pătraş[7], Janka Pénzes[1], Péter L. Pap[1,*], Tomáš Albrecht[3,5,*] and Oldřich Tomášek[3,5,*]

[1]Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj-Napoca, Romania. [2]Department of Tisza Research, MTA Centre for Ecological Research-DRI, Debrecen, Hungary. [3]Institute of Vertebrate Biology of the Czech Academy of Sciences, Brno, Czech Republic. [4]Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic. [5]Department of Zoology, Faculty of Science, Charles University, Prague 2, Czech Republic. [6]Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary. [7]Department of Molecular Biology and Biotechnology, Centre of Systems Biology, Biodiversity and Bioresources, Babeş-Bolyai University, Cluj-Napoca, Romania

*Author for correspondence (C.I.V. csvagasi@gmail.com, P.L.P. peterlpap@gmail.com, T.A. albrecht@ivb.cz, O.T. oltmsk@gmail.com)

ORCID ID: C.I.V. 0000-0002-8736-2391, O.V. 0000-0001-5789-2124, M.A. 0000-0003-2156-9194, T.K. 0000-0003-1247-2497, Á.Z.L. 0000-0002-8953-920X, L. P. 0000-0002-6981-3881, P.L.P. 0000-0002-3659-7684, T.A. 0000-0002-9213-0034, O.T. 0000-0002-2657-5916

Running title: Oxidative costs of glucose in birds

KEY WORDS: Antioxidants, Glucose, Hyperglycaemia, Lipid peroxidation, Phylogenetic comparison, Physiological ecology

Summary statement: High blood glucose levels can harm organisms by causing oxidative stress. We show that birds defy this expectation. Avian physiology could provide solutions for diabetes treatment.

ABSTRACT

Chronically high blood glucose levels (hyperglycaemia) can compromise healthy ageing and lifespan at individual level. Elevated oxidative stress can play a central role in hyperglycaemia-induced pathologies. Nevertheless, the lifespan of birds shows no species-level association with blood glucose. This suggests that the potential pathologies of high blood glucose levels can be avoided at the macroevolutionary scale by adaptations in oxidative physiology. However, this hypothesis remains unexplored. Here we examined this hypothesis using comparative analyses controlled for phylogeny, allometry and fecundity based on data from 51 songbird species. We measured blood glucose at baseline and after stress stimulus and computed glucose stress reactivity as the magnitude of change between the two time points. We also measured three parameters of non-enzymatic antioxidants (uric acid, total antioxidants and glutathione) and a marker of oxidative lipid damage (malondialdehyde). We found no or weak evidence for blood glucose concentration coevolving with either antioxidant or lipid damage levels. The only exceptions were the moderate evidence for species with a stronger stress-induced increase in blood glucose concentration evolving moderately lower investment into antioxidant defence (uric acid and glutathione). Neither baseline nor stress-induced glucose levels were associated with oxidative physiology. Our findings essentially support the hypothesis that birds evolved adaptations preventing the putative (glyc)oxidative costs of high blood glucose. Such adaptations may explain the decoupled evolution of glycaemia and lifespan in birds, and possibly the paradoxical combination of long lifespan and high blood glucose levels in birds relative to mammals.

INTRODUCTION

Living organisms use chemical energy to sustain their vital functions, with carbohydrates, lipids, and proteins as the main metabolic energy substrates. Among carbohydrates, glucose has a prominent role in metabolism as it fulfils multiple metabolic functions. Besides being an important energy carrier, it is a vital precursor in the synthesis of a wide variety of other important biomolecules, including fatty, amino, and nucleic acids, cholesterol, glycoproteins, and glycolipids (Braun and Sweazea, 2008; Sweazea, 2022). The mere existence of several regulatory mechanisms that control glucose metabolism both at baseline conditions and during perturbations (e.g. stress response) provides evidence for the biological significance of glucose homeostasis (Braun and Sweazea, 2008). Deregulation of glucose homeostasis induces metabolic stress with a plethora of deleterious health consequences (Picard et al., 2014).

Birds have outstandingly high blood glucose levels in normal physiological state, in average two- to four times higher than that of similar-sized mammals (Braun and Sweazea, 2008; Holmes et al., 2001; Satoh, 2021). Only frugivorous and nectarivorous bats with sugar-rich diet can have blood glucose values comparable to some birds. Constitutively high blood glucose in birds is evolutionarily coupled with athletic capacity, high metabolic rate and elevated body temperature (Satoh, 2021). Moreover, avian blood glucose levels coevolve positively with fecundity suggesting its importance in sustaining intense reproduction (Tomášek et al., 2019; Tomášek et al., 2022). Directional selection therefore might drive the evolution of high blood glucose until it incurs physiological costs that set an upper limit for circulating glucose levels.

Constitutively high blood glucose levels in birds are coupled, at least partly, with their evolved insulin resistance and lower insulin/insulin-like growth factor 1 signalling (Satoh, 2021). Remarkably, the blood glucose levels at physiological baseline of birds would be considered hyperglycaemia in humans and laboratory mammals, where chronic hyperglycaemia is a major health complication of diabetes and is causally involved in a variety of pathologies associated with this disease. Oxidative stress is believed to be the unifying mechanism behind most of the deleterious consequences of high blood glucose levels (Brownlee, 2001). Oxidative stress is a condition when an excessive production of pro-oxidants (i.e. beyond the low amounts needed for the body's biological functions, such as gene translation or white blood cell function) overwhelms the capacity of antioxidant defence and repair system (reviewed by (Monaghan et al., 2009; Pamplona and Costantini, 2011)). This bias towards pro-oxidants inflicts harm to vital cell

components (lipids, proteins and DNA), and the accumulation of oxidative damage can play a role

in many diseases and ageing-related disorders (Buttemer et al., 2010; Finkel and Holbrook, 2000;

Halliwell and Gutteridge, 2007).

Even though glucose is one of the most stable sugars in terms of oxidation, which probably led to

the selection of glucose as a primary metabolic compound in the animal kingdom (Bunn and

Higgins, 1981), high blood glucose is still a significant risk factor for oxidative stress (Barbieri et

al., 2003; Braun and Sweazea, 2008; Ceriello, 2000; Costantini, 2008; Kristal and Yu, 1992).

Specifically, high circulating levels of glucose cause an overproduction of electron donors in the

tricarboxylic acid cycle, which increases mitochondrial proton gradient and superoxide generation

(Du et al., 2000; Nishikawa et al., 2000). Excess superoxide subsequently inhibits glycolysis,

activating the four main pathways of glucose over-utilisation known to cause diabetes-associated

pathologies (Brownlee, 2001). Two of these pathways, namely the polyol pathway and non-

enzymatic glycation further exacerbate oxidative stress by depleting vital intracellular antioxidants

– NADPH and reduced glutathione (GSH) – and through the generation of advanced glycation end-

products (AGEs; (Brownlee, 2001)). AGEs result from proteins or lipids being cross-linked by

dicarbonyls, which arise either as products of direct autoxidation of glucose or from non-enzymatic

reactions of glucose-derived metabolites (Barbieri et al., 2003; Brownlee, 2001; Holmes et al.,

2001). AGEs can ultimately trigger a cascade of oxidation-mediated damage (Brownlee, 2001;

Holmes et al., 2001; Kristal and Yu, 1992). Oxidative stress induced by chronic high blood glucose

levels or AGEs predisposes cells and tissues to dysfunctions, leading to varied pathologies in both

humans and non-human animals (Braun and Sweazea, 2008; Brownlee, 2001; Ceriello, 1997;

Ceriello, 2000; Picard et al., 2014), and may in part be responsible for ageing (Barbieri et al., 2003;

Cerami, 1985; Holmes et al., 2001; Kristal and Yu, 1992). It has been shown that hyperglycaemia-

induced phenotypes can be ameliorated by antioxidants or overexpression of uncoupling proteins

(UCPs), further supporting the involvement of oxidative physiology in diabetes-associated diseases

(Ceriello, 1997; Du et al., 2000; Nishikawa et al., 2000). Experimentally induced diabetes in rats

increases the level of AGEs, carbonyl stress and oxidative stress (e.g. malondialdehyde), whereas

transgenic individuals that overexpress the glyoxalase-I enzyme, which detoxifies AGE precursors,

avoid hyperglycaemia-induced oxidative stress (Brouwers et al., 2011).

Despite the considerable implications for health and healthy ageing, surprisingly few studies

assessed the functional link between blood glucose and oxidative stress in birds or any other free-

living taxa. At the intraspecific level, baseline blood glucose is positively associated with oxidative

damage to lipids (malondialdehyde) in house sparrows *Passer domesticus* (Vágási et al., 2020).
However, it is still unexplored whether these consequences of high blood glucose levels transcend
to the macroevolutionary scale (i.e. whether they can explain variation in oxidative balance among
species). The only available evidence comes from the comparison of birds and mammals, showing
that birds, on average, evolved lower oxidative stress and longer lifespans, despite their higher
levels of blood glucose and AGEs (Baker et al., 2022; Costantini, 2008; Holmes and Ottinger, 2003;
Holmes et al., 2001; Satoh, 2021). These observations suggest that blood glucose concentrations are
evolutionarily decoupled from oxidative stress levels and thus might not constrain the evolution of
long lifespans in birds (see also (Tomášek et al., 2019)). Therefore, it has been proposed that
species evolving high blood glucose levels concomitantly evolved adaptations protecting them from
(glyc)oxidative damage and resultant cellular dysfunction induced by high blood sugar levels
(Braun and Sweazea, 2008; Holmes and Ottinger, 2003; Holmes et al., 2001; Satoh, 2021), which
allowed birds in general to evolve a 'benign hyperglycaemia' (Baker et al., 2022). Among many
adaptations proposed to keep glucose-induced oxidative stress at bay, antioxidants, such as uric
acid, glutathione, or the total antioxidant capacity of the plasma, belong to the most commonly
considered candidates (Braun and Sweazea, 2008; Brouwers et al., 2011; Ceriello, 1997).

Our understanding of the macroevolutionary association between blood glucose and oxidative stress
is hindered by the lack of phylogenetic comparisons across wide ranges of free-living species with
diverse life histories. Here, we fill this gap using data from 51 free-living European songbird
species. Samples were collected from sexually mature individuals during the breeding season,
which provides the least variable data for temperate birds because all species are physiologically set
for breeding, whereas outside the breeding season species markedly differ in moulting patterns
and/or migratory behaviour. Breeding is an energetically demanding period for birds because of
intense sexual and parental activity; therefore, we expect that associations between oxidative
physiology and blood glucose level, if any, are more prominent during this period of the year. We
collected data on blood glucose levels at physiological baseline and after 30 minutes of standardised
stress exposure and computed the magnitude of stress-induced increase in glucose level (termed
'glucose stress reactivity' (Vágási et al., 2020)). While the baseline levels reflect the routine energy
demands of the species, the stress-induced glucose peaks provide an estimate of the species' ability
to mobilise energy stores depleted during the acute stress response (Romero and Beattie, 2022;
Tomášek et al., 2019; Tomášek et al., 2022). The positive correlation between the stress-induced
change levels of glucose and corticosterone are often reported ((Jimeno et al., 2018; Vágási et al.,
2020); but see (Deviche et al., 2016; Romero and Beattie, 2022)) and may adaptively be modulated

according to life-history strategy (Wingfield and Sapolsky, 2003). Therefore, the macroevolutionary

association of both baseline glucose and glucose stress reactivity with oxidative physiology could

illuminate the potential costs of evolving high levels of circulating glucose. To assess blood

oxidative balance of the species, we measured three non-enzymatic antioxidant parameters (total

antioxidant status, uric acid and total glutathione) and the amount of peroxidative damage to lipids

(quantified as malondialdehyde levels). We predicted that species that evolved high baseline or

stress-induced blood glucose levels or high blood glucose stress reactivity also evolved higher

investment into antioxidant defence or otherwise should experience increased oxidative damage.

Alternatively, if the evolution of blood glucose level and that of oxidative homeostasis is decoupled

in birds by adaptations that circumvent the potential adverse consequences of high blood glucose,

we expect no or only weak evidence for the glucose–redox state relationship.

MATERIALS AND METHODS

Field data collection

163 Samples for blood glucose levels were collected in Czech Republic (details and data in (Tomášek et

164 al., 2019; Tomášek et al., 2022)), and those for oxidative physiology in Romania (details and data in

165 (Vágási et al., 2016; Vincze et al., 2022)). Depending on the oxidative physiology variable, 50 or 51

166 species had both glucose and oxidative values (681 individuals with blood glucose and 1021

167 individuals with oxidative stress data).

Biochemical assays

We measured baseline blood glucose concentrations (hereafter G_0) from samples collected within 3

minutes (baseline glucose levels do not increase within this period following the stress stimulus;

(Tomášek et al., 2019)) and stress-induced glucose levels from samples collected after 30 minutes

of standardised restraint in cloth bags (hereafter G_{30}). G_{30} reflects the ability to mobilise circulating

energy in stressful conditions. Species-specific G_0 and G_{30} values are medians (Tomášek et al.,

2019) as they are less affected by outliers than mean values. We computed 'glucose stress

reactivity' (hereafter ΔG) as a percentage change from G_0 median to G_{30} median (where G_0 =

100%) to express the magnitude of change in blood glucose levels when exposed to an acute stress

stimulus. The absolute change (i.e. difference between G_0 and G_{30} medians) is not informative

about the magnitude of change in a multispecies dataset, where species largely differ in these values

(change from G_0 = 5 to G_{30} = 10 mg/dL and from G_0 = 15 to G_{30} = 20 mg/dL are equal in absolute

change [5 mg/dL], while considerably differ in magnitude [100% vs. 33%, respectively]). To assess

oxidative state, we measured three non-enzymatic antioxidant markers (total antioxidant status, TAS; uric acid, UA; and total glutathione, tGSH) and a marker of peroxidative damage to membrane lipids (malondialdehyde, MDA). Detailed protocols can be found in (Bókony et al., 2014). Due to inter-annual variation in oxidative physiology measures, we computed species-specific values as best linear unbiased estimates (BLUEs; (Vincze et al., 2022)). BLUEs were extracted from linear mixed models (using R function *lmer* from R package *lme4* (Bates et al., 2015)), including each oxidative stress parameter as dependent variables (log (x+1) transformed to ensure model residual normality), as well as species as fixed effect and year of capture as a random grouping variable. There was very strong evidence for positive pairwise associations between species-specific TAS, UA and MDA values (Pearson's correlation, all $n = 112$, TAS vs. UA: $r = 0.45$, $P < 0.0001$; TAS vs. MDA: $r = 0.32$, $P = 0.0001$; UA vs. MDA: $r = 0.62$, $P < 0.0001$), while there was no evidence for tGSH being associated with the other oxidative parameters. These findings agree with our previous results based on a considerably smaller oxidative physiology dataset (Vágási et al., 2016).

Life-history traits

Most biological traits depend on body size; thus, we used body mass to account for allometry. Species-specific body mass was extracted from (Storchová and Hořák, 2018), but values were verified using alternative sources and data for some species were updated using recent primary literature or our field measurements (see Table S2 in (Vincze et al., 2022)). Because both blood glucose and oxidative physiology coevolve with pace-of-life as expressed by reproductive investment (Tomášek et al., 2019; Vágási et al., 2019), we also computed species-specific fecundity (i.e. clutch size × no. clutches per year). Data on clutch size and the number of clutches per year were retrieved from (Vágási et al., 2019).

Statistical analyses

All statistical analyses were carried out in R 4.0.4 (R Core Team, 2021). To explore the glucose–oxidative physiology relationships across species, we built phylogenetic generalised least squares (PGLS) models. The BLUEs of the four oxidative physiology parameters were used as response variables in separate models, and the model always included body mass and the examined glucose measure (G_0 or G_{30} or ΔG) as explanatory variables. To assess the potential confounding effect of pace-of-life, we built two model sets, either including or excluding annual fecundity among predictors. The second set of models for TAS, UA and MDA included as additional predictors those

oxidative parameters (as BLUEs) that correlate with the response variable (e.g. UA and MDA in the

TAS model) to assess whether the relationships between the oxidative parameters and glucose

levels are independent of the intercorrelation among these three oxidative parameters. All

explanatory variables were log-transformed to meet the normality assumption of the model

residuals.

Models were constructed using the *gls* function of package *nlme* (Pinheiro et al., 2015). Phylogeny

was downloaded from http://birdtree.org (Jetz et al., 2012). We downloaded 1,000 random trees

using the Hackett backbone (Hackett et al., 2008) and merged them into a rooted, ultrametric

consensus tree using the SumTrees Python library (Sukumaran and Holder, 2010). The phylogenetic

signal was estimated by maximum likelihood and set to the most appropriate value in each model.

Each model was run with multiple model weights: (1) without weights, (2) weighted by log(sample

size +1) and (3) weighted by raw sample size of the dependent variable to control for the sampling

effort (see also (Vágási et al., 2019)). Akaike's Information Criterion (AIC) was used to compare

models with different weights. Unweighted models performed best in the case of the UA and MDA,

models weighted by log sample size had consistently lower AIC values in case of TAS, while

models weighted by sample size provided the best fit in the case of tGSH. Therefore, plots for TAS

and tGSH were prepared to show point sizes proportional to either log sample size or sample size,

respectively. Fitted lines and associated standard errors on the plots were obtained from the

respective full models. Given that physiological and life-history traits often intercorrelate, we

verified multicollinearity in the models by computing the variance inflation factors (VIFs) within

each full model. All VIFs were below the more conservative < 2 threshold, except the

intercorrelating oxidative parameters had higher VIFs, which were below the < 5 threshold (max

VIF = 4.01 for MDA in TAS model). Therefore, multicollinearity is unlikely to influence our

results. Results are communicated throughout using the language of evidence (i.e. how strong the

evidence is for an association) instead of the significance cut-off at $P \leq 0.05$ as suggested by (Muff

et al., 2022).

Results

244   In the first model set that controlled for allometry only, there was no or weak evidence for an

245   association between blood glucose and oxidative stress parameters (Table 1, Table S1). There was

246   weak evidence for a negative association of G_{30} and ΔG levels with tGSH, and for a negative

247   association of G_0 levels with MDA (Table 1, Table S1).

The second model set that controlled for allometry, pace-of-life (fecundity) and correlations among oxidative parameters also revealed only a few weak associations between blood glucose and oxidative state. The data revealed only weak evidence for negative associations of ΔG with UA and tGSH levels (Fig. 1b–c), and of G_{30} with tGSH (Table 1, Table S2). There was no evidence of any other associations between glucose and oxidative physiology parameters (Table 1, Table S2, Fig. 1). In this second model set, there was no evidence for an association between G_0 and MDA (Table 1, Table S2) as compared with the weak association revealed by the first model set (Table 1, Table S1).

Fig. 1. Phylogenetic relationship between glucose stress reactivity (i.e. ΔG, expressed as % change between G_0 and G_{30}, where $G_0 = 100\%$) and levels of (a) total antioxidant status, (b) uric acid, (c) total glutathione and (d) malondialdehyde. Plots (a) and (c) show point sizes proportional to either log sample size or sample size, respectively (see Methods). Fitted lines were obtained from the respective full models from the second model set (see Table S2).

Table 1. The relationship between oxidative and blood glucose parameters in the first and second

sets of phylogenetically controlled models. The first model set analysed the four oxidative

parameters in separate models as a function of body mass and glucose as continuous predictors,

while the second model set as a function of body mass, fecundity, correlated oxidative parameters

and glucose as predictors. Values of oxidative parameters are best linear unbiased predictions

(BLUEs, see Methods). Glucose was measured either at baseline ($logG_0$), after 30 minutes

standardised acute stress exposure ($logG_{30}$) or calculated as % change between G_0 and G_{30} ($log\Delta G$).

Full models with all the predictors together with model parameters for the first and second models

sets are presented in Table S1 and S2, respectively. *P*-values of associations with moderate

evidence (i.e. $0.01 \leq P \leq 0.05$) are marked in bold, while those of weak evidence (i.e. $0.05 \leq P \leq$

0.1) are marked in italics.

response	predictor	first model set				second model set		
oxidative parameters	glucose	β (SE)	t	P		β (SE)	t	P
total antioxidant status (TAS)	$logG_0$	–0.05 (0.06)	0.82	0.414		–0.05 (0.06)	0.84	0.403
	$logG_{30}$	–0.02 (0.03)	0.66	0.513		–0.01 (0.03)	0.16	0.871
	$log\Delta G$	0.01 (0.01)	0.06	0.956		0.01 (0.01)	0.58	0.563
uric acid (UA)	$logG_0$	–0.02 (0.31)	0.07	0.943		0.28 (0.20)	1.43	0.161
	$logG_{30}$	–0.22 (0.15)	1.43	0.159		–0.10 (0.09)	1.10	0.276
	$log\Delta G$	–0.01 (0.01)	1.49	0.143		–0.01 (0.01)	2.15	**0.037**
total glutathione (tGSH)	$logG_0$	–0.01 (0.39)	0.03	0.979		0.02 (0.34)	0.07	0.943
	$logG_{30}$	–0.34 (0.18)	1.85	*0.071*		–0.27 (0.15)	1.85	*0.071*
	$log\Delta G$	–0.01 (0.01)	1.93	*0.060*		–0.01 (0.01)	2.04	**0.048**
malondialdehyde (MDA)	$logG_0$	–0.33 (0.19)	1.78	*0.081*		–0.18 (0.12)	1.44	0.156
	$logG_{30}$	–0.12 (0.09)	1.25	0.219		0.03 (0.06)	0.52	0.609
	$log\Delta G$	–0.01 (0.01)	0.24	0.814		0.01 (0.01)	1.42	0.163

Discussion

We measured oxidative parameters and blood glucose levels in several songbird species in their

natural physiological ranges. Despite the relatively high blood glucose levels in birds compared

with other vertebrates, we found no (or at best only weak) evidence for the association between

blood glucose level and oxidative physiology. The finding that these systems are decoupled

supports the hypothesis that birds have evolved adaptations to circumvent the potential oxidative

challenges and deleterious consequences imposed by their constitutively high blood glucose levels.

Besides such physiological adaptations, life history constraints or confounding effects might also

explain this lack of glucose–redox state relationship in birds.

Physiological adaptations

Birds abound in physiological adaptations that make them excel in physiological homeostasis and

longevity. Here, we discuss four such adaptations that might explain the evolutionary dissociation

of glucose metabolism and oxidative state. These are higher efficacy of non-enzymatic and

structural antioxidant defences, resistance against AGEs, constitutive activation of nuclear factor

erythroid 2-related factor 2 (NRF2) and augmented activity of mitochondrial uncoupling proteins

(UCPs).

The high blood glucose levels of birds would be regarded as hyperglycaemia in mammals of similar

body size. In humans and mammalian models, hyperglycaemic conditions lead to elevated oxidative

stress, which is the central mechanism inducing activation of all the four main pathways known to

generate diabetes-associated pathologies (Brownlee, 2001). In mammalian tissues, high activity of

one of these pathways, the polyol pathway, depletes NADPH and reduces glutathione (GSH)

reserves, which can augment oxidative stress and associated deleterious consequences of diabetes

(Brownlee, 2001). In accordance with this mechanism, we found that bird species that evolved

higher glucose concentrations or higher glucose stress reactivity during the breeding season also

evolved a concomitant lower level of tGSH. Nevertheless, tGSH was only moderately lower in

these species and, importantly, not accompanied by higher oxidative damage. This suggests that

songbirds with high blood glucose might have evolved higher rates of GSSG-to-GSH regeneration,

thereby preventing GSH depletion despite its lower levels and presumably higher glycoxidative

load. The investment into the non-enzymatic antioxidant capacity in general was not considerably

lower in birds with elevated glucose levels as the relationships between the two physiological

systems were mostly absent or at best weak. Besides, birds might have evolved structural

adaptations making their tissues more resistant to oxidative damage (Pamplona and Costantini,

2011).

High blood glucose levels can induce oxidative stress by means other than decreased GSH levels.

The augmented activity of the polyol pathway in hyperglycaemia can generate AGEs, which in turn

can promote oxidative stress by increasing the rate of free radical and highly reactive dialdehyde

(e.g. MDA) formation (Brownlee, 2001; Fukami et al., 2005). Remarkably, MDA was not higher but either unrelated or slightly lower in species with higher baseline glucose levels. This result is consistent with recent findings showing that birds have lower lipid peroxidation compared with mammals (Jimenez et al., 2019), despite having higher glucose and AGE levels ((Baker et al., 2022); but see (Anthony-Regnitz et al., 2020)). Birds, therefore, defy the expectation that more AGEs should induce more lipid peroxidation, suggesting that high blood glucose levels and lipid peroxidation (e.g. MDA production) are uncoupled at macroevolutionary scales in birds. This uncoupling could be underpinned by either lower rates of formation of particular AGEs (e.g. serum albumin; (Anthony-Regnitz et al., 2020)), or higher rates of other AGEs formation but also better clearance of these AGEs or a lack of receptors for AGEs (Baker et al., 2022). These physiological and molecular traits are considered avian adaptations that can circumvent the potential glycoxidative toll of elevated AGE levels.

It has been suggested that Neoaves have elevated expression of antioxidant enzymes and hence higher resistance to oxidative challenge due to the constitutive activation of NRF2 (Castiglione et al., 2020; Satoh, 2021). The plausibility of this hypothesis is supported by the observed ability of the overexpressed antioxidant enzyme superoxide dismutase to prevent hyperglycaemia-induced diabetic complications in mammalian tissues (Brownlee, 2001; Du et al., 2000; Nishikawa et al., 2000). Comparative studies for testing these relationships in birds are desirable.

Augmented activity of avian UCPs is another adaptation that can help to quench the oxidative costs of high blood glucose in birds. UCPs are mitochondrial proton channels that can reduce proton gradient across the inner mitochondrial membrane by leaking protons back into the mitochondrial matrix. Given that hyperglycaemia induces oxidative stress by increasing the proton gradient, which leads to excessive superoxide generation, higher activity of UCPs may represent and important protective adaptation. Although broad-scale comparative studies are lacking, the observation of a considerably higher UCP expression in canaries compared to mice provided promising initial evidence for this hypothesis (Slocinska et al., 2010). Moreover, UCPs have been demonstrated to prevent oxidative stress in both birds and mammals (Criscuolo et al., 2005; Stier et al., 2014) and prolong lifespan in mammals (Andrews and Horvath, 2009; Conti et al., 2006).

Life history constraints and confounds

Physiological systems are evolutionarily linked to life history and behavioural traits leading to the

emergence of pace-of-life syndrome (Réale et al., 2010; Ricklefs and Wikelski, 2002). The

covariations among the traits integrated into the pace-of-life syndrome pose constraints on each

other. Fecundity is a key component of pace-of-life. Increased fecundity requires high blood

glucose levels to fuel the energetic demands of reproduction, and this might have negative

repercussions for oxidative balance. In accordance with these predictions, species with higher

fecundity have higher blood glucose concentrations (Tomášek et al., 2019). Additionally, high

fecundity may affect oxidative physiology in other ways unrelated to its co-evolution with high

blood glucose levels, for example, due to a lower investment into self-maintenance including

antioxidant defences (Vágási et al., 2019), which led to our decision to control for its possible

confounding effect in the second model set. Nevertheless, our findings suggest that fecundity does

not mediate the glucose–redox state relationship as fecundity had little importance in the models.

Species with high baseline glucose levels (i.e. G_0) showed weakly lower MDA levels indicating a

marginally lower rate of lipid peroxidation. Nevertheless, this association merits future studies on

larger number of bird species or even other taxa. A potential explanation for this weak negative

association, other than the candidate adaptations discussed above, could be that there are

confounding variables that were not considered in our models, such as a shift in metabolic substrate

use in migratory species towards higher utilisation of lipids, rather than glucose, to produce energy

during endurance flights (Jenni and Jenni-Eiermann, 1998). In accordance with this hypothesis,

migratory passerines were shown to have lower blood glucose levels than sedentary ones during the

breeding season (Tomášek et al., 2019; Tomášek et al., 2022), suggesting such a fuel shift might

have indeed evolved and became an inherent property of energy metabolism in migratory species

even outside the migratory period. Therefore, such a potential shift in fuel utilisation in long-

distance migrants could potentially confound the relationship between MDA and G_0. Nonetheless,

adding migration distance as a continuous predictor to the second model set of MDA with G_0 did

not change the relationship between MDA and G_0 and only resulted in reduced model fit (results not

shown). Besides, UA is a product of protein catabolism and proteins are also utilised as substrate to

fuel strenuous non-stop migratory flights. Similarly to MDA models, migration distance did not

change the results of UA (results not shown). These findings indicate that potential fuel shift in

migratory species probably does not confound our results.

Diet might also confound the results because more AGEs are formed at higher protein and amino

acid concentrations (Cerami, 1985). Our previous results support this notion as we showed that

carnivorous birds have higher MDA and lower tGSH levels as compared with herbivorous ones

(Marton et al., 2022). However, in the present study we could not assess whether the interaction

between diet and glucose variables influences markers of oxidative state because our samples only

covered songbirds (see (Tomášek et al., 2019)), a group with less variable diet than in the study by

(Marton et al., 2022). Nevertheless, diet probably has little effect on our results because a recent

analysis on a much larger dataset including 160 temperate and tropical songbird species, showed

that only nectarivory/frugivory elevates blood glucose level (Tomášek et al., 2022). Additionally,

although plasma lipids (triglycerides), which may largely be of dietary origin, might correlate with

MDA in some species (Pérez-Rodríguez et al., 2015), this relationship was clearly non-significant

in our dataset (see Supplementary Information in (Vágási et al., 2019)). Therefore, we believe that

MDA mostly derives from cell membrane lipids rather than from dietary triglycerides in our

samples. Finally, a recent review found that diet macro- and micronutrient manipulations generally

have no effect on blood glucose levels in birds (Basile et al., 2022).

In conclusion, the glucose-induced oxidative stress hypothesis is mostly unsupported by our results

in songbirds, indicating that blood glucose levels and redox state are largely decoupled at the

species level. This suggests that bird species with higher blood glucose levels possess adaptations

that prevent glucose-induced oxidative stress, which would otherwise constrain the evolution of

long lifespans. Such adaptations may also possibly explain the particular case of birds, which

evolve considerably longer lifespans compared to mammals, despite having constitutively high

blood glucose levels (referred to as benign hyperglycaemia; (Baker et al., 2022)), the latter being

evolved to promote the high metabolic rates and hyperathletic capabilities of birds (Satoh, 2021;

Sweazea, 2022). Future comparative studies should explore whether components of the

insulin/insulin-like growth factor 1 signalling do coevolve, across a broad range of taxa, with

oxidative balance, blood glucose and the candidate adaptations potentially preventing their harmful

effects (Satoh, 2021). The implied avian adaptations that prevent glucose-induced oxidative stress

and pathologies make birds attractive models for the diabetes research as they might help discover

treatments to ameliorate diabetes-related health problems found in mammals (Hickey et al., 2012;

Sweazea, 2022). Despite these adaptations, birds still can develop hyperglycaemia (with values

above their already high normal physiological levels) and show signs of diabetes. We should also

emphasize that our results are valid for the blood tissue, while the glucose–redox state relationship

might be different in other tissue types, a topic that merits future attention. Future studies should

also assess whether blood glucose levels are or not associated with oxidative parameters in other

vertebrate groups.

Acknowledgements

We owe a debt of gratitude to Attila Marton, Lőrinc Bărbos, Judit Veres-Szászka, Krisztina Sándor, Attila Fülöp, Lukáš Bobek, Ondřej Kauzál, Jaroslav Cepák, Miroslav Čejka, Kryštof Horák, Martin Janča, Sampath A. Kumar and Pavlína Opatová for their help on the field and Manuela Banciu and Alina Sesarman for biochemical advice. We thank the directorate of the Botanical Garden Cluj-Napoca for permitting bird capturing.

Competing interests

No competing interests declared.

Author contributions

CIV, PLP, TA and OT conceived the project; CIV, OV, MA, TK, JP, PLP, TA and OT collected the blood samples; LP and JP carried out the biochemical assays; OV analysed the data with input from CIV, PLP, TA and OT; CIV wrote the paper with significant input from OV, ÁZL, PLP, TA and OT. All authors gave final approval for publication and agreed to be accountable for the aspects of work that they conducted.

Funding

The research was supported by the Romanian Ministry of Research, Innovation and Digitization (CNCS - UEFISCDI; project no. PN-III-P4-ID-PCE-2016-0404 to C.I.V., O.V., J.P. and P.L.P., and project no. PN-III-P1-1.1-TE-2021-0502 to C.I.V. and O.V.) and by the Czech Science Foundation (project no. 21-22160S to T.A. and O.T.). O.V. and P.L.P. were also financed by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, C.I.V. and Á.Z.L. were also supported by a postdoctoral grant of the Hungarian National Research, Development and Innovation Office (#PD 121166 to C.I.V. and K139021 to Á.Z.L.), and O.V. was also supported by New National Excellence Programme of the Hungarian Ministry of Innovation and Technology.

Data availability

Data will be deposited in Figshare repository upon acceptance. All data can be found in (Tomášek et al., 2019; Vincze et al., 2022).

References

Andrews, Z. B. and Horvath, T. L. (2009). Uncoupling protein-2 regulates lifespan in mice. *Am. J. Physiol. Metab.* **296**, E621–E627.

Anthony-Regnitz, C. M., Wilson, A. E., Sweazea, K. L. and Braun, E. J. (2020). Fewer exposed lysine residues may

explain relative resistance of chicken serum albumin to in vitro protein glycation in comparison to bovine serum
albumin. *J. Mol. Evol.* **88**, 653–661.

Baker, P., Cooper-Mullin, C. M. and Jimenez, A. G. (2022). Differences in advanced glycation endproducts (AGEs)
in plasma from birds and mammals of different body sizes and ages. *Comp. Biochem. Physiol. A* **267**, 111164.

Barbieri, M., Rizzo, M. R., Manzella, D., Grella, R., Ragno, E., Carbonella, M., Abbatecola, A. M. and Paolisso,
G. (2003). Glucose regulation and oxidative stress in healthy centenarians. *Exp. Gerontol.* **38**, 137–143.

Basile, A. J., Singh, K. C., Watson, D. F. and Sweazea, K. L. (2022). Effect of macronutrient and micronutrient
manipulation on avian blood glucose concentration: A systematic review. *Comp. Biochem. Physiol. A* **272**,
111279.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *J. Stat.*
Softw. **67**, 1–48.

Bókony, V., Lendvai, Á. Z., Vágási, C. I., Pătraş, L., Pap, P. L., Németh, J., Vincze, E., Papp, S., Preiszner, B.,
Seress, G., et al. (2014). Necessity or capacity? Physiological state predicts problem-solving performance in
house sparrows. *Behav. Ecol.* **25**, 124–135.

Braun, E. J. and Sweazea, K. L. (2008). Glucose regulation in birds. *Comp. Biochem. Physiol. B* **151**, 1–9.

Brouwers, O., Niessen, P. M., Ferreira, I, Miyata, T., Scheffer, P. G., Teerlink, T., Schrauwen, P., Brownlee, M.,
Stehouwer, C. D. and Schalkwijk, C. G. (2011). Overexpression of glyoxalase-I reduces hyperglycemia-
induced levels of advanced glycation end products and oxidative stress in diabetic rats. *J. Biol. Chem.* **286**, 1374–
1380.

Brownlee, M. (2001). Biochemistry and molecular cell biology of diabetic complications. *Nature* **414**, 813–820.

Bunn, H. F. and Higgins, P. J. (1981). Reaction of monosaccharides with proteins: possible evolutionary significance.
Science (80-.). **213**, 222–224.

Buttemer, W. A., Abele, D. and Costantini, D. (2010). From bivalves to birds: oxidative stress and longevity. *Funct.*
Ecol. **24**, 971–983.

Castiglione, G. M., Xu, Z., Zhou, L. and Duh, E. J. (2020). Adaptation of the master antioxidant response connects
metabolism, lifespan and feather development pathways in birds. *Nat. Commun.* **11**, 2476.

Cerami, A. (1985). Hypothesis: glucose as a mediator of aging. *J. Am. Geriatr. Soc.* **33**, 626–634.

Ceriello, A. (1997). Acute hyperglycaemia and oxidative stress generation. *Diabet. Med.* **14**, 45–49.

Ceriello, A. (2000). Oxidative stress and glycemic regulation. *Metabolism* **49**, 27–29.

Conti, B., Sanchez-Alavez, M., Winsky-Sommerer, R., Morale, M. C., Lucero, J., Brownell, S., Fabre, V.,
Huitron-Resendiz, S., Henriksen, S., Zorrilla, E. P., et al. (2006). Transgenic mice with a reduced core body
temperature have an increased life span. *Science (80-.).* **314**, 825–828.

Costantini, D. (2008). Oxidative stress in ecology and evolution: lessons from avian studies. *Ecol. Lett.* **11**, 1238–1251.

Criscuolo, F., Gonzalez-Barroso, M. del M., Bouillaud, F., Ricquier, D., Miroux, B. and Sorci, G. (2005).
Mitochondrial uncoupling proteins: new perspectives for evolutionary ecologists. *Am. Nat.* **166**, 686–699.

Deviche, P., Valle, S., Gao, S., Davies, S., Bittner, S. and Carpentier, E. (2016). The seasonal glucocorticoid

response of male Rufous-winged Sparrows to acute stress correlates with changes in plasma uric acid, but neither
glucose nor testosterone. *Gen. Comp. Endocrinol.* **235**, 78–88.

Du, X.-L., Edelstein, D., Rossetti, L., Fantus, I. G., Goldberg, H., Ziyadeh, F., Wu, J. and Brownlee, M. (2000).
Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces
plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc. Natl. Acad. Sci. USA* **97**,
12222–12226.

Finkel, T. and Holbrook, N. J. (2000). Oxidants, oxidative stress and the biology of ageing. *Nature* **408**, 239–247.

Fukami, K., Cooper, M. E. and Forbes, J. M. (2005). Agents in development for the treatment of diabetic
nephropathy. *Expert Opin. Investig. Drugs* **14**, 279–294.

Hackett, S. J., Kimball, R. T., Reddy, S., Bowie, R. C. K., Braun, E. L., Braun, M. J., Chojnowski, J. L., Cox, W.
A., Han, K.-L., Harshman, J., et al. (2008). A phylogenomic study of birds reveals their evolutionary history.
Science (80-.). **320**, 1763–1768.

Halliwell, B. and Gutteridge, J. M. C. (2007). *Free radicals in biology and medicine*. 4th ed. Oxford, UK: Oxford
University Press.

Hickey, A. J. R., Jüllig, M., Aitken, J., Loomes, K., Hauber, M. E. and Phillips, A. R. J. (2012). Birds and
longevity: Does flight driven aerobicity provide an oxidative sink? *Ageing Res. Rev.* **11**, 242–253.

Holmes, D. and Ottinger, M. (2003). Birds as long-lived animal models for the study of aging. *Exp. Gerontol.* **38**,
1365–1375.

Holmes, D., Flückiger, R. and Austad, S. (2001). Comparative biology of aging in birds: an update. *Exp. Gerontol.*
36, 869–883.

Jenni, L. and Jenni-Eiermann, S. (1998). Fuel supply and metabolic constraints in migrating birds. *J. Avian Biol.* **29**,
521.

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. and Mooers, A. O. (2012). The global diversity of birds in space
and time. *Nature* **491**, 444–448.

Jimenez, A. G., O'Connor, E. S., Tobin, K. J., Anderson, K. N., Winward, J. D., Fleming, A., Winner, C.,
Chinchilli, E., Maya, A., Carlson, K., et al. (2019). Does cellular metabolism from primary fibroblasts and
oxidative stress in blood differ between mammals and birds? The (lack-thereof) scaling of oxidative stress. *Integr.*
Comp. Biol. **59**, 953–969.

Jimeno, B., Hau, M. and Verhulst, S. (2018). Corticosterone levels reflect variation in metabolic rate, independent of
'stress.' *Sci. Rep.* **8**, 13020.

Kristal, B. S. and Yu, B. P. (1992). An emerging hypothesis: synergistic induction of aging by free radicals and
Maillard reactions. *J. Gerontol.* **47**, B107–B114.

Marton, A., Vágási, C. I., Vincze, O., Bókony, V., Pap, P. L., Pătraş, L., Pénzes, J., Bărbos, L., Fülöp, A., Osváth,
G., et al. (2022). Oxidative physiology is weakly associated with pigmentation in birds. *Ecol. Evol.* **12**, 1–12.

Monaghan, P., Metcalfe, N. B. and Torres, R. (2009). Oxidative stress as a mediator of life history trade-offs:
mechanisms, measurements and interpretation. *Ecol. Lett.* **12**, 75–92.

Muff, S., Nilsen, E. B., O'Hara, R. B. and Nater, C. R. (2022). Rewriting results sections in the language of evidence.

Trends Ecol. Evol. **37**, 203–210.

Nishikawa, T., Edelstein, D., Du, X. L., Yamagishi, S., Matsumura, T., Kaneda, Y., Yorek, M. A., Beebe, D., **Oates, P. J., Hammes, H.-P., et al.** (2000). Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature* **404**, 787–790.

Pamplona, R. and Costantini, D. (2011). Molecular and structural antioxidant defenses against oxidative stress in animals. *Am. J. Physiol. Integr. Comp. Physiol.* **301**, R843–R863.

Pérez-Rodríguez, L., Romero-Haro, A. A., Sternalski, A., Muriel, J., Mougeot, F., Gil, D. and Alonso-Alvarez, C. (2015). Measuring oxidative stress: the confounding effect of lipid concentration in measures of lipid peroxidation. *Physiol. Biochem. Zool.* **88**, 345–351.

Picard, M., Juster, R.-P. and McEwen, B. S. (2014). Mitochondrial allostatic load puts the "gluc" back in glucocorticoids. *Nat. Rev. Endocrinol.* **10**, 303–310.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team (2015). *nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-120*. http://CRAN.R-project.org/package=nlme.

R Core Team (2021). *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V. and Montiglio, P.-O. (2010). Personality and the emergence of the pace-of-life syndrome concept at the population level. *Philos. Trans. R. Soc. London B* **365**, 4051–4063.

Ricklefs, R. E. and Wikelski, M. (2002). The physiology/life-history nexus. *Trends Ecol. Evol.* **17**, 462–468.

Romero, L. M. and Beattie, U. K. (2022). Common myths of glucocorticoid function in ecology and conservation. *J. Exp. Zool. A* **337**, 7–14.

Satoh, T. (2021). Bird evolution by insulin resistance. *Trends Endocrinol. Metab.* **32**, 803–813.

Slocinska, M. B., Almsherqi, Z. A. M., Sluse, F. E., Navet, R. and Deng, Y. (2010). Identification and characterization of uncoupling protein in heart and muscle mitochondria of canary birds. *J. Bioenerg. Biomembr.* **42**, 345–353.

Stier, A., Massemin, S. and Criscuolo, F. (2014). Chronic mitochondrial uncoupling treatment prevents acute cold-induced oxidative stress in birds. *J. Comp. Physiol. B* **184**, 1021–1029.

Storchová, L. and Hořák, D. (2018). Life-history characteristics of European birds. *Glob. Ecol. Biogeogr.* **27**, 400–406.

Sukumaran, J. and Holder, M. T. (2010). DendroPy: a Python library for phylogenetic computing. *Bioinformatics* **26**, 1569–1571.

Sweazea, K. L. (2022). Revisiting glucose regulation in birds – A negative model of diabetes complications. *Comp. Biochem. Physiol. B* **262**, 110778.

Tomášek, O., Bobek, L., Kralova, T., Adámková, M. and Albrecht, T. (2019). Fuel for the pace of life: baseline blood glucose concentration co-evolves with life-history traits in songbirds. *Funct. Ecol.* **33**, 239–249.

Tomášek, O., Bobek, L., Kauzálová, T., Kauzál, O., Adámková, M., Horák, K., Kumar, S. A., Manialeu, J. P.,

Munclinger, P., Nana, E. D., et al. (2022). Latitudinal but not elevational variation in blood glucose level is
linked to life history across passerine birds. *Ecol. Lett.* **25**, 2203–2216.

Vágási, C. I., Vincze, O., Pătraş, L., Osváth, G., Marton, A., Bărbos, L., Sol, D. and Pap, P. L. (2016). Large-
brained birds suffer less oxidative damage. *J. Evol. Biol.* **29**, 1968–1976.

Vágási, C. I., Vincze, O., Pătraş, L., Osváth, G., Pénzes, J., Haussmann, M. F., Barta, Z. and Pap, P. L. (2019).
Longevity and life history coevolve with oxidative stress in birds. *Funct. Ecol.* **33**, 152–161.

Vágási, C. I., Tóth, Z., Pénzes, J., Pap, P. L., Ouyang, J. Q. and Lendvai, Á. Z. (2020). The Relationship between
Hormones, Glucose, and Oxidative Damage Is Condition and Stress Dependent in a Free-Living Passerine Bird.
Physiol. Biochem. Zool. **93**, 466–476.

Vincze, O., Vágási, C. I., Pénzes, J., Szabó, K., Magonyi, N. M., Czirják, G. Á. and Pap, P. L. (2022). Sexual
dimorphism in immune function and oxidative physiology across birds: The role of sexual selection. *Ecol. Lett.*
25, 958–970.

Wingfield, J. C. and Sapolsky, R. M. (2003). Reproduction and resistance to stress: when and how. *J.*
Neuroendocrinol. **15**, 711–724.

Tables S1 and S2

Table S1. First set of phylogenetically controlled models analysing the four oxidative parameters in

separate models (*a–d*) with body mass (log BM) and glucose (log Glu) as continuous predictors.

Values of oxidative parameters are best linear unbiased estimates (BLUEs, see main text Methods).

Different columns show models run with glucose measured either at baseline (G_0, column (1)), after

30 minutes standardised acute stress exposure (G_{30}, column (2)) or measured as % change between

G_0 and G_{30} (ΔG, column (3)). Model parameters are provided in the last row of each model (λ –

Pagel's lambda, AIC – Akaike's Information Criterion, *n* – number of species). *P*-values of

associations with moderate or strong evidence (i.e. $0.001 \leq P \leq 0.05$) are marked in bold, while

those of weak evidence (i.e. $0.05 \leq P \leq 0.1$) are marked in italics.

response	(1) logGlu = logG_0			(2) logGlu = logG_{30}			(3) logGlu = logΔG		
predictor	β (SE)	*t*	*P*	β (SE)	*t*	*P*	β (SE)	*t*	*P*
(a) total antioxidant status (TAS) model									
intercept	0.14 (0.16)	0.92	0.364	0.07 (0.08)	0.82	0.415	0.01 (0.04)	0.33	0.745
logBM	–0.01 (0.01)	0.63	0.533	–0.01 (0.01)	0.48	0.632	–0.01 (0.01)	0.30	0.766
logGlu	–0.05 (0.06)	0.82	0.414	–0.02 (0.03)	0.66	0.513	0.01 (0.01)	0.06	0.956
model: λ = 0.00, AIC = –167.81, *n* = 50				λ = 0.00, AIC = –166.12, *n* = 50			λ = 0.00, AIC = –156.53, *n* = 50		
(b) uric acid (UA) model									
intercept	0.31 (0.86)	0.36	0.720	0.85 (0.45)	1.89	0.065	0.50 (0.23)	2.15	0.037
logBM	–0.06 (0.05)	1.20	0.235	–0.07 (0.05)	1.47	0.148	–0.06 (0.05)	1.26	0.214
logGlu	–0.02 (0.31)	0.07	0.943	–0.22 (0.15)	1.43	0.159	–0.01 (0.01)	1.49	0.143
model: λ = 0.59, AIC = –5.71, *n* = 50				λ = 0.57, AIC = –6.29, *n* = 50			λ = 0.60, AIC = 2.73, *n* = 50		
(c) total glutathione (tGSH) model									
intercept	–0.53 (1.07)	0.49	0.625	0.38 (0.56)	0.68	0.501	–0.17 (0.30)	0.57	0.571
logBM	0.14 (0.07)	1.94	*0.058*	0.13 (0.07)	1.87	*0.067*	0.14 (0.07)	2.19	**0.033**
logGlu	–0.01 (0.39)	0.03	0.979	–0.34 (0.18)	1.85	*0.071*	–0.01 (0.01)	1.93	*0.060*
model: λ = 0.75, AIC = 16.77, *n* = 51				λ = 0.80, AIC = 14.98, *n* = 51			λ = 0.78, AIC = 23.87, *n* = 51		
(d) malondialdehyde (MDA) model									
intercept	1.08 (0.51)	2.12	0.039	0.51 (0.28)	1.83	0.073	0.21 (0.15)	1.43	0.160
logBM	–0.08 (0.03)	2.32	**0.025**	–0.06 (0.03)	2.01	**0.050**	–0.06 (0.03)	1.86	*0.070*
logGlu	–0.33 (0.19)	1.78	*0.081*	–0.12 (0.09)	1.25	0.219	–0.01 (0.01)	0.24	0.814
model: λ = 0.86, AIC = –52.50, *n* = 50				λ = 0.75, AIC = –49.61, *n* = 50			λ = 0.76, AIC = –38.91, *n* = 50		

Table S2. Second set of phylogenetically controlled models analysing the four oxidative parameters
(TAS – total antioxidant status, UA – uric acid, tGSH – total glutathione, and MDA –
malondialdehyde) in separate models (*a–d*) with body mass (log BM), fecundity (log Fec),
correlating oxidative parameters (TAS and/or UA and/or MDA) and glucose (log Glu) as predictors.
Values of oxidative parameters are best linear unbiased estimates (BLUEs, see main text Methods).
Different columns show models run with glucose measured either at baseline (G_0, column (1)), after
30 minutes standardised acute stress exposure (G_{30}, column (2)) or measured as % change between
G_0 and G_{30} (ΔG, column (3)). Model parameters are provided in the last row of each model (λ –
Pagel's lambda, AIC – Akaike's Information Criterion, *n* – number of species). *P*-values of
associations with moderate or strong evidence (i.e. $0.001 \leq P \leq 0.05$) are marked in bold, while
those of weak evidence (i.e. $0.05 \leq P \leq 0.1$) are marked in italics.

response	(1) logGlu = logG_0			(2) logGlu = logG_{30}			(3) logGlu = logΔG		
predictor	β (SE)	*t*	*P*	β (SE)	*t*	*P*	β (SE)	*t*	*P*
(a) total antioxidant status (TAS) model									
intercept	0.12 (0.15)	0.81	0.422	0.01 (0.08)	0.13	0.900	–0.02 (0.05)	0.42	0.679
logBM	0.01 (0.01)	0.06	0.951	0.01 (0.01)	0.43	0.671	0.01 (0.01)	0.52	0.605
logFec	0.01 (0.01)	0.11	0.913	–0.01 (0.01)	0.07	0.944	–0.01 (0.01)	0.17	0.869
UA	0.05 (0.04)	1.13	0.264	0.04 (0.04)	0.94	0.353	0.05 (0.04)	1.11	0.272
MDA	0.04 (0.07)	0.64	0.523	0.06 (0.07)	0.85	0.402	0.05 (0.07)	0.69	0.494
logGlu	–0.05 (0.06)	0.84	0.403	–0.01 (0.03)	0.16	0.871	0.01 (0.01)	0.58	0.563
model: $\lambda = 0.00$, AIC = –160.71, *n* = 50				$\lambda = 0.00$, AIC = –158.48, *n* = 50			$\lambda = 0.00$, AIC = –149.66, *n* = 50		
(b) uric acid (UA) model									
intercept	–0.86 (0.52)	1.65	0.107	0.14 (0.28)	0.50	0.617	0.09 (0.16)	0.58	0.564
logBM	0.04 (0.03)	1.35	0.183	0.01 (0.03)	0.48	0.630	0.02 (0.03)	0.65	0.522
logFec	0.03 (0.04)	0.79	0.433	0.06 (0.04)	1.41	0.166	0.05 (0.04)	1.38	0.174
TAS	0.61 (0.54)	1.13	0.264	0.51 (0.54)	0.94	0.353	0.58 (0.52)	1.11	0.272
MDA	1.27 (0.15)	8.64	**< 0.001**	1.25 (0.15)	8.38	**< 0.001**	1.24 (0.14)	8.67	**< 0.001**
logGlu	0.28 (0.20)	1.43	0.161	–0.10 (0.09)	1.10	0.276	–0.01 (0.01)	2.15	**0.037**
model: $\lambda = 0.00$, AIC = –50.37, *n* = 50				$\lambda = 0.00$, AIC = –48.03, *n* = 50			$\lambda = 0.00$, AIC = –42.05, *n* = 50		
(c) total glutathione (tGSH) model									
intercept	–0.47 (0.86)	0.55	0.588	0.28 (0.43)	0.64	0.523	–0.08 (0.28)	0.29	0.773
logBM	0.12 (0.05)	2.17	**0.035**	0.11 (0.05)	2.10	**0.041**	0.12 (0.05)	2.30	**0.026**
logFec	0.01 (0.08)	0.14	0.893	0.03 (0.07)	0.44	0.662	0.01 (0.07)	0.04	0.968
logGlu	0.02 (0.34)	0.07	0.943	–0.27 (0.15)	1.85	*0.071*	–0.01 (0.01)	2.04	**0.048**
model: $\lambda = 0.82$, AIC = –1.68, *n* = 51				$\lambda = 0.89$, AIC = –3.14, *n* = 51			$\lambda = 0.89$, AIC = 5.54, *n* = 51		
(d) malondialdehyde (MDA) model									
intercept	0.65 (0.32)	2.02	0.050	0.12 (0.18)	0.71	0.482	0.10 (0.10)	1.02	0.311
logBM	–0.04 (0.02)	2.52	**0.016**	–0.03 (0.02)	2.05	**0.046**	–0.03 (0.02)	2.15	**0.037**

logFec	−0.04 (0.03)	1.61	0.114		−0.06 (0.02)	2.37	**0.022**		−0.06 (0.02)	2.41	**0.020**
TAS	0.22 (0.33)	0.66	0.516		0.29 (0.34)	0.85	0.402		0.23 (0.34)	0.69	0.494
UA	0.49 (0.06)	8.49	**< 0.001**		0.49 (0.06)	8.38	**< 0.001**		0.51 (0.06)	8.67	**< 0.001**
logGlu	−0.18 (0.12)	1.44	0.156		0.03 (0.06)	0.52	0.609		0.01 (0.01)	1.42	0.163
model: $\lambda = 0.13$, AIC = −89.10, $n = 50$					$\lambda = 0.00$, AIC = −87.98, $n = 50$				$\lambda = 0.00$, AIC = −80.53, $n = 50$		