Songbirds avoid the oxidative stress costs of high blood glucose levels

Csongor I. Vágási[1,*], Orsolya Vincze[1,2], Marie Adámková[3,4], Tereza Kauzálová[3], Ádám Z. Lendvai[6], Laura Pătraş[7], Janka Pénzes[1], Péter L. Pap[1,*], Tomáš Albrecht[3,5,*] and Oldřich Tomášek[3,5,*]

[1]Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj-Napoca, Romania. [2]Department of Tisza Research, MTA Centre for Ecological Research-DRI, Debrecen, Hungary. [3]Institute of Vertebrate Biology of the Czech Academy of Sciences, Brno, Czech Republic. [4]Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic. [5]Department of Zoology, Faculty of Science, Charles University, Prague 2, Czech Republic. [6]Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary. [7]Department of Molecular Biology and Biotechnology, Centre of Systems Biology, Biodiversity and Bioresources, Babeş-Bolyai University, Cluj-Napoca, Romania

*Author for correspondence (C.I.V. csvagasi@gmail.com, P.L.P. peterlpap@gmail.com, T.A. albrecht@ivb.cz, O.T. oltmsk@gmail.com)

ORCID ID: C.I.V. 0000-0002-8736-2391, O.V. 0000-0001-5789-2124, M.A. 0000-0003-2156-9194, T.K. 0000-0003-1247-2497, Á.Z.L. 0000-0002-8953-920X, L.P. 0000-0002-6981-3881, P.L.P. 0000-0002-3659-7684, T.A. 0000-0002-9213-0034, O.T. 0000-0002-2657-5916

Running title: No oxidative costs of glucose in songbirds

KEY WORDS: Antioxidants, Glucose, Hyperglycaemia, Lipid peroxidation, Phylogenetic comparison, Physiological ecology

Summary statement: High blood glucose levels can harm organisms by causing oxidative stress. We show that, at the macroevolutionary level, songbirds defy this expectation, as their glucose levels and oxidative physiology are uncoupled.

ABSTRACT

Chronically high blood glucose levels (hyperglycaemia) can compromise healthy ageing

and lifespan at the individual level. Elevated oxidative stress can play a central role in

hyperglycaemia-induced pathologies. Nevertheless, the lifespan of birds shows no

species-level association with blood glucose. This suggests that the potential pathologies

of high blood glucose levels can be avoided by adaptations in oxidative physiology at the

macroevolutionary scale. However, this hypothesis remains unexplored. Here, we

examined this hypothesis using comparative analyses controlled for phylogeny, allometry

and fecundity based on data from 51 songbird species (681 individuals with blood glucose

and 1021 individuals with oxidative state data). We measured blood glucose at baseline

and after stress stimulus and computed glucose stress reactivity as the magnitude of

change between the two time points. We also measured three parameters of non-

enzymatic antioxidants (uric acid, total antioxidants and glutathione) and a marker of

oxidative lipid damage (malondialdehyde). We found no clear evidence for blood glucose

concentration being correlated with either antioxidant or lipid damage levels at the

macroevolutionary scale, as opposed to the hypothesis postulating that high blood glucose

levels entail oxidative costs. The only exception was the moderate evidence for species

with a stronger stress-induced increase in blood glucose concentration evolving

moderately lower investment into antioxidant defence (uric acid and glutathione). Neither

baseline nor stress-induced glucose levels were associated with oxidative physiology. Our

findings support the hypothesis that birds evolved adaptations preventing the

(glyc)oxidative costs of high blood glucose observed at the within-species level. Such

adaptations may explain the decoupled evolution of glycaemia and lifespan in birds and

possibly the paradoxical combination of long lifespan and high blood glucose levels

relative to mammals.

INTRODUCTION

Living organisms use chemical energy to sustain their vital functions, with carbohydrates, lipids, and proteins as the main metabolic energy substrates. Among carbohydrates, glucose has a prominent role in metabolism as it fulfils multiple metabolic functions. Besides being an important energy carrier, it is a vital precursor in the synthesis of a wide variety of other important biomolecules, including fatty, amino, and nucleic acids, cholesterol, glycoproteins, and glycolipids (Braun and Sweazea, 2008; Sweazea, 2022). The mere existence of several regulatory mechanisms that control glucose metabolism both at baseline conditions and during perturbations (e.g. stress response) provides evidence for the biological significance of glucose homeostasis (Braun and Sweazea, 2008). Deregulation of glucose homeostasis induces metabolic stress with a plethora of deleterious health consequences (Picard et al., 2014).

Birds have outstandingly high blood glucose levels in normal physiological state, in average two- to four times higher than that of similar-sized mammals (Braun and Sweazea, 2008; Holmes et al., 2001; Satoh, 2021). Only frugivorous and nectarivorous bats with a sugar-rich diet can have blood glucose values comparable to some birds (Kelm et al., 2011; Peng et al., 2017). Constitutively high blood glucose in birds is evolutionarily coupled with athletic capacity, high metabolic rate and elevated body temperature (Satoh, 2021). Moreover, avian blood glucose levels coevolve positively with fecundity, suggesting its importance in sustaining intense reproduction (Tomášek et al., 2019; Tomášek et al., 2022). Directional selection therefore might drive the evolution of high blood glucose until it incurs physiological costs that set an upper limit for circulating glucose levels.

Constitutively high blood glucose levels in birds are coupled, at least partly, with their evolved insulin resistance and lower insulin/insulin-like growth factor 1 signalling (Satoh, 2021). Remarkably, the blood glucose levels at physiological baseline of birds would be considered hyperglycaemia in humans and laboratory mammals, where chronic hyperglycaemia is a major health complication of diabetes and is causally involved in a variety of pathologies associated with this disease (Sweazea, 2022). Oxidative stress is believed to be the unifying mechanism behind most of the deleterious consequences of high blood glucose levels (Brownlee, 2001). Oxidative stress is a condition when an excessive production of pro-oxidants (i.e. beyond the low amounts needed for the body's biological functions, such as gene translation or white blood cell function) overwhelms the capacity of antioxidant defence and repair system (reviewed by (Monaghan et al., 2009; Pamplona and Costantini, 2011)). This bias towards pro-oxidants inflicts harm to vital cell

components (lipids, proteins and DNA), and the accumulation of oxidative damage can play a role

in many diseases and ageing-related disorders (Buttemer et al., 2010; Finkel and Holbrook, 2000;

Halliwell and Gutteridge, 2007).

Even though glucose is one of the most stable sugars in terms of oxidation, which probably led to

the selection of glucose as a primary metabolic compound in the animal kingdom (Bunn and

Higgins, 1981), high blood glucose is still a significant risk factor for oxidative stress (Barbieri et

al., 2003; Braun and Sweazea, 2008; Ceriello, 2000; Costantini, 2008; Kristal and Yu, 1992).

Specifically, high circulating levels of glucose cause an overproduction of electron donors in the

tricarboxylic acid cycle, which increases mitochondrial proton gradient and superoxide generation

(Du et al., 2000; Nishikawa et al., 2000). Excess superoxide subsequently inhibits glycolysis,

activating the four main pathways of glucose over-utilisation known to cause diabetes-associated

pathologies (Brownlee, 2001). Two of these pathways, namely the polyol pathway and non-

enzymatic glycation further exacerbate oxidative stress by depleting vital intracellular antioxidants

– NADPH and reduced glutathione (GSH) – and through the generation of advanced glycation end-

products (AGEs; (Brownlee, 2001)). AGEs result from proteins or lipids being cross-linked by

dicarbonyls, which arise either as products of direct autoxidation of glucose or from non-enzymatic

reactions of glucose-derived metabolites (Barbieri et al., 2003; Brownlee, 2001; Holmes et al.,

2001). AGEs can ultimately trigger a cascade of oxidation-mediated damage (Brownlee, 2001;

Holmes et al., 2001; Kristal and Yu, 1992). Oxidative stress induced by chronic high blood glucose

levels or AGEs predisposes cells and tissues to dysfunctions, leading to varied pathologies in both

humans and non-human animals (Braun and Sweazea, 2008; Brownlee, 2001; Ceriello, 1997;

Ceriello, 2000; Picard et al., 2014), and may in part be responsible for ageing (Barbieri et al., 2003;

Cerami, 1985; Holmes et al., 2001; Kristal and Yu, 1992). It has been shown that hyperglycaemia-

induced phenotypes can be ameliorated by antioxidants or overexpression of uncoupling proteins

(UCPs), further supporting the involvement of oxidative physiology in diabetes-associated diseases

(Ceriello, 1997; Du et al., 2000; Nishikawa et al., 2000). Experimentally induced diabetes in rats

increases the level of AGEs, carbonyl stress and oxidative stress (e.g. malondialdehyde), whereas

transgenic individuals that overexpress the glyoxalase-I enzyme, which detoxifies AGE precursors,

avoid hyperglycaemia-induced oxidative stress (Brouwers et al., 2011).

Despite the considerable implications for health and healthy ageing, surprisingly few studies

assessed the functional link between blood glucose and oxidative stress in birds or any other free-

living taxa. At the intraspecific level, baseline blood glucose is positively associated with oxidative

damage to lipids (malondialdehyde) in house sparrows *Passer domesticus* (Vágási et al., 2020).

However, it is still unexplored whether these consequences of high blood glucose levels transcend

to the macroevolutionary scale (i.e. whether they can explain variation in oxidative balance among

species). The only available evidence comes from the comparison of birds and mammals, showing

that birds, on average, evolved lower oxidative stress and longer lifespans, despite their higher

levels of blood glucose and AGEs (Baker et al., 2022; Costantini, 2008; Holmes and Ottinger, 2003;

Holmes et al., 2001; Satoh, 2021). These observations suggest that blood glucose concentrations are

evolutionarily decoupled from oxidative stress levels and thus might not constrain the evolution of

long lifespans in birds (see also (Tomášek et al., 2019)). Therefore, it has been proposed that

species evolving high blood glucose levels concomitantly evolved adaptations protecting them from

(glyc)oxidative damage and resultant cellular dysfunction induced by high blood sugar levels

(Braun and Sweazea, 2008; Holmes and Ottinger, 2003; Holmes et al., 2001; Satoh, 2021), which

allowed birds in general to evolve a 'benign hyperglycaemia' (Baker et al., 2022). Among many

adaptations proposed to keep glucose-induced oxidative stress at bay, antioxidants, such as uric

acid, glutathione, or the total antioxidant capacity of the plasma, belong to the most commonly

considered candidates (Braun and Sweazea, 2008; Brouwers et al., 2011; Ceriello, 1997).

Our understanding of the macroevolutionary association between blood glucose and oxidative stress

is hindered by the lack of phylogenetic comparisons across wide ranges of free-living species with

diverse life histories. Here, we fill this gap using data from 51 free-living European songbird

species sampled during breeding. We collected data on blood glucose levels at physiological

baseline and after 30 minutes of standardised stress exposure and computed the magnitude of stress-

induced increase in glucose level (termed 'glucose stress reactivity' (Vágási et al., 2020)). While

the baseline levels reflect the routine energy demands of the species, the stress-induced glucose

peaks provide an estimate of the species' ability to mobilise energy stores depleted during the acute

stress response (Romero and Beattie, 2022; Tomášek et al., 2019; Tomášek et al., 2022). The

positive correlation between the stress-induced change levels of glucose and corticosterone are

often reported ((Jimeno et al., 2018; Vágási et al., 2020); but see exceptions (Deviche et al., 2016;

Romero and Beattie, 2022)) and may adaptively be modulated according to life-history strategy

(Wingfield and Sapolsky, 2003). Therefore, the macroevolutionary association of both baseline

glucose and glucose stress reactivity with oxidative physiology could illuminate the potential costs

of evolving high levels of circulating glucose. To assess blood oxidative balance of the species, we

measured three non-enzymatic antioxidant parameters (total antioxidant status, uric acid and total

glutathione) and the amount of peroxidative damage to lipids (quantified as malondialdehyde

levels). Given the hypothesis that high blood glucose levels are detrimental to health partly by

causing oxidative stress, we predicted that species that evolved high baseline also evolved higher

investment into antioxidant defence or otherwise should experience increased oxidative damage. An

alternative hypothesis is that the evolution of blood glucose level and that of oxidative homeostasis

is decoupled in birds by adaptations that circumvent the potential adverse consequences of high

blood glucose. In this case, we expect no or only weak evidence for the glucose–redox state

relationship. There is also a possibility that the stress-induced blood glucose level or glucose stress

reactivity may be more important in defining species-specific oxidative physiology than the

baseline level. To explore this hypothesis, we examined the relationship between the stress-induced

blood glucose parameters and oxidative state variables, despite the fact that the latter are not

influenced by capture stress (Vágási et al., 2016).

MATERIALS AND METHODS

Field data collection

Samples for blood glucose levels were collected from birds captured in Czech Republic from April

to July in 2014 and 2019 (details and data in (Tomášek et al., 2019; Tomášek et al., 2022)), and

those for oxidative physiology from birds captured in Romania from April to July between 2011

and 2013, and between 2016 and 2019 (details and data in (Vágási et al., 2016; Vincze et al.,

2022)). Depending on the oxidative physiology variable, 50 or 51 species had both glucose and

oxidative values (681 individuals with blood glucose and 1021 individuals with oxidative stress

data). Samples were collected from adults during the breeding season. Because small-sized

songbirds rarely skip breeding (Snow et al., 1998), and thus all individuals in reproductive mode,

this sampling period ensures the lowest possible noise in physiological measures. The physiological

readiness for breeding was confirmed by the presence of breeding patch in females and cloacal

protuberances and sperm production in males (for detailed information, see (Horák et al., 2022;

Vágási et al., 2016)). Outside the breeding season, species markedly differ in energy demanding

processes, including substantial variation in timing and extent of moulting and/or migratory

behaviour (ranging from sedentary to short- and long-distance migration). Therefore, during

breeding all small-sized songbird individuals are physiologically in breeding mode, which makes

possible to compare physiological traits across species.

Biochemical parameters

We measured baseline blood glucose concentrations (hereafter G_0) with FreeStyle Freedom Lite

portable glucose meters (Abbott Diabetes Care; linear range: 1.1– 27.8 mM) from samples collected

within 3 minutes (baseline glucose levels do not increase within this period following the stress

stimulus; (Tomášek et al., 2019)) and stress-induced glucose levels from samples collected after 30

minutes of standardised restraint in cloth bags (hereafter G_{30}). G_{30} reflects the ability to mobilise

circulating energy in stressful conditions. Species-specific G_0 and G_{30} values are medians (Tomášek

et al., 2019) as they are less affected by outliers than mean values. We computed 'glucose stress

reactivity' (hereafter ΔG) as a percentage change from G_0 median to G_{30} median (where G_0 =

100%) to express the magnitude of change in blood glucose levels when exposed to an acute stress

stimulus. The absolute change (i.e. difference between G_0 and G_{30} medians) is not informative

about the magnitude of change in a multispecies dataset, where species largely differ in these values

(e.g. change from G_0 = 5 to G_{30} = 10 mg/dL and from G_0 = 15 to G_{30} = 20 mg/dL are equal in

absolute change [5 mg/dL], while considerably differ in magnitude [100% vs. 33%, respectively]).

As expected, mean G_{30} values are significantly higher than G_0 values in our dataset based on 51 bird

species (phylogenetic paired t-test, $t = -5.20$, df = 48, $P < 0.0001$) in accordance with a recent result

from our extended dataset of 160 bird species (Tomášek et al., 2022).

To assess oxidative state, we measured three non-enzymatic antioxidant markers (total antioxidant

status, TAS; uric acid, UA; and total glutathione, tGSH) and a marker of peroxidative damage to

membrane lipids (malondialdehyde, MDA). Detailed protocols can be found in (Bókony et al.,

2014). Briefly, TAS, UA and MDA were measured from plasma, while tGSH from erythrocytes.

TAS level (mM Trolox equivalents) was measured based on a commercial kit (Cayman Chemical,

Ann Arbor, MI) according to the method described by (Erel, 2004), with slight modifications as per

(Sepp et al., 2010). This assay relies on the ability of non-enzymatic antioxidants (e.g. uric acid,

vitamins, sulfhydryl groups of proteins, glutathione) to decolorize the blue-green $ABTS^+$ (2,2'-

azino-bis(3-ethylbenzothiazoline-6-sulfonate)) to a degree proportional to their concentrations,

which can be measured spectrophotometrically at 660 nm (see (Bókony et al., 2014)). UA

concentration (mg/dL plasma) was determined spectrophotometrically from 5 µL of plasma by an

uricase/peroxidase method (Uric Acid liquicolor kit, Human, Wiesbaden, Germany) (see (Bókony

et al., 2014)). MDA is a carbonyl compound that results from the peroxidative degeneration of

membrane polyunsaturated fatty acids by reactive oxygen species, and thus it is a widely used

marker of oxidative stress (Del Rio et al., 2005). MDA concentration (µg/mL plasma) was

determined from 10 µL of plasma by High Performance Liquid Chromatography (HPLC) on a

HPLC SUPELCOSIL^{TM} LC-18 column (5 µm particle size; Sigma-Aldrich) with UV detection at

254 nm (Jasco, UV-2075 Plus, Japan) (see (Bókony et al., 2014)). GSH is the most important

intracellular, endogenous, non-enzymatic antioxidant (Galván and Alonso-Alvarez, 2008). The

concentration of tGSH (nM/mg of erythrocyte pellet) was assayed using a commercial kit (Sigma-

Aldrich, St Louis, MO) and according to (Galván and Alonso-Alvarez, 2008) and (Hõrak et al.,

2010) with the following modifications (see (Bókony et al., 2014)). Blood samples drawn from

small species were usually only enough for one aliquot, and therefore different oxidative state

markers were measured from samples from different individuals of the same species resulting in

variable sample size for different parameters across species (Vágási et al., 2016).

Due to inter-annual variation in oxidative physiology measures, we computed species-specific

values as best linear unbiased estimates (BLUEs; (Vincze et al., 2022)). BLUEs were extracted

from linear mixed models (using R function *lmer* from R package *lme4* (Bates et al., 2015)),

including each oxidative stress parameter as dependent variables (log (x+1) transformed to ensure

model residual normality), as well as species as fixed effect and year of capture as a random

grouping variable. There was very strong evidence for positive pairwise associations between

species-specific TAS, UA and MDA values (Pearson's correlation, all $n = 112$, TAS vs. UA: $r =$

0.45, $P < 0.0001$; TAS vs. MDA: $r = 0.32$, $P = 0.0001$; UA vs. MDA: $r = 0.62$, $P < 0.0001$), while

there was no evidence for tGSH being associated with the other oxidative parameters. These

findings agree with our previous results based on a considerably smaller oxidative physiology

dataset (Vágási et al., 2016). We have previously shown that all blood glucose (G_0, G_{30} and ΔG)

and oxidative state variables measured here are suitable for among-species comparative analyses

because they fulfil the criterion of being species-specific, as demonstrated by their significant

repeatability within species (i.e., conspecifics resemble each other more than members of other

species; see details in (Marton et al., 2022; Tomášek et al., 2022)).

Life-history traits

Most biological traits depend on body size; thus, we used body mass to account for allometry.

Species-specific body mass was extracted from (Storchová and Hořák, 2018), but values were

verified using alternative sources and data for some species were updated using recent primary

literature or our field measurements (see Table S2 in (Vincze et al., 2022)). Because both blood

glucose and oxidative physiology coevolve with pace-of-life as expressed by reproductive

investment (Tomášek et al., 2019; Vágási et al., 2019), we also computed species-specific annual

fecundity (i.e. clutch size × no. clutches per year). Data on clutch size and the number of clutches

per year were retrieved from (Vágási et al., 2019).

Statistical analyses

All statistical analyses were carried out in R 4.0.4 (R Core Team, 2021). To explore the glucose–oxidative physiology relationships across species, we built two sets of phylogenetic generalised least squares (PGLS) models. In the first model set, the BLUEs of the four oxidative physiology parameters were used as response variables in separate models, while body mass and the examined glucose measure (G_0 or G_{30} or ΔG) were continuous explanatory variables. In the second model set, besides body mass, we also controlled for the potential effect of additional confounding variables. Annual fecundity was included in all models of the second set to rule out the potential confounding effect of pace-of-life. In the models of TAS, UA and MDA, we also included as additional predictors those oxidative parameters (as BLUEs) that correlate with the response variable (e.g. UA and MDA in the TAS model) to rule out the potentially confounding effect of intercorrelating oxidative parameters (see above). All explanatory variables were log-transformed to meet the normality assumption of the model residuals.

Models were constructed using the *gls* function of package *nlme* (Pinheiro et al., 2015). Phylogeny was downloaded from http://birdtree.org (Jetz et al., 2012). We downloaded 1000 random trees using the Hackett backbone (Hackett et al., 2008) and merged them into a rooted, ultrametric consensus tree using the SumTrees Python library (Sukumaran and Holder, 2010). The phylogenetic signal was estimated by maximum likelihood and set to the most appropriate value in each model. Each model was run with multiple model weights: (1) without weights, (2) weighted by log(sample size +1) and (3) weighted by raw sample size of the dependent variable to control for the sampling effort (see also (Vágási et al., 2019)). Akaike's Information Criterion (AIC) was used to compare models with different weights. Unweighted models performed best in the case of the UA and MDA, models weighted by log sample size had consistently lower AIC values in case of TAS, while models weighted by sample size provided the best fit in the case of tGSH. Therefore, plots for TAS and tGSH were prepared to show point sizes proportional to either log sample size or sample size, respectively. Fitted lines and associated standard errors on the plots were obtained from the respective full models. Given that physiological and life-history traits often intercorrelate, we verified multicollinearity in the models by computing the variance inflation factors (VIFs) within each full model. All VIFs were below the more conservative < 2 threshold, except the intercorrelating oxidative parameters had higher VIFs, which were below the < 5 threshold (max VIF = 4.01 for MDA in TAS model). Therefore, multicollinearity is unlikely to influence our results. Results are communicated throughout using the language of evidence (i.e. how strong the

287   evidence is for an association) instead of the significance cut-off at $P \le 0.05$ as suggested by (Muff

288   et al., 2022).

Results

In the first model set that controlled for allometry only, there was no or weak evidence for an

association between blood glucose and oxidative stress parameters (Table 1, Table S1). There was

weak evidence for a negative association of G_{30} and ΔG levels with tGSH, and for a negative

association of G_0 levels with MDA (Table 1, Table S1).

The second model set that controlled for allometry, pace-of-life (fecundity) and correlations among

oxidative parameters also revealed only a few weak associations between blood glucose and

oxidative state. The data revealed only weak evidence for negative associations of ΔG with UA and

tGSH levels (Fig. 1b–c), and of G_{30} with tGSH (Table 1, Table S2). There was no evidence of any

other associations between glucose and oxidative physiology parameters (Table 1, Table S2, Fig. 1).

In this second model set, there was no evidence for an association between G_0 and MDA (Table 1,

Table S2) as compared with the weak association revealed by the first model set (Table 1, Table

S1).

Discussion

Our study is the first to examine the oxidative costs (or lack thereof) of high blood glucose levels at

the macroevolutionary level. Despite the relatively high blood glucose levels in birds compared

with other vertebrates, we found no (or at best only weak) evidence for the macroevolutionary

association between blood glucose level and oxidative physiology. The finding that these systems

are decoupled supports the hypothesis that birds have evolved adaptations to circumvent the

potential oxidative challenges and deleterious consequences imposed by their constitutively high

blood glucose levels. Besides such physiological adaptations, life history constraints or confounding

effects might also explain this lack of glucose–redox state relationship in birds.

Physiological adaptations

Birds abound in physiological adaptations that make them excel in physiological homeostasis and

longevity. Here, we discuss four such adaptations that might explain the evolutionary dissociation

of glucose metabolism and oxidative state. These are higher efficacy of non-enzymatic and

structural antioxidant defences (Pamplona and Costantini, 2011), resistance against AGEs (Baker et
al., 2022), constitutive activation of nuclear factor erythroid 2-related factor 2 (NRF2; (Castiglione
et al., 2020; Satoh, 2021)) and augmented activity of mitochondrial uncoupling proteins (UCPs;
(Criscuolo et al., 2005; Stier et al., 2014)).

The high blood glucose levels of birds would be regarded as hyperglycaemia in mammals of similar
body size. In humans and mammalian models, hyperglycaemic conditions lead to elevated oxidative
stress, which is the central mechanism inducing activation of all the four main pathways known to
generate diabetes-associated pathologies (Brownlee, 2001). In mammalian tissues, high activity of
one of these pathways, the polyol pathway, depletes NADPH and reduces glutathione (GSH)
reserves, which can augment oxidative stress and associated deleterious consequences of diabetes
(Brownlee, 2001). In accordance with this mechanism, we found that bird species that evolved
higher glucose concentrations or higher glucose stress reactivity during the breeding season also
evolved a concomitant lower level of tGSH. Nevertheless, tGSH was only moderately lower in
these species and, importantly, not accompanied by higher oxidative damage. This suggests that
songbirds with high blood glucose might have evolved higher rates of GSSG-to-GSH regeneration,
thereby preventing GSH depletion despite its lower levels and presumably higher glycoxidative
load. The investment into the non-enzymatic antioxidant capacity in general was not considerably
lower in birds with elevated glucose levels as the relationships between the two physiological
systems were mostly absent or at best weak. Besides, birds might have evolved structural
adaptations making their tissues more resistant to oxidative damage (Pamplona and Costantini,
2011).

High blood glucose levels can induce oxidative stress by means other than decreased GSH levels.
The augmented activity of the polyol pathway in hyperglycaemia can generate AGEs, which in turn
can promote oxidative stress by increasing the rate of free radical and highly reactive dialdehyde
(e.g. MDA) formation (Brownlee, 2001; Fukami et al., 2005). Remarkably, MDA was not higher
but either unrelated or slightly lower in species with higher baseline glucose levels. This result is
consistent with recent findings showing that birds have lower lipid peroxidation compared with
mammals (Jimenez et al., 2019), despite having higher glucose and AGE levels ((Baker et al.,
2022); but see (Anthony-Regnitz et al., 2020)). Birds, therefore, defy the expectation that more
AGEs should induce more lipid peroxidation, suggesting that high blood glucose levels and lipid
peroxidation (e.g. MDA production) are uncoupled at macroevolutionary scales in birds. This
uncoupling could be underpinned by either lower rates of formation of particular AGEs (e.g. serum

albumin; (Anthony-Regnitz et al., 2020)), or higher rates of other AGEs formation but also better

clearance of these AGEs or a lack of receptors for AGEs (Baker et al., 2022). These physiological

and molecular traits are considered avian adaptations that can circumvent the potential

glycoxidative toll of elevated AGE levels.

It has been suggested that Neoaves have elevated expression of antioxidant enzymes and hence

higher resistance to oxidative challenge due to the constitutive activation of NRF2 (Castiglione et

al., 2020; Satoh, 2021). The plausibility of this hypothesis is supported by the observed ability of

the overexpressed antioxidant enzyme superoxide dismutase to prevent hyperglycaemia-induced

diabetic complications in mammalian tissues (Brownlee, 2001; Du et al., 2000; Nishikawa et al.,

2000). Comparative studies for testing these relationships in birds are desirable.

Augmented activity of avian UCPs is another adaptation that can help to quench the oxidative costs

of high blood glucose in birds. UCPs are mitochondrial proton channels that can reduce proton

gradient across the inner mitochondrial membrane by leaking protons back into the mitochondrial

matrix. Given that hyperglycaemia induces oxidative stress by increasing the proton gradient, which

leads to excessive superoxide generation, higher activity of UCPs may represent and important

protective adaptation. Although broad-scale comparative studies are lacking, the observation of a

considerably higher UCP expression in canaries compared to mice provided promising initial

evidence for this hypothesis (Slocinska et al., 2010). Moreover, UCPs have been demonstrated to

prevent oxidative stress in both birds and mammals (Criscuolo et al., 2005; Stier et al., 2014) and

prolong lifespan in mammals (Andrews and Horvath, 2009; Conti et al., 2006).

Life history constraints and other potential confounding variables

Physiological systems are evolutionarily linked to life history and behavioural traits leading to the

emergence of pace-of-life syndrome (Réale et al., 2010; Ricklefs and Wikelski, 2002). The

covariations among the traits integrated into the pace-of-life syndrome pose constraints on each

other. Fecundity is a key component of pace-of-life. Increased fecundity requires high blood

glucose levels to fuel the energetic demands of reproduction, and this might have negative

repercussions for oxidative balance. In accordance with these predictions, species with higher

fecundity have higher blood glucose concentrations (Tomášek et al., 2019). Additionally, high

fecundity may affect oxidative physiology in other ways unrelated to its co-evolution with high

blood glucose levels, for example, due to a lower investment into self-maintenance including

antioxidant defences (Vágási et al., 2019), which led to our decision to control for its possible

confounding effect in the second model set. Nevertheless, our findings suggest that fecundity does

not mediate or confound the glucose–redox state relationship as fecundity had little importance in

the models.

Species with high baseline glucose levels (i.e. G_0) showed weakly lower MDA levels indicating a

marginally lower rate of lipid peroxidation. This association merits future studies on larger number

of bird species or even other taxa. A potential explanation for this weak negative association, other

than the candidate adaptations discussed above, could be that there are confounding variables that

were not considered in our models, such as a shift in metabolic substrate use in migratory species

towards higher utilisation of lipids, rather than glucose, to produce energy during endurance flights

(Jenni and Jenni-Eiermann, 1998). In accordance with this hypothesis, migratory passerines were

shown to have lower blood glucose levels than sedentary ones during the breeding season (Tomášek

et al., 2019; Tomášek et al., 2022), suggesting such a fuel shift might have indeed evolved and

became an inherent property of energy metabolism in migratory species even outside the migratory

period. Therefore, such a potential shift in fuel utilisation in long-distance migrants could

potentially confound the relationship between MDA and G_0. Nonetheless, adding migration

distance as a continuous predictor to the second model set of MDA with G_0 did not change the

relationship between MDA and G_0 and only resulted in reduced model fit (results not shown).

Besides, UA is a product of protein catabolism and proteins are also utilised as substrate to fuel

strenuous non-stop migratory flights. Similarly to MDA models, migration distance did not change

the results of UA (results not shown). These findings indicate that potential fuel shift in migratory

species probably does not explain our results.

Diet might also confound the results because more AGEs are formed at higher protein and amino

acid concentrations (Cerami, 1985). Our previous results support this notion as we showed that

carnivorous birds have higher MDA and lower tGSH levels as compared with herbivorous ones

(Marton et al., 2022). However, in the present study we could not assess whether the interaction

between diet and glucose variables influences markers of oxidative state because our samples only

covered songbirds (see (Tomášek et al., 2019)), a group with less variable diet than in the study by

(Marton et al., 2022). Nevertheless, diet probably has little effect on our results because a recent

analysis on a much larger dataset including 160 temperate and tropical songbird species, showed

that only nectarivory/frugivory elevates blood glucose level (Tomášek et al., 2022), but this diet

type is missing in our dataset. Additionally, although plasma lipids (triglycerides), which may

largely be of dietary origin, might correlate with MDA in some species (Pérez-Rodríguez et al.,

2015), this relationship was clearly non-significant in our dataset (see Supplementary Information

in (Vágási et al., 2019)). Therefore, we believe that MDA mostly derives from cell membrane lipids

rather than from dietary triglycerides in our samples. Finally, a recent review found that diet macro-

and micronutrient manipulations generally have no effect on blood glucose levels in birds (Basile et

al., 2022).

Besides, any existent causal relationship between blood glucose and oxidative physiology could be

masked under some circumstances because the blood glucose dataset and oxidative physiology

dataset were collected from different individuals in two different countries. However, the general

requirement of any phylogenetic comparative study is that the among-species differences in trait

values that evolved over macroevolutionary timescales are larger than the variation at the within-

species level (e.g., sex- or population-related variation), ensuring that macroevolutionary

relationships among traits can be detected despite the within-species noise. This assumption is

supported when the traits of interest show significant within-species repeatability, which we

demonstrated for both blood glucose parameters and oxidative physiology parameters in our

previous studies (Tomášek et al., 2022; Vágási et al., 2016). Geographical variation, if additive,

should not obscure existing macroevolutionary relationships. To mask such relationships, we would

need to assume interactions between traits and different environments. While we cannot rule out

geographic interactions entirely, they seem unlikely given similar temperate climates and latitudes

in both sample locations (only ca 3° latitude difference). Moreover, analysing data from different

sources is a standard practice in comparative studies, including those focused on physiological

traits. As opposed to many such comparative studies, however, our dataset is based on highly

standardised sampling and lab methodology each physiological trait measured in one lab thereby

avoiding potentially considerable interlaboratory variation (Fanson et al., 2017).

In conclusion, the glucose-induced oxidative stress hypothesis is mostly unsupported by our results

in songbirds, indicating that blood glucose levels and redox state are largely decoupled at the

species level. This suggests that bird species with higher blood glucose levels possess adaptations

that prevent glucose-induced oxidative stress, which would otherwise constrain the evolution of

long lifespans. Such adaptations may also possibly explain the particular case of birds, which

evolve considerably longer lifespans compared to mammals, despite having constitutively high

blood glucose levels (referred to as benign hyperglycaemia; (Baker et al., 2022)), the latter being

evolved to promote the high metabolic rates and hyperathletic capabilities of birds (Satoh, 2021;

Sweazea, 2022). Future comparative studies should explore whether components of the

insulin/insulin-like growth factor 1 signalling do coevolve, across a broad range of taxa, with

oxidative balance, blood glucose and the candidate adaptations potentially preventing their harmful

effects (Satoh, 2021). The implied avian adaptations that prevent glucose-induced oxidative stress

and pathologies make birds attractive models for diabetes research as they might help discover

treatments to ameliorate diabetes-related health problems found in mammals (Hickey et al., 2012;

Sweazea, 2022). Despite these adaptations, birds still can develop hyperglycaemia (with values

above their already high normal physiological levels) and show signs of diabetes (Sweazea, 2022).

We should also emphasize that our results are valid for the blood tissue, while the glucose–redox

state relationship might be different in other tissue types, a topic that merits future attention. Future

studies should also assess the association between blood glucose levels and oxidative parameters in

other vertebrate groups.

Acknowledgements

We owe a debt of gratitude to Attila Marton, Lőrinc Bărbos, Judit Veres-Szászka, Krisztina Sándor, Attila
Fülöp, Lukáš Bobek, Ondřej Kauzál, Jaroslav Cepák, Miroslav Čejka, Kryštof Horák, Martin Janča, Sampath
A. Kumar and Pavlína Opatová for their help on the field and Manuela Banciu and Alina Sesarman for
biochemical advice. We thank the directorate of the Botanical Garden Cluj-Napoca for permitting bird
capturing. Constructive criticism provided by two anonymous reviewers considerably improved the article.

Competing interests

No competing interests declared.

Author contributions

CIV, PLP, TA and OT conceived the project; CIV, OV, MA, TK, JP, PLP, TA and OT collected the blood
samples; LP and JP carried out the biochemical assays; OV analysed the data with input from CIV, PLP, TA
and OT; CIV wrote the paper with significant input from OV, ÁZL, PLP, TA and OT. All authors gave final
approval for publication and agreed to be accountable for the aspects of work that they conducted.

Funding

The research was supported by the Romanian Ministry of Research, Innovation and Digitization (CNCS -
UEFISCDI; project no. PN-III-P1-1.1-TE-2021-0502 to C.I.V. and O.V.) and by the Czech Science
Foundation (project no. 21-22160S to T.A. and O.T.). O.V. and P.L.P. were also financed by the János
Bolyai Research Scholarship of the Hungarian Academy of Sciences, C.I.V. was supported by a postdoctoral
grant (PD 121166) and Á.Z.L. by a research grant (K139021) of the Hungarian National Research,

Development and Innovation Office, and O.V. was also supported by New National Excellence Programme
of the Hungarian Ministry of Innovation and Technology.

Data availability

493 All data used in this study are archived at Figshare (https://doi.org/10.6084/m9.figshare.24242935.v1).

References

Andrews, Z. B. and Horvath, T. L. (2009). Uncoupling protein-2 regulates lifespan in mice. *Am. J. Physiol. Metab.*
296, E621–E627.

Anthony-Regnitz, C. M., Wilson, A. E., Sweazea, K. L. and Braun, E. J. (2020). Fewer exposed lysine residues may
explain relative resistance of chicken serum albumin to in vitro protein glycation in comparison to bovine serum
albumin. *J. Mol. Evol.* **88**, 653–661.

Baker, P., Cooper-Mullin, C. M. and Jimenez, A. G. (2022). Differences in advanced glycation endproducts (AGEs)
in plasma from birds and mammals of different body sizes and ages. *Comp. Biochem. Physiol. A* **267**, 111164.

Barbieri, M., Rizzo, M. R., Manzella, D., Grella, R., Ragno, E., Carbonella, M., Abbatecola, A. M. and Paolisso,
G. (2003). Glucose regulation and oxidative stress in healthy centenarians. *Exp. Gerontol.* **38**, 137–143.

Basile, A. J., Singh, K. C., Watson, D. F. and Sweazea, K. L. (2022). Effect of macronutrient and micronutrient
manipulation on avian blood glucose concentration: A systematic review. *Comp. Biochem. Physiol. A* **272**,
111279.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *J. Stat.*
Softw. **67**, 1–48.

Bókony, V., Lendvai, Á. Z., Vágási, C. I., Pătraş, L., Pap, P. L., Németh, J., Vincze, E., Papp, S., Preiszner, B.,
Seress, G., et al. (2014). Necessity or capacity? Physiological state predicts problem-solving performance in
house sparrows. *Behav. Ecol.* **25**, 124–135.

Braun, E. J. and Sweazea, K. L. (2008). Glucose regulation in birds. *Comp. Biochem. Physiol. B* **151**, 1–9.

Brouwers, O., Niessen, P. M., Ferreira, I., Miyata, T., Scheffer, P. G., Teerlink, T., Schrauwen, P., Brownlee, M.,
Stehouwer, C. D. and Schalkwijk, C. G. (2011). Overexpression of glyoxalase-I reduces hyperglycemia-
induced levels of advanced glycation end products and oxidative stress in diabetic rats. *J. Biol. Chem.* **286**, 1374–
1380.

Brownlee, M. (2001). Biochemistry and molecular cell biology of diabetic complications. *Nature* **414**, 813–820.

Bunn, H. F. and Higgins, P. J. (1981). Reaction of monosaccharides with proteins: possible evolutionary significance.
Science (80-.). **213**, 222–224.

Buttemer, W. A., Abele, D. and Costantini, D. (2010). From bivalves to birds: oxidative stress and longevity. *Funct.*
Ecol. **24**, 971–983.

Castiglione, G. M., Xu, Z., Zhou, L. and Duh, E. J. (2020). Adaptation of the master antioxidant response connects

metabolism, lifespan and feather development pathways in birds. *Nat. Commun.* **11**, 2476.

Cerami, A. (1985). Hypothesis: glucose as a mediator of aging. *J. Am. Geriatr. Soc.* **33**, 626–634.

Ceriello, A. (1997). Acute hyperglycaemia and oxidative stress generation. *Diabet. Med.* **14**, 45–49.

Ceriello, A. (2000). Oxidative stress and glycemic regulation. *Metabolism* **49**, 27–29.

Conti, B., Sanchez-Alavez, M., Winsky-Sommerer, R., Morale, M. C., Lucero, J., Brownell, S., Fabre, V.,
Huitron-Resendiz, S., Henriksen, S., Zorrilla, E. P., et al. (2006). Transgenic mice with a reduced core body
temperature have an increased life span. *Science (80-.).* **314**, 825–828.

Costantini, D. (2008). Oxidative stress in ecology and evolution: lessons from avian studies. *Ecol. Lett.* **11**, 1238–1251.

Criscuolo, F., Gonzalez☐Barroso, M. del M., Bouillaud, F., Ricquier, D., Miroux, B. and Sorci, G. (2005).
Mitochondrial uncoupling proteins: new perspectives for evolutionary ecologists. *Am. Nat.* **166**, 686–699.

Del Rio, D., Stewart, A. J. and Pellegrini, N. (2005). A review of recent studies on malondialdehyde as toxic
molecule and biological marker of oxidative stress. *Nutr. Metab. Cardiovasc. Dis.* **15**, 316–328.

Deviche, P., Valle, S., Gao, S., Davies, S., Bittner, S. and Carpentier, E. (2016). The seasonal glucocorticoid
response of male Rufous-winged Sparrows to acute stress correlates with changes in plasma uric acid, but neither
glucose nor testosterone. *Gen. Comp. Endocrinol.* **235**, 78–88.

Du, X.-L., Edelstein, D., Rossetti, L., Fantus, I. G., Goldberg, H., Ziyadeh, F., Wu, J. and Brownlee, M. (2000).
Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces
plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc. Natl. Acad. Sci. USA* **97**,
12222–12226.

Erel, O. (2004). A novel automated direct measurement method for total antioxidant capacity using a new generation,
more stable ABTS radical cation. *Clin. Biochem.* **37**, 277–285.

Fanson, K. V., Németh, Z., Ramenofsky, M., Wingfield, J. C. and Buchanan, K. L. (2017). Inter☐laboratory
variation in corticosterone measurement: Implications for comparative ecological and evolutionary studies.
Methods Ecol. Evol. **8**, 1745–1754.

Finkel, T. and Holbrook, N. J. (2000). Oxidants, oxidative stress and the biology of ageing. *Nature* **408**, 239–247.

Fukami, K., Cooper, M. E. and Forbes, J. M. (2005). Agents in development for the treatment of diabetic
nephropathy. *Expert Opin. Investig. Drugs* **14**, 279–294.

Galván, I. and Alonso-Alvarez, C. (2008). An intracellular antioxidant determines the expression of a melanin-based
signal in a bird. *PLoS One* **3**, e3335.

Hackett, S. J., Kimball, R. T., Reddy, S., Bowie, R. C. K., Braun, E. L., Braun, M. J., Chojnowski, J. L., Cox, W.
A., Han, K.-L., Harshman, J., et al. (2008). A phylogenomic study of birds reveals their evolutionary history.
Science (80-.). **320**, 1763–1768.

Halliwell, B. and Gutteridge, J. M. C. (2007). *Free radicals in biology and medicine*. 4th ed. Oxford, UK: Oxford
University Press.

Hickey, A. J. R., Jüllig, M., Aitken, J., Loomes, K., Hauber, M. E. and Phillips, A. R. J. (2012). Birds and
longevity: Does flight driven aerobicity provide an oxidative sink? *Ageing Res. Rev.* **11**, 242–253.

Holmes, D. and Ottinger, M. (2003). Birds as long-lived animal models for the study of aging. *Exp. Gerontol.* **38**, 1365–1375.

Holmes, D., Flückiger, R. and Austad, S. (2001). Comparative biology of aging in birds: an update. *Exp. Gerontol.* **36**, 869–883.

Horák, K., Bobek, L., Adámková, M., Kauzál, O., Kauzálová, T., Manialeu, J. P., Nguelefack, T. B., Nana, E. D., Jønsson, K. A., Munclinger, P., et al. (2022). Feather growth and quality across passerines is explained by breeding rather than moulting latitude. *Proc. R. Soc. London B* **289**,.

Hõrak, P., Sild, E., Soomets, U., Sepp, T. and Kilk, K. (2010). Oxidative stress and information content of black and yellow plumage coloration: an experiment with greenfinches. *J. Exp. Biol.* **213**, 2225–2233.

Jenni, L. and Jenni-Eiermann, S. (1998). Fuel supply and metabolic constraints in migrating birds. *J. Avian Biol.* **29**, 521.

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. and Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature* **491**, 444–448.

Jimenez, A. G., O'Connor, E. S., Tobin, K. J., Anderson, K. N., Winward, J. D., Fleming, A., Winner, C., Chinchilli, E., Maya, A., Carlson, K., et al. (2019). Does cellular metabolism from primary fibroblasts and oxidative stress in blood differ between mammals and birds? The (lack-thereof) scaling of oxidative stress. *Integr. Comp. Biol.* **59**, 953–969.

Jimeno, B., Hau, M. and Verhulst, S. (2018). Corticosterone levels reflect variation in metabolic rate, independent of 'stress.' *Sci. Rep.* **8**, 13020.

Kelm, D. H., Simon, R., Kuhlow, D., Voigt, C. C. and Ristow, M. (2011). High activity enables life on a high-sugar diet: blood glucose regulation in nectar-feeding bats. *Proc. R. Soc. London B* **278**, 3490–3496.

Kristal, B. S. and Yu, B. P. (1992). An emerging hypothesis: synergistic induction of aging by free radicals and Maillard reactions. *J. Gerontol.* **47**, B107–B114.

Marton, A., Vágási, C. I., Vincze, O., Bókony, V., Pap, P. L., Pătraş, L., Pénzes, J., Bărbos, L., Fülöp, A., Osváth, G., et al. (2022). Oxidative physiology is weakly associated with pigmentation in birds. *Ecol. Evol.* **12**, 1–12.

Monaghan, P., Metcalfe, N. B. and Torres, R. (2009). Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. *Ecol. Lett.* **12**, 75–92.

Muff, S., Nilsen, E. B., O'Hara, R. B. and Nater, C. R. (2022). Rewriting results sections in the language of evidence. *Trends Ecol. Evol.* **37**, 203–210.

Nishikawa, T., Edelstein, D., Du, X. L., Yamagishi, S., Matsumura, T., Kaneda, Y., Yorek, M. A., Beebe, D., Oates, P. J., Hammes, H.-P., et al. (2000). Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature* **404**, 787–790.

Pamplona, R. and Costantini, D. (2011). Molecular and structural antioxidant defenses against oxidative stress in animals. *Am. J. Physiol. Integr. Comp. Physiol.* **301**, R843–R863.

Peng, X., He, X., Liu, Q., Sun, Y., Liu, H., Zhang, Q., Liang, J., Peng, Z., Liu, Z. and Zhang, L. (2017). Flight is the key to postprandial blood glucose balance in the fruit bats *Eonycteris spelaea* and *Cynopterus sphinx*. *Ecol.*

Evol. **7**, 8804–8811.

Pérez-Rodríguez, L., Romero-Haro, A. A., Sternalski, A., Muriel, J., Mougeot, F., Gil, D. and Alonso-Alvarez, C.
(2015). Measuring oxidative stress: the confounding effect of lipid concentration in measures of lipid
peroxidation. *Physiol. Biochem. Zool.* **88**, 345–351.

Picard, M., Juster, R.-P. and McEwen, B. S. (2014). Mitochondrial allostatic load puts the "gluc" back in
glucocorticoids. *Nat. Rev. Endocrinol.* **10**, 303–310.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team (2015). *nlme: Linear and Nonlinear Mixed Effects*
Models. R package version 3.1-120. http://CRAN.R-project.org/package=nlme.

R Core Team (2021). *R: a language and environment for statistical computing*. R Foundation for Statistical
Computing, Vienna, Austria. http://www.R-project.org/.

Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V. and Montiglio, P.-O. (2010). Personality and
the emergence of the pace-of-life syndrome concept at the population level. *Philos. Trans. R. Soc. London B* **365**,
4051–4063.

Ricklefs, R. E. and Wikelski, M. (2002). The physiology/life-history nexus. *Trends Ecol. Evol.* **17**, 462–468.

Romero, L. M. and Beattie, U. K. (2022). Common myths of glucocorticoid function in ecology and conservation. *J.*
Exp. Zool. A **337**, 7–14.

Satoh, T. (2021). Bird evolution by insulin resistance. *Trends Endocrinol. Metab.* **32**, 803–813.

Sepp, T., Sild, E. and Hõrak, P. (2010). Hematological condition indexes in greenfinches: effects of captivity and
diurnal variation. *Physiol. Biochem. Zool.* **83**, 276–282.

Slocinska, M. B., Almsherqi, Z. A. M., Sluse, F. E., Navet, R. and Deng, Y. (2010). Identification and
characterization of uncoupling protein in heart and muscle mitochondria of canary birds. *J. Bioenerg. Biomembr.*
42, 345–353.

Snow, D., Perrins, C. M. and Cramp, S. (1998). The Complete Birds of the Western Palearctic on CD-ROM.

Stier, A., Massemin, S. and Criscuolo, F. (2014). Chronic mitochondrial uncoupling treatment prevents acute cold-
induced oxidative stress in birds. *J. Comp. Physiol. B* **184**, 1021–1029.

Storchová, L. and Hořák, D. (2018). Life-history characteristics of European birds. *Glob. Ecol. Biogeogr.* **27**, 400–
406.

Sukumaran, J. and Holder, M. T. (2010). DendroPy: a Python library for phylogenetic computing. *Bioinformatics* **26**,
1569–1571.

Sweazea, K. L. (2022). Revisiting glucose regulation in birds – A negative model of diabetes complications. *Comp.*
Biochem. Physiol. B **262**, 110778.

Tomášek, O., Bobek, L., Kralova, T., Adámková, M. and Albrecht, T. (2019). Fuel for the pace of life: baseline
blood glucose concentration co-evolves with life-history traits in songbirds. *Funct. Ecol.* **33**, 239–249.

Tomášek, O., Bobek, L., Kauzálová, T., Kauzál, O., Adámková, M., Horák, K., Kumar, S. A., Manialeu, J. P.,
Munclinger, P., Nana, E. D., et al. (2022). Latitudinal but not elevational variation in blood glucose level is
linked to life history across passerine birds. *Ecol. Lett.* **25**, 2203–2216.

Vágási, C. I., Vincze, O., Pătraş, L., Osváth, G., Marton, A., Bărbos, L., Sol, D. and Pap, P. L. (2016).
Large brained birds suffer less oxidative damage. *J. Evol. Biol.* **29**, 1968–1976.

Vágási, C. I., Vincze, O., Pătra□, L., Osváth, G., Pénzes, J., Haussmann, M. F., Barta, Z. and Pap, P. L. (2019).
Longevity and life history coevolve with oxidative stress in birds. *Funct. Ecol.* **33**, 152–161.

Vágási, C. I., Tóth, Z., Pénzes, J., Pap, P. L., Ouyang, J. Q. and Lendvai, Á. Z. (2020). The Relationship between
Hormones, Glucose, and Oxidative Damage Is Condition and Stress Dependent in a Free-Living Passerine Bird.
Physiol. Biochem. Zool. **93**, 466–476.

Vincze, O., Vágási, C. I., Pénzes, J., Szabó, K., Magonyi, N. M., Czirják, G. Á. and Pap, P. L. (2022). Sexual
dimorphism in immune function and oxidative physiology across birds: The role of sexual selection. *Ecol. Lett.*
25, 958–970.

Wingfield, J. C. and Sapolsky, R. M. (2003). Reproduction and resistance to stress: when and how. *J.*
Neuroendocrinol. **15**, 711–724.

Fig. 1. Phylogenetic relationship between glucose stress reactivity (i.e. ΔG, expressed as % change

between G_0 and G_{30}, where $G_0 = 100\%$) and levels of (a) total antioxidant status, (b) uric acid, (c)

total glutathione and (d) malondialdehyde. Plots (a) and (c) show point sizes proportional to either

log sample size or sample size, respectively (see Methods). Fitted lines were obtained from the

respective full models from the second model set (see Table S2).

Table 1. The relationship between oxidative and blood glucose parameters in the first and second

sets of phylogenetically controlled models. The first model set analysed the four oxidative

parameters in separate models as a function of body mass and glucose as continuous predictors,

while the second model set as a function of body mass, fecundity, correlated oxidative parameters

and glucose as predictors. Values of oxidative parameters are best linear unbiased predictions

(BLUEs, see Methods). Glucose was measured either at baseline ($logG_0$), after 30 minutes

standardised acute stress exposure ($logG_{30}$) or calculated as % change between G_0 and G_{30} ($log\Delta G$).

Full models with all the predictors together with model parameters for the first and second models

sets are presented in Table S1 and S2, respectively. *P*-values of associations with moderate

evidence (i.e. $0.01 \leq P \leq 0.05$) are marked in bold, while those of weak evidence (i.e. $0.05 \leq P \leq$

0.1) are marked in italics.

response	predictor	first model set				second model set		
oxidative parameters	glucose	β (SE)	t	P		β (SE)	t	P
total antioxidant status (TAS)	$logG_0$	–0.05 (0.06)	0.82	0.414		–0.05 (0.06)	0.84	0.403
	$logG_{30}$	–0.02 (0.03)	0.66	0.513		–0.01 (0.03)	0.16	0.871
	$log\Delta G$	0.01 (0.01)	0.06	0.956		0.01 (0.01)	0.58	0.563
uric acid (UA)	$logG_0$	–0.02 (0.31)	0.07	0.943		0.28 (0.20)	1.43	0.161
	$logG_{30}$	–0.22 (0.15)	1.43	0.159		–0.10 (0.09)	1.10	0.276
	$log\Delta G$	–0.01 (0.01)	1.49	0.143		–0.01 (0.01)	2.15	**0.037**
total glutathione (tGSH)	$logG_0$	–0.01 (0.39)	0.03	0.979		0.02 (0.34)	0.07	0.943
	$logG_{30}$	–0.34 (0.18)	1.85	*0.071*		–0.27 (0.15)	1.85	*0.071*
	$log\Delta G$	–0.01 (0.01)	1.93	*0.060*		–0.01 (0.01)	2.04	**0.048**
malondialdehyde (MDA)	$logG_0$	–0.33 (0.19)	1.78	*0.081*		–0.18 (0.12)	1.44	0.156
	$logG_{30}$	–0.12 (0.09)	1.25	0.219		0.03 (0.06)	0.52	0.609
	$log\Delta G$	–0.01 (0.01)	0.24	0.814		0.01 (0.01)	1.42	0.163