Gas extraction under intertidal mudflats is associated with declines in sediment grain size and minor

changes in macrozoobenthic community composition

Paula de la Barra*[1], Geert Aarts[1,2], Allert Bijleveld[1]

[1] NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems

[2] Wageningen Marine Research, Wageningen University and Research

*delabarrapaula@gmail.com

Abstract

1-   In intertidal environments, land subsidence may change the local flooding regime and sediment

composition, two main drivers of the macrozoobenthic community structure. In the Dutch Wadden

Sea, a UNESCO world heritage site, gas extraction has resulted in an average subsidence of up to 2

mm y^{-1} of intertidal mudflats. These mudflats support a highly productive macrozoobenthic

community, which offers important resources for birds and fishes. To what extent land subsidence

due to gas extraction affects sediment and macrozoobenthos remains unknown and increasingly

important given sea level rise.

2-   Taking advantage of a monitoring program, we evaluated the effect of anthropogenically caused

land subsidence on sediment composition and intertidal macrozoobenthos. Nearly 4600 points were

sampled yearly (2008-2020) across the Dutch Wadden Sea, allowing us to compare sediment

composition and macrozoobenthos biomass within and outside the subsidence area while

controlling for the main drivers of these variables. We also compared population trends within and

outside the subsidence area for 31 species with different habitat use in terms of depth and sediment

composition.

3-   Sediment median grain size decreased in the subsided area at 1 µm y^{-1} while on average remaining

24   constant in other intertidal mudflats. Mud fraction was 3% higher within the subsided area

25   throughout the studied period. This had no effect on the total biomass of macrozoobenthos. The

26   biomass of species that use deeper areas increased within the subsidence area compared to outside,

27   and the opposite was true for species using shallower places, but comparable patterns were also

28   found in an area not affected by subsidence.

29   4-   Changes in median grain size could be happening, and minor changes in macrozoobenthic

30   community composition. For a successful implementation of the "hand on the tap" principle in the

31   Wadden Sea, it is necessary to define beforehand the relevant variables that represent the natural

32   values, implement proper monitoring, and define threshold values above which effects are not

33   acceptable. We propose median grain size, mud fraction and macrozoobenthic biomass as good

34   measures of the natural values of the Wadden Sea, and the methods used here as a way for

35   identifying anthropogenic effects on them.

Keywords: anthropogenic impact, benthic invertebrates, inla, land subsidence, macrozoobenthos, sediment

composition, spde, Wadden Sea

Introduction

The extraction of underground oil or gas modifies the structure of subterranean layers and can lead to land

subsidence with consequences for the environment at different scales (Milliman & Haq 1996, Ko & Day

2004). As a consequence, habitat availability for plants and animals is also affected, and local ecosystems can

be transformed (e.g. Li et al. 2019, Turner & Mo 2021). When land subsidence occurs in intertidal areas -

regions that are submerged during high tide and emerge during low tide- it alters the local flooding regime. If

subsidence rates are low relative to the sediment supply by water, subsidence can be partially or completely

compensated by increased sedimentation rates (Wang et al. 2018). In mudflats, the equilibrium mud content

of the sediment primarily depends on sediment deposition fluxes (Colina Alonso et al. 2022), and therefore

subsidence might affect the mud content of the affected area. At high subsidence rates, however, intertidal

areas may almost permanently inundate. This could be aggravated by increased erosion, as deeper areas

permit stronger waves and shear stress (Wang et al. 2018). Both sedimentation and erosion, together with

sediment supply, will ultimately determine the sediment composition of a particular area in a mudflat.

Because subsidence changes depth, which partially determines sedimentation and erosion regimes,

subsidence may ultimately change local sediment composition. Additionally, sediment composition can also

be affected by other factors such as anthropic activities and (plant or animal) ecosystem engineers (Eriksson

et al. 2010, van Katwijk et al. 2010, Donadi et al. 2015b).

The flooding regime and the sediment composition of the seafloor are key drivers of intertidal

macrozoobenthos (Van Colen 2018) -invertebrates larger than 1 mm that live in close association with the

substrate. Areas more frequently inundated will experience different environmental conditions (Beukema

1976), different predation pressure (e.g. Sanchez-Salazar et al. 1987) and may have different access to food

(e.g. filter feeders). Sediment size composition determines the amount of organic matter in the seafloor (e.g.

Yang et al. 2021), which influences the stability of the seafloor (Bartzke et al. 2013; but see Defew et al.

2002), and in turn the amount of food for macroinvertebrates (Christianen et al. 2017), as well as their

behaviour (e.g. Wiesebron et al. 2021).

The Wadden Sea, the largest cohesive intertidal area in the world and a UNESCO world heritage site, is

subject to gas extraction, which has resulted in land subsidence (Fokker et al. 2018). Macroinvertebrates of

the Wadden Sea are an important part of the food web as a feeding resource for birds (e.g. Zwarts & Wanink

1993, Beukema & Dekker 2019) and fishes (Poiesz et al. 2020). Because sediment composition and flooding

regime, two of the key factors that determine the occurrence and abundance of macrozoobenthic species,

could be affected by land subsidence, cascading changes to the macrozoobenthic community are likely.

Gas extraction in the Dutch Wadden Sea is managed following the "hand on the tap" principle (NAM B.V

2007), meaning that if an effect on the ecosystem is detected, the gas extraction should be halted. Because

of this, subsidence and its consequences are monitored through a joint effort of governmental agencies

(Rijkswaterstaat), research institutions (NIOZ) and private parties (the Dutch Petroleum Company, NAM, for

its acronym in Dutch). Deep subsidence occurs at a rate of ~2 mm y^{-1} (NAM B.V 2021). However, changes in

average seafloor elevation have not been detected yet (van der Vegt 2022) as subsidence has been so far

compensated by increased sedimentation (Wang et al. 2018). So far, studies about the impact of gas

extraction on mudflats have mostly focused on the risk of intertidal area loss (e.g. Thienen-Visser et al. 2015,

Wang et al. 2018, Fokker et al. 2018, van der Vegt 2022) but changes in sediment and community

composition have received less attention (but see Beukema 2002).

Detecting and measuring large-scale human impacts on ecosystems, such as those of land subsidence, is one

of the big challenges in applied ecology. Ideally, such effects are examined using a before- after control-

impact assessment (BACI, Green 1979). The approach has been successfully employed in many impact

assessment studies (e.g. Sansom et al. 2016, Smokorowski & Randall 2017), however, there are shortcomings

(e.g. Underwood 1991, Smokorowski & Randall 2017). First, the difficulty in selecting and finding appropriate

and sufficient locations of control sites (Schleicher et al. 2020). Second, and related to the first problem, the

observed variability between sites may be the result of spatially varying environmental variables, which

confounds our ability to attribute observed changes to the anthropogenic impacts of interest (Hewitt et al.

2001). Last, the need for an impact assessment study may only be acknowledged when human activities are

already ongoing, and therefore it is impossible to register the unaffected "before" conditions. Large-scale

monitoring systems, when properly designed (Field et al. 2007), could overcome some of these

shortcomings: sampling locations can be used for comparison if environmental variance can be controlled,

and in this way the problem of confounding can be heavily reduced (discussed by Underwood 1993). This

means we can detect differences between impact and control sites while accounting for differences in

environmental conditions between sites.

The purpose of this study was to investigate potential effects of land subsidence caused by gas extraction on

mudflat's sediment composition and macrozoobenthic communities. To achieve this, we took advantage of

thirteen years (2008-2020) of spatially comprehensive monitoring data of sediment composition and benthic

intertidal macroinvertebrates. The spatial and temporal extent of these data allowed us to use it as an

alternative to a BACI design. The rationale is that using the whole Dutch Wadden Sea as a reference area,

and quantifying the effects of physical and biological conditions, we can control for the main processes

driving sediment composition and macroinvertebrates biomass. In this way, we can detect differences

between 'within' and 'outside' the subsidence area "over and above" (i.e. while controlling for) these

environmental variables. Because land subsidence will change the local sedimentation regime, first, we

asked whether sediment composition (median grain size and mud fraction) was different within compared to

outside the area affected by land subsidence due to gas extraction. Second, as depth and sediment

composition are two of the main drivers of the benthic community composition, we then asked whether the

abundance of macrozoobenthos was also different within compared to outside the area affected by land

subsidence. As total biomass encompasses many different species that favour different abiotic conditions,

biomass might increase or decrease in the subsidence area. To clarify how the abundance of particular

species might have changed, we analysed if the ratio of different species' biomass within and outside the

subsidence area changed in time. We related this ratio to the species' favoured abiotic condition to identify

species that may be benefitted or hindered by land subsidence. Additionally, because the area affected by

subsidence is much smaller than the rest of the Wadden Sea, we replicated these analysis using areas un-

affected by land subsidence to better understand how the natural variability of the Wadden Sea could affect

our results.

Materials and Methods

Study area

The Wadden Sea extends for 8000 km^2 from the north of the Netherlands to the east of Denmark (53° N Lat,

Fig 1). It has a bidaily mesotidal regime with a range of 1.4 to 3.4 m across the system (Postma 1982). The

soil is composed by sand and mud (Oost 1995). Its macrozoobenthic community is dominated by bivalves

and polychaetes, with *Cerastoderma edule, Mya arenaria* and *Arenicola marina* being the most abundant

species it terms of biomass (Compton et al. 2013). Gas extraction started in the area of Ameland in 1986

(Fokker et al. 2018). Since then, measurements of land subsidence have been taken by the NAM through

various methods: leveling (comparing elevation against the national height reference network, Fokker et al.

2018), every 4 years since 1986, surveying with GPS, once a year since 2006, and interferometric synthetic

aperture radar measurements (Insar, on land only) once a year since 1997 (NAM B.V 2021). These measures

were modeled by NAM to determine the contours and depth of deep land subsidence. For this study we use

the contours of 1 cm of deep subsidence registered since 2006 (Fig. 1) (NAM B.V 2021).

Fig 1: Synoptic intertidal benthic survey (SIBES) sampling stations (each point is a station) and subsidence

contour in red (1 cm of subsidence between 2006 and 2020). **A)** Average (2008-2021) median grain size, **B)**

average mud fraction, and **C)** average total macroinvertebrate biomass (measured as ash-free dry mass) at

each sampling station.

Survey

Between 2008 and 2020 the macrozoobenthic community and sediment composition of the Wadden Sea

were sampled for the synoptic intertidal benthic survey (SIBES). Every year, between June and October, we

aimed to visit 4511 stations (Fig. 1) of which 78% were located on a regular (500 m) grid design and the rest

were randomly distributed along the grid (Bijleveld et al. 2012). Due to logistic constraints, each year

between 1441 and 4209 stations were effectively visited (see distribution of samples per station in ESM 1).

528 stations were within the contour of subsidence from which we obtained 3866 samples during the entire

studied period. Stations are visited either by foot or using a small rubber boat, depending on tidal and

sediment conditions.

Sediment samples were taken with a 33 mm diameter core to a depth of 4 cm and conserved at -20° C. In

the laboratory, samples were freeze-dried for 96 h, homogenized with mortar and pestle, weighted and put

in 13 ml auto-sampler tubes with degassed reverse osmosis water. Samples were not treated to remove

calcium or organic material. Sediment size composition was characterized using a particle size analyzer

(Coulter LS 13 320) that measures grain sizes between 0.04 and 2000 μm in 126 size classes through laser

diffraction and polarization intensity differential scattering technology. For each sample, the median grain

size and mud fraction (volume percentage of particles smaller than 63 μm) were obtained.

For the benthic community, a core sample of 0.017 m^2 was taken to a depth of ~25 cm and sieved through a

1 mm round mesh. All animals retained were fixed in formalin (4% solution) except for bivalves larger than

~1 cm that were frozen. In the lab, individuals were identified to the finest taxonomic level possible, and ash

free dry mass (afdm) was measured to 0.1 mg. For a more complete description of the methods see

Compton et al. (2013).

Covariates

We identified 10 environmental variables known to be of importance for the distribution of sediment and/or

macroinvertebrates (Table 1). Elevation and current shear stress determine the amount of energy in a

particular place and therefore are associated with sediment composition. Elevation is also an important

driver of macroinvertebrates distribution (e.g. Kraan et al. 2010, de Jong et al. 2015). Strong shear stress and

waves may dislodge benthic invertebrates, so we also expect these variables to affect their total biomass

(e.g. Anta et al. 2013, Malakauskas et al. 2013, but see Donadi et al. 2015a). We used elevation data from

bathymetric maps of the Dutch Wadden Sea that are updated every five years and made available by

Rijkswaterstaat. The distribution of mean bottom shear stress due to currents in the Dutch Wadden Sea was

obtained from Folmer et al. (2016) who calculated it using the General Estuarine Transport Model (Burchard

& Bolding 2002).

In soft sediment systems, seafloor topographic features are correlated with different sediment composition

(e.g. Trentesaux et al. 1994, Mestdagh et al. 2020). Different benthic communities are also associated with

different topographic features (e.g. Jansen et al. 2018, Mestdagh et al. 2020). For this reason, we included

slope, topographic position index and topographic ruggedness as covariates to the models for median grain

size, mud fraction and biomass. We derived this information from the bathymetry. Slope was calculated

using function 'slope' from R package starsExtra (Dorman 2021), topographic position index using function

'wbt_relative_topographic_position' from R package whitebox (Qiusheng 2019) and a kernel of 79 m.

Ruggedness was calculated through function 'wbt_ruggedness_index' also from R package whitebox

(Qiusheng 2019).

Sediment composition strongly influences the benthic community (e.g. Kraan et al. 2010, de Jong et al.

2015), therefore, median grain size, mud fraction and the standard deviation of sediment size were included

in the biomass model. Day of the year was also included in the biomass model to account for periods

dominated by growth or mortality.

The lugworm, *Arenicola marina*, the blue mussel *Mytilus edulis* and the Pacific oyster *Crassostrea gigas* are

important ecosystem engineers in the Wadden Sea (Volkenborn et al. 2007, Donadi et al. 2015b). *A. marina*

removes the sediment resuspending the smaller particles. Oysters and mussels, when filter-feeding, catch

the smaller particles suspended in the water column and fix them in the sediment when excreting. The

biomass of *A. marina* and the presence of a *M. edulis-C. gigas* bank were used as covariates in the median

grain size and mud fraction models. However, because the effect of *A. marina* over the sediment is small

(~13%) and only surficial (Volkenborn et al. 2007), we also ran additional versions of the models not

including this covariate (ESM 2). *A. marina* biomass was determined through SIBES samples as described in

the 'survey' section. Mussel and oyster banks in the Wadden Sea are mapped every year (Troost et al. 2022).

Because the effects of such bivalve banks on sediment extends beyond the bank itself (Kröncke 1996), we

added a 300 m buffer around the bivalve beds contours and classified samples within these contours as

"within bivalve bank".

Finally, the sampling method (by foot or by rubber boat) was also considered a potential source of variation

for median grain size and mud fraction. Samples are taken by boat when the mudflat is inundated and they

are taken by foot only when the mudflat is exposed. This difference between inundation conditions might

affect the composition of top layer of the sediment, so we controlled for this by considering the sampling

method as a covariate.

Table 1: Alphabetically ordered variables that were included in the models fitted to sediment median grain

size, mud fraction, and total biomass data. Source of the data, year of measurements, frequency

distributions of the data and in which model these variables were included is shown. The colors in the

frequency plots indicate outside (blue) and inside (red) the subsidence area.

Variable name	Model used	Source	Years measured	Histogram
Arenicola marina biomass	Sediment median grain size Mud fraction	This study	2008-2020	

Bivalve bank presence	Sediment median grain size Mud fraction	https://shiny.wur.nl/Schelpdiermonitor_Banken/	2008-2020	 No bank Bank
Current shear stress	Sediment median grain size Mud fraction Total biomass	Folmer et al. (2016)	2009-2011	 0.00 0.75 1.50 Shear stress
Day of the year	Total biomass	-	-	 175 240 300 Day
Elevation	Sediment median grain size Mud fraction Total biomass	From Rijkswaterstaat bathymetry	2005, 2013, 2017	 -300 -100 100 Elevation (cm)
Mud fraction	-	This study	2008-2020	 0 35 70 Mud %
Sampling method	Sediment median grain size Mud fraction	This study	2008-2020	 By boat By foot
Sediment median grain size	Total biomass	This study	2008-2020	 0 150 300 Sediment mgs (µm)
Sediment size's standard deviation	Total biomass	This study	2008-2020	 50 200 350 Sediment sd (µm)
Slope	Sediment median grain size Mud fraction Total biomass	Derived from "Elevation"	2005, 2013, 2017	 0 35 75 Slope (°)

Topographic position index	Sediment median grain size Mud fraction Total biomass	Derived from "Elevation"	2005, 2013, 2017	TPI
Total biomass	-	This study	2008-2020	Biomass (g m^{-2})
Year	Sediment median grain size Mud fraction Total species biomass	-	-	Year

Spatial modelling

We modeled three response variables: sediment median grain size (µm), mud fraction (%) and total

macroinvertebrate biomass (afdm, g m^{-2}), each using an appropriate error distribution. We used a Gaussian

distribution for median grain size and a beta distribution with a logit link function for mud fraction (Douma &

Weedon 2019). As all the samples contained detectable biomass, a Gamma distribution with a log link was

used for total biomass (Zuur et al. 2019).

Covariates described in the previous section and Table 1 were included in the models as fixed-effects. To

avoid numerical estimation problems, all covariates were standardized by subtracting their mean and

dividing by their standard deviation. Collinearity between variables was assessed by computing variance

inflation factors (VIF) with function 'vif' of car R package (Fox et al. 2011). We sequentially dropped the

covariates with the highest VIF until all presented VIF < 2 (Zuur et al. 2010). In this way, topographic

ruggedness was dropped from the models. We tested if the relationship between the link function and each

covariate was linear by running generalized linear mixed models and visually inspecting the plots of

Pearson's residuals against each predictor variable. If the residuals showed a residual trend along the

predictor variable, this indicated that the relationship between the link function and the predictor was non-

linear, and thus, we used a smoother for said covariate in the final model (Zuur et al. 2019).

Subsidence was included in the model as a dichotomous variable: each observation was classified as within

or outside the subsidence area. We did this because 38% of our observations within the subsidence area fell

inside the 1 cm contour, and therefore we did not have enough information to analyze its effect as

continuous. We assessed whether the effect of subsidence changed in time by adding an interaction term

with 'year'. Priors for all fixed effects were set to a mean of zero and a precision of 1. This high prior weight

around zero results in the need of more data support for the posterior of a parameter to be different from

zero, which makes our results more conservative.

To account for spatial autocorrelation between observations that cannot be explained by the available

covariates, a stochastic partial differential equation model (SPDE, Lindgren et al. 2011) was used. In this

approach, spatial pseudoreplication is dealt with by adding a spatially correlated random intercept that

follows the distribution of a Gaussian field with mean zero and the correlation between two locations is

given by a Matérn function (Zuur et al. 2019). To estimate the random intercepts at different points, a mesh

of triangles is constructed for the study area and used to estimate the spatially and temporally structured

random component (SPDE specifications in ESM 3). To prevent the SPDE from absorbing too much of the

variability of the model, penalized complexity priors were added. These priors were set following Fuglstad et

al. (2019) and Zuur et al. (2019). To account for temporal autocorrelation from different survey years, we

assessed three temporal correlation structures for the spatial random field: no correlation, "AR1" and

"replicated" (Zuur et al. 2019). We ran the full models using each of the temporal correlation structures and

kept the one with lower DIC (see ESM 3).

The models were validated by inspecting their Pearson's residuals and checking for homogeneity of variance

along the fitted values, and no residual patterns in the residuals when plotted against covariates (Zuur et al.

2019). Residual spatial autocorrelation was assessed by plotting residuals along x and y coordinates (Zuur et

al. 2019).

Our main objective with this study was to test if there was a difference in the response variables depending

on subsidence, therefore we focused on hypothesis testing rather than on model selection (Fieberg 2022). In

Bayesian statistics the uncertainty around any parameter can be estimated (Fieberg 2022) so we took

advantage of this trait to estimate 95% confidence intervals around the parameter of each covariate,

including subsidence and its interaction with time. If zero lies within the confidence interval of a parameter,

it means we cannot distinguish the effect of that variable from zero with 95% confidence. Therefore, we

considered that subsidence had an effect on sediment or biomass if zero was outside the range of the

confidence intervals for the subsidence or interaction term (Fieberg 2022).

We ran additional analyses to assess if differences similar to those found in the original models arouse by

chance. In these analysis we compared a group of samples not affected by subsidence to the rest of the

Wadden Sea. We called these groups of samples "fake subsidence areas". Because each "fake subsidence

area" should be unique to avoid pseudoreplication, the number of potential areas was limited. We selected

five "fake subsidence areas", each centered in a different tidal basin of the Wadden Sea and with similar

sample sizes as the real subsidence area (see details in ESM 4). We ran models for median grain size, mud

fraction and total biomass with the same specifications as the original ones, but using the "fake subsidence"

area instead of the real subsidence area, and using the rest of the Wadden Sea except for the real

subsidence area as the "no subsidence area". If the effect of "fake subsidence" and the interaction between

"fake subsidence" and year is not different from zero, this would support the differences originally found.

Species-specific effects of land subsidence

Total biomass is an aggregate of different species which might be differently affected by subsidence due to

differences in natural history and ecology. To analyze if particular species are driving the observed biomass

patterns, we identified the more abundant species in terms of biomass within the subsidence area, and

plotted their contribution to total biomass within and outside the subsidence area. These species are

Arenicola marina, Cerastoderma edule, Lanice conchilega, Mya arenaria and *Mytilus edulis* and together they

represent more than 90% of total biomass within the subsidence area each year.

To understand how species with different habitat use are affected by subsidence, we compared population

trends within and outside the subsidence area for a larger group of species (species selection methods in

ESM 5). To classify species' habitat use, for each species, we extracted the median of the 'median grain size'

and the median of the elevation range in which they occurred. Hereafter we call these median values

'sediment use' and 'elevation use'.

To detect differences between population trends, we calculated the ratio between the average biomass

within and outside the subsidence area for each species and year. We used a linear mixed model with

log('biomass ratio') as response variable, 'elevation use' and 'year' as continuous fixed effects, and 'species'

as random slope and random intercept. We did not include 'sediment use' in the model to avoid collinearity

with 'elevation use' (Pearson's correlation: -0.72, *P* < 0.001). We applied a likelihood ratio test to select the

best random structure and AIC together with 95% parameter confidence interval to select the best model

relative to fixed effects (Zuur 2009). As an additional control, we ran the same analysis, but using the "fake

subsidence areas" instead of the subsidence area, in the same way as described for the previous section (see

details in ESM 4). If the log biomass ratio between the "fake subsidence" area and the rest of the Wadden

Sea remains constant between years, this would support the validity of the differences originally found.

Results

Median grain size

The average median grain size of all samples was 154 ± 41 µm (mean ± sd, Fig 1A). The mixed-effects model

fitted to median grain size revealed a positive effect of current shear stress and topographic position index,

and a negative effect of the presence of bivalve banks (Fig 2A). The local elevation and biomass of *Arenicola*

marina had non-linear effects on median grain size (Fig 2A, ESM 6). The model revealed a significant effect of

the interaction between the subsidence term and the year term on median grain size (Fig 2A). This meant

that while there was no evidence for a positive or negative trend in sediment median grain throughout the

Wadden Sea, within the subsidence area, median grain size has been decreasing (Fig 2B). Taking *A. marina*

out of the model did not qualitatively change the results (ESM 2). At the end of the study period, the model

estimated that median grain size within the subsidence area was on average 16 µm smaller than outside (Fig

2B). The "fake subsidence" analysis showed that comparing areas of similar size to the rest of the Wadden

Sea, we do not find differences in median grain size (ESM 4).

Fig 2: Results for the sediment median grain size model. **A.** Parameter estimates for covariates used in the

model (mean ± 95% CI). Covariates for which a smoother was applied have more than one parameter. Points

and lines in black show effects that are different from zero with 95% confidence. **B.** Predictions for median

grain size (mean and 95% CI), within and outside the subsidence area throughout the study period.

Mud fraction

On average, samples contained 0.13 ± 0.13 fraction of mud (mean ± sd, Fig 1B). Mud fraction was negatively

affected by most of the covariates assessed, but positively affected by bivalve bank presence (Fig 3A). We

found a positive effect of subsidence on the mud fraction (Fig 3A) that remained constant in time (Fig 3B).

Taking *A. marina* out of the model did not qualitatively change our results (ESM 2). On average, the mud

fraction within the subsidence area was 0.03 (3%) higher than in the rest of the Wadden Sea. The "fake

subsidence" analysis showed that comparing areas of similar size to the rest of the Wadden Sea, we do not

find differences in mud fraction comparable to those found between the subsidence area and the rest of the

Wadden Sea (ESM 4).

Fig 3: Results for the sediment mud fraction model. **A.** Parameter estimates for covariates used in the model

(mean ± 95% CI). Points and lines in black show effects that are different from zero with 95% confidence. **B.**

Predictions for mud fraction (mean and 95% CI), within and outside the subsidence area throughout the

study period.

Biomass

We found that total biomass was on average 24.5 ± 45.3 g m^{-2} (mean ± sd, Fig 1C). Biomass was negatively

affected by topographic position index current shear stress and slope (Fig 4A). Elevation, day of the year,

median gran size and the standard deviation of sediment grain size had non-linear effects on biomass (Fig

4A, ESM 6). We did not find an effect of subsidence on total biomass of the macroinvertebrate community

(Fig 4A,B). The "fake subsidence" analysis showed differences in the total biomass of two areas compared to

the rest of the Wadden Sea (ESM 4).

Fig 4: Results for the total biomass model. **A.** Parameter estimates for covariates used in the model (mean ±

95% CI). Covariates for which a smoother was applied have more than one parameter. Points and lines in

black show effects that are different from zero with 95% confidence. **B.** Predictions for biomass (mean and

95% CI), within and outside the subsidence area throughout the study period.

Species-specific effects of land subsidence

Fig 5: Species biomass within and outside the subsidence area. The portrayed species are the ones with

higher contribution to biomass within the subsidence area.

The most abundant species (in terms of biomass) within the subsidence area were *Arenicola marina*,

Cerstoderma edule, *Lanice conchilega*, *Mya arenaria* and *Mytilus edulis*. These species also represented an

important fraction of biomass outside the subsidence area (Fig 5), with other species only contributing up to

5 ± 1% of total biomass. From 2008 to 2019, *C. edule* was the species with highest average biomass in both

areas. *C. edule* represented the 56% of total biomass within the subsidence area in 2008, but only the 40%

outside. In 2020, *C. edule* was surpassed by *A. marina* in both areas, and by *M. arenaria* outside the

subsidence area (Fig 5).

With the mixed-effects model, we found differences in population trends within and outside the subsidence

area, and these differences depended on the 'elevation use' of species (Fig 6). For species that use lower

areas, populations have increased in biomass within relative to outside the subsidence area. On the contrary,

for species that use higher areas, populations have been slightly decreasing within relative to outside the

subsidence area (Fig 6). However, when running the "fake subsidence" analysis, we also found differences in

population trends when comparing "fake subsidence" areas with the rest of the Wadden Sea (ESM 4). This

means that the differences in species composition found between the subsidence area and the rest of the

Wadden Sea might be are a reflection of the natural variability of the ecosystem.

Fig 6: Results for the population trends model. **A.** Parameter estimates for covariates (mean ± 95% CI). Points

and lines in black show effects that are different from zero with 95% confidence. **B.** Mean and 95%

confidence intervals of the ratio of biomass within and outside the subsidence area throughout the study

period.

Discussion

We found differences in sediment composition within the land subsidence area relative to outside. After

accounting for confounding environmental variables, median grain size within the subsidence area

decreased by 10% during the study period and the mud fraction was 3 percentage points higher throughout.

These changes do not seem to significantly affect the total biomass of the macroinvertebrates living there.

We detected a change in species composition with an increase in the biomass of species that use deeper

habitat and a decrease of species that use a shallower habitat. However, we could not discard these changes

being part of the greater variability of the Wadden Sea.

Subsidence potentially occurs in an area with unique characteristics within the Wadden Sea, and any change

in the subsidence area could be caused by other natural or anthropogenic processes there. For example, the

land subsidence area is close to the Lauwerszee, that has been closed-off in 1969, changing the sediment

transport dynamics of the system (Elias et al. 2012), and one of the inlets within the subsidence area moves

in a cyclic way, changing and recovering its shape every 20-40 years (Oost 1995, Elias et al. 2012). The

potential uniqueness of the area raises the issue of a lack of a proper control in this study, a problem

inherent to nearly all large-scale ecological field studies. To attribute causation, a BACI design is necessary

(Green 1979), however, subsidence has been going on since the eighties, long before we started collecting

the data. For this reason, using the initial conditions (i.e. data from 2008) as the control would be incorrect.

Median grain size, mud fraction and biomass are, nonetheless, not randomly distributed in space, rather

their distribution results from environmental and biological processes that we aimed to capture in the

statistical model with the covariates (like elevation and current shear stress). Because the range of all the

covariates within the subsidence area fall within the range of the whole Wadden Sea's (see histograms in

Table 1), we can thus attempt to detect differences "over and above" environmental differences between

the subsidence area and the rest of the Wadden Sea assuming that the underlying processes (mechanisms)

are similar. On top of this, the addition of a spatial latent field with a temporal structure allows us to at least

partially control for spatially and temporally structured errors caused by historic events (e.g. the closure of

the Lauwerszee) and missing covariates (e.g. suspended sediment loads). Of course, we cannot discard

alternative explanations to the differences we observe, but even after accounting for confounding

environmental variables and residual spatiotemporal patterns, we still observed differences in median grain

size and mud fraction within and outside the subsidence area. Supporting these results, we could not detect

differences between any other particular area and the rest of the Wadden Sea, showing that the observed

differences are likely not due to the natural variability of the Wadden Sea.

Studies on effects of land subsidence in coastal systems have so far mainly focused on the direct effects of

surface subsidence: loss of elevation and consequent land loss (e.g. Kolker et al. 2011, Al Mukaimi et al.

2018), and its relationship with the sediment accreting capacities of the environment (Reed 2002, Al

Mukaimi et al. 2018). Most of them are carried out on salt marsh ecosystems. We have not found any other

study about the composition of the sediment deposited in intertidal systems after land subsidence (but see

Molinaroli et al. 2009, Al Mukaimi et al. 2018). In terrestrial systems, the effects of land subsidence on soil

composition have been explored generally by comparing soil composition at different areas of the

subsidence bowl (top, slope, bottom) with control areas (Tripathi et al. 2009, Qing-jun et al. 2009, Guo et al.

2018, Ma et al. 2019, Vishwakarma et al. 2020). The lack of measurements from before an area is impacted

is extensive for this kind of studies. We found only one study that characterized the soil composition before

the subsidence occurred (Dejun et al. 2016) and one studying landscape scale effects through maps made

before the impact (e.g. Quanyuan et al. 2009). This speaks about how difficult it still is to achieve a proper

experimental design to assess large scale anthropic impacts on the environment.

Subsidence and sediment composition

Subsidence has not affected the average bathymetry and morphology of the tidal flats (van der Vegt 2022),

which implies that sedimentation was so far able to keep up with subsidence. Sand entering the Wadden Sea

mainly comes from the erosion of the ebb-tidal deltas which are in turn supplied by coastal nourishments

(Elias et al. 2012). Mud is transported along the Dutch Coastal zone by the residual north-easterly current

(van der Hout et al. 2015) and enters the Wadden Sea through the tidal inlets. Within the subsidence area,

median grain size was decreasing, and the overall mud fraction was smaller throughout the study period.

Modelled deep subsidence is bowl-shaped (Fokker et al. 2018), with deeper parts in the centre and

shallower parts on the edges of the subsidence contours. Sediment tends to be smaller in troughs (e.g.

Mestdagh et al. 2020) so we hypothesize that the bowl-shaped area could therefore promote the

sedimentation of particles smaller-than-expected based on the environmental characteristics, resulting in

finer sediment composition in this area. Furthermore, previous studies have shown that the role of mud in

the sediment infilling of the Wadden Sea has increased in the last century (Colina Alonso et al. 2021). We

hypothesize that the larger sedimentation rates that subsidence promotes, together with the more

prominent role of mud in infilling the Wadden Sea might result in the smaller grain sizes observed in this

study. It is possible then, that we are detecting the effects of subsidence on sediment composition, while

changes in bathymetry or morphology might be too subtle or compensated too fast to be detected.

Mud fraction was slightly higher within the subsidence area in the beginning of the study and remained

stable during it, showing a different pattern than median grain size. This indicates that the decrease in

median grain size is possibly due to an increase of small sand rather than an increase in mud fraction. We

hypothesize that the different patterns observed in mud fraction and median grain size might be the result

of mud and sand deposition being governed by independent processes: the settling velocities of the two

types of particles differs by an order of magnitude, and while suspended sand concentration depends on

local hydrological conditions, suspended mud concentration depends on supply (Colina Alonso et al. 2022).

Furthermore, because subsidence has been going on since the eighties (Fokker et al. 2018), we hypothesize

that these differences could be the result of a difference in speed where the effects on mud fraction have

plateaued, while the sand fraction is still being affected.

Subsidence and macrozoobenthos

We did not find changes in macrozoobenthos total biomass within the subsidence area relative to outside,

but we found that species that use deeper areas are increasing within the subsidence area, relative to

outside. However, we also found a significant interaction between depth use and year in one of the five

"fake subsidence" controls. This means that the relative biomass of species is also changing in other areas of

the Wadden Sea. Therefore, we cannot discard the hypothesis that the differences in species composition

found between the subsidence area and the rest of the Wadden Sea are a reflection of the natural variability

of the ecosystem. Due to the interdependency of elevation and sediment composition, species that use

deeper areas are generally those that can be found in coarse sediment. With this study, we therefore cannot

tell if species are choosing for a specific sediment or for a specific elevation. Changes in the elevation of the

mudflats occur throughout the Wadden Sea, with certain areas getting higher and others lower as a result of

natural and anthropic processes (Elias et al. 2012). This is also true for the subsidence area that has gained

up to ~7 cm per year between 2010-2021 in some areas, and lost a similar amount in others, while its

average elevation remained mostly unaffected (van der Vegt 2022). We hypothesize that changes in local

elevation might transform the suitability of areas for different species resulting in the increase and decrease

of species depending on their depth use within the subsidence area, but also in at least one other ("fake

subsidence") area in the Wadden Sea. Further studies focusing on the changes in functional groups

composition might shed light on the process behind this pattern.

Other sources of subsidence in the Wadden Sea

There are other areas that were or are subject of anthropogenic subsidence in the Wadden Sea (Fokker et al.

2018). Gas extraction at the west of the Wadden Sea (in Zuidwal), took place between 1988 and 2020,

resulting in 69.5 mm of subsidence at the deepest point up until 1997 (Rommel 2004) and less than 0.01 mm

y^{-1} by 2018 (Fokker et al. 2018). Salt solution mining (near Harlingen) started in 2020 and is ongoing. It is

expected to result in 106 cm of subsidence by 2052 in the deepest area (Fokker et al. 2018). There is gas

extraction below Groningen since 1963, and the resulting subsidence is partially extending to the Wadden

Sea (Fokker et al. 2018). Because of the difference in rates, start and finish dates, and the way in which these

events were or are being monitored (Fokker et al. 2018), we do not consider the information available

comparable in order to include it in this study. Additionally, subsidence might not have the same effect in

the different areas. For example, while here we address subsidence in a sediment accommodation-limited

basin, subsidence in Zuidwal occurs in a sediment deposition limited basin (Wang et al. 2018), which means

it's likely that it will not be filled out by sedimentation as easily, and changes in sediment composition will

likely be different. The areas of the Wadden Sea affected by subsidence from Zuidwal and Groningen are

much smaller than the subsidence area studied here (e.g. ~1700 samples within Zuidwal, see Hoeksema et

al. 2004) and therefore, we do not expect them to drive the patterns in average sediment compositions we

observe in the rest of the Wadden Sea.

Management implications

The Wadden Sea as a UNESCO heritage site is a protected area. The gas extraction near Ameland is managed

following the "hand on the tap" principle, meaning that "if the natural values of the Wadden Sea are

compromised", gas extraction must be reduced or even ceased. Intertidal area loss is the most evident

threat posed by subsidence, so concrete methods to measure it and thresholds to prevent it have been

established (6 mm y^{-1} for the saltmarsh area, National Project Decree 2005). Other threats posed by

subsidence to "natural values" would also warrant the adjustment of gas production (National Project

Decree 2005), but in those cases three problems might hinder the effective implementation of the "hand on

the tap" principle. First, the lack of definition of what the Wadden Sea's natural values are. We propose that

the variables measured here (sediment composition, biomass and benthic community structure) can be

considered natural values of the Wadden Sea as they constitute essential components of the ecosystem's

function. Sediment composition and the benthic community might not be the charismatic reason why the

Wadden Sea is protected, but these elements are two of the pillars sustaining the ecosystem as we know it

today. The second problem is the lack of guidelines about what constitutes an effect. Here, we showed that

effects on sediment and community composition might occur, but the lack of measures from before the

beginning of gas extraction precludes us from attributing causality. This does not mean that effects do not

exist, but rather that it is not possible to unambiguously show them. To do proper monitoring of potential

effects, BACI studies are essential. Therefore, we call for the enforcement of properly designed monitoring of

the natural values of the Wadden Sea before any activity starts. In cases such as the one studied here, in

which before measurements cannot be obtained anymore, we propose that the methods used in this study

could be implemented as the standard for identifying potential effects. In cases in which it is

methodologically impossible to attribute causation, and given that the Wadden Sea is a protected area, the

weight of proof should be reversed, and those exploiting the area should prove no effects are in place.

Moreover, by law, the burden of proof is already on the companies exploiting the natural resources showing

that their activities have no effect on the Wadden Sea's natural values (e.g. see Eco Advocacy CLG v An Bord

Pleanala. 2023). The third problem that we identify is the lack of a threshold for these effects that, from an

ecological perspective, would warrant management consequences. Does the $1\ \mu m\ y^{-1}$ decrease in median

grain size warrant management consequences? Could this be considered a threat to the natural values of the

Wadden Sea? If we do a stringent interpretation of the decree, any effect is reason to reduce or stop drilling.

In conclusion, for a successful implementation of the "hand on the tap" principle in a protected area like the

Wadden Sea, it is necessary to clearly define beforehand the relevant variables that represent the natural

values, implement proper monitoring for impact assessment, and define threshold values above which

effects are not acceptable, otherwise the water will continue to run regardless of where the hand is.

Author contribution

Allert Bijleveld and Geert Aarts conceived the idea. Allert Bijleveld, Geert Aarts and Paula de la Barra

designed the methodology. Paula de la Barra analyzed the data; interpretation of results was discussed by all

three authors. Paula de la Barra led the writing of the manuscript. All authors contributed critically to the

drafts and gave final approval for publication.

Acknowledgements

We are very grateful to the current SIBES core-team for collecting and processing the thousands of samples.

In alphabetical order they are Leo Boogert, Thomas de Brabander, Anne Dekinga, Job ten Horn, Loran Kleine

Schaars, Jeroen Kooijman, Anita Koolhaas, Franka Lotze, Simone Miguel, Luc de Monte, Dennis Mosk, Amin

Niamir, Dana Nolte, Dorien Oude Luttikhuis, Bianka Rasch, Reyhane Roohi, Juan Schiaffi, Marten Tacoma,

Eveline van Weerlee and Bas de Wit. We also thank all former and current employees and the many

volunteers and students who have ensured that the SIBES program has continued in recent years. The RV

Navicula was essential for collecting the samples and particularly we thank the current crew Wim Jan Boon,

Klaas Jan Daalder, Bram Fey, Hendrik Jan Lokhorst and Hein de Vries. SIBES is currently financed by the

Nederlandsche Aardolie Maatschappij NAM, Rijkswaterstaat RWS and the Royal NIOZ.

Conflict of Interest

SIBES is currently for approximately one-third financed by the Nederlandsche Aardolie Maatschappij (NAM),

which exploits the area for gas production and generating the land subsidence studied here.

Data availability statement

Data will be available at NIOZ Data Archive System upon publication.

References

Al Mukaimi ME, Dellapenna TM, Williams JR (2018) Enhanced land subsidence in Galveston Bay,
Texas: Interaction between sediment accumulation rates and relative sea level rise. Estuar
Coast Shelf Sci 207:183–193.
Anta J, Peña E, Puertas J, Cea L (2013) A bedload transport equation for the Cerastoderma edule
cockle. J Mar Syst 111–112:189–195.
Bartzke G, Bryan KR, Pilditch CA, Huhn K (2013) On the Stabilizing Influence of Silt On Sand Beds.
J Sediment Res 83:691–703.
Beukema JJ (1976) Biomass and species richness of the macro-benthic animals living on the tidal
flats of the Dutch Wadden Sea. Neth J Sea Res 10:236–261.
Beukema JJ (2002) Expected changes in the benthic fauna of Wadden Sea tidal flats as a result of
sea-level rise or bottom subsidence. J Sea Res 47:25–39.
Beukema JJ, Dekker R (2019) Is food supply for shellfish-eating birds in the western Wadden Sea
affected by the between-species synchrony in year-to-year fluctuations of bivalve population
parameters? Helgol Mar Res 73:8.
Bijleveld AI, van Gils JA, van der Meer J, Dekinga A, Kraan C, van der Veer HW, Piersma T (2012)
Designing a benthic monitoring programme with multiple conflicting objectives: *Sampling*
design for conflicting objectives. Methods Ecol Evol 3:526–536.
Burchard H, Bolding K (2002) GETM - a general estuarine transport model.
Christianen MJA, Middelburg JJ, Holthuijsen SJ, Jouta J, Compton TJ, van der Heide T, Piersma T,
Sinninghe Damsté JS, van der Veer HW, Schouten S, Olff H (2017) Benthic primary
producers are key to sustain the Wadden Sea food web: stable carbon isotope analysis at
landscape scale. Ecology 98:1498–1512.
Colina Alonso A, van Maren DS, Elias EPL, Holthuijsen SJ, Wang ZB (2021) The contribution of
sand and mud to infilling of tidal basins in response to a closure dam. Mar Geol 439:106544.
Colina Alonso A, van Maren DS, Herman PMJ, van Weerdenburg RJA, Huismans Y, Holthuijsen SJ,
Govers LL, Bijleveld AI, Wang ZB (2022) The Existence and Origin of Multiple Equilibria in
Sand-Mud Sediment Beds. Geophys Res Lett 49:e2022GL101141.

Compton TJ, Holthuijsen S, Koolhaas A, Dekinga A, ten Horn J, Smith J, Galama Y, Brugge M, van der Wal D, van der Meer J, van der Veer HW, Piersma T (2013) Distinctly variable mudscapes: Distribution gradients of intertidal macrofauna across the Dutch Wadden Sea. J Sea Res 82:103–116.

Defew EC, Tolhurst TJ, Paterson DM (2002) Site-specific features influence sediment stability of intertidal flats. Hydrol Earth Syst Sci 6:971–982.

Dejun Y, Zhengfu B, Shaogang L (2016) Impact on soil physical qualities by the subsidence of coal mining: a case study in Western China. Environ Earth Sci 75:652.

Donadi S, Eriksson BK, Lettmann KA, Hodapp D, Wolff J-O, Hillebrand H (2015a) The body-size structure of macrobenthos changes predictably along gradients of hydrodynamic stress and organic enrichment. Mar Biol 162:675–685.

Donadi S, van der Heide T, Piersma T, van der Zee EM, Weerman EJ, van de Koppel J, Olff H, Devine C, Hernawan UE, Boers M, Planthof L, Klemens Eriksson B (2015b) Multi-scale habitat modification by coexisting ecosystem engineers drives spatial separation of macrobenthic functional groups. Oikos 124:1502–1510.

Dorman M (2021) StarsExtra: Miscellaneous Functions for Working with "stars" Rasters.

Douma JC, Weedon JT (2019) Analysing continuous proportions in ecology and evolution: A practical introduction to beta and Dirichlet regression. Methods Ecol Evol 10:1412–1430.

Eco Advocacy CLG v An Bord Pleanala. (2023) Judgment of the Court in Case C‑721/21. ECLI:EU:C:2023:477.

Elias EPL, Van der Spek AJ, Wang ZB, De Ronde J (2012) Morphodynamic development and sediment budget of the Dutch Wadden Sea over the last century. Neth J Geosci 91:293–310.

Eriksson BK, van der Heide T, van de Koppel J, Piersma T, van der Veer HW, Olff H (2010) Major Changes in the Ecology of the Wadden Sea: Human Impacts, Ecosystem Engineering and Sediment Dynamics. Ecosystems 13:752–764.

Fieberg J (2022) Statistics for Ecologists: A Frequentist and Bayesian Treatment of Modern Regression Models. An open-source online textbook.

Field SA, O'connor PJ, Tyre AJ, Possingham HP (2007) Making monitoring meaningful. Austral Ecol 32:485–491.

Fokker PA, Leijen FJ van, Orlic B, Marel H van der, Hanssen RF (2018) Subsidence in the Dutch Wadden Sea. Neth J Geosci 97:129–181.

Folmer EO, van Beusekom JEE, Dolch T, Gräwe U, van Katwijk MM, Kolbe K, Philippart CJM (2016) Consensus forecasting of intertidal seagrass habitat in the Wadden Sea. J Appl Ecol 53:1800–1813.

Fox J, Weisberg S, Fox J (2011) An R companion to applied regression, 2nd ed. SAGE Publications, Thousand Oaks, Calif.

Fuglstad G-A, Simpson D, Lindgren F, Rue H (2019) Constructing Priors that Penalize the Complexity of Gaussian Random Fields. J Am Stat Assoc 114:445–452.

Green RH (1979) Sampling Design and Statistical Methods for Environmental Biologists. John Wiley & Sons.

Guo X, Zhao T, Chang W, Xiao C, He Y, Guo X, Zhao T, Chang W, Xiao C, He Y (2018) Evaluating the effect of coal mining subsidence on the agricultural soil quality using principal component analysis. Chil J Agric Res 78:173–182.

Hewitt JE, Thrush SE, Cummings VJ (2001) Assessing Environmental Impacts: Effects of Spatial and Temporal Variability at Likely Impact Scales. Ecol Appl 11:1502–1516.

Hoeksema HJ, Mulder HPJ, Rommel MC, Ronde JG, de Vlas J (2004) Bodemdalingstudie Waddenzee 2004. Vragen en onzekerheden opnieuw beschouwd.

van der Hout CM, Gerkema T, Nauw JJ, Ridderinkhof H (2015) Observations of a narrow zone of high suspended particulate matter (SPM) concentrations along the Dutch coast. Cont Shelf Res 95:27–38.

Jansen J, Hill NA, Dunstan PK, Eléaume MP, Johnson CR (2018) Taxonomic Resolution, Functional Traits, and the Influence of Species Groupings on Mapping Antarctic Seafloor Biodiversity. Front Ecol Evol 6.

de Jong MF, Baptist MJ, Lindeboom HJ, Hoekstra P (2015) Relationships between
macrozoobenthos and habitat characteristics in an intensively used area of the Dutch
coastal zone. ICES J Mar Sci 72:2409–2422.
van Katwijk MM, Bos AR, Hermus DCR, Suykerbuyk W (2010) Sediment modification by seagrass
beds: Muddification and sandification induced by plant cover and environmental conditions.
Estuar Coast Shelf Sci 89:175–181.
Ko J-Y, Day JW (2004) A review of ecological impacts of oil and gas development on coastal
ecosystems in the Mississippi Delta. Ocean Coast Manag 47:597–623.
Kolker AS, Allison MA, Hameed S (2011) An evaluation of subsidence rates and sea-level variability
in the northern Gulf of Mexico. Geophys Res Lett 38.
Kraan C, Aarts G, van der Meer J, Piersma T (2010) The role of environmental variables in
structuring landscape-scale species distributions in seafloor habitats. Ecology 91:1583–
1590.
Kröncke I (1996) Impact of Biodeposition on Macrofaunal Communities in Intertidal Sandflats. Mar
Ecol 17:159–174.
Li C, Yang S, Zha D, Zhang Y, Boer WF de (2019) Waterbird Communities in Subsidence Wetlands
Created by Underground Coal Mining in China: Effects of Multi-Scale Environmental and
Anthropogenic Variables. Environ Conserv 46:67–75.
Lindgren F, Rue H, Lindström J (2011) An explicit link between Gaussian fields and Gaussian
Markov random fields: the stochastic partial differential equation approach. J R Stat Soc Ser
B Stat Methodol 73:423–498.
Ma K, Zhang Y, Ruan M, Guo J, Chai T (2019) Land Subsidence in a Coal Mining Area Reduced
Soil Fertility and Led to Soil Degradation in Arid and Semi-Arid Regions. Int J Environ Res
Public Health 16:3929.
Malakauskas DM, Willson SJ, Wilzbach MA, Som NA (2013) Flow variation and substrate type
affect dislodgement of the freshwater polychaete, Manayunkia speciosa. Freshw Sci
32:862–873.
Mestdagh S, Amiri-Simkooei A, van der Reijden KJ, Koop L, O'Flynn S, Snellen M, Van Sluis C,
Govers LL, Simons DG, Herman PMJ, Olff H, Ysebaert T (2020) Linking the morphology and
ecology of subtidal soft-bottom marine benthic habitats: A novel multiscale approach. Estuar
Coast Shelf Sci 238:106687.
Milliman JD, Haq BU (eds) (1996) Sea-level rise and coastal subsidence: causes, consequences,
and strategies. Kluwer Academic Publishers, Dordrecht; Boston.
Molinaroli E, Guerzoni S, Sarretta A, Masiol M, Pistolato M (2009) Thirty-year changes (1970 to
2000) in bathymetry and sediment texture recorded in the Lagoon of Venice sub-basins,
Italy. Mar Geol 258:115–125.
NAM B.V (2021) Gaswinning vanaf de locaties Moddergat, Lauwersoog en Vierhuizen. Resultaten
uitvoering Meet- en regelcyclus 2020.
NAM B.V (2007) Meet- en regel protocol Winning Waddenzeegas vanaf de locaties Moddergat,
Lauwersoog en Vierhuizen.
National Project Decree (2005) Rijksprojectbesluit Gaswinning onder de Waddenzee vanaf de
locaties Moddergat, Lauwersoog en Vierhuizen. Deel 1.
Oost AP (1995) Dynamics and sedimentary developments of the Dutch Wadden Sea with a special
emphasis on the Frisian Inlet: a study of the barrier islands, ebb-tidal deltas, inlets and
drainage basins. Dissertation, Faculteit Aardwetenschappen
Poiesz SSH, Witte JI, Veer HW van der (2020) Only a few key prey species fuel a temperate coastal
fish food web. Mar Ecol Prog Ser 653:153–166.
Postma H (1982) Hydrography of the Wadden sea: movements and properties of water and
particulate matter final report on "hydrography." A. A. Balkema, Rotterdam.
Qing-jun M, Qi-yan F, Qing-qing W, Lei M, Zhi-yang C (2009) Distribution characteristics of nitrogen
and phosphorus in mining induced subsidence wetland in Panbei coal mine, China. Procedia
Earth Planet Sci 1:1237–1241.
Quanyuan W, Jiewu P, Shanzhong Q, Yiping L, Congcong H, Tingxiang L, Limei H (2009) Impacts
of coal mining subsidence on the surface landscape in Longkou city, Shandong Province of
China. Environ Earth Sci 59:783.

Reed DJ (2002) Sea-level rise and coastal marsh sustainability: geological and ecological factors in
the Mississippi delta plain. Geomorphology 48:233–243.
Rommel MC (2004) Morfologische veranderingen als gevolg van bodemdaling door gaswinning.
Rapp 2004004.
Sanchez-Salazar ME, Griffiths CL, Seed R (1987) The interactive roles of predation and tidal
elevation in structuring populations of the edible cockle, Cerastoderma edule. Estuar Coast
Shelf Sci 25:245–260.
Sansom A, Pearce-Higgins JW, Douglas DJT (2016) Negative impact of wind energy development
on a breeding shorebird assessed with a BACI study design. Ibis 158:541–555.
Schleicher J, Eklund J, D. Barnes M, Geldmann J, Oldekop JA, Jones JPG (2020) Statistical
matching for conservation science. Conserv Biol 34:538–549.
Smokorowski KE, Randall RG (2017) Cautions on using the Before-After-Control-Impact design in
environmental effects monitoring programs. FACETS 2:212–232.
Thienen-Visser V, Breunese JN, Muntendam-Bos AG (2015) Subsidence due to gas production in
the Wadden Sea: How to ensure no harm will be done to nature. In: *49th US Rock*
Mechanics/Geomechanics Symposium. OnePetro
Trentesaux A, Stolk A, Tessier B, Chamley H (1994) Surficial sedimentology of the Middelkerke
Bank (southern North Sea). Mar Geol 121:43–55.
Tripathi N, Singh RS, Singh JS (2009) Impact of post-mining subsidence on nitrogen transformation
in southern tropical dry deciduous forest, India. Environ Res 109:258–266.
Troost K, Asch M van, Ende D van den, Es Y van, Perdon KJ, Pool J van der, Suykerbuyk W,
Zweeden C van, Zwol J van (2022) Schelpdierbestanden in de Nederlandse kustzone,
Waddenzee en zoute deltawateren in 2021.
Turner RE, Mo Y (2021) Salt Marsh Elevation Limit Determined after Subsidence from Hydrologic
Change and Hydrocarbon Extraction. Remote Sens 13:49.
Underwood AJ (1991) Beyond BACI: Experimental designs for detecting human environmental
impacts on temporal variations in natural populations. Mar Freshw Res 42:569–587.
Underwood AJ (1993) The mechanics of spatially replicated sampling programmes to detect
environmental impacts in a variable world. Aust J Ecol 18:99–116.
Van Colen C (2018) The Upper Living Levels: Invertebrate Macrofauna. In: *Mudflat Evology*. Aquatic
Ecology Series, Springer International Publishing
van der Vegt H (2022) Monitoring wadplaatareaal Friesche Zeegat met LiDAR (2010-2021).
Vishwakarma AK, Behera T, Rai R, Sonkar AK, Singh AP, Shrivastva BK (2020) Impact assessment
of coal mining induced subsidence on native soil of South Eastern Coal Fields: India.
Geomech Geophys Geo-Energy Geo-Resour 6:31.
Volkenborn N, Hedtkamp SIC, van Beusekom JEE, Reise K (2007) Effects of bioturbation and
bioirrigation by lugworms (Arenicola marina) on physical and chemical sediment properties
and implications for intertidal habitat succession. Estuar Coast Shelf Sci 74:331–343.
Wang ZB, Elias EPL, van der Spek AJ, Lodder QJ (2018) Sediment budget and morphological
development of the Dutch Wadden Sea: impact of accelerated sea-level rise and subsidence
until 2100. Neth J Geosci 97:183–214.
Wiesebron LE, Steiner N, Morys C, Ysebaert T, Bouma TJ (2021) Sediment Bulk Density Effects on
Benthic Macrofauna Burrowing and Bioturbation Behavior. Front Mar Sci 8:707785.
Yang P, Shu Q, Liu Q, Hu Z, Zhang S, Ma Y (2021) Distribution and factors influencing organic and
inorganic carbon in surface sediments of tidal flats in northern Jiangsu, China. Ecol Indic
126:107633.
Zuur AF (ed) (2009) Mixed Effects Models and Extensions in Ecology with R. Springer, New York,
NY.
Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common statistical
problems. Methods Ecol Evol 1:3–14.
Zuur AF, Ieno EN, Savel☐ev AA, Zuur AF (2019) Beginner's guide to spatial, temporal and spatial-
temporal ecological data analysis with R-INLA. Highland Statistics Ltd, Newburgh, United
Kingdom.

Zwarts L, Wanink JH (1993) How the food supply harvestable by waders in the Wadden Sea
719   depends on the variation in energy density, body weight, biomass, burying depth and
720   behaviour of tidal-flat invertebrates. Neth J Sea Res 31:441–476.
721