Gas extraction under intertidal mudflats is associated with changes in sediment structure and

macrozoobenthic invertebrate communities

Paula de la Barra*[1], Geert Aarts[1,2], Allert Bijleveld[1]

[1] NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems

[2] Wageningen Marine Research, Wageningen University and Research

*delabarrapaula@gmail.com

Abstract

1-   Gas withdrawal from subterranean layers may generate land subsidence with consequences for soil

micro-structure, hydrology, and ultimately local biological communities. In intertidal environments,

land subsidence may change the local flooding regime and sediment composition, two main drivers

of the macrozoobenthic community structure. In the Dutch Wadden Sea, a UNESCO world heritage

site, gas extraction has resulted in an average subsidence of up to 2 mm y^{-1}, of sand dunes, salt

marshes and intertidal mudflats. These mudflats support a highly productive macrozoobenthic

community, which offers important feeding grounds for birds and fishes. To what extent land

subsidence due to gas extraction affects the macrozoobenthos remains unknown and increasingly

important given sea level rise.

2-   Taking advantage of an extensive monitoring program of the macrozoobenthic community, we

evaluated the effect of anthropogenically caused land subsidence on sediment composition and

intertidal macrozoobenthos. Nearly 4600 points were sampled yearly (2008-2020) across the Dutch

Wadden Sea, allowing us to compare sediment composition and macrozoobenthos biomass within

and outside the subsidence area while controlling for the main drivers of these variables. With an

INLA-spde approach we could account for spatial and temporal autocorrelation. We also compared

population trends within and outside subsidence area for 31 species with different habitat use in

terms of depth and sediment composition.

3- We found that median grain size of the sediment has decreased in the subsided area at 1 μm y^{-1}

while on average remaining constant in the rest of the intertidal mudflats. Mud fraction was 3%

higher within the subsided area throughout the studied period. This had no effect on the total

biomass of macrozoobenthos. However, the biomass of species that prefer deeper areas increased

within the subsidence area compared to the area outside the subsidence area, and the opposite was

true for species that prefer shallower places.

4- Policy implications: studies suggest that the present subsidence will not result in the drowning of the

mudflat. However, we show that changes in sediment- and macrozoobenthic community

composition might be happening. We call for definitions and thresholds of effects for a better

management of the Wadden Sea

Keywords: anthropogenic impact, inla, land subsidence, sediment composition, soft bottoms, spde, wadden

sea

Introduction

The extraction of underground oil or gas modifies the structure of subterranean layers and can lead to land

subsidence with consequences for the environment at different scales (Milliman & Haq 1996, Ko & Day

2004). As a consequence, habitat availability for plants and animals is also affected, and local ecosystems are

transformed (e.g. Li et al. 2019, Turner & Mo 2021). When land subsidence occurs in intertidal areas -regions

that are submerged during high tide and emerge during low tide- it alters the local flooding regime. If

subsidence rates are low relative to the sediment supply by water, subsidence can be partially or completely

compensated by sedimentation (Wang et al. 2018). At high subsidence rates, however, intertidal areas may

almost permanently inundate. This could be aggravated by increased erosion, as deeper areas permit

stronger waves and shear stress (Wang et al. 2018). Both sedimentation and erosion, together with sediment

supply, will ultimately determine the sediment composition of a particular area in a mudflat. Because

subsidence changes depth, which partially determines sedimentation and erosion regimes, subsidence may

ultimately change local sediment composition. Nevertheless, sediment composition can also be affected by

other factors such as anthropic activities and ecosystem engineers (Eriksson et al. 2010, Donadi et al. 2015b).

The flooding regime and the sediment composition of the seafloor are key drivers of intertidal

macrozoobenthos (Van Colen 2018) -invertebrates larger than 1 mm that live in close association with the

substrate. Areas more frequently inundated will experience different environmental conditions (Beukema

1976), different predation pressure (e.g. Sanchez-Salazar et al. 1987) and may have different access to food

(e.g. filter feeders). Sediment size composition determines the amount of organic matter in the seafloor (e.g.

Yang et al. 2021), which determines the stability of the seafloor (Bartzke et al. 2013; but see Defew et al.

2002), and in turn the amount of food for macroinvertebrates (Christianen et al. 2017), as well as their

behaviour (e.g. Wiesebron et al. 2021).

The Wadden Sea, the largest cohesive intertidal area in the world and a UNESCO world heritage site, is

subject to gas extraction, which has resulted in land subsidence (Fokker et al. 2018). Macroinvertebrates of

the Wadden Sea are an important part of the food web as a feeding resource for birds (e.g. Beukema &

Dekker 2019) and fishes (Poiesz et al. 2020). Because sediment composition and flooding regime, two of the

key factors that determine the occurrence and abundance of macrozoobenthic species, could be affected by

land subsidence, cascading changes to the macrozoobenthic community are likely.

Gas extraction in the Dutch Wadden Sea is managed following the "hand on the tap" principle (NAM B.V

2007), meaning that if an effect on the ecosystem is detected, the gas extraction should be halted. Because

of this, subsidence and its consequences are closely monitored. Deep subsidence occurs at a rate of ~2 mm

y^{-1} (NAM B.V 2021). However, changes in average seafloor elevation have not been detected yet (van der

Vegt 2022) as subsidence has been so far compensated by increased sedimentation (Wang et al. 2018). So

far, studies about the impact of gas extraction on mudflats have mostly focused on the risk of intertidal area

loss (e.g. Thienen-Visser et al. 2015, Wang et al. 2018, Fokker et al. 2018, van der Vegt 2022) but changes in

sediment and community composition have received less attention (but see Beukema 2002).

To detect and measure human impacts on ecosystems, such as those of land subsidence, is one of the big

challenges in applied ecology. Ideally, such effects are examined using a before- after control-impact

assessment (BACI, Green 1979). The approach has been successfully employed in many impact assessment

studies (e.g. Sansom et al. 2016, Smokorowski & Randall 2017), there are shortcomings (e.g. Underwood

1991, Smokorowski & Randall 2017). First, the selection of the location and number of control sites can be

arbitrary. Second, the observed variability between sites may be the result of spatially varying environmental

variables, which confounds our ability to attribute observed changes to the anthropogenic impacts of

interest (Hewitt et al. 2001). Last, the need for an impact assessment study is only acknowledged when

human activities are already ongoing, and therefore it is impossible to register the unaffected "before"

conditions. Large-scale monitoring systems, when properly designed (Field et al. 2007), could overcome

some of these shortcomings: all sampling locations could act as control sites, and by also accounting for the

effect of all other environmental covariates on the variable of interest, the problem of confounding can be

heavily reduced (discussed by Underwood 1993). This means we can detect differences between impact and

control sites while accounting for differences in environmental conditions between sites.

The purpose of this study was to investigate potential effects of land subsidence caused by gas extraction on

mudflat's sediment composition and macrozoobenthic communities. To achieve this, we took advantage of

thirteen years (2008-2020) of spatially comprehensive monitoring data of sediment composition and benthic

intertidal macroinvertebrates. The spatial and temporal extent of these data allowed us to use it as an

alternative to a BACI design, and to quantify the effect of local physical and biological conditions on

sediment composition (median grain size and mud fraction) and macroinvertebrates biomass. In this way, we

can detect differences between 'within' and 'outside' the subsidence area "over and above" (i.e. while

controlling for) these environmental variables. To control for spatiotemporal autocorrelation that cause

pseudo replication, a spatial-temporal latent field was also included using the INLA-SPDE approach (Lindgren

et al. 2011). Because land subsidence will change the local sedimentation regime, first, we asked whether sediment composition was different within compared to outside the area affected by land subsidence due to gas extraction. Second, as depth and sediment composition are two of the main drivers of the benthic community, we then asked whether the abundance of macrozoobenthos was also different within compared to outside the area affected by land subsidence. As total biomass encompasses many different species that favour different abiotic conditions, biomass might increase or decrease in the subsidence area. To clarify how particular species distributions changed, we analysed trends in the ratio of different species' biomass within and outside the subsidence area. We related this ratio to the species' favoured abiotic condition to identify species that may be benefitted or hindered by land subsidence.

Materials and Methods

Study area

The Wadden Sea extends for 8000 km^2 from the north of the Netherlands to the east of Denmark (53° N Lat, Fig 1). It has a bidaily mesotidal regime with a range of 1.4 to 3.4 m across the system (Postma 1982). The soil is composed by sand and mud (Oost 1995). Its macrozoobenthic community is dominated by bivalves and polychaetes, with *Cerastoderma edule, Mya arenaria* and *Arenicola marina* being the most abundant species it terms of biomass (Compton et al. 2013).

Gas extraction started in the area of Ameland in 1986 (Fokker et al. 2018). Since then, measurements of land subsidence have been taken by the Dutch Petroleum Company (NAM, for its acronym in Dutch) through various methods: leveling, every 4 years since 1986, surveying with GPS, once a year since 2006, and Insar measurements (on land only) once a year since 1997 (NAM B.V 2021). These measures were modeled by NAM to determine the contours of deep land subsidence. For this study we use the contours of 1 cm of deep subsidence registered since 2006 (Fig. 1) (NAM B.V 2021).

Fig 1: SIBES sampling stations and subsidence contour in red (1 cm of subsidence between 2006 and 2020).

A) Average (2008-2021) median grain size, **B)** average mud fraction, and **C)** average total macroinvertebrate

biomass (measured as ash-free dry mass) at each sampling station.

Survey

Between 2008 and 2020 the macrozoobenthic community and sediment composition of the Wadden Sea

were sampled for the synoptic intertidal benthic survey (SIBES). Yearly, the survey consisted of 1441 to 4209

stations of which 54 – 91 % were located on a regular (500 m) grid design and the rest were randomly

distributed along the grid (Bijleveld et al. 2012). Of the total of 4580 sampling stations 528 were within the

contour of subsidence. Every year, between June and October, each station was visited, and a sample of

sediment and the benthic community were taken (for methods see Compton et al. 2013). All animals were

fixed in formalin (4% solution) except for bivalves larger than ~1 cm that were frozen. In the lab, individuals

were identified to the finest taxonomic level possible, and ash free dry mass (afdm) was measured to 0.1 mg.

For a more complete description of the methods see Compton et al. (2013).

Sediment samples were taken with a 33 mm diameter core to a depth of 4 cm and conserved at -20° C. In

the laboratory, samples were freeze-dried for 96 h, homogenized with mortar and pestle, weighted and put

in 13 ml auto-sampler tubes with degassed reverse osmosis water. Samples were not treated to remove

calcium or organic material. Sediment size composition was characterized using a particle size analyzer

(Coulter LS 13 320) that measures grain sizes between 0.04 and 2000 µm in 126 size classes through laser

diffraction and polarization intensity differential scattering technology. For each sample, the median grain

size (mgs) and mud fraction (volume percentage of particles smaller than 63 µm) were obtained.

Covariates

We identified a multitude of environmental variables known to be of importance for the distribution of

sediment and macroinvertebrates (Table 1). Depth, current shear stress, wave forcing all determine the

amount of energy in a particular place and therefore are associated with sediment composition. Depth is

also an important driver of macroinvertebrates distribution (Kraan et al. 2010, e.g. de Jong et al. 2015).

Strong shear stress and waves may dislodge benthic invertebrates, so we also expect these variables to

affect their total biomass (e.g. Anta et al. 2013, Malakauskas et al. 2013, but see Donadi et al. 2015a).

In soft sediment systems, seafloor topographic features are correlated with different sediment composition

(Trentesaux et al. 1994, e.g. Mestdagh et al. 2020). Different benthic communities are also associated with

different topographic features (e.g. Jansen et al. 2018, Mestdagh et al. 2020). For this reason, we included

slope, topographic position index (TPI) and topographic ruggedness as covariates to the models for mgs,

mud content and biomass.

Sediment composition determines the benthic community (Kraan et al. 2010, e.g. de Jong et al. 2015),

therefore, mgs, mud fraction and the standard deviation of sediment size were included in the biomass

model. Day of the year was also included in the biomass model to account for periods dominated by growth

or mortality.

Large fresh water discharges through the Afsluitdijk, at the southwestern part of the Wadden Sea, produce

density stratification, which in turn has an important effect on sediment transport (Elias 2006). Salinity is also

an important predictor of habitat type (Baptist et al. 2019) so it was also included in the biomass model.

The lugworm, *Arenicola marina*, the blue mussel *Mytilus edulis* and the Pacific oyster *Crassostrea gigas* are

important ecosystem engineers in the Wadden Sea (Volkenborn et al. 2007, Donadi et al. 2015b). *A. marina*

removes the sediment resuspending the smaller particles. Oysters and mussels, when filter-feeding, catch

the smaller particles suspended in the water column and fix them in the sediment when excreting. The

biomass of *A. marina* and the presence of a *M. edulis-C. gigas* bank were used as covariates in the mgs and

mud fraction models. *A. marina* biomass was determined through SIBES samples as described in the 'survey'

section. Mussel and oyster banks in the Wadden Sea are mapped every year (Troost et al. 2022). Because the

effects of such bivalve banks on sediment extent beyond the bank itself (Kröncke 1996), we used the

position of bivalve banks plus a buffer of 300 m as indication of bivalve bank presence.

Spatial modelling

We modeled three response variables: sediment mgs (μm), mud fraction (%) and total macroinvertebrate

biomass (afdm, g m^{-2}), each using an appropriate error distribution. We used a Gaussian distribution for mgs

and a beta distribution with a logit link function for mud fraction (Douma & Weedon 2019). As all the

samples contained detectable biomass, a Gamma distribution with a log link was used for total biomass

(Zuur et al. 2019).

Covariates described in the previous section and Table 1 were included in the models as fixed-effects.

Collinearity between variables was assessed by computing variance inflation factors (VIF) with function 'vif'

of car R package (Fox et al. 2011). We sequentially dropped the covariates with the highest VIF until all

presented VIF < 2 (Zuur et al. 2010). In this way, topographic ruggedness was dropped from the models. To

avoid numerical estimation problems, all covariates were standardized by subtracting their mean and

dividing by their standard deviation. We used smoothers when Pearson's residuals from generalized linear

mixed models showed patterns relative to covariates.

Subsidence was included in the model as a dichotomous variable: each observation was classified as within

or outside the subsidence area. We did this because 38% of our observations within the subsidence area fell

inside the 1 cm contour, and therefore we did not have enough information to analyze its effect as

continuous. We assessed whether the effect subsidence changed in time by adding an interaction term with

'year'. Priors for all fixed effects were set to a mean of zero and a precision of 1. This high prior weight

around zero results in the need of more data support for the posterior of a parameter to be different from

zero, which makes our results more conservative.

To account for spatial autocorrelation between observations that cannot be explained by the available

covariates, a stochastic partial differential equation (SPDE, Lindgren et al. 2011) model was used. In this

approach, a mesh of triangles is constructed for the study area and used to estimate the spatially- and

temporally-structured random component (SPDE specifications in ESM1).

Our main objective with this study was to test if subsidence affected each of the response variables or not,

therefore we focused on hypothesis testing rather than on model selection (Fieberg 2022). We considered

that subsidence had an effect on sediment or biomass if the confidence interval for its parameter or for the

interaction term was different from zero.

Table 1: Alphabetically ordered variables that were included in the models fitted to sediment median grain

size (sediment mgs), mud fraction, and total biomass data. Source of the data, year of measurements, and

frequency distributions of the data. The colors in the frequency plots indicate outside (blue) and inside (red)

the subsidence area.

Variable name	Model used	Source	Years measured	Histogram
Arenicola marina biomass	Sediment mgs Mud fraction	This study	2008-2020	
Bivalve bank presence	Sediment mgs Mud fraction	https://shiny.wur.nl/Schelpdiermonitor_Banken/	2008-2020	
Depth	Sediment mgs Mud fraction Total biomass	Rijkswaterstaat	2005, 2013, 2017	
Distance to coast	Sediment mgs Mud fraction Total biomass	This study	-	
Distance to tidal inlets	Sediment mgs Mud fraction Total biomass	This study	-	

Day of the year	Total biomass	-	-	 Day
Mud fraction	-	This study	2008-2020	 Mud %
Salinity	Sediment mgs Mud fraction Total biomass	Duran-Matute et al 2014	2013	 Salinity
Sediment mgs	Total biomass	This study	2008-2020	 Sediment mgs (µm)
Sediment size's standard deviation	Total biomass	This study	2008-2020	 Sediment sd (µm)
Current shear stress	Sediment mgs Mud fraction Total biomass	Folmer et al. 2016	2009-2010	 Shear stress
Slope	Sediment mgs Mud fraction Total biomass	Derived from "Depth"	2005, 2013, 2017	 Slope (°)
Topographic position index	Sediment mgs Mud fraction Total biomass	Derived from "Depth"	2005, 2013, 2017	 TPI
Total biomass	-	This study	2008-2020	 Biomass (g m $^{-2}$)
Wave forcing	Sediment mgs Mud fraction Total biomass	Donker et al. 2015	2010	 Wave forcing (ms $^{-1}$)

| Year | Sediment mgs | - | - |

 |
|---|---|---|---|---|
| | Mud fraction | | | |
| | Total species | | | |
| | biomass | | | |

Species-specific effects of land subsidence

Overall biomass is an aggregate of different species which might be differently affected by subsidence. To

analyze if particular species are driving the observed biomass patterns, we identified the more abundant

species (in terms of biomass) within the subsidence area, and plotted their relative contribution to total

biomass within and outside the subsidence area.

To understand how species with different habitat use are affected by subsidence, we compared population

trends within and outside the subsidence area for a larger group of species (ESM2). To classify species'

habitat use, for each species, we extracted the median of the 'median grain size' and the median of the

depth range in which they occurred. Hereafter we call these median values 'sediment use' and 'depth use'.

To detect differences between population trends, we calculated the ratio between the average biomass

within and outside the subsidence area for each species and year. We used a linear mixed model with

log('biomass ratio') as response variable, 'depth use' and 'year' as continuous fixed effects, and 'species' as

random slope and random intercept. We did not include 'sediment use' in the model to avoid collinearity

with 'depth use' (Pearson's correlation: -0.72, $P < 0.001$). We applied a likelihood ratio test to select the best

random structure and AIC together with 95% parameter confidence interval to select the best model relative

to fixed effects (Zuur 2009).

Results

Median grain size

The average mgs of all samples was 154 ± 41 µm (mean ± sd). The mixed-effects model fitted to mgs

revealed a positive effect of distance to coast, current shear stress, topographic position index, wave forcing

and salinity, and a negative effect of the presence of bivalve banks and distance to tidal inlet (Fig 2A). The

local depth and biomass of *Arenicola marina* had non-linear effects on mgs (Fig 2A). The model revealed an

interaction between subsidence and year on sediment size (Fig 2A). This meant that while there was no

evidence for a positive or negative trend in sediment median grain throughout the Wadden Sea, within the

subsidence area, mgs has been decreasing (Fig 2B). At the end of the study period, the model estimated that

mgs within the subsidence area was on average 16 µm smaller than outside (Fig 2B).

Fig 2: Results for the sediment median grain size model. **A.** Parameter estimates for covariates (mean ± 95%

CI). Covariates for which a smoother was applied have more than one parameter. Points and lines in black

show effects that are different from zero with 95% confidence. **B.** Mean and 95% confidence intervals of

median grain size, within and outside the subsidence area throughout the study period.

Mud fraction

On average, samples contained 0.13 ± 0.13 fraction of mud (mean ± sd). Mud fraction was negatively

affected by most of the covariates assessed, but positively affected by bivalve bank presence and wave

forcing (Fig 3 A). We found a positive effect of subsidence on the mud fraction (Fig 3A) that remained

constant in time (Fig 3B). On average, the mud fraction within the subsidence area was 0.03 higher than in

the rest of the Wadden Sea.

Fig 3: Results for the total sediment mud fraction model. **A.** Parameter estimates for covariates (mean ± 95%

CI). Points and lines in black show effects that are different from zero with 95% confidence. **B.** Mean and

95% confidence intervals of mud fraction, within and outside the subsidence area throughout the study

period.

Biomass

We found an average biomass of 24.5 ± 45.3 g m^{-2} (mean ± sd). Biomass was positively affected by wave

forcing and negatively affected by distance to tidal inlet, current shear stress and slope (Fig 3A). Depth and

day of the year had non-linear effects on biomass (Fig 3B). We did not find an effect of subsidence on total

biomass of the macroinvertebrate community throughout the studied period (Fig 3C), but there was a

negative trend within the subsided area and a slightly positive trend outside (Fig 3D).

Fig 4: Results for the total biomass model. **A.** Parameter estimates for covariates (mean ± 95% CI). Covariates

for which a smoother was applied have more than one parameter. Points and lines in black show effects that

are different from zero with 95% confidence. **B.** Mean and 95% confidence intervals of biomass, within and

outside the subsidence area throughout the study period.

Species-specific effects of land subsidence

Fig 5: Relative contribution of species to overall biomass within and outside the subsidence area. The

portrayed species are the ones with higher contribution to biomass within the subsidence area.

The most abundant species (in terms of biomass) within the subsidence area were *Arenicola marina*,

Cerstoderma edule, *Lanice conchilega*, *Mya arenaria* and *Mytilus edulis*. These species also represented an

important fraction of biomass outside the subsidence area, with other species only contributing up to 5 ± 1%

of total biomass (Fig 5). From 2008 to 2019, *C. edule* was the species with highest biomass in both areas,

however, its contribution was higher within the subsidence area. *C. edule* represented the 56% of total

biomass within the subsidence area in 2008, but only the 40% outside. In 2020, *C. edule* was surpassed by *A.*

marina in both areas, and by *M. arenaria* outside the subsidence area (Fig 5).

With the mixed-effects model, we found differences in population trends within and outside the subsidence

area, and these differences depended on the 'depth use' of species (Fig 6). For species that use deeper areas,

populations have increased in biomass within relative to outside the subsidence area. On the contrary, for

species that use shallower areas, populations have been slightly decreasing within relative to outside the

subsidence area (Fig 6).

Fig 6: Results for the population trends model. **A.** Parameter estimates for covariates (mean ± 95% CI). Points

and lines in black show effects that are different from zero with 95% confidence. **B.** Mean and 95%

confidence intervals of the ratio of biomass within and outside the subsidence area throughout the study

period.

Discussion

We found a change in sediment composition within the land subsidence area relative to outside. After

accounting for confounding environmental variables, median grain size within the subsidence area

decreased by 10% during the study period and the mud fraction was 3 percentage points higher throughout.

These changes do not seem to significantly affect the total biomass of the macroinvertebrates living there,

but there was a negative trend. Moreover, we did detect a significant increase of species that use deeper

habitat, suggesting changes at the community level.

The Wadden Sea is a complex and dynamic system, and therefore, it is difficult to establish a cause-and-

effect relationship between changes in sediment (or species) composition and land subsidence. Subsidence

occupies an area with unique characteristics in the Wadden sea, and any change in the subsidence area

could be caused by other natural or anthropogenic processes, raising the issue of a lack of a proper control in

this study, a problem inherent to nearly all ecological field studies. To attribute causation, a BACI design is

necessary (Green 1979). Furthermore, subsidence has been going on long before we started collecting the

data, so using the initial conditions (i.e. data from 2008) as the control would be incorrect. However, median

grain size, mud fraction and biomass are not randomly distributed in space, rather their distribution results

from environmental and biological processes that we aimed to capture through the covariates (like depth

and bed level shear stress of currents) in the statistical model. Because the range of all the covariates within

the subsidence area fall within the range of the whole Wadden Sea's (see histograms in Table 1), we can

attempt to detect differences "over and above" these processes between the subsidence area and the rest

of the Wadden Sea. On top of this, the addition of a spatial latent field with a temporal structure allows us to

control for spatially and temporally structured errors caused by historic events and missing covariates. Even

after accounting for confounding environmental variables and residual spatiotemporal patterns, we still

observed differences in median grain size and mud fraction within and outside the subsidence area.

Subsidence has not affected the average bathymetry and morphology of the tidal flats (van der Vegt 2022),

which implies that sedimentation was so far able to keep up with subsidence. However, within the

subsidence area median grain size was decreasing, and the overall mud fraction was smaller throughout the

study period. Modelled deep subsidence is bowl-shaped, with deeper parts in the centre and shallower parts

on the edges of the subsidence contours. Sediment tends to be smaller in troughs (e.g. Mestdagh et al. 2020)

so the bowl-shaped area could therefore result in a smaller-than-expected sediment pattern. It is possible

then, that we are detecting the effects of subsidence on sediment composition, while changes in bathymetry

or morphology might be too subtle or compensated too fast to be detected.

Mud fraction was higher within the subsidence area in the beginning of the study and remained stable,

showing a different pattern than median grain size. This indicates that the decrease in median grain size is

due to an increase of small sand rather than an increase in mud fraction. The different patterns in mud

fraction and median grain size might be the result of mud and sand deposition being governed by

independent mechanisms (Colina Alonso et al. 2021). Furthermore, because subsidence has been going on

since the eighties (Fokker et al. 2018), these differences could very well be the result of the accumulation of

long-term effects: it is possible that the effects over mud fraction have plateaued, while the sand fraction

has been more recently affected.

We did not find statistical differences in total biomass within and outside the subsidence area, but a

decreasing trend within was observed. It is possible that this trend is due to the prevalence of *Cerastoderma*

edule within the subsidence area. Both, within and outside, this species decreased in biomass during the

study period, but biomass within the subsidence area showed a larger drop. The reasons behind this pattern

need further study, but the habitat-use analysis shows that *C. edule* is one of the species that uses shallower

areas, and that species using shallower areas are decreasing in biomass within the subsidence area relative

to outside.

Species that increase within the subsidence area, relative to the outside, are those that use deeper areas.

Because these are most species, this results in a counter-intuitive pattern, in which most species increase

within the subsidence area relative to outside, while at the same time total biomass within the subsidence

area shows a decreasing trend. There is another counter-intuitive pattern: median grain size is getting

smaller within the subsidence area and, simultaneously, species that use deeper areas – often with coarser

sand – are doing better within the subsidence area relative to outside. Due to the interdependency of depth

and sediment composition, species that use deeper areas are generally those that use coarse sediment. With

this study, we cannot tell if species are choosing for a specific sediment or for a specific depth, but these

patterns might be the result of tidal flats sinking in some places while getting higher in others within the

subsidence area, despite the average elevation being unaffected (van der Vegt 2022).

The Wadden Sea as a UNESCO heritage site is a protected area. Information collected with SIBES together

with statistical tools as those used in this study, could also prove useful to assess the impacts of other human

activities in the area. For example, there are other areas that were or are subject to anthropogenic

subsidence in the Wadden Sea (Fokker et al. 2018). These were not included in the present study because

the information available is not comparable, but applying the methods described here might give us a better

idea of the effects that these and other human activities might have. Furthermore, where human activities

have started after the beginning of the monitoring program, SIBES could be used to perform proper "before,

after – control, impact" studies to study causation.

The gas extraction near Ameland is managed following the "hand on the tap" principle, meaning that "if the

natural values of the Wadden Sea are compromised", gas extraction must be reduced or even cease.

Intertidal area loss is the most evident threat posed by subsidence, so concrete thresholds have been

established to prevent it (6 mm y^{-1} for the saltmarsh area, National Project Decree 2005). Other threats

posed by subsidence to "natural values" would also warrant the adjustment of gas production (National

Project Decree 2005), but in those cases three problems might hinder the effective implementation of the

"hand on the tap" principle. Frist, the lack of definition of what the Wadden Sea's natural values are. Are

sediment and community structure natural values? Second, the lack of guidelines about what constitutes an

effect. Here, we show effects on sediment and community composition might be in place, but the lack of

measures from before the beginning of gas extraction precludes us from attributing causality. This does not

mean that effects do not exist, but rather that it is not possible to unambiguously prove them. To do proper

monitoring of potential effects, before-after control-impact studies are essential. The third problem is the

lack of a threshold for these effects that would warrant management consequences. If the 1 μm y^{-1} decrease

in median grain size is indeed caused by gas production, does this warrant management consequences?

Could this be considered a threat to the natural values of the Wadden Sea? For a successful implementation

of the "hand on the tap" principle it is necessary to clearly define concepts and thresholds beforehand,

otherwise the water will continue to run regardless of where the hand is.

Author contribution

Allert Bijleveld and Geert Aarts conceived the idea. Allert Bijleveld, Geert Aarts and Paula de la Barra

designed the methodology. Paula de la Barra analyzed the data; interpretation of results was discussed by all

three authors. Paula de la Barra led the writing of the manuscript. All authors contributed critically to the

drafts and gave final approval for publication.

Acknowledgements

We are very grateful to the current SIBES core-team for collecting and processing the thousands of samples.

In alphabetical order they are Leo Boogert, Thomas de Brabander, Anne Dekinga, Job ten Horn, Loran Kleine

Schaars, Jeroen Kooijman, Anita Koolhaas, Franka Lotze, Simone Miguel, Luc de Monte, Dennis Mosk, Amin

Niamir, Dana Nolte, Dorien Oude Luttikhuis, Bianka Rasch, Reyhane Roohi, Juan Schiaffi, Marten Tacoma,

Eveline van Weerlee and Bas de Wit. We also thank all former and current employees and the many

volunteers and students who have ensured that the SIBES program has continued in recent years. The RV

Navicula was essential for collecting the samples and particularly we thank the current crew Wim Jan Boon,

Klaas Jan Daalder, Bram Fey, Hendrik Jan Lokhorst and Hein de Vries. SIBES is currently financed by the

Nederlandsche Aardolie Maatschappij NAM, Rijkswaterstaat RWS and the Royal NIOZ.

Conflict of Interest

SIBES is currently partly financed by the Nederlandsche Aardolie Maatschappij NAM, which exploits the area

for gas production generating the land subsidence studied here.

Data availability statement

Data will be available at NIOZ Data Archive System upon publication.

References

Anta J, Peña E, Puertas J, Cea L (2013) A bedload transport equation for the Cerastoderma edule
cockle. J Mar Syst 111–112:189–195.

Baptist MJ, van der Wal JT, Folmer EO, Gräwe U, Elschot K (2019) An ecotope map of the trilateral
Wadden Sea. J Sea Res 152:101761.

Bartzke G, Bryan KR, Pilditch CA, Huhn K (2013) On the Stabilizing Influence of Silt On Sand Beds.
J Sediment Res 83:691–703.

Beukema JJ (1976) Biomass and species richness of the macro-benthic animals living on the tidal
flats of the Dutch Wadden Sea. Neth J Sea Res 10:236–261.

Beukema JJ (2002) Expected changes in the benthic fauna of Wadden Sea tidal flats as a result of
sea-level rise or bottom subsidence. J Sea Res 47:25–39.

Beukema JJ, Dekker R (2019) Is food supply for shellfish-eating birds in the western Wadden Sea
affected by the between-species synchrony in year-to-year fluctuations of bivalve population
parameters? Helgol Mar Res 73:8.

Bijleveld AI, van Gils JA, van der Meer J, Dekinga A, Kraan C, van der Veer HW, Piersma T (2012)
Designing a benthic monitoring programme with multiple conflicting objectives: *Sampling*
design for conflicting objectives. Methods Ecol Evol 3:526–536.

Christianen MJA, Middelburg JJ, Holthuijsen SJ, Jouta J, Compton TJ, van der Heide T, Piersma T,
Sinninghe Damsté JS, van der Veer HW, Schouten S, Olff H (2017) Benthic primary
producers are key to sustain the Wadden Sea food web: stable carbon isotope analysis at
landscape scale. Ecology 98:1498–1512.

Colina Alonso A, van Maren DS, Elias EPL, Holthuijsen SJ, Wang ZB (2021) The contribution of
sand and mud to infilling of tidal basins in response to a closure dam. Mar Geol 439:106544.

Compton TJ, Holthuijsen S, Koolhaas A, Dekinga A, ten Horn J, Smith J, Galama Y, Brugge M, van
der Wal D, van der Meer J, van der Veer HW, Piersma T (2013) Distinctly variable
mudscapes: Distribution gradients of intertidal macrofauna across the Dutch Wadden Sea. J
Sea Res 82:103–116.

Defew EC, Tolhurst TJ, Paterson DM (2002) Site-specific features influence sediment stability of
intertidal flats. Hydrol Earth Syst Sci 6:971–982.

Donadi S, Eriksson BK, Lettmann KA, Hodapp D, Wolff J-O, Hillebrand H (2015a) The body-size
structure of macrobenthos changes predictably along gradients of hydrodynamic stress and
organic enrichment. Mar Biol 162:675–685.

Donadi S, van der Heide T, Piersma T, van der Zee EM, Weerman EJ, van de Koppel J, Olff H,
Devine C, Hernawan UE, Boers M, Planthof L, Klemens Eriksson B (2015b) Multi-scale
habitat modification by coexisting ecosystem engineers drives spatial separation of
macrobenthic functional groups. Oikos 124:1502–1510.

Douma JC, Weedon JT (2019) Analysing continuous proportions in ecology and evolution: A
practical introduction to beta and Dirichlet regression. Methods Ecol Evol 10:1412–1430.

Elias EPL (2006) Morphodynamics of Texel Inlet. Doctoral Thesis, Technische Universiteit Delft

Eriksson BK, van der Heide T, van de Koppel J, Piersma T, van der Veer HW, Olff H (2010) Major
Changes in the Ecology of the Wadden Sea: Human Impacts, Ecosystem Engineering and
Sediment Dynamics. Ecosystems 13:752–764.

Fieberg J (2022) Statistics for Ecologists: A Frequentist and Bayesian Treatment of Modern
Regression Models. An open-source online textbook.

Field SA, O'connor PJ, Tyre AJ, Possingham HP (2007) Making monitoring meaningful. Austral Ecol
32:485–491.

Fokker PA, Leijen FJ van, Orlic B, Marel H van der, Hanssen RF (2018) Subsidence in the Dutch
Wadden Sea. Neth J Geosci 97:129–181.

Fox J, Weisberg S, Fox J (2011) An R companion to applied regression, 2nd ed. SAGE
Publications, Thousand Oaks, Calif.

Green RH (1979) Sampling Design and Statistical Methods for Environmental Biologists. John Wiley
& Sons.

Hewitt JE, Thrush SE, Cummings VJ (2001) Assessing Environmental Impacts: Effects of Spatial
and Temporal Variability at Likely Impact Scales. Ecol Appl 11:1502–1516.
Jansen J, Hill NA, Dunstan PK, Eléaume MP, Johnson CR (2018) Taxonomic Resolution,
Functional Traits, and the Influence of Species Groupings on Mapping Antarctic Seafloor
Biodiversity. Front Ecol Evol 6.
de Jong MF, Baptist MJ, Lindeboom HJ, Hoekstra P (2015) Relationships between
macrozoobenthos and habitat characteristics in an intensively used area of the Dutch
coastal zone. ICES J Mar Sci 72:2409–2422.
Ko J-Y, Day JW (2004) A review of ecological impacts of oil and gas development on coastal
ecosystems in the Mississippi Delta. Ocean Coast Manag 47:597–623.
Kraan C, Aarts G, van der Meer J, Piersma T (2010) The role of environmental variables in
structuring landscape-scale species distributions in seafloor habitats. Ecology 91:1583–
1590.
Kröncke I (1996) Impact of Biodeposition on Macrofaunal Communities in Intertidal Sandflats. Mar
Ecol 17:159–174.
Li C, Yang S, Zha D, Zhang Y, Boer WF de (2019) Waterbird Communities in Subsidence Wetlands
Created by Underground Coal Mining in China: Effects of Multi-Scale Environmental and
Anthropogenic Variables. Environ Conserv 46:67–75.
Lindgren F, Rue H, Lindström J (2011) An explicit link between Gaussian fields and Gaussian
Markov random fields: the stochastic partial differential equation approach. J R Stat Soc Ser
B Stat Methodol 73:423–498.
Malakauskas DM, Willson SJ, Wilzbach MA, Som NA (2013) Flow variation and substrate type
affect dislodgement of the freshwater polychaete, Manayunkia speciosa. Freshw Sci
32:862–873.
Mestdagh S, Amiri-Simkooei A, van der Reijden KJ, Koop L, O'Flynn S, Snellen M, Van Sluis C,
Govers LL, Simons DG, Herman PMJ, Olff H, Ysebaert T (2020) Linking the morphology and
ecology of subtidal soft-bottom marine benthic habitats: A novel multiscale approach. Estuar
Coast Shelf Sci 238:106687.
Milliman JD, Haq BU (eds) (1996) Sea-level rise and coastal subsidence: causes, consequences,
and strategies. Kluwer Academic Publishers, Dordrecht ; Boston.
NAM B.V (2021) Gaswinning vanaf de locaties Moddergat, Lauwersoog en Vierhuizen. Resultaten
uitvoering Meet- en regelcyclus 2020.
NAM B.V (2007) Meet- en regel protocol Winning Waddenzeegas vanaf de locaties Moddergat,
Lauwersoog en Vierhuizen.
National Project Decree (2005) Rijksprojectbesluit Gaswinning onder de Waddenzee vanaf de
locaties Moddergat, Lauwersoog en Vierhuizen. Deel 1.
Oost AP (1995) Dynamics and sedimentary developments of the Dutch Wadden Sea with a special
emphasis on the Frisian Inlet : a study of the barrier islands, ebb-tidal deltas, inlets and
drainage basins. Dissertation, Faculteit Aardwetenschappen
Poiesz SSH, Witte JI, Veer HW van der (2020) Only a few key prey species fuel a temperate coastal
fish food web. Mar Ecol Prog Ser 653:153–166.
Postma H (1982) Hydrography of the Wadden sea: movements and properties of water and
particulate matter final report on "hydrography." A. A. Balkema, Rotterdam.
Sanchez-Salazar ME, Griffiths CL, Seed R (1987) The interactive roles of predation and tidal
elevation in structuring populations of the edible cockle, Cerastoderma edule. Estuar Coast
Shelf Sci 25:245–260.
Sansom A, Pearce-Higgins JW, Douglas DJT (2016) Negative impact of wind energy development
on a breeding shorebird assessed with a BACI study design. Ibis 158:541–555.
Smokorowski KE, Randall RG (2017) Cautions on using the Before-After-Control-Impact design in
environmental effects monitoring programs. FACETS 2:212–232.
Thienen-Visser V, Breunese JN, Muntendam-Bos AG (2015) Subsidence due to gas production in
the Wadden Sea: How to ensure no harm will be done to nature. In: *49th US Rock*
Mechanics/Geomechanics Symposium. OnePetro
Trentesaux A, Stolk A, Tessier B, Chamley H (1994) Surficial sedimentology of the Middelkerke
Bank (southern North Sea). Mar Geol 121:43–55.

Troost K, Asch M van, Ende D van den, Es Y van, Perdon KJ, Pool J van der, Suykerbuyk W,
Zweeden C van, Zwol J van (2022) Schelpdierbestanden in de Nederlandse kustzone,
Waddenzee en zoute deltawateren in 2021.
Turner RE, Mo Y (2021) Salt Marsh Elevation Limit Determined after Subsidence from Hydrologic
Change and Hydrocarbon Extraction. Remote Sens 13:49.
Underwood AJ (1991) Beyond BACI: Experimental designs for detecting human environmental
impacts on temporal variations in natural populations. Mar Freshw Res 42:569–587.
Underwood AJ (1993) The mechanics of spatially replicated sampling programmes to detect
environmental impacts in a variable world. Aust J Ecol 18:99–116.
Van Colen C (2018) The Upper Living Levels: Invertebrate Macrofauna. In: *Mudflat Evology*. Aquatic
Ecology Series, Springer International Publishing
van der Vegt H (2022) Monitoring wadplaatareaal Friesche Zeegat met LiDAR (2010-2021).
Volkenborn N, Hedtkamp SIC, van Beusekom JEE, Reise K (2007) Effects of bioturbation and
bioirrigation by lugworms (Arenicola marina) on physical and chemical sediment properties
and implications for intertidal habitat succession. Estuar Coast Shelf Sci 74:331–343.
Wang ZB, Elias EPL, van der Spek AJ, Lodder QJ (2018) Sediment budget and morphological
development of the Dutch Wadden Sea: impact of accelerated sea-level rise and subsidence
until 2100. Neth J Geosci 97:183–214.
Wiesebron LE, Steiner N, Morys C, Ysebaert T, Bouma TJ (2021) Sediment Bulk Density Effects on
Benthic Macrofauna Burrowing and Bioturbation Behavior. Front Mar Sci 8:707785.
Yang P, Shu Q, Liu Q, Hu Z, Zhang S, Ma Y (2021) Distribution and factors influencing organic and
inorganic carbon in surface sediments of tidal flats in northern Jiangsu, China. Ecol Indic
126:107633.
Zuur AF (ed) (2009) Mixed Effects Models and Extensions in Ecology with R. Springer, New York,
NY.
Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common statistical
problems. Methods Ecol Evol 1:3–14.
Zuur AF, Ieno EN, Savel'ev AA, Zuur AF (2019) Beginner's guide to spatial, temporal and spatial-
temporal ecological data analysis with R-INLA. Highland Statistics Ltd, Newburgh, United
Kingdom.
530