1
2 Number of words in the Abstract: 267
3 Number of words in the Introduction: 630
4 Number of words in the Discussion: 1301
5 Number of pages: 30
6 Number of Figures: 5

8 Systematic characterization of a non-transgenic
9 Aβ_{1-42} amyloidosis model: synaptic plasticity
10 and memory deficits in female and male mice

Raquel Jiménez-Herrera[#], Ana Contreras[#], Guillermo Iborra-Lázaro, Danko Jeremic,

Souhail Djebari, Juan Navarro-López [*], Lydia Jiménez-Díaz [*].

Neurophysiology & Behavior Lab, Centro Regional de Investigaciones Biomédicas

(CRIB), School of Medicine of Ciudad Real, University of Castilla-La Mancha, Ciudad

Real, Spain.

[#]These authors share first authorship

[*]These authors contributed equally to this work and share last authorship.

#Correspondence to:

J.D. Navarro-López, L. Jiménez-Díaz

NeuroPhysiology and Behavior Lab

Universidad Castilla-La Mancha

13071-Ciudad Real

Phone: + 34-926295300 ext 6838

Fax: + 34-926295300

E-mail: Juan.Navarro@uclm.es; mailto:Lydia.Jimenez@uclm.es

Abstract

Background: One of the neuropathological hallmarks of Alzheimer's disease (AD) is amyloid-β (Aβ) accumulation in the hippocampus that causes its dysfunction. This disruption includes excitatory/inhibitory imbalance, synaptic plasticity and oscillatory activity impairments, and memory deficits. Although AD prevalence is higher in women than men, the possible sex difference is scarcely explored and information from amyloidosis transgenic mice models is contradictory. Thus, given the lack of data of the early amyloidosis stages in females, the aim of this study was to systematically characterize the effect of an intracerebroventricular (*icv.*) injection of Aβ_{1-42} on hippocampal-dependent memory, and on associated activity-dependent synaptic plasticity in the hippocampal CA1–CA3 synapse, in both male and female mice.

Methods: To do so, we evaluated long term potentiation (LTP) with *ex vivo* electrophysiological recordings and spatial (working, short- and long-term) and exploratory habituation memory using Barnes maze or open field habituation tasks respectively.

Results: We found that Aβ_{1-42} administration impairs all forms of memory evaluated, regardless the sex, in a long-lasting manner (up to 17 days post-injection). Furthermore, LTP was inhibited at a postsynaptic level, both in males and females, and a long-term depression (LTD) was induced for the same prolonged period, which could underly memory deficits.

Conclusions: In conclusion, our results provide further evidence of the shifting of LTP/LTD threshold due to a single *icv*. Aβ_{1-42} injection, which underly cognitive deficits in early stages of AD. These long-lasting cognitive and functional alterations in males and females validate this model for the study of early amyloidosis in both sexes, thus offering a solid alternative to the inconsistence of amyloidosis transgenic mice models.

Keywords

Alzheimer's disease; amyloid-β; spatial memory; LTP; hippocampus; sex differences.

Background

The role of the hippocampus in learning and memory processes is well

stablished (1, 2). Several types of memory, such as episodic memory, spatial memory,

or contextual fear, are dependent on the hippocampus (3-5). Spatial memory is

essential for the survival of all kind of animals, since it allows the retrieval of the

location of an object, and to place an experience in a specific context in the

environment (6). The encoding of spatial memory relies on the catching of behaviorally

relevant spatial cues on a timescale of seconds, known as spatial working memory (7).

Moreover, the hippocampus is one of the most vulnerable brain areas, prone to

succumb to oxidative stress derived from different diseases (8). This can lead to the

characteristic memory and learning impairments of pathologies like Alzheimer's

disease (AD), addiction or major depression, among others (9, 10).

AD is the main cause of dementia, affecting over 55 million people worldwide

(11). One of the neuropathological hallmarks of AD is amyloid-β (Aβ) plaques

accumulation in the hippocampus, known as senile plaques, seen preferentially in the

late stages of the disease (12). Aβ causes a hippocampal dysfunction, which include

excitatory/inhibitory (E/I) imbalance, impairments of hippocampal oscillatory activity,

synaptic plasticity disruption and memory deficits (13-15). In the early stages of AD,

electrophysiological data have proven that soluble Aβ oligomers can block

hippocampal long-term potentiation (LTP) while enhancing long-term depression (LTD)

(16, 17). These two mechanisms of synaptic strengthening and weakening,

respectively, underlie synaptic plasticity, and the balance between both is highly

important to maintain proper hippocampal functionality, such as learning and memory

processes (18). Indeed, it has been previously demonstrated that LTP inhibition caused

by Aβ correlates with the memory impairments reported in AD (15, 19-23).

As mentioned above, the accumulation of Aβ in the hippocampus is an

important factor for AD development (16) found in both sporadic AD - the most

common form of the disease- and familial inherited AD. Many studies use transgenic

mice which express and accumulate the human Aβ peptide (Amyloid Precursor Protein

-APP- models based on human mutations that represent less than 5% of AD cases

(inherited AD)), *tau* hyperphosphorylation or both (24). However, the physiopathology

of AD starts even decades before the senile plaques are present, when Aβ is soluble

rather than accumulated (25). The Aβ_{1-42} fragment has higher propensity to form

amyloid fibrils than other fragments and is found to be the dominant Aβ specie in the

amyloid plaques of AD patients (24, 26). Moreover, it has been proven that

intracerebroventricular (*icv.*) administration of Aβ diffuses mainly to the dorsal

hippocampal formation (22), which is related specifically to spatial learning and

memory (27, 28). Accordingly, this model seems a better option for investigating early

stages of sporadic AD (29). In this line, previous work from our group have shown an

E/I imbalance in male mice after an *icv.* injection of Aβ_{1-42} both *in vivo* and *in vitro*. This

translated into a deficit in LTP induction, a disruption of the correct neural oscillatory

synchronization in the hippocampus and, ultimately, an impairment in learning and

memory processes (22, 23, 30).

According to the Alzheimer's Association, nearly two-thirds of AD patients in the

US are women (31). Even though AD prevalence is higher in women than men, almost

all the available data comes from studies with male mice (32). Therefore, it is crucial to

include both, male and female animals in this type of studies. A few recent works have

done so, although the results are inconsistent, showing sex-dependent differences (20,

33-36) or not (37, 38) depending on the animal model used, the specific parameters

measured and/or the experiments carried out.

Thus, given the lack of data regarding AD models in females, the aim of the

present study was to characterize the effect of *icv.* Aβ_{1-42} administration on working,

short-term and long-term memory, as well as on synaptic plasticity processes, to

provide a solid model to study early stages of amyloidosis considering sex differences.

Materials and methods

Animals

Female and male C57BL/6 adult mice (12-24 weeks old; 20-30 g) were used

(RRID:MGI:5,656,552; Charles River, US). Animals were kept on 12 h light/dark cycles

with access to food and water *ad libitum*. Temperature (21 ± 1 ºC) and humidity

(50 ± 7%) were controlled. Mice were housed in same-sex groups of 5 per cage before

surgery, and individually afterwards. All experimental procedures were carried out at

the same time interval in both female and male mice to minimize circadian rhythm

interferences.

All experimental procedures were reviewed and approved by the Ethical

Committee for Use of Laboratory Animals of the University of Castilla-La Mancha

(PR-2022-11-04 and PR-2018-05-11) and conducted according to the European Union

guidelines (2010/63/EU) and the Spanish regulations for the use of laboratory animals

in chronic experiments (RD 53/2013 on the care of experimental animals: BOE

08/02/2013).

Surgery for drug injection

Mice were anesthetized with 4% isoflurane (#13400264, ISOFLO, Proyma S.L.,

Spain) using a calibrated R580S vaporizer (RWD Life Science; flow rate: 0.5 L/min O_2).

After induction, 1.5% isoflurane was delivered constantly for the maintenance of

anesthesia. Intramuscular buprenorphine (0.01mg/kg; #062009, BUPRENODALE,

Albet, Spain) and a healing cream (Blastoestimulina®; Almirall, Spain) were

administered as analgesic after surgery, to accelerate recovery and decrease animal

suffering.

As described elsewhere (23), for *icv*. administration of Aβ, animals were

implanted with a blunted, stainless steel, 26-G guide cannula (Plastics One, US) in the

left ventricle (1 mm lateral and 0.5 mm posterior to bregma; depth from brain surface,

1.8 mm) (39). The final position of the cannula was determined by Nissl staining

(Figure 1A).

Mice were allowed at least a week for recovery before any experimental

procedure. After recovery, freely moving animals received a 3 µg/µL *icv.* injection of

either Aβ_{1-42}, Aβ_{42-1} (as a reverse control) or vehicle (control, PBS) through an injection

cannula at a rate of 0.5 µL/min. For this purpose, the injection cannula was inserted

into the guide cannula protruding 0.5 mm into the ventricle and attached to a Hamilton

syringe. All drugs were dissolved in PBS and purchased from Bachem (#4014447 and

#4027991, respectively; Switzerland). Mice were routinely handled to minimize stress

during experimental procedures.

Barnes maze

To evaluate hippocampal-dependent working, short-term and long-term spatial

memory, the Barnes maze (LE851BSW, Panlab, Spain) was used (40). It consists of a

rotating platform disk (92 cm in diameter) with 20 scape holes (5 cm each) around its

periphery, elevated 1 meter above the floor. Spatial cues (circles and squares of

different colors) and mildly aversive white noise were used.

The protocol used was adapted from Suarez et al. (41). Briefly, each trial began

with the mouse inside a starting cylinder (8 x 12.5 cm) positioned in the center of the

maze. After 10 sec, the starting cylinder was removed, and the white noise was

initiated. Between each trial, the maze was cleaned with 70% ethanol to dissipate odor

cues. As shown in Figure 1B, the protocol consisted of a habituation trial the day before

icv. injection (Day−1), four training days of three trials each, starting the day of

treatment administration (Days 0 - 3), and two memory tests (Days 7 and 11). During

the habituation day, mice were able to explore the maze for 90 sec or until they found

the goal box (17.5 x 7.5 x 8 cm) attached to one of the holes. If time ran out, the animal

was gently led to the goal box. In both cases, mice were kept in the box for 1 min

before returning them to their cages. Afterwards, training was carried out on four

consecutive days, with three trials per day and a 15 min interval between trials. During

each trial, mice were allowed to explore the maze for 3 min or until they found the

escape hole, which was changed daily. Latency to find the box, errors made, and

distance traveled were measured. Finally, a single 90 sec trial was performed each

memory testing day, without any scape hole available. Latency and distance to the

target of the latest training day were measured. All sessions were recorded and

analyzed with Barnes-Smart video tracking software (Panlab, Spain).

Open field habituation task

On days 15 and 16 post-*icv.* injection (Figure 1C), to evaluate a non-associative

hippocampal-dependent learning process, such as exploratory habituation to a novel

environment (42), an open field (OF) habituation task was conducted as described

elsewhere (43). Briefly, mice were exposed twice during consecutive days to an OF, to

measure the change in exploration after re-exposure. In the training day (OF1), mice

were placed in the middle of a square acrylic box (23.5 x 17.5 x 4 cm plexiglas base

arena; 26.5 x 21 x 10 cm top) and allowed to freely explore it for 15 min. In the

retention day (OF2), 24 h later, mice were re-exposed to the same environment.

Exploratory behavior was recorded using a LABORAS® apparatus (*Laboratory Animal*

Behavior Observation Registration and Analysis System; Metris, Netherlands), which

transforms mechanical vibrations generated by the movements of the animals into

electrical signals through a sensing platform positioned under the cage.

Spontaneous behaviors

Between days 15 and 17 post-injection, mice went through a battery of

behavioral tests to assess their overall state and spontaneous behaviors (Figure 1D).

Laboratory animal behavior observation registration and analysis system

(LABORAS®) for stereotyped and locomotion behavioral testing

Mice were placed in a rectangular LABORAS® cage for a single 15-min trial, as

previously described (43). Grooming behavior was used as a measurement of

stress-related behavior (44). Locomotion, climbing, and rearing behaviors were used as

measurements of locomotor activity (45). All data were digitized and analyzed using the

LABORAS® software (Metris, Netherlands).

Rotarod performance test

The rotarod apparatus (LE 8500, Panlab, Spain), consisting of a 30 mm

diameter black striated rod positioned 20 cm above the floor, was used to measure

coordination and motor function. Mice were trained to stay for 1 min on the rod at

constant low-speed (6 rpm) rotation. The following day, 5 consecutive trials were

carried out, in which the rod was set to accelerate from 4 to 40 rpm for 2 min, and

mice's time to fall off was recorded.

Elevated plus maze

To assess anxiety-like behaviors (46), the elevated plus maze (LE 842, Panlab,

Spain) was used. It consists of a cross-shaped methacrylate platform with two open

arms (65 x 6 cm) without walls and two arms enclosed (65 x 6 cm) by 15-cm-high

opaque walls, mounted 90° to one another with a central platform (6.3 x 6.3 cm) and

raised 40 cm above the floor. Mice were placed into one of the open arms and were

allowed to freely explore the maze during a single 5 min session. The number of

entries into the open arms and the time spent there were calculated to measure

anxiety-like behavior. Additionally, as a measurement of locomotor activity (47),

number of entries into the closed arms and total entries (open + closed arms) were

recorded.

Tail suspension test

To assess depression-like behaviors, the tail suspension test was performed

(48). The tail suspension apparatus (BIO-TST5, Bioseb, US) consists of three PVC

chambers (50 x 15 x 30 cm each) in which animals were hung by the tail for 6 min

approximately 10 cm away from the ground. The immobility time was registered by the

strain sensors. All data were digitized and analyzed using the BIO-TST5 software

(Bioseb, US).

Ex vivo field EPSP (fEPSP) recordings

Coronal hippocampal slices were prepared as described previously (43). In

brief, animals were deeply anesthetized with halothane (Fluothane, AstraZeneca, UK)

and decapitated. The brain was immediately removed and rapidly immersed in

oxygenated (95% O_2 – 5% CO_2) ice-cold "cutting" solution containing (in mmol/L; all

from Sigma, US): 87 NaCl (#S9888), 10 glucose (#G8270), 75 sucrose (#84100), 1.25

NaH_2PO_4 (#S8282), 3 $C_3H_3NaO_3$ (#P2256), 0.98 $C_6H_7NaO_6$ (#11140), 25 $NaHCO_3$

(#S6014), 2.5 KCl (#P3911), 0.37 $CaCl_2$ (#499609), 3.28 $MgCl_2$ (#208337). Brain was

trimmed and mounted on the stage of a vibratome (7000smz-2; Campden Instruments,

UK) in such a way so that the blade would cut through hemispheres at an angle of

20-30° of their horizontal planes. Coronal slices (300 μm thick) containing the dorsal

hippocampus were then incubated for at least 1.5 h before recording, at room

temperature (22°C) in oxygenated artificial cerebrospinal fluid (aCSF) containing (in

mmol/L; all from Sigma, US): 125.99 NaCl (#S9888), 3 KCl (#P3911), 1.8 $CaCl_2$

(#499609), 1.5 $MgCl_2$ (#208337), 25.99 $NaHCO_3$ (#S6014), 10 glucose (#G8270), and

1.2 NaH_2PO_4 (#S8282).

For electrophysiological recordings, two set-ups consisting of a multi-electrode

array (MEA2100-Mini-System) pre-amplifier and a filter amplifier (gain 1100× or 550×)

were run in parallel by a data acquisition card governed by MC_Experimenter V2.20.0

software. A single slice was transferred to each MEA recording chamber (MEA60; Multi

Channel Systems, Reutlingen, Germany) continually perfused with aCSF (flow rate 2

mL/min) and kept at 32 °C. The MEA was positioned on the platform of an inverted

MEA-VMTC-1 video microscope and it comprised 60 extracellular electrodes

(inter-electrode distance: 200 μm). Each individual electrode from the array could be

used either as a recording or as a stimulation electrode. A nylon mesh was positioned

above the slice to obtain a satisfactory electrical contact between the surface of the

slice and the electrode array. Stimulation was achieved with a stimulus generator unit

integrated in the headstage (Multi Channel Systems, Germany) by applying biphasic

current pulses to one electrode of the array (S1) located in the Schaffer Collateral

pathway of the hippocampus. Field excitatory postsynaptic potentials (fEPSPs) could

then be recorded in the *stratum radiatum* of the CA1 subfield by all the remaining

electrodes of the array at the same time (Figure 1E). A second electrode of the array

(S2) was used to stimulate an independent pathway without High Frequency

Stimulation (HFS, stimulation protocol used to induce LTP; see details below), as

control of synaptic transmission.

Following at least 20 min of equilibration period inside the MEA well, basal

synaptic transmission was studied by input/output (I/O) curves, by applying two stimuli

of increasing intensities (0.02 – 0.4 mA) at 40 ms interstimulus interval. After I/O,

pulses were adjusted to ≈40% of the intensity necessary to evoke a maximum fEPSP

response. To address a typical short-term plasticity phenomenon at a presynaptic level,

the paired-pulse facilitation (PPF) protocol was used. Thus, pairs of stimuli were

delivered at different interstimulus intervals (10, 20, 40, 100, 200, 500 ms). For LTP

induction, a high-frequency stimulation (HFS) protocol was used, consisting of five 1-

sec-long 100-Hz trains delivered at 30 sec intertrain interval. Baseline (BL) values of

fEPSPs amplitude recorded at the CA3-CA1 synapse were collected for 15 min before

LTP induction. After HFS, fEPSPs were recorded during 60 min to evaluate LTP

induction. To pair it with the behavioral tasks, electrophysiology was carried out 1 to 17

days post-*icv.* injection of Aβ_{1-42} or controls: Aβ_{42-1} and vehicle.

Data was analyzed with the Multichannel Analyzer software (V 2.20.0). As

synaptic responses were not contaminated by population spikes, the amplitude (i.e.,

the peak-to-peak value in mV during the rise-time period) of successively evoked

fEPSPs was used. All values represented mean ± SEM with *n* indicating number of

slices. Within each slice, data from 3 different recording electrodes was used.

Statistical analysis

Data was represented as the mean ± SEM and analyzed by two-way ANOVA,

followed by Tukey's *post-hoc* analysis. When comparing only two groups, Student t test

was used. Statistical significance was set at $p < 0.05$. All analyses were performed

using SPSS software v.24 (RRID:SCR_002865; IBM, US) and GraphPad Prism

software v.8.3.1 (RRID:SCR_002798; Dotmatics, US). Final figures were prepared

using CorelDraw X8 Software (RRID:SCR_014235; Corel Corporation, Canada).

280   Results

281   　　　　Firstly, we wondered whether *icv.* injection of Aβ_{1-42} would affect spatial and

282   habituation learning and memory in a sex-specific manner. Thus, female and male

283   mice carried out a protocol to measure spatial learning and memory using a Barnes

284   maze, as well as an exploratory habituation protocol using an OF to assess this type of

285   non-associative memory.

Aβ_{1-42} impairs spatial learning and memory in both female and male mice

For the Barnes Maze task, animals performed 3 trails per day during 4 consecutive days (days 0-3), starting one hour after *icv.* injections (performed in day 0, Figure 1) of either Aβ$_{1-42}$ or both controls: vehicle and reverse Aβ$_{42-1}$ peptide. Differences in the latency to find the open hole between the first and last trial of each day (Figure 2A) is considered an assessment of working memory (41). No differences between males and females within each treatment group were found. Nevertheless, our data showed a significant treatment (Male: $F_{(2,33)}$ = 7.755, p = 0.0017; Female: $F_{(2,35)}$ = 8.719, $p < 0.0008$) and time (Male: $F_{(5.584,169.9)}$ = 13.43, $p < 0.001$, Geisser-Greenhouse's correction; Female: $F_{(5.706,185)}$ = 8.838, $p < 0.001$, Geisser-Greenhouse's correction) effect when comparing the latency to find the escape hole in the first and last trial from each training session, proving that both vehicle (male, n = 14; female, n = 15) and Aβ_{42-1} (male, n = 7; female, n = 6) groups had a normal spatial working memory while Aβ_{1-42} animals (male, n = 14; female, n = 16) showed a deterioration of this kind of memory, regardless of sex. Furthermore, the evolution of the different parameters measured along the four training days allowed us to evaluate short-term memory, since animals should remember how to perform the task and therefore latencies would decrease over time, even if the target hole was changed. Accordingly, our data showed a treatment effect for latency (Figure 2B; Male: $F_{(2,124)}$ = 47.97, $p < 0.001$; Female: $F_{(2,128)}$ = 23.12, $p < 0.001$), number of errors (Figure 2C; Male: $F_{(2,33)}$ = 18.14, $p < 0.001$; Female: $F_{(2,127)}$ = 12.09, $p < 0.001$) and distance traveled (Figure 2D; Male: $F_{(2,34)}$ = 4.59, p = 0.017; Female: $F_{(2,35)}$ = 11.30, p = 0.0002). *Post-hoc* analysis revealed that Aβ_{1-42} disrupted spatial learning in both female and male animals. Finally, to study long-term memory, two memory tests were conducted in days 7 and 11 post-injection, during which all holes were closed (Figure 2E) and latency to reach the target hole of the last training day (day 3), as well as distance traveled, were

quantified. Our results showed a statistically significant difference in latency (Figure 2F;

Male: $F_{(2,62)} = 7.644$, $p = 0.0011$; Female: $F_{(2,62)} = 9.618$, $p < 0.001$) without affecting

distance traveled (Figure 2G). Once again, *post-hoc* analysis revealed that the

differences were due to a worst performance of Aβ_{1-42}-treated mice. Thus, these data

suggest an impairment in hippocampal-dependent spatial working, short- and long-term

memory and learning processes induced by Aβ_{1-42} in both sexes. In contrast, same

concentrations of the control reverse peptide, Aβ_{42-1}, did not differ from the control

vehicle suggesting that the decline induced by Aβ_{1-42} was therefore specific.

Aβ_{1-42} impairs exploratory habituation memory in both female and

male mice

Afterwards, exploratory habituation memory was tested in days 15 and 16

post-*icv.* injection by an OF habituation task (Figure 3A). Data showed no significant

differences in exploration between groups during the training session (OF1). However,

when comparing OF1 and the retention session (OF2) (Figure 3B), a significant

decrease in exploration was observed in both controls, vehicle (Male: $t_{12} = 2.317$,

$p = 0.019$, n = 7; Female: $t_{18} = 4.751$, $p < 0.001$, n = 12) and Aβ_{42-1} (Male: $t_{12} = 6.195$,

$p < 0.001$, n = 7; Female: $t_{10} = 6.433$, $p < 0.001$, n = 6) treated animals, regardless of

sex, proving that they could remember the arena, while Aβ_{1-42} mice (male, n = 9;

female, n = 10) performed slightly better without reaching significance, indicating some

deterioration of memory in this group of animals. Moreover, during the retention

session (OF2), a significant treatment effect was found in both female (Figure 3B;

$F_{(2,22)} = 4.450$, $p = 0.024$) and male (Figure 3B; $F_{(2,20)} = 4.952$, $p = 0.018$) mice.

Post-hoc analyses revealed that Aβ_{1-42} mice traveled a longer distance during this

session than vehicle and Aβ_{42-1} control groups, as illustrated in Figure 3C. Hence, both

results show that Aβ_{1-42} also impairs non-associative habituation memory even 2 weeks

after treatment.

To verify that the deterioration observed after A$\beta_{1\text{-}42}$ injection both in

hippocampal-dependent spatial and habituation memory were due to specific

hippocampal alterations, and rule out general health affections, mice carried out a

battery of tests to evaluate different behavioral aspects. When assessing stereotyped

behaviors using the LABORAS®, data showed that all groups (vehicle male, n = 11;

vehicle female, n = 12; A$\beta_{1\text{-}42}$ male, n = 11; A$\beta_{1\text{-}42}$ female, n = 12; A$\beta_{42\text{-}1}$ male, n = 7;

A$\beta_{42\text{-}1}$ female, n = 6) spent the same amount of time performing the different behaviors

analyzed (Figure 4A): locomotion (Male: $F_{(2,28)}$ = 0.208, p = 0.814; Female:

$F_{(2,29)}$ = 0.493, p = 0.616), climbing (Male: $F_{(2,28)}$ = 0.596, p = 0.558; Female:

$F_{(2,29)}$ = 1.027, p = 0.372), rearing (Male: $F_{(2,28)}$ = 0.962, p = 0.395; Female:

$F_{(2,29)}$ = 0.434, p = 0.653) and grooming (Male: $F_{(2,27)}$ = 0.850, p = 0.439; Female:

$F_{(2,29)}$ = 1.120, p = 0.341). All groups equally improved their performance in the rotarod

(Figure 4B; vehicle male, n = 8; vehicle female, n = 11; A$\beta_{1\text{-}42}$ male, n = 8; A$\beta_{1\text{-}42}$

female, n = 9; A$\beta_{42\text{-}1}$ male, n = 7; A$\beta_{42\text{-}1}$ female, n = 6), as no differences in the latency

to fall off the rod were found between groups along trials (Male: $F_{(2,20)}$ = 0.4095,

p = 0.6694; Female: $F_{(2,23)}$ = 0.09378, p = 0.9108) nor in the whole session (Male:

$F_{(2,20)}$ = 1.783, p = 0.1939; Female: $F_{(2,23)}$ = 0.2739, p = 0.7629). Locomotion was also

tested in all the groups (vehicle male, n = 8; vehicle female, n = 12; A$\beta_{1\text{-}42}$ male, n = 9;

A$\beta_{1\text{-}42}$ female, n = 12; A$\beta_{42\text{-}1}$ male, n = 7; A$\beta_{42\text{-}1}$ female, n = 6) by the number of entries

into closed (Male: $F_{(2,23)}$ = 0.951, p = 0.403; Female: $F_{(2,29)}$ = 0.116, p = 0.891) and total

arms (Male: $F_{(2,23)}$ = 0.255, p = 0.777; Female: $F_{(2,29)}$ = 0.293, p = 0.748) in the elevated

plus maze and, once again, no differences were found (Figure 4C). Regarding

behaviors related to stress, all animals had the same number of entries (Figure 4C;

Male: $F_{(2,23)}$ = 0.663, p = 0.526; Female: $F_{(2,29)}$ = 0.381, p = 0.687) and time spent in

open arms (Male: $F_{(2,23)}$ = 0.273, p = 0.763; Female: $F_{(2,29)}$ = 0.391, p = 0.680) in the

elevated plus maze, and they all had the same immobility time (Figure 4D; vehicle

male, n = 8; vehicle female, n = 12; A$\beta_{1\text{-}42}$ male, n = 9; A$\beta_{1\text{-}42}$ female, n = 12; A$\beta_{42\text{-}1}$

male, n = 7; A$\beta_{42\text{-}1}$ female, n = 6; Male: $F_{(2,23)} = 0.003$, $p = 0.997$; Female:

$F_{(2,29)} = 0.077$, $p = 0.927$) in the tail suspension test, suggesting that the treatment did

not increase depression- nor anxiety-like behaviors.

Thus, this data confirmed that the overall health state and locomotor function

was uniform among the different treated groups and, therefore, all learning and

memory impairments observed were due to a specific hippocampal disruption caused

by A$\beta_{1\text{-}42}$.

A$\beta_{1\text{-}42}$ inhibits *ex vivo* LTP in both female and male mice similarly

Given the deleterious effects of A$\beta_{1\text{-}42}$ on hippocampal-dependent learning and

memory processes in the present amyloidosis model in a similar way regardless the

sex, we wondered whether excitability, presynaptic function, and short- and long-term

plasticity were affected, since they are the underlying physiological mechanisms of

those cognitive capabilities. To pair it with the behavioral tasks, electrophysiology was

carried out 1 to 17 days post-*icv.* injection of A$\beta_{1\text{-}42}$ or controls: A$\beta_{42\text{-}1}$ and vehicle in a

new cohort of mice.

Firstly, I/O curves in all groups (Figures 5A-C) showed a greater amplitude of

both, the first (vehicle male, n = 7; vehicle female, n = 7; A$\beta_{1\text{-}42}$ male, n = 8; A$\beta_{1\text{-}42}$

female, n = 5; A$\beta_{42\text{-}1}$ male, n = 5; A$\beta_{42\text{-}1}$ female, n = 4; Male: $F_{(2.18,159.2)} = 462.3$,

$p < 0.001$; Female: $F_{(2.28,157.3)} = 470.1$, $p < 0.001$, Geisser-Greenhouse's correction) and

the second fEPSP (Male: $F_{(2.221,126.6)} = 242.2$, $p < 0.001$; Female: $F_{(2.259,81.31)} = 234.6$,

$p < 0.001$, Geisser-Greenhouse's correction) with increasing intensities. No differences

between groups were observed for either 1[st] or 2[nd] fEPSP in female slices, however in

male slices a treatment effect was found in fEPSP2 ($F_{(2,57)} = 4.297$, $p = 0.0183$) since

amplitudes evoked by the second pulse in A$\beta_{1\text{-}42}$ injected animals were higher than

those evoked in the other sex-matched groups.

Then, we addressed the short-term plasticity phenomenon, PPF. This protocol

is also related to neurotransmitter release and, therefore, allowed us to evaluate the

presynaptic functionality after A$\beta_{1\text{-}42}$ injection. As shown in Figure 5F, all groups

presented an increased response to the second pulse when the intervals were short

(20, 40 and 100 ms) since the ratio between the second and first EPSPs were above

100%. Nonetheless, treatment caused no significant differences at any of the selected

intervals (vehicle male, n = 6; vehicle female, n = 6; A$\beta_{1\text{-}42}$ male, n = 7; A$\beta_{1\text{-}42}$ female,

n = 5; A$\beta_{42\text{-}1}$ male, n = 3; A$\beta_{42\text{-}1}$ female, n = 3; Male: F$_{(2,44)}$ = 2.930, p = 0.0639; Female:

F$_{(2,39)}$ = 1.159, p = 0.3243). This data indicated a normal short-term plasticity and

presynaptic vesicle release after treatment, suggesting that alterations caused by A$\beta_{1\text{-}42}$

injection could be preferentially acting at a postsynaptic level.

Finally, we measured the effect of A$\beta_{1\text{-}42}$ injection on long-term synaptic

plasticity applying an HFS protocol after a 15-min baseline from day 1 to 17 post-

icv. injection (Figure 5G). Data showed a significant treatment effect (Figure 5H;

vehicle male, n = 6; vehicle female, n = 9; A$\beta_{1\text{-}42}$ male, n = 11; A$\beta_{1\text{-}42}$ female, n = 9;

A$\beta_{42\text{-}1}$ male, n = 10; A$\beta_{42\text{-}1}$ female, n = 10; Male: F$_{(2.78,159.2)}$ = 22.87, p < 0.001; Female:

F$_{(2,81)}$ = 17.67, p < 0.001). *Post-hoc* analysis revealed that the differences were

specifically between the A$\beta_{1\text{-}42}$ and the two control groups, regardless the sex, which

showed an inhibited LTP, right after the HFS and during at least 60 min following it. In

fact, fEPSPs post-HFS were slightly under the BL for this group, suggesting that a

protocol aimed to cause LTP was inducing an LTD instead. As it happened in the

memory assays, the control reverse peptide did not affect LTP, implying the specificity

of A$\beta_{1\text{-}42}$.

Thus, this data indicates that $A\beta_{1-42}$ injection alters long-term synaptic plasticity,

mainly at a postsynaptic level, which could be underlying the hippocampal-dependent

learning and memory deficits observed in our study in both male and female animals.

420 Discussion

The deleterious effect of $A\beta$ - one of the main neuropathological features of

AD -, on hippocampal-dependent learning and memory has been widely reported (14).

However, the possible differences associated to sex are scarcely explored, despite

numerous clinical and epidemiological studies have shown greater prevalence, risk and

severity of AD in women (32). Moreover, animal studies, mainly carried out with

transgenic mice models, have reported contradictory results (34, 37). Here, we aimed

to elucidate whether *icv.* administration of $A\beta_{1-42}$, a local model to study early

hippocampal amyloidosis (49, 50), had a differential effect on hippocampal function in

female and male mice.

Spatial navigation deficit is an early indicator of AD (51). Furthermore, it has

been used to differentiate between various types of dementia, i.e., AD and

frontotemporal dementia (52), since AD shows specific deficits in spatial working

memory compared to others (53). Here, we used the Barnes maze to assess spatial

working, short-term and long-term memory, and the OF habituation test to evaluate a

non-associative hippocampal-dependent memory process. Our results showed a

similar impairment in both male and female mice in all forms of spatial learning and

memory evaluated as well as in the exploratory habituation. It is well stablished that

learning and memory deterioration occurs in both, AD patients and animal models (24,

54). Notwithstanding, transgenic models mostly reflect genetic forms of AD and require

months to develop the cognitive impairments, whereas AD is mainly a sporadic

disorder (55). Moreover, it has been highlighted that AD's prodromal period lasts from 5

years to even decades (25), with the first cognitive symptom appearing as early as 12

years before dementia onset (56), so it is greatly important to study early stages of the

pathology. In agreement with that, A*β icv.* administration mimics the sporadic form of

AD during early stages. Previous works from our group and others have shown that,

even the early amyloidosis caused by a single A*β* injection is sufficient to impair short-

and long-term spatial and habituation memory in male rodents (21-23, 29, 30, 53, 57).

Moreover, the present study proves that working spatial memory is also altered, as it

has been shown in transgenic mice models at later stages of the disease (20, 35).

However, this is of special interest in early amyloidosis since working memory deficits

can be used to predict which patients with mild cognitive impairment (MCI), a

prodromal stage of AD, will eventually develop this type of dementia and the severity of

the cognitive decline (58, 59). What is more, the dorsal hippocampus area is implicated

in spatial processing, while the ventral hippocampus is related to anxiety-related

behaviors (28, 60). Thus, our data show an impairment in spatial learning and memory

without affecting anxiety and depression-like behaviors, proving once again that the *icv.*

administration of A*β* affect mainly the dorsal hippocampus area (22).

On the other hand, our data did not show any difference between female and

male mice neither in the Barnes maze nor in the OF habituation test. Some authors

had described similar results, showing no sex-dependent differences in transgenic

models (37, 38). Other authors, however, had described the opposite, reporting that

females showed more spatial memory deficits than age-matched males also in

transgenic mice (33, 35). Nonetheless, those mice displayed prominent amyloid

plaques as well as neurofibrillary tangles (34) while in our model of early amyloidosis,

A*β* is in its soluble form and not forming senile plaques yet. Thus, it could be

hypothesized that in early stages of amyloidosis there is not a sex-dependent

alteration, whereas in older animals, with a clearer developed AD pathogenesis, sex

becomes an important factor of disease severity. In this line, 3xTg-AD male and female

mice, a widely used model of AD, displayed no memory differences at 4 to 10 months

of age, but female mice performed worse than males at older ages (20, 61, 62).

Interestingly, young females seem to be protected against Aβ toxicity by estrogen, as it

promotes a non-amyloidogenic metabolism of APP and has anti-inflammatory

properties (32). Later on, when females reach menopause, and therefore estrogen

levels decline, this protection is lost, which might underlie the greater susceptibility of

women to suffer AD (33). In fact, human studies have reported a negative correlation

between estrogen levels and spatial cognition (63). Indeed, estrogen treatment in a

menopause-induced 5xFAD mice (another AD mice model) decreases APP and

phosphorylated *tau* levels, compared to untreated 5xFAD females (64).

Regarding activity-dependent synaptic plasticity mechanisms underlying

learning and memory, our data indicates that a single A$\beta_{1\text{-}42}$ injection does not alter

short-term plasticity nor presynaptic vesicle release but does affect long-term plasticity

at postsynaptic level. Several studies concur with the fact that AD pathology affects

mainly at postsynaptic level (65). Early deficits in LTP are associated with the

accumulation of Aβ in the hippocampus of transgenic models, and this weakening in

synaptic strength is associated with an inability to use cues in a spatial learning task

(66). The characteristic hyperexcitability caused by Aβ in the dorsal hippocampus might

be increasing the threshold of LTP induction which, following the Bienenstock, Cooper

and Munro (BCM) theory, would provoke the induction of LTD since the stimulation fails

to properly activate postsynaptic neurons (67). Moreover, Aβ causes an indirect partial

unblocking of the synaptic NMDA receptors (68), known to make an essential

contribution to spatial working memory processing (69), and activates metabotropic

glutamate receptors (70), which result in increased internalization of AMPA receptors,

shifting the signaling cascades toward pathways involved in the induction of LTD and

synaptic loss (29, 71, 72). This, along with the BCM theory, could partially explain that

a protocol aimed to cause LTP induced LTD instead in our female and male A$\beta_{1\text{-}42}$

treated animals. In this line, previous work also showed an HFS-induced LTD both *in*

vivo (23) and *ex vivo* (22, 53) in male mice after A$\beta_{1\text{-}42}$ *icv.* injection. This unbalance in

hippocampal synaptic plasticity processes is underlying the hippocampal-dependent

learning and memory deficits observed both in male and female animals. Interestingly,

LTD, which seems to be increased in AD, is mainly associated with habituation to a

novel environment (73), a non-associative type of learning showed to be altered in our

mice. This shift in the excitability threshold for LTP/LTD induction caused by Aβ has

been shown to be partially governed by G-protein-gated inwardly rectifying potassium

(GIRK) channels (22, 43), which main role in the hippocampus is to maintain the

inhibitory tone. Indeed, it has been found that their activation with the selective opener

ML297 is able to rescue LTP and associated learning and memory processes in this

model of early AD amyloidosis (22, 23, 30). Additionally, GIRK2 subunit expression is

down-regulated in different models of AD (74, 75) and training in an hippocampal

dependent task normalizes its protein level (75). Once again, no differences between

male and female mice were found, as expected since there were no sex-dependent

changes in memory assessments. Remarkably, our synaptic plasticity results correlate

with our behavioral memory experiments, as LTP impairment lasted the same time as

memory deficits were observed (i.e., up to 17 days after *icv.* injection) which highlights

the importance of this Aβ *icv.* administration model to study early acute-induced AD up

to 2 weeks (49, 50). In this line, other authors have shown a progressive degeneration

after A$\beta_{1\text{-}42}$ intrahippocampal administration that last up to a month (76-78), and *icv.*

administration of Aβ mainly diffuses to the dorsal hippocampus (22).

Perspectives and significance

Despite the growing number of publications studying the sex-related differences

underlying the pathogenesis of AD, data regarding females remains inconsistent. In

this regard, our results provide systematic evidence of similar hippocampal deficits

caused by A$\beta_{1\text{-}42}$ regardless the sex, validating this murine model of early amyloidosis.

This work opens numerous research perspectives for the future study of AD

amyloid-related pathogenesis and treatment in both female and male subjects.

526 Conclusions

In summary, our results indicate that a single A$\beta_{1\text{-}42}$ *icv*. injection leads to similar

and robust habituation and spatial working, short- and long-term memory impairments

as well as paired LTP inhibition and LTD facilitation in both sexes. Furthermore, the

cognitive and synaptic alterations were long-lasting (observed up to 17 days after

treatment), evidencing the convenience of using this *in vivo* mouse model to study

early acute stages of amyloidosis regardless the experimental subject's sex.

Declarations

Ethics approval

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by MCIN/AEI/10.13039/501100011033 (grant numbers PID2020-115823-GB100) and JCCM and ERDF A way of making Europe (grant number SBPLY/21/180501/000150) to JDNL and LJD. AC held a *Margarita Salas* Postdoctoral Research Fellow funded by European Union NextGenerationEU/PRTR. DJ and GIL held predoctoral fellowships granted by UCLM/ESF Plan Propio de Investigación Programme

Authors' contributions

LJD and JDNL were responsible for the initial conceptualization; RJH, AC, GIL and DJ performed the experiments; RJH, AC, GIL and SD analyzed the data; RJH and AC were responsible for writing the original draft; RJH, AC, SD, JDNL and LJD did the writing – review and editing. LJD and JDNL were responsible for funding acquisition. All authors read and approved the final manuscript.

560   Acknowledgements

561   Not applicable.

References

1.	Barnes CA. Memory deficits associated with senescence: a neurophysiological and behavioral study in the rat. J Comp Physiol Psychol. 1979;93(1):74-104.

2.	Carrozzo M, Koch G, Turriziani P, Caltagirone C, Carlesimo GA, Lacquaniti F. Integration of cognitive allocentric information in visuospatial short-term memory through the hippocampus. Hippocampus. 2005;15(8):1072-84.

3.	Eichenbaum H. Hippocampus: cognitive processes and neural representations that underlie declarative memory. Neuron. 2004;44(1):109-20.

4.	Scoville WB, Milner B. Loss of recent memory after bilateral hippocampal lesions. 1957. J Neuropsychiatry Clin Neurosci. 1957;12(1):103-13.

5.	Squire LR. Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. Psychol Rev. 1992;99(2):195-231.

6.	Buzsáki G, Moser EI. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. Nat Neurosci. 2013;16(2):130-8.

7.	Spellman T, Rigotti M, Ahmari SE, Fusi S, Gogos JA, Gordon JA. Hippocampal-prefrontal input supports spatial encoding in working memory. Nature. 2015;522(7556):309-14.

8.	Gholipour P, Komaki A, Parsa H, Ramezani M. Therapeutic Effects of High-Intensity Interval Training Exercise Alone and Its Combination with Ecdysterone Against Amyloid Beta-Induced Rat Model of Alzheimer's Disease: A Behavioral, Biochemical, and Histological Study. Neurochem Res. 2022;47(7):2090-108.

9.	Eikelboom WS, Pan M, Ossenkoppele R, Coesmans M, Gatchel JR, Ismail Z, et al. Sex differences in neuropsychiatric symptoms in Alzheimer's disease dementia: a meta-analysis. Alzheimers Res Ther. 2022;14(1):48.

10.	Goodman J, Packard MG. Memory Systems and the Addicted Brain. Front Psychiatry. 2016;7:24.

11.	Gauthier S, Rosa-Neto P, Morais JA, Webster C. World Alzheimer Report 2021: Journey through the diagnosis of dementia.; 2021.

12.	Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. EMBO Mol Med. 2016;8(6):595-608.

13.	Jeremic D, Jiménez-Díaz L, Navarro-López JD. Past, present and future of therapeutic strategies against amyloid-β peptides in Alzheimer's disease: a systematic review. Ageing Res Rev. 2021;72:101496.

14.	Vyas Y, Montgomery JM, Cheyne JE. Hippocampal Deficits in Amyloid-β-Related Rodent Models of Alzheimer's Disease. Front Neurosci. 2020;14:266.

15.	Palop JJ, Mucke L. Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. Nat Neurosci. 2010;13(7):812-8.

16.	Mucke L, Selkoe DJ. Neurotoxicity of amyloid β-protein: synaptic and network dysfunction. Cold Spring Harb Perspect Med. 2012;2(7):a006338.

17.	Kent SA, Spires-Jones TL, Durrant CS. The physiological roles of tau and Aβ: implications for Alzheimer's disease pathology and therapeutics. Acta Neuropathol. 2020;140(4):417-47.

18.	Bliss TVP, Collingridge GL, Morris RGM, Reymann KG. Long-term potentiation in the hippocampus: discovery, mechanisms and function. Neuroforum. 2018;24(3):A103-A20.

19.	Mango D, Saidi A, Cisale GY, Feligioni M, Corbo M, Nisticò R. Targeting Synaptic Plasticity in Experimental Models of Alzheimer's Disease. Front Pharmacol. 2019;10:778.

20.	Stevens LM, Brown RE. Reference and working memory deficits in the 3xTg-AD mouse between 2 and 15-months of age: a cross-sectional study. Behav Brain Res. 2015;278:496-505.

21.	Mayordomo-Cava J, Iborra-Lázaro G, Djebari S, Temprano-Carazo S, Sánchez-Rodríguez I, Jeremic D, et al. Impairments of Synaptic Plasticity Induction Threshold

and Network Oscillatory Activity in the Hippocampus Underlie Memory Deficits in a
Non-Transgenic Mouse Model of Amyloidosis. Biology (Basel). 2020;9(7).

22.   Sanchez-Rodriguez I, Djebari S, Temprano-Carazo S, Vega-Avelaira D,
Jimenez-Herrera R, Iborra-Lazaro G, et al. Hippocampal long-term synaptic depression
and memory deficits induced in early amyloidopathy are prevented by enhancing G-
protein-gated inwardly rectifying potassium channel activity. J Neurochem.
2020;153(3):362-76.

23.   Sánchez-Rodríguez I, Temprano-Carazo S, Nájera A, Djebari S, Yajeya J,
Gruart A, et al. Activation of G-protein-gated inwardly rectifying potassium (Kir3/GirK)
channels rescues hippocampal functions in a mouse model of early amyloid-β
pathology. Sci Rep. 2017;7(1):14658.

24.   Sanchez-Varo R, Mejias-Ortega M, Fernandez-Valenzuela JJ, Nuñez-Diaz C,
Caceres-Palomo L, Vegas-Gomez L, et al. Transgenic Mouse Models of Alzheimer's
Disease: An Integrative Analysis. Int J Mol Sci. 2022;23(10).

25.   Braak H, Thal DR, Ghebremedhin E, Del Tredici K. Stages of the pathologic
process in Alzheimer disease: age categories from 1 to 100 years. J Neuropathol Exp
Neurol. 2011;70(11):960-9.

26.   Lührs T, Ritter C, Adrian M, Riek-Loher D, Bohrmann B, Döbeli H, et al. 3D
structure of Alzheimer's amyloid-β(1–42) fibrils. Proceedings of the
National Academy of Sciences. 2005;102(48):17342-7.

27.   Fanselow MS, Dong HW. Are the dorsal and ventral hippocampus functionally
distinct structures? Neuron. 2010;65(1):7-19.

28.   Moser MB, Moser EI, Forrest E, Andersen P, Morris RG. Spatial learning with a
minislab in the dorsal hippocampus. Proc Natl Acad Sci U S A. 1995;92(21):9697-701.

29.   Morroni F, Sita G, Tarozzi A, Rimondini R, Hrelia P. Early effects of Aβ1-42
oligomers injection in mice: Involvement of PI3K/Akt/GSK3 and MAPK/ERK1/2
pathways. Behav Brain Res. 2016;314:106-15.

30.   Sanchez-Rodriguez I, Gruart A, Delgado-Garcia JM, Jimenez-Diaz L, Navarro-
Lopez JD. Role of GirK Channels in Long-Term Potentiation of Synaptic Inhibition in an
In Vivo Mouse Model of Early Amyloid-beta Pathology. Int J Mol Sci. 2019;20(5).

31.   Alzheimer's_Association. 2022 Alzheimer's disease facts and figures.
Alzheimer's & Dementia. 2022;18(4):700-89.

32.   Viña J, Lloret A. Why women have more Alzheimer's disease than men: gender
and mitochondrial toxicity of amyloid-beta peptide. J Alzheimers Dis. 2010;20 Suppl
2:S527-33.

33.   Yang JT, Wang ZJ, Cai HY, Yuan L, Hu MM, Wu MN, et al. Sex Differences in
Neuropathology and Cognitive Behavior in APP/PS1/tau Triple-Transgenic Mouse
Model of Alzheimer's Disease. Neurosci Bull. 2018;34(5):736-46.

34.   Pang K, Jiang R, Zhang W, Yang Z, Li LL, Shimozawa M, et al. An App knock-
in rat model for Alzheimer's disease exhibiting Aβ and tau pathologies, neuronal death
and cognitive impairments. Cell Res. 2022;32(2):157-75.

35.   Bernaud VE, Bulen HL, Peña VL, Koebele SV, Northup-Smith SN, Manzo AA,
et al. Task-dependent learning and memory deficits in the TgF344-AD rat model of
Alzheimer's disease: three key timepoints through middle-age in females. Sci Rep.
2022;12(1):14596.

36.   Forner S, Kawauchi S, Balderrama-Gutierrez G, Kramar EA, Matheos DP, Phan
J, et al. Systematic phenotyping and characterization of the 5xFAD mouse model of
Alzheimer's disease. Sci Data. 2021;8(1):270.

37.   Hulshof LA, Frajmund LA, van Nuijs D, van der Heijden DCN, Middeldorp J, Hol
EM. Both male and female APPswe/PSEN1dE9 mice are impaired in spatial memory
and cognitive flexibility at 9 months of age. Neurobiol Aging. 2022;113:28-38.

38.   Cohen RM, Rezai-Zadeh K, Weitz TM, Rentsendorj A, Gate D, Spivak I, et al. A
transgenic Alzheimer rat with plaques, tau pathology, behavioral impairment,
oligomeric aβ, and frank neuronal loss. J Neurosci. 2013;33(15):6245-56.

39.   Paxinos G, Franklin KBJ. The Mouse Brain in Stereotaxic Coordinates: Elsevier
Academic Press; 2004.
40.   Gawel K, Gibula E, Marszalek-Grabska M, Filarowska J, Kotlinska JH.
Assessment of spatial learning and memory in the Barnes maze task in rodents-
methodological   consideration.   Naunyn   Schmiedebergs   Arch   Pharmacol.
2019;392(1):1-18.
41.   Suarez AN, Hsu TM, Liu CM, Noble EE, Cortella AM, Nakamoto EM, et al. Gut
vagal sensory signaling regulates hippocampus function through multi-order pathways.
Nat Commun. 2018;9(1):2181.
42.   Leussis MP, Bolivar VJ. Habituation in rodents: a review of behavior,
neurobiology, and genetics. Neurosci Biobehav Rev. 2006;30(7):1045-64.
43.   Djebari S, Iborra-Lázaro G, Temprano-Carazo S, Sánchez-Rodríguez I, Nava-
Mesa MO, Múnera A, et al. G-Protein-Gated Inwardly Rectifying Potassium (Kir3/GIRK)
Channels Govern Synaptic Plasticity That Supports Hippocampal-Dependent Cognitive
Functions in Male Mice. J Neurosci. 2021;41(33):7086-102.
44.   Kalueff AV, Tuohimaa P. Grooming analysis algorithm for neurobehavioural
stress research. Brain Res Brain Res Protoc. 2004;13(3):151-8.
45.   Büttner D. Climbing on the cage lid, a regular component of locomotor activity in
the mouse. J Exp Anim Sci. 1991;34(5-6):165-9.
46.   La-Vu M, Tobias BC, Schuette PJ, Adhikari A. To Approach or Avoid: An
Introductory Overview of the Study of Anxiety Using Rodent Assays. Front Behav
Neurosci. 2020;14:145.
47.   Lopes BC, Medeiros LF, Stein DJ, Cioato SG, de Souza VS, Medeiros HR, et
al. tDCS and exercise improve anxiety-like behavior and locomotion in chronic pain rats
via modulation of neurotrophins and inflammatory mediators. Behav Brain Res.
2021;404:113173.
48.   Cryan JF, Mombereau C, Vassout A. The tail suspension test as a model for
assessing antidepressant activity: review of pharmacological and genetic studies in
mice. Neurosci Biobehav Rev. 2005;29(4-5):571-625.
49.   Kim HY, Lee DK, Chung BR, Kim HV, Kim Y. Intracerebroventricular Injection of
Amyloid-β Peptides in Normal Mice to Acutely Induce Alzheimer-like Cognitive Deficits.
J Vis Exp. 2016(109).
50.   Kasza Á, Penke B, Frank Z, Bozsó Z, Szegedi V, Hunya Á, et al. Studies for
Improving a Rat Model of Alzheimer's Disease: Icv Administration of Well-
Characterized β-Amyloid 1-42 Oligomers Induce Dysfunction in Spatial Memory.
Molecules. 2017;22(11):2007.
51.   Lithfous S, Dufour A, Després O. Spatial navigation in normal aging and the
prodromal stage of Alzheimer's disease: insights from imaging and behavioral studies.
Ageing Res Rev. 2013;12(1):201-13.
52.   Tu S, Wong S, Hodges JR, Irish M, Piguet O, Hornberger M. Lost in spatial
translation - A novel tool to objectively assess spatial disorientation in Alzheimer's
disease and frontotemporal dementia. Cortex. 2015;67:83-94.
53.   Torres-Flores M, Peña-Ortega F. Amyloid Beta Alters Prefrontal-dependent
Functions Along with its Excitability and Synaptic Plasticity in Male Rats. Neuroscience.
2022;498:260-79.
54.   Ma Y, Brettschneider J, Collingwood JF. A Systematic Review and Meta-
Analysis of Cerebrospinal Fluid Amyloid and Tau Levels Identifies Mild Cognitive
Impairment Patients Progressing to Alzheimer's Disease. Biomedicines. 2022;10(7).
55.   LaFerla FM, Oddo S. Alzheimer's disease: Abeta, tau and synaptic dysfunction.
Trends Mol Med. 2005;11(4):170-6.
56.   Amieva H, Le Goff M, Millet X, Orgogozo JM, Pérès K, Barberger-Gateau P, et
al. Prodromal Alzheimer's disease: successive emergence of the clinical symptoms.
Ann Neurol. 2008;64(5):492-8.
57.   Duran-Gonzalez J, Michi ED, Elorza B, Perez-Cordova MG, Pacheco-Otalora
LF, Touhami A, et al. Amyloid beta peptides modify the expression of antioxidant repair

enzymes and a potassium channel in the septohippocampal system. Neurobiol Aging.
2013;34(8):2071-6.
58. Huntley JD, Howard RJ. Working memory in early Alzheimer's disease: a
neuropsychological review. Int J Geriatr Psychiatry. 2010;25(2):121-32.
59. Pillai JA, Bonner-Jackson A, Walker E, Mourany L, Cummings JL. Higher
working memory predicts slower functional decline in autopsy-confirmed Alzheimer's
disease. Dement Geriatr Cogn Disord. 2014;38(3-4):224-33.
60. Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the
hippocampal longitudinal axis. Nat Rev Neurosci. 2014;15(10):655-69.
61. Halagappa VK, Guo Z, Pearson M, Matsuoka Y, Cutler RG, Laferla FM, et al.
Intermittent fasting and caloric restriction ameliorate age-related behavioral deficits in
the triple-transgenic mouse model of Alzheimer's disease. Neurobiol Dis.
2007;26(1):212-20.
62. Clinton LK, Billings LM, Green KN, Caccamo A, Ngo J, Oddo S, et al. Age-
dependent sexual dimorphism in cognition and stress response in the 3xTg-AD mice.
Neurobiol Dis. 2007;28(1):76-82.
63. Locklear MN, Kritzer MF. Assessment of the effects of sex and sex hormones
on spatial cognition in adult rats using the Barnes maze. Horm Behav. 2014;66(2):298-
308.
64. Kim JY, Mo H, Kim J, Kim JW, Nam Y, Rim YA, et al. Mitigating Effect of
Estrogen in Alzheimer's Disease-Mimicking Cerebral Organoid. Front Neurosci.
2022;16:816174.
65. Zhang H, Jiang X, Ma L, Wei W, Li Z, Chang S, et al. Role of Aβ in Alzheimer's-
related synaptic dysfunction. Front Cell Dev Biol. 2022;10:964075.
66. Privitera L, Hogg EL, Lopes M, Domingos LB, Gaestel M, Müller J, et al. The
MK2 cascade mediates transient alteration in mGluR-LTD and spatial learning in a
murine model of Alzheimer's disease. Aging Cell. 2022:e13717.
67. Keck T, Hubener M, Bonhoeffer T. Interactions between synaptic homeostatic
mechanisms: an attempt to reconcile BCM theory, synaptic scaling, and changing
excitation/inhibition balance. Curr Opin Neurobiol. 2017;43:87-93.
68. Liu J, Chang L, Song Y, Li H, Wu Y. The Role of NMDA Receptors in
Alzheimer's Disease. Front Neurosci. 2019;13:43.
69. McHugh SB, Niewoehner B, Rawlins JN, Bannerman DM. Dorsal hippocampal
N-methyl-D-aspartate receptors underlie spatial working memory performance during
non-matching to place testing on the T-maze. Behav Brain Res. 2008;186(1):41-7.
70. Campos-Peña V, Meraz-Ríos MA. Alzheimer Disease: The Role of Aβ in the
Glutamatergic System. Rijeka: IntechOpen; 2014. p. Ch. 10.
71. Wiatrak B, Mieszała P, Gąsiorowski K. Impact of NMDA receptor activation on
DNA damage in PC12 neuron-like cell cultures in the presence of β-amyloid peptides.
Mol Biol Rep. 2022.
72. Liu L, Wong TP, Pozza MF, Lingenhoehl K, Wang Y, Sheng M, et al. Role of
NMDA receptor subtypes in governing the direction of hippocampal synaptic plasticity.
Science. 2004;304(5673):1021-4.
73. Collingridge GL, Peineau S, Howland JG, Wang YT. Long-term depression in
the CNS. Nat Rev Neurosci. 2010;11(7):459-73.
74. Alfaro-Ruiz R, Martín-Belmonte A, Aguado C, Hernández F, Moreno-Martínez
AE, Ávila J, et al. The Expression and Localisation of G-Protein-Coupled Inwardly
Rectifying Potassium (GIRK) Channels Is Differentially Altered in the Hippocampus of
Two Mouse Models of Alzheimer's Disease. Int J Mol Sci. 2021;22(20).
75. Temprano-Carazo S, Contreras A, Saura CA, Navarro-López JD, Jiménez-Díaz
L. Spatial Memory Training Counteracts Hippocampal GIRK Channel Decrease in the
Transgenic APPSw,Ind J9 Alzheimer’s Disease Mouse Model. International
Journal of Molecular Sciences. 2022;23(21):13444.
76. Calvo-Flores Guzmán B, Elizabeth Chaffey T, Hansika Palpagama T, Waters S,
Boix J, Tate WP, et al. The Interplay Between Beta-Amyloid 1-42 (Aβ(1-42))-Induced

Hippocampal Inflammatory Response, p-tau, Vascular Pathology, and Their Synergistic Contributions to Neuronal Death and Behavioral Deficits. Front Mol Neurosci. 2020;13:522073.

77.   Durán-González J, Michi ED, Elorza B, Perez-Córdova MG, Pacheco-Otalora LF, Touhami A, et al. Amyloid β peptides modify the expression of antioxidant repair enzymes and a potassium channel in the septohippocampal system. Neurobiol Aging. 2013;34(8):2071-6.

78.   Faucher P, Mons N, Micheau J, Louis C, Beracochea DJ. Hippocampal Injections of Oligomeric Amyloid β-peptide (1-42) Induce Selective Working Memory Deficits and Long-lasting Alterations of ERK Signaling Pathway. Front Aging Neurosci. 2015;7:245.

792 Figures

Figure 1. Experimental design showing the timeline and corresponding figure numbers. (A) A stainless-steel guide cannula was implanted for *icv.* drug administration on the left ventricle, as shown in the histologic verification image on the right. A minimum of 8 days after surgery (day 0), either Aβ$_{1-42}$ or controls: reverse Aβ$_{42-1}$ peptide or vehicle, were administered *icv.* Scale bars: 500 mm. **(B)** The Barnes maze task habituation phase was carried out the day before *icv.* administration (day -1). One hour after *icv.* injection of the corresponding treatment (day 0), training in the Barnes maze started. The training phase consisted of 3 trails per day, during 4 consecutive days (days 0 to 3). Two memory tests were applied, on days 7 and 11 post-injection. **(C)** The Open field habituation test was carried out on days 15 and 16 post injection. **(D)** Finally, mice went through a test battery to assess their overall spontaneous behavior and health, that consisted of testing of stereotyped and locomotor behaviors by an automated LABORAS® System, as well as the rotarod test, the elevated plus maze and the tail suspension test (days 15 to 17). **(E)** Another cohort of animals was used to evaluate the effect of Aβ$_{1-42}$ on *ex vivo* hippocampal LTP by multi-electrode arrays (MEAs) electrophysiology (days 1 to 17 post-injection). Representative location of stimulation (Stm, green) and recording (Rec, red) electrodes in a hippocampal

811   coronal slice. D, dorsal; DG, dentate gyrus; *icv.*, intracerebroventricular; L, lateral; LV,

812   lateral ventricle; M, medial.

813
Figure 2. **Aβ₁₋₄₂ impairs spatial learning and memory in both female and male**
mice. (A) Representative traces of the path traveled during the first and last trial of the
last training day (day 3) for Aβ$_{1-42}$ and controls: vehicle and reverse Aβ$_{42-1}$ peptide. **(B-**
D) Escape latency (**B**; in s), number of errors (**C**) and distance traveled (**D**; in cm)
during the four training days. **(E)** Overview image of the test phase on the Barnes
maze, with all holes closed. **(F)** Latency (in s) to reach the target hole of the latest
training day for the first time, during the two test sessions. **(G)** Distance traveled (in cm)
during the two test sessions. Data is expressed as the mean ± SEM of the 3 trials per
day. Aβ, Amyloid-β; cm, centimeters; s, seconds. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$
vs. vehicle of the corresponding sex; # $p < 0.05$, ## $p < 0.01$ *vs.* Aβ$_{42-1}$ of the
corresponding sex.

Figure 3. **Aβ₁₋₄₂ administration similarly alters non-associative exploratory**
habituation memory in both female and male mice. (A) An OF habituation test was
carried out by submitting the animals to the same OF arena twice, on consecutive days
15 and 16 post-*icv.* injection. **(B)** Total distance traveled during the two OF sessions
(training -OF1- and retention -OF2- sessions). Data is expressed as the percentage
(%) of the distance traveled in the training session (OF1). **(C)** Examples of mice
movement tracked during OF1 and OF2 for the different treatment groups. Aβ,
Amyloid-β; *icv.*, intracerebroventricular; OF, open field. * $p < 0.05$, ** $p < 0.01$ *vs.*
vehicle of the corresponding sex; # $p < 0.05$, ## $p < 0.01$ *vs.* A$\beta_{42\text{-}1}$ of the
corresponding sex; ‡ $p < 0.05$, ‡‡ $p < 0.01$, ‡‡‡ $p < 0.001$ *vs.* OF1.

Figure 4. **A*β*₁₋₄₂ administration does not induce alterations in locomotor activity,**
anxiety, and depression-like behavior. Behavioral tasks to evaluate general health
state were carried out on days 15 - 17 post-*icv.* injection. **(A)** Stereotyped behaviors
were assessed using a LABORAS® system, measuring the time (in s) spent performing
each type of activity (locomotion, climbing, rearing, and grooming). **(B)** Latency (in s) to
fall off the Rotarod during the six trials (left) and the whole session (right). **(C)** Number
of entries in closed and total arms were used as measure of locomotor activity (left),
while anxiety levels were assessed by the percentage (%) of entries and time spent on
open arms in an elevated plus maze (right). **(D)** Depression-like behavior was
assessed by measuring the immobility time (in s) during a single session in the tail
suspension test. A*β*, Amyloid-*β*; s, seconds.

Figure 5. **A$\beta_{1\text{-}42}$ inhibits ex vivo hippocampal LTP and induces LTD in both female and male mice. (A-C)** I/O curve with paired fEPSPs collected at increasing stimulus intensities (from 0.075 to 0.4 mA) from control vehicle **(A)**, A$\beta_{1\text{-}42}$ **(B)** and A$\beta_{42\text{-}1}$ reverse control **(C)** slices, respectively. Data is expressed as a percentage (%) of the maximum amplitude obtained. **(D)** PPF curve with paired fEPSPs collected at interstimulus intervals of 10, 20, 40, 100, 200 and 500 ms. Data is expressed as mean ± SEM amplitude of the second fEPSP as a percentage of the first [(second/first) x 100] for each inter-pulse interval used. **(E)** Representative averaged (n = 5) traces of fEPSPs recorded in the CA1 area, collected during the baseline (1) and ≈50 min post-HFS (2)

in hippocampal slices from the different groups. Left traces are from female mice and

right traces from male mice. **(F)** Time course of LTP evoked in the CA1 area after HFS

in hippocampal slice from the different groups. Recordings were obtained from day 1 to

17 post-*icv.* injection. Aβ, amyloid-β; HFS, High frequency stimulation; LTP, long-term

potentiation; mA, milliamperes; ms, milliseconds; min, minutes. *** $p < 0.001$ *vs.*

vehicle of the corresponding sex; ## $p < 0.01$ *vs.* A$\beta_{42\text{-}1}$ of the corresponding sex.