Nap1 and Kap114 co-chaperone H2A-H2B and facilitate targeted histone release in the nucleus

Ho Yee Joyce Fung[1,2], Jenny Jiou[1,3], Ashley B. Neisman[1,2], Natalia E. Bernardes[1], and Yuh Min Chook[1,4]

[1]Department of Pharmacology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

[2]Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

[3]Current address: The Walter and Eliza Hall Institute of Medical Research, 1G, Royal Parade, Parkville, Victoria, 302, Australia.

[4]Correspondence: yuhmin.chook@utsouthwestern.edu

Author Contributions: J.J. expressed, purified, and assembled all protein complexes for AUC, SEC-MALS, SAXS, cryo-EM and MP experiments, designed the experiments, and collected, processed, and interpreted the data. J.J. prepared all the cryo-EM samples. J.J. prepared cryo-EM grids, processed cryo-EM data and solved the structures of the assembly without Ran^{GTP}. H.Y.J.F. prepared cryo-EM grids, processed cryo-EM data and solved the structures of the assembly containing Ran^{GTP} and oversaw deposition of all PDBs. H.Y.J.F. cloned and A.B.N. expressed all MBP-Kap114 and Nap1 (WT and mutants) used in the pull-down, DNA competition and nucleosome assembly assays. A.B.N. purified WT MBP-Kap114, J.J. generated H2A-H2B, N.E.B. generated H3-H4 tetramers, and H.Y.J.F. purified the remaining proteins and DNA used in these assays. H.Y.J.F. designed and performed all pull-down and FP binding assays, DNA competition

and nucleosome assembly assays. Y.M.C. conceptualized the research and supervised

experiments. H.Y.J.F. and Y.M.C. prepared figures and wrote the manuscript with input

from the other authors.

Competing Interest Statement: No competing interest declared.

Classification: Biological Sciences: Biophysics and Computational Biology

Keywords: Karyopherin, Histones, Chaperone, Importin, Nucleosome

This PDF file includes:

Main Text

Figures 1 to 5

Abstract

Core histones are synthesized and processed in the cytoplasm before transport into the nucleus for assembly into nucleosomes; however, they must also be chaperoned as free histones are toxic. The importin Kap114 binds and transports histone H2A-H2B into the yeast nucleus, where Ran^{GTP} facilitates H2A-H2B release. Kap114 and H2A-H2B also bind the Nap1 histone chaperone, which is found in both the cytoplasm and the nucleus, but how Nap1 and Kap114 cooperate in H2A-H2B processing and nucleosome assembly has been unclear. To understand these mechanisms, we used biochemical and structural analyses to reveal how Nap1, Kap114, H2A-H2B and Ran^{GTP} interact. We show that Kap114, H2A-H2B and a Nap1 dimer ($Nap1_2$) assemble into a 1:1:1 ternary complex. Cryogenic electron microscopy revealed two distinct $Kap114/Nap1_2/H2A\text{-}H2B$ structures: one of H2A-H2B sandwiched between $Nap1_2$ and Kap114, and another in which $Nap1_2$ bound to the Kap114•H2A-H2B complex without contacting H2A-H2B. Another $Nap1_2$•H2A-H2B•Kap114•Ran^{GTP} structure reveals the nuclear complex. Mutagenesis revealed shared critical interfaces in all three structures. Consistent with structural findings, DNA competition experiments demonstrated that Kap114 and $Nap1_2$ together chaperone H2A-H2B better than either protein alone. When Ran^{GTP} is present, Kap114's chaperoning activity diminishes. However, the presence of $Nap1_2$ within the $Nap1_2$•H2A-H2B•Kap114•Ran^{GTP} quaternary complex restores its ability to chaperone H2A-H2B. This complex effectively deposits H2A-H2B into nucleosomes. Together, these findings suggest that Kap114 and $Nap1_2$ provide a sheltered path from cytoplasm to nucleus, facilitating the transfer of H2A-H2B from Kap114 to $Nap1_2$, ultimately directing its specific deposition into nucleosomes.

Significance Statement

Free histones are toxic and must be sequestered by other macromolecules in the cell. Nuclear import receptor Kap114 imports H2A-H2B into the nucleus while also chaperoning it. The histone chaperone Nap1 also chaperones H2A-H2B, but it is unclear how Nap1 and Kap114 cooperate to process H2A-H2B. We present biochemical and structural results that explain how Kap114, Nap1 and H2A-H2B assemble in the absence and presence of Ran^{GTP}, how Nap1 and Kap114 co-chaperone H2A-H2B, and how Ran^{GTP} and Nap1 coordinate the transfer of H2A-H2B from Kap114 to assembling nucleosomes in the nucleus.

Introduction

Formation of new nucleosomes to pack newly replicated DNA during the S-phase of the cell cycle begins with rapid synthesis of core histones H3, H4, H2A and H2B followed by their swift transport into the nucleus. Exposure of these very basic polypeptides to the cellular environment is deeply deleterious as they form toxic aggregates easily (1, 2). Therefore, core histones are thought to be always chaperoned and never free (3, 4). As H3 and H4 emerge from translating ribosomes, they are folded into H3-H4 heterodimers by heat shock proteins and folding chaperones, acetylated at several lysine side chains (5-7) and then passed to histone chaperone ASF1 and *H. sapiens* (*Hs*) Importin-4 or its *S. cerevisiae* (*Sc*) homolog Kap123 for transport across the nuclear pore complex (NPC) into the nucleus (8-11). Unfortunately, the steps of H2A-H2B biosynthesis and processing in the cytoplasm have not been delineated. However, selective ribosome profiling studies showed no association of *Sc* importins with the nascent chains of H2A and H2B (12),

suggesting that H2A-H2B may follow a processing path like H3-H4's that involves heat

shock protein mediated folding and association with particular histone chaperones prior to

importin-binding for nuclear import.

The heat shock proteins/folding chaperones for H2A and H2B have not been

identified, but it is well-established that the folded H2A-H2B heterodimer binds the

Nucleosome Assembly Protein 1 (Nap1) histone chaperone in the *Sc* cytoplasm (13-15). In

human cells, H2A-H2B also binds several *Hs* Nap1 homologs (16, 17). The importin

primarily responsible for H2A-H2B nuclear import is also known - *Sc* Karyopherin-114

(Kap114) or the homologous *Hs* Importin-9 (IMP9/IPO9).

Sc Nap1 is mostly localized to the yeast cytoplasm where it chaperones newly

synthesized and folded H2A-H2B (13, 18). It also has well-known roles in the nucleus,

including nucleosome assembly/remodeling, transcription, DNA repair and mitosis (14, 19-

33). Genetic interaction between Nap1 and Kap114 as well as interactions with Ran

binding proteins Yrb1 and Yrb2 have been reported (34). *Sc* Nap1 was also reported to be

a cofactor for H2A-H2B nuclear import by Kap114 (13, 35). Both immunoprecipitation (IP)

and size exclusion chromatography (SEC) of yeast cytosolic extract showed Kap114

association with both Nap1 and H2A-H2B while IP from nuclear extract showed persisting

association of the three proteins in the Ran^{GTP}-rich nucleus (13, 36). Therefore, Kap114,

H2A-H2B and Nap1 proteins likely form a complex in the cytoplasm and remain associated

in the nucleus prior to unloading the histones onto assembling nucleosomes (36). Nap1's

functions in both compartments, its interaction with Kap114 and its mostly cytoplasmic

localization suggest that it may shuttle between the nucleus and the cytoplasm. According

to the Saccharomyces Genome Database (SGD), Nap1, which often forms dimers or

Nap1$_2$, is three-fold more abundant than Kap114 (37). However, there is a lack of

information about its temporal-spatial distribution in the cell, including how much and when

it remains in the cytoplasm, enters the nucleus, or subsequently returns to the cytoplasm.

H2A and H2B contain disordered N- and C-terminal tails and central alpha helices

that fold together into the globular H2A-H2B histone-fold domain (Fig. 1A) (38). The H2A-

H2B domain has very basic surfaces that bind DNA in the nucleosome. Outside of the

nucleosome, these basic surfaces participate in non-specific/aberrant interactions that

produce toxic aggregates unless they are shielded (2). Crystal structures show Nap1$_2$

binding asymmetrically to one H2A-H2B domain to shield its DNA-binding surfaces,

explaining its function as a histone chaperone (22, 39).

The superhelical Kap114 wraps around the H2A-H2B histone-fold domain (Fig. 1B),

occluding the histone's DNA-binding surfaces and thus also acting as a histone chaperone

(40, 41). The Kap114•H2A-H2B complex is unusual in its interactions with the GTPase

Ran^{GTP}. Most importin•cargo complexes are dissociated by Ran^{GTP} but Kap114•H2A-H2B

binds Ran^{GTP} to form a stable Ran^{GTP}•Kap114•H2A-H2B complex, which alters interactions

between Kap114 and H2A-H2B, and facilitates histone transfer to other nuclear

chaperones or to the assembling nucleosome (41).

While we understand how Nap1$_2$ binds H2A-H2B and how Kap114 binds H2A-H2B,

we do not know how the three proteins interact for Nap1$_2$ to act as a co-factor in H2A-H2B

import. We also do not know how Ran^{GTP} interacts with Kap114, Nap1$_2$ and H2A-H2B in

the nucleus or how their interactions affect nucleosome assembly. Here, to address these

questions, we used biochemical analyses and cryo-EM structure determination to

understand how Kap114, Nap1$_2$ and H2A-H2B might assemble in the cytoplasm and

nucleus. We showed by cryogenic electron microscopy (cryo-EM) their assembly into a 1:1:1 Kap114/Nap1$_2$/H2A-H2B ternary complex that adopts two different arrangements, both showing extensive sequestration and occlusion of H2A-H2B DNA-binding interfaces by Kap114 and Nap1$_2$. The structural data is consistent with co-chaperoning indicated by DNA competition assays. Analysis in the presence of Ran^{GTP} to mimic conditions in the nucleus showed the GTPase binding to form a quaternary Nap1$_2$•H2A-H2B•Kap114•Ran^{GTP} complex where H2A-H2B remains sequestered by Kap114 and Nap1$_2$. DNA competition and nucleosome assembly assays confirmed that in the presence of Ran^{GTP}, Kap114 and Nap1$_2$ continue to shield H2A-H2B from non-specific aggregation with DNA while they cooperate to transfer H2A-H2B effectively and specifically into nucleosomes.

Results and Discussion

Kap114 binds simultaneously to a H2A-H2B heterodimer and a Nap1 dimer

Biochemical and structural studies of binary Kap114-H2A-H2B and Nap1$_2$-H2A-H2B interactions, and pull-down studies of Kap114 and Nap1 from yeast lysates have been reported (13, 22, 36, 41). First, we confirmed the previous findings using analytical ultracentrifugation (AUC), size exclusion chromatography multi-angle light scattering (SEC-MALS) and pull-down binding assays with recombinant Kap114, Nap1 and H2A-H2B (SI Appendix, Fig. S1). We showed that Nap1 alone is a mixture of dimers and tetramers (SI Appendix, Fig. S1B), and Nap1 binds H2A-H2B to form a mixture of Nap1 dimer and tetramers bound to one to two copies of H2A-H2B (SI Appendix, Fig. S1C). Nap1 binds Kap114 to form a 1:1 Kap114:Nap1$_2$ complex (SI Appendix, Fig. S1A and B).

Figure 1. Equimolar Kap114, Nap1₂ and H2A-H2B form a 1:1:1 complex but excess histone disrupts the ternary complex. (A) Organization schematics of the Nap1 (cyan), H2A (yellow), H2B (magenta) and Kap114 (gray, long loops labeled) polypeptides. **(B)** The Kap114•H2A-H2B binary complex structure (8F0X). **(C)** Pull-down assay with immobilized MBP-Kap114 (1 μM) and equimolar Nap1₂, ± 1 or 2 molar ratio H2A-H2B; Coomassie-stained SDS-PAGE with quantified Nap1₂ bands on the right. Controls in SI Appendix, Fig. S2. **(D)** SEC-MALS analysis of 1:1:1 (orange) or 1:1:2 (lilac) molar ratio Kap114, Nap1₂ and H2A-H2B mixtures. Left, differential refractive index (dRI, left y-axis, thin lines) and molecular mass (kDa, right y-axis, thick lines) traces and theoretical masses of Kap114•H2A-H2B (KH), Nap1₂•H2A-H2B (N₂H and 2(N₂)2H) and Kap114/Nap1₂/H2A-H2B (K/N₂/H) complexes marked with dashed lines. Middle panel, Coomassie-stained SDS-PAGE of peak fractions. Right, tabulated SEC-MALS results. At 1:1:1 molar ratio, the three proteins assemble into a 1 Kap114/1 Nap1₂/1 H2A-H2B complex. No ternary complex is observed when H2A-H2B is in excess; only Kap114•H2A-H2B, Nap1₂•H2A-H2B and 2Nap1₂•2H2A-H2B binary complexes formed. **(E)** A summary schematic of the results in **(D)**.

150   Next, we focused on interactions between the three proteins. Immobilized MBP-

Kap114 pulled down Nap1 in the absence/presence of H2A-H2B; Kap114 also pulled down H2A-H2B in the absence/presence of Nap1 (Fig. 1C and SI Appendix, Fig. S2A-C). SEC-MALS analysis showed the three proteins forming a 1:1:1 Kap114/Nap1$_2$/H2A-H2B complex (Fig. 1D, and SI Appendix, Fig. S2D). Interestingly, molar excess of H2A-H2B slightly decreased the amount of Nap1 pulled down by MBP-Kap114 beads (Fig. 1C and SI Appendix, Fig. S2B). SEC-MALS analysis of 1:1 Kap114:Nap1$_2$ in the presence of two-fold excess H2A-H2B showed two peaks that match the binary Nap1•H2A-H2B and Kap114•H2A-H2B complexes (Fig. 1D, and SI Appendix, Fig. S1C), consistent with destabilization of the ternary complex seen in the pull-down experiment.

In summary, equimolar Kap114, Nap1$_2$ and H2A-H2B assemble into a stable 1:1:1 ternary complex (Fig. 1E). Excess H2A-H2B destabilizes the ternary complex, dissociating it into binary Kap114•H2A-H2B and Nap1$_2$•H2A-H2B complexes (Fig. 1E). Ternary Kap114/Nap1$_2$/H2A-H2B interactions, such as in the cytoplasm, may be most stable when all H2A-H2B heterodimers are adequately chaperoned.

Structure determination of the Kap114/Nap1$_2$/H2A:H2B ternary complex

Because the Nap1$_2$•H2A-H2B complex likely forms before encountering Kap114 (5, 11), we prepared a cryo-EM sample by first assembling Nap1$_2$ core•H2A-H2B with excess histones to ensure binding, then adding Kap114, immediately followed by mild crosslinking (SI Appendix, Fig. S3A). After SEC, the sample remained heterogenous, a characteristic evident in the cryo-EM dataset that produced ~1 million particles: 38.7% are Nap1$_2$, 36.0% Kap114•H2A-H2B, and 25.3% ternary complex of Kap114, Nap1$_2$ and H2A-H2B (SI Appendix, Fig. S3B). The smaller population of ternary complex is likely due to

destabilization by excess H2A-H2B.

The $Nap1_2$ and Kap114•H2A-H2B particles produced cryo-EM maps/structures that

are very similar to the respective crystal and cryo-EM structures in the Protein Data Bank

or PDB (SI Appendix, Fig. S4 and Table S1) (41). Particles of the ternary

Kap114/$Nap1_2$/H2A-H2B complex were classified into two evenly divided classes that

produced two high-resolution maps and structures (Fig. 2A-F and SI Appendix, Fig. S5 and

Table S1). The first is a 3.2 Å resolution map of a $Nap1_2$ bound to a Kap114•H2A-H2B

complex where $Nap1_2$ contacts the importin but not the histone; we termed this structure

$Nap1_2$•Kap114•H2A-H2B (Fig. 2A and B). The second is a 3.5 Å resolution map of Kap114

and $Nap1_2$ interacting with the same H2A-H2B; we termed the structure $Nap1_2$•H2A-

H2B•Kap114 (Fig. 2C and D). Local resolution of the $Nap1_2$ region in both maps is poor, at

~ 6-9 Å, suggesting dynamic $Nap1_2$ in both structures (SI Appendix, Fig. S5A and E).

Nevertheless, local refinement improved the focused maps at the $Nap1_2$ regions, allowing

us to dock in the histone chaperone. We emphasize the distinct protein arrangements in

the two structures by highlighting 'Kap114' in bold: $Nap1_2$•**Kap114**•H2A-H2B versus

$Nap1_2$•H2A-H2B•**Kap114**.

Two ternary complex structures: the $Nap1_2$•*Kap114*•H2A-H2B structure

One of the two ternary complex structures, $Nap1_2$•**Kap114**•H2A-H2B, shows

Kap114 wrapping around H2A-H2B while $Nap1_2$ contacts only Kap114 (Fig. 2A and SI

Appendix, Fig. S6A and B). The H2A-H2B-bound Kap114 here is very similar to the binary

Kap114•H2A-H2B complex (Fig. 1B and SI Appendix, Fig. S7A); root mean square

deviation (RMSD) of the two structures is ~0.5 Å (41). The extensive Kap114-histone

interactions in the two structures, that bury ~2200 Å^2 of H2A-H2B, are virtually the same

(Figs. 1B, 2B and SI Appendix, Fig. S8).

Figure 2. Cryo-EM structures reveal two distinct configurations of the Kap114/Nap1₂/H2A-H2B complex. (A) Left, the Nap1₂•**Kap114**•H2A-H2B (cyan•gray•yellow-pink) structure with overall/local refined maps (transparent gray/cyan surfaces)**.** The β-hairpin and AS of the Nap1₂ subunit contacting Kap114 are labeled while those of the other subunit are labeled β-hairpin' and AS'. Right, a schematic of the structure. Note the similarity of Kap114 and H2A-H2B here with the binary Kap114•H2A-H2B complex in Figure 1B. **(B)** Nap1₂•**Kap114**•H2A-H2B: H2A-H2B (orange; right view same as in **(A)**) and Nap1₂ (cyan) surfaces, showing the Kap114 interfaces in gray. **(C)** Left, the Nap1₂•H2A-H2B•**Kap114** structure with its consensus and local refined maps, aligned to H2A-H2B of **(A)**. Right, schematic. **(D)** Nap1₂•H2A-H2B•**Kap114**: the histone (orange), Nap1₂ (cyan) and Kap114 (gray) interfaces shown on Nap1₂ and histone surfaces. The Nap1₂ here is related to that in Nap1₂•**Kap114**•H2A-H2B **(B)** by a 50°rotation. **(E)** The Nap1₂•H2A-H2B (5G2E) crystal structure, with Nap1₂ aligned to that in **(C)**. The Nap1₂ β-hairpins here are disordered and here shown here in gray are hairpins from the structure in **(C)**; Nap1₂ alignment in SI Appendix, Fig. S4E. Alignment of (**C**) and (**E**) in SI Appendix, Figs. S7 and 17. **(F)** Nap1₂ and H2A-H2B interfaces of the 5G2E structure.

198   Kap114 binds $Nap1_2$ at two interfaces that involve the same $Nap1_2$ subunit (Figs.

199   2B and 3A). Interface 1, formed by residues 931-942 in the acidic Kap114 h19loop binding

200   to the basic β-hairpin of one $Nap1_2$ subunit, covers 192 $Å^2$ of $Nap1_2$. Interface 2, formed by

201   the basic Kap114 h6-h7loop being close to the negatively charged $Nap1_2$ accessory

202   subdomain (AS), covers 82 $Å^2$ of $Nap1_2$.

Two ternary complex structures: the $Nap1_2$•H2A-H2B•*Kap114* structure

The $Nap1_2$•H2A-H2B•**Kap114** structure is very different. Here, H2A-H2B is

sandwiched between one subunit of the $Nap1_2$ and the C-terminal HEAT repeats of

Kap114 (Fig. 2C and SI Appendix, Fig. S6C and D). The 1555 $Å^2$ interface of H2A-H2B

with the C-terminal HEAT repeats of Kap114 is like the equivalent interfaces in

$Nap1_2$•**Kap114**•H2A-H2B and in other H2A-H2B-bound Kap114 structures (SI Appendix,

Fig. S8). On the contrary, in the $Nap1_2$•H2A-H2B•**Kap114** structure, the N-terminal half of

Kap114 assumes a distinct, open conformation where repeats h1-h2 are ~20 Å apart from

the bound H2A-H2B, twice the distance observed in the structure of Ran^{GTP}•Kap114•H2A-

H2B (41) (SI Appendix, Fig. S7B).

$Nap1_2$ in $Nap1_2$•H2A-H2B•**Kap114** is tethered at two sites: one interface is with

Kap114 and the other with H2A-H2B (Figs. 2D and 3B). The 271 $Å^2$ Kap114-$Nap1_2$

interface here involves the same $Nap1_2$ β-hairpin and Kap114 h19loop that form Interface

1 of $Nap1_2$•**Kap114**•H2A-H2B (Fig. 2B and D). However, $Nap1_2$ in the two structures are

oriented differently relative to the importins (Fig. 2A and C, and SI Appendix, Fig. S9). In

$Nap1_2$•**Kap114**•H2A-H2B, $Nap1_2$ residues K291, R293 and K295, on one face of the β5

strand, contact the Kap114 h19loop (Fig. 3A). In $Nap1_2$•H2A-H2B•**Kap114**, $Nap1_2$ residues

E288, R290 and Q292, on the opposite face of β5 strand, contact the Kap114 h19loop and

the h20b helix (Fig. 3B).

Figure 3. The two Kap114/Nap1$_2$/H2A-H2B structures: interface details and mutagenesis. Dashed lines in **(A)** and **(B)** show intermolecular contacts <4.0 Å (yellow) and <8.0 Å (light grey). **(A)** Details of the Kap114 (grey)-Nap1$_2$ (cyan) interfaces in the Nap1$_2$•**Kap114**•H2A-H2B structure. Contacts are not drawn for Interface 2 because the Nap1$_2$ AS is not observed in the final map. **(B)** Kap114-Nap1$_2$ contacts (left) and H2A-H2B-Nap1$_2$ contacts (right) in the Nap1$_2$•H2A-H2B•**Kap114** structure. **(C)** Pull-down assay with 1 µM immobilized MBP-Kap114 and Nap1 FL WT or interface mutants β$_{KRK}$ (K291A/R293A/K295A), AS$_{mut}$ (S177A/E178A), β$_{ERQ}$ (E288A/R290A/Q292A), α5/6$_{mut}$ (E194A/D201A/D205A), ± equimolar H2A-H2B. Mutated residues are in bold in **(A)** and **(B)**. Both sets of Nap1$_2$ β-hairpin mutations disrupted Kap114/Nap1$_2$/H2A-H2B ternary complex formation. Neither Nap1$_2$ AS nor histone-binding mutations had any effect. **(D)** Pull-down assay using 1 µM MBP-Kap114 WT or mutants Δh19L (h19loop deleted), YD$_{AA}$ (Y939A/D942A) and DD$_{AA}$ (D928A/D929A), with Nap1 FL ± H2A-H2B. Kap114 h19loop deletion disrupted ternary complex formation but double loop mutations did not. **(C)** and **(D)**, Coomassie-stained SDS-PAGE (controls in SI Appendix, Figs. S10).

The Nap1$_2$-histone interface of Nap1$_2$•H2A-H2B•**Kap114** involves acidic residues of

the α4-6 helices of one $Nap1_2$ subunit contacting basic residues of the H2A αN and α1

helix (Fig. 3B). This 207 $Å^2$ interface is located at the histone interface in the binary

$Nap1_2 \cdot$H2A-H2B complex but is only ~25% of the binary complex interface (Fig. 2D-F).

However, the histone is far from exposed as more than 1500 $Å^2$ of its surface is buried by

Kap114 (Fig. 2E).

$Nap1_2$ β-hairpin is the hotspot for Kap114-binding

We mutated residues in both Kap114 and Nap1 that participate in Kap114-$Nap1_2$

interactions to assess their importance in the two different Kap114/$Nap1_2$/H2A-H2B

structures. First, we mutated Nap1 residues in the two Kap114-$Nap1_2$ interfaces of the

$Nap1_2 \cdot$**Kap114**$\cdot$H2A-H2B structure (Fig. 3A). Alanine mutations of $Nap1_2$ β-hairpin

residues K291, R293 and K295 in Interface 1 (mutant β_{KRK}) abolished pull-down by

immobilized MBP-Kap114 in the absence and presence of H2A-H2B, while alanine

mutations of the AS residues S177 and E178 in Interface 2 (mutant AS_{mut}) did not (Fig.

3C). These results validated the $Nap1_2$ β-hairpin as an energetically important Kap114-

binding site and showed that the AS is less important for ternary complex formation.

Next, we mutated contact residues in the Kap114-$Nap1_2$ and the $Nap1_2$-H2A-H2B

interfaces that are unique to the $Nap1_2 \cdot$H2A-H2B$\cdot$**Kap114** structure (Fig. 3B). Alanine

mutations of the $Nap1_2$ β-hairpin residues E288, R290 and Q292 (mutant β_{ERQ}) that

contact Kap114 abolished Nap1 pull-down by MBP-Kap114 in the absence and presence

of H2A-H2B, while mutations of residues E194, D201 and D205 (mutant $\alpha5/6_{mut}$) that

contact H2A-H2B did not (Fig. 3C). These results support the importance of the $Nap1_2$ β-

hairpin for Kap114-binding, in contrast to Nap1$_2$-H2A-H2B interactions, which are less

important for ternary complex formation.

We also mutated contact residues in Kap114. The Kap114 h19loop binds the

Nap1$_2$ β-hairpin in both ternary complex structures (Fig. 3A and B). It is therefore not

surprising that truncating the Kap114 h19loop to remove all contact residues in both

structures (mutant Kap114 Δh19L) abolished Kap114-Nap1 pull-down and formation of the

ternary complex (Fig. 3D). However, mutating just Kap114 Y939 and D942 (mutant

Kap114 YD$_{AA}$) that contact Nap1$_2$ solely in Nap1$_2$•**Kap114**•H2A-H2B or mutating just D928

and D929 (mutant Kap114 DD$_{AA}$) that contact Nap1$_2$ solely in Nap1$_2$•H2A-H2B•**Kap114**,

did not affect Kap114-Nap1 pull-down (Fig. 3D). Many nearby acidic/electronegative

Kap114 side chains may also interact with the Nap1$_2$ β-hairpin (Fig. 3A and B).

In summary, mutagenesis findings indicate that although the interfaces between the

Nap1$_2$ β-hairpin and Kap114 h19loop are relatively small in both the Nap1$_2$•Kap114•H2A-

H2B and Nap1$_2$•H2A-H2B•Kap114 structures, these interactions are essential for binding.

Thus, these results validate the cryo-EM structures of both complexes. Our structural and

biochemical results are also supported by previous studies that suggested the importance

of the Nap1 b-hairpin for localization to the yeast nuclei (13, 36, 42).

Nap1$_2$•*Kap114*•H2A-H2B *versus* Nap1$_2$•H2A-H2B•*Kap114*

The cryo-EM structures and mutagenic analysis above inform on how Kap114,

Nap1$_2$ and H2A-H2B may associate in the absence of Ran^{GTP}, such as in the cytoplasm.

Mutagenesis clearly shows that interactions between Nap1$_2$ β-hairpin and Kap114 h19loop

are critical for ternary complex formation even though cryo-EM analysis suggests the

interactions are dynamic (Figs. 2 and 3). While challenging for structure determination,

$Nap1_2$ motions might influence interactions with its binding partners. Various $Nap1_2$

structures suggest variations in protein dynamics. The cryo-EM map of unliganded $Nap1_2$

has features for the entire homodimer (SI Appendix, Fig. S4A), contrasting with the crystal

structure of the $Nap1_2$•H2A-H2B complex where electron density for both sets of β-hairpins

and AS regions was absent (Fig. 2E). In our ternary complex structures, the β-hairpins that

bind Kap114 are resolved in the cryo-EM maps, whereas those of the other $Nap1_2$ subunit

are not (Fig. 2A and C). These structural observations may hint at dynamics-driven

allostery and asymmetry within the conformationally symmetric $Nap1_2$. Ligand binding,

such as of histone and/or Kap114, does not alter $Nap1_2$ conformation but may amplify its

movements, resulting in an entropic penalty that prevents symmetric binding of the second

ligand (43, 44). Increased $Nap1_2$ motions upon Kap114 binding could contribute

entropically to binding energy, stabilizing interactions, despite the small interfaces.

The interfaces between the $Nap1_2$ β-hairpin and Kap114 differ in

$Nap1_2$•**Kap114**•H2A-H2B and $Nap1_2$•H2A-H2B•**Kap114**. However, mutating either

interface prevents the association of the three proteins. This implies that dynamic

switching between the two configurations is essential for forming and maintaining the

stability of the ternary complex. The $Nap1_2$•**Kap114**•H2A-H2B and $Nap1_2$•H2A-

H2B•**Kap114** structures likely represent components within a dynamic ensemble of

interconverting Kap114/$Nap1_2$/H2A-H2B states in the cytoplasm. It is unclear which

components of this ensemble will subsequently translocate through the NPC into the

nucleus. The physical and chemical nature of the permeability barrier of the NPC and the

mechanisms importin-cargo translocation across that barrier, including knowledge of

importin-cargo conformations in the barrier, are far from understood at this time (45-48).

We need to consider another scenario. Our cryo-EM sample was prepared with

Kap114, $Nap1_2$ and excess H2A-H2B, which has been observed to destabilize the ternary

Kap114/$Nap1_2$/H2A-H2B complex (Fig. 1C-E). It is plausible that the structures of

$Nap1_2$•**Kap114**•H2A-H2B and/or $Nap1_2$•H2A-H2B•**Kap114** structures could represent

intermediates during disassembly of the ternary complex into binary $Nap1_2$•H2A-H2B and

Kap114•H2A-H2B   complexes.   Among   these   potential   intermediates,   the

$Nap1_2$•**Kap114**•H2A-H2B structure, characterized by a substantially smaller $Nap1_2$

interface, might be a more favorable candidate for an intermediate state in the disassembly

process.

Both $Nap1_2$•**Kap114**•H2A-H2B and $Nap1_2$•H2A-H2B•**Kap114** likely represent

configurations of the Kap114/$Nap1_2$/H2A-H2B ternary complex that could exist in the

cytoplasm. However, their specific roles in the process of importing H2A-H2B into the

nucleus   are   unclear   and   warrant   deeper   investigation.   Understanding   how

Kap114/$Nap1_2$/H2A-H2B interacts with Ran^{GTP}, as is expected in the Ran^{GTP}-rich nucleus,

may provide insights into the distinct functions of the $Nap1_2$•**Kap114**•H2A-H2B *versus*

$Nap1_2$•H2A-H2B•**Kap114** structures in the cytoplasmic millieu.

Interactions between Kap114, Nap1 and H2A-H2B in the presence of Ran^{GTP}

The Kap114/$Nap1_2$/H2A-H2B ternary complex will encounter Ran^{GTP} upon entering

the nucleus. All importins bind Ran^{GTP} tightly, almost always causing importin•cargo

dissociation to release cargo in the nucleus (49). H2A-H2B import is an exception because

Ran^{GTP} binds Kap114•H2A-H2B to form a stable ternary Ran^{GTP}•Kap114•H2A-H2B

complex (Fig. 4A) (41). Similarly, Kap114-Nap1 interaction persists in the presence of Ran^{GTP} as shown by pull-down assays and AUC analysis (Fig. 4B and SI Appendix, Fig. S12). SEC- MALS analysis showed that an equimolar mix of Kap114, Nap1$_2$, H2A-H2B and Ran^{GTP} produced a major peak consistent with a complex that contains one molecule each of the four proteins (Fig. 4C).

We solved the cryo-EM structure of this quaternary Kap114/Nap1$_2$/H2A-H2B/Ran^{GTP} complex to 2.9 Å resolution (Fig. 4D and Si Appendix, Figs. S13 and S14 and Table S2). The final cryo-EM map is well-defined for Kap114 and Ran^{GTP} but the local resolution and map feature for Nap1$_2$ and H2A-H2B are poor. Local refinement produced a 4.0 Å map that allowed docking of Nap1$_2$ and H2A-H2B. We built the structure of the quaternary complex using Ran^{GTP} bound to the N-terminal HEAT repeats of Kap114 from the Ran^{GTP}•Kap114•H2A-H2B structure (PDB: 8F1E) (41) and Nap1$_2$, H2A-H2B and Kap114 repeats h17-h20 from the Nap1$_2$•H2A-H2B•**Kap114** structure. We named this assembly the Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} complex (Fig. 4D).

In the Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} quaternary complex structure, Ran^{GTP} binds Kap114 the same as in the Ran^{GTP}•Kap114•H2A-H2B structure (41) (Fig. 4A and D). Comparing Fig. 4D and E shows that Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} is essentially Nap1$_2$•H2A-H2B•**Kap114** with Ran^{GTP} bound at the N-terminal half of Kap114. Ran-binding in the quaternary complex stabilizes a Kap114 conformation where its N-terminus

Figure 4. Structure of the Nap1₂•H2A-H2B•Kap114•Ran^GTP complex. (A) The Ran^GTP•Kap114•H2A-H2B structure (8F1E). **(B)** Pull-down assay with equimolar MBP-Kap114 (1 µM), Nap1₂, H2A-H2B and Ran^GTP; controls in SI Appendix, Fig. S12A-C. **(C)** SEC-MALS of Kap114, Nap1₂, H2A-H2B and Ran^GTP (green curve), plotted like Fig. 1C, with gel of peak fractions. The major peak at 270-293 kDa has mass like the 1:1:1:1 Kap114/Nap1₂/H2A-H2B/Ran^GTP complex (269 kDa). Kap114/Nap12/H2A-H2B SEC-MALS (orange, from Fig. 1D) is shown for comparison. Control in SI appendix, Fig. S12E. **(D)** The Nap1₂•H2A-H2B•**Kap114**•Ran^GTP structure, with the consensus/local refined maps (gray/cyan) overlayed, shown next to the Nap1₂•H2A-H2B•**Kap114** structure in **(E)**. **(A)** and **(D)** are aligned at Ran^GTP, and **(D)** and **(E)** are aligned at H2A-H2B. Pull-down assays with **(F)** Nap1₂ WT, β-hairpin mutants (β_KRK and β_ERQ) and histone-binding site mutant α5/6_mut, and **(G)** Kap114 h19loop mutants Δh19L, YD_AA and DD_AA. Proteins in **(B)**, **(C)**, **(F)** and **(G)** visualized by Coomassie-stained SDS-PAGE. **(H)** Summary of all pull-down assays in Figs. 1, 3 and 4. iK refers to immobilized MBP-Kap114.

331
is closer to the bound H2A-H2B than in the ternary Nap1₂•H2A-H2B•**Kap114** structure

(Fig. 4D and E and SI Appendix, Fig. S7C). It is interesting that $Nap1_2$•H2A-

H2B•**Kap114**•Ran^{GTP} looks like $Nap1_2$•H2A-H2B•**Kap114** rather than $Nap1_2$•**Kap114**•H2A-

H2B (compare Fig. 4D and E with Fig. 2A). This similarity may suggest that $Nap1_2$•H2A-

H2B•**Kap114** is the ternary configuration that enters the NPC for translocation into the

nucleus.

Like the cytoplasmic $Kap114/Nap1_2/H2A$-H2B ternary complex, mutation of the

$Nap1_2$ β-hairpin or truncation of the Kap114 h19loop abolished Kap114-$Nap1_2$ pull-down in

the presence of Ran^{GTP} and H2A-H2B (Fig. 4F-H), validating the $Nap1_2$•H2A-

H2B•**Kap114**•Ran^{GTP} cryo-EM structure and revealing that the $Nap1_2$ β-hairpin remains the

key Kap114-binding determinant in the presence of Ran^{GTP}.

Kap114 and $Nap1_2$: co-chaperoning H2A-H2B in the absence of Ran^{GTP}

Nap1 and Kap114/*Hs* IMP9 were previously identified as highly effective histone

chaperones that prevent aggregation of H2A-H2B with DNA (13-16, 41, 50). Consistent

with their histone chaperone function, the $Nap1_2$ and Kap114 in both $Nap1_2$•**Kap114**•H2A-

H2B and $Nap1_2$•H2A-H2B•**Kap114** shield substantial portions (~1700-2200 Å^2) of the

nucleosomal H3-H4 and DNA binding regions of H2A-H2B (Fig. 2 and SI Appendix, Fig.

S15).

We performed DNA competition assays to probe the chaperoning of H2A-H2B by

Kap114 and $Nap1_2$. In this assay, H2A-H2B chaperoning would result in disappearance of

low-mobility DNA•H2A-H2B bands and appearance of high-mobility free DNA bands. We

performed the experiments by either assembling DNA•H2A-H2B first and then titrating in

$Nap1_2$ and/or Kap114 (Fig. 5A) or assembling complexes of H2A-H2B with Kap114 and/or

Nap1$_2$ first before titrating them into DNA (SI Appendix, Fig. S16A). Regardless of the

order of addition, Kap114 and Nap1$_2$ together chaperone H2A-H2B from aggregation with

DNA more effectively than either Kap114 or Nap1 alone.

To summarize, when Ran^{GTP} is absent, as in the cytoplasm, Kap114 and Nap1$_2$

within Kap114/Nap1$_2$/H2A-H2B work together to enhance H2A-H2B sequestration

compared to Kap114 alone. This effectively shields the histone from aberrant interactions.

361 Kap114/Nap1 co-chaperoning and nucleosome assembly in the presence of Ran^{GTP}

The quaternary Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} structure buries 1847 Å^2 of H2A-

H2B. Here, Nap1$_2$ and Kap114 together shield the DNA binding interfaces of H2A-H2B,

reminiscent of their role in the cytosolic/ternary Nap1$_2$•H2A-H2B•**Kap114** structure (Figs.

2D, 4D and SI Appendix, Fig. S15). The similarity in their protein arrangements suggests

that, even in the presence of Ran^{GTP}, Kap114 and Nap1$_2$ are likely to prevent H2A-H2B

from aggregating with DNA. We tested this hypothesis with DNA competition assays using

Kap114, H2A-H2B, with/without Nap1$_2$ and titrating in Ran^{GTP} (Fig. 5B and C, SI Appendix,

Fig. S16B). When Ran^{GTP} is absent, Kap114 alone effectively chaperones H2A-H2B;

however, its efficacy as a histone chaperone decreases substantially in the presence of

the GTPase. Conversely, when both Nap1$_2$ and Kap114 are present, they efficiently

prevent non-specifically aggregation of H2A-H2B with DNA irrespective of whether Ran^{GTP}

is present or absent. Therefore, within the quaternary/nuclear Nap1$_2$•H2A-

H2B•**Kap114**•Ran^{GTP} complex, Kap114 and Nap1$_2$ cooperate to successfully chaperone

H2A-H2B. SI Appendix Fig. S15 compares H2A-H2B surfaces in various complexes.

Figure 5. Kap114 and Nap1₂ co-chaperone H2A-H2B in the presence of RanGTP for targeted release to tetrasomes. (A, B) DNA competition assays of DNA•H2A-H2B complex (D•H) added to 1-4 μM Nap1 (N₂), Kap114 (K) or Kap114+Nap1₂ mixture (K+N₂), in the absence of RanGTP **(A)** or added to 4 μM K or K + N₂ and 1-8 μM RanGTP (R) **(B)**. **(C)** Summary of DNA competition assay results. **(D)** Nucleosome assembly assay. Pre-assembled K•H or K/N₂/H complexes without or with 0.125-1 μM R were added to tetrasomes (TET); **TET added last and nucleosome (NUC) formation tracked (details in methods). **(E)** Nucleosome assembly assay results. **(A, B** and **D)** visualized using ethidium bromide- and Coomassie-stained native PAGE gels. **(F)** Schematic of key findings: without RanGTP (e.g. cytoplasm), Kap114, Nap1₂ and H2A-H2B assemble into a ternary complex that adopts at least two different configurations where Kap114 and Nap1₂ co-chaperone H2A-H2B. In the presence of RanGTP (e.g. nucleus), Kap114 binds RanGTP to form the quaternary complex Nap1₂•H2A-H2B•**Kap114**•RanGTP that inhibits histone aggregation and transfers H2A-H2B to assemble nucleosome (bottom right). Without Nap1₂, H2A-H2B•Kap114•RanGTP does not inhibit histone aggregation (bottom left).

We performed nucleosome assembly assays to assess the effects of Kap114,

Nap1$_2$ and Ran^{GTP} on depositing H2A-H2B to form nucleosomes (Fig. 5D, E and SI

Appendix, Fig. S16). As shown previously, Kap114 alone does not deposit H2A-H2B onto

tetrasomes (41), but Ran^{GTP} facilitates efficient histone transfer from Kap114 to form

nucleosomes. Nap1$_2$ facilitates efficient histone transfer from Kap114 to form nucleosomes

when Ran^{GTP} absent. However, nucleosome assembly is most efficient when Kap114,

Nap1$_2$ and Ran^{GTP} are all present, as evidenced by the almost complete disappearance of

tetrasomes in lanes 9 – 11 of Fig. 5D.

In summary, Kap114 and Nap1$_2$ in Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} cooperate to

chaperone H2A-H2B (Fig. 5F). Without Nap1$_2$, Ran^{GTP} would bind to the Kap114•H2A-

H2B complex to reorient the bound histone and increase its exposure to DNA. However,

Nap1$_2$ binding H2A-H2B alongside Kap114 in the nuclear/quaternary Nap1$_2$•H2A-

H2B•**Kap114**•Ran^{GTP} complex shields the histone from DNA and positions it for effective

transfer into nucleosomes (Fig. 5F).

H2A-H2B transfer from Nap1$_2$•H2A-H2B•Kap114•Ran^{GTP} to nucleosome

We used a Kap114 mutant that cannot bind Nap1$_2$ to understand how Kap114,

Nap1$_2$ and Ran^{GTP} might work together to transfer H2A-H2B onto the assembling

nucleosome. Pull-down binding assays were conducted with equimolar Kap114 Δh19L,

Nap1$_2$, H2A-H2B and Ran^{GTP} (Figs. 3D, 4G and H). The Kap114 Δh19L mutant, unable to

bind Nap1$_2$, still managed to capture H2A-H2B in the presence of Nap1$_2$, forming the

Kap114 Δh19L•H2A-H2B complex (Figs. 3D and 4H), consistent with H2A-H2B's high

affinity for the importin (41). Kap114 Δh19L also pulled down H2A-H2B in the presence of

Ran^{GTP} to form Ran^{GTP}•Kap114 Δh19L•H2A-H2B (Fig. 4G and H). However, Kap114

Δh19L failed to capture H2A-H2B in the presence of both Ran^{GTP} and Nap1$_2$, suggesting

that H2A-H2B may be binding to unbound Nap1 instead (Fig. 4G and H). When Ran^{GTP}

was present, free Nap1, unable to bind Kap114 Δh19L, effectively displaced histones from

the importin. These results imply that Ran^{GTP} in the Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP}

complex likely promotes the transfer of H2A-H2B from Kap114 to Nap1$_2$, facilitating

optimal deposition of H2A-H2B into nucleosomes. The quaternary assembly may provide a

path to specifically move H2A-H2B from Kap114 to Nap1$_2$ to the tetrasome without losing

the histone to non-specific/aberrant interactions in the nucleus.

Conclusion

In an environment devoid of Ran^{GTP}, such as in the cytoplasm, equimolar Kap114,

Nap1$_2$ and H2A-H2B assemble into a 1:1:1 Kap114/Nap1$_2$/H2A-H2B ternary complex. Two

cryo-EM structures, Nap1$_2$•**Kap114**•H2A-H2B and Nap1$_2$•H2A-H2B•**Kap114**, revealed two

distinct configurations of the three proteins, potentially depicting various states within a

dynamic equilibrium of Kap114/Nap1$_2$/H2A-H2B complexes in the cytoplasm. The DNA-

binding surfaces of H2A-H2B in both structures are shielded and DNA competition assays

showed that Kap114 and Nap1$_2$ together co-chaperone the histone from aberrant

interactions more effectively than either protein alone. The binding of Ran^{GTP}, simulating

condition within the nucleus, results in the Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} quaternary

complex, which assumes the protein arrangement observed in the Nap1$_2$•H2A-

H2B•**Kap114** ternary structure, with Ran^{GTP} binding to the N-terminal HEAT repeats of

Kap114. The Nap1$_2$•H2A-H2B•**Kap114**•Ran^{GTP} complex relies on Nap1$_2$'s critical role for

effective H2A-H2B chaperoning in the presence of Ran^{GTP}. This complex efficiently

deposits H2A-H2B to form nucleosomes. Through delineating the biochemical mechanism

of Nap1$_2$ as a co-factor in H2A-H2B nuclear import, together with Kap114, this study

explains how histones H2A-H2B are co-chaperoned and seamlessly transferred into

assembling nucleosomes.

Materials and Methods

Pull-down binding Assays

Details for protein constructs, expression, and purification are in SI methods. All

biochemical and biophysical studies were conducted at an ionic strength close to

physiological salt levels of 150 mM NaCl. *In vitro* pull-down binding assays were performed

in triplicate by incubating proteins at their indicated concentrations with 20 µL amylose

resin (bead bed volume; New England Biolabs) in a 200 µL reaction in Assay Buffer (20

mM HEPES pH 7.4, 150 mM NaCl, 2 mM MgCl2, 10% (v/v) glycerol and 2 mM DTT) for 30

min at 4°C, followed by washing with 500 µL buffer twice. Bound proteins were separated

and visualized by SDS-PAGE and Coomassie Blue staining. Nap1 band intensities were

quantified using Bio Rad Image Lab software, adjusted for the intensity of the MBP-

Kap114 band and normalized to Nap1 WT control.

Analytical Ultracentrifugation

439 Individual proteins were dialyzed overnight into AUC buffer containing 20 mM Tris-

440 HCl pH 7.5, 150 mM NaCl, 2 mM MgCl$_2$ and 2 mM TCEP, at 4°C and assembled in the

441 AUC sample chamber at the indicated concentrations in 400 µL. Sedimentation

coefficients were measured by monitoring absorbance at 280 nm in a Beckman-Coulter

Optima XL-1 Analytical Ultracentrifuge. Time stamps were corrected using REDATE (51).

SEDNTERP was used to calculate the buffer density, buffer viscosity, and protein partial-

specific volumes (52). SEDFIT was used to calculate sedimentation coefficient

distributions c(s) where the regularization calculated a confidence level of 0.68 was used,

time-independent noise elements were accounted for, and at a resolution of 100 (53).

SEDFIT was also used to estimate molecular weight by obtaining the sedimentation

coefficient through integration of c(s) and the frictional ratios by refining the fit of the model.

The c(s) distribution and isotherm integration was done with GUSSI (54). Further isotherm

fitting was performed using SEDPHAT (55).

Size Exclusion Chromatography Multi-Angle Light Scattering

SEC-MALS experiments were performed in Assay buffer without glycerol with

proteins at the indicated molar ratios to 80 µM Kap114 following established protocol,

including sample dialysis and buffer filtration (56). Concentration of proteins are high in the

small injection of 100 µL to ensure complete complex formation when it is diluted in the

SEC column. The data was collected and processed using ASTRA software using default

settings with no de-spiking (Wyatt Technology). The run for 80 µM Kap114 + 80 µM H2A-

H2B + 80 µM Nap1$_2$ was interrupted at the void volume due to fractionation issues causing

a total pause of 1.895 minutes. A correction of 1.38933 mL was applied to match other

runs.

Cryo-EM Sample preparation and data collection and analysis

To assemble the Kap114/H2A-H2B/Nap1 complex, *Sc* H2A-H2B, Nap1 core, and

Kap114 were dialyzed overnight into cryo-EM buffer containing 20 mM Tris pH 7.5, 150

mM NaCl, and 2 mM DTT. A 1:2 molar ratio mixture of Nap1 core dimer:H2A-H2B was

incubated at room temperature for 10 minutes followed by addition of 1 molar ratio of

Kap114, followed by rapid addition of glutaraldehyde to a final concentration of 0.05%.

Crosslinking proceeded for one minute before quenching and removal of glutaraldehyde by

SEC using a Superdex200 10/300 Increase column (Cytiva) that was equilibrated with

cryo-EM buffer containing TCEP (instead of DTT). 0.5 mL fractions were collected for

mass   photometry   (MP)   analysis   (details   in   SI   Methods).   To   assemble   the

Ran^{GTP}/Kap114/Nap1/H2A-H2B complex, a 10 mg/mL mix of a stoichiometric molar ratio

of 1 Kap114 to 1 H2A-H2B to 2 Nap1$_2$ FL (His tag intact) to 1 Ran^{GTP} was dialyzed

overnight before crosslinking with 0.05% glutaraldehyde for 1 minute and separation by

SEC in Assay buffer without glycerol. Fractions of 0.25 mL were collected for MP analysis.

Fractions with the least aggregated species and most enriched with the relevant

complexes (described in SI Appendix, Figs. S3A and S13A) were selected for grid

preparation.

For Kap114/H2A-H2B/Nap1, the sample was diluted to ~1.2 mg/mL and

supplemented with 0.003125% (w/v) Triton X-100. For Ran^{GTP}/Kap114/Nap1/H2A-H2B,

sample was diluted to a ~1.5 mg/mL with 0.003125% (w/v) Tyloxapol. Quantifoil grids were

glow-discharged, blotted, and plunge-frozen. Grids were screened and data was collected

and analyzed as described in SI Methods.

DNA competition assays

601 Widom DNA was purified from pUC19-30X601 grown in C2925 dam⁻dcm⁻ cells

(gift from Dr. Mike Rosen) using Qiagen Giga prep kit following manufacturer's protocol.

The DNA was resuspended in 10 mM Tris pH 7.4, 1 mM EDTA in the last step of

purification. EcoRV (New England Biolabs) cleavage was performed in CutSmart buffer

overnight, and the DNA extracted using PCIAA and precipitated in sodium acetate pH 5.2

and ethanol. The DNA was then rinsed with ethanol, dried, and resolubilized in 10 mM Tris

pH7.4, 10 mM $MgCl_2$. The cut insert was separated from the vector using PEG8000

precipitation and subjected to a final ion exchange purification on a DEAE-FF column

before it was concentrated and frozen.

DNA was mixed with H2A-H2B, Kap114 and/or $Nap1_2$ FL with and without Ran^{GTP}

at the indicated concentrations in 15 µL reactions in Assay Buffer. DNA was either pre-

mixed with H2A-H2B in a 3X concentration master-mix, or alternatively, added at the end.

Reactions were incubated at room temperature for 30 min and then 5 µL of sample buffer

containing 20% (v/v) glycerol and bromophenol blue was added. 8 µL each of this gel

sample was ran on two separate Native 5% PAGE gels with 0.5X TBE at 150V for 40 min.

One gel was stained with EtBr in water and the other with Coomassie Blue. Assays were

performed in duplicates.

Nucleosome assembly assays

503   Tetrasomes were made by mixing *Xl* H3-H4 tetramers and DNA at 1:1 ratio (DNA

504   conc = 0.7 mg/mL) in 10 mM HEPES pH7.4, 1 mM EDTA, 2 M NaCl, 1 mM DTT and then

505   dialyzed step-wise from 2 M NaCl (0.5 hr) to 1M (1 hr), 0.85 M (1.5 hr), 0.65 M (1.5 hr), 0.2

506   M (1.5 hr) and 0.15 NaCl M (overnight) at 4°C. The volume was measured at the end and

reaction was diluted to 2 µM with Assay buffer to 2X working concentration immediately before use, assuming 100% formation of tetrasomes. 12 µL reactions with *Sc* H2A-H2B, Kap114, Nap1$_2$ FL and tetrasomes with or without Ran^{GTP} were assembled at indicated concentrations in figure legends of Figure 5 and incubated together at room temperature for 1 hr before addition of 4 µL of sample buffer. 7 µL of the resulting samples were visualized the same way as DNA competition assays. Assays were performed in duplicates.

The input sample in lane 1 of Fig. 5D shows that tetrasome formation is not 100% complete and some free DNA was present. Free DNA was present in all reactions and not incorporated into nucleosomes, suggesting that any free H3-H4 not incorporated into tetrasomes was inactive/degraded. The control experiment without Ran^{GTP} (SI Appendix, Fig. S16C) shows that no tetrasomes remain after addition of 1 µM H2A-H2B, suggesting that the tetrasome concentration was likely < 0.5 µM. This is consistent with the results of lane 10 Fig. 5D where nucleosome formation is not 100% complete with the addition of just 0.5 µM H2A-H2B. The concentration of 0.5 µM H2A-H2B used in the experiment shown in Fig 5D was selected to capture differences in the effectiveness of the Kap114/Nap1$_2$/H2A-H2B ternary complex and the binary Kap114•H2A-H2B complex to deposit H2A-H2B onto tetrasomes, in the presence and absence of Ran^{GTP}.

Acknowledgments

We thank all members of the Structural Biology Laboratory (SBL) and the CEMF in UTSW for their expert assistance with cryo-EM data collection. We greatly appreciate Macromolecular Biophysics Resource (MBR) and Chad Brautigam for discussion and use

of the AUC and Mass Photometer. We also thank the Erzberger lab for using their plate

reader for FP assays. We thank Rosen Lab for help with the Widom DNA purification. We

also thank Casey Wing, Mike Rosen and Mike Rout for critical reading of the manuscript.

SBL and CEMF at UTSW are partially supported by grant RP220582 from the Cancer

Prevention & Research Institute of Texas (CPRIT). The Refeyn Mass Photometer in MBR

was supported by grant S10-OD030312 from the National Institute of Health. We thank the

D'Arcy lab for the use of their SEC-MALS equipment and Joy Shaffer for training.

Molecular graphics and analyses performed with UCSF ChimeraX, developed by the

Resource for Biocomputing, Visualization, and Informatics at the University of California,

San Francisco, with support from National Institutes of Health R01-GM129325 and the

Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and

Infectious Diseases. This research used resources of the Advanced Photon Source, a U.S.

Department of Energy (DOE) Office of Science user facility operated for the DOE Office of

Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. This

work was funded by NIGMS of NIH under Awards R35GM141461 (Y.M.C.),

R01GM069909 (Y.M.C.), T32GM008203 (J.J), the Welch Foundation Grants I-1532

(Y.M.C.), support from the Alfred and Mabel Gilman Chair in Molecular Pharmacology,

Eugene McDermott Scholar in Biomedical Research (Y.M.C.) and the Gilman Special

Opportunities Award (H.Y.J.F.).

References

1.   A. K. Hogan, D. R. Foltz, Reduce, Retain, Recycle: Mechanisms for Promoting
Histone Protein Degradation versus Stability and Retention. *Mol Cell Biol* **41**,
e0000721 (2021).

2.   R. K. Singh *et al.*, Excess histone levels mediate cytotoxicity via multiple
mechanisms. *Cell Cycle* **9**, 4236-4244 (2010).

3.   S. J. Elsasser, S. D'Arcy, Towards a mechanism for histone chaperones. *Biochim
Biophys Acta* **1819**, 211-221 (2013).

4.   C. M. Hammond, C. B. Stromme, H. Huang, D. J. Patel, A. Groth, Histone
chaperone networks shaping chromatin function. *Nat Rev Mol Cell Biol* **18**, 141-158
(2017).

5.   E. I. Campos *et al.*, The program for processing newly synthesized histones H3.1
and H4. *Nat Struct Mol Biol* **17**, 1343-1351 (2010).

6.   H. Tagami, D. Ray-Gallet, G. Almouzni, Y. Nakatani, Histone H3.1 and H3.3
complexes mediate nucleosome assembly pathways dependent or independent of
DNA synthesis. *Cell* **116**, 51-61 (2004).

7.   L. J. Benson *et al.*, Modifications of H3 and H4 during chromatin replication,
nucleosome assembly, and histone exchange. *J Biol Chem* **281**, 9287-9296 (2006).

8.   M. Baake, D. Doenecke, W. Albig, Characterisation of nuclear localisation signals of
the four human core histones. *J Cell Biochem* **81**, 333-346 (2001).

9.   P. Muhlhausser, E. C. Muller, A. Otto, U. Kutay, Multiple pathways contribute to
nuclear import of core histones. *EMBO Rep* **2**, 690-696 (2001).

10.   K. Schwamborn, W. Albig, D. Doenecke, The histone H1(0) contains multiple
sequence elements for nuclear targeting. *Exp Cell Res* **244**, 206-217 (1998).

11.   N. E. Bernardes, Y. M. Chook, Nuclear import of histones. *Biochem Soc Trans* **48**,
2753-2767 (2020).

12.   M. Seidel *et al.*, Co-translational binding of importins to nascent proteins. *Nat
Commun* **14**, 3418 (2023).

13.   N. Mosammaparast, C. S. Ewart, L. F. Pemberton, A role for nucleosome assembly
protein 1 in the nuclear transport of histones H2A and H2B. *EMBO J* **21**, 6527-6538
(2002).

14.   X. Chen *et al.*, Histone Chaperone Nap1 Is a Major Regulator of Histone H2A-H2B
Dynamics at the Inducible GAL Locus. *Mol Cell Biol* **36**, 1287-1296 (2016).

15.   Y. Huang, Y. Dai, Z. Zhou, Mechanistic and structural insights into histone H2A-H2B
chaperone in chromatin regulation. *Biochem J* **477**, 3367-3386 (2020).

16.   L. Chang *et al.*, Histones in transit: cytosolic histone complexes and diacetylation of
H4 during nucleosome assembly in human cells. *Biochemistry* **36**, 469-480 (1997).

17.   M. Okuwaki, K. Kato, K. Nagata, Functional characterization of human nucleosome
assembly protein 1-like proteins as histone chaperones. *Genes Cells* **15**, 13-27
(2010).

18.   M. E. Calvert *et al.*, Phosphorylation by casein kinase 2 regulates Nap1 localization
and function. *Mol Cell Biol* **28**, 1313-1325 (2008).

19.   A. J. Andrews, X. Chen, A. Zevin, L. A. Stargell, K. Luger, The histone chaperone
Nap1 promotes nucleosome assembly by eliminating nonnucleosomal histone DNA
interactions. *Mol Cell* **37**, 834-842 (2010).

20.   R. Vlijm, J. S. Smitshuijzen, A. Lusser, C. Dekker, NAP1-assisted nucleosome
assembly on DNA measured in real time by single-molecule magnetic tweezers.
PLoS One **7**, e46306 (2012).

21.   J. Mazurkiewicz, J. F. Kepert, K. Rippe, On the mechanism of nucleosome
assembly by histone chaperone NAP1. *J Biol Chem* **281**, 16462-16472 (2006).

22.   C. Aguilar-Gurrieri *et al.*, Structural evidence for Nap1-dependent H2A-H2B
deposition and nucleosome assembly. *EMBO J* **35**, 1465-1482 (2016).

23.   F. Nagae, S. Takada, T. Terakawa, Histone chaperone Nap1 dismantles an
H2A/H2B dimer from a partially unwrapped nucleosome. *Nucleic Acids Res* **51**,
5351-5363 (2023).

24.   R. Dronamraju *et al.*, Redundant Functions for Nap1 and Chz1 in H2A.Z Deposition.
Sci Rep **7**, 10791 (2017).

25.   K. W. Hsu *et al.*, The synergy between RSC, Nap1 and adjacent nucleosome in
nucleosome remodeling. *Biochim Biophys Acta Gene Regul Mech* **1862**, 129-140
(2019).

26.   W. A. Krajewski, The intrinsic stability of H2B-ubiquitylated nucleosomes and their
in vitro assembly/disassembly by histone chaperone NAP1. *Biochim Biophys Acta*
Gen Subj **1864**, 129497 (2020).

27. B. C. Del Rosario, L. F. Pemberton, Nap1 links transcription elongation, chromatin
assembly, and messenger RNP complex biogenesis. *Mol Cell Biol* **28**, 2113-2124
(2008).

28. N. Sharma, J. K. Nyborg, The coactivators CBP/p300 and the histone chaperone
NAP1 promote transcription-independent nucleosome eviction at the HTLV-1
promoter. *Proc Natl Acad Sci U S A* **105**, 7959-7963 (2008).

29. V. Levchenko, V. Jackson, Histone release during transcription: NAP1 forms a
complex with H2A and H2B and facilitates a topologically dependent release of H3
and H4 from the nucleosome. *Biochemistry* **43**, 2359-2372 (2004).

30. Y. M. Moshkin *et al.*, Histone chaperone NAP1 mediates sister chromatid resolution
by counteracting protein phosphatase 2A. *PLoS Genet* **9**, e1003719 (2013).

31. R. Altman, D. Kellogg, Control of mitotic events by Nap1 and the Gin4 kinase. *J Cell*
Biol **138**, 119-130 (1997).

32. D. R. Kellogg, A. W. Murray, NAP1 acts with Clb1 to perform mitotic functions and
to suppress polar bud growth in budding yeast. *J Cell Biol* **130**, 675-685 (1995).

33. Y. J. Park, J. V. Chodaparambil, Y. Bao, S. J. McBryant, K. Luger, Nucleosome
assembly protein 1 exchanges histone H2A-H2B dimers and assists nucleosome
sliding. *J Biol Chem* **280**, 1817-1825 (2005).

34. J. Zlatanova, C. Seebart, M. Tomschik, Nap1: taking a closer look at a juggler
protein of extraordinary skills. *FASEB J* **21**, 1294-1310 (2007).

35. K. Straube, J. S. Blackwell, Jr., L. F. Pemberton, Nap1 and Chz1 have separate
Htz1 nuclear import and assembly functions. *Traffic* **11**, 185-197 (2010).

36. N. Mosammaparast, B. C. Del Rosario, L. F. Pemberton, Modulation of histone
deposition by the karyopherin kap114. *Mol Cell Biol* **25**, 1764-1778 (2005).

37. B. Ho, A. Baryshnikova, G. W. Brown, Unification of Protein Abundance Datasets
Yields a Quantitative Saccharomyces cerevisiae Proteome. *Cell Syst* **6**, 192-205
e193 (2018).

38. K. Luger, A. W. Mader, R. K. Richmond, D. F. Sargent, T. J. Richmond, Crystal
structure of the nucleosome core particle at 2.8 A resolution. *Nature* **389**, 251-260
(1997).

39. Y. J. Park, S. J. McBryant, K. Luger, A beta-hairpin comprising the nuclear
localization sequence sustains the self-associated states of nucleosome assembly
protein 1. *J Mol Biol* **375**, 1076-1085 (2008).

40.   C. C. Liao *et al.*, Karyopherin Kap114p-mediated trans-repression controls
ribosomal gene expression under saline stress. *EMBO Rep* **21**, e48324 (2020).

41.   J. Jiou *et al.*, Mechanism of RanGTP priming H2A-H2B release from Kap114 in an
atypical RanGTP*Kap114*H2A-H2B complex. *Proc Natl Acad Sci U S A* **120**,
e2301199120 (2023).

42.   J. L. Hodges *et al.*, Nuclear import of TFIIB is mediated by Kap114p, a karyopherin
with multiple cargo-binding domains. *Mol Biol Cell* **16**, 3200-3210 (2005).

43.   N. Popovych, S. Sun, R. H. Ebright, C. G. Kalodimos, Dynamically driven protein
allostery. *Nat Struct Mol Biol* **13**, 831-838 (2006).

44.   S. R. Tzeng, C. G. Kalodimos, Dynamic activation of an allosteric regulatory protein.
Nature **462**, 368-372 (2009).

45.   C. Li, A. Goryaynov, W. Yang, The selective permeability barrier in the nuclear pore
complex. *Nucleus* **7**, 430-446 (2016).

46.   G. Paci, J. Caria, E. A. Lemke, Cargo transport through the nuclear pore complex at
a glance. *J Cell Sci* **134** (2021).

47.   T. Kozai *et al.*, Dynamic molecular mechanism of the nuclear pore complex
permeability barrier. *bioRxiv* 10.1101/2023.03.31.535055 (2023).

48.   H. B. Schmidt, D. Gorlich, Transport Selectivity of Nuclear Pores, Phase
Separation, and Membraneless Organelles. *Trends Biochem Sci* **41**, 46-61 (2016).

49.   C. E. Wing, H. Y. J. Fung, Y. M. Chook, Karyopherin-mediated nucleocytoplasmic
transport. *Nat Rev Mol Cell Biol* **23**, 307-328 (2022).

50.   A. Padavannil *et al.*, Importin-9 wraps around the H2A-H2B core to act as nuclear
importer and histone chaperone. *Elife* **8** (2019).

51.   H. Zhao *et al.*, Recorded scan times can limit the accuracy of sedimentation
coefficients in analytical ultracentrifugation. *Anal Biochem* **437**, 104-108 (2013).

52.   T. M. S. Laue, B.; Ridgeway, T.M.; Pelletier, S.L. , "Computer-aided Interpretation of
Sedimentation Data for Proteins" in Analytical Ultracentrifugation in Biochemistry
and Polymer Science. (Royal Society of Chemistry, Cambridge, U.K. , 1992), pp.
90-125.

53.   P. Schuck, Size-distribution analysis of macromolecules by sedimentation velocity
ultracentrifugation and lamm equation modeling. *Biophys J* **78**, 1606-1619 (2000).

54.   C. A. Brautigam, Calculations and Publication-Quality Illustrations for Analytical
Ultracentrifugation Data. *Methods Enzymol* **562**, 109-133 (2015).

55.   H. Zhao, G. Piszczek, P. Schuck, SEDPHAT--a platform for global ITC analysis and
679   global multi-method analysis of molecular interactions. *Methods* **76**, 137-148 (2015).

680   56.   P. Sarkar, N. Akhavantabib, S. D'Arcy, Comprehensive analysis of histone-binding
681   proteins with multi-angle light scattering. *Methods* **184**, 93-101 (2020).