Integrating biodiversity assessments into local conservation planning: the importance of assessing suitable data sources

Thibaut Ferraille*[1,2], Christian Kerbiriou[3,4], Charlotte Bigard[5], Fabien Claireau[3,4], John D. Thompson[1]

[1] CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France
[2] Naturalia environnement, Avignon, France
[3] Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum national d'Histoire naturelle, Centre National de la Recherche Scientifique, Sorbonne Université, Paris, France
[4] Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum national d'Histoire naturelle, Station de Biologie Marine, Concarneau, France
[5] Agence Bretonne de la Biodiversité, Brest, France

*Corresponding author
Correspondence: thibaut.ferraille@gmail.com

ABSTRACT

Strategic Environmental Assessment (SEA) of land-use planning is a fundamental tool to minimize environmental impacts of artificialization. In this context, Systematic Conservation Planning (SCP) tools based on Species Distribution Models (SDM) are frequently used for the elaboration of spatially exhaustive biodiversity diagnostics. Despite the paradigm of "garbage in – garbage out" that emphasises the importance of testing the suitability of data for SDM and priority conservation areas, the assessment of database sources remains relatively rare. In addition, the lack of practical recommendations for the use of open-access databases by SEA stakeholders remains a problem. The aim of this study is to explore the quality of data sources that can be used in SEA to assess priority conservation areas in SEA. The study used data for nine taxonomic groups (commonly used in inventories for environmental impact assessment) and three databases available to SEA stakeholders. Three local administrative entities in very different socio-ecological contexts were used to examine three main issues : (i) the suitability of local versus regional or country databases for assessing conservation priorities, (ii) differences among taxonomic groups or territories in terms of the suitability of databases, (iii) the importance of the quality of databases for the application of SDM to assess priority conservation areas. Our study provides several clear messages for potential users of open-access databases. First, the need for prudence in the interpretation of biodiversity maps. Second, the collection of individual databases at the country scale is necessary to complete local data and ensure the suitability of SDM in a local context. Third, a data driven approach can lead to the use of notably different species communities to identify priority conservation areas when compared to the community in the original database. Finally, we propose a workflow to guide SEA stakeholders through the process of data rationalization and use in conservation planning.

Keywords: Data-driven approach, Species Distribution Models (SDM), Strategic Environmental Assessment (SEA), Systematic Conservation Planning (SCP)

1 - Introduction

Land-use change, in particular urban land expansion, leads to artificialization of habitats and soils and is one of the major causes of the loss of biodiversity (Maxwell et al., 2016; IPBES, 2019). The reduction and fragmentation of natural habitats leads to population declines and species extinction (Fahrig, 1997; Horváth et al., 2019; Lino et al., 2019), as well as biotic homogenization, i.e. mostly the extinction of specialist species and the introduction of exotic species, which involves an increase in genetic, taxonomic and functional similarity (Olden and Rooney, 2006; Zambrano et al., 2019).

A major tool to limit artificialization is the mitigation hierarchy used in environmental assessments studies. This approach consists of three sequential steps: "avoid" impacts, "reduce or minimize" impacts not avoided and "offset" residual impacts (Bull et al., 2016; Maron et al., 2016). However, in the current application of the mitigation hierarchy several weaknesses prevent it from achieving the goal of "No Net Loss" of biodiversity (Quétier et al., 2014; Bezombes et al., 2019). Avoidance is poorly implemented despite the fact that it is the first and most efficient step of the hierarchy (Bigard et al., 2017; Phalan et al., 2018). What is more, the mitigation hierarchy is mostly applied in a project-by-project approach without scaling up (Pope et al., 2013; Bigard et al., 2017), which limits proper consideration of fragmentation issues (Gontier et al., 2006) and cumulative impacts (Whitehead et al., 2017), including those of multiple small projects (Bigard et al., 2017).

To anticipate avoidance measures, Strategic Environmental Assessment (SEA) of land-use planning is a global and fundamental tool to minimize environmental impacts (Baker et al., 2005). SEA provides for the integration of avoidance measures early in the land-use planning process through environmental assessment of policies, plans and programs (Fundingsland Tetlow and Hanusch, 2012; Bigard et al., 2020). However, the implementation of SEA is often based on biodiversity diagnostic maps that are rarely complete and exhaustive. Indeed, biodiversity diagnostics are rarely exclusively based on empirical observations from field surveys (Phalan et al., 2018) and usually use areas and documents already known (e.g., protected areas and green infrastructures).

Spatial modelling provides a tool for the elaboration of spatially exhaustive diagnostics of biodiversity maps for land-use and conservation planning (Almenar et al., 2019; Tarabon et al., 2019; Bigard et al., 2020; Tulloch et al., 2019; Baker et al., 2021; Boileau et al., 2022). Among these methods of biodiversity modelling, Species Distribution Models (SDM) are widely used to predict suitable habitat for species based empirical observations (Guisan et al., 2017; Zurell et al., 2020) and are increasingly used in conservation planning (Guisan et al., 2017; Domisch et al., 2019; Baker et al., 2021). Systematic Conservation Planning (SCP) tools are also particular pertinent to identify priority biodiversity stakes and avoid the adoption of an *ad hoc* approach (Margules and Pressey, 2000; Pressey and Bottrill, 2008) in order to inform SEA (Tulloch et al., 2019).

The management of databases and their use for conservation planning is a critical issue for the application of such methods to practical conservation planning. The databases available for SEA stakeholders (i.e. decision makers, environmental consultants and conservation managers) are often limited because of data sensitivity or ownership issues, although more and more programs contain data that are publicly available and use of them can be made without any particular attention to their quality (Costello and Wieczorek, 2014; Tittensor et al., 2014) and they are generally unfamiliar to SEA stakeholders. Surprisingly however, despite the prevailing recognition of the "garbage in – garbage out" that emphasises the critical importance of the quality of data (Sanders and Saxe, 2017; Canbek, 2022), an examination of data suitability is relatively rare in local conservation planning (Rondinini et al., 2006; Hermoso et al., 2015a). In this context, some authors argue the necessity of examining the sensitivity of model results to the nature of the datasets that are used (Sanders and Saxe, 2017; Clare et al., 2019; Velazco et al., 2020). SDM studies generally use data that has not been designed specifically for this type of analysis, and is often comprised of presence-only data, hence the need for a rigorous assessment of sampling biases (Beck et al., 2014; Botella et al., 2018; Guisan et al., 2017). Another particularly important point that can influence distribution modelling is the spatial extent of the data, and in particular the question of whether to use only local data or those collected on a larger scale (Baker et al., 2021; Meyer, 2007). The choice and possible combination of data sources is part of this problem due to the fact that they often vary considerably in their design, the gradients covered, and potential sampling biases (Fletcher et al., 2019; Boyd et al., 2023). Basically, the use of available databases requires a rigorous test of their quality and pertinence (Zuckerberg et al., 2011), especially when

used for analyses such as SDM (Tulloch et al., 2016; Domisch et al., 2019). Confidence in the models must be
assessed through the use of metrics adapted to the data (Guisan et al., 2017; Leroy et al., 2018). As
recognised by Clare et al., (2019), the lack of practical recommendations for the use of databases that differ
in terms of their quality and pertinence by public authorities or other institutions remains a serious problem.
The overall goal of this study is to test the influence of different database sources that can be used by
SEA stakeholders to map priority conservation areas in SEAs based on SCP. To do so, we studied three local
administrative territories that occur in different socio-ecological contexts in France. The study has three main
objectives. First, we assess the content of three open-access databases for nine taxonomic groups commonly
used in naturalist inventories in environmental assessment studies. We evaluate their suitability in terms of
data quantity for SDM application, at three scales (local, regional and country). SDM and SCP analyses were
performed for two taxonomic groups (Aves and Papilionidae) to test the hypothesis that sampling bias and
differences in ecological response scales of species may influence the identification priority conservation
areas. Second, we explore the influence of databases on the application of SDM to assess priority
conservation areas. Third, we analyse the influence of this data-driven approach on the composition of
species communities that are ultimately used in the identification of priority conservation areas relative to
the actual communities in the original databases.

2 - Methods

2.1 - Study sites

116   To assess the availability and suitability of pertinent data sources, we selected three French local
117 administrative entities in charge of land-use planning: Lodévois-Larzac (T1), Brocéliande (T2), La Rochelle
118 (T3). We selected these study sites on the basis their contrasting social, ecological and geographical contexts
119 in order to examine patterns of variation of data suitability among sites (Figure 1,

Table **1**). For example, each of the three study sites have different ecosystems and bioclimates and the sites vary in terms of urbanization pressures from 3% to 28% artificial land-use cover for sites T1 and T2 respectively. This is due to the presence of a major city (La Rochelle) in the T2 study site. The major towns in the other two study sites are smaller however there is a major city less than 50 km away for both of them (Montpellier and Rennes respectively). Only site T1 has an important cover of protected areas (Natura 2000) area.

Figure 1 - Localization of the study sites in French administrative regions: T1 is Lodévois-Larzac, T2 is La Rochelle, T3 is Brocéliande. Source: IGN, Google, 2023.

Table 1 - Description of three study sites in France

Study site		Lodévois-Larzac (T1)	La Rochelle (T2)	Brocéliande (T3)
Country / Administrative region		France / Occitanie	France / Nouvelle-Aquitaine	France / Bretagne
Main city coordinates		43°43'57"N 3°19'02"E	46°09'35"N 1°09'05"W	48°00'08"N 2°05'48"W
Area		554 km^2	327 km^2	298 km^2
Climate		Mediterranean	Oceanic	Oceanic
Major habitats / land-use		Mediterranean forest and scrubland	Intensive agricultural field, swamps, few forests	Pastures, temperate forest
% land-use [1]	Artificial – urban areas	3%	28%	7%
	Farmland	3%	55%	29%
	Semi-natural	94%	17%	64%
Landscape and dynamics		Rural, woodland expansion	Coastal, intensive agriculture, urban expansion	Rural, agricultural intensification
Urbanization context		Close to a major city (Montpellier), linked by a highway	Economic and tourist dynamism, major city presence	Close to a major city (Rennes), linked by a highway
% protected area [2]	Regulatory	0%	327 ha – 0.99%	0%
	Land control	1618 ha – 0.6%	148 ha – 0.45%	80 ha – 0.25%
	Contractual	38096 ha – 68.8%	3534 ha – 10.7%	586 ha – 2%
	Total	38602 ha – 69.7%	3546 ha – 10.8%	592 ha – 2%

(1) source: OSO Land Cover (Inglada et al., 2019). Artificial is impermeable surfaces; crop field is annual crop and orchards and vineyards; semi-natural is hardwood and soft wood and grasslands and water.
(2) source: https://inpn.mnhn.fr/

2.2 – Workflow of analysis

131
132 A methodological framework was developed to test the influence of different database sources in
133 mapping priority conservation areas in SEAs thanks to a SCP approach, all steps are summarized in Figure 2
134 and in the following text.

2.3 – Databases available for SEA stakeholders

135
2.3.1 – Content of available databases

136
We focused on nine taxonomic groups commonly used for naturalist inventories for environmental
impact assessment studies: Amphibia, nesting Aves (hereafter name Aves), Chiroptera, Flora, Mammalia
aptera (hereafter name Mammalia), Orthoptera, Odonata, Papilionidae and Reptilia (Bigard et al., 2017;
Guillet et al., 2019; Iorio et al., 2022). The three spatial scales used for data collection are depicted in Figure
1: the local scale (i.e. study site with a 10km buffer around), the regional scale (i.e. French administrative
regions); and the country scale (i.e. continental France). We selected three open-access databases containing
these groups that can be widely used by SEA stakeholders for the assessment and hierarchy of conservation
priorities (Figure 2, step 1.1). This study thus directly addresses SEA stakeholders (i.e. decision makers,
environmental consultants and conservation managers) using the databases available to them.
The first of these databases concerns the French Natural and Landscape Information System (SINP) that is
structured at the scale of French administrative regions in charge of data extraction requests. Each site has
its own database, can be collected only at local scale due to the limited extent of data requests, without the
need for a special request (maximum 2000 km^2, i.e. nearby 10km buffer zone around the study site). This
database is composed of opportunist observations and only contains presence data for taxa for which
identification is confirmed by experts (Jomier et al., 2018) (see Appendix A.1).
The second database is the Global Biodiversity Information Facility (GBIF) an international platform for
the provision of biodiversity data that is based on information collected from various databases (Telenius,
2011). It is composed of observation data that are not based on protocols and for which presence data and
identification are not subject to expert confirmation. The data downloading is autonomous from the website
(see Appendix A.1).
The third database is a French biodiversity monitoring scheme (Vigie Nature) dedicated to assess spatio-
temporal populations trends. Within this monitoring scheme data collection is based on a standardized
biodiversity survey. Despite local spatial distribution heterogeneities, the sampling plan ensures a
representation of the current distribution of habitats and landscapes across France (Julliard and Jiguet, 2002;

Mariton et al., 2022). Homogeneity in identification criteria and compliance with the protocol are ensured by
offering training to volunteers. This database is composed of presence/absence and abundance data. Access
to these databases requires a data extraction request to the person in charge (see Appendix A.1).
These three databases were combined in two ways: "All databases" (i.e. the combination of SINP, GBIF
and Vigie Nature) used in section 2.3 (i.e. for assess which taxa are enough documented within each dataset)
and "National databases" (i.e. the combination of GBIF and Vigie Nature , which are available at country
scale) used specifically in section 2.4 (i.e. to test the effect of database sources on SDM performance).
The databases were collected for continental France except for the SINP that was collected in 10km
buffer zone around the study sites due to the access restrictions explained above. The databases were
collected over a period of 11 years (i.e. from 01/01/2010 to 31/12/2020) and data with spatial inaccuracy
greater than 50 meters were not considered. We made a series of operations to standardize, correct and
homogenize taxa names at the specific taxonomic level using the French taxonomic reference "TAXREF.V14"
(Gargominy et al., 2021). We transformed data into occurrences and limited their sampling biases by
geographical filtering using "spThin" package (Aiello-Lammens et al., 2015). In each study site, we identify
the presence of one species at least five observations from "All databases" combined at local scale and
defined as present species in France by TAXREF.V14 (Appendix A.1).

2.3.2 – Quantitative description of databases for SDM
For the nine taxonomic groups, four metrics were selected to quantitatively describe the amount and
thus the suitability of each database (i.e. SINP, GBIF, Vigie Nature and "All databases") for the realization of
SDM in presence-only (Figure 2, step 1.2): (i) the number of species observed in the study site; (ii) the
proportion of species with < 15 observations which represents the minimal threshold for the utility of SDM
with more accurate predictions than in a random model (Støa et al., 2019); (iii) the proportion of species with
between 15 and 50 observations, i.e. the minimum number of presences recommended for SDM (Merow et
al., 2014 in Guisan et al., 2017); (iv) the proportion of species with > 50 observations, i.e. highly suitable for
modelling (Støa et al., 2019). These metrics were calculated at three different scales: (a) for each study site
including a 10km buffer zone (local scale) that is the maximum extent for a SINP data request; (b) on a
regional scale that is used for the structure of biodiversity data in France; (c) for continental France. SINP
database is analysed individually only at the local scale due to the previously mentioned restriction of access,
nevertheless it integrated the three scales of the "All databases".

2.4 - Systematic conservation planning process
For the three study sites, we identified priority conservation areas with a Systematic Conservation
Planning (SCP) tool based on Species Distribution Models (SDM) (Figure 2, steps 2.1 and 2.2). Several variants
of SDM were made using different database sources (i.e. GBIF, Vigie Nature and "National databases") and
two methods of generating pseudo-absences (named individual database or mixed databases). Among the
nine taxonomic groups studied above, only Aves and Papilionidae taxa were analysed to compare the tests in
the SCP process. The data available in France for these groups is sufficient in quantity to realize SDM with
each database. The use of these two groups allows for a comparison between one group of highly mobile
taxa with a large home range (Aves) and another group with a smaller home range and whose movement
closely tracks local environmental variation (Papilionidae). These two taxonomic groups thus have different
biological traits associated with their dispersal and function, hence we predict differences in in terms of the
spatial resolution of their distribution.

2.4.1 - Species distribution modelling (SDM)
We modelled favourable habitats for birds and butterflies in the three study sites using SDM (Figure 2,
step 2.1). A resolution of 50m was used to meet the needs of the SEA of land-use planning. A buffer zone of
10km around each of the study sites (i.e. local scale) was used for the SDM prediction to limit any border
edge effects and to increase the number of species that could be modelled and evaluated. Indeed, species
with less than 15 data points for the calibration (threshold explain above, Støa et al., 2019) and/or less than
10 data for performance evaluation (threshold defined by expert opinion) were not modelling.
Biodiversity data used for the SDM came from the databases described above at the country scale,
according to the results of section 2.3 (Table 2). These data were separated into two independent datasets
that allow for robust validations with independent data (Matutini et al., 2021). Country data without local

data were used for model calibration and local data were used only for model performance evaluation.
Therefore, the SINP database, which is only available at local scale, was not used for model calibration.
For model calibration, pseudo-absences were generated with two methods, separately for each
taxonomic group. First, methods to generate pseudo-absences in the individual databases for all their data
(i.e. GBIF, Vigie Nature and "National databases"). For databases with a protocol for sampling (i.e. Vigie
Nature) to optimize species detection (day and year periods), the absence points were defined as all the
points without the observed species. For databases without such sampling protocols for all their data (i.e.
GBIF and "National databases"), pseudo-absence data were generated with the target-group (TG) approach,
which infers the sampling bias from the aggregated occurrences of (TG) species, i.e. the respective taxonomic
groups (Ponder et al., 2001; Anderson, 2003; Phillips et al., 2009). Second, a method to mix the presence
data in the GBIF and "National databases" with the absence from Vigie Nature (named mixed databases) was
applied (Hermoso et al., 2015a).
Three types of environmental variables were used for SDM: geographic, human occupancy and pollution
and fragmentation (Appendix A.2).
SDM were calibrated by Random Forest down-sample (Valavi et al., 2021a) which according to Valavi et
al. (2021b) is among the best performing models for presence-only data. Although the Random Forest is not
very sensitive to the non-independence of the variables and over-parametrization (Matsuki et al., 2016;
Srisa-An, 2021), in order to be parsimonious, the collinear variables were removed (Pearson >0.7, Appendix
A.2, Brun et al., 2020). Thirty bootstraps were performed for each SDM (Guisan et al., 2017) using a
calibration for 70% of the data at the country scale outside of local scale. The thirty Random Forest
bootstraps were combined with mean to provide an ensemble prediction of habitat suitability for all species.
The performance evaluation of the models was done using the Boyce's index (CBI), the most suitable
metric for model in presence-only (Boyce et al., 2002; Leroy et al., 2018), with local data (i.e. "All databases"
combining SINP, GBIF and Vigie Nature). Dubos et al. (2022) reveal the CBI turns out to be misleading in some
cases, thus we used a threshold of 0.3 to define good or poor model quality.

2.4.2 – Systematic conservation planning (SCP) tool
Priority conservation areas in the three study sites were analysed from SDMs for each database source in
using a SCP tool (Figure 2, step 3.1). To meet the needs of SEAs, the study site was restricted to
administrative boundaries with a buffer zone of 1km to maintain coherence between administrative entities
and a resolution of 50m. The aim of SEA biodiversity conservation strategies is to establish priorities for the
whole study site as a whole and all the cells have the same cost value of 1. The objective was a maximum
coverage objective that seeks to maximize the number of features, i.e. the SDMs (Church et al., 1996). The
features were only the good quality models defined previously. The priority conservation areas decisions
were between 0 and 1. To obtain a priority gradient, we cumulated Ferrier importance scores (Ferrier et al.,
2000) from nine targets of the total amount of each feature (from 0.1 to 0.9 every 0.1). We used the package
"prioritizr" (Hanson et al., 2021) with the open-source solver SYMPHONY (Kim et al., 2023).

2.5 - Comparative analysis of SDM and priority conservation areas
249   We analysed the influence of database sources on SDM predictions and priority conservation areas
250 (Figure 2, step 2.2 and 3.2). SDM performance evaluations were analysed between database sources. The
251 SDM prediction and priority conservation areas maps were compared with the Spearman's rank coefficient
252 (Phillips et al., 2009) and the Schoener's D index as a measure of projection overlap (Schoener, 1968) which
253 was calculated with the ENMTool R package (Warren et al., 2008; Warren and Dinnage, 2022).

2.6 - Species community analysis
We assessed the influence of the complete data driven workflow on the composition of species
communities, i.e. differences between the original community (i.e. all species observed in study site) in the
database and the final community used to identify priority conservation areas (Figure 2, step 4).
To do so, we developed a workflow resistance score for each of the methodological filters for all species.
A score of 1 is allocated to species observed in the study area that did not cross any of the stepwise filters. A
score of 2 is allocated to species with sufficient data to calibrate SDM, i.e. > 15 country observations, or to
evaluate the performance of SDM, i.e. > 10 local observations. A score of 3 is allocated to species with
sufficient data to calibrate and evaluate the performance of the SDM. A score of 4 is for species that were

present in the final analysis as a priority species for conservation planning (i.e. with the two previous filters
and a Boyce's index > 0.3). In order to assess species composition bias ultimately considered in priority
conservation areas, species communities were analysed through traits that can influence species detection
(mass, displacement capacity, period activity) and ecological traits (habitats, specialisation) (Appendix A.3).
Missing data were completed with a trait imputation procedure generated using the R package "missForest"
(Stekhoven and Bühlmann, 2012) by considering evolutionary relationships in the imputation process (see
Carmona et al., 2021) using the R script of Toussaint et al. (2021). Due to the nature of the response variable
(i.e. ordinal scoring including four modalities), we used ordinal regression mixed models with cumulative link
using the clmm function of "ordinal" R package (Christensen, 2022). We adapted the link function to the data
distribution, using a "cauchit" link for Aves and a "logit" link for Papilionidae. Species traits were used as
fixed effects, while the random effects selected were the study sites for Aves and the combination for study
sites and database sources for Papilionidae. Finally, we evaluated the quality of the full model by comparing
to the null model with Akaike's information criterion (AIC) (Mac Nally et al., 2018).
Thus, our models were structured in the following way:

For Aves $Resistance \sim \beta_0 + \beta_i \cdot 'trait'_i + (1|StudySites)$

For Papilionidae $Resistance \sim \beta_0 + \beta_i \cdot 'traits'_i + (1|StudySites/Databases)$

Where β is the parameter estimates, i correspond to the variables of 'traits' using in fix effect, 0 is the
shift between ordinal class of resistance (i.e. 1|2, 2|3 and 3|4) and "1|" is the random effect.

Figure 2 - Methodological framework applied to test the influence of different database sources to map conservation planning areas in SEAs based on SCP. The green and blue boxes are the appendices detailing the methodology and the results of the analyses, respectively. The red boxes "R." are the methodological filters used for generate workflow resistance score.

 3 - Results

3.1 - Quantitative description of databases for SDM
The use of individual observation databases provided a limited number of observation data for
distribution modelling of many species, regardless of the taxonomic group (Table 2, Appendix B.1). None of
the GBIF and SINP individual databases have more than 50 observations for all nine taxonomic groups, and
there are currently no programs for three of the nine taxonomic groups in the Vigie Nature databases. Local
databases and, to a lesser extent, regional databases are not equivalent in terms of the amount of data
available for different taxa for the three study sites. The difference is particularly pronounced for the SINP
databases, with no data available for five taxonomic groups in the T3 study site, whereas in the T1 study site,
six taxonomic groups have sufficient data to model over 50% of the species. At the regional scale, the
number of observations per species is highly variable among study sites for the GBIF databases and is more
similar for the Vigie Nature databases.
The use of combined databases (i.e. "All databases") increased the total number of species that can be
studied. At the local scale, the proportion of species in each suitability class showed only small changes,
while at the country scale of France, it allowed a significant gain in species with suitable data (Table 2).
Indeed, on a country scale, the GBIF and Vigie Nature databases are complementary with each other. For
example, the GBIF has few Chiroptera data, which is complemented by Vigie Nature data, and vice versa for
Amphibian data.
At the country scale, aggregation of the databases seem to provide the most suitable setup (databases
and scale) for SDM analysis. Using these compiled, country databases provides a large amount of data for a
large number of species present in the three study sites (Table 2 and Appendix B.1).

3.2 - SDM and priority conservation areas analysis
The evaluation of SDM revealed differences among the database sources; none of which produced more
than 87% of satisfactory models for the two studied taxonomic groups and some had less than 20% of
satisfactory models (Table 3, Appendix B.2). Use of the GBIF data led to a higher proportion of well-evaluated
SDM, ranging between 48 and 79% of satisfactory models for the species in the two taxonomic groups. GBIF
data are also more suitable than Vigie Nature data, they produced between 11% and 37% more satisfactory
models than the latter database (Table 3). Nevertheless, between 4% and 9% of species provide well-
evaluated models from Vigie Nature and poorly evaluated by GBIF database. The combination of "National
databases" (i.e. GBIF and Vigie Nature) decreased the performance of SDMs with GBIF data, but still yield
better results than SDM based on the Vigie Nature database. The substitution in individual databases of
pseudo-absences for the absences from Vigie Nature (i.e. mixed databases) reduced the performance of
models based on GBIF, but increased the performance of the combined "National databases" (Figure 3,
Appendix B.2). Regardless of the database used, our analyses revealed significant differences between study
sites (Table 3). No SDMs for butterflies in T2 could be evaluated due to insufficient local data. For the T3
study site, over 50% of the Papilionidae and Aves models perform poorly, whereas for the T1 study site poor
models occur in less than 50% of the evaluations.
Although important differences in model performance between the database sources used for SDM
showed a high degree of overlap, as indicated by Shoener's D index with values above 0.8, the ranking of
habitat suitability was highly variable. This was in particularly the case for the GBIF and Vigie Nature
databases that had median spearman's rank coefficient values between 0.2 and 0.5 and a very wide
distribution (Appendix B.3). The substitution of pseudo-absence data in GBIF and "National databases" by
absence data from Vigie Nature, showed a similar situation (Appendix B.4).
For priority conservation areas, whatever the individual databases used, the overlaps with Schoener's D
index were above 0.72 and similar in each study site. Nevertheless, Sperman's rank coefficients showed a
greater difference in prioritization ranks in particular between GBIF and Vigie Nature and for the T3 study
site (Table 4). Between maps of priority conservation areas, we observed similarities in overlap, despite a
significant difference in the hierarchy of areas to be prioritized (Figure 4, Appendix B.5). The list of species is
presented in Appendix B.7, where it can be seen that there are no difficulties with respect to invasive species
which are very few in the data sets.

3.3 - Species community analysis

The distribution of workflow resistance scores showed that only 30% and 42% respectively of Aves and Papilionidae species were integrated in priority conservation areas maps. Among species not integrated, the workflow steps filtering the most species concern the amount of suitable data for model evaluation followed by the quality of the models (Figure 5). The analysis of the species community composition observed in each of the three study sites in comparison with the species community integrated in priority conservation area identification revealed significant differences for all three study sites. For Aves communities, the differences concern an under-representation of nocturnal species, large species with high dispersal capacity, and species of swamp habitats and deciduous forests in relation to the observed species community in the databases for the three study sites. Conversely, species that favour urban habitats, shrubland, grassland and coniferous forests are over-represented in the species community of the final maps, as are species with specialized diets and foraging strata (Table 5). For Papilionidae communities, common species with long flight periods are over-represented in the final community used for analysis. Species related to anthropogenic and thermo/meso Mediterranean habitats, and species that use a wide range of hostplants are over-represented in relation to the original species community, while supra-Mediterranean species and those of montane environments are under-represented (Table 5).

Table 2 - Quantitative description of the observation databases (Local SINP, GBIF, Vigie Nature) for nine taxonomic groups in the three study sites ("T1": Lodévois-Larzac, "T2": La Rochelle, "T3": Brocéliande) for "local" (study site with buffer of 10km), "Regional" (administrative region) and "Country" (continental France). "nSp" is the number of species observed at least 3 times in a given study site within a 10 km buffer zone, ">15obs" is the percentage of species with more than 15 observation and ">50obs" the percentage with more than 50 observations (red < 25%, orange 25-75%, green >75%, grey - no data).

Taxa		GBIF Local T1	T2	T3	GBIF Regional T1	T2	T3	GBIF Country T1	T2	T3	SINP Local T1	T2	T3	VN Local T1	T2	T3	VN Regional T1	T2	T3	VN Country T1	T2	T3	All Local T1	T2	T3	All Regional T1	T2	T3	All Country T1	T2	T3
Amp	nSp	11	1	11	13	2	11	14	2	11	13	2	0	\multicolumn{9}{No program}										14	2	11	14	2	11	14	2	11
	>15obs																															
	>50obs																															
Ave	nSp	100	73	77	198	159	120	204	170	131	197	94	82	114	155	101	182	143	120	201	170	132	205	171	132	205	166	132	205	171	132
	>15obs																															
	>50obs																															
Chi	nSp	4	0	5	13	13	5	19	19	21	21	6	18	25	22	24	25	22	24	25	22	25	26	22	26	26	22	26	26	22	26
	>15obs																															
	>50obs																															
Flo	nSp	786	476	639	1428	1093	825	1573	1261	995	1703	1335	843	286	177	136	825	657	413	1336	1105	848	1844	1460	1093	1844	1158	1093	1844	1460	1093
	>15obs																															
	>50obs																															
Mam	nSp	10	1	14	23	6	21	23	6	28	23	7	30	No program									23	7	30	23	6	30	23	7	30
	>15obs																															
	>50obs																															
Odo	nSp	38	5	23	56	7	28	57	7	30	57	1	0	0	1	24	56	7	27	57	7	30	57	7	30	57	7	30	57	7	30
	>15obs																															
	>50obs																															
Ort	nSp	53	3	24	93	25	26	95	25	34	91	8	0	25	19	19	25	19	19	25	19	19	98	25	34	98	25	34	98	25	34
	>15obs																															
	>50obs																															
Pap	nSp	115	15	33	144	18	43	147	18	52	154	11	0	0	0	50	132	18	51	149	18	52	154	18	52	154	18	52	154	18	52
	>15obs																															
	>50obs																															
Rep	nSp	18	2	7	20	4	7	21	4	7	21	4	0	No program									22	4	7	22	4	7	22	4	7
	>15obs																															
	>50obs																															

Table 3 - Proportion of well-evaluated SDM corresponding to Boyce's index greater than 0.3. "National" combines GBIF and Vigie Nature databases.

Taxa	Study site	Number of species	Individual database GBIF	Vigie Nature	"National"	SDM with Vigie Nature absence GBIF	"National"
Aves	T1	114	79 %	61 %	70 %	87 %	82 %
	T2	87	68 %	31 %	47 %	57 %	46 %
	T3	56	48 %	38 %	29 %	52 %	43 %
Papilionidae	T1	127	75 %	52 %	64 %	75 %	73 %
	T3	25	52 %	16 %	20 %	12 %	20 %

Figure 3 - Proportion of SDM with Boyce's index (CBI) greater than 0.3 by database source
("Individual" and "Mixed" with Vigie Nature absence), by combining study sites (T1, T2, T3) and
taxonomic groups (Aves, Papilionidae). "National" combines GBIF and Vigie Nature databases.

Table 4 - Overlap of priority conservation areas between database sources using two metrics:
Schoener's D index (D) and Spearman's rank coefficient (S cor). VN is Vigie Nature database and
"National" combines GBIF and Vigie Nature databases.

| Taxa | Study sites | Individual database | | | | | | Pseudo-absence of individual database - absence from VN | | | |
| | | "National" – GBIF | | "National" – VN | | GBIF - VN | | GBIF | | "National" | |
		D	S cor	D	S cor	D	S cor	D	S cor	D	S cor
Aves	T1	0.82	0.56	0.81	0.53	0.78	0.36	0.75	0.88	0.79	0.38
	T2	0.82	0.74	0.76	0.49	0.72	0.42	0.76	0.64	0.79	0.64
	T3	0.77	0.34	0.81	0.51	0.78	0.29	0.73	0.19	0.79	0.39
Papilionidae	T1	0.85	0.64	0.84	0.65	0.83	0.59	0.72	0.66	0.82	0.66
	T3	0.80	0.49	0.77	0.31	0.78	0.22	0.76	0.17	0.72	0.06

369　**Figure 4** - Maps of priority conservation areas for Aves and Papilionidae of tree study sites (T1, T2,
370　T3) from different individual database source (All, GBIF, Vigie Nature).

Figure 5 - Workflow resistance scores for Papilionidae and Aves. 1 - species observed in the study area that did not cross any of the stepwise filters. 2 - species with sufficient data to calibrate SDM, i.e. > 15 country observations, or evaluate the performance of SDM, i.e. > 10 local observations. 3 - species with sufficient data to calibrate and evaluate the performance of the SDM. 4 - species present in the final analysis, i.e. Boyce's index > 0.3.

Table 5 - Parameter estimates (β), standard error (se) and P-values for the full model of Aves and Papilionidae species resistance to the workflow. Appendix B.6, the evaluation of the quality of the model.

		Aves				Papilioniadae		
	Variables	β	se	P-value	Variables	β	se	P-value
Full model	β_0 1|2	-7.84	1.12	/	β_0 1|2	-2.54	1.28	/
	β_0 2|3	0.14	0.33	/	β_0 2|3	1.39	1.26	/
	β_0 3|4	1.4	0.34	/	β_0 3|4	2.97	1.27	/
	Mass	-0.21	0.13	.	WingspanM	0.01	0.09	
	Avian hand-wing index	-0.31	0.07	***	FMoMean	0.38	0.11	***
	Nocturn	-2.28	0.59	***	Hostplant N	0.23	0.11	*
	Deciduous	-0.27	0.15	.	Hostplant Spe	0.16	0.11	
	Coniferous	0.38	0.15	*	HPG Bi	0.28	0.50	
	Woodland	0.17	0.14		HPG Th	-0.21	0.22	
	Shrub	0.55	0.13	***	HPG Sb	-0.32	0.23	
	Grassland	0.47	0.13	**	HPG Tr	0.22	0.45	
	Mountain meadows	0.23	0.24		HPG Li	0.02	0.36	
	Reed	0.08	0.33		SSI	0	0	
	Swamps	-0.84	0.28	**	AltVeg 'A'	0.06	1.12	
	Rocks	-0.14	0.21		AltVeg 'Mo'	-0.93	0.31	**
	Urban	0.83	0.14	***	AltVeg 'SupMed'	-0.79	0.46	.
	Spe. Diet	0.92	0.35	**	AltVeg 'ThMeMed'	1.37	0.37	***
	Spe. Foraging behav.	-0.08	0.4		Rarity '2'	0.86	0.29	**
	Spe. Diet strat	1.49	0.4	***	Rarity '3'	1.42	0.34	***
	Spe. Habitat	-0.14	0.38					
	Spe. Nest	-0.28	0.72					
	Spe. Mean	-2.26	1.63					

P-value: *** P < 0.001, ** P < 0.01, * P < 0.05, . P < 0.1, / P-value not applicable
Trait description: Spe. is the specialization, FMoMean is duration of yearly flight period, Hostplant N is hostplant specificity, HostplantSpe is hostplant specificity index, HPG is hostplang growth form (Bi: short herb, Th: tall herb, Sb: shrub, Tr: tree, Li: Liana), SSi is Species Specialization Index, AltVeg is altitudinal vegetation (A=Anthropogenic, Co: Foothill, TheMeMed: Thermo/Meso-Mediterranean, Med: Mediterranean, SupMed: Supra-Mediterranean , Mo: Montane, Asa: Alpine and Subalpine). For more details on traits see Appendix A.3

4 - Discussion

The absence of recommendations for the use of available databases that differ in terms of their quality and pertinence by public authorities or other institutions remains a serious problem for local conservation planning (Clare et al. 2019). The goal of this study was to test the suitability of different database sources that can be used by public stakeholders to map priorities for biodiversity stakes in SEAs and SCP. We found that the compilation of databases at the country scale is the most suitable procedure to apply SDM to a large number of species. For Aves and Papilionidae, the GBIF database provided the highest proportion of well-assessed SDM. We detected a significant overlap in species distributions in different database sources despite significant variability in the order of habitat suitability and similar spatial predictions for priority conservation areas. We showed that the composition of the species community used for priority conservation areas in all three study sites were clearly not representative of the observed species communities in the original database (in terms of species and ecological traits). Finally, despite important differences among the study sites in terms of the proportion of artificial land cover and protected areas we found no particular differences between the three study sites. Clearly, the data sources are the most important factor influencing the results.

Open-access biodiversity data provide a valuable source of information for decision makers, environmental consultants and conservation managers (i.e. SEA stakeholders); they contain vital information on species locations compared to expert knowledge and unshared datasets that are inaccessible to most users (Sousa-Baena et al., 2014; Meyer et al., 2015). Through their use and careful application of SDM, they contribute to the estimation of relative habitat suitability in a given study site (Baker et al., 2021). Our study showed however that SDM for a large majority of species observed locally requires their compilation on a country scale. Local and regional data are not suitable for model calibration but remain important for assessing the suitability of models in a local context. Indeed, this performance evaluation step is one of the most restrictive filters in the workflow we proposed, as evidenced by SDM's for butterflies in the T2 study site, where no species could be evaluated. The spatial extent of data collection can influence distribution modelling (Meyer, 2007), and our study emphasises this importance for SDM and the use of data available on country scale in France.

Different types of databases are constructed in different way – with opportunistic data collection or, in some cases, as part of a scientific monitoring scheme – allowing the use of as complementary data sources (Beck et al., 2013; Shirey et al., 2021). The three individual databases examined in this study are indeed complementary in that when they are combined they provide a suitable source for modelling the distribution of many species. Nevertheless, some groups commonly have a low amount of data in such bases, e.g. Insecta (Troudet et al., 2017). The study of data with a fine spatial grain, as required for SEA and SCP (Guisan et al., 2013), reveals information gaps for more taxonomic groups at the global scale, e.g. Amphibians and Mammals (Witté and Touroult, 2017). The construction of an overall database at country scale is therefore the most appropriate way to have suitable data for SDM of different taxonomic groups. Furthermore, there is a dilemma between protocolized and opportunistic data. Although protocolized data are recommended for SDM (Guisan et al., 2017; Guillera-Arroita et al., 2015), very often the amount of such data is low, which can be detrimental at the local scale, particularly for model evaluation with data having the same sampling bias. For opportunistic data, their large number is of course a positive point, however the estimating their sampling bias can be a real challenge (Botella et al., 2018; Fithian et al., 2015; Matutini et al., 2021) to ensure the reliability of the results.

SDM of Aves and Papilionidae species clearly revealed differences between the databases used for modelling, with differential impacts on the identification of conservation priorities. Indeed, the high overlap in species distribution between data sources, as indicated by Schoener's D index, indicates that, regardless of data source, species are predicted in similar environments (Warren et al., 2008). However, Spearman's ranking of habitat suitability between data sources was highly variable, indicative that species' responses to environments are highly variable, as are the location of favourable habitats (Warren et al., 2008). Although the use of presence-absence data is advocated for SDM (Guillera-Arroita et al., 2015; Valavi et al., 2021b; Dubos et al., 2022), we showed that opportunist data from GBIF provided a greater number of well-assessed models at the local scale. Models using opportunist data with a target-group approach to generate pseudo-absences provides a sufficient quality of information on species distribution (Phillips et al., 2009; Barber et al., 2022) and can be correctly used in SCP (Sofaer et al., 2019;

Baker et al., 2021). The lack of data at the local scale, whatever the database, does not allow us to explain
a better fit of models using GBIF data. Evaluating the models with a large proportion of opportunist data
could however bias the evaluation, but only independent data were used, which provides robust
validation of SDM (Matutini et al., 2021). Moreover, in contrast to Hermoso et al. (2015a), we found that
mixing presence-only data with absence data increased the number of misjudged models. In addition to
the use of the ROC curve (AUC) as a presence-only model evaluation metric by Hermoso et al. (2015a),
the different results can be explained by different sampling biases between the two data types (Baker et
al., 2022; Barber et al., 2022). Finally, the GBIF data seem to be more adapted to model the distribution
of a large number of species.
The notion of "garbage in – garbage out" emphasises the critical importance of the quality of data
(Sanders and Saxe, 2017), nevertheless, the examination of data suitability for conservation planning
remains rare. In addition to the above issues our study revealed the importance of attention that should
be paid to the representativeness of the species communities used in the models compared with the
actual species communities observed in the study sites. This is particularly important in the light of the
finding that there are marked differences between conservation priorities when different database
sources are employed. Indeed, the number and composition of species in the community used can
influence conservation priorities. Elsewhere it has been shown the difference will decline as the number
of species increases (Kujala et al., 2018). The methodology tested in our study is based on a data-driven
approach that attempts to use all available biodiversity data. This approach is data intensive, but is
necessary to ensure the best representation of the observed local biodiversity. We revealed that such an
approach can nevertheless induce a significant bias in the species community that is ultimately studied.
Indeed, the prevalence of data affects the composition of the modelled species as well as the accuracy of
the models and the evaluation of the species response (Fukuda and De Baets, 2016). Particular attention
should thus be paid to the representativeness of the species communities used in the models in relation
to the actual species communities observed in the study site.
Our study presents a workflow (Figure 2) for identifying biodiversity stakes using a data-driven
approach from open-access database sources. SEA stakeholders can use this workflow as a step towards
the rationalization of data in order to reduce the biases mentioned above. The confrontation of the limits
of such a workflow with the needs of SEA stakeholders could illustrate how to precisely target new
sources of database that should be collected according to the suitability of current databases for priority
groups. This workflow could be compared with the data context of another country to compare our
findings. Hermoso et al. (2015b) revealed that evaluation models using a new collection of field data does
not necessarily reduce the problems of model uncertainty. However, other databases can be examined
by SEA stakeholders as well as other monitoring schemes (e.g. "PopAmphibien" for Reptilian and
Amphibian populations in France http://lashf.org/popamphibien-2/) or negotiate the use of databases
that are not yet shared. To overcome this data sharing problem, the structuring of networks of different
contributors of data and users of the databases and ambitious regional policies is necessary. As
evidenced by our three study sites, the quantity of local data available is correlated with the number of
years the SINP has been implemented. An important issue is thus the integration of SEA stakeholders in
the workflow we propose, and their appropriation of the procedure. This could be done by a form of
participatory modelling (Lagabrielle et al., 2010; Lees et al., 2021), where stakeholders are consulted for
issues and choices such as the species to be examined. In such participatory modelling it is important to
avoid arbitrary choices that are neither reproducible nor representative of local diversity, but rather the
result of administrative or political interest. Finally, it is currently recommended to use these tools to
elaborate a more holistic approach to SCP (Cadotte and Tucker, 2018).

5 – Perspectives: operational implementation by SEA stakeholders

Spatially exhaustive and ecologically representative priority conservation areas are crucial for the
elaboration of SEAs that aim to limit artificialization impacts as early as possible in the planning process.
Empirical observations are major sources of information on biodiversity that are still rarely used by SEAs.
The collection of open-access databases for SEA territories provides important but incomplete knowledge
on species occurrence. Furthermore, their use is particularly interesting to help strategically direct
inventory campaigns (especially for under sampled taxa and areas) that go beyond the emphasis on rare,

threatened and emblematic species. These additional data would clearly improve the assessment of the
SDMs suitability in administrative entities as our study sites. What is also interesting here for SEA
stakeholders is that the process of filtering species and attributing them a score allows for the
identification of different groups of species in terms of their needs for additional data in order to
undertake SDM.
The influence of database sources on the identification of priority conservation areas reveals the
importance of examining their suitability. In our study this is true for three highly contrasting study areas
that differ markedly in terms of the cover of protected areas and artificialisation. The problem of data
sources is thus typical of many areas. Thus, it is necessary to be prudent in the interpretation of
biodiversity maps. The integration of local experts may help limit any misjudgements in the workflow
procedure. Indeed, the integration of "expert" knowledge and local studies is valuable information, which
is important to share, and which it is important to consider in order to complete our proposal. In future
studies, species conservation issues for spatial prioritization could be considered by focusing on (for
example) the issues associated with threatened and/or invasive species. The multiple dimensions of
biodiversity could be analyzed within a context of limited data access and the complementarity of
different facets (functional and phylogenetic) in addition to a classical species-based approach (Brumm et
al., 2021; Cadotte and Tucker, 2018).
A data-driven approach that considers as many species as possible requires a large amount of data,
biases the species communities considered and does not highlight species of particular interest as their
threats and regulatory protections. It is therefore necessary to rationalize this approach, by integrating
the needs and issues of local SEA stakeholders.
511

Acknowledgements

This work would not have been possible without the help of many voluntary field workers who
continually provide open-access data. We are grateful to Benoit Fontaine for his work as administrator
and for providing access to the Vigie Nature data, to Annegret Nicolai for collection of all open-access
data on "Communauté de commune de Brocéliande (i.e. the study site "T3"), to Paul Fromage for
providing access for Fauna data, to Mathieu Largarde for OEB data access, to Solène Robert for the
exchanges about SINP database and Joel Kamdoum Ngueuko for access to SINP Occitanie data. We also
thank Josselin Giffard and Samuel Alleaume for access to DHI NDVI, to Karine Princé for discussion on
agricultural intensification variables and to Jean-Pierre Moussus for access to Papilionidae trait databases
and Jeremy Froidevaux for the preview access to the EuroBaTrait https://doi.org/10.1038/s41597-023-
02157-4. We thank Nicolas Dubos and Boris Leroy for discussion on the use of SDM and Eric Durand for
more general discussion of this study. Finally, we also acknowledge the reviewers which improved this
version.

Data, scripts, code, and supplementary information availability

526   Observation data from SINP and Vigie nature databases are not available for confidentiality reasons,
527   but their link and the request process are detailed in Appendix A1. The data of GBIF are available online:
528   https://doi.org/10.15468/dl.ry6uw7.
529   The data, R scripts, outputs of steps 1.2 to 4 of figure 2 are available online:
530   https://doi.org/10.5281/zenodo.7883973.

Conflict of interest disclosure

532   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
533   relation to the content of the article.

Funding

535   This study was funded by "Naturalia Environnement" and "Association Nationale de la Recherche et
536   de la Technologie" (grant number: 2020/0584).

References

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B., Anderson, R.P., 2015. spThin: an R package
for spatial thinning of species occurrence records for use in ecological niche models. Ecography 38,
541–545. https://doi.org/10.1111/ecog.01132
Almenar, J.B., Bolowich, A., Elliot, T., Geneletti, D., Sonnemann, G., Rugani, B., 2019. Assessing habitat
loss, fragmentation and ecological connectivity in Luxembourg to support spatial planning. Landscape
and Urban Planning 189, 335–351. https://doi.org/10.1016/j.landurbplan.2019.05.004
Anderson, R.P., 2003. Real vs. artefactual absences in species distributions: tests for Oryzomys albigularis
(Rodentia: Muridae)   in   Venezuela.   Journal   of   Biogeography   30,   591–605.
https://doi.org/10.1046/j.1365-2699.2003.00867.x
Baker, D.J., Maclean, I.M.D., Goodall, M., Gaston, K.J., 2022. Correlations between spatial sampling biases
and environmental niches affect species distribution models. Global Ecology and Biogeography 31,
1038–1050. https://doi.org/10.1111/geb.13491
Baker, D.J., Maclean, I.M.D., Goodall, M., Gaston, K.J., 2021. Species distribution modelling is needed to
support   ecological   impact   assessments.   Journal   of   Applied   Ecology   21–26.
https://doi.org/10.1111/1365-2664.13782
Baker, M., Carter, J., Short, M., Jay, S., 2005. Strategic Environmental Assessment and Land Use Planning:
An International Evaluation. Routledge.

Barber, R.A., Ball, S.G., Morris, R.K.A., Gilbert, F., 2022. Target-group backgrounds prove effective at correcting sampling bias in Maxent models. Diversity and Distributions 28, 128–141. https://doi.org/10.1111/ddi.13442

Beck, J., Ballesteros-Mejia, L., Nagel, P., Kitching, I.J., 2013. Online solutions and the 'Wallacean shortfall': what does GBIF contribute to our knowledge of species' ranges? Diversity and Distributions 19, 1043–1050. https://doi.org/10.1111/ddi.12083

Beck, J., Böller, M., Erhardt, A., Schwanghart, W., 2014. Spatial bias in the GBIF database and its effect on modeling species' geographic distributions. Ecological Informatics 19, 10–15. https://doi.org/10.1016/j.ecoinf.2013.11.002

Bezombes, L., Kerbiriou, C., Spiegelberger, T., 2019. Do biodiversity offsets achieve No Net Loss? An evaluation of offsets in a French department. Biological Conservation 231, 24–29. https://doi.org/10.1016/j.biocon.2019.01.004

Bigard, C., Pioch, S., Thompson, J.D., 2017. The inclusion of biodiversity in environmental impact assessment: Policy-related progress limited by gaps and semantic confusion. Journal of Environmental Management 200, 35–45. https://doi.org/10.1016/j.jenvman.2017.05.057

Bigard, C., Thiriet, P., Pioch, S., Thompson, J.D., 2020. Strategic landscape-scale planning to improve mitigation hierarchy implementation: An empirical case study in Mediterranean France. Land Use Policy 90, 104286. https://doi.org/10.1016/j.landusepol.2019.104286

Boileau, J., Calvet, C., Pioch, S., Moulherat, S., 2022. Ecological equivalence assessment: The potential of genetic tools, remote sensing and metapopulation models to better apply the mitigation hierarchy. Journal of Environmental Management 305, 114415. https://doi.org/10.1016/j.jenvman.2021.114415

Botella, C., Joly, A., Bonnet, P., Monestiez, P., Munoz, F., 2018. A Deep Learning Approach to Species Distribution Modelling, in: Joly, A., Vrochidis, S., Karatzas, K., Karppinen, A., Bonnet, P. (Eds.), Multimedia Tools and Applications for Environmental & Biodiversity Informatics. Springer International Publishing, Cham, pp. 169–199. https://doi.org/10.1007/978-3-319-76445-0_10

Boyce, M.S., Vernier, P.R., Nielsen, S.E., Schmiegelow, F.K.A., 2002. Evaluating resource selection functions. Ecological Modelling 157, 281–300. https://doi.org/10.1016/S0304-3800(02)00200-4

Boyd, R.J., Stewart, G., Pescott, O., 2023. Descriptive inference using large, unrepresentative nonprobability samples: An introduction for ecologists.

Brumm, K.J., Hanks, R.D., Baldwin, R.F., Peoples, B.K., 2021. Accounting for multiple dimensions of biodiversity to assess surrogate performance in a freshwater conservation prioritization. Ecological Indicators 122, 107320. https://doi.org/10.1016/j.ecolind.2020.107320

Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R.O., Wang, Z., Zimmermann, N.E., 2020. Model complexity affects species distribution projections under climate change. Journal of Biogeography 47, 130–142. https://doi.org/10.1111/jbi.13734

Bull, J.W., Gordon, A., Watson, J.E.M., Maron, M., 2016. Seeking convergence on the key concepts in 'no net loss' policy. Journal of Applied Ecology 53, 1686–1693. https://doi.org/10.1111/1365-2664.12726

Cadotte, M.W., Tucker, C.M., 2018. Difficult decisions: Strategies for conservation prioritization when taxonomic, phylogenetic and functional diversity are not spatially congruent. Biological Conservation 225, 128–133. https://doi.org/10.1016/j.biocon.2018.06.014

Canbek, G., 2022. Gaining insights in datasets in the shade of "garbage in, garbage out" rationale: Feature space distribution fitting. WIREs Data Mining and Knowledge Discovery 12, e1456. https://doi.org/10.1002/widm.1456

Carmona, C.P., Tamme, R., Pärtel, M., de Bello, F., Brosse, S., Capdevila, P., González-M., R., González-Suárez, M., Salguero-Gómez, R., Vásquez-Valderrama, M., Toussaint, A., 2021. Erosion of global functional diversity across the tree of life. Science Advances 7, eabf2675. https://doi.org/10.1126/sciadv.abf2675

CESBIO, 2022. Dynamic Habitat Indices of NDVI - Available data [WWW Document]. URL https://files.coinduf.eu/AFSfaJ2Kymt8iYWw/ (accessed 4.3.23).

CESBIO, 2021. Variables for Biodiversity – Theia [WWW Document]. URL https://www.theia-land.fr/en/ceslist/variables-for-biodiversity-sec/ (accessed 4.3.23).

Christensen, R., 2022. ordinal: Regression Models for Ordinal Data version 2022.11-16 from CRAN.

Church, R.L., Stoms, D.M., Davis, F.W., 1996. Reserve selection as a maximal covering location problem. Biological Conservation 76, 105–112. https://doi.org/10.1016/0006-3207(95)00102-6

Clare, J.D.J., Townsend, P.A., Anhalt-Depies, C., Locke, C., Stenglein, J.L., Frett, S., Martin, K.J., Singh, A.,
Van Deelen, T.R., Zuckerberg, B., 2019. Making inference with messy (citizen science) data: when are
data accurate enough and how can they be improved? Ecological Applications 29, e01849.
https://doi.org/10.1002/eap.1849
Costello, M.J., Wieczorek, J., 2014. Best practice for biodiversity data management and publication.
Biological Conservation 173, 68–73. https://doi.org/10.1016/j.biocon.2013.10.018
Domisch, S., Friedrichs, M., Hein, T., Borgwardt, F., Wetzig, A., Jähnig, S.C., Langhans, S.D., 2019. Spatially
explicit species distribution models: A missed opportunity in conservation planning? Diversity and
Distributions 25, 758–769. https://doi.org/10.1111/ddi.12891
Dubos, N., Préau, C., Lenormand, M., Papuga, G., Monsarrat, S., Denelle, P., Louarn, M.L., Heremans, S.,
May, R., Roche, P., Luque, S., 2022. Assessing the effect of sample bias correction in species
distribution models. Ecological Indicators 145, 109487. https://doi.org/10.1016/j.ecolind.2022.109487
Dupont, P., Demerges, D., Drouet, E., Luquet, G.Chr., 2013. Révision systématique, taxinomique et
nomenclaturale des Rhopalocera et des Zygaenidae de France métropolitaine. Conséquences sur
l'acquisition et la gestion des données d'inventaire (No. 19). MNHN-SPN.
Elvidge, C.D., Zhizhin, M., Ghosh, T., Hsu, F.-C., Taneja, J., 2021. Annual Time Series of Global VIIRS
Nighttime Lights Derived from Monthly Averages: 2012 to 2019. Remote Sensing 13, 922.
https://doi.org/10.3390/rs13050922
Essens, T., van Langevelde, F., Vos, R.A., Van Swaay, C.A.M., WallisDeVries, M.F., 2017. Ecological
determinants of butterfly vulnerability across the European continent. Journal of Insect Conservation
21, 439–450. https://doi.org/10.1007/s10841-017-9972-4
Fahrig, L., 1997. Relative effects of habitat loss and fragmentation on population extinction. The Journal
of wildlife management 61, 603–610. https://doi.org/10.2307/3802168
Ferrier, S., Pressey, R.L., Barrett, T.W., 2000. A new predictor of the irreplaceability of areas for achieving
a conservation goal, its application to real-world planning, and a research agenda for further
refinement. Biological Conservation 93, 303–325. https://doi.org/10.1016/S0006-3207(99)00149-4
Fithian, W., Elith, J., Hastie, T., Keith, D.A., 2015. Bias correction in species distribution models: pooling
survey and collection data for multiple species. Methods in Ecology and Evolution 6, 424–438.
https://doi.org/10.1111/2041-210X.12242
Fletcher, R.J., Hefley, T.J., Robertson, E.P., Zuckerberg, B., McCleery, R.A., Dorazio, R.M., 2019. A practical
guide for combining data to model species distributions. Ecology 100, e02710.
https://doi.org/10.1002/ecy.2710
Fukuda, S., De Baets, B., 2016. Data prevalence matters when assessing species' responses using data-
driven species distribution models. Ecological Informatics 32, 69–78.
https://doi.org/10.1016/j.ecoinf.2016.01.005
Fundingsland Tetlow, M., Hanusch, M., 2012. Strategic environmental assessment: the state of the art.
Impact Assessment and Project Appraisal 30, 15–24. https://doi.org/10.1080/14615517.2012.666400
Gargominy, O., Tercerie, S., Régnier, C., Ramage, T., Dupont, T., Daszkiewicz, P., Poncet, L., 2021. TAXREF
v14, référentiel taxonomique pour la France : méthodologie, mise en oeuvre et diffusion, Rapport
Patrinat. UMS PatriNat (OFB-CNRS-MNHN), Paris, France.
Gontier, M., Balfors, B., Mörtberg, U., 2006. Biodiversity in environmental assessment - current practice
and tools for prediction. Environmental Impact Assessment Review 26, 268–286.
https://doi.org/10.1016/j.eiar.2005.09.001
Guillera-Arroita, G., Lahoz-Monfort, J.J., Elith, J., Gordon, A., Kujala, H., Lentini, P.E., McCarthy, M.A.,
Tingley, R., Wintle, B.A., 2015. Is my species distribution model fit for purpose? Matching data and
models to applications. Global Ecology and Biogeography 24, 276–292.
https://doi.org/10.1111/geb.12268
Guillet, F., Le Floch, C., Julliard, R., 2019. Séquence Eviter-Réduire-Compenser : quelle biodiversité est
visée par les mesures d'évitement ? Sciences Eaux et Territoires 1–8. https://doi.org/10.14758/SET-
REVUE.2019.HS.07
Guisan, A., Thuiller, W., Zimmermann, N.E., 2017. Habitat Suitability and Distribution Models: with
Applications in R. Cambridge University Press.
Guisan, A., Tingley, R., Baumgartner, J.B., Naujokaitis-Lewis, I., Sutcliffe, P.R., Tulloch, A.I.T., Regan, T.J.,
Brotons, L., McDonald-Madden, E., Mantyka-Pringle, C., Martin, T.G., Rhodes, J.R., Maggini, R.,

Setterfield, S.A., Elith, J., Schwartz, M.W., Wintle, B.A., Broennimann, O., Austin, M., Ferrier, S.,
Kearney, M.R., Possingham, H.P., Buckley, Y.M., 2013. Predicting species distributions for conservation
decisions. Ecology Letters 16, 1424–1435. https://doi.org/10.1111/ele.12189
Hanson, J.O., Schuster, R., Morrell, N., Strimas-Mackey, M., Edwards, B.P.M., Watts, M.E., Arcese, P.,
Bennett, J., Possingham, H.P., 2021. prioritizr: Systematic Conservation Prioritization in R.
Hermoso, V., Kennard, M.J., Linke, S., 2015a. Assessing the risks and opportunities of presence-only data
for conservation planning. Journal of Biogeography 42, 218–228. https://doi.org/10.1111/jbi.12393
Hermoso, V., Kennard, M.J., Linke, S., 2015b. Evaluating the costs and benefits of systematic data
acquisition   for   conservation   assessments.   Ecography   38,   283–292.
https://doi.org/10.1111/ecog.00792
Horváth, Z., Ptacnik, R., Vad, C.F., Chase, J.M., 2019. Habitat loss over six decades accelerates regional
and local biodiversity loss via changing landscape connectance. Ecology Letters 22, 1019–1027.
https://doi.org/10.1111/ele.13260
IGN, 2022. Géoservices - Téléchargement de données [WWW Document]. URL
https://geoservices.ign.fr/telechargement (accessed 4.3.23).
Inglada, J., Vincent, A., Thierion, V., 2019. Theia OSO Land Cover Map 2019 [Data set]. Zenodo.
https://doi.org/10.5281/zenodo.6538321
Iorio, É., Dusoulier, F., Soldati, F., Noël, F., Guilloton, J.-A., Doucet, G., Ponel, P., Dupont, P., Krieg-
Jacquier, R., Chemin, S., Tillier, P., Touroult, J., 2022. Les Arthropodes terrestres dans les études
d'impact : limites actuelles et propositions pour une meilleure prise en compte des enjeux de
conservation. Naturae 43–99. https://doi.org/10.5852/naturae2022a4
IPBES, 2019. Global assessment report on biodiversity and ecosystem services of the Intergovernmental
Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES secretariat.
https://doi.org/10.5281/zenodo.6417333
Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K., Mooers, A.O., 2012. The global diversity of birds in space
and time. Nature 491, 444–448. https://doi.org/10.1038/nature11631
Jomier, R., Solène, R., Dupont, P., 2018. National Scientific Validation for Data in the Information System
on Nature and Landscapes (Système d'Information sur la Nature et les Paysages - SINP). Biodiversity
Information Science and Standards e25881. https://doi.org/10.3897/biss.2.25881
Julliard, R., Jiguet, F., 2002. Un suivi intégré des populations d'oiseaux communs en France, in: Alauda.
Presented at the Colloque Francophone Ornithologique, Limoges, France, pp. 137–147.
Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann, N.E., Linder,
H.P., Kessler, M., 2021. Climatologies at high resolution for the earth's land surface areas. EnviDat
170122. https://doi.org/ 10.16904/envidat.228.v2.1
Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann, N.E., Linder,
H.P., Kessler, M., 2017. Climatologies at high resolution for the earth's land surface areas. Scientific
Data 4, 170122. https://doi.org/10.1038/sdata.2017.122
Kim, V., Ralphs, T., Guzelsoy, M., Mahajan, A., Reinhard, H., Hornik, K., Szymanski, C., Theussl, S., 2023.
lpsymphony: Symphony integer linear programming solver in R.
Kujala, H., Moilanen, A., Gordon, A., 2018. Spatial characteristics of species distributions as drivers in
conservation   prioritization.   Methods   in   Ecology   and   Evolution   9,   1121–1132.
https://doi.org/10.1111/2041-210X.12939
Lagabrielle, E., Botta, A., Daré, W., David, D., Aubert, S., Fabricius, C., 2010. Modelling with stakeholders
to integrate biodiversity into land-use planning - Lessons learned in Réunion Island (Western Indian
Ocean).   Environmental   Modelling   &   Software   25,   1413–1427.
https://doi.org/10.1016/j.envsoft.2010.01.011
Lees, C.M., Rutschmann, A., Santure, A.W., Beggs, J.R., 2021. Science-based, stakeholder-inclusive and
participatory conservation planning helps reverse the decline of threatened species. Biological
Conservation 260, 109194. https://doi.org/10.1016/j.biocon.2021.109194
Leroy, B., Delsol, R., Hugueny, B., Meynard, C.N., Barhoumi, C., Barbet-Massin, M., Bellard, C., 2018.
Without quality presence–absence data, discrimination metrics such as TSS can be misleading
measures   of   model   performance.   Journal   of   Biogeography   45,   1994–2002.
https://doi.org/10.1111/jbi.13402

Lino, A., Fonseca, C., Rojas, D., Fischer, E., Pereira, M.J.R., 2019. A meta-analysis of the effects of habitat
loss and fragmentation on genetic diversity in mammals. Mammalian Biology 94, 69–76.
https://doi.org/10.1016/j.mambio.2018.09.006
Mac Nally, R., Duncan, R.P., Thomson, J.R., Yen, J.D.L., 2018. Model selection using information criteria,
but is the "best" model any good? Journal of Applied Ecology 55, 1441–1444.
https://doi.org/10.1111/1365-2664.13060
Margules, C.R., Pressey, R.L., 2000. Systematic conservation planning. Nature 405, 243–253.
https://doi.org/10.1038/35012251
Mariton, L., Kerbiriou, C., Bas, Y., Zanda, B., Le Viol, I., 2022. Even low light pollution levels affect the
spatial distribution and timing of activity of a "light tolerant" bat species. Environmental Pollution
305, 119267. https://doi.org/10.1016/j.envpol.2022.119267
Maron, M., Ives, C.D., Kujala, H., Bull, J.W., Maseyk, F.J.F., Bekessy, S., Gordon, A., Watson, J.E.M., Lentini,
P.E., Gibbons, P., Possingham, H.P., Hobbs, R.J., Keith, D.A., Wintle, B.A., Evans, M.C., 2016. Taming a
Wicked Problem: Resolving Controversies in Biodiversity Offsetting. BioScience 66, 489–498.
https://doi.org/10.1093/biosci/biw038
Matsuki, K., Kuperman, V., Van Dyke, J.A., 2016. The Random Forests statistical technique: An
examination of its value for the study of reading. Scientific Studies of Reading 20, 20–33.
https://doi.org/10.1080/10888438.2015.1107073
Matutini, F., Baudry, J., Pain, G., Sineau, M., Pithon, J., 2021. How citizen science could improve species
distribution models and their independent assessment. Ecology and Evolution 11, 3028–3039.
https://doi.org/10.1101/2020.06.02.129536
Maxwell, S.L., Fuller, R.A., Brooks, T.M., Watson, J.E.M., 2016. Biodiversity: The ravages of guns, nets and
bulldozers. Nature News 536, 143–145. https://doi.org/10.1038/536143a
MEDDE, GIS Sol, 2014. . Enveloppes des milieux potentiellement humides de la France métropolitaine.
Notice d'accompagnement. Programme de modélisation des milieux potentiellement humides de
France. Ministère d'Ecologie, du Développement Durable et de l'Energie, Groupement d'Intérêt
Scientifique Sol.
Meyer, C., Kreft, H., Guralnick, R., Jetz, W., 2015. Global priorities for an effective information basis of
biodiversity distributions. Nature Communications 6, 1–8. https://doi.org/10.1038/ncomms9221
Meyer, C.B., 2007. Does Scale Matter in Predicting Species Distributions? Case Study with the Marbled
Murrelet. Ecological Applications 17, 1474–1483. https://doi.org/10.1890/06-1410.1
Middleton-Welling, J., Dapporto, L., García-Barros, E., Wiemers, M., Nowicki, P., Plazio, E., Bonelli, S.,
Zaccagno, M., Šašić, M., Liparova, J., Schweiger, O., Harpke, A., Musche, M., Settele, J., Schmucki, R.,
Shreeve, T., 2020. A new comprehensive trait database of European and Maghreb butterflies,
Papilionoidea. Scientific Data 7, 351. https://doi.org/10.1038/s41597-020-00697-7
Morelli, F., Benedetti, Y., Callaghan, C.T., 2020. Ecological specialization and population trends in
European breeding birds. Global Ecology and Conservation 22, e00996.
https://doi.org/10.1016/j.gecco.2020.e00996
Moussus, J.-P., Lorin, T., Cooper, A., 2019. Guide pratique des papillons de France, Delachaux et Niestlé.
ed. France.
Olden, J.D., Rooney, T.P., 2006. On defining and quantifying biotic homogenization. Global Ecology and
Biogeography 15, 113–120. https://doi.org/10.1111/j.1466-822X.2006.00214.x
Phalan, B., Hayes, G., Brooks, S., Marsh, D., Howard, P., Costelloe, B., Vira, B., Kowalska, A., Whitaker, S.,
2018. Avoiding impacts on biodiversity through strengthening the first stage of the mitigation
hierarchy. Oryx 52, 316–324. https://doi.org/10.1017/S0030605316001034
Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J., Ferrier, S., 2009. Sample
selection bias and presence-only distribution models: implications for background and pseudo-
absence data. Ecological Applications 19, 181–197. https://doi.org/10.1890/07-2153.1
Ponder, W.F., Carter, G.A., Flemons, P., Chapman, R.R., 2001. Evaluation of Museum Collection Data for
Use in Biodiversity Assessment. Conservation Biology 15, 648–657. https://doi.org/10.1046/j.1523-
1739.2001.015003648.x
Pope, J., Bond, A., Morrison-Saunders, A., Retief, F., 2013. Advancing the theory and practice of impact
assessment: Setting the research agenda. Environmental Impact Assessment Review 41, 1–9.
https://doi.org/10.1016/j.eiar.2013.01.008

Pressey, R.L., Bottrill, M.C., 2008. Opportunism, Threats, and the Evolution of Systematic Conservation
Planning. Conservation Biology 22, 1340–1345. https://doi.org/10.1111/j.1523-1739.2008.01032.x
Quétier, F., Regnery, B., Levrel, H., 2014. No net loss of biodiversity or paper offsets? A critical review of
the French no net loss policy. Environmental Science & Policy 38, 120–131.
https://doi.org/10.1016/j.envsci.2013.11.009
Rondinini, C., Wilson, K.A., Boitani, L., Grantham, H., Possingham, H.P., 2006. Tradeoffs of different types
of species occurrence data for use in systematic conservation planning. Ecology Letters 9, 1136–1145.
https://doi.org/10.1111/j.1461-0248.2006.00970.x
Sanders, H., Saxe, J., 2017. Garbage in, garbage out: how purportedly great ML models can be screwed up
by bad data. Presented at the Blackhat, Las Vegas, USA, p. 6.
Schoener, T.W., 1968. The Anolis Lizards of Bimini: Resource Partitioning in a Complex Fauna. Ecology 49,
704–726. https://doi.org/10.2307/1935534
Sheard, C., Neate-Clegg, M.H.C., Alioravainen, N., Jones, S.E.I., Vincent, C., MacGregor, H.E.A., Bregman,
T.P., Claramunt, S., Tobias, J.A., 2020. Ecological drivers of global gradients in avian dispersal inferred
from wing morphology. Nature Communications 11, 2463. https://doi.org/10.1038/s41467-020-
16313-6
Shirey, V., Belitz, M.W., Barve, V., Guralnick, R., 2021. A complete inventory of North American butterfly
occurrence data: narrowing data gaps, but increasing bias. Ecography 44, 537–547.
https://doi.org/10.1111/ecog.05396
Sofaer, H.R., Jarnevich, C.S., Pearse, I.S., Smyth, R.L., Auer, S., Cook, G.L., Edwards, T.C., Jr, Guala, G.F.,
Howard, T.G., Morisette, J.T., Hamilton, H., 2019. Development and Delivery of Species Distribution
Models to Inform Decision-Making. BioScience 69, 544–557. https://doi.org/10.1093/biosci/biz045
Sousa-Baena, M.S., Garcia, L.C., Peterson, A.T., 2014. Completeness of digital accessible knowledge of the
plants of Brazil and priorities for survey and inventory. Diversity and Distributions 20, 369–381.
https://doi.org/10.1111/ddi.12136
Srisa-An, C., 2021. Guideline of Collinearity - Avoidable Regression Models on Time-series Analysis, in:
2021 2nd International Conference on Big Data Analytics and Practices (IBDAP). Presented at the 2021
2nd International Conference on Big Data Analytics and Practices (IBDAP), Bangkok, Thailand, pp. 28–
32. https://doi.org/10.1109/IBDAP52511.2021.9552165
Stekhoven, D.J., Bühlmann, P., 2012. MissForest--non-parametric missing value imputation for mixed-
type data. Bioinformatics 28, 112–118. https://doi.org/10.1093/bioinformatics/btr597
Støa, B., Halvorsen, R., Stokland, J.N., Gusarov, V.I., 2019. How much is enough? Influence of number of
presence observations on the performance of species distribution models. Sommerfeltia 39, 1–28.
https://doi.org/10.2478/som-2019-0001
Storchová, L., Hořák, D., 2018. Life-history characteristics of European birds. Global Ecology and
Biogeography 27, 400–406. https://doi.org/10.1111/geb.12709
Tarabon, S., Bergès, L., Dutoit, T., Isselin-Nondedeu, F., 2019. Environmental impact assessment of
development projects improved by merging species distribution and habitat connectivity modelling.
Journal of Environmental Management 241, 439–449. https://doi.org/10.1016/j.jenvman.2019.02.031
Telenius, A., 2011. Biodiversity information goes public: GBIF at your service. Nordic Journal of Botany 29,
378–381. https://doi.org/10.1111/j.1756-1051.2011.01167.x
Tittensor, D.P., Walpole, M., Hill, S.L.L., Boyce, D.G., Britten, G.L., Burgess, N.D., Butchart, S.H.M., Leadley,
P.W., Regan, E.C., Alkemade, R., Baumung, R., Bellard, C., Bouwman, L., Bowles-Newark, N.J., Chenery,
A.M., Cheung, W.W.L., Christensen, V., Cooper, H.D., Crowther, A.R., Dixon, M.J.R., Galli, A., Gaveau,
V., Gregory, R.D., Gutierrez, N.L., Hirsch, T.L., Höft, R., Januchowski-Hartley, S.R., Karmann, M., Krug,
C.B., Leverington, F.J., Loh, J., Lojenga, R.K., Malsch, K., Marques, A., Morgan, D.H.W., Mumby, P.J.,
Newbold, T., Noonan-Mooney, K., Pagad, S.N., Parks, B.C., Pereira, H.M., Robertson, T., Rondinini, C.,
Santini, L., Scharlemann, J.P.W., Schindler, S., Sumaila, U.R., Teh, L.S.L., van Kolck, J., Visconti, P., Ye,
Y., 2014. A mid-term analysis of progress toward international biodiversity targets. Science 346, 241–
244. https://doi.org/10.1126/science.1257484
Toussaint, A., Brosse, S., Bueno, C.G., Pärtel, M., Tamme, R., Carmona, C.P., 2021. Extinction of
threatened vertebrates will lead to idiosyncratic changes in functional diversity across the world.
Nature Communications 12, 5162. https://doi.org/10.1038/s41467-021-25293-0

Troudet, J., Grandcolas, P., Blin, A., Vignes-Lebbe, R., Legendre, F., 2017. Taxonomic bias in biodiversity
data and societal preferences. Scientific Reports 7, 9132. https://doi.org/10.1038/s41598-017-09084-
6

Tulloch, A.I.T., Gordon, A., Runge, C.A., Rhodes, J.R., 2019. Integrating spatially realistic infrastructure
impacts into conservation planning to inform strategic environmental assessment. Conservation
Letters 12, e12648. https://doi.org/10.1111/conl.12648

Tulloch, A.I.T., Sutcliffe, P., Naujokaitis-Lewis, I., Tingley, R., Brotons, L., Ferraz, K.M.P.M.B., Possingham,
H., Guisan, A., Rhodes, J.R., 2016. Conservation planners tend to ignore improved accuracy of
modelled species distributions to focus on multiple threats and ecological processes. Biological
Conservation 199, 157–171. https://doi.org/10.1016/j.biocon.2016.04.023

Valavi, R., Elith, J., Lahoz-Monfort, J.J., Guillera-Arroita, G., 2021a. Modelling species presence-only data
with random forests. Ecography 44, 1731–1742. https://doi.org/10.1111/ecog.05615

Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J.J., Elith, J., 2021b. Predictive performance of presence-
only species distribution models: a benchmark study with reproducible code. Ecological Monographs.
https://doi.org/10.1002/ecm.1486

Velazco, S.J.E., Ribeiro, B.R., Laureto, L.M.O., De Marco Júnior, P., 2020. Overprediction of species
distribution models in conservation planning: A still neglected issue with strong effects. Biological
Conservation 252, 108822. https://doi.org/10.1016/j.biocon.2020.108822

Warren, D., Dinnage, R., 2022. ENMTools: Analysis of Niche Evolution using Niche and Distribution
Models.

Warren, D.L., Glor, R.E., Turelli, M., 2008. Environmental Niche Equivalency Versus Conservatism:
Quantitative Approaches to Niche Evolution. Evolution 62, 2868–2883.
https://doi.org/10.1111/j.1558-5646.2008.00482.x

Whitehead, A.L., Kujala, H., Wintle, B.A., 2017. Dealing with Cumulative Biodiversity Impacts in Strategic
Environmental Assessment: A New Frontier for Conservation Planning. Conservation Letters 10, 195–
204. https://doi.org/10.1111/conl.12260

Wiemers, M., Chazot, N., Wheat, C.W., Schweiger, O., Wahlberg, N., 2020. A complete time-calibrated
multi-gene phylogeny of the European butterflies. ZooKeys 938, 97–124.
https://doi.org/10.3897/zookeys.938.50878

Wiemers, M., Chazot, N., Wheat, C.W., Schweiger, O., Wahlberg, N., 2019. A complete time-calibrated
multi-gene phylogeny of the European butterflies - data [Data set]. Zenodo.
https://doi.org/10.5281/zenodo.3531555

Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M.M., Jetz, W., 2014. EltonTraits 1.0:
Species-level foraging attributes of the world's birds and mammals. Ecology 95, 2027–2027.
https://doi.org/10.1890/13-1917.1

Witté, I., Touroult, J., 2017. Identification et cartographie des zones de méconnaissance naturaliste à
l'échelle nationale (métropole) à partir des données partagées (No. 2017–6). Muséum National
d'Histoire Naturel (MNHN) - Service du Patrimoine Naturel (SPN), France.

Zambrano, J., Garzon-Lopez, C.X., Yeager, L., Fortunel, C., Cordeiro, N.J., Beckman, N.G., 2019. The effects
of habitat loss and fragmentation on plant functional traits and functional diversity: what do we know
so far? Oecologia 191, 505–518. https://doi.org/10.1007/s00442-019-04505-x

Zuckerberg, B., Huettmann, F., Frair, J., 2011. Proper Data Management as a Scientific Foundation for
Reliable Species Distribution Modeling, in: Drew, C.A., Wiersma, Y.F., Huettmann, F. (Eds.), Predictive
Species and Habitat Modeling in Landscape Ecology: Concepts and Applications. Springer, New York,
NY, pp. 45–70. https://doi.org/10.1007/978-1-4419-7390-0_4

Zurell, D., Franklin, J., König, C., Bouchet, P., Dormann, C., Elith, J., Fandos, G., Feng, X., Guillera-Arroita,
G., Guisan, A., Lahoz-Monfort, J.J., Leitão, P.J., Park, D., Peterson, A.T., Rapacciuolo, G., Schmatz, D.,
Schröder, B., Serra-Diaz, J.M., Thuiller, W., Yates, K., Zimmermann, N., Merow, C., 2020. A standard
protocol for reporting species distribution models. https://doi.org/10.18452/22268

Appendix

Appendix A - Additional material and methods

A.1 - Collect and series of operation on species occurrences

Databases collection:
- Access to source databases:
  o French Natural and Landscape Information System (SINP, https://inpn.mnhn.fr), each of the three study sites has its own database: https://sinp-occitanie.fr/atlas/ for Lodévois-Larzac (T1), https://obv-na.fr/ and https://observatoire-fauna.fr/ for La Rochelle (T2), https://data.biodiversite-bretagne.fr for Brocéliande (T3).
  o GBIF: GBIF.org (04 October 2022) GBIF Occurrence Download https://doi.org/10.15468/dl.ry6uw7
  o French biodiversity monitoring scheme (Vigie Nature): https://www.vigienature.fr/, the database used are the STOC for Aves, the Vigie-Chiro for Chiroptera and Orthoptera, the Vigie-Flore for Flora, the STELI for Odonata, the STERF for Papilionidae
- Period of 11 years, from 01/01/2010 to 31/12/2020
- Extent: continental France (without island, Corsica and overseas regions)
- Spatial accuracy of 0 to 50m

Standardization of species names : French taxonomic reference TAXREF.V14 (Gargominy et al., 2021).
- Keep only the species and subspecies
- Groups data at the species level

Taxonomic filtering: from TAXREF.V14
- Keep only species in nine taxonomic groups selected: Amphibia, Aves, Chiroptera, Flora, Mammalia aptera (Mammalia), Orthoptera, Odonata, Papilionidae and Reptilia
- Delete species not present in France so keep: "Present", "Endemic", "Subendemic", "Cryptogenic", "Introduced", "Invasive introduced"

Data transformation and filtering:
- Transformation data into occurrence, i.e. transform data into presence only
- Geographical filtering for reduce sampling biases and delete duplicate data: spatial thinning at a distance of 50m with "spThin" package (Aiello-Lammens et al., 2015)

Identify the presence of species into each study site: at least five observations from all databases in the study site to certify the presence of the species

Standardization of the observation period between databases: for Aves filtering observations between 1 March and 15 July corresponding to the STOC period.

906 *A.2 - Environmental variables selected for Species Distribution Models (SDM) of Aves and Papilionidae species. Variables used for SDM are the variables with*
907 *collinearity removed (Pearson >0.7).*

Type of variables		Variables	Database	Year	Raw precision	Source	Variables used for SDM
Geography	Climatic	19 bioclimatic variables	Chelsa	1981-2010	1 km	Karger et al., 2021, 2017	Bio 2 (temp. range), Bio 5 (temp. warmest month), Bio 6 (min. temp. coldest month), Bio 9 (temp. driest quarter), Bio 13 (prec. wettest month), Bio 14 (prec. driest month), Bio 15(prec. seasonality)
	Elevation	Elevation, mean in a buffer of fifty meters	BD ALTI, IGN	2014	25 m	IGN, 2022	Elevation
		Slope, mean in a buffer of 50 meters					Slope
		Topographic Position Index (TPI), mean in a buffer of 50 meters					TPI
	Wetland	Area of wetland potential: very high in a buffer of 50, 500, 2000 meters	INRAE Wetland potential	2014	50 m	Berthier et al., 2014	Wetland in buffer of 50 and 500 m
	Waterway	Linear meters of waterway in a buffer of 50, 500, 2000 meters	BD TOPO, IGN	2022	Vector	IGN, 2022	Waterway in buffer of 50 and 500m
Human occupancy	Land-use	Area of artificial, conifer forest, crop, deciduous forest, land, lawn, mineral, prairie, water in buffer of 50, 500 and 2000 meters	OSO Land Cover	2018	25 m	Inglada et al., 2019	Artificial, conifer, crops, deciduous, fruit culture, land, mineral, prairie, water in a buffer of 50 and 500 m, Lawn in a buffer of 500 m, Sand in a buffer of 50, 500, 2000 m
	Hedge line	Linear meters of hedge in a buffer of 50, 500, 2000 meters	BD TOPO, IGN	2022	Vector	IGN, 2022	Hedge line in buffer of 50, 500 m
	NDVI mean	NDVI mean in buffer of 50, 500, 2000 meters	DHI NDVI	2022	10 m	CESBIO, 2021, 2022	DHI NDVI in buffer of 50 and 500m
Pollution / Fragmentation	Light pollution	Mean value in buffer of 500 meters of night time light average masked of 2015	VIIRS night time lights	2015	500 m	Elvidge et al., 2021	Night time light
	Road fragmentation	Linear meters of fragmenting of main road in a buffer of 50, 500, 2000 meters	BD TOPO, IGN	2022	Vector	IGN, 2022	Main road in a buffer of 50, 500, 2000 m

909 *A.3 - Species traits of Aves and Papilionidae. Missing traits were imputed by considering evolutionary*
910 *relationships following the process of Carmona et al. (2021) with the help the R scripts of Toussaint et*
911 *al. (2021).*

	Traits	Description	Reference
Aves	Mass	Body mass using data from males, females, and/or unspecified adults in g	Storchová and Hořák, 2018
	HWI	Avian hand-wing index (HWI), an estimate of wing shape as a proxy for dispersal ability in birds	Sheard et al., 2020
	Nocturn	Nocturnal activity	Wilman et al., 2014
	Habitats (Deciduous Coniferous, Woodland, Shrub, Grassland, Mountain meadows, Reed, Swamps, Rocks, Urban)	Species occupies habitat in breeding area, 1: yes; 0: no	Storchová and Hořák, 2018
	Spe. Diet, Spe. Foraging behav., Spe. Diet strat, Spe. Hab., Spe. Nest, Spe. Mean	Specialization respectively of diet, foraging behaviour, foraging substrate, habitat, nesting site and mean of five specializations	Morelli et al., 2020
Papilionidae	Wingspan	Average length of male wingspan in mm	Middleton-Welling et al., 2020
	FMoMean	Average duration of yearly flight period	
	HostplantN	Hostplant specificity, number of hostplants	
	HostplantSpe	Hostplant specificity index	
	HPG	Hostplant growth form: Short herb/grass (<1 m) (Bi), tall herb/grass (>1 m) (Th), shrub (Sb), tree (Tr), liana (Li)	
	SSI	Species Specialization Index	Essens et al., 2017
	AltVeg	Altitudinal vegetation: A=Anthropogenic, Co=Foothill, TheMeMed=Thermo/Meso-Mediterranean, Med=Mediterranean, SupMed=Supra-Mediterranean, Mo=Montane, ASa=Alpine and Subalpine	Dupont et al., 2013
	Rarity	Rarity in France between 1 very rare to 3 common species	In Moussus et al., 2019 from oreina.org and faune.france.org

The phylogenetic trees used to consider evolutionary relationships were from Jetz et al. (2012) and Wiemers et al. (2019, 2020) for the Aves and Papilionidae, respectively.

 Appendix B - Supplementary results

B.1 - Description of observation databases (Local SINP, GBIF, Vigie Nature) available nine taxonomic
groups in the three study sites (T1: Lodévois-Larzac, T2: Brocéliande, T3: La Rochelle) for "Local"
(study site with a buffer zone 10 km), "Regional" (administrative region) and "Country" (continental
France) scales. Statistics: "nSp" is the number of species with at least five observations on the study
site within a 10 km buffer; "pSp<15obs", "pSp15.50obs" and "pSp>50obs" are the percentage of
species with <15, 15 to 50 and >50 observations respectively and "-" represents no data available.

B.1.A - Local scale (10km buffer zone).

Taxa	Statistc	GBIF			SINP			Vigie Nature			All databases combined		
		T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3
Amphibia	nSp	11	1	11	13	2	0				14	2	11
	pSp<15obs	82%	100%	100%	23%	50%	-				29%	50%	100%
	pSp15.50obs	18%	0%	0%	15%	50%	-		No program		7%	50%	0%
	pSp.>50obs	0%	0%	0%	62%	0%	-				64%	0%	0%
Aves	nSp	100	73	77	197	94	82	114	155	101	205	171	132
	pSp<15obs	87%	93%	95%	37%	96%	85%	56%	50%	54%	37%	47%	59%
	pSp15.50obs	10%	6%	4%	23%	3%	15%	32%	17%	20%	22%	22%	13%
	pSp.>50obs	3%	1%	1%	40%	1%	0%	12%	33%	27%	41%	31%	28%
Chiroptera	nSp	4	0	5	21	6	18	25	22	24	26	22	26
	pSp<15obs	100%	-	100%	10%	83%	50%	12%	41%	21%	12%	41%	27%
	pSp15.50obs	0%	-	0%	38%	0%	44%	16%	36%	46%	19%	36%	34%
	pSp.>50obs	0%	-	0%	52%	17%	6%	72%	23%	33%	69%	23%	39%
Flora	nSp	786	476	639	1703	1335	843	286	177	136	1844	1460	1093
	pSp<15obs	81%	85%	81%	52%	48%	74%	100%	100%	100%	52%	50%	72%
	pSp15.50obs	16%	14%	17%	30%	25%	14%	0%	0%	0%	28%	24%	17%
	pSp.>50obs	3%	1%	3%	18%	27%	12%	0%	0%	0%	20%	25%	11%
Mammalia	nSp	10	1	14	23	7	30				23	7	30
	pSp<15obs	100%	100%	100%	78%	86%	60%				74%	86%	57%
	pSp15.50obs	0%	0%	0%	17%	14%	20%		No program		22%	14%	20%
	pSp.>50obs	0%	0%	0%	4%	0%	20%				4%	0%	23%
Odonata	nSp	38	5	23	57	1	0	0	1	24	57	7	30
	pSp<15obs	74%	100%	100%	28%	100%	-	-	100%	100%	28%	100%	97%
	pSp15.50obs	26%	0%	0%	37%	0%	-	-	0%	0%	30%	0%	3%
	pSp.>50obs	0%	0%	0%	35%	0%	-	-	0%	0%	42%	0%	0%
Orthoptera	nSp	53	3	24	91	8	0	25	19	19	98	25	34
	pSp<15obs	92%	100%	100%	71%	100%	-	24%	58%	16%	56%	68%	53%
	pSp15.50obs	8%	0%	0%	23%	0%	-	24%	32%	52%	18%	24%	29%
	pSp.>50obs	0%	0%	0%	6%	0%	-	52%	10%	32%	26%	8%	18%
Papilionidae	nSp	115	15	33	154	11	0	0	0	50	154	18	52
	pSp<15obs	60%	100%	94%	21%	100%	-	-	-	88	21%	100%	73%
	pSp15.50obs	30%	0%	6%	19%	0%	-	-	-	12	18%	0%	27%
	pSp.>50obs	10%	0%	0%	60%	0%	-	-	-	0	60%	0%	0%
Reptilia	nSp	18	2	7	21	4	0				22	4	7
	pSp<15obs	67%	100%	100%	24%	100%	-				27%	75%	100%
	pSp15.50obs	28%	0%	0%	33%	0%	-		No program		32%	25%	0%
	pSp.>50obs	6%	0%	0%	43%	0%	-				41%	0%	0%

922 *B.1.A - Regional scale.*

Taxa	Statistic	GBIF			Vigie Nature			All databases combined		
		T1	T2	T3	T1	T2	T3	T1	T2	T3
Amphibia	nSp	13	2	11				14	2	11
	pSp<15obs	8%	0%	73%				7%	0%	73%
	pSp15.50obs	23%	50%	18%		No program		14%	50%	18%
	pSp.>50obs	69%	50%	9%				79%	50%	9%
Aves	nSp	198	159	120	182	143	120	205	166	132
	pSp<15obs	21%	21%	40%	31%	29%	32%	12%	16%	19%
	pSp15.50obs	12%	12%	22%	21%	15%	27%	11%	10%	28%
	pSp.>50obs	67%	67%	38%	48%	56%	41%	77%	74%	53%
Chiroptera	nSp	13	13	5	25	22	24	26	22	26
	pSp<15obs	100%	100%	100%	0%	0%	8%	0%	0%	15%
	pSp15.50obs	0%	0%	0%	4%	0%	17%	8%	0%	15%
	pSp.>50obs	0%	0%	0%	96%	100%	75%	92%	100%	69%
Flora	nSp	1428	1093	825	825	657	413	1844	1158	1093
	pSp<15obs	43%	38%	47%	95%	94%	94%	32%	40%	47%
	pSp15.50obs	19%	19%	20%	5%	6%	6%	26%	19%	23%
	pSp.>50obs	38%	49%	33%	0%	0%	0%	42%	41%	31%
Mammalia	nSp	23	6	21				23	6	30
	pSp<15obs	39%	33%	62%				30%	33%	50%
	pSp15.50obs	26%	0%	33%		No program		22%	0%	23%
	pSp.>50obs	35%	67%	5%				48%	67%	27%
Odonata	nSp	56	7	28	56	7	27	57	7	30
	pSp<15obs	18%	14%	64%	77%	71%	100%	9%	14%	53%
	pSp15.50obs	30%	14%	25%	23%	29%	0%	21%	14%	37%
	pSp.>50obs	52%	71%	11%	0%	0%	0%	70%	72%	10%
Orthoptera	nSp	93	25	26	25	19	19	98	25	34
	pSp<15obs	42%	24%	58%	4%	0%	11%	30%	4%	27%
	pSp15.50obs	39%	40%	42%	4%	0%	16%	22%	0%	29%
	pSp.>50obs	19%	36%	0%	92%	100%	74%	48%	96%	44%
Papilionidae	nSp	144	18	43	132	18	51	154	18	52
	pSp<15obs	12%	0%	37%	42%	11%	55%	7%	0%	34%
	pSp15.50obs	19%	11%	42%	30%	22%	41%	13%	6%	31%
	pSp.>50obs	69%	89%	21%	28%	67%	4%	81%	94%	35%
Reptilia	nSp	20	4	7				22	4	7
	pSp<15obs	20%	0%	14%				18%	0%	14%
	pSp15.50obs	30%	25%	57%		No program		28%	25%	57%
	pSp.>50obs	50%	75%	29%				59%	75%	29%

923   *B.1.C - Country scale.*

Taxa	Statistic	GBIF			Country Vigie Nature			All databases combined		
		T1	T2	T3	T1	T2	T3	T1	T2	T3
Amphibia	nSp	14	2	11	No program			14	2	11
	pSp<15obs	7%	0%	0%				0%	0%	0%
	pSp15.50obs	0%	0%	9%				7%	0%	9%
	pSp.>50obs	93%	100%	91%				93%	100%	91%
Aves	nSp	204	170	131	201	170	132	205	171	132
	pSp<15obs	2%	1%	1%	8%	9%	4%	1%	0%	0%
	pSp15.50obs	3%	1%	1%	11%	9%	7%	1%	1%	0%
	pSp.>50obs	95%	98%	98%	81%	82%	89%	98%	99%	100%
Chiroptera	nSp	19	19	21	25	22	25	26	22	26
	pSp<15obs	90%	89%	90%	0%	0%	0%	0%	0%	4%
	pSp15.50obs	10%	11%	10%	0%	0%	0%	4%	0%	0%
	pSp.>50obs	0%	0%	0%	100%	100%	100%	96%	100%	96%
Flora	nSp	1573	1261	995	1336	1105	848	1844	1460	1093
	pSp<15obs	25%	20%	17%	68%	61%	53%	19%	14%	16%
	pSp15.50obs	18%	18%	16%	19%	23%	27%	21%	17%	16%
	pSp.>50obs	57%	62%	67%	13%	16%	20%	60%	69%	68%
Mammalia	nSp	23	6	28	No program			23	7	30
	pSp<15obs	13%	16%	18%				9%	14%	23%
	pSp15.50obs	17%	17%	28%				13%	29%	17%
	pSp.>50obs	70%	67%	54%				78%	57%	60%
Odonata	nSp	57	7	30	57	7	30	57	7	30
	pSp<15obs	2%	0%	0%	11%	0%	0%	0%	0%	0%
	pSp15.50obs	10%	14%	0%	26%	0%	13%	3%	0%	0%
	pSp.>50obs	88%	86%	100%	63%	100%	87%	97%	100%	100%
Orthoptera	nSp	95	25	34	25	19	19	98	25	34
	pSp<15obs	22%	12%	6%	4%	0%	0%	15%	0%	0%
	pSp15.50obs	28%	20%	21%	0%	0%	0%	25%	4%	9%
	pSp.>50obs	50%	68%	73%	96%	100%	100%	60%	96%	91%
Papilionidae	nSp	147	18	52	149	18	52	154	18	52
	pSp<15obs	4%	0%	0%	13%	5%	0%	1%	0%	0%
	pSp15.50obs	8%	0%	0%	18%	6%	0%	7%	0%	0%
	pSp.>50obs	88%	100%	100%	69%	89%	100%	92%	100%	100%
Reptilia	nSp	21	4	7	No program			22	4	7
	pSp<15obs	0%	0%	0%				4%	0%	0%
	pSp15.50obs	14%	0%	0%				5%	0%	0%
	pSp.>50obs	86%	100%	100%				91%	100%	100%

B.2 - *SDM evaluation of Aves and Papilionidae taxonomic groups for (A) three individual database sources*
and (B) between SDM using different pseudo-absence sources. Each graphic is a pair of databases
(National combined GBIF and Vigie Nature). Each point is an individual species that is listed in both
databases, some species are assessed only in one database (missing). A Boyce's index threshold of 0.3
is used to assess the proportion of satisfactory and unsatisfactory models conjointly for each of the
two database sources. The blue quadrant contains species whose SDMs are evaluated in a
satisfactory manner in both data bases (Boyce index > 0.3 for both databases) ; the red quadrant
contains the species for which SDMs are evaluated in an unsatisfactory manner in both databases
(Boyce index < 0.3 for both databases). In the top-left quadrant and the bottom-right quadrants, SDM
model evaluation was satisfactory for only one database (on the in y and x axes respectively).

936
937

B.3 - Overlap between SDM from individual database sources. VN is Vigie Nature database and National
combined GBIF and Vigie Nature.

B.4 - Overlap between SDM from individual database sources and SDM from mixed databases (i.e. using
absence from Vigie Nature). National combine of GBIF and Vigie Nature.

B.5 - Maps of conservation planning area for Aves and Papilionidae of tree study sites (T1, T2, T3) from
different individual database source (GBIF, Vigie Nature and "National databases" combining the two
databases).

947
948

949 *B.6 - Evaluation of the quality of the full model (Table 5) compared to null model using AIC.*

Model	Aves	Papilionidae
Full	3057	1322
Null	3357	1574

 B.7 – Species list observed in each study site (T1, T2, T3) and their French status from TAXREF.V14
952   (Presence, Introduced, Invasive introduced) and IUCN regional red list from BD.STATUT.V16 (LC, NT,
953   VU, EN, CR, DD, NA) for A) Aves species and B) Papilionidae Species.
954 **B.7.A – Aves species**

Species	Observed			French status	IUCN regional red list		
	T1	T2	T3		T1	T2	T3
Accipiter gentilis	Yes	Yes	Yes	Present	LC	VU	EN
Accipiter nisus	Yes	Yes	Yes	Present	LC	LC	LC
Acrocephalus arundinaceus	Yes	Yes	-	Present	VU	CR	-
Acrocephalus schoenobaenus	-	Yes	Yes	Present	-	VU	LC
Acrocephalus scirpaceus	Yes	Yes	-	Present	NT	VU	-
Actitis hypoleucos	Yes	Yes	Yes	Present	EN	CR	NA
Aegithalos caudatus	Yes	Yes	Yes	Present	LC	LC	LC
Aegypius monachus	Yes	-	-	Present	CR	-	-
Aix galericulata	Yes	-	-	Introduced	NA	-	-
Alauda arvensis	Yes	Yes	Yes	Present	LC	VU	LC
Alcedo atthis	Yes	Yes	Yes	Present	NT	NT	LC
Alectoris rufa	Yes	Yes	Yes	Present	DD	DD	DD
Anas acuta	-	-	Yes	Present	-	-	VU
Anas crecca	-	-	Yes	Present	-	-	LC
Anas platyrhynchos	Yes	Yes	Yes	Present	DD	LC	LC
Anser anser	-	Yes	Yes	Present	-	VU	NA
Anthus campestris	Yes	Yes	-	Present	VU	EN	-
Anthus pratensis	Yes	Yes	Yes	Present	VU	EN	VU
Anthus spinoletta	-	Yes	Yes	Present	-	LC	LC
Anthus trivialis	Yes	Yes	Yes	Present	LC	LC	LC
Apus apus	Yes	Yes	Yes	Present	LC	NT	LC
Aquila chrysaetos	Yes	-	-	Present	VU	-	-
Aquila fasciata	Yes	-	-	Present	CR	-	-
Ardea alba	Yes	Yes	Yes	Present	VU	NT	EN
Ardea cinerea	Yes	Yes	Yes	Present	LC	LC	LC
Ardea purpurea	Yes	Yes	-	Present	EN	VU	-
Ardeola ralloides	Yes	-	-	Present	VU	-	-
Arenaria interpres	-	Yes	-	Present	-	NA	-
Asio otus	Yes	Yes	-	Present	LC	LC	-
Athene noctua	Yes	Yes	Yes	Present	NT	NT	VU
Aythya ferina	-	-	Yes	Present	-	-	EN
Aythya fuligula	-	-	Yes	Present	-	-	LC
Botaurus stellaris	Yes	-	-	Present	EN	-	-
Branta bernicla	-	Yes	Yes	Present	-	NA	LC
Bubo bubo	Yes	-	-	Present	LC	-	-
Bubulcus ibis	Yes	Yes	Yes	Present	LC	LC	NT
Burhinus oedicnemus	Yes	Yes	-	Present	LC	NT	-
Buteo buteo	Yes	Yes	Yes	Present	LC	LC	LC
Calandrella brachydactyla	Yes	-	-	Present	EN	-	-
Calidris alba	-	Yes	-	Present	-	NA	-
Calidris alpina	-	Yes	-	Present	-	NA	-
Caprimulgus europaeus	Yes	-	Yes	Present	LC	-	LC
Carduelis carduelis	Yes	Yes	Yes	Present	VU	NT	LC
Cecropis daurica	Yes	-	-	Present	VU	-	-
Certhia brachydactyla	Yes	Yes	Yes	Present	LC	LC	LC
Certhia familiaris	Yes	-	-	Present	LC	-	-
Cettia cetti	Yes	Yes	Yes	Present	LC	LC	LC
Charadrius alexandrinus	-	Yes	-	Present	-	EN	-
Charadrius dubius	Yes	Yes	-	Present	NT	VU	-
Charadrius hiaticula	Yes	Yes	-	Present	VU	VU	-
Chlidonias niger	Yes	Yes	-	Present	NA	CR	-
Chloris chloris	Yes	Yes	Yes	Present	NT	NT	LC
Chroicocephalus ridibundus	Yes	Yes	Yes	Present	LC	VU	LC
Ciconia ciconia	Yes	Yes	-	Present	NT	NT	-
Ciconia nigra	Yes	-	-	Present	EN	-	-
Cinclus cinclus	Yes	-	-	Present	LC	-	-
Circaetus gallicus	Yes	Yes	-	Present	LC	EN	-
Circus aeruginosus	Yes	Yes	-	Present	VU	VU	-
Circus cyaneus	Yes	Yes	Yes	Present	EN	NT	EN
Circus pygargus	Yes	Yes	-	Present	EN	NT	-
Cisticola juncidis	Yes	Yes	Yes	Present	LC	NT	LC
Clamator glandarius	Yes	-	-	Present	NT	-	-

Coccothraustes coccothraustes	Yes	Yes	Yes	Present	LC	NT	VU
Columba livia	Yes	Yes	Yes	Present	DD	DD	DD
Columba oenas	Yes	Yes	Yes	Present	VU	EN	LC
Columba palumbus	Yes	Yes	Yes	Present	LC	LC	LC
Coracias garrulus	Yes	-	-	Present	NT	-	-
Corvus corax	Yes	-	-	Present	LC	-	-
Corvus corone	Yes	Yes	Yes	Present	LC	LC	LC
Corvus frugilegus	-	Yes	Yes	Present	-	LC	LC
Corvus monedula	Yes	Yes	Yes	Present	LC	NT	LC
Coturnix coturnix	Yes	Yes	Yes	Present	NT	VU	LC
Cuculus canorus	Yes	Yes	Yes	Present	LC	LC	LC
Cyanistes caeruleus	Yes	Yes	Yes	Present	LC	LC	LC
Cygnus olor	Yes	Yes	-	Present	NA	LC	-
Delichon urbicum	Yes	Yes	Yes	Present	LC	NT	LC
Dendrocopos major	Yes	Yes	Yes	Present	LC	LC	LC
Dendrocopos minor	Yes	Yes	Yes	Present	LC	NT	LC
Dryocopus martius	Yes	-	Yes	Present	LC	-	LC
Egretta garzetta	Yes	Yes	Yes	Present	LC	LC	NT
Elanus caeruleus	-	Yes	-	Present	-	VU	-
Emberiza calandra	Yes	Yes	Yes	Present	LC	VU	EN
Emberiza cirlus	Yes	Yes	Yes	Present	LC	LC	LC
Emberiza citrinella	Yes	Yes	Yes	Present	NT	NT	NT
Emberiza hortulana	Yes	-	-	Present	VU	-	-
Emberiza schoeniclus	Yes	Yes	Yes	Present	EN	EN	VU
Erithacus rubecula	Yes	Yes	Yes	Present	LC	LC	LC
Falco columbarius	-	Yes	-	Present	-	NA	-
Falco naumanni	Yes	-	-	Present	VU	-	-
Falco peregrinus	Yes	Yes	-	Present	VU	CR	-
Falco subbuteo	Yes	Yes	Yes	Present	NT	NT	NT
Falco tinnunculus	Yes	Yes	Yes	Present	LC	NT	LC
Falco vespertinus	Yes	Yes	-	Present	NA	NA	-
Ficedula albicollis	Yes	-	-	Present	NT	-	-
Ficedula hypoleuca	Yes	-	-	Present	EN	-	-
Fringilla coelebs	Yes	Yes	Yes	Present	LC	LC	LC
Fringilla montifringilla	-	-	Yes	Present	-	-	DD
Fulica atra	Yes	Yes	Yes	Present	LC	LC	LC
Galerida cristata	Yes	Yes	-	Present	LC	LC	-
Gallinago gallinago	Yes	Yes	Yes	Present	CR	CR	NA
Gallinula chloropus	Yes	Yes	Yes	Present	LC	NT	LC
Garrulus glandarius	Yes	Yes	Yes	Present	LC	LC	LC
Grus grus	Yes	Yes	-	Present	CR	CR	-
Gyps fulvus	Yes	-	-	Present	VU	-	-
Hieraaetus pennatus	Yes	-	-	Present	VU	-	-
Himantopus himantopus	Yes	Yes	-	Present	LC	NT	-
Hippolais polyglotta	Yes	Yes	Yes	Present	LC	LC	LC
Hirundo rustica	Yes	Yes	Yes	Present	NT	NT	LC
Ichthyaetus melanocephalus	-	Yes	-	Present	-	CR	-
Jynx torquilla	Yes	Yes	-	Present	NT	VU	-
Lanius collurio	Yes	Yes	Yes	Present	NT	NT	EN
Lanius excubitor	-	-	Yes	Present	-	-	NA
Lanius meridionalis	Yes	-	-	Present	EN	-	-
Lanius senator	Yes	-	-	Present	NT	-	-
Larus argentatus	-	Yes	Yes	Present	-	VU	VU
Larus canus	-	Yes	-	Present	-	EN	-
Larus fuscus	Yes	Yes	Yes	Present	NA	LC	LC
Larus marinus	-	Yes	-	Present	-	EN	-
Larus michahellis	Yes	Yes	-	Present	LC	VU	-
Limosa lapponica	-	Yes	-	Present	-	NA	-
Limosa limosa	-	Yes	-	Present	-	CR	-
Linaria cannabina	Yes	Yes	Yes	Present	NT	NT	LC
Locustella luscinioides	-	Yes	-	Present	-	EN	-
Locustella naevia	Yes	Yes	Yes	Present	DD	VU	LC
Lophophanes cristatus	Yes	Yes	Yes	Present	LC	VU	LC
Loxia curvirostra	Yes	-	-	Present	LC	-	-
Lullula arborea	Yes	Yes	Yes	Present	LC	NT	LC
Luscinia megarhynchos	Yes	Yes	Yes	Present	LC	LC	VU
Luscinia svecica	Yes	Yes	-	Present	LC	LC	-
Mareca penelope	-	-	Yes	Present	-	-	LC
Mareca strepera	-	-	Yes	Present	-	-	LC
Mergus merganser	-	-	Yes	Present	-	-	NA
Merops apiaster	Yes	Yes	-	Present	NT	VU	-

Milvus migrans	Yes	Yes	-	Present	LC	LC	-
Milvus milvus	Yes	-	-	Present	EN	-	-
Monticola saxatilis	Yes	-	-	Present	VU	-	-
Monticola solitarius	Yes	-	-	Present	VU	-	-
Motacilla alba	Yes	Yes	Yes	Present	LC	LC	LC
Motacilla cinerea	Yes	-	Yes	Present	LC	-	LC
Motacilla flava	Yes	Yes	-	Present	NT	LC	-
Muscicapa striata	Yes	Yes	Yes	Present	LC	NT	LC
Neophron percnopterus	-	Yes	-	Present	-	EN	-
Numenius arquata	Yes	Yes	-	Present	CR	EN	-
Numenius phaeopus	-	Yes	-	Present	-	NA	-
Nycticorax nycticorax	Yes	Yes	-	Present	NT	VU	-
Oenanthe oenanthe	Yes	Yes	Yes	Present	NT	EN	EN
Oriolus oriolus	Yes	Yes	-	Present	LC	LC	-
Otus scops	Yes	-	-	Present	NT	-	-
Pandion haliaetus	Yes	-	-	Present	VU	-	-
Parus major	Yes	Yes	Yes	Present	LC	LC	LC
Passer domesticus	Yes	Yes	Yes	Present	LC	NT	LC
Passer montanus	Yes	Yes	Yes	Present	NT	EN	EN
Perdix perdix	Yes	Yes	Yes	Present	NT	DD	DD
Periparus ater	Yes	-	Yes	Present	LC	-	NT
Pernis apivorus	Yes	Yes	-	Present	LC	VU	-
Petronia petronia	Yes	-	-	Present	LC	-	-
Phalacrocorax carbo	Yes	Yes	Yes	Present	NA	VU	LC
Phasianus colchicus	Yes	Yes	Yes	Introduced	NA	DD	DD
Phoenicurus ochruros	Yes	Yes	Yes	Present	LC	LC	LC
Phoenicurus phoenicurus	Yes	Yes	Yes	Present	LC	LC	VU
Phylloscopus bonelli	Yes	Yes	-	Present	LC	NT	-
Phylloscopus collybita	Yes	Yes	Yes	Present	LC	LC	LC
Phylloscopus sibilatrix	Yes	-	Yes	Present	EN	-	NT
Phylloscopus trochilus	Yes	Yes	Yes	Present	NA	CR	EN
Pica pica	Yes	Yes	Yes	Present	LC	LC	LC
Picus canus	Yes	Yes	-	Present	EN	CR	-
Picus viridis	Yes	Yes	Yes	Present	LC	LC	LC
Platalea leucorodia	-	Yes	-	Present	-	EN	-
Pluvialis apricaria	-	-	Yes	Present	-	-	LC
Pluvialis squatarola	-	Yes	-	Present	-	NA	-
Podiceps cristatus	Yes	-	Yes	Present	LC	-	LC
Poecile palustris	Yes	Yes	Yes	Present	LC	VU	NT
Prunella collaris	Yes	-	-	Present	EN	-	-
Prunella modularis	Yes	Yes	Yes	Present	LC	LC	LC
Ptyonoprogne rupestris	Yes	-	-	Present	LC	-	-
Pyrrhocorax pyrrhocorax	Yes	-	-	Present	VU	-	-
Pyrrhula pyrrhula	Yes	-	Yes	Present	VU	-	VU
Rallus aquaticus	Yes	Yes	-	Present	LC	VU	-
Recurvirostra avosetta	-	Yes	-	Present	-	VU	-
Regulus ignicapilla	Yes	Yes	Yes	Present	LC	LC	LC
Regulus regulus	Yes	-	Yes	Present	LC	-	LC
Riparia riparia	Yes	Yes	Yes	Present	EN	NT	LC
Saxicola rubetra	Yes	Yes	-	Present	EN	CR	-
Saxicola rubicola	Yes	Yes	Yes	Present	VU	NT	LC
Scolopax rusticola	Yes	-	-	Present	DD	-	-
Serinus serinus	Yes	Yes	Yes	Present	LC	NT	LC
Sitta europaea	Yes	Yes	Yes	Present	LC	LC	LC
Spatula clypeata	Yes	Yes	Yes	Present	DD	VU	LC
Spatula querquedula	-	Yes	-	Present	-	CR	-
Spinus spinus	Yes	-	-	Present	VU	-	-
Sterna hirundo	Yes	Yes	-	Present	LC	VU	-
Streptopelia decaocto	Yes	Yes	Yes	Present	LC	LC	LC
Streptopelia turtur	Yes	Yes	Yes	Present	LC	VU	LC
Strix aluco	Yes	Yes	Yes	Present	LC	LC	DD
Sturnus vulgaris	Yes	Yes	Yes	Present	LC	LC	LC
Sylvia atricapilla	Yes	Yes	Yes	Present	LC	LC	LC
Sylvia borin	Yes	Yes	Yes	Present	LC	NT	LC
Sylvia cantillans	Yes	Yes	-	Present	LC	LC	-
Sylvia communis	Yes	Yes	Yes	Present	LC	NT	LC
Sylvia conspicillata	-	Yes	-	Present	-	EN	-
Sylvia hortensis	Yes	-	-	Present	LC	-	-
Sylvia melanocephala	Yes	-	-	Present	LC	-	-
Sylvia undata	Yes	-	Yes	Present	VU	-	LC
Tachybaptus ruficollis	Yes	Yes	Yes	Present	LC	LC	LC

Species				French status			
Tachymarptis melba	Yes	-	-	Present	VU	-	-
Tadorna ferruginea	-	Yes	-	Present	-	NA	-
Tadorna tadorna	Yes	Yes	-	Present	LC	LC	-
Tetrax tetrax	Yes	-	-	Present	NT	-	-
Thalasseus sandvicensis	Yes	Yes	-	Present	VU	NT	-
Tichodroma muraria	Yes	-	-	Present	CR	-	-
Tringa erythropus	-	Yes	-	Present	-	NA	-
Tringa nebularia	-	Yes	-	Present	-	NA	-
Tringa ochropus	Yes	Yes	Yes	Present	NA	NA	DD
Tringa totanus	Yes	Yes	-	Present	EN	VU	-
Troglodytes troglodytes	Yes	Yes	Yes	Present	LC	LC	LC
Turdus iliacus	Yes	-	Yes	Present	NA	-	DD
Turdus merula	Yes	Yes	Yes	Present	LC	LC	LC
Turdus philomelos	Yes	Yes	Yes	Present	LC	LC	LC
Turdus pilaris	Yes	-	Yes	Present	VU	-	DD
Turdus torquatus	-	Yes	-	Present	-	LC	-
Turdus viscivorus	Yes	Yes	Yes	Present	LC	NT	LC
Tyto alba	-	Yes	Yes	Present	-	VU	DD
Upupa epops	Yes	Yes	Yes	Present	LC	LC	LC
Vanellus vanellus	-	Yes	Yes	Present	-	VU	VU

B.7.A – Papilionidae species

Species	Observed		French status	IUCN regional red list	
	T1	T3		T1	T3
Aglais io	Yes	Yes	Present	LC	LC
Anthocharis cardamines	Yes	Yes	Present	LC	LC
Anthocharis euphenoides	Yes	-	Present	LC	-
Apatura ilia	Yes	-	Present	LC	-
Apatura iris	Yes	Yes	Present	NT	NT
Aphantopus hyperantus	Yes	Yes	Present	LC	NT
Aporia crataegi	Yes	Yes	Present	LC	LC
Araschnia levana	Yes	Yes	Present	LC	LC
Arethusana arethusa	Yes	-	Present	LC	-
Argynnis pandora	Yes	-	Present	LC	-
Argynnis paphia	Yes	Yes	Present	LC	LC
Aricia agestis	Yes	Yes	Present	LC	LC
Aricia montensis	Yes	-	Present	DD	-
Boloria dia	Yes	Yes	Present	LC	LC
Boloria euphrosyne	Yes	-	Present	LC	-
Boloria selene	Yes	-	Present	NT	-
Brenthis daphne	Yes	-	Present	LC	-
Brenthis hecate	Yes	-	Present	VU	-
Brintesia circe	Yes	-	Present	LC	-
Cacyreus marshalli	Yes	-	Invasive introduced	NA	-
Callophrys avis	Yes	-	Present	LC	-
Callophrys rubi	Yes	Yes	Present	LC	LC
Carcharodus alceae	Yes	Yes	Present	LC	LC
Celastrina argiolus	Yes	Yes	Present	LC	LC
Charaxes jasius	Yes	-	Present	LC	-
Chazara briseis	Yes	-	Present	VU	-
Coenonympha arcania	Yes	Yes	Present	LC	LC
Coenonympha dorus	Yes	-	Present	LC	-
Coenonympha pamphilus	Yes	Yes	Present	LC	LC
Colias alfacariensis	Yes	Yes	Present	LC	LC
Colias crocea	Yes	Yes	Present	LC	LC
Cupido alcetas	Yes	-	Present	LC	-
Cupido argiades	-	Yes	Present	-	NT
Cupido minimus	Yes	-	Present	LC	-
Cupido osiris	Yes	-	Present	NT	-
Cyaniris semiargus	Yes	Yes	Present	LC	NT
Erebia aethiops	Yes	-	Present	NT	-
Erebia epistygne	Yes	-	Present	EN	-
Erebia meolans	Yes	-	Present	LC	-
Erebia neoridas	Yes	-	Present	NT	-
Erynnis tages	Yes	Yes	Present	LC	LC
Euchloe crameri	Yes	-	Present	LC	-
Euphydryas aurinia	Yes	-	Present	NT	-
Fabriciana adippe	Yes	-	Present	NT	-

Fabriciana niobe	Yes	-	Present	NT	-
Glaucopsyche alexis	Yes	-	Present	LC	-
Glaucopsyche melanops	Yes	-	Present	LC	-
Gonepteryx cleopatra	Yes	-	Present	LC	-
Gonepteryx rhamni	Yes	Yes	Present	LC	LC
Hamearis lucina	Yes	-	Present	LC	-
Hesperia comma	Yes	-	Present	LC	-
Hipparchia alcyone	Yes	-	Present	DD	-
Hipparchia fagi	Yes	-	Present	LC	-
Hipparchia fidia	Yes	Yes	Present	LC	LC
Hipparchia genava	Yes	-	Present	NT	-
Hipparchia semele	Yes	Yes	Present	LC	EN
Hipparchia statilinus	Yes	Yes	Present	LC	EN
Hyponephele lupina	Yes	-	Present	EN	-
Hyponephele lycaon	Yes	-	Present	EN	-
Iberochloe tagis	Yes	-	Present	EN	-
Iphiclides podalirius	Yes	-	Present	LC	-
Issoria lathonia	Yes	Yes	Present	LC	LC
Laeosopis roboris	Yes	-	Present	LC	-
Lampides boeticus	Yes	-	Present	LC	-
Lasiommata maera	Yes	Yes	Present	LC	CR
Lasiommata megera	Yes	Yes	Present	LC	LC
Leptidea sinapis	Yes	Yes	Present	LC	LC
Leptotes pirithous	Yes	-	Present	LC	-
Libythea celtis	Yes	-	Introduced	LC	-
Limenitis camilla	Yes	Yes	Present	LC	LC
Limenitis reducta	Yes	-	Present	LC	-
Lycaena alciphron	Yes	-	Present	LC	-
Lycaena dispar	Yes	-	Present	NT	-
Lycaena phlaeas	Yes	Yes	Present	LC	LC
Lycaena tityrus	Yes	Yes	Present	LC	LC
Lysandra bellargus	Yes	-	Present	LC	-
Lysandra coridon	Yes	-	Present	LC	-
Lysandra hispana	Yes	-	Present	LC	-
Maniola jurtina	Yes	Yes	Present	LC	LC
Melanargia galathea	Yes	Yes	Present	LC	LC
Melanargia lachesis	Yes	-	Present	LC	-
Melanargia occitanica	Yes	-	Present	LC	-
Melanargia russiae	Yes	-	Present	VU	-
Melitaea athalia	Yes	-	Present	DD	-
Melitaea celadussa	Yes	-	Present	LC	-
Melitaea cinxia	Yes	Yes	Present	LC	LC
Melitaea deione	Yes	-	Present	DD	-
Melitaea diamina	Yes	-	Present	NT	-
Melitaea didyma	Yes	-	Present	LC	-
Melitaea parthenoides	Yes	-	Present	LC	-
Melitaea phoebe	Yes	Yes	Present	LC	LC
Minois dryas	Yes	-	Present	LC	-
Muschampia floccifera	Yes	-	Present	NT	-
Muschampia lavatherae	Yes	-	Present	NT	-
Muschampia proto	Yes	-	Present	NT	-
Nymphalis antiopa	Yes	-	Present	NT	-
Nymphalis polychloros	Yes	Yes	Present	LC	LC
Ochlodes sylvanus	Yes	Yes	Present	LC	LC
Papilio machaon	Yes	Yes	Present	LC	LC
Pararge aegeria	Yes	Yes	Present	LC	LC
Parnassius apollo	Yes	-	Present	VU	-
Phengaris alcon	Yes	-	Present	VU	-
Phengaris arion	Yes	-	Present	NT	-
Pieris brassicae	Yes	Yes	Present	LC	LC
Pieris mannii	Yes	-	Present	DD	-
Pieris napi	Yes	Yes	Present	LC	LC
Pieris rapae	Yes	Yes	Present	LC	LC
Plebejus argus	Yes	Yes	Present	LC	NT
Plebejus idas	Yes	Yes	Present	NT	CR
Polygonia c-album	Yes	Yes	Present	LC	LC
Polyommatus amandus	Yes	-	Present	VU	-
Polyommatus daphnis	Yes	-	Present	VU	-
Polyommatus dolus	Yes	-	Present	VU	-
Polyommatus dorylas	Yes	-	Present	VU	-
Polyommatus escheri	Yes	-	Present	LC	-

Polyommatus icarus	Yes	Yes	Present	LC	LC
Polyommatus thersites	Yes	-	Present	DD	-
Pontia daplidice	Yes	-	Present	LC	-
Pseudophilotes baton	Yes	-	Present	LC	-
Pyrgus alveus	Yes	-	Present	NT	-
Pyrgus armoricanus	Yes	-	Present	LC	-
Pyrgus carthami	Yes	-	Present	NT	-
Pyrgus cirsii	Yes	-	Present	VU	-
Pyrgus foulquieri	Yes	-	Present	EN	-
Pyrgus malvae	Yes	-	Present	DD	-
Pyrgus malvoides	Yes	-	Present	LC	-
Pyrgus onopordi	Yes	-	Present	DD	-
Pyronia bathseba	Yes	-	Present	LC	-
Pyronia cecilia	Yes	-	Present	LC	-
Pyronia tithonus	Yes	Yes	Present	LC	LC
Quercusia quercus	Yes	Yes	Present	LC	LC
Satyrium acaciae	Yes	-	Present	LC	-
Satyrium esculi	Yes	-	Present	LC	-
Satyrium ilicis	Yes	Yes	Present	LC	LC
Satyrium pruni	Yes	-	Present	DD	-
Satyrium spini	Yes	-	Present	LC	-
Satyrium w-album	Yes	-	Present	LC	-
Satyrus actaea	Yes	-	Present	VU	-
Satyrus ferula	Yes	-	Present	VU	-
Speyeria aglaja	Yes	-	Present	LC	-
Spialia sertorius	Yes	-	Present	LC	-
Thecla betulae	Yes	-	Present	LC	-
Thymelicus acteon	Yes	-	Present	LC	-
Thymelicus lineola	Yes	Yes	Present	LC	LC
Thymelicus sylvestris	Yes	Yes	Present	LC	LC
Vanessa atalanta	Yes	Yes	Present	LC	LC
Vanessa cardui	Yes	Yes	Present	LC	LC
Zerynthia polyxena	Yes	-	Present	LC	-
Zerynthia rumina	Yes	-	Present	LC	-

957