Deletion of miR-146a enhances therapeutic protein restoration in model of dystrophin exon skipping

Nikki M. McCormack[1], Kelsey A. Calabrese[1], Christina M. Sun[1], Christopher B. Tully[1], Christopher R. Heier[1,2], Alyson A. Fiorillo[1,2]

[1]Center for Genetic Medicine Research, Children's National Hospital, Washington, District of Columbia, USA

[2]Department of Genomics and Precision Medicine, George Washington University School of Medicine and Health Sciences, Washington, District of Columbia, USA

Correspondence should be addressed to:

Alyson A. Fiorillo, PhD

Center for Genetic Medicine Research, Children's Research Institute
7144 13th Place NW, Washington DC, 20012
Email: afiorillo@childrensnational.org

Tel: 202-545-2813

Fax:

Short title: miR-146a deletion improves dystrophin restoration

Abstract (200 words max, no abbreviations)

Duchenne muscular dystrophy (DMD) is a progressive muscle disease caused by the absence

of dystrophin protein. One current DMD therapeutic strategy, exon skipping, produces a

truncated dystrophin isoform using phosphorodiamidate morpholino oligomers (PMOs).

However, the potential of exon skipping therapeutics has not been fully realized as increases in

dystrophin protein have been minimal in clinical trials. Here, we investigate how miR-146a-5p,

which is highly elevated in dystrophic muscle, impacts dystrophin protein levels. We find

inflammation strongly induces miR-146a in dystrophic, but not wild-type myotubes.

Bioinformatics analysis reveals that the dystrophin 3'UTR harbors a miR-146a binding site, and

subsequent luciferase assays demonstrate miR-146a binding inhibits dystrophin translation. In

dystrophin-null *mdx52* mice, co-injection of miR-146a reduces dystrophin restoration by an exon

51 skipping PMO. To directly investigate how miR-146a impacts therapeutic dystrophin rescue,

we generated *mdx52* with body-wide miR-146a deletion (*146aX*). Administration of an exon

skipping PMO via intramuscular or intravenous injection markedly increases dystrophin protein

levels in *146aX* versus *mdx52* muscles; skipped dystrophin transcript levels are unchanged,

suggesting a post-transcriptional mechanism-of-action. Together, these data show that miR-

146a expression opposes therapeutic dystrophin restoration, suggesting miR-146a inhibition

warrants further research as a potential DMD exon skipping co-therapy.

Introduction (no longer than 2 pages**)**

Duchenne muscular dystrophy (DMD) is a progressive muscle disease caused by out-of-frame mutations in the dystrophin gene resulting in the complete loss of dystrophin protein[1]. DMD is characterized by severe muscle weakness and chronic muscle inflammation leading to loss of ambulation and a shortened lifespan. Becker muscular dystrophy (BMD) is an allelic disease where the reading frame is preserved, resulting in production of a truncated dystrophin isoform that is expressed at lower and more variable levels[2-4]. BMD is less severe and shows high variability both in clinical presentation and in levels of dystrophin protein even when comparing subjects harboring the same in-frame mutation[5-8]. Interestingly, while dystrophin protein levels are variable and lower-than-normal in BMD, there are no differences in the amount of dystrophin mRNA detected in BMD as compared to unaffected/healthy muscles[7, 8].

Precision medicine strategies for DMD such as gene therapies and exon skipping antisense oligonucleotides (AOs) seek to convert a severe DMD genotype (dystrophin null) into a milder BMD (truncated dystrophin) phenotype. Specifically, exon skipping AOs act by binding to a complementary sequence on the dystrophin pre-mRNA to promote the exclusion of the frame-shifting exon. Currently four exon skipping phosphorodiamidate morpholino oligomers (PMOs) have received accelerated FDA approval; these target dystrophin exons 51 (Eteplirsen), 53 (Golodirsen, Viltepso) and 45 (Casimersen)[9-12]. However, results from clinical trial data show low and variable dystrophin rescue. As an example, for the first approved exon skipping drug (Eteplirsen), the Western blot analysis showed an average of 0.93% dystrophin in treated muscles compared with a baseline value of 0.08% in biopsies of untreated Duchenne muscles[13]. In the later Golodirsen clinical trial, mean dystrophin protein levels measured via Western blot were 1.019% of normal dystrophin levels despite showing greater levels of skipped dystrophin transcripts (16.363%)[10, 14].

The disconnect between the extent of dystrophin exon skipping (mRNA) and dystrophin protein levels point to post-transcriptional factors in the muscle that affect total dystrophin protein

production. microRNAs (miRNAs) are elevated in a variety of muscle diseases; their canonical

role is to fine-tune gene expression by binding to the 3ʹUTR of target genes and most often

function to regulate protein levels post-transcriptionally via translation inhibition[7, 15-17].

Previously, we identified several miRNAs that bind the dystrophin 3'UTR and inhibit dystrophin

protein production (termed dystrophin targeting miRNAs or DTMs)[7]. These include miR-146a-

5p, which is elevated in DMD and BMD muscle biopsies and is induced by inflammation *in vitro*,

suggesting regulation by the pro-inflammatory transcription factor NF-κB. Indeed, others have

demonstrated NF-κB-specific regulation of miR-146a using cell culture studies[18] and

bioinformatic analysis from ENCODE data[19]. Importantly, our lab previously found that levels of

miR-146a were inversely associated with dystrophin restoration in the muscles of DMD model

(*mdx)* mice treated with an exon skipping PMO[7]. In a separate study, we identified a panel of 9

inflammatory miRNAs that are elevated in the *mdx* diaphragm and are reduced by treatment

with the corticosteroid prednisone or the dissociative steroid vamorolone; included in this panel

is the DTM miR-146a[20]. Finally, we recently reported that treating BMD model (*bmx*) mice with

prednisone or vamorolone dampens expression of DTMs including miR-146a and increases

dystrophin protein levels in the gastrocnemius and heart of bmx mice by ~50%[21]. Collectively,

these data suggest that miR-146a is regulated by inflammation, is elevated in dystrophic

muscle, and inhibits dystrophin protein production.

In the present study, we hypothesized that genetic deletion of miR-146a specifically

would improve dystrophin restoration in dystrophin-deficient mice harboring a deletion of *Dmd*

exon 52 (*mdx52*). To test this, we generated double knockout mice and carried out a proof-of-

concept study, performing high dose systemic administration of an exon 51 skipping PMO to

mdx52 and *mdx52*;*miR-146a*[-/-] (*146aX*) mice, where our primary readout was extent of

dystrophin protein restoration.   Excitingly, following both intramuscular and systemic delivery of

PMO, we show deletion of miR-146a significantly increases dystrophin protein levels.

Methods

Animal care and maintenance

All animal studies were done in adherence to the NIH Guide for the Care and Use of Laboratory

Animals. All experiments were conducted according to protocols within the guidelines and under

approval of the Institutional Animal Care and Use Committee of Children's National Hospital.

mdx52 mice contain a deletion of exon 52 of the *Dmd* gene, resulting in the absence of full-

length dystrophin. These mice on a C57/BL6 background are maintained in our own facility and

were originally a kind gift from Dr. Shin'ichi Takeda. Male C57/BL6 mice were purchased from

JAX laboratory (Bar Harbor, ME). *bmx* mice contain an endogenous deletion of murine

dystrophin exons 45-47, are maintained on a C57/BL6J background, and were previously

characterized [8].

Muscle biopsies

Human biopsies used here were banked diagnostic biopsies obtained from the vastus lateralis

muscle. Six BMD muscle biopsies were obtained from the Eurobiobank, Telethon Network

Genetic Biobanks (GTB 12001D) (ID #: 2518, 3010, 3419, 4228, 7560, 7827). The remaining 4

BMD biopsies were obtained from our muscle biopsy bank as previously reported (ID#: 1141,

3212, 3255, 3461)[4, 7].

Generation of mdx/146a$^{-/-}$ mice

mdx52 mice (*dmd$^{-/-}$* or *dmd$^{-/Y}$*) on a C57/BL/6 background are bred in house[22]. *miR-146a-/-*

(B6.Cg-*Mir146$^{tm1.1Bal}$*/J) have been previously described[23] and were purchased from The

Jackson Laboratory (Strain # 016239) and are on a C57/BL/6J background. To generate *mdx52*

mice with deletion of miR-146a, dmd$^{-/-}$;146a$^{+/+}$ females were crossed with dmd$^{+/Y}$;146a$^{-/-}$ males

to generate dmd$^{-/+}$:146a$^{+/-}$ females and dmd$^{-/Y}$;146a$^{+/-}$ males. dmd$^{-/+}$;146a$^{+/-}$ females and dmd^{-}

/Y;146a+/- males were then crossed to generate dmd-/-;146a-/- and dmd-/Y;146a-/- breeders.

Breeders were crossed to generate dmd-/Y;146a-/- mice (*146aX*). For all experiments only males

were utilized as DMD is an X-linked disorder. Genotyping for *mdx52* and miR-146a KO alleles

was performed using TransnetYX with custom primers in *Dmd* exon 3 and a neomycin cassette

for *mdx52* and with the standard JAX protocol for miR-146a-/- with forward primer 24664 (5′-GCT

TAT GAA CTT GCC TAT CTT GTG-3′) and reverse primer 24665 (5′-CAG CAG TTC CAC GCT

TCA-3′).

Cell culture and treatment of H2K myotubes

H2K (H2Kb-tsA58) WT and *mdx* myoblasts were cultured as previously described[24]. H2K *mdx*

or WT myoblasts were differentiated into myotubes in 12-well plates (1.25 × 10^5 cells/well)

with Matrigel at 37°C. For TNF-α inductions, after 4 days of differentiation, myotubes were

induced with TNF-α (10 ng/ml) for 24 hr. After 24 h, cells were lysed for RNA using TRIzol and

expression of miR-146a and control RNAs was quantified by TaqMan Assay (Thermo Fisher

Scientific). Comparison of miR-146a induction between groups was performed using a Two-

Way ANOVA with a Holms-Sidak post-hoc analysis. For LPS-treatment, seeded myoblasts were

differentiated into myotubes in 12-well plates (1.25 × 10^5 cells/well) with Matrigel at 37°C. After

4 days of differentiation, myotubes were then treated with LPS to induce inflammation (Thermo

Fisher Scientific) at a dilution of 1:1,000. After 24 h, cells were lysed for RNA using TRIzol and

expression of miR-146a and control RNAs was quantified by TaqMan Assay (Thermo Fisher

Scientific).

Cloning

Cloning was performed as previously reported[7]. Briefly, the entire 2.7kb of the dystrophin

3′UTR was amplified from normal muscle using the Flex cDNA synthesis kit (Quanta) using

primers that added 5′ XhoI and 3′ NotI restriction enzyme sequences (Forward primer "017 DMD

3ʹUTR"- 5ʹCCGCTCGAGGCG AAGTCTTTTCCACATGGCAGAT3ʹ Reverse primer "018 DMD

3ʹUTR"- 5ʹCGGCGGCCGC GTAACATAACTGCGTGCTTTATT3ʹ. The amplified product was gel

purified, digested with XhoI and NotI, and ligated into the pSiCheck2 plasmid (Promega Cat

#C8021) where renilla luciferase is upstream of the 3'UTR and firefly luciferase is driven by a

separate promoter as a reference control. Ligation products were transformed into TOP10 cells

(Life Technologies) using ampicillin selection. Positive clones were verified by diagnostic digests

and Sanger sequencing.

Luciferase in vitro Reporter Assay. HEK293 cells were seeded in 24-well plates at a density of

4×10^4 or 8×10^4 cells/well and co-transfected 24 hours later with 200 ng of the PsiCheck2

plasmid (Promega, cat #C8021) with the human dystrophin 3ʹ UTR or mutant 3ʹ UTR and with

50 nM miR-146a mimic (Life Technologies) with Lipofectamine 2000 according to the

manufacturer's protocol. Cells were harvested 24 h later according to Dual-Glow Luciferase

Reporter Assay System protocol (Promega) where renilla luciferase was the main readout

(3ʹUTR) and firefly luciferase served as the internal control.

miR-146a mimic in vivo injections

mdx52 mice were anesthetized with isoflurane and 2 µg of PMO was injected into each tibialis

anterior. Mice were co-injected with 10 µg of either a miR-146a mimic or a control miRNA

sequence (ThermoFisher scientific). Mice were euthanized two weeks after intramuscular

injection and muscles were harvested.

PMO treatment of dystrophic mice

The *Dmd* exon 51 skipping PMO utilized here is the sequence for Eteplirsen modified for the

mouse genome and has been previously reported [25]. The sequence is: (5ʹ-

CTCCAACAGCAAAGAAGATGGCATTTCTAG-3ʹ) and was synthesized by Gene Tools. For

intramuscular injections, 12- week- old mice were anesthetized with isoflurane and 2 µg of PMO

was administered into each tibialis anterior. Mice were euthanized two weeks after

intramuscular injection and muscles were harvested and frozen.

For intravenous injections, 15-week-old mice were anesthetized with isoflurane and 400 mg/kg

PMO was administered via retroorbital injection. A second IV dose of PMO (400 mg/kg) was

given seven days later. Mice were euthanized two weeks after the second injection and muscles

were harvested and frozen.

Capillary electrophoresis (Wes)

Muscles were dissected and frozen in liquid-nitrogen cooled isopentane. 8 µm sections were

lysed in 5% SDS buffer containing 10mM EDTA (pH 8.0), 75mM Tris-HCL (pH 6.8), and

protease inhibitors. Capillary Western immunoassay (Wes) analysis was performed according to

manufacturer's instructions using 66–440 kDa Separation Modules (ProteinSimple). In each

capillary 0.2mg/mL protein was loaded for analysis with antibodies to dystrophin (Abcam #

ab15277, dilution 1:15) as in [26], or vinculin (Abcam #ab130007, dilution 1:100), and anti-rabbit

secondary (ProteinSimple #042-206). Compass for SW software was used to quantify

chemiluminescence data.

qRT-PCR

Dmd exon skipping. Total RNA was extracted from muscle samples by standard TRIzol (Life

Technologies) isolation. RNA quantity and quality was determined using a nanodrop 2000.

400 ng of RNA was reverse-transcribed using Random Hexamers and High-Capacity cDNA

Reverse Transcription Kit (Thermo Fisher). Skipped and non-skipped *Dmd* transcripts were

measured by qRT-PCR using TaqMan assays and Taqman Fast Advanced MasterMix (Thermo

Fisher) on an ABI QuantStudio 7 Real-Time PCR machine (Applied Biosystems). A custom

TaqMan probe for the skipped *Dmd* product (Thermo Fisher) was designed to amplify the splice

junction spanning *Dmd* exons 50-53. TaqMan probe Mm01216492_m1 (Thermo Fisher) that

amplifies the region spanning *Dmd* exons 2–3 was utilized to calculate non-

skipped *Dmd* transcript levels. Percent exon skipping was calculated based on the

corresponding ΔCt values for skipped and total *Dmd* transcript normalized

to *Hprt* (Mm01545399_m1; Thermo Fisher) and *18S* (Mm03928990_g1; Thermo Fisher) mRNA,

using the following equation: [(Average of triplicate reactions of skipped *Dmd*)/(Average of

triplicate reactions of skipped *Dmd* + Average of triplicate reactions of non-

skipped *Dmd*)] * 100.

miRNAs. Total RNA was converted to cDNA using multiplexed RT primers and High Capacity

cDNA Reverse Transcription Kit (ThermoFisher; Carlsbad, CA). miRNAs were then quantified

using individual TaqMan assays on an ABI QuantStudio 7 real time PCR machine (Applied

Biosystems). Assay IDs include: miR-146a 000468 and controls sno202 001232, RNU48

001006. miR-146a levels were normalized to the geometric mean of the two control genes[27].

All assay IDs are listed below in Table 1:

Oligonucleotide Name	Sequence	Purpose
Dmd exons 50-53	Thermo Fisher Scientifc Custom Assay ID ARKA43N	qRT-PCR
Dmd exons 2-3	Thermo Fisher Scientific Assay ID Mm01216492_m1	qRT-PCR
Hprt	Thermo Fisher Scientific Assay ID Mm01545399_m1	qRT-PCR
18S	Thermo Fisher Scientific Assay ID Mm03928990_g1	qRT-PCR
miR-146a	Thermo Fisher Scientific Assay ID 000468	qRT-PCR
sno202	Thermo Fisher Scientific Assay ID 001232	qRT-PCR
RNU48	Thermo Fisher Scientific Assay ID 001006	qRT-PCR

Immunofluorescence of muscle samples

Muscles were mounted on cork, flash frozen, and sectioned (8 μm) onto slides. For dystrophin

and F4/80 immunofluorescence experiments muscle sections were fixed in ice cold acetone for

10 minutes. Slides were washed with 1X PBST (0.1% Tween 20), blocked for 1 hour (1X PBST

230 with 0.1% Triton X-100, 1% BSA, 10% goat serum, and 10% horse serum), washed 3

times, then exposed to primary antibodies overnight at 4ºC. For dystrophin immunostaining

these included: anti-dystrophin 1:150 (Abcam, #ab154168) as in[28] and rat anti-laminin-2 1:100

233 (Sigma-Aldrich, Cat. #L0663). For macrophage staining these included: anti-F4/80 which

recognizes the mouse F4/80 antigen, a 160kD glycoprotein expressed by murine macrophages

235 (1:100, Bio-Rad clone C1:A3-1, Cat. MCA497) and rabbit anti-laminin (1:50, Sigma-Aldrich

Cat. 236 #L9393). Secondary antibodies included: goat anti-rabbit 568 1:400 (ThermoFisher,

#A-11036), 237 donkey anti-rat 488 1:400 (ThermoFisher, #A-21208), and goat anti-rabbit 647

1:400 238 (ThermoFisher, #A-21245). Wheat germ agglutinin (WGA) conjugated to Alexa Fluor

647 (Life Technologies, MA) was prepared as 1 mg/ml stock solutions and used at a 1:500

dilution in PBS. Coverslips were mounted using Prolong Gold Mounting Medium 239 with DAPI.

Slides were imaged using an Olympus VS-120 scanning microscope at 20X.

Dystrophin immunofluorescence image analysis

Images in each experiment were thresholded to the same levels, and then exported as PNG

files and blinded using randomly assigned number IDs. To determine % dystrophin-positive

fibers, the total number of myofibers (marked by either wheat-germ agglutinin or laminin

staining) were counted manually using ImageJ. Dystrophin positive fibers were additionally

counted, and the percentage of dystrophin-positive fibers were determined by the following

formula: dystrophin-positive fibers/ total fibers x 100. For secondary analysis dystrophin high

and low fiber percentages were determined based on a visual standard (refer to **Figure 6C**).

Histological staining

Hematoxylin and eosin was performed to analyze necrosis and inflammation. Briefly, tissue

sections were stained for 10 minutes with hematoxylin and excess stain was removed in running

tap water until nuclei turned blue. Slides were incubated in 70% ethanol for 3 minutes and then

stained with eosin for 3 minutes. Tissue sections were dehydrated with ethanol and cleared with

xylene. Coverslips were mounted with permount. All slides were imaged using an Olympus VS-

120 scanning microscope at 20X.

Muscle histology analysis

Cross-sectional area (CSA) and minimum Feret's diameter were determined using the MuscleJ

macro for Fiji.[62] The variance coefficients for CSA and minimum Feret's diameter were

calculated by dividing standard deviation by the average and multiplying by 1000. Hematoxylin

and Eosin staining was performed to assay necrosis and inflammation. ImageJ was used to

outline areas of necrosis and inflammation.

Motor function tests

Motor function tests were performed in PMO treated mice 1 week after second injection (17

weeks of age). Forelimb and hindlimb grip strength was assessed using a grip strength meter

(Columbus Instruments) daily for 2 consecutive days according to Treat NMD protocols

(DMD_M.2.2.001), with data interpreted as average maximum daily values. Two-limb wire hang

and four-limb grid hang tests were performed in accordance with Treat NMD protocols

(DMD_M.2.1.005). For two-limb wire hang, a wire hanger was suspended placed ~35cm above

a cage with soft bedding. Mice were hung using only their forelimbs; however, they were

allowed to swing and hang with all four limbs if able. Hang time was recorded, with 600 seconds

used as a cutoff.

Statistical Analysis

For assays with greater than two groups, measurements were compared between groups using

one-way ANOVA unless otherwise indicated. Post hoc linear tests between each group were

performed; the resulting p value reported in figures was adjusted for multiple testing via the

Holmes-Sidak method. The contrasting groups in all post hoc comparisons are indicated in each

figure. For assays with two groups where the null hypothesis was testing a change in one

direction, a one-tailed, Student's t test was utilized, whereas in assays where the potential

change between groups was + or –, a two-tailed Student's t test was utilized. For all bar graphs,

data are presented as mean ± standard error of the mean (S.E.M). Details of statistical tests are

specified in the figure legends.

Results

Inflammation-regulated miR-146a is elevated in dystrophic muscle

We previously reported that DTMs, including miR-146a, inhibit dystrophin protein

production *in vitro* and are inversely related to exon skipping-mediated dystrophin restoration *in*

vivo[7]. One particular miRNA, miR-146a, was found to be highly elevated in BMD, DMD and *mdx*

muscle[7] and analysis of ENCODE data shows it is regulated by the pro-inflammatory

transcription factor NF-κB[19].

The miR-146a locus is found on chromosome 5q33.3 and is located in a cluster along

with the long non-coding RNA MIR3142HG. miR-146a is evolutionarily conserved across

species, as demonstrated by alignment of the miR-146a sequence from human, chimpanzee,

baboon, dog, mouse, xenopus and zebrafish **(Figure 1A**). Evaluating levels of miR-146a from

Becker muscular dystrophy human biopsies with deletion of *DMD* exons 45-47, we previously

found miR-146a levels are higher in biopsies from patients with low dystrophin levels (less than

20% of unaffected control biopsies, **Figure 1B**, data from[7].) Plotting miR-146a levels versus

dystrophin protein levels (as measured by Western blot[7]) in these same BMD 45-47 biopsies,

we observed a significant inverse correlation between miR-146a levels and the percent of

dystrophin protein in BMD biopsies (Spearman correlation, r=-0.6365, p<0.01, **Figure 1B).**

Elevated miR-146a levels in dystrophic muscle are corroborated by our analysis of previously

published data from Eisenberg et. al[15] showing miR-146a is upregulated in DMD, and BMD

patient biopsies as compared to unaffected control biopsies (**Figure 1C**). We analyzed the

diaphragm, tibialis anterior, quadriceps, gastrocnemius, and triceps muscles from 3-month-old

mdx52 mice and found elevated levels of miR-146a in all muscles analyzed (**Figure 1D**). We

also analyzed miR-146a in the *bmx* mouse model of BMD[8] with deletion of murine *Dmd* exon

45-47 at 5 months of age, and saw significantly elevated levels in the quadriceps,

gastrocnemius and triceps (**Figure 1E**).

Previous studies have demonstrated that miR-146a is regulated by NF-κB in THP-1

macrophage cells[18]. We wanted to determine how induction of inflammation affects miR-146a

levels in both *mdx* and wild-type (WT) myotubes (*mdx and WT H-2K*). Interestingly, in both WT

and mdx myotubes TNF-α increases miR-146a, however this increase is only significant in *mdx*

myotubes (**Figure 1F**). Analysis of qRT-PCR data of miR-146a also showed that 1) basal levels

of miR-146a are significantly higher in *mdx* myotubes (373% increase over WT, p<0.01), 2) the

overall induction of miR-146a in response to TNF-α is greater in mdx vs. WT (111% vs. 193%

increase in miR-146a expression p<0.0001) and 3) the absolute levels of induced miR-146a

were higher in *mdx* (555% increase, p<0.0001). We also observed high induction of miR-146a

in *mdx* myotubes upon lipopolysaccharide (LPS)-induced inflammation (**Figure S1**). Overall,

these findings show that miR-146a is upregulated in the muscle of BMD and DMD patients, as

well as in mouse models of these diseases, and that dystrophic myotubes produce higher levels

of miR-146a as compared to WT myotubes *in vitro*.

miR-146a inhibits dystrophin translation

To validate our previous finding that miR-146a regulates dystrophin translation, we

utilized the scanMiR toolkit[29] to identify putative miR-146a binding sites or microRNA response

elements (MREs) within the human dystrophin 3′UTR (**Figure 2A, B**). scanMiR identified

multiple types of binding sites relative to the miR-146a seed sequence which is located at

nucleotides 2-7 of the miRNA sequence[30]. These site types included a 7mer-m8 (perfect

complementarity of the seed sequence to the UTR and complementarity extending to nucleotide

8), a 6mer-A1, (seed sequence complementarity at positions 3-7 and additional

complementarity at nucleotide 8), a 6mer-A1 (seed sequences at positions 2-6 and an A at

position 1), a wobbled 8mer (A at position 1, seed sequence complementarity interrupted by a

single mismatch and additional complementarity at position 8), and several non-canonical

binding sites[30]. In total, scanMiR predicted 17 binding sites, although only two showed strong

dissociation constants (-log K_d <-3). The miR-146a MRE with the strongest predicted

dissociation constant (log K_d -4.425) is located at position 1281-1288 of the 3′UTR (**Figure 2B,**

2C).

Using this information, we next utilized a dystrophin 3′UTR luciferase reporter (WT) or a

reporter where the 3'UTR MRE was mutated (Mut) to eliminate base-pairing of miR-146a at

positions 2,3 and 6-8 (**Figure 2C**). WT and Mut constructs were transfected into HEK293T cells

with either a control or miR-146a mimic sequence. Analogous to our previously reported

results[7], miR-146a significantly reduced WT 3′UTR luciferase expression while mutation of the

miR-146a MRE attenuated this inhibition (**Figure 2D**). To support this *in vitro* data, we utilized

the *mdx52* mouse, which is amenable to exon 51 skipping. We performed intramuscular tibialis

anterior (TA) injections with 2 μg of an exon 51 skipping PMO and 10 μg of either a miR-146a

mimic or a control sequence (n=10 per group). Fourteen days post-injection, muscles were

harvested, and dystrophin immunofluorescence was performed. Using qRT-PCR, we observed

significantly increased miR-146a in muscles co-injected with the miR-146a mimic sequence

(**Figure S2**). Additionally, we found a significant decrease in dystrophin positive fibers in

muscles co-injected with miR-146a as compared to muscle injected with a control sequence

(36.1% decrease, $p<0.05$, **Figure 2E, E′**). Together these data suggest that miR-146a binds to

the dystrophin 3′UTR to regulate its translation and high levels of miR-146a are inhibitory to

successful dystrophin rescue via exon skipping.

Generation of mdx/146a$^{-/-}$ (146aX) mice

To investigate the potential therapeutic implications of miR-146a, we generated *mdx52*

mice with body-wide deletion of miR-146a (*146aX*) by crossing several generations of *mdx52*

and miR-*146a-/-* mice both *on* a C57/BL/6 background **(Figure 3A)**. Deletion of miR-146a was

determined by genotyping and validated by qRT-PCR in the gastrocnemius of 3-month-old mice

(**Figure 3B**). The primary goal of generating these mice was to perform a proof-of-concept study

to determine how deletion of miR-146a affects dystrophin restoration in *mdx52* mice via exon

skipping. It has previously been shown that in dystrophin-deficient *mdx* mice with a nonsense

mutation in exon 23 (*mdx23*), muscles contain sporadic clusters of dystrophin-expressing

revertant fibers[31]. Revertant fibers occur due to spontaneous exon skipping (alternative splicing)

and tend to accumulate as a function of age. Previous work has demonstrated that *mdx52*

muscles have less revertant fibers as compared to *mdx23* mice[32]. We performed dystrophin

immunofluorescence on muscle sections from wild-type, *mdx52* and *146aX* quadricep muscles

to observe the baseline levels of revertant fibers in *mdx52* versus *146aX* muscles. As shown in

Figure 3C there were minimal revertant fibers in both genotypes which totaled 0.16% in mdx

and 0.14% of total fibers in *146aX* muscles (ns, *p*=0.6830). As typical high dose exon skipping

experiments show 5-80% dystrophin positive fibers[33] (depending on the specific muscle) the

basal levels of revertant fibers observed in both *mdx* and 146aX here are negligible in

comparison.

Body-wide deletion of miR-146a increases dystrophin rescue after intramuscular PMO delivery

We next injected an exon 51 skipping PMO (2µg) into the TAs of *mdx52* and *146aX* mice

(**Figure 4A**). Fourteen days post-injection, muscles were harvested and analyzed for extent of

exon skipping and dystrophin protein. To analyze exon skipping we performed qRT-PCR on

both *Dmd* exon 2 (total *Dmd* transcript levels) and the novel junction between *Dmd* exon 50 and

53 (skipped *Dmd* transcript levels) to determine the percent exon skipping achieved in each

muscle as we previously described[34]. Levels of exon skipping ranged between 0.5% and 7%,

and there were no significant differences observed in the extent of exon skipping when

comparing *mdx52* and *146aX* TAs (p=0.48, **Figure 4B**). We next performed capillary Western

immunoassays (Wes), as dystrophin quantification using this method has proven to be highly

sensitive, reproducible, and quantitative over a large dynamic range[26]. While no differences in

exon skipped *Dmd* transcript levels were observed, Wes protein quantification revealed a

significant increase in dystrophin levels in *146aX* muscles (~19% increase, p<0.05, **Figure 4C,**

C′). Supporting this, dystrophin immunofluorescence showed a greater than 2-fold increase in

dystrophin positive fibers in *146aX* TAs (p<0.01, **Figure 4D, D′**).

Body-wide deletion of miR-146a increases dystrophin rescue after systemic PMO treatment

We next performed systemic injections of a high dose PMO into 15-week-old *mdx52* or

146aX mice (400 mg/kg) n=4 mice per group. One week later we performed a second high dose

injection (400 mg/kg). Mice were phenotyped one week after the second dose of PMO and then

sacrificed 2 weeks after the final dose. Diaphragm (Dia), tibialis anterior (TA), quadriceps (Quad),

gastrocnemius (Gastroc), and Triceps (Tri) were collected for RNA and protein analysis (**Figure**

5A). When we analyzed the extent of exon skipping across muscles at the RNA level via qRT-

PCR, no significant differences were observed between groups (**Figure 5B**), however, there

was a non-significant trend towards increased exon skipping in the *mdx52* versus *146aX*

muscles. Conversely, analysis of dystrophin protein levels via Wes showed significantly

increased dystrophin in *146aX* muscles as compared to *mdx52* (~55% increase, p<0.05, **Figure**

5C, C′ Figure S3). To account for differences in PMO delivery between muscles and genotypes,

we also normalized dystrophin protein levels as measured by Wes, to exon skipping RNA levels

measured by qRT-PCR. Interestingly, using this method to normalize dystrophin protein we

found that on average, *146aX* muscles showed a 2.4-fold increase in dystrophin protein per

"skipped" transcript (**Figure 5D**).

To support this data, we additionally performed quantification of dystrophin positive

fibers using immunofluorescence analysis. Analysis showed a greater dystrophin restoration via

an increase in dystrophin positive fibers in *146aX* muscles (43.5% increase p<0.05, **Figure 6A,**

A′, Figure S4). This difference persisted after normalizing to the extent of skipped *Dmd*

transcripts, showing a 3.5-fold increase in dystrophin positive fibers per "skipped" transcript

(**Figure 6B**). In our analysis we observed high variability in the "brightness" or intensity of

dystrophin positive fibers. To investigate this further we generated a visual standard to label the

intensity of dystrophin positive fibers as either "high" or "low" (**Figure 6C**) and then quantified

each group. Interestingly, we found that while there was a modest increase in the extent of

"high" intensity fibers in *146aX* muscle (35.6% increase, p<0.1), "low" intensity fibers were

60.52% higher in *146aX* muscles (p<0.05, **Figure 6D**). A potential interpretation of this

difference is that the absence of miR-146 stabilizes "skipped" dystrophin transcripts and more

protein continues to be produced from these skipped transcripts. Collectively, this data supports

the idea that reducing miR-146a can markedly improve exon-skipping-mediated dystrophin

restoration in dystrophin-deficient mice.

Improved muscle function in PMO-treated 146aX

We performed functional testing on 17-week-old, treated *mdx52* and *146aX* mice (1

week after second PMO injection). *146aX* mice showed a 24.4% increase in hindlimb grip

strength versus *mdx52* mice (p<0.05, **Figure 7A**), while forelimb grip strength showed similar

measurements for treated *mdx52* and *146aX* (**Figure S5A**). Suspension time in the two-limb

wire hang in treated *146aX* mice showed an increase (39.4% increase) but did not reach

statistical significance (p=0.163, **Figure 7A**).

Reduced pathology in treated 146aX muscles

Given the increases we saw in dystrophin rescue in *146aX*, most notably in the

diaphragm and triceps muscles, we investigated the therapeutic benefit of systemic PMO

treatment on these muscles by performing additional endpoint analysis on tissues. In *mdx* mice,

limb muscles undergo dramatic necrosis, which is followed by regeneration, increased

hypertrophy and increased muscle mass[35]. Analyzing tissue weights, the mass of both triceps

and diaphragm muscles showed reductions in *146aX* mice (**Figure 7B**), indicating reduced

pseudohypertrophy. The mass of gastrocnemius muscle was also significantly reduced, while

other muscles showed non-significant reductions in *146aX* versus *mdx52* genotypes (**Figure**

S5B). We observed no marked differences in endpoint body mass between treated *mdx52* and

146aX mice, however, triceps showed significant reduction in *146aX* mice when normalizing to

body weight, while other muscles showed non-significant reductions (**Figure S5C,D**).

Additionally, laminin staining showed visually less myofiber size variability in diaphragm and

triceps muscles (**Figure 7C**) and analysis of minimal Feret's diameter in laminin-stained muscle

confirmed this reduction (variance coefficient, **Figure 7C′**) consistent with reduced

asynchronous regeneration and a "healthier" muscle state in treated *146aX*[36]. These data were

corroborated with hematoxylin and eosin (H&E) analysis which showed visually healthier

muscle in treated *146aX* diaphragms muscles; quantification showed reduced inflammation and

necrosis in these muscles (**Figure S5E**). Additionally, immunofluorescence analyses of the

mouse macrophage marker F4/80 showed visually reduced macrophage infiltration in treated

146aX diaphragms (**Figure S5F**). We additionally analyzed the dystrophin associated protein, α-

sarcoglycan in treated diaphragms. Interestingly, we observed stronger α-sarcoglycan staining

in treated *146aX* diaphragms visible as increased yellow areas of α-sarcoglycan (green) and

laminin (red) co-localization, and this staining was especially pronounced in myofibers with

dystrophin positivity (**Figure 7D**)

Discussion

In the study presented here, we find miR-146a inhibits dystrophin protein levels *in vitro* and *in vivo* and is induced by inflammation in muscle. We also find that genetic deletion of miR-146a increases local dystrophin protein levels after intramuscular injection of an exon 51 skipping PMO in *mdx52* mice and increases body-wide dystrophin protein levels after systemic injection. Increased dystrophin rescue in systemically treated *146aX* mice also corresponds to increased muscle function, reduced histopathology and reduced myofiber size variability. To our knowledge, this is the first report showing that genetic deletion of a single miRNA can increase dystrophin protein in combination with administration of an exon skipping PMO.

We show here that PMO-treated *146aX* mice exhibit increased dystrophin protein levels compared to *mdx52*, while the extent of skipped *Dmd* transcript levels remains similar between groups. The observed increase in dystrophin protein but not dystrophin transcript levels is in line with our previous data in BMD biopsies and in BMD model mice showing that, while dystrophin transcript levels are the same, dystrophin protein levels are significantly reduced and negatively correlate with levels of miR-146a and other DTMs[7, 8], and that most often miRNAs inhibit translation but do not promote mRNA decay[17].

In our previous work, we identified and characterized 7 miRNAs that are elevated in BMD muscle biopsies, are inversely correlated with dystrophin protein levels, and found that these miRNAs target the dystrophin 3'UTR *in vitro*[7]. We collectively refer to this group of miRNAs as dystrophin targeting miRNAs or DTMs. This previous report also showed that treatment with the corticosteroid prednisone or the dissociative steroid vamorolone reduced expression of 3 DTMs including miR-146a in *mdx* mice[7]. In a separate report, we demonstrated that administration of

either prednisone or vamorolone to low-dystrophin expressing BMD model (*bmx*) mice reduces

expression of miR-146a and other DTMs (miR-31, miR-146b, miR-223) and increases dystrophin

levels by 50%[21]. The goal of the previous report was to determine if reducing expression of

multiple miRNAs could increase dystrophin protein levels. In the present study, our goal was to

determine if targeting a single dystrophin-targeting miRNA could improve dystrophin restoration

via exon skipping PMOs.

By using a genetic deletion model to ablate miR-146a in *mdx52* mice, this initial study

provides proof-of-principal that targeting miR-146a could potentially lead to increased dystrophin

rescue in DMD exon skipping. Upon systemic delivery of PMO, we found increased dystrophin

protein levels in dystrophic mice lacking miR-146a. Specifically, Wes immunoassay

quantification showed a 56.6% increase in dystrophin levels in *146aX* muscles. Additionally,

using dystrophin immunofluorescence, we show that extent of dystrophin positive fibers

increases from ~15 to 25% in 146aX muscles and that on average, deletion of miR-146a enable

~2.5 to 3.5-fold more dystrophin protein to be expressed from each "skipped transcript."

Together these data suggest that genetic deletion of miR-146a results in more efficient

dystrophin protein production. We observed more myofibers expressing a "low" level of

dystrophin via immunofluorescence. This could potentially be explained by the fact that mice

were sacrificed two weeks after the final PMO injection and that fibers with "low" dystrophin

were just starting to accumulate sufficient dystrophin protein from skipped transcripts.

Potentially, if mice have been sacrificed 4-6 weeks after the final PMO injection, it would have

allowed more time for dystrophin protein to accumulate from "skipped" transcripts which would

have yielded a greater difference between *mdx52* and *146aX* genotypes. Future studies will

determine how miR-146a deficiency affects the extent of *mdx52* muscle turnover, the

persistence of skipped transcripts and the overall accumulation of dystrophin protein after PMO

administration.

Our data show that deletion of miR-146a results in significant increases in dystrophin

protein as measured via Wes and immunofluorescence. The extent of dystrophin rescue

reported from DMD exon skipping clinical trials for Eteplirsen, golodirsen and casimersen is, on

average, about 0.9-1.9% of unaffected levels[10, 13] (NCT02500381). The target amount of

dystrophin restoration needed to show a functional benefit in DMD is still unclear. Some reports

using severe mouse models show 4% dystrophin is sufficient for improved muscle function and

survival [37-39]. Other reports show that 20% dystrophin improves symptoms[40] and 15% could be

sufficient to protect against contraction-induced injury in DMD model mice but more than 40%

dystrophin is needed to also improve muscle force[41]. Additionally, we previously reported that

even in BMD model (*bmx*) mice that express up to 30-50% dystrophin, muscles show

susceptibility to contraction-induced injury and have reduced function as measured by grip

strength and wire hang measures[8]. The most recent clinical data for the NS Pharma drug

Viltolarsen shows more promising preliminary data, with 5.9% dystrophin measured in muscle

biopsies[42]. Even with the increased efficacy of viltolarsen, improvement in the extent of

dystrophin restoration will be important to yield the full potential of exon skipping therapeutics in

DMD. Our study here suggests that targeting miR-146a could potentially contribute to that

overall goal. We should caveat this statement, however, by noting that our proof-of-concept

study here is preliminary and that additional pre-clinical studies to target miR-146a and to

measure potential side effects of miR-146a inhibition are required before determining the true

clinical translatability.

Cacchiarelli et. al[16] was the first to show that a miRNA that is significantly upregulated in

mdx muscles, miR-31, could target the dystrophin 3'UTR and inhibit dystrophin protein

production. Our lab later built on this data using BMD del45-47 biopsies and described seven

additional miRNAs that target dystrophin, including miR-146a. Following the initial miR-31

publication, Wells and Hildyard performed intraperitoneal injections of an exon 23 skipping vivo-

PMO alone or in combination with a miR-31 protector sequence (to prevent miR-31 binding to

the dystrophin 3'UTR) or a "scrambled protector" sequence[43]. While they observed increased

dystrophin in cell culture experiments by inhibiting or blocking miR-31, this result did not

translate *in vivo*[43]. There were, however, limitations to this study as the authors discuss.

Specifically, they postulated that delivery of the vivo-morpholinos overall was low, with almost

no delivery to hindlimbs, modest delivery to the diaphragm and less to the abdominal wall, and

therefore it wasn't clear whether the miR-31 target blocking strategy failed or the co-delivery of

the exon skipping PMO, and target blocker was too low to observe any measurable differences

in dystrophin protein. Based on their report and interpretation, here, we chose to use a genetic

approach to ablate expression of miR-146a so that the efficiency of delivery of an inhibitor or

blocking sequence was not a confounding factor in analysis. Moving forward we plan on

performing studies where we will conjugate a miR-146a inhibitor to a cell penetrating peptide or

antibody to determine how co-delivery of a miR-146a inhibitor or target blocking sequence with

an exon skipping PMO affects dystrophin protein restoration in dystrophic mice.

Elevated levels of miR-146a have been described in numerous other chronic inflammatory

and muscle disorders including myositis[44], Miyoshi myopathy[15], limb girdle muscular dystrophy[15],

FSHD[45], rheumatoid arthritis[46], Sjogren's syndrome[47], inflammatory bowel disease[19, 48, 49], and

asthma[50] suggesting it plays a role in diseases where inflammation persists. Previous studies

have reported that miR-146a is rapidly elevated in response to inflammatory stimuli, but then

"turns off" the NF-κB signaling pathway via binding to the *Traf6* and *Irak1* 3'UTRs[18, 51, 52].

Because of this, it is generally regarded as "anti-inflammatory."   However, more recent work

has described a mechanism where exogenous miR-146a-5p, but not its duplex precursor, is

"pro-inflammatory" and induces inflammation via activation of TLR7 while it's duplex precursor

functions as part of a negative feedback loop to "turn off" NF-κB[53]. The work of Wang et. al

described above therefore suggest that the narrow definition of a miRNA such as miR-146a as

either anti- or pro-inflammatory is perhaps too reductive and more work is required to fully

delineate the comprehensive role of each microRNA in both a healthy and a disease state.

It should be noted that another group also generated *mdx:miR-146a-/-* mice[54], however,

the overall aim of that work was significantly different from the work we present here. The

previous study sought to test the hypothesis that miR-146a is anti-inflammatory and that genetic

deletion of miR-146a in *mdx* would worsen the natural etiology of disease; after analysis they

reported no signficant differences between the *mdx* mice with or without miR-146a expression.

The data we present here demonstrates that miR-146a deletion benefits dystrophin rescue in

mdx52 mice treated with an exon 51 skipping PMO.

Interestingly, we found elevated miR-146a in murine myositis muscle and human muscle

biopsies of inclusion body myositis[44]. In these same muscles, we observed reduced dystrophin

staining via Western blot as well as reduced and discontinuous dystrophin staining[44].

Additionally, another group recently reported reduced dystrophin and increased dystrophin-

targeting microRNAs (miR-146a, miR-223) in immune-mediated necrotizing myopathy[55]. Thus, it

is possible that DTMs such as miR-146a contribute to a secondary dystrophin deficiency that

would impair overall muscle function and the therapeutic targeting of this inflammation-regulated

miRNA could be beneficial in the context of other muscle disorders.

One potential way to improve DMD exon skipping outcomes is to harness miRNA

therapeutics as a combination therapy with the goal of inhibiting miR-146a-5p levels to enable

increased dystrophin protein production from skipped dystrophin transcripts. This strategy could

also potentially be applied as a stand-alone therapy in BMD where lower-than-normal levels of

dystrophin are observed in muscle. There are a few different ways in which miR-146a inhibition

or reduction can be achieved. These include the use of 1) a miR-146a inhibitor (anti-miR), 2) an

oligonucleotide that prevents miR-146a from binding to the dystrophin 3′UTR (target blockers or

blockmiRs), or 3) a small molecule inhibitor. Currently, there are several anti-miR therapeutics in

clinical trials that show promising preliminary results (reviewed in[56]) including CDR132L, a

locked nucleic acid (LNA) targeting miR-132 in heart failure[57], Miravirsen/SPC3649, an LNA

targeting miR-122 in patients with chronic hepatitis C virus[58], and MRG-110, an LNA targeting

miR-92a to benefit wound repair failure[59]. Additionally myotonic dystrophy (DM1) preclinical

studies, using either a miR-23b antagomiR[60] or a peptide-conjugated blockmiR to prevent miR-

23 binding to MBNL[61], have shown these strategies improve muscle function in mouse models.

We have also observed reduced levels of miR-146a through treatment with anti-inflammatory

small molecules, either vamorolone or prednisone, in *mdx* mice[7, 20]. A recent report from our lab

shows that while both prednisone and vamorolone reduce miR-146a and increase dystrophin in

the gastrocnemius muscle of *bmx* mice, only vamorolone significantly increases dystrophin

levels in the heart[21]. While corticosteroids will likely always be given as a combination therapy

along with exon skipping PMOs or other dystrophin restoration therapies, pre-clinical studies

that investigate which corticosteroids (prednisone vs. deflazacort vs. vamorolone) yield the

greatest increases in dystrophin restoration will be informative to current and future DMD

treatment regimens.

In conclusion, the data presented here shows that genetic deletion of miR-146a in

mdx52 mice increases levels of therapeutic dystrophin protein restoration without affecting the

extent of skipped dystrophin transcript levels. These data provide a strong rationale for

investigating how therapeutic targeting of miR-146a and other DTMs may impact dystrophin

protein levels in DMD exon skipping, as well as in other muscle diseases more broadly.

Figures and Figure Legends

Figure 1. miR-146a is significantly elevated in BMD and DMD patients and model mice.

(**A**) The sequence of miR-146a-5p is conserved across species. Note that in mammals there is

100% conservation. (**B**) Left; miR-146a levels are elevated in BMD patients with low dystrophin

(<20% dystrophin of unaffected individuals). n=5 normal, 6 "BMD high" and 4 "BMD low";

ANOVA, **p<0.001, adapted from [7]. Right; Dystrophin protein and miR-146a levels are inversely

correlated in BMD patients. Spearman correlation, r=-0.6365, **p<0.01. Adapted from[7]. (**C**) miR-

146a levels are significantly elevated in both BMD and DMD patients (taken from raw data in

Eisenberg et. al[15]) n=9 unaffected (UA), 6 BMD and 8 DMD biopsies; ANOVA, #p<0.1, *p<0.05.

(**D**) miR-146a levels are significantly increased in *mdx52* mouse diaphragm (D), tibialis anterior

(TA), quadriceps (Q), gastrocnemius (G), and triceps (Tri). n=6/group; Two-way ANOVA,

p<0.01, *p<0.05. (E**) miR-146a levels in *bmx* mouse diaphragm (D), tibialis anterior (TA),

quadriceps (Q), gastrocnemius (G), and triceps (Tri). miR-146a is significantly increased in the

latter 3 muscles. n=6/group, Two-way ANOVA, *p<0.05. (**F**) miR-146a is more highly expressed

and induced in mdx vs. WT H2K myotubes treated with 10 ng/mL TNF-α (n=4 replicates per

group) Student's t-test, ****p<0.0001. Data are represented as mean ± S.E.M. See also Figure

S1.

Figure 2. miR-146a targets the dystrophin 3′UTR and reduces exon skipping-mediated dystrophin restoration. (**A-B**) Predicted miR-146a-5p binding sites in the human *DMD* 3'UTR. (**A**) Scan-miR was used to search for putative miR-146a-5p binding sites within the human dystrophin 3′UTR sequence. The y axis denotes the dissociation constant (-log Kd), and the x axis denotes the location on the dystrophin 3′UTR in the 5′ to 3′ direction. (**B**) List of 17 putative miR-146a binding sites in the *DMD* 3' UTR; orange denotes the strongest binding site and the one that was mutated in the luciferase construct in (**C**). The predicted binding sites include: a 7mer-m8 where there is perfect complementarity of the seed sequence with the UTR and this

complementarity extends to nucleotide 8; a 6mer-A1 where there is seed sequence

complementarity at nucleotide positions 3-7 and additional complementarity at nucleotide 8; a

6mer-A1, where the seed sequence shows complementarity to the UTR at positions 2-6 and

additionally possesses an adenosine (A) at position 1; an wobbled 8mer, where there is an

adenosine position 1, seed sequence complementarity at position 2-7 that is interrupted by a

single mismatch and additional complementarity at position 8 and several non-canonical binding

sites [30]. (**C**) Schematic showing the binding site of miR-146a on the *DMD* 3' UTR (top) and the

base pairing of miR-146a with the *DMD* 3' UTR when the binding site is mutated (bottom). (**D**) A

luciferase reporter assay was used to examine miR-146a binding to the WT and mutant *DMD* 3'

UTR in HEK-293T cells. Transfection with miR-146a significantly reduces reporter activity in

cells with WT *DMD* 3' UTR whereas there is no reduction in reporter activity in cells with the

mutated *DMD* 3' UTR. n=3-4, ANOVA, *p<0.05. (**E**) The TAs of *mdx52* mice were injected with

2µg of exon skipping PMO and 10µg of either control or miR-146a. TA cross sections were

stained for dystrophin (red) and wheat germ agglutinin (white). (**E'**) Quantification of dystrophin

positive fibers; significantly fewer dystrophin-positive myofibers were observed in muscles co-

injected with miR-146a compared to control. n=9-10, Student's t-test, *p<0.05. Data represented

as mean ± S.E.M. See also Figure S2.

Figure 3. Generation of dmd$^{-/-}$; 146a$^{-/-}$ (*146aX* mice). (**A**) Breeding schematic to develop

146aX mice. (**B**) qRT-PCR showing miR-146a levels in the gastrocnemius muscles of WT, *mdx*,

and *146aX* mice. *146aX* mice have no detectable levels of miR-146a via qRT-PCR. n=6/group,

ANOVA, ****p<0.0001. (**C**)Left; Representative dystrophin immunofluorescence (red) and

laminin staining in *wild-type (WT), mdx52* and *146aX* quadriceps showing strong dystrophin

staining in WT muscle and no staining except for a few revertant fibers in *mdx52* and *146aX*

muscles (white boxes with asterisks). Right; Quantification of revertant fibers in *mdx52* and

146aX muscles shows that they represent only a small fraction of all fibers. There is no

significant difference in revertant dystrophin-positive fibers between genotypes. n=6-7/group,

students t-test, ns p=0.6830.

Figure 4. Deletion of miR-146a significantly increases dystrophin restoration in *mdx52*

mice treated with intramuscular PMO. (**A**) Schematic of intramuscular PMO injection

experimental design. *mdx* and *146aX* tibialis anterior muscles (TAs) were injected with 2µg of

an exon 51 skipping PMO. 2 weeks post-injection muscles were harvested for analysis. n=10

muscles/group and 5 mice per group. Wes = Western capillary immunoassay, IF =

immunofluorescence. (**B**) qRT-PCR was performed to quantify % exon skipping in PMO-injected

mdx52 and *146aX* TAs as previously described[34]. Exon skipping efficiency at the RNA level is

not significantly between *mdx52* and *146aX* TAs. (**C**) Capillary-based immunoassay (Wes) was

used to quantify levels of dystrophin protein. (**C'**) Dystrophin protein is significantly increased in

146aX mice injected with PMO compared to *mdx* mice. (**D**) Immunofluorescence showing

dystrophin staining in the TA of mdx and *146aX* mice with intramuscular PMO. (i) shows a

zoomed in area of high dystrophin rescue and (ii) shows a zoomed in area of low dystrophin

rescue. Bars = 500μm and 50 μm respectively.   (**D'**) Quantification of dystrophin-positive

myofibers. The percentage of dystrophin-positive myofibers following PMO injection is

significantly increased in *146aX* mice compared to mdx mice. Images were blinded during

quantification. Data represented as mean ± S.E.M. ns $p>0.05$, *$p≤0.05$, **$p≤0.01$, ****$p≤0.0001$.

See also, Figure S3.

Figure 5. Deletion of miR-146a significantly increases dystrophin protein in *mdx52* mice

treated with systemic PMO. (**A**) Schematic of experimental design for systemic PMO

injections in *mdx* and *146aX* mice. *mdx52* and *146aX* mice were administered systemic PMO

via the retroorbital sinus (400 mg/kg, exon 51 skipping PMO) at 15 weeks of age. One week

later a second 400mg/kg PMO injection was performed. Muscle function tests were performed

one week after the second injection and mice were sacrificed 2 weeks after the second

injection. Muscles were analyzed by qRT-PCR to determine extent of skipped *Dmd* mRNA and

by Western capillary immunoassay (Wes) and immunofluorescence (IF) to determine dystrophin

protein levels (IF). n=4 mice and 5 muscles per group. (**B**) Exon skipping efficiency at the RNA

level via systemic injection of PMO in *mdx* and *146aX* mice is not significantly different.

Student's t-test. (**C**) Capillary-based immunoassay (Wes) was used to quantify dystrophin

protein levels in skeletal muscle of *mdx* and *146aX* mice systemically treated with exon-skipping

PMO via retroorbital injection. Depicted is a virtual Wes blot . (**C'**) Wes quantification of

dystrophin protein levels in skeletal muscle. Dystrophin protein is significantly increased in

748   *146aX* mice versus *mdx52* mice. Student's t-test, *p<0.05. Note: % Dystrophin was calculated

749   by normalizing to the average *mdx* intensity for each muscle and setting it to 100%. (**D**)

750   Dystrophin protein levels from Wes normalized to % *Dmd* exon skipped transcripts. There is a

751   higher ratio of dystrophin restoration in *146aX* mice compared to *mdx* mice. Data represented

752   as mean ± S.E.M. Student's t-test, *p<0.05.

769
Figure 6. Deletion of miR-146a significantly increases dystrophin positive fibers in *mdx52*

mice. (**A**) Dystrophin immunofluorescence (red) in PMO-injected *mdx* and *146aX* diaphragm

and triceps muscles. Bar= 200mM and 100 mM, respectively. Muscles were counterstained with

DAPI (nuclei) and laminin (green, not shown) to count total muscle fibers). (**A'**) Quantification of

dystrophin-positive myofibers, (n=4 mice and 5 muscles per group). The percentage of

dystrophin-positive myofibers following PMO injection is significantly increased in *146aX* mice

compared to mdx mice. (**B**)The percentage of dystrophin positive fibers normalized to the

percentage of *Dmd* exon skipped transcripts. There is a higher ratio of dystrophin restoration in

146aX mice compared to *mdx* mice. Data represented as mean ± S.E.M Student's t-test,

*p<0.05. (**C**) Visual standard for classification of dystrophin high, low, and negative fibers. (**D**)

Left; Quantification of "high" intensity dystrophin-positive fibers. Right; quantification of "low"

intensity dystrophin positive fibers. Images were blinded during quantification. Student's t-test,

#p<0.1, *p<0.05. All data represented as mean ± S.E.M. Dia, diaphragm; TA, tibialis anterior;

Quad, quadriceps; Gastroc, gastrocnemius; Tri, triceps. See also Figure S4.

Figure 7. Functional and histological analysis of PMO-treated *mdx52* and *146aX* mice. (**A**)

Muscle function tests in *mdx52* and *146aX* mice. Left; Hindlimb grip strength was performed and

normalized to body weight. Right; Wire hang-time assay. n=4 mice per group; Student's t-test.

(**B**) Terminal tissue masses of triceps (n=8) and diaphragms (n=4). Student's t-test, #$p<0.1$,

*$p<0.05$. All data represented as mean ± S.E.M.(**C**) Immunofluorescence of muscle sections

using laminin (green), dystrophin (red) and DAPI (blue) staining of treated *mdx52* versus *146aX*

muscles to visualize both dystrophin restoration and myofiber size. Bar=100μm. (**C'**)

Quantification of myofiber size variability via the variance coefficient of minimal Feret's diameter.

(**D**) To visualize how dystrophin restoration affects localization of a dystrophin-associated

protein (α-sarcoglycan), laminin (red), a-sarcoglycan (green), and dystrophin (blue) were

visualized via immunofluorescence in *mdx52* and *146aX* diaphragms. Note that α-sarcoglycan is

brighter in treated *146aX* diaphragms which is apparent by the yellow laminin/a-sarcoglycan co-

localization (merge) image versus the more orange hue in the laminin/α-sarcoglycan image in

treated *mdx52* diaphragms. Bar = 50μm. See also Figure S5.

Conflict of Interest

AAF has an issued patent on intellectual property relating to the manuscript (United States Patent #10266824).

Author Contributions

NM contributed to experiments, data analysis, manuscript writing and interpretation; KC performed data analysis; CS performed F4/80 immunofluorescence, CT performed experiments and analysis, CH contributed to experimental design, writing and interpretation, AF contributed to experimental design, experiments, analysis, and manuscript writing.

Funding

CRH receives support from the Foundation to Eradicate Duchenne and the NIH (R01HL153054, R00HL130035, K99HL130035, L40AR068727). AAF receives support from the Foundation to Eradicate Duchenne, the Department of Defense (W81XWH-17-1-0475), and the NIH (L40AR070539 and L40NS127380).

Acknowledgements

We would like to thank Dr. Shin'ichi Takeda for the generous gift of the *mdx52* mice.

Keywords

 antisense oligonucleotides, RNA therapeutics, Duchenne Muscular Dystrophy (DMD), Becker Muscular Dystrophy (BMD), dystrophin, exon skipping, microRNA, miR-146a-5p

References

1. Hoffman, EP, Brown, RH, Jr., and Kunkel, LM (1987). Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* **51**: 919-928.
2. Beggs, AH, Hoffman, EP, Snyder, JR, Arahata, K, Specht, L, Shapiro, F*, et al.* (1991). Exploring the molecular basis for variability among patients with Becker muscular dystrophy: dystrophin gene and protein studies. *Am J Hum Genet* **49**: 54-67.
3. Hoffman, EP, Kunkel, LM, Angelini, C, Clarke, A, Johnson, M, and Harris, JB (1989). Improved diagnosis of Becker muscular dystrophy by dystrophin testing. *Neurology* **39**: 1011-1017.
4. Kesari, A, Pirra, LN, Bremadesam, L, McIntyre, O, Gordon, E, Dubrovsky, AL*, et al.* (2008). Integrated DNA, cDNA, and protein studies in Becker muscular dystrophy show high exception to the reading frame rule. *Hum Mutat* **29**: 728-737.
5. van den Bergen, JC, Wokke, BH, Janson, AA, van Duinen, SG, Hulsker, MA, Ginjaar, HB*, et al.* (2014). Dystrophin levels and clinical severity in Becker muscular dystrophy patients. *J Neurol Neurosurg Psychiatry* **85**: 747-753.
6. Barp, A, Bello, L, Caumo, L, Campadello, P, Semplicini, C, Lazzarotto, A*, et al.* (2017). Muscle MRI and functional outcome measures in Becker muscular dystrophy. *Sci Rep* **7**: 16060.
7. Fiorillo, AA, Heier, CR, Novak, JS, Tully, CB, Brown, KJ, Uaesoontrachoon, K*, et al.* (2015). TNF-alpha-Induced microRNAs Control Dystrophin Expression in Becker Muscular Dystrophy. *Cell Rep* **12**: 1678-1690.
8. Heier, CR, McCormack, NM, Tully, CB, Novak, JS, Newell-Stamper, BL, Russell, AJ*, et al.* (2023). The X-linked Becker muscular dystrophy (bmx) mouse models Becker muscular dystrophy via deletion of murine dystrophin exons 45-47. *J Cachexia Sarcopenia Muscle*.
9. Cirak, S, Arechavala-Gomeza, V, Guglieri, M, Feng, L, Torelli, S, Anthony, K*, et al.* (2011). Exon skipping and dystrophin restoration in patients with Duchenne muscular dystrophy after systemic phosphorodiamidate morpholino oligomer treatment: an open-label, phase 2, dose-escalation study. *Lancet* **378**: 595-605.
10. Frank, DE, Schnell, FJ, Akana, C, El-Husayni, SH, Desjardins, CA, Morgan, J*, et al.* (2020). Increased dystrophin production with golodirsen in patients with Duchenne muscular dystrophy. *Neurology* **94**: e2270-e2282.
11. Wagner, KR, Kuntz, NL, Koenig, E, East, L, Upadhyay, S, Han, B*, et al.* (2021). Safety, tolerability, and pharmacokinetics of casimersen in patients with Duchenne muscular dystrophy amenable to exon 45 skipping: A randomized, double-blind, placebo-controlled, dose-titration trial. *Muscle Nerve* **64**: 285-292.
12. Clemens, PR, Rao, VK, Connolly, AM, Harper, AD, Mah, JK, McDonald, CM*, et al.* (2023). Efficacy and Safety of Viltolarsen in Boys With Duchenne Muscular Dystrophy: Results From the Phase 2, Open-Label, 4-Year Extension Study. *J Neuromuscul Dis*.
13. . Study of Eteplirsen in DMD Patients (PROMOVI).
14. Servais, L, Mercuri, E, Straub, V, Guglieri, M, Seferian, AM, Scoto, M*, et al.* (2022). Long-Term Safety and Efficacy Data of Golodirsen in Ambulatory Patients with Duchenne Muscular Dystrophy Amenable to Exon 53 Skipping: A First-in-human, Multicenter, Two-Part, Open-Label, Phase 1/2 Trial. *Nucleic Acid Ther* **32**: 29-39.
15. Eisenberg, I, Eran, A, Nishino, I, Moggio, M, Lamperti, C, Amato, AA*, et al.* (2007). Distinctive patterns of microRNA expression in primary muscular disorders. *Proc Natl Acad Sci U S A* **104**: 17016-17021.
16. Cacchiarelli, D, Incitti, T, Martone, J, Cesana, M, Cazzella, V, Santini, T*, et al.* (2011). miR-31 modulates dystrophin expression: new implications for Duchenne muscular dystrophy therapy. *EMBO Rep* **12**: 136-141.

17. Humphreys, DT, Westman, BJ, Martin, DI, and Preiss, T (2005). MicroRNAs control
translation initiation by inhibiting eukaryotic initiation factor 4E/cap and poly(A) tail
function. *Proc Natl Acad Sci U S A* **102**: 16961-16966.
18. Taganov, KD, Boldin, MP, Chang, KJ, and Baltimore, D (2006). NF-kappaB-dependent
induction of microRNA miR-146, an inhibitor targeted to signaling proteins of innate
immune responses. *Proc Natl Acad Sci U S A* **103**: 12481-12486.
19. Heier, CR, Fiorillo, AA, Chaisson, E, Gordish-Dressman, H, Hathout, Y, Damsker, JM*, et*
al. (2016). Identification of Pathway-Specific Serum Biomarkers of Response to
Glucocorticoid and Infliximab Treatment in Children with Inflammatory Bowel Disease.
Clin Transl Gastroenterol **7**: e192.
20. Fiorillo, AA, Tully, CB, Damsker, JM, Nagaraju, K, Hoffman, EP, and Heier, CR (2018).
Muscle miRNAome shows suppression of chronic inflammatory miRNAs with both
prednisone and vamorolone. *Physiol Genomics* **50**: 735-745.
21. McCormack, NM, Nguyen, NY, Tully, CB, Oliver, T, Fiorillo, AA, and Heier, CR (2023).
Vamorolone improves Becker muscular dystrophy and increases dystrophin protein in
bmx model mice. *iScience* **26**: 107161 %U
https://www.sciencedirect.com/science/article/pii/S2589004223012385.
22. Araki, E, Nakamura, K, Nakao, K, Kameya, S, Kobayashi, O, Nonaka, I*, et al.* (1997).
Targeted disruption of exon 52 in the mouse dystrophin gene induced muscle
degeneration similar to that observed in Duchenne muscular dystrophy. *Biochem*
Biophys Res Commun **238**: 492-497.
23. Lu, LF, Boldin, MP, Chaudhry, A, Lin, LL, Taganov, KD, Hanada, T*, et al.* (2010).
Function of miR-146a in controlling Treg cell-mediated regulation of Th1 responses. *Cell*
142: 914-929.
24. Mann, CJ, Honeyman, K, Cheng, AJ, Ly, T, Lloyd, F, Fletcher, S*, et al.* (2001).
Antisense-induced exon skipping and synthesis of dystrophin in the mdx mouse. *Proc*
Natl Acad Sci U S A **98**: 42-47.
25. Aoki, Y, Nakamura, A, Yokota, T, Saito, T, Okazawa, H, Nagata, T*, et al.* (2010). In-
frame dystrophin following exon 51-skipping improves muscle pathology and function in
the exon 52-deficient mdx mouse. *Mol Ther* **18**: 1995-2005.
26. Beekman, C, Janson, AA, Baghat, A, van Deutekom, JC, and Datson, NA (2018). Use of
capillary Western immunoassay (Wes) for quantification of dystrophin levels in skeletal
muscle of healthy controls and individuals with Becker and Duchenne muscular
dystrophy. *PLoS One* **13**: e0195850.
27. Vandesompele, J, De Preter, K, Pattyn, F, Poppe, B, Van Roy, N, De Paepe, A*, et al.*
(2002). Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. *Genome Biol* **3**: RESEARCH0034.
28. Wein, N, Vetter, TA, Vulin, A, Simmons, TR, Frair, EC, Bradley, AJ*, et al.* (2022).
Systemic delivery of an AAV9 exon-skipping vector significantly improves or prevents
features of Duchenne muscular dystrophy in the Dup2 mouse. *Mol Ther Methods Clin*
Dev **26**: 279-293.
29. Soutschek, M, Gross, F, Schratt, G, and Germain, PL (2022). scanMiR: a biochemically
based toolkit for versatile and efficient microRNA target prediction. *Bioinformatics* **38**:
2466-2473.
30. Grimson, A, Farh, KK, Johnston, WK, Garrett-Engele, P, Lim, LP, and Bartel, DP (2007).
MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol Cell*
27: 91-105.
31. Lu, QL, Morris, GE, Wilton, SD, Ly, T, Artem'yeva, OV, Strong, P*, et al.* (2000). Massive
idiosyncratic exon skipping corrects the nonsense mutation in dystrophic mouse muscle
and produces functional revertant fibers by clonal expansion. *J Cell Biol* **148**: 985-996.

32.   Echigoya, Y, Lee, J, Rodrigues, M, Nagata, T, Tanihata, J, Nozohourmehrabad, A*, et al.*
946   (2013). Mutation types and aging differently affect revertant fiber expansion in dystrophic
947   mdx and mdx52 mice. *PLoS One* **8**: e69194.
948   33.   Vila, MC, Klimek, MB, Novak, JS, Rayavarapu, S, Uaesoontrachoon, K, Boehler, JF*, et
949   al.* (2015). Elusive sources of variability of dystrophin rescue by exon skipping. *Skelet
950   Muscle* **5**: 44.
951   34.   Novak, JS, Spathis, R, Dang, UJ, Fiorillo, AA, Hindupur, R, Tully, CB*, et al.* (2021).
952   Interrogation of Dystrophin and Dystroglycan Complex Protein Turnover After Exon
953   Skipping Therapy. *J Neuromuscul Dis* **8**: S383-S402.
954   35.   Coulton, GR, Morgan, JE, Partridge, TA, and Sloper, JC (1988). The mdx mouse
955   skeletal muscle myopathy: I. A histological, morphometric and biochemical investigation.
956   *Neuropathol Appl Neurobiol* **14**: 53-70.
957   36.   Dadgar, S, Wang, Z, Johnston, H, Kesari, A, Nagaraju, K, Chen, YW*, et al.* (2014).
958   Asynchronous remodeling is a driver of failed regeneration in Duchenne muscular
959   dystrophy. *J Cell Biol* **207**: 139-158.
960   37.   Li, D, Yue, Y, and Duan, D (2008). Preservation of muscle force in Mdx3cv mice
961   correlates with low-level expression of a near full-length dystrophin protein. *Am J Pathol*
962   **172**: 1332-1341.
963   38.   Li, D, Yue, Y, and Duan, D (2010). Marginal level dystrophin expression improves
964   clinical outcome in a strain of dystrophin/utrophin double knockout mice. *PLoS One* **5**:
965   e15286.
966   39.   van Putten, M, Hulsker, M, Nadarajah, VD, van Heiningen, SH, van Huizen, E, van
967   Iterson, M*, et al.* (2012). The effects of low levels of dystrophin on mouse muscle
968   function and pathology. *PLoS One* **7**: e31937.
969   40.   Phelps, SF, Hauser, MA, Cole, NM, Rafael, JA, Hinkle, RT, Faulkner, JA*, et al.* (1995).
970   Expression of full-length and truncated dystrophin mini-genes in transgenic mdx mice.
971   *Hum Mol Genet* **4**: 1251-1258.
972   41.   Godfrey, C, Muses, S, McClorey, G, Wells, KE, Coursindel, T, Terry, RL*, et al.* (2015).
973   How much dystrophin is enough: the physiological consequences of different levels of
974   dystrophin in the mdx mouse. *Hum Mol Genet* **24**: 4225-4237.
975   42.   Clemens, PR, Rao, VK, Connolly, AM, Harper, AD, Mah, JK, Smith, EC*, et al.* (2020).
976   Safety, Tolerability, and Efficacy of Viltolarsen in Boys With Duchenne Muscular
977   Dystrophy Amenable to Exon 53 Skipping: A Phase 2 Randomized Clinical Trial. *JAMA
978   Neurol* **77**: 982-991.
979   43.   Hildyard, JC, and Wells, DJ (2016). Investigating Synthetic Oligonucleotide Targeting of
980   Mir31 in Duchenne Muscular Dystrophy. *PLoS Curr* **8**.
981   44.   Kinder, TB, Heier, CR, Tully, CB, Van der Muelen, JH, Hoffman, EP, Nagaraju, K*, et al.*
982   (2020). Muscle Weakness in Myositis: MicroRNA-Mediated Dystrophin Reduction in a
983   Myositis Mouse Model and Human Muscle Biopsies. *Arthritis Rheumatol* **72**: 1170-1183.
984   45.   Nunes, AM, Ramirez, M, Jones, TI, and Jones, PL (2021). Identification of candidate
985   miRNA biomarkers for facioscapulohumeral muscular dystrophy using DUX4-based
986   mouse models. *Dis Model Mech* **14**.
987   46.   Moran-Moguel, MC, Petarra-Del Rio, S, Mayorquin-Galvan, EE, and Zavala-Cerna, MG
988   (2018). Rheumatoid Arthritis and miRNAs: A Critical Review through a Functional View.
989   *J Immunol Res* **2018**: 2474529.
990   47.   Reale, M, D'Angelo, C, Costantini, E, Laus, M, Moretti, A, and Croce, A (2018).
991   MicroRNA in Sjogren's Syndrome: Their Potential Roles in Pathogenesis and Diagnosis.
992   *J Immunol Res* **2018**: 7510174.
993   48.   Chapman, CG, and Pekow, J (2015). The emerging role of miRNAs in inflammatory
994   bowel disease: a review. *Therap Adv Gastroenterol* **8**: 4-22.

49. Batra, SK, Heier, CR, Diaz-Calderon, L, Tully, CB, Fiorillo, AA, van den Anker, J, *et al.* (2020). Serum miRNAs Are Pharmacodynamic Biomarkers Associated With Therapeutic Response in Pediatric Inflammatory Bowel Disease. *Inflamm Bowel Dis* **26**: 1597-1606.

50. Elnady, HG, Sherif, LS, Kholoussi, NM, Ali Azzam, M, Foda, AR, Helwa, I, *et al.* (2020). Aberrant Expression of Immune-related MicroRNAs in Pediatric Patients with Asthma. *Int J Mol Cell Med* **9**: 246-255.

51. Boldin, MP, Taganov, KD, Rao, DS, Yang, L, Zhao, JL, Kalwani, M, *et al.* (2011). miR-146a is a significant brake on autoimmunity, myeloproliferation, and cancer in mice. *J Exp Med* **208**: 1189-1201.

52. Nahid, MA, Pauley, KM, Satoh, M, and Chan, EK (2009). miR-146a is critical for endotoxin-induced tolerance: IMPLICATION IN INNATE IMMUNITY. *J Biol Chem* **284**: 34590-34599.

53. Wang, S, Yang, Y, Suen, A, Zhu, J, Williams, B, Hu, J, *et al.* (2021). Role of extracellular microRNA-146a-5p in host innate immunity and bacterial sepsis. *iScience* **24**: 103441.

54. Bronisz-Budzynska, I, Chwalenia, K, Mucha, O, Podkalicka, P, Karolina Bukowska, S, Jozkowicz, A, *et al.* (2019). miR-146a deficiency does not aggravate muscular dystrophy in mdx mice. *Skelet Muscle* **9**: 22.

55. Marmen, MB, Orfi, Z, Dort, J, Proulx-Gauthier, JP, Chrestian, N, Dumont, NA, *et al.* (2023). Decreased dystrophin expression and elevated dystrophin-targeting miRNAs in anti-HMGCR immune-mediated necrotizing myopathy. *Acta Neuropathol*.

56. Diener, C, Keller, A, and Meese, E (2022). Emerging concepts of miRNA therapeutics: from cells to clinic. *Trends Genet* **38**: 613-626.

57. Taubel, J, Hauke, W, Rump, S, Viereck, J, Batkai, S, Poetzsch, J, *et al.* (2021). Novel antisense therapy targeting microRNA-132 in patients with heart failure: results of a first-in-human Phase 1b randomized, double-blind, placebo-controlled study. *Eur Heart J* **42**: 178-188.

58. van der Ree, MH, van der Meer, AJ, van Nuenen, AC, de Bruijne, J, Ottosen, S, Janssen, HL, *et al.* (2016). Miravirsen dosing in chronic hepatitis C patients results in decreased microRNA-122 levels without affecting other microRNAs in plasma. *Aliment Pharmacol Ther* **43**: 102-113.

59. Abplanalp, WT, Fischer, A, John, D, Zeiher, AM, Gosgnach, W, Darville, H, *et al.* (2020). Efficiency and Target Derepression of Anti-miR-92a: Results of a First in Human Study. *Nucleic Acid Ther* **30**: 335-345.

60. Cerro-Herreros, E, Gonzalez-Martinez, I, Moreno-Cervera, N, Overby, S, Perez-Alonso, M, Llamusi, B, *et al.* (2020). Therapeutic Potential of AntagomiR-23b for Treating Myotonic Dystrophy. *Mol Ther Nucleic Acids* **21**: 837-849.

61. Overby, SJ, Cerro-Herreros, E, Gonzalez-Martinez, I, Varela, MA, Seoane-Miraz, D, Jad, Y, *et al.* (2022). Proof of concept of peptide-linked blockmiR-induced MBNL functional rescue in myotonic dystrophy type 1 mouse model. *Mol Ther Nucleic Acids* **27**: 1146-1155.