1

5 **Serine arginine-rich splicing factor (SRSF7) cooperates with the histone methyltransferase**

6 **KMT5a to promote the type I interferon response via transcriptional activation of IRF7**

8 Haley M. Scott[1], Mackenzie H. Smith[1], Aja K. Coleman[1], Summer L. Apostalo[1], Allison R. Wag-

9 ner[1], Robert O. Watson[1], Kristin L. Patrick[1]*#

16 [1]Department of Microbial Pathogenesis and Immunology, Texas A&M Health, School of Medicine

23 *Corresponding author: kpatrick03@tamu.edu

24 #Lead contact

25 Phone: (979) 436-0354

26 Twitter: @The_PW_Lab

SUMMARY

Tight regulation of macrophage immune gene expression is required to fight infection without risking harmful inflammation. The contribution of RNA binding proteins (RBPs) to shaping the macrophage response to pathogens remains poorly understood. Transcriptomic analysis revealed that a member of the serine/arginine-rich (SR) family of mRNA processing factors, SRSF7, is required for optimal expression of a cohort of interferon stimulated genes (ISGs) in macrophages. Using genetic and biochemical assays, we discovered that in addition to its canonical role in regulating alternative splicing, SRSF7 drives transcription of interferon regulatory transcription factor 7 (IRF7) to promote antiviral immunity. At the *Irf7* promoter, SRSF7 maximizes STAT1 transcription factor binding and RNA polymerase II elongation via cooperation with the H4K20me1 histone methyltransferase KMT5a (SET8). These studies define an unorthodox role for an SR protein in activating transcription and reveal an unappreciated RNA binding protein-chromatin network that orchestrates macrophage antiviral gene expression.

INTRODUCTION

Macrophages (MΦ) are innate immune cells that serve as the body's first line of defense against pathogens. They express a panoply of pattern recognition receptors that detect pathogen- and damage-associated molecular patterns (PAMPs and DAMPs) and transmit these "danger signals" to the nucleus though complex signal transduction cascades. Generally, these pathogen sensing cascades terminate with phosphorylation and nuclear translocation of specialized immune transcription factors (e.g. NFκB, IRF3, IRF7). Because innate immune transcription factors exert control over hundreds of genes with massive inflammatory potential, their misregulation can have dire consequences for a host: innate responses that are too weak enable pathogen survival and replication, whereas responses that are too strong can trigger a cytokine storm, chronic inflammation, and autoimmunity. While the mechanisms through which innate immune transcription factors are regulated post-translationally—via phosphorylation, dimerization, and nuclear translocation—are well-established (Caamano and Hunter, 2002; Jefferies, 2019; Ning et al., 2011; Smale, 2012), how cells control expression of these master regulators at the transcriptional and post-transcriptional levels remains less well-understood.

A growing literature has built a strong case for RNA binding proteins (RBPs) in shaping the transcriptomes of specialized cells like macrophages. SR (serine-arginine rich) proteins are a 15-member family of RBPs that are broadly defined as regulators of alternative splicing. By virtue of distinct RNA recognition motifs, SRs bind to unique exonic splicing enhancers to help direct the spliceosome to cis-splicing signals in target transcripts (Wagner and Frye, 2021). While best known for their role in pre-mRNA splicing, SRs can participate in virtually any step of RNA processing, from transcription all the way to translation (Jeong, 2017). These studies of SR function have mostly been carried out in genetically tractable "model" cell lines (e.g. K562, HEK293T, and HepG2), but we are beginning to appreciate specialized roles for SR proteins in immune cells. SRSF1 is required for regulatory T cell homeostasis and function (Katsuyama and Moulton, 2021) and several SR proteins have been implicated in T cell development via controlling alternative splicing of CD45 (Lemaire et al., 1999; ten Dam et al., 2000; Wang et al., 2001). We recently implicated SRSF6 in maintaining mitochondrial homeostasis and antiviral immunity in macrophages by controlling alternative splicing of the pro-apoptotic factor BAX (Wagner et al., 2022).

Intriguingly, differential phosphorylation of SR proteins has been reported in macrophages infected with intracellular pathogens like the bacterium *Mycobacterium tuberculosis* (Budzik et al., 2020) and the fungal pathogen *Cryptococcus neoformans* (Pandey et al., 2017). As phosphorylation of SR proteins regulates their localization and function (Misteli et al., 1998; Shin et al., 2004; Zhong et al., 2009; Zhou and Fu, 2013), these macrophage phosphoproteomics studies suggest

that functionalization of SR proteins can occur downstream of pattern recognition receptor en-
gagement. We hypothesized that these SR proteins play privileged roles in tuning the macro-
phage response to pathogens. Indeed, we found that stable knockdown of several SR proteins in
RAW 264.7 macrophage-like cells (RAW MΦs) profoundly altered the transcriptome in resting and
activated macrophages (4h post-*Salmonella* Typhimurium infection) (Wagner et al., 2021). The
absence of one of these proteins, SRSF7, generated some particularly interesting phenotypes.
SRSF7 (formerly known as 9G8) is a canonical member of the SR protein family. It encodes an
RNA recognition motif (RRM) and an eponymic SR domain. It is the only SR family protein to also
encode a CCHC-type Zn knuckle domain, which is thought to contribute to its RNA-binding spec-
ificity (Cavaloc et al., 1999; Cavaloc et al., 1994; Wang et al., 2021)). To date, its function is best
studied in the context of controlling its own expression levels via poison cassette exon inclusion
(Konigs et al., 2020), in regulating mRNA export (Muller-McNicoll et al., 2016), and in alternative
splicing (Kadota et al., 2020). Importantly, altered expression of SRSF7 is associated with a vari-
ety of human cancers (Boguslawska et al., 2016; Cun et al., 2021; Fu and Wang, 2018; Song et
al., 2020; Xu et al., 2022).

In RAW MΦs lacking SRSF7, we identified a large cohort of genes whose expression
relied on SRSF7 at rest and after *Salmonella* infection. Curiously, in both conditions, many
SRSF7-impacted genes fell into the category of interferon stimulated genes (ISGs). After compu-
tational quantification of alternatively spliced genes ruled out a role for SRSF7 in mediating alter-
native splicing of ISG pre-mRNAs, we set out to elucidate the mechanism though which SRSF7
controls the type I IFN response in macrophages. Herein, we report a novel role for SRSF7 in
controlling the transcription of interferon responsive factor 7 (IRF7), a critical player in activating
the type I IFN response (Ning et al., 2011). Specifically, we demonstrate that SRSF7 enables
recruitment of signal transducer and activator of transcription 1 (STAT1) to the *Irf7* promoter and
relieves RNA polymerase II pausing. SRSF7 accomplishes this through interactions with the his-
tone methyltransferase KMT5a (a.k.a. SET8), which deposits H4K20me1 at the *Irf7* promoter fol-
lowing LPS treatment to maximize STAT1 association. By revealing a complex regulatory network
at the *Irf7* promoter that controls homeostatic and pathogen-induced type I interferon signaling,
our work identifies an unexpected mechanism through which SR proteins can orchestrate mac-
rophage innate immune gene expression and dictate infection outcomes.
126

RESULTS
SRSF7-dependent alternative splicing is independent of macrophage activation state

As differential phosphorylation of SRSF7 was detected over the course of *M. tuberculosis*
infection in macrophages (Budzik et al., 2020), we were motivated to examine how macrophage
activation might impact its function. To test the hypothesis that alternative splicing (AS) events
might be distinct between resting and activated macrophages, signifying functionalization of
SRSF7, we generated two *Srsf7* knockdown cell lines by transducing RAW MΦs (*Srsf7* KD #1
and *Srsf7* KD #2) with lentivirus containing shRNA hairpin constructs designed against the *Srsf7*
3'UTR and selected these cell lines alongside a scramble (SCR) control (**Fig. 1A-B**). RNA-seq
was performed comparing SCR and *Srsf7* knockdown (KD) #2 in two conditions: uninfected (UN)
or infected with the gram-negative bacterial pathogen *Salmonella enterica* serovar Typhimurium
(4h post-infection; +SAL). *Salmonella* induces robust innate immune responses in macrophages
though TLR4, which activates NFκB downstream of the MyD88 adapter and IRF3/IRF7 down-
stream of TRIF (Fitzgerald et al., 2003). Differential expression analysis was conducted using the
Qiagen CLC Genomics Workbench and quantification of alternative splicing changes was carried
out using MAJIQ (Modeling Alternative Junction Inclusion Quantification) (Vaquero-Garcia et al.,
2016). Consistent with SRSF7's designation as a splicing factor and previous studies implicating
it in alternative splicing specific pre-mRNAs (Gu et al., 2012; Kadota et al., 2020), we identified
1857 total SRSF7-dependent alternative splicing events in resting RAW MΦs and 1909 SRSF7-
dependent events in *Salmonella*-infected RAW MΦs (4h post-infection) (**Fig. 1C**) (p<0.05; >10%
PSI) (**Supplemental Table 1**). As expected for an SR protein that promotes exon recognition/in-
clusion, most changes were at the level of exon skipping. Of the 964 genes subject to SRSF7-
dependent alternative splicing in resting MΦs and the 993 genes in *Salmonella*-infected macro-
phages, 493 of these genes overlapped (**Fig. 1D**). Notably, only a handful of the detected alter-
native splicing changes occurred in genes that were also differentially expressed in *Srsf7* knock-
down macrophages, suggesting that *Srsf7* impacts steady state RNA levels in a splicing-inde-
pendent fashion (**Fig. 1E**).
Because our lists of alternatively spliced genes in resting and *Salmonella*-infected MΦs
did not completely overlap, we hypothesized that SRSF7 mediates distinct alternative splicing
events in the two MΦ states. To begin to test this, we prioritized a set of genes with the most
significant SRSF7-dependent AS changes (>20% PSI; confidence threshold 0.95) in each condi-
tion (UN and +SAL) (**Fig. S1A**). Manual annotation of these 36 genes did not reveal significant
enrichment for any functional category and none of the top alternative splicing events occurred in
genes with characterized roles in the MΦ innate immune response (**Fig. S1A**).
We first looked at SRSF7-dependent exon 6 skipping in *Dcun1d2* (Ensembl Transcript ID:
ENSMUST00000045366.10), a gene associated with Cullin neddylation (Kajigaya et al., 2016).

Figure 1

Figure 1: Macrophage activation does not regulate SRSF7 control of alternative splicing

A. RT-qPCR of *Srsf7* in *Srsf7* KD RAW MΦs relative to SCR controls. **B.** Immunoblot of SRSF7 in shSCR, sh*Srsf7* #1, sh*Srsf7* #2 RAW MΦs. Right, quantification. **C.** Local splicing variations (AS) in *Srsf7* RAW MΦs compared to SCR at baseline (UN; left) and +*Salmonella* Typhimurium (4h) (+SAL; right). ΔPSI > 0.1 confidence threshold set at 0.90. **D.** Venn diagram of genes containing one or more AS event in UN vs. +SAL samples. **E.** Venn diagram of differentially expressed genes (DEGs) vs. alternatively spliced genes (AS) in UN and +SAL samples. **F.** MAJIQ PSI quantification of junctions with SCR (left) and *Srsf7* KD (right) in UN samples. Violin plot of ΔPSI *Dcun1d2* SRSF7-dependent local splicing variation. Splice graph of *Dcun1d2* in SCR (top) and *Srsf7* KD (bottom) generated by MAJIQ/VOILA **G.** As in F but for +SAL samples. **H.** Semi-quantitative RT-PCR of *Dcun1d2* in SCR, *Srsf7* KD #1 and KD #2 RAW MΦs. Each sample in biological triplicate. Right, quantification. **I.** As in H but for *Dcun1d2* +LPS (4h) **J.** As in F but *Mphosph9.* **K.** As in H but *Mphosph9* (+4h LPS). **L.** As in H but *Mphosph9* in UN samples. Data are expressed as a mean of three or more biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

S1

Figure S1: Supplemental data for Figure 1

A. Functional categorization of alternatively spliced transcripts in *Srsf7* knockdown RAW MΦs vs.

SCR (UN, +SAL, and ΔPSI > 0.2 confidence threshold set at 0.95. **B.** MAJIQ PSI quantification

of *Gripap1* junctions with SCR (left) and *Srsf7* KD (right). Violin plot of ΔPSI *Gripap1* SRSF7-

dependent local splicing variation. Splice graph of *Gripap1* in SCR (bottom left) and *Srsf7* KD

(bottom right) generated by MAJIQ/VOILA at baseline. **C.** Semiquantitative PCR of *Gripap1* in

SCR, *Srsf7* KD1 and KD2 RAW MΦs, UN (top) and 4h +LPS (bottom). Each sample in biological

triplicate. Right, quantification. **D.** MAJIQ PSI quantification of *Dop1a* junctions with SCR (left)

and *Srsf7* KD (right). Violin plot of ΔPSI *Dop1a* SRSF7-dependent local splicing variation. Splice

graph of *Dop1a* in SCR (bottom left) and *Srsf7* KD (bottom right) generated by MAJIQ/VOILA at

baseline. **E** Semiquantitative PCR of *Gripap1* in SCR, *Srsf7* KD #1 and KD #2 RAW MΦs, 4h

+LPS (top) and UN (bottom). Each sample in biological triplicate. Right, quantification. Data are

expressed as a mean of three biological replicates with error bars depicting SEM. Statistical sig-

nificance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01,

***=p<0.001, ****=p<0.0001.

MAJIQ measured significant enrichment of this exon skipping event in both untreated and *Salmo-*
nella-infected *Srsf7* knockdown MΦs (ΔPSI 39.2% in UN and 43.4% in +SAL) (**Fig. 1F-G**). We
confirmed this via semi-quantitative RT-PCR using RNA isolated from uninfected MΦs and MΦs
treated with lipopolysaccharide (LPS). LPS activates TLR4 in MΦs in a manner analogous to
Salmonella infection but provides tighter kinetic control experimentally. Consistent with our MAJIQ
analysis, we measured significantly higher levels of *Dcun1d2* exon 6 skipping in *Srsf7* knockdown
MΦs in both conditions (**Fig. 1H-I**).

Next, we looked at an alternative splicing event computationally predicted to be "activa-
tion-dependent" (i.e. only detected in +SAL MΦs) (**Fig. 1D**). In +SAL samples, MAJIQ quantified
SRSF7-dependent alternative 3'SS usage in exon 3 of the *Mphosph9* gene (Ensembl Transcript
ID: ENSMUST00000184951.8), which encodes an M-phase phosphoprotein (*Srsf7* knockdown
#2 (ΔPSI -35.7%) (**Fig. 1J**)). Again, by RT-PCR, we confirmed that this event occurs preferentially
in *Srsf7* knockdown MΦs (**Fig. 1K**). Curiously however, we also measured preferential alternative
3'SS usage in untreated *Srsf7* knockdown MΦs (**Fig. 1L**), even though this event was not com-
putationally identified by RNA-seq/MAJIQ. Similarly, SRSF7-dependent inclusion of exon 8 in
Gripap1 (Ensembl Transcript ID ENSMUST00000065932.14) (**Fig. S1B**), and inclusion of exon
35 in *Dop1a* (Ensembl Transcript ID ENSMUST00000190957.7) (**Fig. S1D)** only met our thresh-
old in UN and +SAL conditions, respectively, but could be detected in both macrophage states by
RT-PCR (**Fig. S1C, S1E**). These data argue that although SRSF7 is important for making alter-
native splicing decisions in RAW MΦs, it largely mediates the same events in the presence and
absence of LPS. Based on the nature of the genes whose alternative splicing relies on SRSF7
and the lack of specificity for alternative splicing in resting vs. activated MΦs, we can conclude
that SRSF7 does not dramatically influence the macrophage innate immune transcriptome at the
level of splicing.

SRSF7 is a positive regulator of interferon stimulated genes (ISGs) in macrophages

In addition to the alternative splicing events reported in Fig. 1, we also measured hundreds
of differentially expressed genes in resting and *Salmonella*-infected *Srsf7* knockdown RAW MΦs
compared to controls (**Supplementary Table 2**). In resting MΦs, *Srsf7* knockdown impacted ex-
pression of many housekeeping genes in resting MΦs, particularly in pathways related to transla-
tion regulation (EIF2 signaling and Regulation of eIF4 and p70S6K signaling) and phagosome
maturation (**Fig. S2A**). Interestingly, in resting MΦs, we also noted enrichment for SRSF7-de-
pendent genes in pathways like "Communication between innate and adaptive immune cells"

Figure 2

Figure 2: SRSF7 promotes the type I interferon response

A. Scatterplot of genes differentially expressed in *Srsf7* KD RAW MΦs relative to SCR in each of the conditions queried (X axis = UN; Y axis = +SAL). Select interferon stimulated genes depicted in black. **B.** Integrated Genomics Viewer track of *Zbp1* SCR (grey) and *Srsf7* KD (teal) reads from *Salmonella*-infected cells. **C.** As in B but for *Ifit3*. **D.** RT-qPCR of *Zbp1* in SCR, *Srsf7* KD#1, and *Srsf7* KD#2 RAW MΦs at baseline (UN). **E.** As in D but for *Ifit3*. **F.** RT-qPCR of *Zbp1* in SCR, *Srsf7* KD#1, and *Srsf7* KD#2 RAW MΦs 4h post-stimulation with 100 ng/ml LPS. **G.** As in F but for *Ifit3*. **H.** RT-qPCR of *Zbp1* in SCR, *Srsf7* KD#1, and *Srsf7* KD#2 RAW MΦs 4h post-transfection with 1 µg/ml ISD. **I.** As in F but for *Ifit3*. **J.** RT-qPCR of *Zbp1* in SCR and *Srsf7* KD #2 RAW MΦs at 4 and 8h post-infection with *M. tuberculosis* (Erdman strain; MOI=5). **K.** As in J but for *Ifit3*. **L.** Immunoblot of ZBP1 protein levels in SCR and *Srsf7* KD #2 RAW MΦs at baseline, 4, and 8h, post-stimulation with 100 ng/ml LPS. Right, quantification. **M.** RT-qPCR of *Srsf7* in primary MEFs at baseline or 4h post stimulation with 100ng LPS. **N.** As in M but for *Zbp1*. **O.** As in M but for *Ifit3*. **P.** As in M but for *Irf7*. **Q.** RT-qPCR of *Vsvg* in SCR, *Srsf7* KD #1, and *Srsf7* KD #2 RAW MΦs at 4, 8, 12h post-infection with VSV (MOI=0.1). **R.** As in M but for *Vsvm*. **S.** RT-qPCR of *Zbp1* in SCR, *Srsf7* KD #1, *Srsf7* KD #2 RAW MΦs at 1, 4, 8, 12h post-infection with VSV (MOI=0.1). **T.** As in S but for *Ifit3*. Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001, ****=p<0.0001.

A

Srsf7 knockdown
Uninfected

EIF2 signaling
Phagosome maturation
Regulation of eIF4 and p70S6K signaling
Antigen presentation pathway
Communication between innate and adaptive immune cells
Polyamine regulation in colon cancer
Virus entry via endocytic pathways
mTOR signaling

0   5   10   15   20

B

Srsf7 knockdown
+*Salmonella*

Communication between innate and adaptive immune cells
Role of pattern recognition receptors in recognition of bacteria and viruses
Systemic lupus erythematosus in B cell signaling pathway
Role of PKR in interferon induction and antiviral response
Interferon signaling
Activation of IRF by cytosolic pattern recognition receptors
Granulocyte adhesion and diapedesis
Agranulocyte adhesion and diapedesis

0   5   10   15

C

+*Salmonella*

D

+*M. tuberculosis*

SCR
Srsf7 #2

E

+LPS
MEFs

SCR
Srsf7 #2

F

+VSV

SCR
Srsf7 #1
Srsf7 #2

S2

Figure S2: Supplemental data for Figure 2

A. Ingenuity Pathway Analysis (IPA; Qiagen) of pathways enriched for differentially expressed

genes in *Srsf7* KD RAW MΦs (UN). **B.** As in A for *Srsf7* KD RAW MΦs +SAL. **C.** Infection of SCR

and *Srsf7* KD RAW MΦs with *Salmonella* enterica (MOI=1) in non-SPI-1 inducing conditions. Fold-

replication represented as 22h over 2h post-infection. **D.** Mtb luxBCADE growth in SCR or *Srsf7*

KD RAW MΦs measured by relative light units (RLUs) over a 4-day time course (MOI=1). **E.** RT-

qPCR of *Rsad2* in siRNA *Srsf7-* or siControl-transfected primary MEFs at baseline or 4h post

stimulation with 100ng/ml of LPS. **F.** RT-qPCR of *Il1b* in SCR or *Srsf7* KD RAW MΦs over a time

course of VSV infection. Data are expressed as a mean of three or more biological replicates with

error bars depicting SEM. Statistical significance was determined using two tailed unpaired stu-

dent's t test. *=$p<0.05$, **=$p<0.01$.

196   (**Fig. S2A**), suggesting that loss of SRSF7 dysregulates innate immune gene expression even in

197   the absence of a stimulus.

198   Enrichment for pathways related to innate immunity was also evident in *Salmonella*-in-

fected macrophages, where knockdown of *Srsf7* dysregulated genes in pathways related to "role

of pattern recognition receptors in recognition of bacteria and viruses," "interferon signaling," "role

of PKR in interferon induction and antiviral response," and "activation of IRF by cytosolic pattern

recognition receptors" (**Fig. S2B**). Many of these categories contain genes related to the type I

interferon (IFN) response, a ubiquitous antiviral response characterized by expression of inter-

feron alpha and beta and interferon stimulated genes (ISGs) (Schoggins, 2019; Schoggins and

Rice, 2011). In MΦs, the type I IFN response is activated by the transcription factors IRF3 and

IRF7 in response to several immune agonists including LPS/*Salmonella* infection, cytosolic

dsDNA, and cytosolic dsRNA (Honda and Taniguchi, 2006; Jefferies, 2019; Schoggins et al.,

2011).

209   For ISGs that are appreciably expressed in resting MΦs (Barrat et al., 2019; Gough et al.,

2012), e.g. *Apol9a*, *Apol9b*, *Ifit3*, and *Ifi202b*, *Srsf7* knockdown decreased their abundance in

both untreated and +*Salmonella* conditions (**Fig. 2A**; bottom left quadrant). Genes like *Mx1, Zbp1,*

and *Ifi209*, which are virtually undetectable in resting cells, only displayed SRSF7-dependence in

our *Salmonella*-infected samples (**Fig. 2A**; lower y-axis). Visualizing RNA reads from *Srsf7* knock-

down versus control MΦs for two such ISGs, *Zbp1* and *Ifit3*, on the Integrated Genomics Viewer,

revealed no changes in intronic reads (**Fig. 2B-C**). This was consistent with our MAJIQ analysis,

which did not detect many statistically significant splicing changes in genes annotated as ISGs in

Srsf7 knockdown RAW MΦs (**Fig. 1E, Supplemental Table 1)**.

218   Failure to phagocytose *Salmonella* or defects in *Salmonella* replication could result in

lower ISG expression in *Srsf7* knockdown cells, but enumeration of colony forming units (CFUs)

did not reveal any differences in bacterial burdens between SCR and *Srsf7* knockdown RAW MΦs

over a 22h infection time-course (**Fig. S2C**). To determine if SRSF7's ability to promote ISG ex-

pression was unique to *Salmonella* infection, we measured ISG expression in SCR and *Srsf7*

knockdown MΦs at 4h post-treatment with LPS (100 ng/ml), after dsDNA transfection, which ac-

tivates type I IFN gene expression downstream of cGAS (1 μg ISD), and after treatment with

recombinant IFN-β, which activates ISG expression in MΦs by binding to IFNAR and engaging

JAK/STAT cascades (200 IU, 4h). In all cases, loss of *Srsf7* led to impaired activation of *Zbp1*

and *Ifit3* expression (**Fig. 2D-I**). We also measured lower ISG expression in *Srsf7* knockdown

MΦs during infection with the intracellular bacterial pathogen, *Mycobacterium tuberculosis* (MOI

= 5) (**Fig. 2J-K**), which elicits *Ifnb1* and ISG expression via the cGAS/STING pathway (Watson et

al., 2015). Again, this defect in ISG expression was independent of bacterial internalization/repli-

cation (**Fig. S2D**). Over an 8h time-course of LPS-treatment, lower *Zbp1* transcript levels corre-

lated with lower protein accumulation (**Fig. 2L**), suggesting that failure to fully upregulate ISG

expression in *Srsf7* knockdown MΦs is likely to functionally impact the innate immune response

to infection. This phenotype was not unique to RAW MΦs, as siRNA knockdown of SRSF7 in

primary mouse embryonic fibroblasts (MEFs)(**Fig. 2M**), a cell type commonly used for studies of

type I interferon signaling, also resulted in low basal ISG expression and a failure to fully induce

ISGs upon LPS stimulation (**Fig. 2N-P and S2E**).

To test whether SRSF7 is a *bona fide* player in activating antimicrobial immunity in MΦs,

we infected SCR and *Srsf7* knockdown MΦs with vesicular stomatitis virus (VSV), a negative-

strand RNA virus that stimulates type I IFN responses via TLR3 and RIG-I and is extremely sen-

sitive to ISG expression (Gao et al., 2021; Wagner et al., 2021). We measured a dramatic increase

in VSV replication in *Srsf7* knockdown MΦs, measured by RT-qPCR of two VSV genes, *Vsvg* and

Vsvm (**Fig. 2Q-R**), that was concomitant with depressed ISG transcript induction over a 12-hour

time course of VSV infection (MOI = 0.1) (**Fig. 2S-T**). We did not detect SRSF7-dependent differ-

ences in *Il1b* expression in response to VSV infection, arguing against a role for SRSF7 in con-

trolling innate gene expression downstream of NFκB (**Fig. S2F**). Together, these data identify

SRSF7 as a positive regulator of the type I IFN response in MΦs and suggest that the mechanism

of SRSF7-dependent ISG expression is shared by several pathogen sensing cascades (i.e. oc-

curs at one or more nodes where the TLR4, cGAS, and RIG-I cascades converge).

SRSF7 drives ISG expression by promoting expression of the transcription factor IRF7

Given that loss of SRSF7 results in a failure to fully induce many ISGs, we hypothesized

that SRSF7 may act directly on one or more critical innate signaling proteins or transcription fac-

tors. To pinpoint where in the type I IFN signaling response SRSF7 acts, we treated *Srsf7*

knockdown or SCR RAW MΦs with LPS (100 ng/ml) and measured secretion of type I interferon

(IFN-β/α) over a 6h time-course via an ISRE-luciferase reporter cell line (Hoffpauir et al., 2020;

Wagner et al., 2022). *Srsf7* knockdown MΦs produced significantly less IFN-β/α compared to

SCR controls at all time points measured (**Fig. 3A**). We next asked if impaired ISG expression

in *Srsf7* knockdown MΦs could be rescued if *Srsf7* knockdown and SCR control cells received

equal amounts of recombinant IFN-β. We found that even post treatment of IFN-β, *Srsf7* knock-

down MΦs fail to express ISGs like *Ifi202b* and *Zbp1* to the same level as SCR controls, at both

the transcript (**Fig. 3B-C**) and protein level (**Fig. 3D**). These results suggest that loss of SRSF7

impacts some aspect of the type I IFN response that is engaged upstream and downstream of

Figure 3

Figure 3: SRSF7 is required for IRF7 accumulation in macrophages

A. Secreted IFNα/β in SCR and *Srsf7* KD#2 RAW MΦs, as a measure of luciferase expressed by

ISRE reporter cells. **B.** RT-qPCR of *Ifi202b* in SCR, *Srsf7* KD#1 and *Srsf7* KD#2 in RAW MΦs 4h

post-stimulation with 200 IU IFN-β. **C.** As in B for *Zbp1.* **D.** Immunoblot of ZBP1 and ACTIN over

a time course of IFN-β (200 IU) treatment in SCR and *Srsf7* KD#2 RAW MΦs. Quantification,

right. **E.** Immunoblot of Y701 STAT1, STAT1, S396 IRF3, IRF3, and ACTIN in SCR and *Srsf7* KD

RAW MΦs over a time course of LPS treatment (100 ng/ml). **F.** Differential expression of tran-

scription factor genes in untreated *Srsf7* knockdown vs. SCR RAW MΦs as fold change and -

log(p-value). Gene of interest (*Irf7*) highlighted in red. **G.** As in F but for +SAL samples. **H.** Inte-

grated Genomics Viewer with *Irf7* reads from SCR (grey) and *Srsf7* KD#2 (teal) RAW MΦs, un-

treated (top) and +*Salmonella* (bottom). **I.** RT-qPCR of *Irf7* over a time course of LPS stimulation

(100 ng/ml) in SCR and *Srsf7* KD RAW MΦs. **J.** Immunoblot of IRF7 and ACTIN protein levels

over a time course of LPS stimulation (100 ng/ml) in SCR and *Srsf7* KD RAW MΦs. Quantification,

right. **K.** RT-qPCR of *Irf7* over a time course of Mtb infection (Erdman strain; MOI=5) in SCR and

Srsf7 KD#2 MΦs. **L.** RT-qPCR of *Irf7* over a time course of VSV infection (MOI=0.1) in SCR, *Srsf7*

KD#1, *Srsf7* KD#2 RAW MΦs. **M.** Immunoblot of SRSF7, IRF7, and ACTIN in siControl- or si*Srsf7*-

transfected BMDMs at baseline. Quantifications, right. **N.** RT-qPCR of *Irf7* in siControl or si*Srsf7*

BMDMs at baseline. **O.** Immunoblot of IRF7 protein levels in GFP gRNA, *Irf7* KO1 and *Irf7* KO2

RAW MΦs at baseline. ACTIN used as loading control. **P.** RT-qPCR of *Ifit3b, Zbp1,* or *Ifnb1* at

baseline in *Irf7* KO1 (blue) vs. GFP gRNA control (grey) RAW MΦs. **Q.** Integrated Genomics

Viewer with *Ifi44* and *Ifi202b* reads from GFP gRNA (grey) or *Irf7* KO (blue) RAW MΦs at baseline

(UN). Data expressed as a mean of three biological replicates with error bars depicting SEM

unless otherwise noted. Statistical significance was determined using two tailed unpaired stu-

dent's t test. *=p<0.05, **=p<0.01, ***=p<0.001, ****=p<0.0001.

IFNAR and that loss of IFN-β production is not the sole driver of ISG defects in *Srsf7* knock-

down MΦs.

We then asked whether altered signaling downstream of TLR4 could explain defective

ISG expression in *Srsf7* knockdown RAW MΦs. We treated *Srsf7* knockdown and SCR RAW

MΦs with LPS and measured phosphorylation of the transcription factor IRF3 (activated down-

stream of TLR4-TRIF signaling) and STAT1 (activated downstream of IFNAR) over a 6h time

course. We did not measure significant changes in phosphorylation of IRF3 at S396 or phos-

phorylation of STAT1 at Y701 (**Fig. 3E**; quantification in **Fig. S3A**), effectively ruling out defec-

tive signal transduction as the culprit for dampened ISG expression in *Srsf7* knockdown MΦs.

Next, we entertained the possibility that SRSF7 is required for the expression of one or

more transcription factors needed to activate the type I IFN response downstream of pattern

recognition receptor and/or IFNAR engagement. To test this, we generated a list of transcription

factors associated with macrophage innate immunity in the STAT, IRF, and NFκB families, and

turned back to our RNA-seq data to see how expression of each factor was impacted by loss of

SRSF7. While expression of most transcription factors was unaffected, we observed a signifi-

cant defect in *Irf7* expression in *Srsf7* knockdown RAW MΦs at baseline and after *Salmonella*

infection (**Fig. 3F-G**). Visualization of *Irf7* gene expression data on the IGV highlights fewer *Irf7*

exonic reads in *Srsf7* knockdown (teal) compared to SCR (grey) RAW MΦs in both conditions

(**Fig. 3H**). We did not detect an increase in intron reads in *Srsf7* knockdown MΦs nor did our

MAJIQ analysis identify SRSF7-dependent *Irf7* alternative splicing events (**Fig. 1D**), arguing

against a role for SRSF7 in regulating *Irf7* transcript abundance via splicing.

Initially discovered in the context of Epstein Barr virus, IRF7 (interferon regulatory factor

7) is a master regulator of antiviral type I IFN responses (Honda et al., 2005). Importantly, IRF7

is itself an ISG whose expression is induced downstream of IFNAR engagement. Although ba-

sal levels of IRF7 vary between immune cell types, we know that dendritic cells are heavily reli-

ant upon IRF7 at baseline and several studies have implicated IRF7 in priming early responses

in macrophages downstream of LPS (Fitzgerald et al., 2003; Gu et al., 2022; Jefferies, 2019;

Sin et al., 2020). We can readily measure IRF7 transcript and protein in resting wild-type RAW

MΦs (**Fig. 3I-J**) and in wild-type immortalized BMDMs (**Fig. S3B**), confirming their utility for

studies of IRF7 in resting and activated states. IRF7 protein levels are low in WT resting

BMDMs and are upregulated in response to LPS (**Fig. S3C**).

To better appreciate the kinetics of SRSF7-dependent control of *Irf7* expression, we

treated *Srsf7* knockdown and SCR control RAW MΦs with LPS and measured *Irf7* transcript and

protein over a 12h time course. Levels of *Irf7* mRNA were significantly lower in *Srsf7* knockdown

S3

Figure S3: Supplemental data for Figure 3

A. Quantification of Figure 3E; n=2 **B.** Immunoblot of IRF7 and ACTIN protein levels over a time course of LPS stimulation (100 ng/ml) in WT iBMDMs. **C.** As in B but in WT BMDMs. **D.** Venn diagram depicting differential gene expression of *Irf7* KO1 at baseline vs. 4h +LPS (100ng/ml). Cut off: fold change < -1.5 and p-value <0.05. **E.** Top: BioPlanet Pathway Analysis of untreated *Irf7* KO gene expression. Bottom: BioPlanet Pathway Analysis of 4h +LPS *Irf7* KO gene expression. **F.** Heatmap depicting *Irf7* KO baseline gene expression from BioPlanet pathway analysis categories: "Interferon alpha/beta signaling," "Interferon signaling," and "Interferon-gamma signaling pathway." Duplicate terms removed. **G.** Left: Venn diagram depicting differentially expressed genes (DEGs) in *Irf7* KO1 (compared to GFP gRNA) and *Srsf7* KD (compared to SCR) RAW MΦs at baseline. Right: Heatmap of 26 genes in common between *Irf7* KO1 and *Srsf7* KD at baseline. Data are expressed as a mean of three biological replicates with error bars depicting SEM unless otherwise noted. Statistical significance was determined using two tailed unpaired student's t test.

RAW MΦs at every time-point queried (**Fig. 3I**). IRF7 protein levels were lower in *Srsf7* knock-
down RAW MΦs at 0 and 4h but appeared to rebound at later time points (8 and 12h post-LPS)
(**Fig. 3J**), consistent with earlier data suggesting that IRF7 protein can be stabilization during
the innate response to infection (Ning et al., 2011). We also measured significantly less *Irf7*
transcript in *Srsf7* knockdown RAW MΦs in response to *M. tuberculosis* (**Fig. 3K**) and VSV in-
fection (**Fig. 3L**). Although we were only able to achieve modest knockdown of SRSF7 in
BMDMs transfected with an siRNA directed against *Srsf7* (alongside a non-targeting control;
siNC) (**Fig. 3M**), we can detect less IRF7 protein (**Fig. 3M**) and transcript (**Fig. 3N**) in these pri-
mary MΦs as well. Together, our data demonstrate a previously unappreciated role for SRSF7
in controlling *Irf7* transcript and protein levels in macrophage cell lines, primary macrophages,
(**Fig. 3I-J** and **3M-N**) and MEFs (**Fig. 2P**).
309
IRF7 controls basal and induced expression of ISGs in RAW MΦs
Our findings argue that loss of ISG expression in *Srsf7* knockdown cells stems primarily
from dampened levels of IRF7. However, as most literature points to a role for IRF7 in activated
immune cells, we were initially cautious to attribute basal ISG phenotypes in *Srsf7* knockdown
MΦs to IRF7. To better understand how IRF7 controls type I IFN responses in resting and acti-
vated RAW MΦs, we generated *Irf7* KO RAW MΦs using CRISPR-Cas9 editing with a gRNA de-
signed to target *Irf7* exon 3, and negative control cells expressing a GFP-specific (non-targeting)
gRNA. Two clonally expanded *Irf7* KO cell lines and a GFP gRNA control cell line were se-
lected, and loss of IRF7 protein was confirmed by immunoblot (**Fig. 3O**). Total RNA from un-
treated *Irf7* KO #1 and control cell lines, as well as from cells stimulated with LPS for 4h (100
ng/ml) were sent for RNA-seq analysis. We measured differential expression of 896 transcripts
(fold-change < -1.5; p<0.05) in *Irf7* KO vs. control cells at baseline and 1575 genes were differ-
entially expressed in *Irf7* KO MΦs at 4h post-LPS treatment (**Supplemental Table 3** and **Fig.**
S3D). Pathway analysis (BioPlanet Pathway Enrichment Analysis, NCATS (Huang et al., 2019))
identified a number of type I interferon-related pathways enriched for differentially expressed
genes in *Irf7* KO cells at baseline and at 4h post-LPS treatment (**Fig. S3E**). Consistent with our
hypothesis that SRSF7 controls ISG expression in an IRF7-dependent fashion, we found that
ISG expression was dramatically dampened in *Irf7* KO RAW MΦs at rest and after LPS stimula-
tion (**Figs. 3P-Q, S3F,** and **Supplemental Table 3**) and that differentially expressed ISGs were
largely shared between *Srsf7* KD and *Irf7* KO cells (**Fig. S3G**). Together, these data provide
strong evidence for IRF7 maintaining homeostatic expression of type I IFN and ISGs in macro-
phages and suggest that SRSF7 contributes to this circuit by mediating IRF7 expression.

SRSF7 overexpression is sufficient to drive IRF7 and ISG expression

We hypothesized that if SRSF7 can directly activate *Irf7* transcription, then overexpression of SRSF7 should be sufficient to drive *Irf7* expression. To test this model, we generated doxycycline-inducible RAW MΦ cell lines that express 3xFLAG-SRSF7 or 3xFLAG-GFP. To ensure comparable levels of 3xFLAG-GFP and 3xFLAG-SRSF7 protein expression and to best mimic endogenous levels of SRSF7, we tested and optimized treatment times, deciding on 2h +DOX for 3xFLAG-GFP and 22h +DOX for 3xFLAG-SRSF7 (**Fig. 4A**). Notably, we found that expression of SRSF7 alone was sufficient to increase the abundance of *Irf7* transcript (**Fig. 4B**), IRF7 protein levels (**Fig. 4C**), and ISG transcripts (**Fig. 4D**) in otherwise unstimulated RAW MΦs. Doxycycline-inducible expression of 3xFLAG-SRSF7 was also sufficient to drive expression of additional genes in the IRF7 baseline transcriptional regulon (*Xaf*, *Irak2*, *Ccl2*, and *Isg20* (**Figs. 4E and S3G**)), lending further support to our hypothesis that SRSF7 controls ISG expression at the level of IRF7. We observed a similar phenotype in RAW MΦ cell lines that stably express 3xFLAG-SRSF7 compared to 3xFLAG-GFP controls (**Fig. 4F**). While overexpression of 3xFLAG-SRSF7 had a less pronounced impact on IRF7 transcript and protein expression after LPS stimulation (**Fig. S4A-B)**, we measured a dramatic increase in *Irf7* and *Ifit3* transcript levels in 3xFLAG-SRSF7-expressing stable cells upon infection with *M. tuberculosis* (**Fig. 4G**). Because expression of interferon stimulated genes has been associated with cell-intrinsic antibacterial control mechanisms and cell death pathways that impact bacterial spread (Banks et al., 2019; Hoffpauir et al., 2020; MacMicking et al., 1997; MacMicking et al., 2003), we also asked whether overexpression of SRSF7 affected *M. tuberculosis* replication. Briefly, we infected cells with Mtb *luxBCADE* and followed bacterial replication as a measure of relative light units (as in (Bell et al., 2021; Wagner et al., 2022)). We measured a significant increase in RLUs at Day 3 post-infection (**Fig. 4H**) suggesting that SRSF7 expression may help control *M. tuberculosis* replication and/or spread. Together, these data suggest a direct role for SRSF7 in activating expression of *Irf7* and promoting type I interferon responses in MΦs.

SRSF7 promotes STAT1 association and RNAPII elongation at the *Irf7* promoter

Having established that decreased levels of IRF7 is the likely culprit for the type I IFN defect in *Srsf7* knockdown MΦs (**Fig. 3**) and demonstrated that SRSF7 overexpression is sufficient to promote *Irf7* expression (**Fig. 4**), we set out to determine the mechanism though which SRSF7 controls *Irf7* expression. Earlier work has implicated SR proteins in coupling transcription and splicing and have been shown to influence transcription by interacting with histone tails

Figure 4

Figure 4: SRSF7 overexpression is sufficient to drive basal expression of IRF7 and ISGs

A. Immunoblot of 3xFLAG-GFP or 3xFLAG-SRSF7 tetracycline inducible RAW MΦs treated with doxycycline (DOX) (1 µg/ml). 3xFLAG-GFP MΦs received DOX for 2h and 3xFLAG-SRSF7 MΦs received DOX for 22h. ACTIN shown as loading control. **B.** RT-qPCR of *Irf7* in 3xFLAG-GFP (2h DOX treatment) and 3xFLAG-SRSF7 (22h DOX treatment) inducible RAW MΦs at baseline. **C.** Immunoblot of IRF7 in 3xFLAG-GFP (2h DOX treatment) and 3xFLAG-SRSF7 (22h dox treatment) inducible RAW MΦs at baseline. Quantification right. **D.** As in B but for *Ifit3*. **E.** As in B but for *Xaf, Irak2, Ccl2,* and *Isg20*. **F.** RT-qPCR of *Zbp1*, *Ifi44*, or *Oasl2* in 3xFLAG-GFP and 3xFLAG-SRSF7 stable overexpression RAW MΦs at baseline. **G.** RT-qPCR of *Irf7* or *Ifit3* at 0, 4, and 8h post-infection with Mtb (Erdman; MOI=5). **H.** Mtb *luxBCADE* replication in 3xFLAG-GFP and 3xFL-SRSF7 doxycycline-inducible RAW MΦs measured by relative light units (RLUs) over time course (Erdman; MOI=1). Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

A

+LPS
■ 3xFL-GFP
days ■ 3xFL-SRSF7

Irf7/Actb

**

hours

B

+LPS
100ng

2h   4h   6h

3xFL-GFP 3xFL-SR7 3xFL-GFP 3xFL-SR7 3xFL-GFP 3xFL-SR7

55—   α-IRF7

35—   α-ACTIN

■ 3xFL-GFP ■ 3xFL-SRSF7

IRF7/ACTIN

hours

S4

Figure S4: Supplemental data for Figure 4

A. RT-qPCR of *Irf7* in 3xFLAG-GFP (2h dox treatment) and 3xFLAG-SRSF7 (22h dox treatment)

inducible RAW MΦs over a time course of LPS stimulation (100 ng/ml). **B.** Immunoblot of IRF7 in

3xFLAG-GFP (2h dox treatment) and 3xFLAG-SRSF7 (22h dox treatment) inducible RAW MΦs

over a time course of LPS stimulation (100 ng/ml) (quantification right). Data are expressed as a

mean of three biological replicates with error bars depicting SEM. Statistical significance was

determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

(Loomis et al., 2009; Luco et al., 2011; Luco et al., 2010) or by directly mediating transcriptional

elongation via pTEFb (Ji et al., 2013). To begin to investigate a role for SRSF7 in activating *Irf7*

transcription, we performed chromatin immunoprecipitation (ChIP)-qPCR for RNA polymerase II

(RNAPII) in *Srsf7* knockdown and SCR control MΦs at 4h post-LPS treatment, using primers de-

signed to tile the *Irf7* promoter region (100bp amplicon, 100 bp apart from each other (**Fig. 5A**))

and an antibody that recognizes both phosphorylated and unphosphorylated forms of the C-ter-

minal domain of the largest subunit of RNAPII. We observed dramatic accumulation of RNAPII

at/around the *Irf7* transcription start site (TSS) (-400 to +250) in *Srsf7* knockdown MΦs (**Fig. 5B**,

teal). This pattern of RNAPII in *Srsf7* knockdown MΦs is suggestive of a pile-up resulting from a

failure to convert to a successful elongation complex. We next asked whether a defect in

RNAPII clearance could be explained by defects in recruitment or retention of transcription fac-

tors like STAT1 to the *Irf7* promoter. Using ChIP-qPCR with an antibody against Y701 STAT1,

we measured a near total loss of STAT1 signal in *Srsf7* knockdown MΦs (**Fig. 5C**). An analo-

gous set of ChIP-qPCRs at the promoter of *Ifit3*, another ISG whose expression is SRSF7-de-

pendent, showed no differences in RNAPII or STAT1 occupancy in *Srsf7* knockdown cells

(**Figs. 5D-F**). Likewise, we saw no difference in RNAPII occupancy in *Srsf7* knockdown MΦs at

the promoter of *Stat1* itself (**Figs. 5G-H**). Consistent with upregulation of *Irf7* transcript and pro-

tein levels (**Figs. 4B-C**), overexpression of 3xFLAG-SRSF7 was sufficient to recruit significant

levels of STAT1 to the *Irf7* promoter in resting macrophages (**Fig. 5I**). These data support a

role for SRSF7 in promoting transcription of *Irf7* and suggest that dampened expression of ISGs

like *Ifit3* in *Srsf7* knockdown MΦs occurs indirectly, downstream of *Irf7* transcriptional defects.

Having linked STAT1 binding and RNAPII elongation at the *Irf7* promoter to SRSF7, we

next asked whether SRSF7 carries out this function by directly associating with the *Irf7* genomic

locus. Lacking ChIP-grade antibodies for SRSF7, we turned back to our doxycycline-inducible

3xFLAG-SRSF7 and 3xFLAG-GFP cell lines (**Fig. 4A**) and performed anti-FLAG ChIP-qPCR at

the *Irf7* promoter using the same primer sets shown in Fig. 6A. Consistent with a direct role for

SRSF7 in facilitating STAT1 recruitment and RNAPII elongation at *Irf7*, we observed significant

enrichment of 3xFLAG-SRSF7 at the *Irf7* promoter (-500 to 0) in both resting (**Fig. 5J**) and LPS-

treated macrophages (4h post-LPS) (**Fig. 5K**) with no SRSF7 enrichment at *Ifit3* (**Fig. 5L**). To

determine the extent to which SRSF7 globally associates with promoter regions, we performed

ChIP-seq. Briefly, reads were trimmed by Cutadapt and aligned to the *Mus musculus* genome

build mm10 using Bowtie. After quality control, peaks were called using MACS2 with input/IgG

controls background subtracted. We detected 548 unique peaks for SRSF7 across the genome,

with approximately half of these peaks detected at transcription start sites (**Figs. 5M and S5A-**

Figure 5

**Figure 5: SRSF7 recruitment to the *Irf7* promoter maximizes STAT1 association and pro-
motes RNA polymerase II elongation**

A. Diagram of the *Irf7* gene with tiling primers designed around the transcriptional start site (or-
ange). **B.** RNA polymerase II ChIP-qPCR (antibody recognizes both modified and unmodified
CTD) at the *Irf7* promoter in SCR and *Srsf7* KD RAW MΦs 4h post-LPS stimulation (100ng/ml).
Data represented as elution/input normalized to that of an intergenic control primer set. **C.** Y701
STAT1 ChIP-qPCR at the *Irf7* promoter in SCR or *Srsf7* KD RAW MΦs 4h post-LPS stimulation
(100 ng/ml). Data represented as elution/input normalized to that of an intergenic control. **D.** Dia-
gram of the *Ifit3* gene with tiling primers designed around the transcriptional start site (orange). **E.**
As in B but at the *Ifit3* promoter. **F.** As in C but at the *Ifit3* promoter. **G.** Diagram of the *Stat1* gene
with tiling primers designed around the transcriptional start site (orange). **H.** As in B but at the
Stat1 promoter. **I.** Y701 STAT1 ChIP-qPCR at the *Irf7* promoter in 3xFLAG-GFP and 3xFLAG-
SRSF7 dox-inducible RAW MΦs at baseline. **J.** 3xFLAG ChIP-qPCR of 3xFLAG-GFP and
3xFLAG-SRSF7 in dox-inducible RAW MΦs at baseline. **K.** 3xFLAG ChIP-qPCR of 3xFLAG-GFP
and 3xFLAG-SRSF7 in dox-inducible RAW MΦs at the *Irf7* promoter 4h post LPS stimulation (100
ng/ml). **L.** As in K at the *Ifit3* promoter. **M.** Categorization of unique 3xFLAG ChIP peaks in 3xFL-
SRSF7 dox-inducible RAW MΦs 4h post-LPS (100 ng/ml). **N.** Enrichment of 3xFLAG-SRSF7 and
3xFLAG-GFP at the *Irf7* promoter by ChIP-seq: IP (top) and input (bottom). **O.** As in M but at the
Ifit3 promoter. Unless otherwise noted, data are expressed as a mean of three or more biological
replicates with error bars depicting SEM. ChIP-qPCRs are shown as the mean of three or more
technical replicates with error bars depicting SEM. Statistical significance was determined using
two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

E

Motif	Name	P-value	Percent of Targets	Percent of Background
	Etv2(ETS)/ES-ER71-ChIP-Seq(GSE59402)/Homer	1e-6	16.79%	9.68%
	ZNF41(Zf)/HEK293-ZNF41.GFP-ChIP-Seq(GSE58341)/Homer	1e-6	1.46%	0.11%
	Fli1(ETS)/CD8-FLI-ChIP-Seq(GSE20898)/Homer	1e-6	25.36%	17.02%
	AT1G76870(Trihelix)/col-AT1G76870-DAP-Seq(GSE60143)/Homer	1e-6	4.93%	1.63%
	Elk4(ETS)/Hela-Elk4-ChIP-Seq(GSE31477)/Homer	1e-5	19.34%	12.62%
	ETV4(ETS)/HepG2-ETV4-ChIP-Seq(ENCODE)/Homer	1e-4	25.36%	18.08%
	ETS(ETS)/Promoter/Homer	1e-4	12.04%	7.09%
	ELF1(ETS)/Jurkat-ELF1-ChIP-Seq(SRA014231)/Homer	1e-4	17.34%	11.50%
	NAM(NAC)/col-NAM-DAP-Seq(GSE60143)/Homer	1e-4	16.06%	10.71%

S5

Figure S5: Supplemental data for Figure 5

A. Heatmap indicating read coverage density at transcriptional start sites (TSS) in 3xFL-GFP or 3xFL-SRSF7 dox-inducible RAW MΦs, 4h post-LPS treatment. **B.** Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Nfkbia* promoter by ChIP-seq: IP (top) and input (bottom). **C.** Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Elk4* promoter by ChIP-seq: IP (top) and input (bottom). **D.** Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Stat1* promoter by ChIP-seq: IP (top) and input (bottom). **E.** Putative SRSF7 motifs predicted by HOMER from 3xFLAG-SRSF7 ChIP-seq data.

C). By ChIP-seq we again saw clear enrichment of 3xFLAG-SRSF7 at the *Irf7* promoter (**Fig. 5N**),

but not at *Ifit3* (**Fig. 5O**) or *Stat1* (**Fig. S5D**). Although SRSF7 peaks were called at several im-

mune associated genes, including *Fos*, *Myd88*, *Ifitm3*, and *Dusp5*, there was no evidence for

SRSF7 enrichment at the promoter of any other type I IFN response-associated transcription fac-

tors (**Supplemental Table 4**). Taken together, our ChIP-seq data argue against universal asso-

ciation of SRSF7 with gene promoters, as was reported for factors like SRSF1 and SRSF2 (Ji et

al., 2013), and instead support a privileged role for SRSF7 in facilitating transcription at specific

genes like *Irf7*.

SRSF7 interacts with the H4K20me1 histone methyltransferase KMT5a/SET8 to promote

***Irf7* transcription**

ChIP-seq did not reveal a strong consensus sequence for SRSF7 and even weak consen-

sus sequence enrichment was only detected at subsets of genes (**Fig. S5E**). As this argues

against recruitment of SRSF7 to the *Irf7* promoter via a specific DNA-encoded motif, we hypoth-

esized that SRSF7 could be brought to/retained at the *Irf7* promoter via one or more protein-

protein interactions. To begin to explore this possibility, we generated RAW MΦ cell lines stably

expressing 3xFLAG-SRSF1, 3xFLAG-SRSF3, 3xFLAG-SRSF6, 3xFLAG-SRSF9, in addition to

the previously described 3xFLAG-SRSF7 and 3xFLAG-GFP cell lines. We performed a series of

immunoprecipitation experiments followed by mass spectrometry to identify peptides enriched in

each pulldown. Although many protein binding partners were shared between SRSF7 and other

SR protein family members (e.g. SR proteins, hnRNPs, RRM domain-containing proteins) (**Fig.**

S6A), peptides derived from components of the KMT5a/SET8 histone methyltransferase complex

were unique to SRSF7 (**Fig. 6A**). KMT5a is the sole histone methyltransferase responsible for

depositing monomethylation at H4K20. H4K20me1 levels are linked to the cell cycle and during

mitosis H4K20me1 mediates chromatin condensation (van Nuland and Gozani, 2016). Several

studies report enrichment of H4K20me1 at actively transcribed genes (Barski et al., 2007),

(Veloso et al., 2014) and it has been implicated in facilitating RNAPII escape from promoter-prox-

imal regions (Nikolaou et al., 2017). We confirmed interaction between 3xFLAG-SRSF7 and

KMT5a in macrophages by immunoprecipitation/immunoblot, using antibodies against endoge-

nous KMT5a (**Fig. 6B**).

To determine a role for KMT5a in mediating *Irf7* expression, we silenced its expression

via siRNA transfection in RAW MΦs, alongside a non-targeting control. We found that loss of

KMT5a dampened *Irf7* transcript levels and those of downstream ISGs, phenocopying SRSF7

knockdown (**Fig. 6C**). Expression of inflammatory genes outside the *Irf7* regulon, like *Tnf*, was

Figure 6

Figure 6: SRSF7 interacts with the H4K20me1 histone methyltransferase KMT5a/SET8 to

promote *Irf7* transcription

A. Proteins identified as unique interactors of 3xFL-SRSF7 in RAW MΦs at baseline by IP-LC/MS.

B. Immunoblot of immunoprecipitation of 3xFL-SRSF7 (+DOX 1 μg/ml 22h), 3xFL-GFP (+DOX 1

μg/ml 2h) and endogenous KMT5a/SET8. **C.** RT-qPCR of *Kmt5a, Irf7*, or *Ifit3b* in RAW MΦs trans-

fected with siControl or si*Kmt5a* siRNA at baseline. **D.** Y701 STAT1 ChIP-qPCR at the *Irf7* pro-

moter in siControl or si*Kmt5a/Set8* RAW MΦs at baseline. Data represented as elution/input nor-

malized to that of an intergenic control. **E.** RT-qPCR of *Kmt5a* and *Irf7* in bone marrow derived

macrophages (BMDMs) transfected with siControl or si*Kmt5a* siRNA. **F.** As in E but at 4h post-

stimulation with LPS (100 ng/ml). **G.** H4K20me1 ChIP-qPCR in RAW MΦs at baseline or 4h post

stimulation with 100 ng/ml LPS at the *Irf7* promoter. Data represented as elution/input normalized

to IgG. **H.** As in G but in BMDMs. **I.** Proposed model for cooperation between SRSF7 and KMT5a

promoting transcriptional activation of *Irf7*. Unless otherwise noted, data are expressed as a mean

of three or more biological replicates with error bars depicting SEM. ChIP-qPCRs are shown as

the mean of three or more technical replicates with error bars depicting SEM. Statistical signifi-

cance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01,

***=p<0.001.

A

B
UN
RAW 264.7
Tnf/Actb

C
UN
BMDMs
Irf3/Actb

D
+LPS (4h)
BMDMs
Ifit3b/Actb

+LPS (4h)
BMDMs
Irf3/Actb

E
Irf7
+4h LPS
BMDMs
elution/input normalized to NC
Irf7 primer set

— WT BMDMs IgG pulldown
— WT BMDMs RNA Pol II pulldown
— IFNAR KO BMDMs IgG pulldown
— IFNAR KO BMDMs RNA Pol II pulldown

F
BMDM, *Ifit3*
— 0h H4K20me1
— 4h H4K20me1
elution/input normalized to IgG
Ifit3 primer set

G
RAW 264.7, *Zbp1*
— 0h H4K20me1
— 4h H4K20me1
elution/input normalized to IgG
Zbp1 primers

Zbp1
Reverse Strand
TSS

S6

Figure S6: Supplemental data for Figure 6

A. Proteins identified as shared interactors of 3xFL-SRSF7, 3xFL-SRSF1, 3xFL-SRSF3, 3xFL-SRSF6, and 3xFL-SRSF9 in RAW MΦs at baseline by IP-LC/MS. **B.** RT-qPCR of *Tnf* in RAW MΦs transfected with siControl or si*Kmt5a* siRNA at baseline. **C.** RT-qPCR of *Irf3* in BMDMs transfected with siControl or si*Kmt5a* siRNA at baseline. **D.** As in C but 4h post-stimulation with 100 ng/ml LPS. **E.** RNA polymerase II ChIP-qPCR 4h post-stimulation with LPS in WT and *Ifnar* -/- BMDMs at the *Irf7* promoter **F.** H4K20me1 ChIP-qPCR at the *Ifit3* promoter in RAW MΦs at baseline or 4h post- stimulation with 100 ng/ml LPS. **G.** As in F at the *Zbp1* promoter. Right: Diagram of the *Zbp1* gene with tiling primers designed around the transcriptional start site (orange). Unless otherwise noted, data are expressed as a mean of three or more biological replicates with error bars depicting SEM. ChIP-qPCRs are shown as the mean of three or more technical replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

not impacted by loss of KMT5a (**Fig. S6B**), hinting at a privileged role for KMT5a in modulating

ISGs. Notably, STAT1 failed to associate with the *Irf7* promoter in *Kmt5a* knockdown RAW MΦs

(**Fig. 6D**), strongly supporting a role for KMT5a in promoting a pro-transcription chromatin envi-

ronment at *Irf7*. We also detected lower *Irf7* transcript levels both at rest (**Fig. 6E**) and at 4h post-

LPS treatment (**Fig. 6F**) when we knocked down KMT5a in BMDMs. Together, these data impli-

cate KMT5a in stimulating *Irf7* expression in macrophages.

Lastly, we sought to implicate the histone mark deposited by KMT5a, H4K20me1, in acti-

vating *Irf7* expression. We found that LPS treatment triggered significant accumulation of

H4K20me1 at the *Irf7* promoter in both RAW MΦs (**Fig. 6G**) and BMDMs (**Fig. 6H**) that was

concomitant with IFNAR-dependent transcriptional activation and recruitment of RNAPII (**Fig.**

S6E). LPS treatment also elevated H4K20me1 levels at ISGs like *Ifit3* and *Zbp1* (**Fig. S6F-G**).

Because we did not see SRSF7 enrichment at the promoters of these or other ISGs by ChIP-

qPCR or ChIP-seq (**Fig. 5L**), our data argue that H4K20me1 does not recruit SRSF7 to promot-

ers. Instead, our data support a model wherein interaction between SRSF7 and KMT5a confers

tight control over the chromatin landscape at *Irf7* promoter, which is required to maximize STAT1

recruitment and promote RNAPII elongation (**Fig. 6I**).

DISCUSSION

Because of their potential to activate large gene regulons, cells need to exert tight con-

trol over transcription factors themselves. This is particularly true for transcription factors that

activate inflammatory immune gene expression, which are subject to multiple layers of post-

translational regulation (cytoplasmic sequestration, phosphorylation, proteasomal degradation)

(Caamano and Hunter, 2002; Jefferies, 2019; Ning et al., 2011; Smale, 2012). Here, we de-

scribe a novel role for the splicing regulatory protein SRSF7 in stimulating the type I IFN re-

sponse by activating transcription of the *Irf7* gene. We demonstrate that enrichment of SRSF7

at the *Irf7* promoter facilitates association of the transcription factor STAT1 and promotes

RNAPII elongation and initiate expression of *Irf7*. SRSF7 accomplishes this by establishing a

pro-transcription chromatin landscape at the *Irf7* promoter via interaction with the H4K20me1

methyltransferase KMT5a. By controlling IRF7 abundance, SRSF7 plays a privileged role in reg-

ulating both antiviral (**Fig. 2Q-T**) and antibacterial (**Fig. 5H**) innate immune responses in macro-

phages. In addition to implicating SRSF7 in activating *Irf7* expression, this work highlights a criti-

cal and underappreciated role for both SRSF7 and IRF7 in maintaining tonic and early interferon

signaling in MΦs.

A major finding of this study relates to the ability of SRSF7—a canonical RNA binding
protein—to promote transcription of *Irf7*. Over the past two decades, the idea that transcription
and RNA processing occur in a linear, stepwise fashion has been repeatedly challenged. It is
now generally accepted that many steps in RNA processing (i.e. capping, pre-mRNA splicing,
RNA editing) occur co-transcriptionally while nascent transcripts are still bound to RNAPII. Ac-
cordingly, splicing can influence transcription and transcription can impact splicing (Tellier et al.,
2020). Several studies have linked SR proteins to RNAPII dynamics and co-transcriptional
splicing. SRSF2, for example, releases the cyclin-dependent kinase pTEFb from an inhibitory
7SK-containing complex to relieve stalled polymerase at promoters (Ji et al., 2013) and SRSF3
regulates exon inclusion via the C-terminal domain of RNAPII (de la Mata and Kornblihtt, 2006).
Although our data suggest an analogous role for SRSF7 in mediating transcription factor binding
and RNAPII elongation at the *Irf7* promoter, several aspects of this mechanism still need to be
explored.
One outstanding question relates to why we see SRSF7 at certain promoters, like *Irf7*,
but not others (**Fig. 6**). Owing to its ability to interact with histone modifying enzymes like
KMT5a, it is possible that the local chromatin environment dictates the specificity of SRSF7 pro-
moter enrichment. Our data demonstrate that H4K20me1 levels are significantly enhanced at
the promoters of innate genes including *Irf7* following LPS treatment, linking this mark to macro-
phage activation and innate immune responses for the first time. However, because H4K20me1
is enriched at innate genes like *Ifit3* and *Zbp1* where SRSF7 promoter association was not de-
tected (**Fig. S6F-G**), we currently do not favor a "chromatin recruits SRSF7" model. Instead, we
propose that specificity is dictated by sequences encoded in exons/introns of genes down-
stream of SRSF7-sensitive promoters, such that nascent RNAs serve as the platform for SRSF7
recruitment. The reported consensus sequence for SRSF7 is GA(C/T)GA(C/T) (Cavaloc et al.,
1999) and SRSF7 was shown to bind TGA(C/T)C sequences in its own pre-mRNA to regulate
poison cassette exon inclusion (Konigs et al., 2020). We can detect several potential SRSF7
bindings sites in the first coding exon of *Irf7* (TGACC). Thus, it is possible that SRSF7's associ-
ation at the *Irf7* promoter occurs by virtue of co-transcriptional binding of exonic splicing enhanc-
ers (ESEs) in nascent transcripts.
It is also conceivable that SRSF7 associates with the *Irf7* promoter in an RNA-independ-
ent, DNA-dependent fashion. SRSF7 is the only SR protein that encodes a Zn-knuckle, and to
date, this domain has been shown to enhance the RNA binding activity of SRSF7 (Cavaloc et
al., 1999; Cavaloc et al., 1994; Wang et al., 2021)). However, in other proteins, for example
TAF1, the largest subunit of the general transcription factor TFIIID, a Zn knuckle encourages

DNA binding and promoter occupancy (Curran et al., 2018). It is tempting to speculate that the

Zn finger domain in SRSF7 could promote its association directly with DNA. Future experiments

designed to implicate the Zn knuckle in SRSF7 association with the *Irf7* promoter and/or test the

ability of SRSF7 to directly bind DNA will help resolve the molecular mechanisms of SRSF7 pro-

moter enrichment.

In addition to linking SRSF7 to *Irf7* expression, our studies revealed an unexpected role

for SRSF7 in controlling tonic type I IFN expression in resting RAW cells (**Fig. 2A, E**) and MEFs

(**Fig. 2M-P**, time 0). These observations challenge the paradigm that IRF7 needs to be dramati-

cally upregulated before it can contribute to type I IFN responses. Indeed, differences in ISG ex-

pression were more robust in IRF7 KO RAW MΦs at rest than after LPS stimulation (**Fig. S4B**).

These findings suggest that IRF7 is in fact the major mediator of the type I IFN response in rest-

ing macrophages, but *both* IRF3 and IRF7 contribute once they are activated downstream of

pattern recognition receptor engagement. To date, the role for IRF7 in maintaining tonic type I

IFN signaling has not been fully explored. Our RNA-seq data identify a specific regulon of ISGs

that is particularly sensitive to loss of IRF7 at baseline (**Fig. S3F, G**). Among the most downreg-

ulated genes in resting *Irf7* KO macrophages are several encoding guanylate binding proteins

(GBPs) and members of the oligoadenylate synthetase (OAS) family. Regulated expression of

these factors at baseline may reflect their role in directly mediating anti-viral and anti-bacterial

killing, with low level expression of these ISGs in resting cells providing an early advantage

upon infection. By demonstrating a role for IRF7 in maintaining tonic levels of these and other

ISGs, our results hint at additional roles for IRF7 in cellular homeostasis and disease that have

not been appreciated to date.

Consistent with this idea, growing evidence suggests that IRF7 expression in immune

cells is subject to multiple levels of control mediated by RBPs. Both mouse and human IRF7

genes encode very small introns (<100nt), an unusual adaptation in mammalian genomes. Reg-

ulated *IRF7* intron retention has been reported in THP-1 cells, wherein intron 2 is efficiently re-

moved upon monocyte-to-macrophage differentiation but not in resting monocytes (Green et al.,

2020). In response to treatment with TNF-α and IFN-α, as well as polyI:C transfection, regulated

removal of IRF7 intron 4 has been reported and was shown to be controlled by the RBP BUD13,

which is enriched on *Irf7* transcripts in BMDMs (Frankiw et al., 2019). Such regulation is pro-

posed to provide innate cells a way to limit the inappropriate expression of genes with high in-

flammatory potential like IRF7, while leaving cells poised to upregulate these genes when cer-

tain signals are received. We did not observe differences in the number of intron reads in *Srsf7*

KD vs. SCR RAW cells (**Fig. 3H**), nor did the MAJIQ algorithm identify SRSF7-dependent

alternative splicing changes in *Irf7* (**Fig. 1**). We propose that intron retention/alternative splicing

of *Irf7* by factors like BUD13 adds an additional layer of regulation on top of the transcriptional

control conferred by SRSF7. The fact that IRF7 is subject to regulation at multiple nodes argues

that it plays a larger than appreciated role in mediating inflammatory responses in immune cells.

Together, these previously published studies and our work presented here, argue that IRF7 is

required for basal maintenance and ramping up of type I interferon responses and highlights a

preeminent role for RNA binding proteins in regulating innate immune outcomes.

ACKNOWLEDGEMENTS

We would like the thank members of Patrick and Watson labs for critical reading of the

manuscript and invaluable feedback. We would also like to thank Phillip West and members of

his lab at Texas A&M School of Medicine for sharing reagents and protocols related to VSV in-

fection. Thanks to Susan Carpenter and her lab for sharing reagents and protocols related to

iBMDMs. and to the Genomic and RNA Profiling Core (GARP) at Baylor College of Medicine for

RNA-seq and ChIP-seq library generation and sequencing. Funding was provided by

NIH/NIGMS to KP (R35GM133720), NIH/NIAID to ROW and KP (R01AI155621), and

NIH/NIGMS to HMS (F31GM143893).

DECLARATION OF INTERESTS

The authors have no competing interests.

Materials and Methods

Cell culture

RAW 264.7 macrophage-like cells (RAW MΦs) (TIB-71™ ATCC) (originally isolated from male
BALB/c mice), L929 ISRE (Hoffpauir et al., 2020), *Irf7* KO RAW MΦs (this paper), and Lenti-X
cells (TaKaRa Bio) were cultured at 37 °C with a humidified atmosphere of 5% CO_2 in complete
media containing high glucose, DMEM (ThermoFisher) with 10% FBS (Millipore) 0.2% HEPES
(Thermo Fisher).

shRNA knockdowns

For RAW MΦs stably expressing scramble knockdown and *Srsf7* knockdown, Lenti-X cells were
transfected with a pSICO scramble non-targeting shRNA construct and pSICO *Srsf7* shRNA
constructs targeted at exon 3 and exon 4 of *Srsf7* using Polyjet (SignaGen Laboratories). Virus
was collected 24 and 48 h post transfection. RAW MΦs were transduced using Lipofectamine
2000 (Thermo Fischer). After 48 h, media was supplemented with Hygromycin (Invitrogen) to
select for cells containing the shRNA plasmid.

S. enterica (ser. Typhimurium)

Salmonella enterica serovar Typhimurium (SL1344) was obtained from Dr. Denise Monack,
Stanford. *S.* Typhimurium were streaked out on LB agar plates and incubated at 37°C overnight.

S. enterica (ser. Typhimurium) infections:

RAW MΦs were seeded in 12-well tissue culture-treated plates at a density of 7×10^5 cells per cell,
and cultures of *S.* Typhimurium were grown at 37°C in LB broth containing 0.3M NaCl 16h prior
to infection. Bacterial cultures were diluted 1:20 3h prior to infection to reach mid-log phase OD_{600}
0.6-0.8. Mid-log phase bacteria were washed 3x with PBS and pelleted at 5,000 rpm for 3 min.
Bacteria were then added to DMEM (Hyclone) with 10% FBS (Sigma) and 2% HEPES (HyClone)
at an MOI of 5 for transcripts and MOI of 1 for replication. Infected cells were spun at 1,000 rpm
for 5 min then incubated for 10 min at 37 °C prior to gentamycin washes adding fresh media. At
indicated times post infection, cells were harvested with TRIzol for RNA isolation described below.

M. tuberculosis

The Erdman strain was used for all *M. tuberculosis* (Mtb) infections as well as Mtb luxBCADE a
luciferase expressing strain. Low passage lab stocks were thawed for each experiment to

997   ensure virulence was preserved. Mtb was cultured in roller bottles at 37 °C in Middlebrook 7H9

998   broth (BD Biosciences) supplemented with 10% OADC (BD Biosciences), 0.5% glycerol

999   (Fisher), and 0.1% Tween-80 (Fisher). All work with Mtb was performed under Biosafety Level 3

1000   containment using procedures approved by the Texas A&M University Institutional Biosafety

1001   Committee.

1002

Mtb in vitro infections

To prepare the inoculum, bacteria were grown to mid log phase (OD 0.6–0.8), spun at low

speed (500 rcf) to remove clumps, and then pelleted and washed with PBS 2X. Resuspended

bacteria were sonicated and spun at low speed again to further remove clumps. Then Mtb was

then diluted in DMEM plus 10% horse serum (Gibco) and added to cells at a multiplicity of infec-

tion (MOI) of 10 for RNA and protein analysis, a MOI of 5 for cell death studies, and a MOI of 1

for bacterial growth assays. The day before the infection, RAW MΦs were plated on 12-well tis-

sue culture–treated plates at a density of 3×10^5 cells per well or plated in corning 96 well black

plates at 2.5×10^4 cells/well and allowed to rest overnight. Cells were spun with bacteria for 10

min at 1,000 rcf to synchronize infection, washed 2X with PBS, and then incubated in fresh me-

dia. Where applicable, RNA was harvested from infected cells using 0.25 ml TRIzol reagent at

each time point. For bacterial growth assays, RAW MΦs were plated on 12-well tissue culture–

treated plates at a density of 2.5×10^5 cells per well. Luminescence was read for *M. tuberculosis*

luxBCADE by lysing in 250 µl 0.5% Triton X-100 and dividing sample into duplicate wells of a

96-well white bottomed plate (Costar). Luminescence was measured and normalized to back-

ground using the luminescence feature of the INFINITE 200 PRO (Tecan) at 0, 48, 72 and 92 h

post infection.

1020

1021   Vesicular stomatitis virus

1022   Recombinant *Vesicular stomatitis* virus (VSV; Indiana serotype) containing a GFP reporter

1023   cloned downstream of the VSV G-glycoprotein (VSV-G/GFP) was obtained collaboratively from

1024   Dr. John Rose at Yale School of Medicine though Dr. A. Phillip West at Texas A&M Health Sci-

1025   ence Center.

1026

Vesicular stomatitis Virus (VSV) infections

RAW MΦs were seeded in 12-well tissue culture-treated plates at a density of 7×10^5 cells per

well 16h before infection. The next day cells were infected with VSV-GFP virus (Dalton and

Rose, 2001) at an MOI of 0.1 in serum-free DMEM (HyClone). After 1h of incubation with media

containing virus, supernatant was removed, and fresh DMEM plus 10% FBS (Millipore) was

added to each well. At indicated times post infection, cells were harvested with TRIzol for RNA

isolation described below.

1034

M. musculus

Mice used in this study were C57BL6/J (Stock #00064) initially purchased from Jackson Labra-

tory and afterward maintained with filial breeding. All mice used in experiments were compared

to age- and sex- matched controls. Littermates were used for experiments. Mice used to gener-

ate BMDMs were males between 10-16 weeks old. Embryos used to make primary MEFs were

14.5 days post-coitum. All animals were housed, bred, and studied at Texas A&M Health Sci-

ence Center under approved Institutional Care and Use Committee guidelines. All experiments

for this study were reviewed and approved by the Texas A&M University Institutional Animal

Care and Use Committee (AUP# 2019–0083). Mice were fed 4% standard chow and were kept

on a 12 h light/dark cycle and provided food and water *ad libitum*. Mice were group housed

(maximum 5 per cage) by sex on ventilated racks in temperature-controlled rooms.

1046

Primary cell culture

Bone marrow derived macrophages (BMDMs) were differentiated from bone marrow (BM) cells

isolated by washing mouse femurs with 10 ml DMEM 1 mM sodium pyruvate. Cells were then

centrifuged for 5 min at 400 rcf and resuspended in BMDM media (DMEM, 20% FBS (Millipore),

1 mM sodium pyruvate (Lonza), 10% MCSF conditioned media (Watson lab)). BM cells were

counted and plated at 5×10^6 cells per 15 cm non-tissue culture treated dishes in 30 ml complete

BMDM media. Cells were fed with an additional 15 ml of BMDM media on day 3. Cells were har-

vested on day 7 with 1X PBS EDTA (Lonza).

1055

Mouse embryonic fibroblasts (MEFs) were isolated from embryos. Briefly, embryos were dis-

sected from yolk sacs, washed two times with cold 1X PBS, decapitated, and stripped of perito-

neal contents. Headless embryos were disaggregated in cold 0.05% trypsin-EDTA (Lonza) and

incubated on ice for 20 min., followed by incubation at 37 °C for an additional 20 min. Cells were

then DNase treated with 4 ml disaggregation media (DMEM, 10% FBS, 100 µg/ml DNASE I

(Worthington)) for 20 min at 37 °C. Isolated supernatants were spun down at 1000 rpm for 5

min. Cells were resuspended in complete MEF media (DMEM, 10% FBS, 1 mM sodium py-

ruvate), and plated in 15 cm tissue culture-treated dishes 1 dish per embryo in 25 ml of media.

MEFs were allowed to expand for 2-3 days before harvest with 0.05% trypsin-EDTA (Lonza).

iBMDMs were immortalized with J2 virus to generate iBMDMs.

1066

<u>siRNA knockdowns in MEFs</u>

Knockdown of mRNA transcripts was performed by plating $3x10^6$ MEFs in 10 cm plates and

rested overnight. The following day complete media was replaced with 5 ml fresh complete me-

dia 30 min prior to transfection. Cells were transfected using Fugene SI reagent 10 µM of siRNA

stock against either *Srsf7* (ThermoFisher, s105011) or negative control, Silencer® Select Nega-

tive Control #1 (4390843). Cells were incubated for 48h in transfection media at 37 °C with 5%

CO_2 prior to downstream experiments.

1074

<u>siRNA knockdowns in BMDMs</u>

The day prior to transfection, $6x10^5$ BMDMs were seeded into 12-well plates. The day of trans-

fection, 30nM of siRNA against *Srsf7* (ThermoFisher, s105011), against *Kmt5a/Setd8* (Ther-

moFisher, s85986) or negative control, Silencer® Select Negative Control #1 (4390843) were

mixed with 3 µl of LipoRNAiMAX (13778150, Thermo Fisher Scientific), incubated for 10m at

room temperature, and then added into wells with 500 µl plain DMEM. Transfection proceeded

for 6h and then wells were replenished with complete media.

1082

<u>Tetracycline inducible cell line generation</u>

For generation of tetracycline inducible RAW MΦs, pLenti CMV rtTA3 Blast (Addgene w756-1)

stably expressing clonal RAW MΦs were transduced with pLenti CMV Puro DEST (Addgene

w118-1) constructs containing 3xFL-GFP, 3xFL-SRSF7, and al 3xFL-SRSF7 phosphorylation

mutants. After 48h, construct containing cells were selected through addition of puromycin (Invi-

vogen). 1 µg/ml doxycycline (Sigma-Aldrich) treatment was used to activate construct expres-

sion.

1090

<u>Cell stimulation for immune activation</u>

$7.5x10^5$ RAW MΦs were plated on 12-well tissue-culture treated plates 16h before cellular stim-

ulation assays. The next day, cells were treated with lipopolysaccharide (LPS) from *E. coli* (Invi-

vogen) at 100 ng/ml, or ISD (IDT, annealed in house) at 1 µg/ml, or IFN-β (PBL Assay Science)

at 200 I/U per ml for the respective time points. Cells were collected for RNA isolation using Tri-

zol reagent.

1097

<u>*Srsf7* KD mRNA sequencing and analysis</u>

As reported in Wagner, Scott and West et al. 2021, RNA-seq analysis was performed on RAW

MΦs containing shRNA knockdowns of *Srsf6*, *Srsf2*, *Srsf1*, *Srsf7*, and *Srsf9* compared to SCR

control with biological triplicates of each cell line. RNA-Seq and library preparation was per-

formed by Texas A&M AgriLife Genomics and Bioinformatics Service. RNA samples were se-

quenced on Illumina 4000 using 2 x 150-bp paired-end reads. Raw reads were filtered and

trimmed and Fastq data was mapped to the Mus musculus Reference genome (RefSeq) using

CLC Genomics Workbench 8.0.1. Differential expression analyses were performed using CLC

Genomics Workbench. Relative transcript expression was calculated by counting Reads Per

Kilobase of exon model per Million mapped reads (RPKM). Statistical significance was deter-

mined by the EDGE test via CLC Genomics Workbench. Differentially expressed genes (DEGs)

were selected as those with p value threshold < 0.05. Critical pathways affected by SR protein

knockdown compared to control RAW MΦs, were analyzed by canonical pathway analysis using

Ingenuity Pathway Analysis software from QIAGEN Bioinformatics. Genes that were differen-

tially expressed with a p-value < 0.05 from our RNA-SEQ analysis were used as input. Inte-

grated genomics viewer (IGV) visualization tracks were generated using the RNA-SEQ filtered

and trimmed reads.

1115

Irf7 KO mRNA sequencing and analysis

RNA- seq analysis was performed on control GFP KO or *Irf7* KO1 RAW MΦs in biological tripli-

cate of each cell line. Data was analyzed by ROSALIND® (https://rosalind.bio/), with a

HyperScale architecture developed by ROSALIND, Inc. (San Diego, CA). Reads were trimmed

using cutadapt. Quality scores were assessed using FastQC. Reads were aligned to the Mus

musculus genome build GRCm39 using STAR[3]. Individual sample reads were quantified using

HTseq[4] and normalized via Relative Log Expression (RLE) using DESeq2 R library. Read Distri-

bution percentages, violin plots, identity heatmaps, and sample MDS plots were generated as

part of the QC step using RSeQC. DEseq2 was also used to calculate fold changes and p-val-

ues and perform optional covariate correction. Clustering of genes for the final heatmap of dif-

ferentially expressed genes was done using the PAM (Partitioning Around Medoids) method us-

ing the fpc R library. Hypergeometric distribution was used to analyze the enrichment of path-

ways, gene ontology, domain structure, and other ontologies. The topGO R library was used to

determine local similarities and dependencies between GO terms in order to perform Elim prun-

ing correction.

1131

Alternative splicing analysis

Modeling Alternative Junction Inclusion Quantification (MAJIQ) and the Voila visualization pack-

age were used to model alternative splicing events with default parameters described by

Vaquero-Garcia et al., 2016. Junction-spanning reads were used to construct splice graphs for

transcripts by using the RefSeq annotation supplemented with de-novo (junctions not in the Ref-

Seq transcriptome database but had sufficient evidence in the RNAseq data) detected junctions.

Resulting splice graphs were analyzed for all identified local splice variations (LSVs). MAJIQ

then quantified expected percent spliced in (PSI) value in control and knockdown samples for

every junction in each LSV. Resulted viewed on VOILA were filtered for highest confidence

change in LSVs (one or more junctions had at least a 95% probability of expected dPSI of at

least an absolute value of 0.2 PSI units between control and knockdown.

1143

RNA isolation and qPCR analysis

For transcript analysis, cells and tissue were harvested in TRIzol and RNA was isolated using

Direct-zol RNA Miniprep kits (Zymo Research) with 1 h DNase treatment. cDNA was synthe-

sized with iScript cDNA Synthesis Kit (Bio-Rad). CDNA was diluted as appropriate for treatment,

and a pool of cDNA was used to make a 1:3 standard curve with each standard sample diluted

1:5 to produce a linear curve. RT-qPCR was performed using Power-Up SYBR Green Master

Mix (Thermo Fisher) using a Quant Studio Flex6 (Applied Biosystems). Samples were run in

triplicate wells in a 384-well plate. Averages of the raw values were normalized to average val-

ues for the same sample with the control gene, *Actb*.

1153

Semiquantitative qPCR analysis

cDNA was synthesized by iScript cDNA Synthesis Kit (Bio-Rad) using an extended 1h amplifica-

tion. Q5 high fidelity 2X Master mix (New England Biolabs) was used for PCR amplification us-

ing targeted primers. Loading dye was added to PCR products and samples were run on 2%

agarose gel containing ethidium bromide at 100 volts for 1 h. Gels were imaged on LiCOR Od-

yssey Fc Dual-Mode Imaging System and bands were quantified.

1160

Chromatin immunoprepitation

$6x10^6$ RAW MΦs or BMDMs were seeded in a 10 cm dish, and 16h later stimulated for the indi-

cated time. After stimulation cells were scraped, transferred to a 50 ml conical and fixed with 1%

(v/v) formaldehyde for 10 min with gentle shaking. Formaldehyde was quenched with 0.125 M

glycine for 5 min. Cells were pelleted and washed 1x with PBS, and then cells were lysed in 1

ml of ChIP lysis buffer (150 mM NaCl, 5 mM EDTA, 0.5% NP-40 and 1% Triton X-100, 50 mM

Tris-HCl, and protease/phosphatase inhibitor (added immediately before lysis) pH 7.5). Lysates

were flash frozen, or ChIP proceeded by sonicating with for 40 min (30 sec on/off 10 min cycles)

in a Bioruptor UCD-200 (Diagenode). Lysates were cleared and 10% taken for input samples.

For RNA polymerase II, Y701 STAT1, IgG (isotype control) or H4K20me1 samples: 2 µg

of the anti-pol II (Active Motif) or 2 µg of isotype control (IgG, Cell Signaling), 5ul Y701 STAT1

(Cell Signaling), or 2 µg H4K20me1 (Invitrogen) (were incubated overnight at 4 with rotation.

The following day, Pierce beads (Invitrogen) were washed 3x in ChIP buffer, blocked for 30 min

with 100 µg salmon sperm DNA (Invitrogen) and 0.5 mg BSA, and washed once more with ChIP

lysis buffer. Blocked beads were incubated with cleared chromatin immunocomplexes for 1h at

4 °C with rotation. Beads were washed five times with ChIP buffer and then eluted with 250 µl of

elution buffer (1% SDS and 0.1 M NaHCO$_3$) at 65 °C for 15 mins. The eluate and input were

centrifuged and reverse the cross-links by adding 0.4 M NaCl for overnight at 55 °C. The next

day, 1 µl of RNASE A (10 mg/ml, ThermoFisher) and 2 µl proteinase K (20 mg/ml, Ther-

moFisher) were added to the samples and incubated for 1h at 55 °C while shaking. Samples

were then subjected to Genomic Clean up and Concentrator (Zymo) to extract DNA. Purified in-

put and IP DNA were analyzed by qRT-PCR using primers specific for the promoter region of

target genes. Results were normalized to input. Data is presented as fold enrichment relative to

intergenic control. Flag ChIP: As above except cleared lysates were added to blocked (as

above) M2 anti-3xFLAG beads and protocol proceeded as above.

1186

3x-FLAG ChIP-seq

Data was analyzed by ROSALIND® (https://rosalind.bio/), with a HyperScale architecture devel-

oped by ROSALIND, Inc. (San Diego, CA). Reads were trimmed using cutadapt[1]. Quality scores

were assessed using FastQC[2]. Reads were aligned to the Mus musculus genome build mm10

using bowtie2[3]. Per-sample quality assessment plots were generated with HOMER[4] and Mosa-

ics[5]. Peaks were called using MACS2[6] (with input/IgG controls background subtracted, if pro-

vided). Peak overlaps and differential binding were calculated using the DiffBind R library[7]. Dif-

ferential binding was calculated at gene promoter sites. Read distribution percentages, identity

heatmaps, and FRiP plots were generated as part of the QC step using ChIPQC R library[8] and

HOMER. HOMER was also used to generate known and de novo motifs and perform functional

enrichment analysis of pathways, gene ontology, domain structure and other ontologies.

1198

CRISPR/Cas9 KO of *Irf7*

Guide RNAs were generated to target the 3rd exon of *Irf7* using the Broad online tool

(https://portals.broadinstitute.org/gppx/crispick/public) and synthesized by IDT. Primers were

cloned into digested vector by phosphorylating, annealing, and ligating primers into LentiGuide-

Puro (Addgene, plasmid 52963). Colonies were validated by Sanger sequencing using

pLK0.1/hU6 promoter primer (Eton Biosciences, San Diego, CA). Lentivirus with sgRNAs were

produced and used to transduce RAW MΦs stably expressing FL-Cas9 (Addgene, plasmid

52962). After selection with 5 μg/ml puromycin knockout efficiency was analyzed by immunoblot

and then sequencing for mutants by amplifying a 300 bp region of genomic DNA around exon 3.

These PCR fragments were sequenced (as above) and compared to controls using TIDE analy-

sis. The two best, most efficient sgRNA populations were clonally selected, expanded, and as-

sayed for clones with total KO of IRF7 by immunoblot.

1211

<u>Protein quantification by immunoblot</u>

Cells were washed with PBS and lysed in 1X RIPA buffer with protease and phosphatase EDTA

free inhibitors (Thermo Scientific), with the addition of 1 U/ml Benzonase (Millipore) to degrade

genomic DNA. Proteins were separated by SDS-PAGE on Any kD mini-PROTEAN TGX precast

gel (Bio-Rad) and transferred to 0.45μm nitrocellulose membranes (GE Healthcare). Mem-

branes were blocked for 1h at RT in TBS +5% BSA. Blots were incubated overnight at 4 °C with

the following primary antibodies: β-ACTIN (Abcam, 1:5000), Y701 STAT1 (Cell Signaling,

1:1000), STAT1 (Cell Signaling, 1:1000), IRF7 (Cell Signaling, 1:1000), SRSF7 (Bethyl 1:3000)

ZBP1 (Santa Cruz Biotechnologies, 1:1000), p-IRF3(S396) (Cell Signaling, 1:1000); IRF3

(Bethyl, 1:1000); ANTI-FLAG M2 (Sigma Aldrich, 1:5000); Membranes were washed 3 x 5 min

in PBS-Tween20 and incubated with appropriate secondary antibodies (LI-COR) for 1h at RT

prior to imaging on a LiCOR Odyssey Fc Dual-Mode Imaging System.

1224

1225   <u>Quantitation and Statistical Analysis</u>

1226   Statistical analysis of data was performed using GraphPad Prism software. Two-tailed unpaired

1227   Student's t tests were used for statistical analyses, and unless otherwise noted, all results are

1228   representative of at least three biological samples (mean +/- SEM (n = 3 per group)).

1229

1230

1231

REFERENCES

Banks, D.A., Ahlbrand, S.E., Hughitt, V.K., Shah, S., Mayer-Barber, K.D., Vogel, S.N., El-Sayed, N.M., and Briken, V. (2019). Mycobacterium tuberculosis Inhibits Autocrine Type I IFN Signaling to Increase Intracellular Survival. J Immunol *202*, 2348-2359.

Barrat, F.J., Crow, M.K., and Ivashkiv, L.B. (2019). Interferon target-gene expression and epigenomic signatures in health and disease. Nat Immunol *20*, 1574-1583.

Barski, A., Cuddapah, S., Cui, K., Roh, T.Y., Schones, D.E., Wang, Z., Wei, G., Chepelev, I., and Zhao, K. (2007). High-resolution profiling of histone methylations in the human genome. Cell *129*, 823-837.

Bell, S.L., Lopez, K.L., Cox, J.S., Patrick, K.L., and Watson, R.O. (2021). Galectin-8 Senses Phagosomal Damage and Recruits Selective Autophagy Adapter TAX1BP1 To Control Mycobacterium tuberculosis Infection in Macrophages. mBio *12*, e0187120.

Boguslawska, J., Sokol, E., Rybicka, B., Czubaty, A., Rodzik, K., and Piekielko-Witkowska, A. (2016). microRNAs target SRSF7 splicing factor to modulate the expression of osteopontin splice variants in renal cancer cells. Gene *595*, 142-149.

Budzik, J.M., Swaney, D.L., Jimenez-Morales, D., Johnson, J.R., Garelis, N.E., Repasy, T., Roberts, A.W., Popov, L.M., Parry, T.J., Pratt, D.*, et al.* (2020). Dynamic post-translational modification profiling of Mycobacterium tuberculosis-infected primary macrophages. Elife *9*.

Caamano, J., and Hunter, C.A. (2002). NF-kappaB family of transcription factors: central regulators of innate and adaptive immune functions. Clin Microbiol Rev *15*, 414-429.

Cavaloc, Y., Bourgeois, C.F., Kister, L., and Stevenin, J. (1999). The splicing factors 9G8 and SRp20 transactivate splicing through different and specific enhancers. RNA *5*, 468-483.

Cavaloc, Y., Popielarz, M., Fuchs, J.P., Gattoni, R., and Stevenin, J. (1994). Characterization and cloning of the human splicing factor 9G8: a novel 35 kDa factor of the serine/arginine protein family. EMBO J *13*, 2639-2649.

Cun, Y., An, S., Zheng, H., Lan, J., Chen, W., Luo, W., Yao, C., Li, X., Huang, X., Sun, X.*, et al.* (2021). Specific Regulation of m(6)A by SRSF7 Promotes the Progression of Glioblastoma. Genomics Proteomics Bioinformatics.

Curran, E.C., Wang, H., Hinds, T.R., Zheng, N., and Wang, E.H. (2018). Zinc knuckle of TAF1 is a DNA binding module critical for TFIID promoter occupancy. Sci Rep *8*, 4630.

de la Mata, M., and Kornblihtt, A.R. (2006). RNA polymerase II C-terminal domain mediates regulation of alternative splicing by SRp20. Nat Struct Mol Biol *13*, 973-980.

Fitzgerald, K.A., Rowe, D.C., Barnes, B.J., Caffrey, D.R., Visintin, A., Latz, E., Monks, B., Pitha,
P.M., and Golenbock, D.T. (2003). LPS-TLR4 signaling to IRF-3/7 and NF-kappaB involves the
toll adapters TRAM and TRIF. J Exp Med *198*, 1043-1055.
Frankiw, L., Majumdar, D., Burns, C., Vlach, L., Moradian, A., Sweredoski, M.J., and Baltimore,
D. (2019). BUD13 Promotes a Type I Interferon Response by Countering Intron Retention in
Irf7. Mol Cell *73*, 803-814 e806.
Fu, Y., and Wang, Y. (2018). SRSF7 knockdown promotes apoptosis of colon and lung cancer
cells. Oncol Lett *15*, 5545-5552.
Gao, D., Ciancanelli, M.J., Zhang, P., Harschnitz, O., Bondet, V., Hasek, M., Chen, J., Mu, X.,
Itan, Y., Cobat, A.*, et al.* (2021). TLR3 controls constitutive IFN-beta antiviral immunity in human
fibroblasts and cortical neurons. J Clin Invest *131*.
Gough, D.J., Messina, N.L., Clarke, C.J., Johnstone, R.W., and Levy, D.E. (2012). Constitutive
type I interferon modulates homeostatic balance through tonic signaling. Immunity *36*, 166-174.
Green, I.D., Pinello, N., Song, R., Lee, Q., Halstead, J.M., Kwok, C.T., Wong, A.C.H., Nair, S.S.,
Clark, S.J., Roediger, B.*, et al.* (2020). Macrophage development and activation involve
coordinated intron retention in key inflammatory regulators. Nucleic Acids Res *48*, 6513-6529.
Gu, J., Shi, J., Wu, S., Jin, N., Qian, W., Zhou, J., Iqbal, I.G., Iqbal, K., Gong, C.X., and Liu, F.
(2012). Cyclic AMP-dependent protein kinase regulates 9G8-mediated alternative splicing of tau
exon 10. FEBS Lett *586*, 2239-2244.
Gu, L., Casserly, D., Brady, G., Carpenter, S., Bracken, A.P., Fitzgerald, K.A., Unterholzner, L.,
and Bowie, A.G. (2022). Myeloid cell nuclear differentiation antigen controls the pathogen-
stimulated type I interferon cascade in human monocytes by transcriptional regulation of IRF7.
Nat Commun *13*, 14.
Hoffpauir, C.T., Bell, S.L., West, K.O., Jing, T., Wagner, A.R., Torres-Odio, S., Cox, J.S., West,
A.P., Li, P., Patrick, K.L.*, et al.* (2020). TRIM14 Is a Key Regulator of the Type I IFN Response
during Mycobacterium tuberculosis Infection. J Immunol *205*, 153-167.
Honda, K., and Taniguchi, T. (2006). IRFs: master regulators of signalling by Toll-like receptors
and cytosolic pattern-recognition receptors. Nat Rev Immunol *6*, 644-658.
Honda, K., Yanai, H., Negishi, H., Asagiri, M., Sato, M., Mizutani, T., Shimada, N., Ohba, Y.,
Takaoka, A., Yoshida, N.*, et al.* (2005). IRF-7 is the master regulator of type-I interferon-
dependent immune responses. Nature *434*, 772-777.
Huang, R., Grishagin, I., Wang, Y., Zhao, T., Greene, J., Obenauer, J.C., Ngan, D., Nguyen,
D.T., Guha, R., Jadhav, A.*, et al.* (2019). The NCATS BioPlanet - An Integrated Platform for

Exploring the Universe of Cellular Signaling Pathways for Toxicology, Systems Biology, and

Chemical Genomics. Front Pharmacol *10*, 445.

Jefferies, C.A. (2019). Regulating IRFs in IFN Driven Disease. Front Immunol *10*, 325.

Jeong, S. (2017). SR Proteins: Binders, Regulators, and Connectors of RNA. Mol Cells *40*, 1-9.

Ji, X., Zhou, Y., Pandit, S., Huang, J., Li, H., Lin, C.Y., Xiao, R., Burge, C.B., and Fu, X.D.

(2013). SR proteins collaborate with 7SK and promoter-associated nascent RNA to release

paused polymerase. Cell *153*, 855-868.

Kadota, Y., Jam, F.A., Yukiue, H., Terakado, I., Morimune, T., Tano, A., Tanaka, Y., Akahane,

S., Fukumura, M., Tooyama, I.*, et al.* (2020). Srsf7 Establishes the Juvenile Transcriptome

through Age-Dependent Alternative Splicing in Mice. iScience *23*, 100929.

Kajigaya, H., Ishibashi, T., Hayashi, A., Yamaguchi, Y., and Baba, H. (2016). Concentration of

neddylation-related molecules in paranodal myelin of the peripheral nervous system. Proc Jpn

Acad Ser B Phys Biol Sci *92*, 56-68.

Katsuyama, T., and Moulton, V.R. (2021). Splicing factor SRSF1 is indispensable for regulatory

T cell homeostasis and function. Cell Rep *36*, 109339.

Konigs, V., de Oliveira Freitas Machado, C., Arnold, B., Blumel, N., Solovyeva, A., Lobbert, S.,

Schafranek, M., Ruiz De Los Mozos, I., Wittig, I., McNicoll, F.*, et al.* (2020). SRSF7 maintains its

homeostasis through the expression of Split-ORFs and nuclear body assembly. Nat Struct Mol

Biol *27*, 260-273.

Lemaire, R., Winne, A., Sarkissian, M., and Lafyatis, R. (1999). SF2 and SRp55 regulation of

CD45 exon 4 skipping during T cell activation. Eur J Immunol *29*, 823-837.

Loomis, R.J., Naoe, Y., Parker, J.B., Savic, V., Bozovsky, M.R., Macfarlan, T., Manley, J.L., and

Chakravarti, D. (2009). Chromatin binding of SRp20 and ASF/SF2 and dissociation from mitotic

chromosomes is modulated by histone H3 serine 10 phosphorylation. Mol Cell *33*, 450-461.

Luco, R.F., Allo, M., Schor, I.E., Kornblihtt, A.R., and Misteli, T. (2011). Epigenetics in

alternative pre-mRNA splicing. Cell *144*, 16-26.

Luco, R.F., Pan, Q., Tominaga, K., Blencowe, B.J., Pereira-Smith, O.M., and Misteli, T. (2010).

Regulation of alternative splicing by histone modifications. Science *327*, 996-1000.

MacMicking, J.D., North, R.J., LaCourse, R., Mudgett, J.S., Shah, S.K., and Nathan, C.F.

(1997). Identification of nitric oxide synthase as a protective locus against tuberculosis. Proc

Natl Acad Sci U S A *94*, 5243-5248.

MacMicking, J.D., Taylor, G.A., and McKinney, J.D. (2003). Immune control of tuberculosis by

IFN-gamma-inducible LRG-47. Science *302*, 654-659.

Misteli, T., Caceres, J.F., Clement, J.Q., Krainer, A.R., Wilkinson, M.F., and Spector, D.L.

(1998). Serine phosphorylation of SR proteins is required for their recruitment to sites of

transcription in vivo. J Cell Biol *143*, 297-307.

Muller-McNicoll, M., Botti, V., de Jesus Domingues, A.M., Brandl, H., Schwich, O.D., Steiner,

M.C., Curk, T., Poser, I., Zarnack, K., and Neugebauer, K.M. (2016). SR proteins are NXF1

adaptors that link alternative RNA processing to mRNA export. Genes Dev *30*, 553-566.

Nikolaou, K.C., Moulos, P., Harokopos, V., Chalepakis, G., and Talianidis, I. (2017). Kmt5a

Controls Hepatic Metabolic Pathways by Facilitating RNA Pol II Release from Promoter-

Proximal Regions. Cell Rep *20*, 909-922.

Ning, S., Pagano, J.S., and Barber, G.N. (2011). IRF7: activation, regulation, modification and

function. Genes Immun *12*, 399-414.

Pandey, A., Ding, S.L., Qin, Q.M., Gupta, R., Gomez, G., Lin, F., Feng, X., Fachini da Costa, L.,

Chaki, S.P., Katepalli, M*., et al.* (2017). Global Reprogramming of Host Kinase Signaling in

Response to Fungal Infection. Cell Host Microbe *21*, 637-649 e636.

Schoggins, J.W. (2019). Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol *6*,

567-584.

Schoggins, J.W., and Rice, C.M. (2011). Interferon-stimulated genes and their antiviral effector

functions. Curr Opin Virol *1*, 519-525.

Schoggins, J.W., Wilson, S.J., Panis, M., Murphy, M.Y., Jones, C.T., Bieniasz, P., and Rice,

C.M. (2011). A diverse range of gene products are effectors of the type I interferon antiviral

response. Nature *472*, 481-485.

Shin, C., Feng, Y., and Manley, J.L. (2004). Dephosphorylated SRp38 acts as a splicing

repressor in response to heat shock. Nature *427*, 553-558.

Sin, W.X., Yeong, J.P., Lim, T.J.F., Su, I.H., Connolly, J.E., and Chin, K.C. (2020). IRF-7

Mediates Type I IFN Responses in Endotoxin-Challenged Mice. Front Immunol *11*, 640.

Smale, S.T. (2012). Transcriptional regulation in the innate immune system. Curr Opin Immunol

24, 51-57.

Song, J., Su, Z.Z., and Shen, Q.M. (2020). Long non-coding RNA MALAT1 regulates

proliferation, apoptosis, migration and invasion via miR-374b-5p/SRSF7 axis in non-small cell

lung cancer. Eur Rev Med Pharmacol Sci *24*, 1853-1862.

Tellier, M., Maudlin, I., and Murphy, S. (2020). Transcription and splicing: A two-way street.

Wiley Interdiscip Rev RNA *11*, e1593.

ten Dam, G.B., Zilch, C.F., Wallace, D., Wieringa, B., Beverley, P.C., Poels, L.G., and Screaton,

G.R. (2000). Regulation of alternative splicing of CD45 by antagonistic effects of SR protein

splicing factors. J Immunol *164*, 5287-5295.

van Nuland, R., and Gozani, O. (2016). Histone H4 Lysine 20 (H4K20) Methylation, Expanding

the Signaling Potential of the Proteome One Methyl Moiety at a Time. Mol Cell Proteomics *15*,

755-764.

Vaquero-Garcia, J., Barrera, A., Gazzara, M.R., Gonzalez-Vallinas, J., Lahens, N.F.,

Hogenesch, J.B., Lynch, K.W., and Barash, Y. (2016). A new view of transcriptome complexity

and regulation through the lens of local splicing variations. Elife *5*, e11752.

Veloso, A., Kirkconnell, K.S., Magnuson, B., Biewen, B., Paulsen, M.T., Wilson, T.E., and

Ljungman, M. (2014). Rate of elongation by RNA polymerase II is associated with specific gene

features and epigenetic modifications. Genome Res *24*, 896-905.

Wagner, A.R., Scott, H.M., West, K.O., Vail, K.J., Fitzsimons, T.C., Coleman, A.K., Carter, K.E.,

Watson, R.O., and Patrick, K.L. (2021). Global Transcriptomics Uncovers Distinct Contributions

From Splicing Regulatory Proteins to the Macrophage Innate Immune Response. Front Immunol

12, 656885.

Wagner, A.R., Weindel, C.G., West, K.O., Scott, H.M., Watson, R.O., and Patrick, K.L. (2022).

SRSF6 balances mitochondrial-driven innate immune outcomes through alternative splicing of

BAX. Elife *11*.

Wagner, R.E., and Frye, M. (2021). Noncanonical functions of the serine-arginine-rich splicing

factor (SR) family of proteins in development and disease. Bioessays *43*, e2000242.

Wang, H.Y., Xu, X., Ding, J.H., Bermingham, J.R., Jr., and Fu, X.D. (2001). SC35 plays a role in

T cell development and alternative splicing of CD45. Mol Cell *7*, 331-342.

Wang, Y., Yu, Y., Pang, Y., Yu, H., Zhang, W., Zhao, X., and Yu, J. (2021). The distinct roles of

zinc finger CCHC-type (ZCCHC) superfamily proteins in the regulation of RNA metabolism.

RNA Biol *18*, 2107-2126.

Watson, R.O., Bell, S.L., MacDuff, D.A., Kimmey, J.M., Diner, E.J., Olivas, J., Vance, R.E.,

Stallings, C.L., Virgin, H.W., and Cox, J.S. (2015). The Cytosolic Sensor cGAS Detects

Mycobacterium tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. Cell

Host Microbe *17*, 811-819.

Xu, Q., Shi, J., Zhang, L., Sheng, Y., Zhang, Y., Chu, D., and Xu, A. (2022). Circ_0006006

facilitates non-small cell lung cancer progression by modulating miR-924/SRSF7 axis. J Gene

Med *24*, e3411.

Zhong, X.Y., Ding, J.H., Adams, J.A., Ghosh, G., and Fu, X.D. (2009). Regulation of SR protein

phosphorylation and alternative splicing by modulating kinetic interactions of SRPK1 with

molecular chaperones. Genes Dev *23*, 482-495.

Zhou, Z., and Fu, X.D. (2013). Regulation of splicing by SR proteins and SR protein-specific

kinases. Chromosoma *122*, 191-207.

1400