1

5 **Type I interferon expression requires transcriptional activation of IRF7 by the RNA binding**

6 **protein SRSF7**

8 Haley M. Scott[1], Mackenzie H. Smith[1], Aja K. Coleman[1], Allison R. Wagner[1,2], Robert O. Watson[1],

9 Kristin L. Patrick[1]*#

16 [1]Department of Microbial Pathogenesis and Immunology, Texas A&M University School of Medicine

17 [2]Department of Immunology, UT Southwestern Medical Center

23 *Corresponding author: kpatrick03@tamu.edu

24 #Lead contact

25 Phone: (979) 436-0354

26 Twitter: @The_PW_Lab

Abstract

While the signaling cascades and transcription factors that turn on expression of innate immune genes are well-characterized, the role of RNA binding proteins in activating and sustaining the innate immune response is poorly understood. Members of the serine/arginine-rich (SR) family of mRNA processing factors play diverse roles in transcription, pre-mRNA splicing, export, and translation. Transcriptomic analysis of murine macrophage cell lines revealed that one such SR protein, SRSF7, is required for optimal expression of interferon stimulated genes (ISGs) at rest and in response to a variety of innate immune stimuli in a splicing-independent fashion. We demonstrate that SRSF7 is necessary and sufficient to drive expression of interferon regulatory transcription factor 7 (IRF7), an important activator of the type I IFN response in resting and LPS-treated macrophages. By associating with the *Irf7* promoter, SRSF7 maximizes binding of the STAT1 transcription factor and promotes RNA polymerase II elongation. These studies define an unorthodox role for an SR protein in activating transcription and highlight the importance of RNA binding proteins in shaping the macrophage innate immune response to pathogens.

Introduction

In mammals, the SR (serine-arginine rich) family of proteins is comprised of >10 RNA

binding proteins that are broadly defined as regulators of alternative splicing. By virtue of distinct

RNA recognition motifs, SRs bind to unique exonic splicing enhancers to help direct the spliceo-

some to cis-splicing signals in target transcripts (Wagner and Frye, 2021). While best known for

their role in pre-mRNA splicing, SRs can participate in virtually any step of RNA processing, from

transcription all the way to translation (Jeong, 2017). To date, studies of SR function have mostly

been carried out in genetically tractable "model" cell lines (e.g. K562, HEK293T, and HepG2), but

there is a growing appreciation for the importance of SR proteins in responding to stress and

shaping the transcriptome in specialized cell types. Specialization of SR protein function has been

repeatedly linked to post-translational modification, which can alter the subcellular localization,

protein binding partners, and RNA recognition abilities of SRs, ultimately enabling them to carry

out different functions in different cell types (Colwill et al., 1996; Duncan et al., 1995; Gui et al.,

1994).

We are just beginning to appreciate the potential roles for SR protein functionalization in

immune cells. Recent proteomics analyses identified multiple SR proteins that are differentially

phosphorylated in macrophages upon infection with intracellular pathogens like the bacterium

Mycobacterium tuberculosis (Budzik et al., 2020) and the fungal pathogen *Cryptococcus neofor-*

mans (Pandey et al., 2017). Macrophages (MΦ) are innate immune cells that serve as the body's

first line of defense against pathogens. They express a variety of extracellular, endosomal, and

cytosolic pattern recognition receptors that detect pathogen- and damage-associated molecular

patterns (PAMPs and DAMPs) and transmit these "danger signals" to the nucleus though complex

signal transduction cascades. Activation of these cascades turns on expression of hundreds of

cytokines, chemokines, and antimicrobial mediators, many of which are transcriptionally and

translationally upregulated >100-fold in a matter of minutes (Nau et al., 2002). While a rapid and

robust innate immune response is needed to limit infection, a response that is too strong carries

risks (e.g. cytokine storm, autoimmunity, chronic inflammation, etc.). Therefore, MΦ gene expres-

sion needs to be tightly regulated. Despite their critical role in shaping mammalian transcriptomes,

molecular mechanisms though which SR proteins regulate macrophage gene expression—both

in terms of maintaining homeostatic levels of antimicrobial mediators and in inducing innate genes

following a stimulus—remain poorly understood.

We hypothesized that SR proteins with the potential to undergo modification downstream

of pattern recognition receptor engagement play a privileged role in tuning the macrophage re-

sponse to pathogens. We found that stable knockdown of several of these SR proteins in RAW

264.7 macrophage-like cells (RAW MΦs) profoundly altered the macrophage transcriptome in

resting and activated macrophages (4h post-*Salmonella* Typhimurium infection) (Wagner et al.,

2021). The absence of one of these proteins, SRSF7, generated some particularly interesting

phenotypes. SRSF7 (formerly known as 9G8) is a canonical member of the SR protein family. It

encodes an RNA recognition motif (RRM) and an eponymic SR domain. It is the only SR family

protein to also encode a CCHC-type Zn knuckle domain, which is thought to contribute to its RNA-

binding specificity (Cavaloc et al., 1999; Cavaloc et al., 1994; Wang et al., 2021)). To date, its

function is best studied in the context of mRNA export (Muller-McNicoll et al., 2016) and in con-

trolling its own expression levels via poison cassette exon inclusion (Konigs et al., 2020).

In RAW MΦs lacking SRSF7, we identified a large cohort of transcripts whose alternative

splicing (AS) was dependent on SRSF7. We also identified genes whose expression relied on

SRSF7 at rest and after *Salmonella* infection. Curiously, in both conditions, many SRSF7-im-

pacted genes fell into the category of interferon stimulated genes (ISGs). Studies suggest that as

much as 10% of the human genome has the potential to be regulated by one of the three types

of IFNs (I, II, III). Macrophages robustly express type I IFNs and hundreds of ISGs in response to

innate stimuli like lipopolysaccharide (LPS) and cytosolic nucleic acids. After ruling out a role for

SRSF7 in alternative splicing of these ISGs, we set out to elucidate the mechanism though which

SRSF7 controls the type I IFN response in macrophages. We report a novel role for SRSF7 in

controlling the abundance of interferon responsive factor 7 (IRF7), a critical player in activating

the type I IFN response (Ning et al., 2011). Specifically, we demonstrate that SRSF7 promotes

Irf7 transcription by enabling recruitment of signal transducer and activator of transcription 1

(STAT1) to the *Irf7* promoter and relieving RNAPII pausing. Together, these data identify SRSF7

as a transcriptional activator and provide important insight into how SR proteins can regulate

innate immune responses in macrophages in unexpected ways.

Results

SRSF7 control of alternative splicing is independent of macrophage activation state

Motivated by aforementioned phosphoproteomics analyses (Budzik et al., 2020; Pandey

et al., 2017) and previous studies of hnRNP M (West et al., 2019) and SRSF6 (Wagner et al.,

2022), we sought to identify additional splicing regulatory proteins that are functionally regulated

in the context of macrophage activation. To test the hypothesis that AS events might be distinct

between resting and activated macrophages, signifying functionalization of SRSF7, we generated

two *Srsf7* knockdown cell lines by transducing RAW 264.7 MΦs (*Srsf7* KD #1 and *Srsf7* KD #2)

with lentivirus containing shRNA hairpin constructs designed against the *Srsf7* 3'UTR. These cell

lines were selected alongside a scramble (SCR) control. We generated macrophage cell lines

with stable knockdown of *Srsf7* at the RNA (**Fig. 1A**) and protein level (**Fig. 1B**) with ~70-80%

efficiency. We performed RNA-seq (150 bp paired-end reads; Illumina 4000; ~20 million reads

per sample in triplicate) on RNA isolated from *Srsf7* KD #2 and SCR control cells that were resting

(uninfected; UN) or infected with the gram-negative bacterial pathogen *Salmonella enterica*

serovar Typhimurium (4h post-infection; +SAL). *Salmonella* induces robust innate immune re-

sponses in macrophages though TLR4, which activates NFκB downstream of the MyD88 adapter

and IRF3/IRF7 downstream of TRIF (Fitzgerald et al., 2003). Differential expression analysis was

conducted using the Qiagen CLC Genomics Workbench and quantification of alternative splicing

changes was carried out using MAJIQ (Modeling Alternative Junction Inclusion Quantification)

(Vaquero-Garcia et al., 2016). Consistent with SRSF7's designation as a splicing factor and pre-

vious studies implicating it in alternative splicing of *Eif4a2*, *Rbm7* (Kadota et al., 2020), and *Tau*

(Gu et al., 2012)), we identified 1857 total SRSF7-dependent AS events in resting RAW 264.7

MΦs and 1909 SRSF7-dependent events in RAW 264.7 MΦs 4h post-*Salmonella* infection (**Fig.**

1C) (p<0.05; >10% PSI). As expected for an SR protein that promotes exon recognition/inclusion,

most changes were at the level of exon skipping. Of the 964 genes for which local splicing varia-

tion was measured in resting MΦs and the 993 genes in *Salmonella*-infected macrophages, 493

of these genes overlapped (**Fig. 1D**). Notably, only a handful of the detected alternative splicing

changes occurred in genes that were also differentially expressed in *Srsf7* knockdown macro-

phages, suggesting that *Srsf7* impacts steady state RNA levels in a splicing-independent fashion

(**Fig. 1E**).

Based on these findings, we hypothesized that SRSF7 mediates distinct alternative splic-

ing events in the two macrophage states. To begin to test this, we prioritized a set of genes with

the most significant SRSF7-dependent AS changes (>20% PSI; confidence threshold 0.95) in

each condition (resting (UN) and 4h post-*Salmonella* infection (+SAL)) (**Fig. S1A**). Manual anno-

tation of these 36 genes did not reveal significant enrichment for any functional category and none

of the top AS events occurred in genes with characterized roles in the MΦ innate immune re-

sponse (**Fig. S1A**). To visualize and validate SRSF7-dependent AS in resting and *Salmonella*-

infected macrophages, we looked at exon 6 skipping in *Dcun1d2* (Ensembl Transcript ID:

ENSMUST00000045366.10), a gene associated with Cullin neddylation (Kajigaya et al., 2016).

MAJIQ measured significant enrichment of this exon skipping event in both untreated and *Salmo-*

nella-infected *Srsf7* knockdown MΦs (ΔPSI 39.2% in UN and 43.4% in +SAL) (**Fig. 1F-G**). To

confirm this event, we performed semi-quantitative RT-PCR, using RNA isolated from SCR, *Srsf7*

KD #1 and #2 MΦs, untreated and 4h post-stimulation with lipopolysaccharide (LPS), which acti-

vates TLR4 in MΦs in a manner analogous to *Salmonella* infection but provides tighter kinetic

Figure 1

Figure 1: SRSF7 controls alternative splicing in macrophages in an activation-independent fashion

A RT-qPCR of *Srsf7* in *Srsf7* KD RAW MΦs relative to SCR controls. **B.** Immunoblot of SRSF7 in shSCR, sh*Srsf7* #1, sh*Srsf7* #2 RAW 264.7 macrophages. Right, quantification. **C.** Local splicing variations (AS) in *Srsf7* RAW 264.7 cells compared to SCR at baseline (UN; left) and +*Salmonella* Typhimurium (4h) (+SAL; right). ΔPSI > 0.1 confidence threshold set at 0.90. **D.** Venn diagram of genes containing one or more AS event in UN vs. +SAL samples. **E.** Venn diagram of differentially expressed genes (DEGs) vs. alternatively spliced genes (AS) in UN and +SAL samples. **F.** MAJIQ PSI quantification of junctions with SCR (left) and *Srsf7* KD (right) in UN samples. Violin plot of ΔPSI *Dcun1d2* SRSF7-dependent local splicing variation. Splice graph of *Dcun1d2* in SCR (top) and *Srsf7* KD (bottom) generated by MAJIQ/VOILA **G.** As in F but for +SAL samples. **H.** Semi-quantitative PCR of *Dcun1d2* in SCR, *Srsf7* KD #1 and KD #2 RAW 264.7 cells. Each sample in biological triplicate. Right, quantification. **I.** As in H but for *Dcun1d2* +LPS (4h) **J.** As in F but *Mphosph9.* **K.** As in H but *Mphosph9* (+4h LPS). **L.** As in H but *Mphosph9* in UN samples. Data are expressed as a mean of three or more biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

S1

Figure S1: Supplemental data for Figure 1

A. Functional categorization of transcripts differentially spliced in *Srsf7* knockdown RAW 264.7 vs. SCR cells (UN, +SAL, and ΔPSI > 0.2 confidence threshold set at 0.95. **B.** MAJIQ PSI quantification of junctions with SCR (left) and *Srsf7* KD (right). Violin plot of ΔPSI *Gripap1* SRSF7-dependent local splicing variation. Splice graph of *Gripap1* in SCR (bottom left) and *Srsf7* KD (bottom right) generated by MAJIQ/VOILA at baseline. **C.** Semiquantitative PCR of *Gripap1* in SCR, *Srsf7* KD1 and KD2 RAW 264.7 cells, UN (top) and 4h +LPS (bottom). Each sample in biological triplicate. Right, quantification. **D.** MAJIQ PSI quantification of junctions with SCR (left) and *Srsf7* KD (right). Violin plot of ΔPSI *Dop1a* SRSF7-dependent local splicing variation. Splice graph of *Dop1a* in SCR (bottom left) and *Srsf7* KD (bottom right) generated by MAJIQ/VOILA at baseline. **E** Semiquantitative PCR of *Gripap1* in SCR, *Srsf7* KD #1 and KD #2 RAW 264.7 cells, UN (top) and 4h +LPS (bottom). Each sample in biological triplicate. Right, quantification. Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=$p<0.05$, **=$p<0.01$, ***=$p<0.001$, ****=$p<0.0001$.

control experimentally. Consistent with our MAJIQ analysis, we measured significantly higher lev-

els of *Dcun1d2* exon 6 skipping in *Srsf7* knockdown MΦs in both conditions (**Fig. 1H-I**).

We then measured condition-specific SRSF7-dependent splicing events, based on pre-

dictions made by the MAJIQ algorithm (**Fig. 1D**). In +SAL samples, MAJIQ quantified SRSF7-

dependent alternative 3'SS usage in exon 3 of the *Mphosph9* gene (Ensembl Transcript ID:

ENSMUST00000184951.8) that encodes an M-phase phosphoprotein (*Srsf7* knockdown #2

(ΔPSI -35.7%) (**Fig. 1J**). RT-PCR confirmed this SRSF7-dependent AS event in LPS-treated MΦs

(**Fig. 1K**). Curiously, we measured similar levels of alternative 3'SS usage in untreated *Srsf7*

knockdown MΦs (**Fig. 1L**), even though this event was not called by the splicing algorithm.

SRSF7-dependent inclusion of exon 8 in *Gripap1* (Ensembl Transcript ID

ENSMUST00000065932.14) (**Fig. S1B**), and inclusion of exon 35 in *Dop1a* (Ensembl Transcript

ID ENSMUST00000190957.7) (**Fig. S1D)** only met our threshold in UN and +SAL conditions,

respectively, but could be detected in both macrophage states by RT-PCR (**Fig. S1C, S1E**).

These data argue that SRSF7 mediates the same AS events in resting and activated MΦs. Based

on the nature of the genes whose AS relies on SRSF7 and the lack of specificity for events be-

tween resting and activated MΦs, we can conclude that SRSF7 is unlikely to dramatically influ-

ence the macrophage innate immune transcriptome at the level of splicing.

SRSF7 is a positive regulator of type I interferon transcript accumulation in macrophages

In addition to our AS analysis, we also measured differentially expressed genes in *Srsf7* knock-

down vs. SCR control RAW MΦs, both in untreated cells and at 4h post-*Salmonella* infection

(Wagner et al., 2021). Consistent with our studies of other SR proteins (Wagner et al., 2021;

Wagner et al., 2022)), *Srsf7* knockdown impacted expression of many housekeeping genes in

resting MΦs, particularly in pathways related to translation regulation (EIF2 signaling and Regu-

lation of eIF4 and p70S6K signaling) and phagosome maturation (**Fig. S2A**). Interestingly, in

resting MΦs, we also saw enrichment for SRSF7-dependent genes in pathways like "Communi-

cation between innate and adaptive immune cells" (**Fig. S2A**), suggesting a dysregulation of in-

nate immune genes in *Srsf7* knockdown MΦs even in the absence of an innate stimulus. Enrich-

ment for pathways related to innate immunity was also evident in *Salmonella*-infected samples,

where *Srsf7* knockdown dysregulated genes in pathways related to "role of pattern recognition

receptors in recognition of bacteria and viruses," "interferon signaling," "role of PKR in interferon

induction and antiviral response," and "activation of IRF by cytosolic pattern recognition receptors"

(**Fig. S2B**). Many of these categories contain genes related to the type I interferon (IFN) response,

a ubiquitous antiviral response characterized by expression of *Ifnb1* and interferon stimulated

Figure 2

Figure 2: SRSF7 promotes type I interferon expression.

A. Scatterplot of genes differentially expressed in *Srsf7* KD RAW MΦs relative to SCR in each of the conditions queried (X axis = UN; Y axis = +SAL). **B.** Integrated Genomics Viewer track of *Zbp1* reads in +SAL samples. SCR (grey) and *Srsf7* KD reads (teal). **C.** As in B but for *Ifit3.* **D.** RT-qPCR of *Zbp1* in SCR and *Srsf7* KD#1 or KD#2 RAW 264.7 MΦs at baseline (UN). **E.** As in D but for *Ifit3.* **F.** RT-qPCR of *Zbp1* in SCR and *Srsf7* KD#1 or KD#2 RAW 264.7 MΦs 4h post stimulation with 100 ng/ml LPS. **G.** As in F but for *Ifit3.* **H.** RT-qPCR of *Zbp1* in SCR and *Srsf7* KD #1 or KD #2 RAW 264.7 MΦs 4h post-transfection with 1 µg/ml ISD. **I.** As in F but for *Ifit3.* **J.** RT-qPCR of *Zbp1* in SCR and *Srsf7* KD #2 RAW 264.7 MΦs 4 or 8h post infection with *M. tuberculosis* (Erdman strain; MOI=5). **K.** As in J but for *Ifit3.* **L.** Immunoblot of ZBP1 protein levels in SCR and *Srsf7* KD #2 cells at baseline, 4, and 8h, post stimulation with 100 ng/ml LPS. Right, quantification. **M.** RT-qPCR of *Srsf7* in primary MEFs at baseline or 4h post stimulation with 100ng LPS. **N.** As in M but for *Zbp1*. **O.** As in M but for *Ifit3*. **P.** As in M but for *Irf7*. **Q.** RT-qPCR of *Vsvg* in SCR, *Srsf7* KD #1, and *Srsf7* KD #2 RAW 264.7 MΦs over a time course (4, 8, 12h) post infection with VSV (MOI=0.1). **R.** As in M but for *Vsvm.* **S.** RT-qPCR of *Zbp1* in SCR, *Srsf7* KD #1, *Srsf7* KD #2 RAW 264.7 MΦs over a time course (1, 4, 8, 12h) post infection with VSV MOI=0.1. **T.** As in S but for *Ifit3.* Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001, ****=p<0.0001.

A

Srsf7 knockdown
Uninfected

EIF2 signaling
Phagosome maturation
Regulation of eIF4 and p70S6K signaling
Antigen presentation pathway
Communication between innate and adaptive immune cells
Polyamine regulation in colon cancer
Virus entry via endocytic pathways
mTOR signaling

0   5   10   15   20

B

Srsf7 knockdown
+*Salmonella*

Communication between innate and adaptive immune cells
Role of pattern recognition receptors in recognition of bacteria and viruses
Systemic lupus erythematosus in B cell signaling pathway
Role of PKR in interferon induction and antiviral response
Interferon signaling
Activation of IRF by cytosolic pattern recognition receptors
Granulocyte adhesion and diapedesis
Agranulocyte adhesion and diapedesis

0   5   10   15

C

+*S. Typhimurium*

ns
ns

CFUs 22h/0h

SCR   *Srsf7* KD1   *Srsf7* KD2

D

+*M. tuberculosis*

SCR
Srsf7 #2

RLU

0   2   3   4
days post-infection

E

+LPS
MEFs

SCR
Srsf7 #2

Rsad2/*Actb*

**   **

0   4
hours

F

+VSV

SCR
Srsf7 #1
Srsf7 #2

Il1b/*Actb*

ns
ns

0   1   4   8   12
hours

S2

Figure S2: Supplemental data for Figure 2
A. Ingenuity Pathway Analysis (IPA; Qiagen) of pathways enriched for differentially expressed
genes in *Srsf7* KD RAW 264.7 MΦs (UN). **B.** As in A for *Srsf7* KD RAW 264.7 MΦs +SAL. **C.**
Infection of SCR or *Srsf7* KD RAW 264.7 MΦs with *Salmonella* enterica (MOI=1) in non-SPI-1
inducing conditions. Replication represented as fold 22h over 2h post infection. **D.** Mtb luxBCADE
growth in SCR or *Srsf7* KD RAW 264.7 MΦs measured by relative light units (RLUs) over a 4-day
time course (MOI=1). **F.** RT-qPCR of *Rsad2* in siRNA *Srsf7-* or siControl-transfected primary
MEFs at baseline or 4h post stimulation with 100ng/ml of LPS. **G.** RT-qPCR of *Il1b* in SCR or
Srsf7 KD RAW 264.7 MΦs over a time course of VSV infection. Data are expressed as a mean
of three or more biological replicates with error bars depicting SEM. Statistical significance was
determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01.

genes (ISGs) (Schoggins, 2019; Schoggins and Rice, 2011). In MΦs, the type I IFN response is
activated by the transcription factors IRF3 and IRF7 in response to several immune agonists in-
cluding LPS/*Salmonella* infection, cytosolic dsDNA, and cytosolic dsRNA (21478870 (Honda and
Taniguchi, 2006; Jefferies, 2019; Schoggins et al., 2011).

Certain ISGs are maintained in MΦs at low basal levels (Barrat et al., 2019; Gough et al.,
2012)). We observed that *Srsf7* knockdown decreased abundance of ISGs like *Apol9a*, *Apol9b*,
Ifit3, and *Ifi202b*, which are appreciably expressed in resting MΦs (**Fig. 2A**; bottom left quadrant).
On the other hand, genes like *Mx1, Zbp1,* and *Ifi209*, which are virtually undetectable in resting
cells, only displayed *Srsf7*-dependence in our *Salmonella*-infected samples (**Fig. 2A**; lower y-
axis). Visualizing RNA reads from two such ISGs, *Zbp1* and *Ifit3*, on the Integrated Genomics
Viewer (Broad Institute), we did not notice any changes in intronic reads in *Srsf7* knockdown
versus control cells (**Fig. 2B-C**). This was consistent with our MAJIQ analysis, which did not detect
many statistically significant splicing changes in genes annotated as ISGs in *Srsf7* knockdown
RAW MΦs (**Fig. 1E**).

We predicted that failure to phagocytose *Salmonella* or defects in *Salmonella* replication
could result in lower ISG expression in *Srsf7* knockdown cells, but enumeration of colony forming
units (CFUs) did not reveal any differences in bacterial burdens between SCR and *Srsf7* knock-
down RAW MΦs over a 22h infection time-course (**Fig. S2C**). To determine if SRSF7's ability to
promote ISG expression was unique to *Salmonella* infection, we measured ISG expression in
SCR and *Srsf7* knockdown MΦs at 4h post-treatment with LPS (100 ng/ml), after dsDNA trans-
fection (1 μg ISD), which activates type I IFN gene expression downstream of cGAS, and after
treatment with recombinant IFN-β (200 IU, 4h), which activates ISG expression in MΦs by binding
to IFNAR and engaging JAK/STAT cascades. In all cases, loss of *Srsf7* led to impaired activation
of *Zbp1* and *Ifit3* expression (**Fig. 2D-I**). We also measured lower ISG expression in *Srsf7* knock-
down MΦs during infection with the intracellular bacterial pathogen, *Mycobacterium tuberculosis*
(MOI = 5) (**Fig. 2J-K**), which elicits *Ifnb1* and ISG expression via the cGAS/STING pathway
(Watson et al., 2015). Again, this defect in ISG expression was independent of bacterial internal-
ization/replication (**Fig. S2D**). Importantly, over an 8h time-course of LPS-treatment, lower *Zbp1*
transcript levels correlated with lower protein accumulation (**Fig. 2L**), suggesting that failure to
fully upregulate ISG expression in *Srsf7* knockdown MΦs is likely to functionally impact the innate
immune response to infection. This phenotype was not unique to RAW MΦs, as loss of SRSF7 in
primary mouse embryonic fibroblasts (MEFs) via siRNA transfection (alongside siNC transfected
controls) (**Fig. 2M**) also resulted in low basal ISG expression and a failure to fully induce ISGs
upon LPS stimulation (**Fig. 2N-P and S2E)**

To test whether SRSF7 is a *bona fide* player in activating antimicrobial immunity in MΦs,
we infected SCR and *Srsf7* knockdown MΦs with vesicular stomatitis virus (VSV), a negative-
strand RNA virus that stimulates type I IFN responses via TLR3 and RIG-I and is extremely sen-
sitive to type I IFN expression (Gao et al., 2021; Wagner et al., 2021)). While genes like *Ifnb1* and
ISGs are robustly induced by bacterial pathogens (e.g. *Salmonella* and *M. tuberculosis*), they are
most studied during viral infection, where they function to inhibit different steps of the viral life
cycle (Schoggins, 2019). We measured a dramatic increase in VSV replication in *Srsf7* knock-
down MΦs, measured by RT-qPCR of two VSV genes, *Vsvg* and *Vsvm* (**Fig. 2Q-R**) that was
concomitant with depressed ISG transcript induction over a 12-hour time course of VSV infection
(MOI = 0.1) (**Fig. 2S-T**). We did not detect SRSF7-dependent differences in *Il1b* expression in
response to VSV infection, arguing against a role for SRSF7 in controlling innate gene expression
downstream of NFκB (**Fig. S2F**). Together, these data identify SRSF7 as a positive regulator of
the type I IFN response in MΦs and suggest that the mechanism of SRSF7-dependent ISG ex-
pression is shared by several pathogen sensing cascades (i.e. occurs at one or more nodes where
the TLR4, cGAS, and RIG-I cascades converge).
254
SRSF7 drives ISG expression without interfering with innate immune signaling.
Given that loss of SRSF7 results in a failure to fully induce many ISGs, we hypothesized
that SRSF7 may act directly on one or more critical innate signaling proteins or transcription fac-
tors. To pinpoint where in the type I IFN signaling response SRSF7 acts, we treated *Srsf7*
knockdown or SCR RAW MΦs with LPS (100 ng/ml) and measured secretion of type I interferon
(IFN-β/α) over a 6h time-course via a luciferase ISRE reporter cell line (Hoffpauir et al., 2020;
Wagner et al., 2022)). *Srsf7* knockdown MΦs produced significantly less IFN-β/α compared to
SCR controls at all time points measured (**Fig. 3A**). We next asked if impaired ISG expression
in *Srsf7* knockdown MΦs could be rescued if *Srsf7* knockdown and SCR control cells received
equal amounts of recombinant IFN-β, a potent type I interferon driver. We found that even in the
presence of high amounts of IFN-β, *Srsf7* knockdown MΦs fail to express ISGs like *Ifi202b* and
Zbp1 to the same level as SCR controls, at both the transcript (**Fig. 3B-C**) and protein level
(**Fig. 3D**). These results suggest that loss of SRSF7 impacts some aspect of the type I IFN re-
sponse that is engaged upstream and downstream of IFNAR and that loss of IFN-β production
is not the sole driver of ISG defects in *Srsf7* knockdown MΦs.
We then asked whether altered signaling downstream of IFNAR could explain defective
ISG expression in *Srsf7* knockdown RAW MΦs. We treated *Srsf7* knockdown and SCR RAW
MΦs with LPS and measured phosphorylation of the transcription factor IRF3 (activated

Figure 3

Figure 3: SRSF7 is required for IRF7 accumulation in macrophages

A. Secreted IFNα/β in SCR and *Srsf7* KD RAW 264.7 MΦs, as a measure of luciferase expressed by ISRE reporter cells. **B.** RT-qPCR of *Ifi202b* in SCR and *Srsf7* KD RAW 264.7 MΦs 4h post stimulation with 200 IU IFN-β. **C.** As in B for *Zbp1*. **D.** Immunoblot of ZBP1 and ACTIN over a time course of IFN-β (200 IU) treatment in SCR and *Srsf7* KD RAW 264.7 MΦs. Quantification, right. **E.** Immunoblot of Y701 STAT1, STAT1, S396 IRF3, IRF3, and ACTIN in SCR and *Srsf7* KD RAW 264.7 MΦs over a time course of LPS treatment (100ng/ml). **F.** Differential expression of transcription factor genes in untreated *Srsf7* knockdown vs. SCR RAW 264.7 MΦs as fold change and -log(pvalue). **G.** As in F but for +SAL samples. **H.** Integrated Genomics Viewer track of *Irf7* reads in +SAL samples. SCR (grey) and *Srsf7* KD reads (teal). **I.** RT-qPCR of *Irf7* over a time course of LPS stimulation (100ng/ml) in SCR and *Srsf7* KD RAW 264.7 MΦs. **J.** Immunoblot of IRF7 and ACTIN protein levels over a time course of LPS stimulation (100ng/ml) in SCR and *Srsf7* KD RAW 264.7 MΦs. Quantification, right. **K.** RT-qPCR of *Irf7* over a time course of Mtb infection (Erdman strain; MOI=5) in SCR and *Srsf7* KD RAW 264.7 MΦs. **L.** RT-qPCR of *Irf7* over a time course of VSV infection (MOI=0.1) in SCR or *Srsf7* KD RAW 264.7 MΦs. **M.** Immunoblot of SRSF7, IRF7, and ACTIN in siControl- or si*Srsf7*-transfected BMDMs at baseline. Quantifications, right. **N.** RT-qPCR of *Irf7* in siControl or si*Srsf7* BMDMs at baseline. Data expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001, ****=p<0.0001.

A

B

iBMDMs
+LPS   0   2   4   6   hrs

α-IRF7

α-ACTIN

C

BMDMs
+LPS   0   2   4   6   hrs

α-IRF7

α-ACTIN

S3

Figure S3: Supplemental data for Figure 3

A. Quantification of Figure 3E. n=2 **B.** Immunoblot of IRF7 and ACTIN protein levels over a time course of LPS stimulation (100ng/ml) in WT BMDMs. **C.** As in B but in WT iBMDMs. Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test.

downstream of TLR4-TRIF signaling) and STAT1 (activated downstream of IFNAR) over a 6h

time course. We did not measure significant changes in phosphorylation of IRF3 at S396 or

phosphorylation of STAT1 at Y701 (**Fig. 3E**; quantification in **Fig. S3A**) effectively ruling out de-

fective signal transduction as the culprit for dampened ISG expression in *Srsf7* knockdown

MΦs.

Next, we entertained the possibility that SRSF7 is required for the expression of one or

more transcription factors needed to activate the type I IFN response downstream of pattern

recognition receptor engagement. To test this, we generated a list of transcription factors asso-

ciated with macrophage innate immune gene expression in the STAT, IRF, and NFκB families,

and turned back to our RNA-seq data to see how expression of each factor was impacted by

loss of SRSF7. While expression of most TFs was unaffected, we observed a significant defect

in *Irf7* expression in *Srsf7* knockdown RAW MΦs compared to SCR controls at both baseline

and after *Salmonella* infection (**Fig. 3F-G**). Visualization of *Irf7* gene expression data on the IGV

highlights fewer *Irf7* exonic reads in *Srsf7* knockdown (teal) compared to SCR (grey) RAW MΦs

in both conditions (**Fig. 3H**). We did not detect an increase in intron reads in *Srsf7* knockdown

MΦs nor did our MAJIQ analysis identify SRSF7-dependent *Irf7* alternative splicing events (**Fig.**

1D), arguing against a role for SRSF7 in regulating *Irf7* transcript abundance via splicing.

Initially discovered in the context of Epstein Barr virus, IRF7 (interferon regulatory factor

7) acts as both a master regulator of antiviral type I IFN responses and an ISG (Honda et al.,

2005). Basal amounts of IRF7 vary between immune cell types. Dendritic cells are heavily reli-

ant upon IRF7 at baseline and several recent reports identify IRF7 as important for priming bone

marrow derived macrophages (BMDMs) downstream of LPS (Fitzgerald et al., 2003; Gu et al.,

2022; Jefferies, 2019; Sin et al., 2020). We can readily measure IRF7 transcript and protein in

resting WT RAW MΦs (**Fig. 3I-J**) and in WT immortalized BMDMs (**Fig. S3B**). IRF7 protein lev-

els are low in WT resting BMDMs but are upregulated in response to LPS (**Fig. S3C**).

To better appreciate the kinetics of SRSF7 control of *Irf7* expression, we treated *Srsf7*

knockdown and SCR control RAW MΦs with LPS and measured *Irf7* transcript and protein over

a 12h time course. We measured significantly lower levels of *Irf7* mRNA in *Srsf7* knockdown

RAW MΦs at every time-point queried (**Fig. 3I**). IRF7 protein levels were lower in *Srsf7* knock-

down RAW MΦs at 0 and 4h but appeared to rebound at later time points (8 and 12h post-LPS)

(**Fig. 3J**), consistent with previous reports of IRF7 stabilization during the innate response to in-

fection (Sin et al., 2020). We also measured significantly less *Irf7* transcript in *Srsf7* knockdown

RAW MΦs in response to *M. tuberculosis* (**Fig. 3K**) and VSV infection (**Fig. 3L**). Although we

were only able to achieve modest knockdown of SRSF7 in BMDMs transfected with an siRNA

directed against *Srsf7* (alongside a non-targeting control; siNC) (**Fig. 3M**), we can detect less

IRF7 protein (**Fig. 3M**) and transcript (**Fig. 3N**) in these primary cells as well. Together, these

data demonstrate a previously unappreciated role for SRSF7 in controlling *Irf7* transcript and

protein levels in primary macrophages and in the RAW MΦ cell line.

IRF7 controls basal and induced expression of ISGs in RAW MΦs

Our findings argue that loss of ISG expression in *Srsf7* knockdown cells stems primarily

from loss of IRF7. However, as most literature points to a role for IRF7 in activated immune

cells, we were initially cautious to attribute basal ISG phenotypes in *Srsf7* knockdown MΦs to

IRF7. To better understand how IRF7 controls type I IFN responses in resting and activated

RAW MΦs, we generated *Irf7* KO RAW MΦs using CRISPR-Cas9 editing. *Irf7*-specific guides

were designed to target *Irf7* exon 3 alongside a GFP-specific gRNA as a negative control. Two

clonally expanded *Irf7* KO cell lines and a GFP gRNA control cell line were selected and loss of

IRF7 protein was confirmed by western blot (**Fig. 4A**). Total RNA from untreated *Irf7* KO #1 and

control cell lines as well as from cells stimulated with LPS for 4h (100ng/ml) were sent for RNA-

seq analysis (150bp paired end reads; Illumina NovaSeq S4, ~80,000,000 reads per sample un-

treated and ~20,000,000 reads per sample +4h LPS). We measured differential expression of

896 transcripts (fold-change < -1.5; p<0.05) in *Irf7* KO vs. control cells at baseline and 1575

genes were differentially expressed in *Irf7* KO MΦs at 4h post-LPS treatment (**Fig. 4B**). Pathway

analysis (BioPlanet Pathway Enrichment Analysis, NCATS (Huang et al., 2019) identified inter-

feron-related pathways enriched for differentially expressed genes in *Irf7* KO cells at baseline

and at 4h post-LPS treatment (**Fig. 4C**). Cataloging these lists of genes, we identified >30 ISGs

whose expression was impacted by loss of IRF7 in resting RAW MΦs (**Fig. 4D**), and confirmed

significant downregulation of *Ifit3b*, *Zbp1*, and *Ifnb1* in untreated *Irf7* KOs compared to control

cells by RT-qPCR (**Fig. 4E**). Visualization of reads for select ISGs on the IGV showed lower lev-

els of exonic reads in *Irf7* KO RAW MΦs (purple vs. grey) (**Fig. 4F**). We also measured impaired

ISG induction in *Irf7* KO RAW MΦs after LPS treatment, but curiously, while the overall number

of genes impacted was higher after LPS treatment, the degree to which their expression was

impacted by loss of IRF7, compared to GFP gRNA controls, was less (**Fig. S4B**). We predict

this is due to activation of IRF3 that can compensate for loss of IRF7 after stimulation. Im-

portantly, we found that differentially expressed ISGs were shared between *Srsf7* KD and *Irf7*

KO cells: most of the downregulated ISGs in *Srsf7* knockdown cells are also downregulated in

Irf7 KO MΦs (**Fig. 4G**). Any lack of overlap between these lists likely stems from the fact that

complete ablation of IRF7 and a 2-3-fold downregulation of IRF7 in *Srsf7* knockdown cells

Figure 4

Figure 4: RAW 264.7 macrophage cell lines are reliant on IRF7 for basal ISG expression
A. Immunoblot of IRF7 protein levels in GFP gRNA, *Irf7* KO1 and *Irf7* KO2 RAW 264.7 MΦs at baseline. ACTIN used as loading control. **B.** Venn diagram depicting differential gene expression of *Irf7* KO1 at baseline vs. 4h +LPS (100ng/ml). Cut off: fold change < -1.5 and p-value=<0.05. **C.** Top: BioPlanet Pathway Analysis of untreated *Irf7* KO gene expression. Bottom: BioPlanet Pathway Analysis of 4h +LPS *Irf7* KO gene expression. **D.** Heatmap depicting *Irf7* KO baseline gene expression from BioPlanet pathway analysis categories: "Interferon alpha/beta signaling," "Interferon signaling," and "Interferon-gamma signaling pathway." Duplicate terms removed. **E.** RT-qPCR of *Ifit3, Zbp1,* or *Ifnb1* at baseline in *Irf7* KO vs. GFP gRNA control cells. **F.** Integrated Genomics Viewer track of *Ifi44, Ifi202b, and Oasl1* reads in baseline (UN) samples in GFP gRNA (light grey) or *Irf7* KO (dark blue). **G.** Top: Venn diagram depicting DEGs in *Irf7* KO1 (compared to GFP gRNA) and *Srsf7* KD (compared to SCR) RAW 264.7 MΦs at baseline. Bottom: Heatmap of 26 genes in common between *Irf7* KO1 and *Srsf7* KD at baseline. Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

843 **Figure S4: Supplemental data for Figure 4**
844 **A.** RT-qPCR of *Stat1* at baseline (UN) in *Irf7* KO cells. **B.** Heatmap depicting BioPlanet Analysis
845 of genes in the n "interferon a/b signaling" category at baseline or 4h post stimulation with LPS in
846 GFP gRNA or *Irf7* KO RAW 264.7 cells.

341   differentially impact ISG expression in MΦs. Together, these data provide strong evidence for a
342   role for IRF7 in maintaining basal levels of type I IFNs and ISGs in MΦs.

SRSF7 is sufficient for IRF7 and ISG expression

Together, our data support a model wherein SRSF7 directly activates *Irf7* transcription. If this
were the case, then overexpression of SRSF7 should be sufficient to drive *Irf7* expression. To
test this model, we generated doxycycline-inducible MΦs cell lines that express either 3xFLAG-
SRSF7 or 3xFLAG-GFP negative control after addition of doxycycline. To ensure comparable
levels of 3xFLAG-GFP and 3xFLAG-SRSF7 protein expression, we tested and optimized treat-
ment times, deciding on 2h +DOX for GFP and 22h +DOX for SRSF7 (**Fig. 5A**). We found that
expression of SRSF7 alone was sufficient to increase the abundance of *Irf7* transcript (**Fig. 5B**)
and IRF7 protein levels (**Fig. 5C**) in otherwise unstimulated RAW MΦs. We also measured in-
creased expression of ISGs like *Ifit3* in 3xFLAG-SRSF7 MΦs compared to GFP controls (**Fig.**
5D). To ensure that different lengths of doxycycline treatment was not contributing to our pheno-
type, we also measured ISG expression in MΦs stably expressing 3xFLAG-SRSF7 or 3xFLAG-
GFP. We found that stable overexpression of 3xFLAG-SRSF7 was also sufficient to drive ISG
expression in resting MΦs (**Fig. 5E-G**). The impact of SRSF7 overexpression on *Irf7* following
LPS stimulation was less pronounced. We detected increased *Irf7* transcript expression in
3xFLAG-SRSF7 expressing cells at 6h post LPS, but this did not correspond to differences in
IRF7 protein accumulation (**Fig. S5A,B**). Given how strong an inducer LPS is of the type I IFN
response, it follows that SRSF7 overexpression may not further induce *Irf7* expression on top of
these already high levels. We did, however, measure a dramatic increase in *Irf7* and *Ifit3* tran-
script levels in 3xFLAG-SRSF7-expressing stable cells upon infection with *M. tuberculosis* (**Fig.**
5H-I), suggesting that the nature of the stimulus may impact how SRSF7 controls IRF7 and ISG
expression. Because expression of type I IFN genes has been associated with cell-intrinsic anti-
bacterial control mechanisms and cell death pathways that impact bacterial spread (Patrick et
al., 2016), we also asked whether overexpression of SRSF7 affected Mtb replication. Briefly, we
infected cells with Mtb luxBCADE and followed bacterial replication as a measure of relative
light units (as in (Bell et al., 2021; Wagner et al., 2022)). Remarkably, we measured a significant
increase in Mtb RLUs at Day 3 post-infection (**Fig. 5J**) suggesting that SRSF7 expression may
help control Mtb replication and/or spread. Together, these data suggest a direct role for the
RNA binding protein SRSF7 in promoting expression of *Irf7*.

SRSF7 promotes STAT1 association and promoter clearance at the *Irf7* promoter

Figure 5

Figure 5: Overexpression of 3xFL-SRSF7 drives basal ISG expression and accumulation of IRF7

A. Immunoblot of 3xFLAG-GFP or 3xFLAG-SRSF7 tetracycline inducible RAW 264.7 MΦs treated with doxycycline (DOX) (1 µg/ml). 3xFLAG-GFP MΦs received DOX for 2h and 3xFLAG-SRSF7 MΦs received DOX for 22h. ACTIN shown as loading control. **B.** RT-qPCR of *Irf7* in 3xFLAG-GFP (2h DOX treatment) or 3xFLAG-SRSF7 (22h DOX treatment) inducible RAW 264.7 cells at baseline. **C.** Immunoblot of IRF7 in 3xFLAG-GFP (2h DOX treatment) or 3xFLAG-SRSF7 (22h dox treatment) inducible RAW 264.7 MΦs at baseline (quantification right). **D.** As in B but for *Ifit3*. **E.** RT-qPCR of *Zbp1* in 3xFLAG-GFP or 3xFLAG-SRSF7 stable overexpression RAW 264.7 MΦs. **F.** As in E but for *Ifi44l* **G.** As in E but for *Oasl2.* **H.** RT-qPCR of *Irf7* over a time course of infection with Mtb (Erdman; MOI=5). **I.** As in H but for *Ifit3* **J.** Mtb luxBCADE replication in 3xFLAG-GFP (2h pretreatment with 1 µg/ml doxycycline) or 3xFL-SRSF7 (22h pretreatment with 1µg/ml doxycycline)-inducible RAW 264.7 macrophages measured by relative light units (RLUs) over time course (Erdman; MOI=1). Data are expressed as a mean of three biological replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

A

B

Figure S5: Supplemental data for Figure 5
A. RT-qPCR of *Irf7* in 3xFLAG-GFP (2h dox treatment) or 3xFLAG-SRSF7 (22h dox treatment) inducible RAW 264.7 MΦs over a time course of LPS stimulation (100 ng/ml). **B.** Immunoblot of IRF7 in 3xFLAG-GFP (2h dox treatment) or 3xFLAG-SRSF7 (22h dox treatment) inducible RAW 264.7 cells over a time course of LPS stimulation (100 ng/ml) (quantification right).

Having established that decreased levels of IRF7 is the likely culprit for the type I IFN

defect in *Srsf7* knockdown MΦs (**Figs. 3-4**) and demonstrated that SRSF7 overexpression is

sufficient to promote *Irf7* expression (**Fig. 5**), we set out to determine the mechanism though

which SRSF7 controls IRF7 expression. SR proteins play a well-established role in coupling

transcription and splicing and have been implicated in transcription by interacting with histone

tails (Loomis et al., 2009; Luco et al., 2011; Luco et al., 2010) or by directly mediating transcrip-

tional elongation via pTEFb (Ji et al., 2013). To begin to investigate a role for SRSF7 in activat-

ing *Irf7* transcription, we performed chromatin immunoprecipitation (ChIP)-qPCR for RNAPII

(anti-CTD tail) in *Srsf7* knockdown and SCR control MΦs. We designed primers to tile the *Irf7*

promoter region (100bp size amplicon, 100 bp apart from each other (**Fig. 6A**)). We performed

ChIP at a 4h time point post-LPS treatment as *Irf7* transcription is induced at this time point and

we know the *Srsf7* knockdown phenotype is in play. We observed dramatic accumulation of

RNAPII at/around the *Irf7* transcription start site (TSS) (-400 to +250) in *Srsf7* knockdown MΦs

(**Fig. 6B**, teal). This pattern of RNAPII in *Srsf7* knockdown MΦs is suggestive of a pile-up result-

ing from a failure to convert to a successful elongation complex. We next asked whether this de-

fect in RNAPII clearance could be explained by defects in recruitment or retention of transcrip-

tion factors like STAT1 to the *Irf7* promoter. Using chromatin immunoprecipitation-qPCR with an

antibody against Y701 STAT1, we measured a massive enrichment of STAT1 at the *Irf7* pro-

moter in SCR cells at 4h post-LPS (**Fig. 6C**, grey). This STAT1 signal was completely lost in

Srsf7 knockdown MΦs (**Fig. 6C**, teal). An analogous set of ChIP-qPCRs at the promoter of *Ifit3*,

another ISG whose expression is SRSF7-dependent, showed no differences in RNAPII or

STAT1 occupancy (**Fig. 6D-F**). Likewise, we saw no difference in RNAPII occupancy in *Srsf7*

knockdown MΦs versus SCR controls at the *Stat1* promoter (**Fig. 6G-H**). These data support a

role for SRSF7 in promoting transcription of *Irf7* and suggest that dampened expression of ISGs

like *Ifit3* in *Srsf7* knockdown MΦs occurs indirectly, downstream of IRF7 defects.

Having linked STAT1 binding and RNAPII elongation at the *Irf7* promoter to SRSF7, we

next asked whether SRSF7 carries out this function by directly associating with the *Irf7* genomic

locus. As ChIP-grade antibodies are not available for SRSF7, we turned back to our 3xFLAG-

SRSF7 and 3xFLAG-GFP doxycycline-inducible cell lines (**Fig. 5A**) and performed ChIP-qPCR

of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Irf7* promoter using the same primer sets shown in

Fig. 6A. We observed significant enrichment of SRSF7 at the *Irf7* promoter (-500 to 0) (3 to 4-fold

over 3xFLAG-GFP control) (**Fig. 6I**) and no SRSF7 enrichment at *Ifit3* (**Fig. 6J**). To take an unbi-

ased global look at SRSF7 association with the murine genome, we performed ChIP-seq. Data

was analyzed by ROSALIND. Briefly, Cutadapt trimmed reads, quality scores were determined

Figure 6

Figure 6: SRSF7 promotes STAT1 association and RNA polymerase II elongation by associating with the *Irf7* promoter

A. Schematic diagram of the *Irf7* gene with tiling primers designed around the transcriptional start site (orange). **B.** RNA polymerase II ChIP-qPCR at the *Irf7* promoter in SCR and *Srsf7* KD RAW MΦs 4h post-LPS stimulation (100ng/ml). Data represented as elution/input normalized to that of an intergenic control primer set. **C.** Y701 STAT1 ChIP-qPCR at the *Irf7* promoter in SCR or *Srsf7* KD RAW MΦs 4h post-LPS stimulation (100 ng/ml). Data represented as elution/input normalized to that of an intergenic control. **D.** Schematic diagram of the *Ifit3* gene with tiling primers designed around the transcriptional start site (orange). **E.** As in B but at the *Ifit3* promoter. **F.** As in C but at the *Ifit3* promoter. **G.** Schematic diagram of the *Stat1* gene with tiling primers designed around the transcriptional start site (orange). **H.** As in B but at the *Stat1* promoter. **I.** ChIP-qPCR at the *Irf7* promoter 4h post-LPS (100 ng/ml) in 3xFLAG-GFP and 3xFLAG-SRSF7 dox-inducible RAW 264.7 MΦs. **J.** As in I at the *Ifit3* promoter. **K.** Functional categorization of unique ChIP peaks in 3xFL-SRSF7 dox-inducible RAW 264.7 MΦs 4h post-LPS (100 ng/ml). **L.** Heatmap indicating read coverage density at transcriptional start sites (TSS) in 3xFL-GFP or 3xFL-SRSF7 inducible RAW 264.7 MΦs. **M.** Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Irf7* promoter by ChIP-seq: IP (top) and input (bottom). **N.** As in M but at the *Ifit3* promoter. Data are expressed as a mean of three or more technical replicates with error bars depicting SEM. Statistical significance was determined using two tailed unpaired student's t test. *=p<0.05, **=p<0.01, ***=p<0.001.

Figure S6: Supplemental data for Figure 6

A. Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Nfkbia* promoter by ChIP-seq: IP (top)

and input (bottom). **B.** Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the *Elk4* promoter by

ChIP-seq: IP (top) and input (bottom). **C.** Enrichment of 3xFLAG-SRSF7 and 3xFLAG-GFP at the

Irf7 promoter by ChIP-seq: IP (top) and input (bottom). **D.** Putative SRSF7 motifs predicted by

HOMER from 3xFLAG-SRSF7 ChIP-seq data.

by FASTQC, and reads were aligned to the *Mus musculus* genome build mm10 using Bowtie.

After quality control, peaks were called using MACS2 with input/IgG controls background sub-

tracted. We detected 548 unique peaks for SRSF7 across the genome, with almost half of these

peaks being at transcription start sites (**Fig. 6K-L**), for example at *Nfkbia* (**Fig. S6A**) and *Elk4*

(**Fig. S6B**). By ChIP-seq we again saw a clear SRSF7 peak at the *Irf7* promoter compared to GFP

(**Fig. 6M**), but not at *Ifit3* (**Fig. 6N**) or *Stat1* (**Fig. S6A**). ChIP-seq data revealed 3xFL-SRSF7

enrichment at several immune-associated genes such as *Fos*, *Myd88, Ifitm3*, and *Dusp5*. Inter-

estingly, and in support of our model that lack of ISG induction in *Srsf7* knockdown MΦs is driven

by defects in *Irf7* expression, we did not see SRSF7 peaks at any other interferon-associated

transcription factors. ChIP-seq did not reveal a strong consensus sequence for SRSF7 and con-

sensus sequence enrichment was only detected at subsets of genes: Etv2 (ETS) sites (P-value

= 1e-6 and 16.79% of targets), ZNF41 (P-value = 1e-6 and 1.46% of targets), Fili1 (ETS) (P-value

= 1e-6 and 25.36% of targets), and AT1G76870 (Trihelix) (P-value = 1e-6 and 4.93% of targets)

(**Fig. S6D**). These data reveal unexpected roles for the RNA binding protein SRSF7 in associating

with gene promoters and in facilitating STAT1 association and RNAPII elongation at the *Irf7* pro-

moter.

DISCUSSION

Considering the critical role SR/hnRNP family proteins play in regulating many aspects

of gene expression, we know surprisingly little about how they shape the transcriptomes of spe-

cialized cell types. Here, we describe a novel role for the splicing regulatory protein SRSF7 in

stimulating the type I IFN response by activating transcription of the *Irf7* gene. Our data demon-

strate that enrichment of SRSF7 at the *Irf7* promoter facilitates association of the transcription

factor STAT1 and promotes RNAPII elongation to activate *Irf7* gene expression. By controlling

IRF7 abundance, SRSF7 plays a privileged role in regulating both antiviral (**Fig. 2Q-T**) and pro-

bacterial (**Fig. 5J**) innate immune responses in macrophages. In addition to implicating SRSF7

in activating *Irf7* expression, this work demonstrates a critical and underappreciated role for both

SRSF7 and IRF7 in maintaining tonic and early interferon signaling in MΦs.

A major finding of this study relates to the ability of SRSF7—a canonical RNA binding

protein—to promote transcription of *Irf7*. Over the past two decades, the idea that transcription

and RNA processing occur in a linear, stepwise fashion has been repeatedly challenged. It is

now generally accepted that many steps in RNA processing (i.e. capping, pre-mRNA splicing,

RNA editing) occur co-transcriptionally while nascent transcripts are still bound to RNAPII. Ac-

cordingly, splicing can influence transcription and transcription can impact splicing (Tellier et al.,

2020). Several studies have linked SR proteins to RNAPII dynamics and co-transcriptional

splicing. SRSF2, for example, releases the cyclin-dependent kinase pTEFb from an inhibitory

7SK-containing complex to relieve stalled polymerase at promoters (Ji et al., 2013) and SRSF3

regulates exon inclusion via the C-terminal domain of RNAPII (de la Mata and Kornblihtt, 2006).

Although our data suggest an analogous role for SRSF7 in mediating transcription factor binding

and RNAPII elongation at the *Irf7* promoter, several aspects of this mechanism still need to be

explored.

One outstanding question relates to why we see SRSF7 at certain promoters, like *Irf7*,

but not others (**Fig. 6**). We currently do not know whether SRSF7 association with the *Irf7* pro-

moter is RNA-dependent. If so, it is possible that specificity is dictated by sequences encoded in

exons/introns of genes downstream of SRSF7-sensitive promoters such that nascent RNAs

serve as the platform for SRSF7 recruitment. The reported consensus sequence for SRSF7 is

GA(C/T)GA(C/T) (Cavaloc et al., 1999) and SRSF7 was shown to bind TGA(T/C)C sequences

in its own transcript to regulate poison cassette exon inclusion (Konigs et al., 2020). We can de-

tect several potential SRSF7 bindings sites in the first coding exon of *Irf7* (TGACC). Thus, it is

possible that SRSF7's association at the *Irf7* promoter occurs by virtue of co-transcriptional

binding of exonic splicing enhancers (ESEs) in nascent transcripts. It is also conceivable that

SRSF7 associates with the *Irf7* promoter in an RNA-independent, DNA-dependent fashion.

SRSF7 is the only SR protein that encodes a Zn-knuckle, and to date, this domain has been

shown to enhance the RNA binding activity of SRSF7 (Cavaloc et al., 1999; Cavaloc et al.,

1994; Wang et al., 2021)). However, in other proteins, for example TAF1, the largest subunit of

the general transcription factor TFIIID, a Zn knuckle encourages DNA binding and promoter oc-

cupancy (Curran et al., 2018). It is tempting to speculate that the Zn finger domain in SRSF7

could promote its association directly with DNA. Future experiments designed to implicate the

Zn knuckle in SRSF7 association with the *Irf7* promoter and/or test the ability of SRSF7 to di-

rectly bind DNA will help resolve the molecular mechanisms of SRSF7 promoter enrichment.

In addition to linking SRSF7 to *Irf7* expression, our studies revealed an unexpected role

for IRF7 in controlling tonic type I IFN expression in resting RAW cells (**Fig. 2A, E**) and MEFs

(**Fig. 2M-P**, time 0). These observations challenge the paradigm that IRF7 needs to be upregu-

lated before it can contribute to type I IFN responses. Indeed, differences in ISG expression

were more robust in IRF7 KO RAW 264.7 MΦs at rest than after LPS stimulation (**Fig. S4B**).

These findings suggest that IRF7 is in fact the major mediator of the type I IFN response in rest-

ing macrophages, but *both* IRF3 and IRF7 contribute once they are activated downstream of

pattern recognition receptor engagement. To date, the role for IRF7 in maintaining tonic type I

IFN signaling has not been fully explored. Our RNA-seq data identify a specific regulon of ISGs

that is particularly sensitive to loss of IRF7 at baseline (**Fig. 4D, G**). Among the most downregu-

lated genes in resting *Irf7* KO macrophages are several encoding guanylate binding proteins

(GBPs) and members of the oligoadenylate synthetase (OAS) family. Regulated expression of

these factors at baseline may reflect their role in directly mediating anti-viral and anti-bacterial

killing, with low level expression of these ISGs in resting cells providing an early advantage

upon infection. By demonstrating a role for IRF7 in maintaining tonic levels of these and other

ISGs, our results hint at roles for IRF7 in cellular homeostasis and disease have not been ap-

preciated to date.

Consistent with this idea, growing evidence suggests that IRF7 expression in immune

cells is subject to multiple levels of control mediated by RBPs. Both mouse and human IRF7

genes encode very small introns (<100nt), an unusual adaptation in mammalian genomes. Reg-

ulated *IRF7* intron retention has been reported in THP-1 cells, wherein intron 2 is efficiently re-

moved upon monocyte-to-macrophage differentiation but not in resting monocytes (Green et al.,

2020). In response to treatment with TNF-α and IFN-α, as well as polyl:C transfection, regulated

removal of IRF7 intron 4 has been reported and was shown to be controlled by the RBP BUD13,

which is enriched on *Irf7* transcripts in BMDMs (Frankiw et al., 2019). Such regulation is pro-

posed to provide innate cells a way to limit the inappropriate expression of genes with high in-

flammatory potential like IRF7, while leaving cells poised to upregulate these genes when cer-

tain signals are received. We did not observe differences in the number of intron reads in *Srsf7*

KD vs. SCR RAW cells (**Fig. 3H**), nor did the MAJIQ algorithm identify SRSF7-dependent alter-

native splicing changes in *Irf7* (**Fig. 1**). We propose that intron retention/alternative splicing of

Irf7 by factors like BUD13 adds an additional layer of regulation on top of the transcriptional con-

trol conferred by SRSF7. The fact that IRF7 is subject to regulation at multiple nodes argues

that it plays a larger than appreciated role in mediating inflammatory responses in immune cells.

Together, these previously published studies and our work presented here, argue that IRF7

does more than amplify late interferon responses and highlight a preeminent role for RNA bind-

ing proteins in regulating innate immune outcomes.

ACKNOWLEDGEMENTS

We would like the thank members of Patrick and Watson labs for critical reading of the manu-

script and invaluable feedback. We would also like to thank Phillip West and members of his lab

at Texas A&M School of Medicine for sharing reagents and protocols related to VSV infection.

Thanks to Susan Carpenter and her lab for sharing reagents and protocols related to iBMDMs.

and to the Genomic and RNA Profiling Core at Baylor, College of Medicine for RNA-seq and

ChIP-seq library generation and sequencing.

513
514

Materials and Methods

M. tuberculosis
The Erdman strain was used for all *M. tuberculosis* (Mtb) infections as well as Mtb luxBCADE a luciferase expressing strain. Low passage lab stocks were thawed for each experiment to ensure virulence was preserved. Mtb was cultured in roller bottles at 37 °C in Middlebrook 7H9 broth (BD Biosciences) supplemented with 10% OADC (BD Biosciences), 0.5% glycerol (Fisher), and 0.1% Tween-80 (Fisher). All work with Mtb was performed under Biosafety Level 3 containment using procedures approved by the Texas A&M University Institutional Biosafety Committee.

Mtb in vitro infections
To prepare the inoculum, bacteria were grown to mid log phase (OD 0.6–0.8), spun at low speed (500 rcf) to remove clumps, and then pelleted and washed with PBS 2X. Resuspended bacteria were sonicated and spun at low speed again to further remove clumps. Then Mtb was then diluted in DMEM plus 10% horse serum (Gibco) and added to cells at a multiplicity of infection (MOI) of 10 for RNA and protein analysis, a MOI of 5 for cell death studies, and a MOI of 1 for bacterial growth assays. The day before the infection, RAW 264.7 macrophages were plated on 12-well tissue culture–treated plates at a density of 3×10^5 cells per well or plated in corning 96 well black plates at 2.5×10^4 cells/well and allowed to rest overnight. Cells were spun with bacteria for 10 min at 1,000 rcf to synchronize infection, washed 2X with PBS, and then incubated in fresh media. Where applicable, RNA was harvested from infected cells using 0.25 ml TRIzol reagent at each time point. For bacterial growth assays RAW 264.7 macrophages were plated on 12-well tissue culture–treated plates at a density of 2.5×10^5 cells per well. Luminescence was read for *M. tuberculosis luxBCADE* by lysing in 250 µl 0.5% Triton X-100 and dividing sample into duplicate wells of a 96-well white bottomed plate (Costar). Luminescence was measured and normalized to background using the luminescence feature of the INFINITE 200 PRO (Tecan) at 0, 48, 72 and 92 h post infection.

S. enterica (ser. Typhimirium)
Salmonella enterica serovar Typhimurium (SL1344) was obtained from Dr. Denise Monack, Stanford. S. T. stocks were streaked out on LB agar plates and incubated at 37°C overnight.

S. enterica (ser. Typhimirium) infections:
RAW 264.7 macrophages were seeded in 12-well tissue culture-treated plates at a density of 7×10^5 cells per cell, and cultures of *S. Typhimurium* were grown at 37°C in LB broth containing 0.3M NaCl 16h prior to infection. Bacterial cultures were diluted 1:20 3h prior to infection to reach mid-log phase OD_{600} 0.6-0.8. Mid-log phase bacteria were washed 3x with PBS and pelleted at 5,000 rpm for 3 min. Bacteria were then added to DMEM (Hyclone) with 10% FBS (Sigma) and 2% HEPES (HyClone) at an MOI of 5 for transcripts and MOI of 1 for replication. Infected cells were spun at 1,000 rpm for 5 min then incubated for 10 min at 37 °C prior to gentamycin washes adding fresh media. At indicated times post infection, cells were harvested with TRIzol for RNA isolation described below.

Vesicular stomatitis virus
Recombinant *Vesicular stomatitis* virus (VSV; Indiana serotype) containing a GFP reporter cloned downstream of the VSV G-glycoprotein (VSV-G/GFP) was obtained collaboratively from Dr. John Rose at Yale School of Medicine though Dr. A. Phillip West at Texas A&M Health Science Center.

Vesicular stomatitis Virus (VSV) infections

RAW 264.7 macrophages were seeded in 12-well tissue culture-treated plates at a density of
$7x10^5$ cells per well 16h before infection. The next day cells were infected with VSV-GFP virus
(Dalton and Rose, 2001) at an MOI of 0.1 in serum-free DMEM (HyClone). After 1h of incuba-
tion with media containing virus, supernatant was removed, and fresh DMEM plus 10% FBS
(Millipore) was added to each well. At indicated times post infection, cells were harvested with
TRIzol for RNA isolation described below.
1106
M. musculus
Mice used in this study were C57BL6/J (Stock #00064) initially purchased from Jackson Labra-
tory and afterward maintained with filial breeding. All mice used in experiments were compared
to age- and sex- matched controls. Littermates were used for experiments. Mice used to gener-
ate BMDMs were males between 10-16 weeks old. Embryos used to make primary MEFs were
14.5 days post-coitum. All animals were housed, bred, and studied at Texas A&M Health Sci-
ence Center under approved Institutional Care and Use Committee guidelines. All experiments
for this study were reviewed and approved by the Texas A&M University Institutional Animal
Care and Use Committee (AUP# 2019–0083). Mice were fed 4% standard chow and were kept
on a 12 h light/dark cycle and provided food and water _ad libitum_. Mice were group housed
(maximum 5 per cage) by sex on ventilated racks in temperature-controlled rooms.
1118
Primary cell culture
Bone marrow derived macrophages (BMDMs) were differentiated from bone marrow (BM) cells
isolated by washing mouse femurs with 10 ml DMEM 1 mM sodium pyruvate. Cells were then
centrifuged for 5 min at 400 rcf and resuspended in BMDM media (DMEM, 20% FBS (Millipore),
1 mM sodium pyruvate (Lonza), 10% MCSF conditioned media (Watson lab)). BM cells were
counted and plated at $5x10^6$ cells per 15 cm non-tissue culture treated dishes in 30 ml complete
BMDM media. Cells were fed with an additional 15 ml of BMDM media on day 3. Cells were har-
vested on day 7 with 1X PBS EDTA (Lonza).
Mouse embryonic fibroblasts (MEFs) were isolated from embryos. Briefly, embryos were dis-
sected from yolk sacs, washed two times with cold 1X PBS, decapitated, and stripped of perito-
neal contents. Headless embryos were disaggregated in cold 0.05% trypsin-EDTA (Lonza) and
incubated on ice for 20 min., followed by incubation at 37 °C for an additional 20 min. Cells were
then DNase treated with 4 ml disaggregation media (DMEM, 10% FBS, 100 µg/ml DNASE I
(Worthington)) for 20 min at 37 °C. Isolated supernatants were spun down at 1000 rpm for 5
min. Cells were resuspended in complete MEF media (DMEM, 10% FBS, 1 mM sodium py-
ruvate), and plated in 15 cm tissue culture-treated dishes 1 dish per embryo in 25 ml of media.
MEFs were allowed to expand for 2-3 days before harvest with 0.05% trypsin-EDTA (Lonza).
iBMDMs were immortalized with J2 virus to generate iBMDMs.
1137
Cell culture
RAW 264.7 macrophages (TIB-71™ ATCC) (originally isolated from male BALB/c mice), L929
ISRE (Hoffpauir et al., 2020), _Irf7_ KO RAW 264.7 macrophages (this paper), and Lenti-X cells
(TaKaRa Bio) were cultured at 37 °C with a humidified atmosphere of 5% CO_2 in complete me-
dia containing high glucose, DMEM (ThermoFisher) with 10% FBS (Millipore) 0.2% HEPES
(Thermo Fisher).
1144
shRNA knockdowns
For RAW 264.7 macrophages stably expressing scramble knockdown and _Srsf7_ knockdown,
Lenti-X cells were transfected with a pSICO scramble non-targeting shRNA construct and
pSICO _Srsf7_ shRNA constructs targeted at exon 3 and exon 4 of _Srsf7_ using Polyjet (SignaGen
Laboratories). Virus was collected 24 and 48 h post transfection. RAW 264.6 macrophages

were transduced using Lipofectamine 2000 (Thermo Fischer). After 48 h, media was supple-
mented with Hygromycin (Invitrogen) to select for cells containing the shRNA plasmid.
1152
<u>siRNA knockdowns in MEFs</u>
Knockdown of mRNA transcripts was performed by plating $3x10^6$ MEFs in 10 cm plates and
rested overnight. The following day complete media was replaced with 5 ml fresh complete me-
dia 30 min prior to transfection. Cells were transfected using Fugene SI reagent 10 µM of siRNA
stock against either *Srsf7* (ThermoFisher, s105011) or negative control, Silencer® Select Nega-
tive Control #1 (4390843). Cells were incubated for 48h in transfection media at 37 °C with 5%
CO_2 prior to downstream experiments.
1160
<u>siRNA knockdowns in BMDMs</u>
The day prior to transfection, 6x10e5 BMDMs were seeded into 12-well plates. The day of trans-
fection, 50nM of siRNA against *Srsf7* (ThermoFisher, s105011) or negative control, Silencer®
Select Negative Control #1 (4390843) were mixed with 3 µl of LipoRNAiMAX (13778150,
Thermo Fisher Scientific), incubated for 10m at room temperature, and then added into wells
with 500 µl plain DMEM. Transfection proceeded for 6h and then wells were replenished with
complete media.
1168
<u>Tetracycline inducible cell line generation</u>
For generation of tetracycline inducible RAW 264.7 macrophages, pLenti CMV rtTA3 Blast
(Addgene w756-1) stably expressing clonal RAW 264.7 macrophages were transduced with
pLenti CMV Puro DEST (Addgene w118-1) constructs containing 3xFL-GFP, 3xFL-SRSF7, and
al 3xFL-SRSF7 phosphorylation mutants. After 48 h construct containing cells were selected
through addition of puromycin (Invivogen). 1 mg/ml doxycycline (Sigma-Aldrich) treatment was
used to activate construct expression.
1176
<u>Cell stimulation for immune activation</u>
$7.5x10^5$ RAW 264.7 macrophage were plated on 12-well tissue-culture treated plates 16h before
cellular stimulation assays. The next day, cells were treated with lipopolysaccharide (LPS) from
E. coli (Invivogen) at 100 ng/ml, or ISD (IDT, annealed in house) at 1 µg/ml, or IFNβ (PBL Assay
Science) at 200 I/U per ml for the respective time points. Cells were collected for RNA isolation
using Trizol reagent.
1183
<u>*Srsf7* KD mRNA sequencing and analysis</u>
As reported in Wagner, Scott and West et al. 2021, RNA-seq analysis was performed on RAW
264.7 cells containing shRNA knockdowns of *Srsf6*, *Srsf2*, *Srsf1*, *Srsf7*, and *Srsf9* compared to
SCR control with biological triplicates of each cell line. RNA-Seq and library preparation was
performed by Texas A&M AgriLife Genomics and Bioinformatics Service. RNA samples were
sequenced on Illumina 4000 using 2 × 150-bp paired-end reads. Raw reads were filtered and
trimmed and Fastq data was mapped to the Mus musculus Reference genome (RefSeq) using
CLC Genomics Workbench 8.0.1. Differential expression analyses were performed using CLC
Genomics Workbench. Relative transcript expression was calculated by counting Reads Per
Kilobase of exon model per Million mapped reads (RPKM). Statistical significance was deter-
mined by the EDGE test *via* CLC Genomics Workbench. Differentially expressed genes (DEGs)
were selected as those with p value threshold < 0.05. Critical pathways affected by SR protein
knockdown compared to control RAW 264.7 macrophages, were analyzed by canonical path-
way analysis using Ingenuity Pathway Analysis software from QIAGEN Bioinformatics. Genes
that were differentially expressed with a p-value < 0.05 from our RNA-SEQ analysis were used
as input. Integrated genomics viewer (IGV) visualization tracks were generated using the RNA-
SEQ filtered and trimmed reads.

1201
Irf7 KO mRNA sequencing and analysis
RNA- seq analysis was performed on control GFP KO or *Irf7* KO1 RAW 264.7 cells in biological
triplicate of each cell line. Data was analyzed by ROSALIND® (https://rosalind.bio/), with a
HyperScale architecture developed by ROSALIND, Inc. (San Diego, CA). Reads were trimmed
using cutadapt[1]. Quality scores were assessed using FastQC[2]. Reads were aligned to the Mus
musculus genome build GRCm39 using STAR[3]. Individual sample reads were quantified using
HTseq[4] and normalized via Relative Log Expression (RLE) using DESeq2 R library[5]. Read Dis-
tribution percentages, violin plots, identity heatmaps, and sample MDS plots were generated as
part of the QC step using RSeQC[6]. DEseq2 was also used to calculate fold changes and p-val-
ues and perform optional covariate correction. Clustering of genes for the final heatmap of dif-
ferentially expressed genes was done using the PAM (Partitioning Around Medoids) method us-
ing the fpc R library[7]. Hypergeometric distribution was used to analyze the enrichment of path-
ways, gene ontology, domain structure, and other ontologies. The topGO R library[8,] was used to
determine local similarities and dependencies between GO terms in order to perform Elim prun-
ing correction.
1217
Alternative splicing analysis
Modeling Alternative Junction Inclusion Quantification (MAJIQ) and the Voila visualization pack-
age were used to model alternative splicing events with default parameters described by
Vaquero-Garcia et al., 2016. Junction-spanning reads were used to construct splice graphs for
transcripts by using the RefSeq annotation supplemented with de-novo (junctions not in the Ref-
Seq transcriptome database but had sufficient evidence in the RNAseq data) detected junctions.
Resulting splice graphs were analyzed for all identified local splice variations (LSVs). MAJIQ
then quantified expected percent spliced in (PSI) value in control and knockdown samples for
every junction in each LSV. Resulted viewed on VOILA were filtered for highest confidence
change in LSVs (one or more junctions had at least a 95% probability of expected dPSI of at
least an absolute value of 0.2 PSI units between control and knockdown.
1229
RNA isolation and qPCR analysis
For transcript analysis, cells and tissue were harvested in TRIzol and RNA was isolated using
Direct-zol RNA Miniprep kits (Zymo Research) with 1 h DNase treatment. cDNA was synthe-
sized with iScript cDNA Synthesis Kit (Bio-Rad). CDNA was diluted as appropriate for treatment,
and a pool of cDNA was used to make a 1:3 standard curve with each standard sample diluted
1:5 to produce a linear curve. RT-qPCR was performed using Power-Up SYBR Green Master
Mix (Thermo Fisher) using a Quant Studio Flex6 (Applied Biosystems). Samples were run in
triplicate wells in a 384-well plate. Averages of the raw values were normalized to average val-
ues for the same sample with the control gene, *Actb*.
1239
Semiquantitative qPCR analysis
cDNA was synthesized by iScript cDNA Synthesis Kit (Bio-Rad) using an extended 1h amplifica-
tion. Q5 high fidelity 2X Master mix (New England Biolabs) was used for PCR amplification us-
ing targeted primers. Loading dye was added to PCR products and samples were run on 2%
agarose gel containing ethidium bromide at 100 volts for 1 h. Gels were imaged on LiCOR Od-
yssey Fc Dual-Mode Imaging System and bands were quantified.
1246
Chromatin immunoprepitation
6×10^6 RAW 264.7 cells or BMDMs were seeded in a 10 cm dish, and 16h later stimulated for the
indicated time. After stimulation cells were scraped, transferred to a 50 ml conical and fixed with
1% (v/v) formaldehyde for 10 min with gentle shaking. Formaldehyde was quenched with 0.125
M glycine for 5 min. Cells were pelleted and washed 1x with PBS, and then cells were lysed in 1

ml of ChIP lysis buffer (150 mM NaCl, 5 mM EDTA, 0.5% NP-40 and 1% Triton X-100, 50 mM
Tris-HCl, and protease/phosphatase inhibitor (added immediately before lysis) pH 7.5). Lysates
were flash frozen, or ChIP proceeded by sonicating with for 40 min (30 sec on/off 10 min cycles)
in a Bioruptor UCD-200 (Diagenode). Lysates were cleared and 10% taken for input samples.
Endogneous ChIP: For RNA polymerase 2 samples: 2 µg of the anti-pol II (Active Motif) or 2 µg
of isotype control (IgG, Cell Signaling) were incubated overnight at 4 with rotation. The following
day, Piece beads (Invitrogen) were washed 3x in ChIP buffer, blocked for 30 min with 100 µg
salmon sperm DNA (Invitrogen) and 0.5 mg BSA, and washed once more with ChIP lysis buffer.
Blocked beads were incubated with cleared chromatin immunocomplexes for 1h at 4 °C with ro-
tation. Beads were washed five times with ChIP buffer and then eluted with 250 µl of elution
buffer (1% SDS and 0.1 M NaHCO$_3$) at 65 °C for 15 mins. The eluate and input were centrifuged
and reverse the cross-links by adding 0.4 M NaCl for overnight at 55 °C. The next day, 1 µl of
RNASE A (10 mg/ml, ThermoFisher) and 2 µl proteinase K (20 mg/ml, ThermoFisher) were
added to the samples and incubated for 1h at 55 °C while shaking. Samples were then sub-
jected to Genomic Clean up and Concentrator (Zymo) to extract DNA. Purified input and IP DNA
were analyzed by qRT-PCR using primers specific for the promoter region of target genes. Re-
sults were normalized to input. Data is presented as fold enrichment relative to intergenic con-
trol. Flag ChIP: As above except cleared lysates were added to blocked (as above) M2 anti-
3xFLAG beads and protocol proceeded as above.
1271
3x-FLAG ChIPseq
Data was analyzed by ROSALIND® (https://rosalind.bio/), with a HyperScale architecture devel-
oped by ROSALIND, Inc. (San Diego, CA). Reads were trimmed using cutadapt[1]. Quality scores
were assessed using FastQC[2]. Reads were aligned to the Mus musculus genome build mm10
using bowtie2[3]. Per-sample quality assessment plots were generated with HOMER[4] and Mosa-
ics[5]. Peaks were called using MACS2[6] (with input/IgG controls background subtracted, if pro-
vided). Peak overlaps and differential binding were calculated using the DiffBind R library[7]. Dif-
ferential binding was calculated at gene promoter sites. Read distribution percentages, identity
heatmaps, and FRiP plots were generated as part of the QC step using ChIPQC R library[8] and
HOMER. HOMER was also used to generate known and de novo motifs and perform functional
enrichment analysis of pathways, gene ontology, domain structure and other ontologies.
1283
CRISPR/Cas9 KO of Irf7
Guide RNAs were generated to target the 3rd exon of Irf7 using the Broad online tool
(https://portals.broadinstitute.org/gppx/crispick/public) and synthesized by IDT. Primers were
cloned into digested vector by phosphorylating, annealing, and ligating primers into LentiGuide-
Puro (Addgene, plasmid 52963). Colonies were validated by Sanger sequencing using
pLK0.1/hU6 promoter primer (Eton Biosciences, San Diego, CA). Lentivirus with sgRNAs were
produced and used to transduce RAW 264.7 cells stably expressing FL-Cas9 (Addgene, plas-
mid 52962). After selection with 5 µg/ml puromycin knockout efficiency was analyzed by im-
munoblot and then sequencing for mutants by amplifying a 300 base pair region of genomic
DNA around exon 3. These PCR fragments were sequenced (as above) and compared to con-
trols using TIDE analysis. The two best, most efficient sgRNA populations were clonally se-
lected, expanded, and assayed for clones with total KO of IRF7 by immunoblot.
1296
Protein quantification by immunoblot
Cells were washed with PBS and lysed in 1X RIPA buffer with protease and phosphatase EDTA
free inhibitors (Thermo Scientific), with the addition of 1 U/ml Benzonase (Millipore) to degrade
genomic DNA. Proteins were separated by SDS-PAGE on AnykD mini -PROTEAN TGX precast
gel (Bio-Rad) and transferred to 0.45µm nitrocellulose membranes (GE Healthcare). Mem-
branes were blocked for 1h at RT in TBS +5% BSA. Blots were incubated overnight at 4 °C with

the following primary antibodies: β-ACTIN (Abcam, 1:5000), Y701 STAT1 (Cell Signaling,
1:1000), STAT1 (Cell Signaling, 1:1000), IRF7 (Cell Signaling, 1:1000), SRSF7 (Bethyl 1:3000)
ZBP1 (Santa Cruz Biotechnologies, 1:1000), p-IRF3(S396) (Cell Signaling, 1:1000); IRF3
(Bethyl, 1:1000); ANTI-FLAG M2 (Sigma Aldrich, 1:5000); Membranes were washed 3X for 5
min in PBS-Tween20 and incubated with appropriate secondary antibodies (LI-COR) for 1h at
RT prior to imaging on a LiCOR Odyssey Fc Dual-Mode Imaging System.
1309
1310   <u>Quantitation and Statistical Analysis</u>
1311   Statistical analysis of data was performed using GraphPad Prism software. Two-tailed unpaired
1312   Student's t tests were used for statistical analyses, and unless otherwise noted, all results are
1313   representative of at least three biological samples (mean +/- SEM (n = 3 per group)).
1314
1315
1316

REFERENCES

Barrat, F.J., Crow, M.K., and Ivashkiv, L.B. (2019). Interferon target-gene expression and epigenomic signatures in health and disease. Nat Immunol *20*, 1574-1583.

Bell, S.L., Lopez, K.L., Cox, J.S., Patrick, K.L., and Watson, R.O. (2021). Galectin-8 Senses Phagosomal Damage and Recruits Selective Autophagy Adapter TAX1BP1 To Control Mycobacterium tuberculosis Infection in Macrophages. mBio *12*, e0187120.

Budzik, J.M., Swaney, D.L., Jimenez-Morales, D., Johnson, J.R., Garelis, N.E., Repasy, T., Roberts, A.W., Popov, L.M., Parry, T.J., Pratt, D.*, et al.* (2020). Dynamic post-translational modification profiling of Mycobacterium tuberculosis-infected primary macrophages. Elife *9*.

Cavaloc, Y., Bourgeois, C.F., Kister, L., and Stevenin, J. (1999). The splicing factors 9G8 and SRp20 transactivate splicing through different and specific enhancers. RNA *5*, 468-483.

Cavaloc, Y., Popielarz, M., Fuchs, J.P., Gattoni, R., and Stevenin, J. (1994). Characterization and cloning of the human splicing factor 9G8: a novel 35 kDa factor of the serine/arginine protein family. EMBO J *13*, 2639-2649.

Colwill, K., Pawson, T., Andrews, B., Prasad, J., Manley, J.L., Bell, J.C., and Duncan, P.I. (1996). The Clk/Sty protein kinase phosphorylates SR splicing factors and regulates their intranuclear distribution. EMBO J *15*, 265-275.

Curran, E.C., Wang, H., Hinds, T.R., Zheng, N., and Wang, E.H. (2018). Zinc knuckle of TAF1 is a DNA binding module critical for TFIID promoter occupancy. Sci Rep *8*, 4630.

de la Mata, M., and Kornblihtt, A.R. (2006). RNA polymerase II C-terminal domain mediates regulation of alternative splicing by SRp20. Nat Struct Mol Biol *13*, 973-980.

Duncan, P.I., Howell, B.W., Marius, R.M., Drmanic, S., Douville, E.M., and Bell, J.C. (1995). Alternative splicing of STY, a nuclear dual specificity kinase. J Biol Chem *270*, 21524-21531.

Fitzgerald, K.A., Rowe, D.C., Barnes, B.J., Caffrey, D.R., Visintin, A., Latz, E., Monks, B., Pitha, P.M., and Golenbock, D.T. (2003). LPS-TLR4 signaling to IRF-3/7 and NF-kappaB involves the toll adapters TRAM and TRIF. J Exp Med *198*, 1043-1055.

Frankiw, L., Majumdar, D., Burns, C., Vlach, L., Moradian, A., Sweredoski, M.J., and Baltimore, D. (2019). BUD13 Promotes a Type I Interferon Response by Countering Intron Retention in Irf7. Mol Cell *73*, 803-814 e806.

Gao, D., Ciancanelli, M.J., Zhang, P., Harschnitz, O., Bondet, V., Hasek, M., Chen, J., Mu, X., Itan, Y., Cobat, A.*, et al.* (2021). TLR3 controls constitutive IFN-beta antiviral immunity in human fibroblasts and cortical neurons. J Clin Invest *131*.

Gough, D.J., Messina, N.L., Clarke, C.J., Johnstone, R.W., and Levy, D.E. (2012). Constitutive type I interferon modulates homeostatic balance through tonic signaling. Immunity *36*, 166-174.

Green, I.D., Pinello, N., Song, R., Lee, Q., Halstead, J.M., Kwok, C.T., Wong, A.C.H., Nair, S.S., Clark, S.J., Roediger, B.*, et al.* (2020). Macrophage development and activation involve coordinated intron retention in key inflammatory regulators. Nucleic Acids Res *48*, 6513-6529.

1367 Gu, J., Shi, J., Wu, S., Jin, N., Qian, W., Zhou, J., Iqbal, I.G., Iqbal, K., Gong, C.X., and Liu, F.
1368 (2012). Cyclic AMP-dependent protein kinase regulates 9G8-mediated alternative splicing of tau
1369 exon 10. FEBS Lett *586*, 2239-2244.
1370
1371 Gu, L., Casserly, D., Brady, G., Carpenter, S., Bracken, A.P., Fitzgerald, K.A., Unterholzner, L.,
1372 and Bowie, A.G. (2022). Myeloid cell nuclear differentiation antigen controls the pathogen-
1373 stimulated type I interferon cascade in human monocytes by transcriptional regulation of IRF7.
1374 Nat Commun *13*, 14.
1375
1376 Gui, J.F., Lane, W.S., and Fu, X.D. (1994). A serine kinase regulates intracellular localization of
1377 splicing factors in the cell cycle. Nature *369*, 678-682.
1378
1379 Hoffpauir, C.T., Bell, S.L., West, K.O., Jing, T., Wagner, A.R., Torres-Odio, S., Cox, J.S., West,
1380 A.P., Li, P., Patrick, K.L.*, et al.* (2020). TRIM14 Is a Key Regulator of the Type I IFN Response
1381 during Mycobacterium tuberculosis Infection. J Immunol *205*, 153-167.
1382
1383 Honda, K., and Taniguchi, T. (2006). IRFs: master regulators of signalling by Toll-like receptors
1384 and cytosolic pattern-recognition receptors. Nat Rev Immunol *6*, 644-658.
1385
1386 Honda, K., Yanai, H., Negishi, H., Asagiri, M., Sato, M., Mizutani, T., Shimada, N., Ohba, Y.,
1387 Takaoka, A., Yoshida, N.*, et al.* (2005). IRF-7 is the master regulator of type-I interferon-
1388 dependent immune responses. Nature *434*, 772-777.
1389
1390 Huang, R., Grishagin, I., Wang, Y., Zhao, T., Greene, J., Obenauer, J.C., Ngan, D., Nguyen,
1391 D.T., Guha, R., Jadhav, A.*, et al.* (2019). The NCATS BioPlanet - An Integrated Platform for
1392 Exploring the Universe of Cellular Signaling Pathways for Toxicology, Systems Biology, and
1393 Chemical Genomics. Front Pharmacol *10*, 445.
1394
1395 Jefferies, C.A. (2019). Regulating IRFs in IFN Driven Disease. Front Immunol *10*, 325.
1396 Jeong, S. (2017). SR Proteins: Binders, Regulators, and Connectors of RNA. Mol Cells *40*, 1-9.
1397
1398 Ji, X., Zhou, Y., Pandit, S., Huang, J., Li, H., Lin, C.Y., Xiao, R., Burge, C.B., and Fu, X.D.
1399 (2013). SR proteins collaborate with 7SK and promoter-associated nascent RNA to release
1400 paused polymerase. Cell *153*, 855-868.
1401
1402 Kadota, Y., Jam, F.A., Yukiue, H., Terakado, I., Morimune, T., Tano, A., Tanaka, Y., Akahane,
1403 S., Fukumura, M., Tooyama, I.*, et al.* (2020). Srsf7 Establishes the Juvenile Transcriptome
1404 through Age-Dependent Alternative Splicing in Mice. iScience *23*, 100929.
1405
1406 Kajigaya, H., Ishibashi, T., Hayashi, A., Yamaguchi, Y., and Baba, H. (2016). Concentration of
1407 neddylation-related molecules in paranodal myelin of the peripheral nervous system. Proc Jpn
1408 Acad Ser B Phys Biol Sci *92*, 56-68.
1409
1410 Konigs, V., de Oliveira Freitas Machado, C., Arnold, B., Blumel, N., Solovyeva, A., Lobbert, S.,
1411 Schafranek, M., Ruiz De Los Mozos, I., Wittig, I., McNicoll, F.*, et al.* (2020). SRSF7 maintains its
1412 homeostasis through the expression of Split-ORFs and nuclear body assembly. Nat Struct Mol
1413 Biol *27*, 260-273.
1414
1415 Loomis, R.J., Naoe, Y., Parker, J.B., Savic, V., Bozovsky, M.R., Macfarlan, T., Manley, J.L., and
1416 Chakravarti, D. (2009). Chromatin binding of SRp20 and ASF/SF2 and dissociation from mitotic
1417 chromosomes is modulated by histone H3 serine 10 phosphorylation. Mol Cell *33*, 450-461.

Luco, R.F., Allo, M., Schor, I.E., Kornblihtt, A.R., and Misteli, T. (2011). Epigenetics in alternative pre-mRNA splicing. Cell *144*, 16-26.

Luco, R.F., Pan, Q., Tominaga, K., Blencowe, B.J., Pereira-Smith, O.M., and Misteli, T. (2010). Regulation of alternative splicing by histone modifications. Science *327*, 996-1000.

Muller-McNicoll, M., Botti, V., de Jesus Domingues, A.M., Brandl, H., Schwich, O.D., Steiner, M.C., Curk, T., Poser, I., Zarnack, K., and Neugebauer, K.M. (2016). SR proteins are NXF1 adaptors that link alternative RNA processing to mRNA export. Genes Dev *30*, 553-566.

Nau, G.J., Richmond, J.F., Schlesinger, A., Jennings, E.G., Lander, E.S., and Young, R.A. (2002). Human macrophage activation programs induced by bacterial pathogens. Proc Natl Acad Sci U S A *99*, 1503-1508.

Ning, S., Pagano, J.S., and Barber, G.N. (2011). IRF7: activation, regulation, modification and function. Genes Immun *12*, 399-414.

Pandey, A., Ding, S.L., Qin, Q.M., Gupta, R., Gomez, G., Lin, F., Feng, X., Fachini da Costa, L., Chaki, S.P., Katepalli, M.*, et al.* (2017). Global Reprogramming of Host Kinase Signaling in Response to Fungal Infection. Cell Host Microbe *21*, 637-649 e636.

Patrick, K.L., Bell, S.L., and Watson, R.O. (2016). For Better or Worse: Cytosolic DNA Sensing during Intracellular Bacterial Infection Induces Potent Innate Immune Responses. J Mol Biol *428*, 3372-3386.

Schoggins, J.W. (2019). Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol *6*, 567-584.

Schoggins, J.W., and Rice, C.M. (2011). Interferon-stimulated genes and their antiviral effector functions. Curr Opin Virol *1*, 519-525.

Schoggins, J.W., Wilson, S.J., Panis, M., Murphy, M.Y., Jones, C.T., Bieniasz, P., and Rice, C.M. (2011). A diverse range of gene products are effectors of the type I interferon antiviral response. Nature *472*, 481-485.

Sin, W.X., Yeong, J.P., Lim, T.J.F., Su, I.H., Connolly, J.E., and Chin, K.C. (2020). IRF-7 Mediates Type I IFN Responses in Endotoxin-Challenged Mice. Front Immunol *11*, 640.
Tellier, M., Maudlin, I., and Murphy, S. (2020). Transcription and splicing: A two-way street. Wiley Interdiscip Rev RNA *11*, e1593.

Vaquero-Garcia, J., Barrera, A., Gazzara, M.R., Gonzalez-Vallinas, J., Lahens, N.F., Hogenesch, J.B., Lynch, K.W., and Barash, Y. (2016). A new view of transcriptome complexity and regulation through the lens of local splicing variations. Elife *5*, e11752.

Wagner, A.R., Scott, H.M., West, K.O., Vail, K.J., Fitzsimons, T.C., Coleman, A.K., Carter, K.E., Watson, R.O., and Patrick, K.L. (2021). Global Transcriptomics Uncovers Distinct Contributions From Splicing Regulatory Proteins to the Macrophage Innate Immune Response. Front Immunol *12*, 656885.

1467   Wagner, A.R., Weindel, C.G., West, K.O., Scott, H.M., Watson, R.O., and Patrick, K.L. (2022).
1468   SRSF6 balances mitochondrial-driven innate immune outcomes through alternative splicing of
1469   BAX. Elife *11*.
1470
1471   Wagner, R.E., and Frye, M. (2021). Noncanonical functions of the serine-arginine-rich splicing
1472   factor (SR) family of proteins in development and disease. Bioessays *43*, e2000242.
1473
1474   Wang, Y., Yu, Y., Pang, Y., Yu, H., Zhang, W., Zhao, X., and Yu, J. (2021). The distinct roles of
1475   zinc finger CCHC-type (ZCCHC) superfamily proteins in the regulation of RNA metabolism.
1476   RNA Biol *18*, 2107-2126.
1477
1478   Watson, R.O., Bell, S.L., MacDuff, D.A., Kimmey, J.M., Diner, E.J., Olivas, J., Vance, R.E.,
1479   Stallings, C.L., Virgin, H.W., and Cox, J.S. (2015). The Cytosolic Sensor cGAS Detects
1480   Mycobacterium tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. Cell
1481   Host Microbe *17*, 811-819.
1482
1483   West, K.O., Scott, H.M., Torres-Odio, S., West, A.P., Patrick, K.L., and Watson, R.O. (2019).
1484   The Splicing Factor hnRNP M Is a Critical Regulator of Innate Immune Gene Expression in
1485   Macrophages. Cell Rep *29*, 1594-1609 e1595.
1486