E2F1 promotes, JNK and DIAP1 inhibit, and chromosomal position has little effect on

radiation-induced Loss of Heterozygosity in *Drosophila*

Jeremy Brown[1] and Tin Tin Su[2,3]

Department of Molecular, Cellular and Developmental Biology, 347 UCB,

University of Colorado, Boulder, CO 80309-0347

[1] University of Colorado, Anschutz Medical Campus, 13001 E. 17th Pl., Aurora, CO 80045

[2] University of Colorado Cancer Center, Anschutz Medical Campus, 13001 E. 17th Pl., Aurora, CO

80045

[3]Corresponding author: tin.su@colorado.edu

Short title: Determinants of radiation-induced genome instability

Keywords: *Drosophila*, radiation, loss of heterozygosity

Abstract

Loss of Heterozygosity (LOH) can occur when a heterozygous mutant cell loses the remaining

wild type allele to become a homozygous mutant. LOH can have physiological consequences if,

for example, the affected gene encodes a tumor suppressor. We used two fluorescent

reporters to study mechanisms of LOH induction by X-rays, a type of ionizing radiation (IR), in

Drosophila larval wing discs. IR is used to treat more than half of cancer patients, so

understanding its effects is of biomedical relevance. We report that IR-induced LOH does not

correlate with the chromosomal position of the LOH locus, unlike previously shown for

spontaneously occurring LOH. Like spontaneous LOH, however, IR-induced LOH of X-linked loci

shows a sex-dependence, occurring predominately in females. A focused genetic screen

identified E2F1 as a key promotor of LOH and further testing suggests a mechanism involving its

role in cell cycle regulation rather than apoptosis. We combined the QF/QS LOH reporter with

QUAS-transgenes to manipulate gene function after LOH induction. This approach identified

JNK signaling and apoptosis as key determinants of LOH maintenance. These studies reveal

previously unknown mechanisms for generation and maintenance of cells with chromosome

aberrations after exposure to IR.

Introduction

Loss of Heterozygosity (LOH) occurs when a heterozygous cell loses one of the alleles to become homozygous. LOH can have physiological consequences if, for example, the affected allele encodes a tumor suppressor. Possible mechanisms for the loss of a genetic locus are many and include mutations in the gene, aneuploidy involving loss of whole or segments of chromosomes, and mitotic recombination between the locus and the centromere (producing -/- and +/+ sisters from a -/+ mother cell). We have been using *Drosophila melanogaster* as a model to study LOH that follows exposure to X-rays, a type of ionizing radiation (IR). IR causes single and double strand breaks in the DNA and its ability to induce apoptosis is the reason IR is used to treat half of cancer patients. Understanding IR's effects on genome integrity is a crucial step towards rendering radiation therapy safe by reducing side effects.

Classical and recent studies of IR-induced LOH in *Drosophila* employed visible adult markers such as *multiple wing hair* (*mwh*) and *forked (f)* that alter the number or morphology of actin-based hairs on the cell surface. Starting with heterozygotes of recessive alleles, induced homozygosity is detected by the presence of mutant hairs and scored on a cell-by-cell basis. In these studies, exposure to IR occurred in the larval stages while the resulting LOH was scored many days later in the adult, with intervening steps poorly understood. To monitor these steps, we have been developing fluorescent reporters for LOH that can be used at all stages of development. We focus our studies on larval wing imaginal discs that develop into the adult wing, the body wall and the hinge that connects the two. Unlike the bulk of the larvae that is made up of polyploid cells that do not undergo mitosis, imaginal discs are comprised of diploid epithelial cells that proliferate by mitotic division.

In our published work with a fluorescent LOH reporter [1], we used the QF/QS module from the bread mold *Neurospora crassa* wherein QF binds its recognition sequence QUAS to promote transcription while QS represses QF [2]. psc-QF can drive the expression of QUAS-tdTomato (Tom), a fluorescent protein from coral, throughout the wing disc (Fig 1A) unless repressed by ubiquitously-expressed QS^{9B} inserted at 100E1 on chromosome 3 (Fig 1F) [3]. Loss of QS^{9B} by

LOH results in de-repression of QF and cell-autonomous Tom expression (Fig 1C-C'). Using this

system, we identified two phases of culling of LOH cells, one in late larval stages and another

during the pupa stage, that show differential genetic requirements [1].

Our published study produced two surprising results. First, the incidence of IR-induced LOH as

detected by the adult wing marker *mwh* was reproducibly lower by greater than 10-fold when

compared to IR-induced LOH detected by QS^{9B} under the same experimental conditions in adult

wings [1]. Second, p53, which is required to reduce *mwh* LOH both in our study and others' [1,

4], had little effect on QS^{9B} LOH [1]. Classical studies using adult markers suggest a possible

explanation. These studies found that 60% of homozygous mutant LOH clones induced by IR

lacked a wild type (+/+) twin clone that is expected if mitotic recombination was the

mechanism [5]. The same study found that in spontaneously arising LOH clones in un-irradiated

animals, 20% of the cells showed a *Minute* bristle phenotype, which is caused by heterozygosity

of ribosomal protein loci that are scattered throughout the genome. These data led to the

proposal that segmental aneuploidy (loss of a chromosome segment) is the primary mechanism

for IR-induced LOH. If breakage and loss of the corresponding chromosome tip is indeed the

mechanism, *mwh* LOH would accompany the loss of more genes compared to QS^{9B} LOH

because *mwh* is further from the telomere than QS^{9B} (Fig 1F). Resulting *mwh* LOH cells may be

less fit than QS^{9B} LOH cells, explaining the difference in IR-induced LOH incidence and p53-

dependence we saw. Alternatively, cells that are homozygous mutant for *mwh*, which encodes

an actin-bundling protein, may be less fit while the loss of QS, which has no known role in

Drosophila, may be neutral and it is this difference in gene function that explains the

differences in IR-induced LOH incidence and p53-dependence.

To address these possibilities and to identify regulators of IR-induced LOH, we studied QS^{5A},

another QS insertion that is further from the telomere than QS^{9B}, along with five GAL80

insertions that span the distance from the telomere to the centromere along the X

chromosome (Fig 1F) and performed a focused genetic screen.

Results

Distance from the telomere makes little difference in IR-induced LOH

We used 4000 Rads (R) of X-rays to induce LOH, a dose used by many labs (for example, [6, 7]) because it kills more than half of the cells in wing discs by apoptosis, yet most larvae develop into pupae and eclose into viable adults. Thus, this dose of IR causes significant damage but is still compatible with recovery, regeneration, and development. To ask if the distance from the chromosome tip matters, we studied the IR-induced loss of QS[5A] that is further from the telomere than QS[9B] or *mwh* (Fig 1F). As we reported for QS[9B] [1], LOH for QS[5A] is IR-dependent, with little Tom signal in discs from unirradiated larvae (Fig 2A-B). More important, we saw no statistically significant difference between QS[9B] and QS[5A] in terms of IR-induced LOH clone number (Fig 2A) or clone area (Fig 2B). At the developmental stages we study, cells of the wing disc that will ultimately form adult structures are in a single layer columnar epithelium with uniform cell size. Therefore, clone area serves as a proxy for cell number here, as in previous studies [1]. We normalize clone number and area to the total disc area so that we can control for variations in disc size/total cell number. The results suggest that distance from the chromosome tip did not affect LOH incidence.

Another critical difference we reported between *mwh* and QS[9B] concerned the role of p53. Using the same RNAi construct and identical experimental conditions to deplete p53, we found that p53 was required to cull cells with *mwh* LOH whereas QS[9B] LOH was insensitive to p53 depletion [1]. We reproduce these findings for QS[9B] by expressing the same p53 RNAi construct used before or p53 mutants that lack the DNA binding domain and function as dominant negatives [8]. en-GAL4>UAS-GFP marks the posterior (P) compartment where GAL4-driven transgene expression occurs (Fig 1C') while the anterior (A) compartment serves as an internal control. Clone number and area were normalized to compartment area to control for differences in A/P compartment sizes. Transgene expression was initiated with a temperature shift 24h prior to irradiation with 0 R (-IR control) or 4000 R of X-rays, and LOH was quantified after 2 (0 R) or 3 (4000 R) days after irradiation (protocol in Fig 1E). The results confirmed the

previous finding that QS^{9B} LOH is insensitive to p53 depletion (Fig 2C-D) and allowed us to test

the same on QS^{5A} LOH, which is further from the chromosome tip.

To assess the sensitivity of QS^{5A} to p53 depletion, we generated a recombinant chromosome 2

that bears QS^{5A}, psc-QF>Tom, and en-GAL4>UAS-GFP (this was not needed for QS^{9B} which is on

chromosome 3). We characterized ten independent recombinant QS^{5A} lines and found that they

consistently produced lower LOH than the parental QS^{5A} (data for two recombinants shown in

Fig 2A-B). Perhaps chromosome rearrangements that occurred during recombination interfered

with QS^{5A} or psc>Tom expression. More importantly, two recombinant lines with highest level

of IR-induced LOH showed no change upon p53 depletion (Fig 2C-D). Collectively, these data

suggest that (1) LOH incidence does not correlate with the distance from the telomere and that

(2) not all LOH events are sensitive to p53. We speculate that p53 may respond to LOH events

differently depending on their consequence on cellular physiology, which is a new idea (see

Discussion).

IR-induced LOH shows sex-dependence and locus-independence

To more rigorously test the idea that IR-induced LOH is not dependent on chromosomal

position, we turned to a published GAL4/GAL80 system with GAL80 inserted at 5 different

locations along the X chromosome that has been used to detect spontaneous (without IR) LOH

(Fig 1B,D,F) [9]. For four of the five insertions, wing discs from un-irradiated larvae show very

few cells with GFP signal that is as bright as what is seen after IR (Fig 3A-B, Supplemental Fig 1).

Occasional cells with faint GFP (arrow in Supplemental Fig 1A-A'') are consistent with a recent

report of spontaneous GAL80 'loss' in larval brains through poorly understood mechanisms

[10]. Importantly, we noticed in early experiments with GAL80 that approximately half of the

irradiated discs lacked GFP expressing clones. This is different from what we saw with QS, for

example, 32 of 38 QS^{9B} discs had Tom+ cells in Fig 2A '9B+IR' dataset. QS insertions are on

autosomes while GAL80 insertions are on the X chromosome. This led us to hypothesize a sex-

dependence in GAL80 LOH, similar to what was reported previously for spontaneous LOH of X-

linked loci [9]. To address this possibility, we separated males and female larvae before

dissection and found that IR-induced increases in GFP+ cells were confined almost exclusively to

females (Fig 3C-D).

The exception is the 12C6 insertion, which shows the greatest incidence of GFP+ cells without

IR via unknown mechanisms, and a further increase after IR (Fig 3E-F). Even in this case, IR

induced increase was significant only in the females (Fig 3G-J, quantified in E and F). Thus, IR-

induced GAL80 LOH shows a sex-dependence for all five insertions, similar to the case for

spontaneous LOH [9]. To our surprise, however, IR-induced LOH in terms of clone number or

clone area did not decrease with proximity to the centromere (Fig 3C-D). This is unlike what was

observed before for spontaneous LOH, which decreased with increasing proximity to the

centromere for both X and autosomal loci [9]; GAL80^{5B} showed spontaneous LOH in 75% of

adult *Drosophila* intestines while the same parameter for GAL80^{19E} was 40%. In contrast, IR-

induced LOH clone number and area were nearly identical for GAL80^{5B} and GAL80^{19E} (Fig 3C-D,

p=0.193 for clone number and 0.861 for clone area). We conclude that IR-induced LOH shows

sex-dependence but chromosomal position independence. Possible explanations for these

results are provided in the Discussion.

A focused screen identified E2F1 as a regulator of LOH

Results so far address cis-acting determinants, namely, chromosome position of the LOH loci

and the presence of a homologous X chromosome. To identify trans-acting regulators of LOH,

we performed a focused screen through known regulators of DNA damage and stress

responses, cell growth, cell survival and cell competition (see Supplemental Table 1 for the full

list), using published constructs that have been shown to produce a phenotype, when possible.

We used QS rather than GAL80 in this screen so that we can use the GAL4/UAS system in

parallel to knock down/overexpress genes. We used QS^{9B} rather than QS^{5A} because the former

shows more robust IR-induced LOH incidence.

Genes were knocked down or over-expressed in the posterior (P) half of wing discs using the

protocol in Fig 1E. As stated above, we saw no Tom clones without IR; therefore, only +IR data

are shown (Fig 4 and Supplemental Fig 2). Three controls were used: the parental TRiP III line

that was the recipient of many RNAi constructs and RNAi for the w gene (both as controls for

RNAi) and w[1118] (control for overexpression or OE). IR-induced LOH clone number or area are

not significantly different between A and P compartments in all three controls (Fig 4A-B and

Supplemental Fig 2). Most of the genes tested did not alter IR-induced LOH significantly

(Supplemental Fig 2), including those with known roles in cell competition such as duox,

Myo31D and Xrp1. This group is of interest because a prior publication suggested that cells with

segmental aneuploidy, a possible mechanism for LOH generation, may be eliminated by cell

competition [11]. We interpret negative results with caution, however, because our protocol

for conditional and transient expression (Fig 1E) may not deplete gene function sufficiently

even if the same construct produced a phenotype under different expression conditions before.

The screen found mild effects with RNAi against Chk2 or Myc and with overexpression of

oncogenic Ras^{V12} (Fig 4A-B). Specifically, Chk2 RNAi increased LOH clone number while Myc

RNAi and Ras^{V12} decreased LOH clone number. The effect sizes, however, were small (1.5 to 2-

fold) and clone area was not significantly affected, suggesting compensating changes in the size

of individual clones. In addition, expressing Ras^{V12} in the P compartment, we found, produced

epithelial folds in both P and A compartments in irradiated discs (Fig 4D-D'), making the

quantification of clone area challenging. Because of this and small effect sizes, we have not

followed up on Chk2, Myc and Ras further.

In contrast, we observed a profound effect on both LOH clone number and clone area with

RNAi against E2F1, a reduction of the former by 6-fold and the latter by more than 10-fold (Fig

4C, quantified in A-B). *Drosophila* E2F1, like its mammalian homologs, functions in cell cycle

regulation where it promotes G1/S transition via activation of cyclin E expression and G2/M

transition via activation of String (*Drosophila* Cdc25) [12]. It is also known to promote IR-

induced apoptosis in the wing discs when p53 has been depleted; that is, its role in IR-induced

apoptosis is negligible in a wild type background [13]. Therefore, we addressed the more likely

possibility that the requirement for E2F1 in LOH clone formation/maintenance is through its

role in the cell cycle. To this end, we expressed Cdk inhibitors Dap and Rux to inhibit G1/S and

G2/M transitions, respectively, using the same temperature shift protocol as in Fig 1E. Both Dap

and Rux phenocopied E2F1 RNAi (Fig 4A-B), supporting the idea that the role of E2F1 is to

promote the proliferation of cells with LOH.

As explained above, we normalize LOH clone number and LOH area to the area of the

compartment. If E2F1 RNAi or overexpression of Dap and Rux in the P compartment affected all

cells within this compartment equally regardless of their LOH status, we should see little change

in normalized LOH clone number or area. Yet, we see multiple-fold reductions by both

measures. We infer that manipulations of cell cycle regulators have a more severe effect on

cells with LOH than on cells in the rest of the P compartment. Greater sensitivity of LOH cells to

cell cycle inhibition is a new finding.

The manipulations thus far alter gene expression in the whole compartment and identified

differential effects on LOH versus non-LOH cells. To address cell-autonomous needs in LOH cells

more directly, we took advantage of QS loss and consequent QF de-repression to express

QUAS-transgenes and assayed their effect. This protocol has the added advantage of identifying

needs AFTER the clones have formed and have initiated QF-mediated gene expression. As

proof-of-concept, we used a published QUAS-rpr transgene [14] to express this pro-apoptotic

gene specifically in LOH clones, with the expectation that such clones will be lost to cell death.

The results show that the system works as expected; the addition of QUAS-rpr reduced both

clone number and clone area to near completion (Fig 5A-B, quantified in E-F).

Using QUAS-transgenes to express inhibitors, we tested the role of JNK and apoptotic signaling

in maintaining cells with LOH. We first assessed the functionality of each transgene by

expressing them ubiquitously with the psc-QF promoter (without QS) and assaying for expected

phenotypes. For rpr, dominant negative bsk (*Drosophila* JNK) and puc (phosphatase and JNK

inhibitor), we were able to identify lines that produced significant lethality (Supplemental Table

2). For *Drosophila* Inhibitor of Apoptosis Protein 1 (DIAP1), even the strongest and statistically

significant effect on lethality was modest. Therefore, we assayed for inhibition of apoptosis

instead. Of three lines tested, QUAS-DIAP1 12M reduced IR-induced cleaved Dcp1 signal to near

completion (Fig 5H), 14F showed a modest effect (Fig 5I), and 3M showed variable effect from

disc to disc. Therefore, QUAS-DIAP1 12M line was used in subsequent experiments. The results

show that LOH cell-autonomous inhibition of JNK or apoptosis led to a significant increase in

both Tom clone number and area. We conclude that in cells that survived LOH long enough to

activate QF-mediated transgene expression, stress signaling continues to operate to eliminate

them.

Discussion

We report here studies on the role of chromosomal position and trans-acting regulators on IR-

induced LOH in *Drosophila* larval wing discs. We found that position along the chromosome

makes little difference while the presence of a homologous chromosome is required. We

identified E2F1 as a positive regulator of LOH clone incidence and propose that cells with LOH

are more sensitive to inhibition of cell proliferation than their neighbors. And we identified JNK

signaling and apoptosis as cell-autonomous mechanisms that eliminate cells that survived LOH

induction.

Spontaneous LOH in the adult *Drosophila* intestine is attributed to mitotic recombination,

based on several lines of evidence. First, sex-dependence is interpreted as arising from

recombination-based mechanisms that require the presence of a homologous chromosome

such as crossing over, gene conversion or break-Induced replication [9]. Second, adult guts with

spontaneous GAL80[5B] LOH was about twice as frequent as those with spontaneous GAL80[19E]

LOH. An increase in LOH with distance from the centromere is predicted for recombination-

based mechanisms because a distal locus would experience more frequent exchanges between

homologous chromosomes. This study noted, however, that the frequency of spontaneous LOH

at 19E was higher than predicted from meiotic maps and higher than observed for a

centromere-proximal locus on chromosome 3R, suggesting additional factors at play. Third,

whole genome sequencing confirmed that spontaneous LOH of chromosome 2 loci in the adult

intestine occur through mitotic recombination [15].

Our finding that IR-induced LOH clone number and area were nearly identical for GAL80^{5B} and

GAL80^{19E} cannot be explained by recombination-based mechanisms. Classical studies by Baker

and others monitored several adult visible markers such as *yellow (y)* and *forked (f)* in

heterozygous adult females that were irradiated as larvae or pupae with 1000 R from a Co-60

source [5]. In adult female abdomens monitored for *y* or *f*, both X-linked genes, 40% of LOH

clones induced by IR (i.e., the difference between -/+IR samples) consisted of 40% twin clones

(likely products of mitotic recombination) and the remaining 60% were single clones.

Examination of single clones that arose spontaneously (without radiation) in the same study

found that 20% of cells in such clones showed *Minute* bristles, which result from

haploinsufficiency of loci encoding ribosomal proteins. The authors suggest that single LOH

clones arose by aneuploidy that turned wild type (+/+) cells into heterozygotes for ribosomal

protein genes (M/+). In another set of studies, McNamee and Brodsky used the *Minute* bristle

phenotype and assayed gain-of-heterozygosity, from +/+ to M/+, rather than loss-of-

heterozygosity, in bristles from adults that result from irradiated larvae [11]. Recombination-

based mechanisms induced by IR would not be detectable in this assay as it would change +/+

to +/+. Regardless, taken together, these results suggest that both aneuploidy and

recombination-based mechanisms operate to produce IR-induced LOH in tissues that produce

the adult body wall. Our data suggest a greater contribution by non-recombination-based

mechanisms.

How then do we explain our finding that LOH of X-linked loci requires a homologous

chromosome? We suggest that without part or whole of the X chromosome in XY males, a cell

may not survive to express GFP, reducing the level of LOH observed in males. In other words,

presence of one X chromosome allows cells to survive with partial or complete loss of the other

X chromosome. The results of studies by Baker, Brodsky and colleagues did not distinguish

among various types of aneuploidy, which can include whole chromosome loss, segmental

aneuploidy (loss of a chromosomal segment) and terminal deletions, as specific mechanisms for

IR-induced genomic rearrangements. Over-expression of telomere capping protein HipHop has

been shown to increase the survival of cells with terminal deletions that result from dividing

experimentally induced dicentric chromosomes [16]. Overexpression of HipHop, however, did

not increase the incidence of IR-induced LOH (Supplemental Fig 2), suggesting that terminal

deletions may not be the mechanism. The finding that distance from the telomere did not

affect the QS LOH incidence supports this idea. Instead, relevant types of aneuploidy may

include X-ray-induced segmental losses that are well-documented in numerous studies in

Drosophila and mammals (for example, [17-20]).

McNamee and Brodsky found that p53 single mutants, *hid* single mutants, and p53 *hid* double

mutants show progressively higher incidence of IR-induced *Minute* (M/+) bristles, in that order

[11]. The authors interpreted the data to mean that while p53 makes a partial contribution to

removing M/+ cells, there is also a p53-independent mechanism and that *hid* is involved in

both. Additional data implicate JNK signaling in activating *hid* but only when p53 is absent. In

other words, p53 and JNK play primary and back-up roles, respectively, in eliminating cells with

chromosome aberrations that affect the Minute loci. In contrast, we detect a minimal effect of

p53 depletion and a significant effect of JNK inhibition (even when p53 is wild type) on QS LOH

(this study), and a significant effect of p53 depletion on *mwh* LOH [1]. We propose that the

effect of p53 on cells with LOH depends on whether the LOH affects the fitness of cells (*mwh*

and M) or not (QS and GAL80). The requirement for JNK may be the reverse because we detect

a need for it in repressing the less deleterious QS LOH (this study) while it plays a secondary

role to p53 in culling M/+ cells that experience a lowered fitness [11].

Materials and Methods

Drosophila stocks and methods

Drosophila stocks used in this work are listed in Supplemental Table 1 and include UAS-puc [21],

UAS-Wg [22], and stocks bearing recombinant chromosome 2 of the genotype psc-QF, QUAS-

Tom, en-GAL4, UAS-GFP, tub-QS^{5A} balanced over CyO-GFP and recombinant chromosome 3 of

the genotype tub-QS^{9B}, tub-GAL80^{ts} [1]. Virgin females and males were crossed and cultured for

three days on Nutri-Fly German Formula food (Genesee Scientific) at 25°C before egg

collections.

Larvae culture and irradiation

Eggs were collected and larvae were raised on Nutri-Fly Bloomington Formula food (Genesee

Scientific) at 25°C unless otherwise noted. The cultures were monitored daily for signs of

crowding, typically seen as 'dimples' in the food surface as larvae try to increase the surface

area for access to air. Cultures were split at the first sign of crowding. Larvae in food were

placed in petri dishes and irradiated in a Faxitron Cabinet X-ray System Model RX-650

(Lincolnshire, IL) at 115 kVp and 5.33 rad/sec. Larval sex was determined using gonad size

observed under a dissecting microscope; males have large, clearly visible gonads free of the fat

body while female gonads are hard to find [23]. For eclosion (Supplemental Table 2), empty and

full pupal cases were counted 12-14 days after collection.

Tissue preparation and imaging

Larval wing discs were dissected in PBS, fixed in 4% para-formaldehyde (PFA) in PBS for 30 min,

and washed three times with PBTx (0.1% Triton X-100). The discs were stained with 10 μg/ml

Hoechst33342 in PBTx for 5 min, washed 3 times, and mounted on glass slides in Fluoromount

G (SouthernBiotech). Wing discs were imaged on a Leica DMR compound microscope using a Q-

Imaging R6 CCD camera and Ocular or Micro-Manager software.

Lethality test with QUAS lines

QUAS lines are a kind gift from Ainhoa Pérez-Garijo and Hermann Steller [14]. For insertions on

chromosome 2, CyO balancer was floating so CyO+ males were sorted and used. Male QUAS

flies were crossed to virgins of the genotype psc-QF>UAS-Tomato/CyO-GFP. Embryos were

collected overnight. CyO and CyO+ progenies were counted until no new adults eclosed for a 24

h period. X^2 test was used to assess deviations from the 1:1 expected Mendelian ratio. For

insertions on chromosome 3, Tb+ pupae were sorted and scored as GFP+ or GFP- before

eclosion was monitored. X^2 test was used to assess deviations from the 2GFP- (Sco/QF>Tom

and QF>Tom/CyO):1GFP+ (Sco/CyO-GFP) expected Mendelian ratio (See Supplemental Table 2

for genotypes of the crosses).

Statistical Analysis

For sample size justifications, we used a simplified resource equation from [24]; E = total

number of animals – total number of groups, where E value of 10-20 is considered adequate.

When we compare two groups (-IR and +IR, for example, where the former is the control group

and the latter is the treated group), n = 6 per group, or E = 11, would be adequate. All samples

subjected to statistical analysis meet or exceed this criterion. To compute p-values, 2-tailed

Student t-tests were used.

Acknowledgments

We thank Alison Bardin and Herman Steller for GAL80 and QUAS stocks, respectively. Additional

stocks from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) and Vienna

Drosophila Resource Center were used in this study.

Funding

This work was supported by the National Institutes of Health (grant R35 GM130374 to TTS).

Figure and Table Legends

Fig 1. Fluorescent reporters for loss-of-heterozygosity

A. Loss of QS results in de-repression of psc-QF-driven Tom expression.

B. Loss of GAL80 results in de-repression of act-GAL4-driven GFP expression.

C-C'. An example of a wing disc with a Tom clone. This disc also expresses en-GAL4>GFP

that was used to express RNAi and transgenes in the posterior compartment of the disc. The

genotype was en-GAL4>UAS-GFP, psc-QF>Tom, QS^{5A}/+ from the cross of en-GAL4>UAS-GFP,

psc-QF>Tom, QS^{5A}/CyO-GFP with w^{1118}. Larvae were treated as in panel E but without the

temperature shift.

D. An example of a wing disc with GFP clones. The genotype was GAL80^{5B}/+; Act-GAL4,

UAS-GFP/+ from the cross of GAL80^{5B}/GAL80^{5B} with Act-GAL4, UAS-GFP/CyO RFP Tb. Larvae

were treated as in panel E but without the temperature shift.

E. The experimental protocol used to express RNAi and transgenes conditionally in the

focused screen. For experiments that do not involve transgene expression, the same protocol

was followed but larvae were maintained at 25°C without a temperature shift.

F. Chromosomal location of QS^{5A}, QS^{9B}, *mwh* and GAL80 studied here (not to scale).

Scale bar = 120 microns in C-D.

Fig 2. Distance from the telomere makes little difference in LOH incidence

Larvae were treated as in Fig 1E but without the temperature shift and wing discs were fixed

and stained for DNA. The graphs show data are from at least two biological replicates per

sample, with the total number of wing discs examined for each sample is shown in parentheses.

Error bars represent mean±1 SEM.

(A-B) IR-induced LOH incidence for QS^{9B} and QS^{5A} are not significantly different but two

recombinant QS^{5A} lines show reduced LOH compared to parental QS^{5A}. Larvae were treated as

in Fig 1E but without the temperature shift. p-values were calculated using a 2-tailed t-test.

(C-D) Knocking down of p53 did not affect QS LOH. Larvae were treated as in Fig 1E. p-

values were calculated using a paired 2-tailed t-test.

The genotypes are as shown below. Stock information is in Supplemental Table 1.

A-B: 9B = psc-QF>Tom/+; QS^{9B}/+ that results from psc-QF>Tom; QS^{9B} X to w^{1118}

5A = psc-QF>Tom/QS^{5A} that results from psc-QF>Tom/CyO-GFP X QS^{5A}/CyO-GFP

5A #6 or #11 = en-GAL4>UAS-GFP, psc-QF>Tom, QS^{5A}/+ from the cross of en-GAL4>UAS-

GFP, psc-QF>Tom, QS^{5A}/CyO-GFP with w^{1118}

C-D: 9B p53 RNAi = UAS-Dcr/+; en>GFP, UAS-p53^{RNAi}/psc-QF>Tom; QS^{9B}, GAL80^{ts}/+

9B p53 H159N or R115H = en>GFP, psc-QF>Tom/+; QS^{9B}, GAL80^{ts}/UAS-p53$^{H159N \text{ or } R115H}$

5A p53 RNAi = UAS-Dcr/+; en>GFP, UAS-p53^{RNAi}/psc-QF>Tom, en>GFP, QS^{5A}; GAL80^{ts}/+

Fig 3. IR-induced LOH shows sex-dependence and locus-independence

Larvae were treated as in Fig 1E but without the temperature shift and wing discs were fixed

and stained for DNA. The genotype of the larvae was GAL80/+; Act-GAL4, UAS-GFP/+ from the

cross of GAL80/GAL80 with Act-GAL4, UAS-GFP/CyO RFP Tb. The graphs show data from at least

two biological replicates per sample, with the total number of wing discs examined per sample

shown in parentheses. The mean±1 SEM is shown for each sample. p-values were calculated

using a 2-tailed t-test.

(A-D) Quantification of GFP+ clone number (A-B) and clone area (C-D) with and without

irradiation for four GAL80 insertions on the X chromosome. Male and female larvae were

separated before dissection and processed separately.

(E-F) Quantification of GFP+ clone number (E) and clone area (F) with and without

irradiation for 12C6 GAL80 insertion on the X chromosome. Male and female larvae were

separated before dissection and processed separately.

(G-J) Representative examples of wing discs for the 12C insertion. Scale bar = 120

microns.

Fig 4. A focused screen identified E2F1 as a regulator of LOH

Wing discs from 3rd instar larvae were dissected 3 days after irradiation, fixed and stained for

DNA. Larvae were treated as in Fig 1E. The graphs show data from at least two biological

replicates per sample, with the total number of wing discs examined per sample shown in

parentheses. The mean±1 SEM is shown for each sample. p-values were calculated using a

paired 2-tailed t-test.

(A-B) Results of the focused screen showing clone number (A) and clone area (B). RNAi

against w is shown as the control for RNAi constructs and w^{1118} is shown as the control for

overexpression constructs.

(C-D') Representative examples of wing discs expressing E2F1^{RNAi} or RAS^{V12} in the P

compartment (GFP+).

Fig 5. Cell autonomous requirements for LOH

(A-D) Larvae were treated as in Fig 1E but without the temperature shift and wing discs

were fixed and stained to visualize DNA. The discs shown are from irradiated larvae.

(E-F) Quantification of Tom+ clone number (E) and clone area (F) in irradiated discs. The

graphs show data from at least two biological replicates per sample, with the total number of

wing discs examined per sample shown in parentheses. The mean±1 SEM is shown for each

sample. p-values relative to w^{1118} controls were calculated using a 2-tailed t-test.

(G-I) Larvae were cultured and irradiated as in (A-F) but wing discs were dissected 4h

after IR, fixed and stained for DNA and with an antibody against cleaved active Dcp1 as a

surrogate marker for apoptosis.

Scale bars = 120 microns.

The genotypes are as shown below. Stock information is in Supplemental Table 1.

(A,G) Control = psc-QF>Tom/+; QS^{9B}/+ that results from psc-QF>Tom; QS^{9B} X to w^{1118}

(H-I) psc-QF>Tom/QUAS-transgene; QS^{9B}/+

(B-D) psc-QF>Tom/+; QS^{9B}/QUAS-transgene

References

1.   Brown J, Bush I, Bozon J, Su TT. Cells with loss-of-heterozygosity after exposure to ionizing radiation in Drosophila are culled by p53-dependent and p53-independent mechanisms. PLoS Genet. 2020;16(10):e1009056. Epub 2020/10/20. doi: 10.1371/journal.pgen.1009056. PubMed PMID: 33075096; PubMed Central PMCID: PMCPMC7595702.

2.   Riabinina O, Luginbuhl D, Marr E, Liu S, Wu MN, Luo L, et al. Improved and expanded Q-system reagents for genetic manipulations. Nat Methods. 2015;12(3):219-22, 5 p following 22. Epub 2015/01/13. doi: 10.1038/nmeth.3250. PubMed PMID: 25581800; PubMed Central PMCID: PMCPMC4344399.

3.   Potter CJ, Tasic B, Russler EV, Liang L, Luo L. The Q system: a repressible binary system for transgene expression, lineage tracing, and mosaic analysis. Cell. 2010;141(3):536-48. Epub 2010/05/04. doi: 10.1016/j.cell.2010.02.025. PubMed PMID: 20434990; PubMed Central PMCID: PMCPMC2883883.

4.   Sogame N, Kim M, Abrams JM. Drosophila p53 preserves genomic stability by regulating cell death. Proc Natl Acad Sci U S A. 2003;100(8):4696-701. Epub 2003/04/04. doi: 10.1073/pnas.0736384100. PubMed PMID: 12672954; PubMed Central PMCID: PMCPMC153618.

5.   Baker BS, Carpenter AT, Ripoll P. The Utilization during Mitotic Cell Division of Loci Controlling Meiotic Recombination and Disjunction in DROSOPHILA MELANOGASTER. Genetics. 1978;90(3):531-78. Epub 1978/11/01. PubMed PMID: 17248870; PubMed Central PMCID: PMCPMC1213905.

6.   Brodsky MH, Weinert BT, Tsang G, Rong YS, McGinnis NM, Golic KG, et al. Drosophila melanogaster MNK/Chk2 and p53 regulate multiple DNA repair and apoptotic pathways following DNA damage. Mol Cell Biol. 2004;24(3):1219-31. Epub 2004/01/20. doi: 10.1128/mcb.24.3.1219-1231.2004. PubMed PMID: 14729967; PubMed Central PMCID: PMCPMC321428.

7.   Wells BS, Johnston LA. Maintenance of imaginal disc plasticity and regenerative potential in Drosophila by p53. Dev Biol. 2012;361(2):263-76. Epub 2011/11/01. doi:

10.1016/j.ydbio.2011.10.012. PubMed PMID: 22036477; PubMed Central PMCID:

PMCPMC3296280.

8.   Ollmann M, Young LM, Di Como CJ, Karim F, Belvin M, Robertson S, et al. Drosophila p53

is a structural and functional homolog of the tumor suppressor p53. Cell. 2000;101(1):91-101.

Epub 2000/04/25. doi: 10.1016/S0092-8674(00)80626-1. PubMed PMID: 10778859.

9.   Siudeja K, Nassari S, Gervais L, Skorski P, Lameiras S, Stolfa D, et al. Frequent Somatic

Mutation in Adult Intestinal Stem Cells Drives Neoplasia and Genetic Mosaicism during Aging.

Cell Stem Cell. 2015;17(6):663-74. Epub 2015/11/27. doi: 10.1016/j.stem.2015.09.016. PubMed

PMID: 26607382; PubMed Central PMCID: PMCPMC5138153.

10.   Goupil A, Heinen JP, Salame R, Rossi F, Reina J, Pennetier C, et al. Illuminati: a form of

gene expression plasticity in Drosophila neural stem cells. Development. 2022;149(22). Epub

2022/11/19. doi: 10.1242/dev.200808. PubMed PMID: 36399062; PubMed Central PMCID:

PMCPMC9845751.

11.   McNamee LM, Brodsky MH. p53-independent apoptosis limits DNA damage-induced

aneuploidy. Genetics. 2009;182(2):423-35. Epub 2009/04/15. doi:

10.1534/genetics.109.102327. PubMed PMID: 19364807; PubMed Central PMCID:

PMCPMC2691752.

12.   Reis T, Edgar BA. Negative regulation of dE2F1 by cyclin-dependent kinases controls cell

cycle timing. Cell. 2004;117(2):253-64. Epub 2004/04/16. doi: 10.1016/s0092-8674(04)00247-8.

PubMed PMID: 15084262.

13.   Wichmann A, Jaklevic B, Su TT. Ionizing radiation induces caspase-dependent but Chk2-

and p53-independent cell death in Drosophila melanogaster. Proc Natl Acad Sci U S A.

2006;103(26):9952-7. doi: 10.1073/pnas.0510528103. PubMed PMID: 16785441; PubMed

Central PMCID: PMCPMC1502560.

14.   Perez-Garijo A, Fuchs Y, Steller H. Apoptotic cells can induce non-autonomous apoptosis

through the TNF pathway. Elife. 2013;2:e01004. Epub 2013/09/26. doi: 10.7554/eLife.01004.

PubMed PMID: 24066226; PubMed Central PMCID: PMCPMC3779319.

15.   zouabi LA, Stefanutti M, Riddiford N, Rubanova N, Bohec M, Servant N, et al. Molecular

Underpinnings and Environmental Drivers of Spontaneous Loss of Heterozygosity in

Drosophila Intestinal Stem Cells. bioRxiv. 2022:2022.07.21.500951. doi:

10.1101/2022.07.21.500951.

16.   Kurzhals RL, Fanti L, Ebsen ACG, Rong YS, Pimpinelli S, Golic KG. Chromosome Healing Is

Promoted by the Telomere Cap Component Hiphop in Drosophila. Genetics. 2017;207(3):949-

59. Epub 2017/09/25. doi: 10.1534/genetics.117.300317. PubMed PMID: 28942425; PubMed

Central PMCID: PMCPMC5676247.

17.   Chick WS, Mentzer SE, Carpenter DA, Rinchik EM, Johnson D, You Y. X-ray-induced

deletion complexes in embryonic stem cells on mouse chromosome 15. Mamm Genome.

2005;16(9):661-71. Epub 2005/10/26. doi: 10.1007/s00335-005-0011-5. PubMed PMID:

16245023.

18.   Eeken JC, de Jong AW, Loos M, Vreeken C, Romeyn R, Pastink A, et al. The nature of X-

ray-induced mutations in mature sperm and spermatogonial cells of Drosophila melanogaster.

Mutat Res. 1994;307(1):201-12. Epub 1994/05/01. doi: 10.1016/0027-5107(94)90293-3.

PubMed PMID: 7513798.

19.   Pastink A, Vreeken C, Schalet AP, Eeken JC. DNA sequence analysis of X-ray-induced

deletions at the white locus of Drosophila melanogaster. Mutat Res. 1988;207(1):23-8. Epub

1988/01/01. doi: 10.1016/0165-7992(88)90006-1. PubMed PMID: 3122035.

20.   Pastink A, Schalet AP, Vreeken C, Paradi E, Eeken JC. The nature of radiation-induced

mutations at the white locus of Drosophila melanogaster. Mutat Res. 1987;177(1):101-15. Epub

1987/03/01. doi: 10.1016/0027-5107(87)90026-1. PubMed PMID: 2881201.

21.   Martin-Blanco E, Gampel A, Ring J, Virdee K, Kirov N, Tolkovsky AM, et al. puckered

encodes a phosphatase that mediates a feedback loop regulating JNK activity during dorsal

closure in Drosophila. Genes Dev. 1998;12(4):557-70. Epub 1998/03/21. doi:

10.1101/gad.12.4.557. PubMed PMID: 9472024; PubMed Central PMCID: PMCPMC316530.

22.   Lawrence PA, Sanson B, Vincent JP. Compartments, wingless and engrailed: patterning

the ventral epidermis of Drosophila embryos. Development. 1996;122(12):4095-103. Epub

1996/12/01. doi: 10.1242/dev.122.12.4095. PubMed PMID: 9012529.

23.   Kerkis J. The Growth of the Gonads in DROSOPHILA MELANOGASTER. Genetics.

1931;16(3):212-24. Epub 1931/05/01. doi: 10.1093/genetics/16.3.212. PubMed PMID:

17246617; PubMed Central PMCID: PMCPMC1201094.

24.   Charan J, Kantharia ND. How to calculate sample size in animal studies? J Pharmacol

Pharmacother. 2013;4(4):303-6. doi: 10.4103/0976-500X.119726. PubMed PMID: 24250214;

PubMed Central PMCID: PMCPMC3826013.

Figure 1

Figure 2

Figure 3

Figure 4

A clone number_hits

B clone area_hits

C DNA Tom GFP

E2F1 RNAi

D DNA Tom GFP **D'** DNA Tom

RAS^{V12}

Figure 5

Supplemental Figure 1

Supplemental Figure 1. Spontaneous GFP is less bright than IR-induced GFP.
Larvae were treated as in Fig 1E but without the temperature shift and wing discs were fixed and stained to visualize DNA. The genotype of the larvae was GAL80^{5B}/+; Act-GAL4, UAS-GFP/+ from the cross of GAL80^{5B}/GAL80^{5B} with Act-GAL4, UAS-GFP/CyO RFP Tb.

(A-A''') A disc from an unirradiated larva showing one faint GFP spot (arrow).

(B-B') A disc from an irradiated larva showing bright GFP spots. A' and B' are shown with similar brightness and contrast range in ImageJ so that signals would be directly comparable. A'' is shown with a narrower range to allow the viewer to see the faint GFP spot.

Supplemental Figure 2

Supplemental Figure 2. A focused screen for modulators of IR-induced QS^{9B} LOH.
Wing discs from 3rd instar larvae were dissected 3 days after irradiation, fixed, and stained for DNA. Larvae were treated as in Fig 1E. The graphs show data from at least two biological replicates per sample, with the total number of wing discs examined per sample shown in parentheses. The mean±1 SEM is shown for each sample. Because A and P compartments are of different sizes, clone numbers and clone areas have been normalized to compartment area. p-values between A and P samples for each genotype (none <0.05) were calculated using a paired 2-tailed t-test. TRiP III (background stock for most RNAi lines used) and RNAi against w served as controls for RNAi constructs and w^{1118} served as the control for overexpression constructs.

 (A-B) LOH clone numbers for RNAi (A) and overexpression (OE,B) constructs.
 (C-D) LOH clone areas for RNAi (C) and overexpression (OE,D) constructs.

Supplemental Table 1

Supplemental Table 1. Drosophila stocks used in this work					
I. General stocks (BL = Bloomington Stock Center)					
stock #/source	used in Fig.	used for		genotype	
BL30042	Fig 2, 4, 5, S1	monitor LOH of QS{9B}		y[1] w[1118]; P{w[+mC]=M2ET-QF}ET40, P{w[+mC]=QUAS-mtdTomato-3xHA}CP; P{ry[+t7.2]=neoFRT}82B P{w[+mC]=tub-QS.P}9B	
BL30133	Fig 1, 2	monitor LOH of QS{5A}		y[1] w[1118]; P{w[+mW.hs]=FRT(w[hs])}G13 P{w[+mC]=tub-QS.P}5A	
BL30043	Fig 2, 4, 5, S1, Table S2	expresses Tom throughout the wing disc		y[1] w[1118]; P{w[+mC]=QUAS-mtdTomato-3xHA}14 P{w[+mC]=M2ET-QF}ET40/CyO	
BL24645	Fig 2	enhance p53 RNAi		P{w[+mC]=UAS-Dcr-2.D}1, w[1118]; Df(3L)Ly, sens[Ly-1]/TM3, Sb[1]	
unknown	Fig 1, 2, 4, S1	P compartment expression		engrailed-GAL4 on chromosome II	
BL5431	Fig 1, 2, 4, S1	mark the P compartment		UAS-eGFP on chromosome II	
Bardin lab	Fig 3	detecting loss of GAL80		Actin-GAL4, UAS-GFP	
II. Stocks used in the focused screen and for functional testing (VDRC = Vienna Drosophila Resource Center)					
gene	function	construct	manipulation	stock #	ref./source
p35	anti-apoptotic	UAS-p35	overexpression	BL5073	Bloomington Stock Center
Diap1	anti-apoptotic	QUAS-DIAP1	overexpression	N/A	Steller lab
p53	pro-apoptotic	UAS-dsRNA	RNAi	V38235	VDRC
p53	pro-apoptotic	UAS-p53R115H	dominant negative	BL8419	Bloomington Stock Center, ref.9
p53	pro-apoptotic	UAS-p53H159N	dominant negative	BL8421	Bloomington Stock Center, ref.9
rpr	pro-apoptotic	QUAS-rpr	overexpression	N/A	Steller lab, ref.14
E2F1	cell cycle, apoptosis	UAS-dsRNA	RNAi	BL27564	Bloomington Stock Center
rux	inhibits mitotic Cdk	UAS-Rux	overexpression	BL9166	Bloomington Stock Center
dap	inhibits G1/S Cdk	UAS-Dap	overexpression	BL83334	Bloomington Stock Center
tefu/ATM	DNA damage response	UAS-dsRNA	RNAi	V22502	VDRC
mei-41/ATR	DNA damage response	UAS-dsRNA	RNAi	v11251	VDRC
grp/Chk1	DNA damage response	UAS-dsRNA	RNAi (GL00067)	N/A	Perrimon lab
grp/Chk1#2	DNA damage response	UAS-dsRNA	RNAi (HMS01573)	BL36685	Bloomington Stock Center
loki/Chk2	DNA damage response	UAS-dsRNA	RNAi	BL35152	Bloomington Stock Center
hiphop	chromosome end protection	P{EPgy2} insertion	overexpression (with GAL4)	BL17618	Bloomington Stock Center
Puc	represses JNK signaling	UAS-puc	overexpression	N/A	ref.21
Puc	represses JNK signaling	QUAS-puc	overexpression	N/A	Steller lab
Bsk/JNK	stress response	QUAS-JNK DN	dominant negative	N/A	Steller lab
hep/JNKK	Stress response	UAS-hep ACT	constitutively active	BL9306	Bloomington Stock Center
Akt	Stress response	UAS-dsRNA	RNAi	BL31701	Bloomington Stock Center
myc	Growth	UAS-dsRNA	RNAi	v2947	VDRC
Ras	Growth, apoptosis	UAS-RAS V12 mutant	constitutively active	BL64195	Bloomington Stock Center
wg/Wnt1	Developmental signaling	UAS	overexpression	N/A	ref.22
Axin	inhibits Wg signaling	UAS-Axin	overexpression	BL7225	Bloomington Stock Center
duox	Cell competition	UAS-dsRNA	RNAi	BL33975	Bloomington Stock Center
duox	Cell competition	UAS-shRNA	shRNA	v330085	VDRC
Myo31DF	Cell competition	UAS-dsRNA	RNAi	BL33971	Bloomington Stock Center
Xrp1	Cell competition	UAS-dsRNA	RNAi	v107860	VDRC
glut1	p53 target	UAS-dsRNA	RNAi	v13326	VDRC
glut3	p53 target	UAS-dsRNA	RNAi	v13302	VDRC
w	control	UAS-dsRNA	RNAi	BL33623	Bloomington Stock Center
TRiP III	control stock for RNAi lines	N/A	none	BL36303	Bloomington Stock Center
w1118	control	none	none	BL3605	Bloomington Stock Center

Supplemental Table 2

Supplemental Table 2. X2 test for survival after ubiquitous expression of QUAS transgenes.													
Psc-QF, mtdTomato/CyO-GFP virgins crossed to	tested stock (stock#)	inserted Chromosome	Cy	Cy+	GFP-	GFP+	total	expected Cy or GFP-	expected Cy+ or GFP-	(o-e)^2	(o-e)^2	X2	p
N/A	**w1118 control**		**38**	**50**			**88**	**44**	**44**	**36.0**	**36.0**	**1.6**	**n.s.**
QUAS-X/CyO; Sb/TM6 Tb	QUAS-rpr (11M)	2	88	0	N/A	N/A	88	44	44	1936.0	1936.0	88.0	<0.001
QUAS-X/CyO; Sb/TM6 Tb	QUAS-rpr (12M)	2	58	0	N/A	N/A	58	29	29	841.0	841.0	58.0	<0.001
Sco/CyO; QUAS-X/TM6 Tb	**QUAS-rpr (2M)**	**3**	**N/A**	**N/A**	**0**	**47**	**47**	**31**	**16**	**981.8**	**981.8**	**94.0**	**<0.001**
QUAS-X/CyO; Sb/TM6 Tb	QUAS-DIAP1 (3M)	2	34	14	N/A	N/A	48	24	24	100.0	100.0	8.3	<0.005
QUAS-X/CyO; Sb/TM6 Tb	QUAS-DIAP1 (8M)	2	64	24	N/A	N/A	88	44	44	400.0	400.0	18.2	<0.001
QUAS-X/CyO; Sb/TM6 Tb	**QUAS-DIAP1 (12M)**	**2**	**49**	**61**	**N/A**	**N/A**	**110**	**55**	**55**	**36.0**	**36.0**	**1.3**	**n.s.**
QUAS-X/CyO; Sb/TM6 Tb	QUAS-bskDN (19)	2	38	0	N/A	N/A	38	19	19	361.0	361.0	38.0	<0.001
Sco/CyO; QUAS-X/TM6 Tb	**QUAS-bskDN (15A)***	**3**	**N/A**	**N/A**	**0**	**25**	**25**	**17**	**8**	**277.8**	**277.8**	**50.0**	**<0.001**
Sco/CyO; QUAS-X/TM6 Tb	QUAS-bskDN (30)	3	N/A	N/A	4	19	23	15	8	128.4	128.4	25.1	<0.001
Sco/CyO; QUAS-X/TM6 Tb	QUAS-bskDN (43)	3	N/A	N/A	2	13	15	10	5	64.0	64.0	19.2	<0.001
QUAS-X/CyO; Sb/TM6 Tb	QUAS-puc (21)	2	30	0	N/A	N/A	30	15	15	225.0	225.0	30.0	<0.001
QUAS-X/CyO; Sb/TM6 Tb	QUAS-puc (66)	2	72	10	N/A	N/A	82	41	41	961.0	961.0	46.9	<0.001
Sco/CyO; QUAS-X/TM6 Tb	QUAS-puc (59)	3	N/A	N/A	0	12	12	8	4	64.0	64.0	24.0	<0.001
Sco/CyO; QUAS-X/TM6 Tb	**QUAS-puc (60)**	**3**	**N/A**	**N/A**	**0**	**16**	**16**	**11**	**5**	**113.8**	**113.8**	**32.0**	**<0.001**
*counted as pupa													