Title: Increasing environmental fluctuations can dampen variability of endogenously cycling populations

Authors:

1. Nicholas Kortessis.

Department of Biology, Wake Forest University, Winston Salem, NC 27109 and Department of Biology, University of Florida, Gainesville, FL 32611. email: kortessn@wfu.edu;

2. José Miguel Ponciano.

Department of Biology, University of Florida, Gainesville, FL 32611. email: josemi@ufl.edu;

3. Franz W. Simon

Department of Biology, University of Kentucky, Lexington, KY 40506. email: fs365@cornell.edu;

4. Jake M. Ferguson

Department of Biology, University of Hawaii at Manoa, Honolulu, HI 96822. email: jake.ferguson@uky.edu;

Corresponding Author Mailing Address: Jake Ferguson, 101 T.H. Morgan Building, Lexington KY 40506, email: jake.ferguson@uky.edu

Running Title: Environmental Variability Dampens Cycles

Keywords: Long transient | Nonlinear dynamics | Extinction risk | Stability | Stochasticity

Article Type: Major Article

Word Count: Abstract: 200; Main Text: 4431

Manuscript elements: References 66; Figures: 6; Tables: 0

Author Contributions: JMF and JMP conceived of the original idea and secured funding. NK constructed and analyzed models with input and discussion from all authors. FWS analyzed models. NK wrote the original draft, and all authors contributed to revisions.

Abstract

Understanding how populations respond to increasingly variable conditions is a major objective for natural resource managers attempting to forecast extinction risk. The lesson from current modeling is clear: Increasing environmental variability increases population abundance variability. We show that this paradigm fails to describe a broad class of empirically observed dynamics, namely endogenously-driven population cycles. In contrast to the dominant paradigm, these populations can exhibit reduced long-run population variance under increasing environmental variability. We hypothesize that this paradox arises from interactions between environmental stochasticity and nonlinear density dependence. Such interactions violate the oft-assumed additivity of stochastic and deterministic drivers of population fluctuations present in many models that forecast population size. We show evidence for the interaction in two canonical cyclical populations: flour beetles and Canadian Lynx. To help identify the interaction, we develop new theory to quantify the strength of these interactions by partitioning the effects of nonlinear dynamics and stochastic variation on dynamical systems. In both empirical examples, the partitioning shows that the interaction between deterministic and stochastic dynamics reduces the overall variance in population size. Our results highlight that previous predictions about extinction under environmental variability may prove inadequate to understand the effects of climate change in many populations.

Introduction

Most projections of future biodiversity under climate change present a bleak outlook [1, 2, 3, 4, 5]. Models predict that the warmer average temperatures predicted by climate models will be outside the current thermal limits of some species [6, 7]. While changes in average conditions are likely to have significant consequences for many species, more frequent extreme events are also predicted to occur with climatic change. Examples include the projected increases in variability of annual temperature [8] and rainfall [9]. In a landmark study, Vasseur et al. [10] used thermal performance curves to show that future changes in annual temperate variation may impact ectotherms' thermal performance as much or more than predicted changes in average temperature. While suggestive, theoreticians and practitioners alike are still far from understanding the many ways that increasingly variable environmental conditions impact the ability of populations to persist.

Theory points to two ways in which environmental variability affects population fluctuations. The first mechanism is through changes in a population's long-term average growth rate through nonlinear averaging. Thus, assuming fixed growth under average conditions, increasing environmental variability lowers long-term population growth [11], and the same nonlinear averaging effects apply at the level of demographic rates (see Real and Ellner [12] for a graphical application of this idea). This effect has been demonstrated empirically in the growth rate of experimental populations of green algae (*Tetraselmis tetrahele*) [13], in herbivore feeding rates [14], and in predator feeding rates [15] (although, in some cases, environmental variability can elevate average population growth, especially when the environment has some predictability [16] or populations are spatially distributed [17, 18]).

The second-way environmental variability puts species at greater risk of extinction is by causing

short-term population fluctuations. To a first approximation, greater fluctuations in population size enhance extinction risk because populations dip to lower numbers more frequently. At low numbers, populations are at heightened risk of extinction from the increased chance of below-average survival and reproduction among the remaining individuals (i.e., demographic stochasticity; [19, 20]). Taken together, current theory predicts that increases in environmental fluctuations in single-species population models will always drive increases in the long-run population variance [21, 22, 23].

Stochastic population models can predict extinction in both stable [24] and declining populations [25], but do so by assuming relatively simple forms of density-dependence. By simple forms of density-dependence, we mean that which leads to a single stable equilibrium, as in the logistic model with a carrying capacity. Under these models, populations may approach and persist at this equilibrium indefinitely or, below a critical reproductive rate, decline to extinction. This behavior is found in many commonly used models in ecology beyond the logistic model, including the Gompertz, Beverton-Holt, and theta-Ricker models [26, 27]. Ecologists have made progress integrating the effect of environmental variability into our understanding of population and community dynamics that exhibit a single equilibrium [28]. However, these models are ill-suited to describe populations that exhibit more complex dynamics.

In many biological systems, population biologists have recognized the need for models that include more complex forms of density-dependence that generate, for example, multiple equilibria. For example, systems with tipping points between alternative stable states are one case where models of simple density-dependence are insufficient. These systems can exhibit a much richer portfolio of responses to environmental variability than simple density-dependence. For example, in models of alternative stable states, environmental variability can push such a system between

these alternative stable states [29], a phenomenon termed stochastic flickering [30, 31]. In addition, stochastic resonance, a sudden increase in system variation, can arise when increases in the extrinsic variance push the system to periodically cycle between alternative stable states when it would otherwise stay in one of the states [32]. This resonance has been hypothesized to play a role in seasonal influenza outbreaks [33] and in the apparent periodic behavior of ice ages [34]. Flickering and resonance predict sudden jumps in the system variance in response to small increases in the environmental variance.

Here, we explore how the variability in the abundance of cycling populations responds to changes in environmental variability. Population cycles are common in nature and serve as a valuable lens for studying dynamical systems with complex feedbacks. We start by analyzing a stochastic, time-dependent Ricker model, which is sufficiently flexible to model both simple density-dependence and cycling populations in a single framework under two cycling mechanisms (following [35]): exogenous cycles that are caused by external factors (e.g., weather [36]) and endogenous cycles that are caused by delayed density feedback loops that can be thought of as self-regulating mechanisms (e.g., stage structure; [37, 38]) or interpopulation mechanisms (e.g., predator effects; [39] or pathogen effects; [40]). Differences between simple density-dependence and density-dependence in cycling populations are visualized in Fig. 1a-c. The most crucial feature is how the qualitative features of density-dependence (illustrated by the contours of the surface) change with density. We use simulations (illustrated in Fig. 1d-f) to qualitatively describe how these models respond to increases in environmental variability and quantify how variation in the environment interacts with the density-dependent dynamics using a novel variance partitioning analysis. We apply this partitioning framework to empirical examples of delayed density dependence in experimental populations of flour beetles (*Tribolium castaneum*) and in predator-prey cycles with a Canadian lynx (*Lynx canadensis*)

 - snowshoe hare dataset (*Lepus americanus*).

The impact of variable environments on cycling populations

We start by investigating the properties of the Ricker model, originally developed to qualitatively account for density dependence in the young members of a population [41], for example due to predation. The model has since been derived under additional scenarios [42, 43, 44], and its general applicability to a wide array of empirical conditions has led the Ricker model to be used by many fisheries and wildlife agencies as a management tool. Aiding the adoption of this modeling framework in applied ecology is the ease with which estimation and forecasting can be done, even when observations are contaminated with measurement error [45]. The choice to use the Ricker here, in addition to its wide application in fisheries and wildlife management, is the fact that multiple dynamical regimes can be explored in a single, simple model where environmental variability can be easily incorporated. This allows for direct comparisons of the effect of environmental variability in different dynamical regimes without having to make comparisons between different modeling frameworks.

If n_t is the density of individuals in the population at time t, the Ricker model defines the density of individuals in the following time step, n_{t+1}, as $n_{t+1} = n_t e^{a_t - b n_t}$. Here, a_t is the (possibly time-dependent) density-independent per-capita growth rate while b is a coefficient scaling the effect of density. A typical interpretation is that the species under consideration has discrete generations, and the time t is units of generations such that each individual produces e^{a_t} offspring in year t and $e^{-b n_t}$ is the fraction that survive.

The Ricker model can exhibit either a single stable equilibrium, exogenous cycles, or endogenous cycles depending on the time-dependent parameter a_t. A stable equilibrium occurs when a_t is a

Figure 1: Conceptualization of environmental variability on population dynamics in different models. Under simple models of density-dependence (a), population fluctuations reflect a ball (blue lines) bouncing in a bowl (heavy black lines), where the depth of the bowl (z-axis in all plots) reflects the action of density-dependence. Environmental variation drives random perturbations of the ball away from the bottom of the bowl, which is the carrying capacity in this framework. A conception of exogenous cycles (b) is that the position of the bottom of the bowl moves with the exogenous driver, but the shape of the bowl is unchanging. A conception of endogenous cycles (c) is that the shape and action of density-dependence (black lines) changes over time, meaning that when the population is at low density, the equilibrium moves to higher density, but at high density, the equilibrium moves again to lower density. The topography of the landscape of population change – the bowl itself – is thus changing with the state of the population.

constant less than 2 for all t. Exogenous cycles occur when a_t is a time-dependent cyclic function (for the simulations below, we chose a sine function), and endogenous cycles occur when a_t is a constant slightly larger than 2, which models overcompensatory density-dependence [e.g., 42, 46]. Examples of the three dynamical regimes are given in Fig. 1d-f. The parameter b scales population density without affecting dynamical regimes and so can be re-scaled out of the model; see SI S2.2 for details. The stable-Ricker and similar models have been used to to understand the effects of environmental variability on populations [e.g., 47, 48].

To include environmental variation into the Ricker model we assume that the per-capita growth rate includes random variation over time:

$$N_{t+1} = N_t e^{a_t - bN_t + \sigma Z_t}, \tag{1}$$

where Z_t is a random variable representing variable environmental conditions in year t, and σ is a measure of the magnitude of environmental variation over time. Following standard modeling assumptions, Z_t is an i.i.d. standard normal random variable (i.e., $Z_t \sim \mathcal{N}(0,1)$ for all t, and $E[Z_t Z_s] = 0$ for all $t \neq s$). The effect of the environment is given by the term σZ_t, which has variance σ^2.

To understand the impact of environmental variation on cycling population dynamics we first consider a small variance approximation [49, 11] in the case of exogenous cycles, then show how this approximation breaks down in the case of endogenous cycles. Small variance approximations simplify the shape of density-dependence around a point (typically an equilibrium) and, as such, describe population dynamics with a linear forms of density-dependence near this point.

 Exogenous cycles

129 The approximation for variance in population size in the stochastic Ricker model with an exogenous

130 driver is

131
$$\text{Var}(N) \approx \underbrace{\frac{\bar{N}^2}{\bar{a}(2-\bar{a})}\sigma_a^2(1+c)}_{\substack{\text{Variance from} \\ \text{exogenous driver}}} + \underbrace{\frac{\bar{N}^2}{\bar{a}(2-\bar{a})}\sigma^2}_{\substack{\text{Variance from} \\ \text{environmental} \\ \text{stochasticity}}}, \tag{2}$$

where $\bar{N} = \bar{a}/b$ is the average population size (details in S1), $\bar{a}$ is the average low density growth

rate, σ_a^2 is the (approximate) variance when a_t cycles exogenously, σ^2 is the variance in the stochastic

component of the environment, and $c = \lim_{T\to\infty} \sum_{u=1}^{T} 2(1-\bar{a})^u \rho_a(u)$ is the effect of autocorrelated

environmental variation on population fluctuations ($\rho_a(u)$ is the autocorrelation function of a_t with

time lag u; full derivation in supplementary material S3). Equation (2) states that the long-run

variance in population size can be partitioned into a sum of two components: variability from a pe-

riodic exogenous factor (σ_a^2) and from environmental stochasticity (σ^2). Approximation (2) predicts

a linear increase in population variability with increases in either the periodic exogenous factor or

in the random environmental stochasticity. This prediction matches well the observed qualitative

linear increase in $\text{Var}(N)$ with increasing σ^2 for both models of standard density-dependence when

$\sigma_a^2 = 0$ (Fig. 2a) and exogenous cycles when when $\sigma_a^2 > 0$ (Fig. 2b). Moreover, this approximation

does a good job matching the quantitative relationship between $\text{Var}(N)$ and σ_a^2 (Fig. S3) and σ^2

(Fig. S4) in the model with exogneous cycles.

Figure 2: Effect of changing environmental variability for three qualitatively different kinds of dynamics. Each plot shows the average variance of population size over the time series across 1000 replicates (solid black line) as well as the the 95% upper and lower quantiles across replicates. Parameters: In (a), $\bar{a} = 1$, $\sigma_a^2 = 0$, and $b = 1$. In (b), $\bar{a} = 1$, $\sigma_a^2 = 0.25$, and $b = 1$. In (c), $\bar{a} = 2\ln\{3\} \approx 2.2$, $\sigma_a^2 = 0$, and $b = 2\ln\{3\}$. The value of a in (c) is chosen so that $\text{Var}(N)$ in the deterministic cycling models are approximately the same, thereby facilitating comparisons of the two different mechanisms of cycles (details in S2).

Endogenous cycles

The small variance approximation fails in the endogenous cycle case as eq. (2) predicts negative variance when a_t is a constant greater than 2 (and so $\bar{a} > 2$). Simulations of the stochastic endogenously cycling model show that the variance in population size initially declines as the environment becomes more variable (Fig. 2c). The implication here is that, for some levels of environmental variation, increasing environmental variability reduces population fluctuations.

The shape of density-dependence suggests a possible mechanism of this paradoxical decline in population variability with increasing environmental variability. One way to understand density-dependence for the 2-cycle is to consider the second iterate map [50]. If $F(n_t)$ is the deterministic Ricker map (i.e., $n_{t+1} = F(n_t)$), then the second iterate map is $F_2(n_t) = F(F(n_t))$ which projects population size 2 time-steps in the future. For a 2-cycle, the second iterate map has 3 non-trivial fixed points: 2 stable fixed points corresponding to the two asymptotic cycling values (illustrated with the dashed lines in Fig. 3), and an unstable fixed point exactly between the two (Fig. S1). This unstable fixed point is given by the value $\bar{a}/b$ (supplementary material S2), and has weak local stability as measured by the local rate of change in the 2-step per-capita growth rate (i.e., the slope of $\ln\{n_{t+2}/n_t\}$ with respect to n_t at each fixed point). Populations that begin near this weak unstable fixed point can take a long time before exhibiting asymptotic dynamics (Fig. S2), a phenomenon termed a long transient dynamic [51].

The fact that the system can take a long time to escape the unstable fixed point means that the dynamical consequences of stochastic variation depend on both the strength of environmental fluctuations and on the nature of density-dependence. Near stable fixed points, stochasticity has minimal effects due to strong density-dependence. Near the weak unstable fixed point, stochastic

Figure 3: Sample path for the stochastic Ricker model in the case of endogenous cycles showing the origin of stochastic variance induced population fluctuations reduction. Periodically, the population gets "stuck" near the unstable equilibrium separating the two points of the deterministic cycle. In this example, the deterministic model cycles between the two fixed points $n_t = 50$ and $n_t = 150$ (dashed lines). The mean population density (here $\bar{n} = 100$) is an unstable fixed point. In this example $a = 2\ln\{3\}$, $b = a/100$, and $\sigma^2 = 0.02$.

effects are relative strong in comparison with the weak effect of density-dependence. To visualize this effect, consider the sample path of the stochastic endogenously cycling model in Fig. 3. Occasionally, stochastic variation pushes the population towards the unstable fixed point (located at $\bar{n} = 100$ in Fig. 3). As a consequence, the population gets temporarily "stuck" and spends extended time in a transient state where fluctuations in population size are mostly stochastic, until

the population eventually exits the weak transient phase. Exit of the transient can occur either

because of the slow deterministic dynamics back to the cycle or due to large chance perturbations.

This dynamical phenomenon keeps the population near the long-run average ($\bar{n} = 100$ in Fig. 3),

thereby reducing overall population size variability in comparison to the system without environ-

mental stochasticity. The inset of Fig. 3 shows one particularly acute example of this effect in a

representative sample path of the stochastic Ricker model with endogenous cycles.

Variance decomposition of time series

179 To quantitatively assess the roles of stochasticity, deterministic dynamics, and their potential in-

180 teraction, we partitioned the overall variance in population fluctuations (denoted as $\text{Var}(N_t)$) as

181 follows:

$$\text{Var}(N_t) = \sigma_{\text{Det}}^2 + \sigma_{\text{Stoc}}^2 + \sigma_{\text{Nonlin}}^2 + \sigma_{\text{Int}}. \tag{3}$$

This formula partitions the variance in population size as a sum of four components: (i) the variance

in population size in a purely deterministic model, $\sigma_{\text{Det}}^2 = \text{E}[(n_t - E(n_t))^2]$, where n_t is the predicted

deterministic trajectory in the absence of stochasticity, (ii) the contribution of stochastic variation

on top of the deterministic variation, $\sigma_{\text{Stoc}}^2 = \text{E}[(N_t - n_t)^2]$, where N_t is the observed stochastic

process, the effects of stochasticity on average population density caused by nonlinear averaging

(i.e., mean shifts from Jensen's inequality) $\sigma_{\text{Nonlin}}^2 = -(\text{E}(N_t) - \text{E}(n_t))^2$, and (iv) and covariation

between stochastic and deterministic variation, $\sigma_{\text{Int}} = 2\text{E}(n_t - \text{E}(n_t))(N_t - n_t)$. The interaction

component σ_{Int} can be positive or negative, which differs from all other components, which are

all strictly non-negative. Its sign measures whether the direction of deterministic fluctuations

$(n_t - E(n_t))$ are on average in the same direction as stochastic fluctuations $(N_t - n_t)$. Thus,

$\sigma_{\text{Int}} > 0$ when environmental stochasticity amplifies deterministic fluctuations while $\sigma_{\text{Int}} < 0$ when

environmental stochasticity damps deterministic fluctuations. The derivation of eq. 3 is provided

in supplementary material S4.

We applied eq. (3) to simulations from the Ricker model under simple density dependence,

exogenous cycles, and endogenous cycles. We found that the Ricker behaves as expected under

simple density-dependence and exogenous cycles with $\sigma_{\text{Int}} = 0$. Under endogenous cycling, $\sigma_{\text{Int}} < 0$

(Fig. 4), reflecting damping of deterministic fluctuations by environmental stochasticity. This

demonstrates that the variance partition accurately measures the novel variance damping behavior

shown in the relationship between population variability and environmental variability for cycling

populations (Fig. 2c).

Empirical example of lagged density dependence: flour beetle population

dynamics

We applied the variance partitioning framework to cyclic populations of flour beetles to test for any

evidence of interactions between density-dependence and environmental stochasticity. Experimental

data were reported in [52] and are illustrated in Fig. 5. We fit the stage-based larvae-pupae-adult

(hereafter LPA) stochastic model described by [53], given as

$$L_{t+1} = bA_t \exp\left(-c_{el}L_t - c_{ea}A_t + \varepsilon_{1t}\right)$$
$$P_{t+1} = L_t(1 - \mu_l)\exp(\varepsilon_{2t}) \tag{4}$$
$$A_{t+1} = [P_t \exp\left(-c_{pa}A_t\right) + A_t(1 - \mu_a)]\exp(\varepsilon_{3t}),$$

209 Here, the ε_{it}'s are random values drawn from a multivariate normal distribution where the ϵ_{it}'s are

210 all i.i.d. over time and across life stages. While the LPA model is not the Ricker model, it does

Figure 4: Interaction between environmental variability and deterministic processes as measured by the interaction covariance σ_{Int} (eq. [3]) for three different cases of the Ricker model. In the simple density-dependence model, $a_t = 1$. In the exongenously cycling model, $a_t = 1 + \sin(t/\pi)$. In the endogenous cycle mdoel, $a_t = 2.2$. The interaction term σ_{Int} is rescaled by the total population variance $\text{Var}(N_t)$ to provide a relative measure of the strength of the effect. The lines for simple density-dependence and the exogenous cycle lie exactly on top of each other but are offset slightly in the figure for clarity.

have similar qualitative behavior, exhibiting cycles and chaos in some regions of parameter space

due to lagged density dependence [52].

All parameters were estimated with maximum likelihood using one-step predictions following

the methods described in [53]. We used the fitted model to estimate the variance components for

the total population abundance, $N_t = L_t + P_t + A_t$, in eq. (3). Further details on estimating

the variance components are given in S4.1. Uncertainty in each component was estimated using a

parametric bootstrap with 10,000 draws from the sampling distribution. To combine the estimated

variance components from each of the four experimental replicates, we used the sum of the variance

components of each replicate for each bootstrap draw, which assumes replicates are independent.

Using our partitioning scheme, we estimated that variance reduction from interactions between

environmental variability and density-dependence reduced population fluctuations by 50%. The

sampling distributions of each variance component (reported as mean [95% confidence interval]),

normalized by the overall population variance, are shown in Fig. 5. We estimated population fluctu-

ations arising from complex density-dependence alone $\sigma^2_{\mathrm{Det}}/\mathrm{Var}(N_t) = 0.74$ [0.02, 2.28], population

fluctuations from stochasticity $\sigma^2_{\mathrm{Stoch}}/\mathrm{Var}(N_t) = 1.76$ [1.34, 2.96], moderate effects of nonlinear

averaging $\sigma^2_{\mathrm{Nonlin}}/\mathrm{Var}(N_t) = -0.47$ [-1.87, -0.09], and variance damping sufficient to cancel more

than half of the stochastic population fluctuations $\sigma_{\mathrm{Int}}/\mathrm{Var}(N_t) = -1.02$ [-2.4, -0.1]. The interac-

tion estimate indicates that the covariance between stochastic and deterministic dynamics plays

a sizeable role in damping the overall variance, and that, without this damping effect, the overall

population variance would be about double what is observed.

Figure 5: Data for the flour beetle experiment reported in [52] (left panels). Colored lines give weekly counts of each stage, black line gives the total count ($L + P + A$) used in the variance component analysis. Estimated variance components from eq. (3) (right panel), scaled by the overall population variance.

Empirical example of predator-prey dynamics: lynx-hare interactions

While intraspecific interactions are one mechanism that generates endogenous cycles, trophic interactions, such as predation, can also create population cycles. Here, we used ninety-one years of data on Lynx and snowshoe hare (*Lepus americanus*) pelts from the Hudson Bay Company (data from [54]) to model predator-prey dynamics. We then applied the variance partitioning framework to the fitted model of the Canadian Lynx to estimate the interaction component in eq. (3). We fit a discrete-time version of the Lotka-Volterra predator-prey model where Lynx (L) and hare (H) dynamics follow

$$H(t_{i+1}) = \left(\int_{t_i}^{t_{i+1}} \alpha H(u) - \beta H(u) L(u) \, du \right) e^{\varepsilon_{H,t_i}}$$

$$L(t_{i+1}) = \left(\int_{t_i}^{t_{i+1}} -\gamma L(u) + \delta H(u) L(u) \, du \right) e^{\varepsilon_{L,t_i}},$$

$$(5)$$

where the $\varepsilon_{\cdot,t_i}$'s are random values drawn from normal distributions that are i.i.d over time and

species. Parameters from eq. (5) were estimated using maximum likelihood in the template model

builder R package [55]. We assumed observations of hare and Lynx abundance (Fig. 6A) were

lognormally distributed. We also assumed that the first observations in the time series were drawn

from a lognormal distribution with unknown mean and the same variance as subsequent observa-

tions. Using the fitted model, we estimated the variance components and their uncertainty in the

same manner as above. Further details on the estimation of the variance components and their

uncertainty are given in S4.1.

As with the flour beetle case studied above, we find again that the interaction between density-

dependence and environmental variability has effects on population fluctuations of similar orders

of magnitude to either operating alone. The sampling distributions of each variance component,

normalized by the overall population variance, are shown in Fig. 6. We estimated a dominant

contribution of deterministic dynamics to overall population fluctuations $\sigma_{\text{Det}}^2/\text{Var}(N_t) = 0.92$

[0.56, 1.41], smaller effects of environmental stochasticity $\sigma_{\text{Stoch}}^2/\text{Var}(N_t) = 0.45$ [0.37, 0.66], weak

nonlinear averaging $\sigma_{\text{Nonlin}}^2/\text{Var}(N_t) = -0.06$ [-0.30, -4.66×10^{-5}], and sizeable variance damping

$\sigma_{\text{Int}}/\text{Var}(N_t) = -0.31$ [-0.92, 0.03]. Here, we estimate that population fluctuations would be $\approx 31\%$

more variable without the interaction between complex density-dependence and environmental

fluctuations.

Figure 6: Lynx population data (left panel). This times series was modeled using eq. (5). Estimated variance components from eq. (3) (rescaled by the overall population variance) are illustrated in the right panel.

Discussion

As global change continues, one expectation is that conditions will become more variable over time. With increasing environmental variability, a common worry is that many populations will experience greater abundance fluctuations and a corresponding increase in the probability of extinction. Such a perspective derives from models assuming populations fluctuate around a steady state and that increasing variability in the environment will drive increases in population fluctuations [56, 57, 58, 47]. Here, we considered the response of non-equilibrium dynamics, namely cycles, to increasing environmental variability. We found that increasing environmental variability in an exogenous cycle (i.e., caused by a dynamically unlinked cycling factor) leads to proportional increases in population fluctuations, as expected from prior models assuming simple density dependence (i.e., a point equilibrium). However, endogenous cycles driven by lagged density-dependence respond to environmental variability qualitatively differently; increasing environmental variability can diminish

overall population fluctuations. The mechanism diminishing population fluctuations reflects an interaction between environmental variation and density-dependence that relies on the presence of long-transients in the deterministic system. A long transient is a dynamic that is not represented in the asymptotic behavior of a dynamical system, but lasts long enough to be relevant on the timescales of typical ecological observations [51].

In many domains of ecology and evolution, researchers have recognized the potential for stochastic factors to interact in non-intuitive ways with deterministic processes. Coulson et al. [59] cataloged a number of notable examples of these interactions in ecology and epidemiology and subsequent work has noted additional ways environmental variability can interact in complex ways with deterministic dynamics. Nearly all these studies have emphasized how environmental variability can amplify deterministic fluctuations, or generate new fluctuations (i.e., a positive interaction in eq. [3]). For example, environmental variability has been shown to push a system currently in one stable state to another [29] or to "stochastically flicker" between alternative stable states when it would otherwise remain in one of the stable states [30]. In addition to stochastic flickering, stochastic resonance [32] and stochastic amplification [60] both describe interactions between stochastic variation and deterministic dynamics that increase system variability. Our study shows that these examples fall into a larger class of interactions that also include the paradoxical phenomenon whereby increases in stochastic variation reduce overall system variability.

The variance damping explored here appears to be most similar to a phenomenon noted in the literature as inverse stochastic resonance. As first described in models of neural circuits, the oscillatory firing of neurons collapses to a fixed point with increases in stochastic variation [61]. Touboul et al. [31] also described inverse resonance arising in an ecosystem model of savanna-forest dynamics. Another potential example of this phenomenon may be related to work on whooping

cough, where Rohani et al. [62] found that increasing levels of stochasticity in disease models slowed down annual epidemic cycles. More work needs to be done to determine if the mechanism driving all of these phenomenon is equivalent to the damping process we describe here. In any case, all would be undoubtedly classified as negative interactions in our variance partitioning scheme, and together suggest an interesting avenue for future research and synthesis.

The partitioning scheme we develop has broad applications to any ecological system in which a deterministic trajectory can be described and researchers want to understand the relative contributions of stochastic and deterministic factors to population fluctuations. Recent work by Mutshinda et al. [63] determined the contribution of density-dependent dynamics to the long-run population variance by comparing the predicted long-run variance to a density-independent model under the same stochastic forces. The partitioning scheme we present here (eq. (3)) addresses a similar problem, but differs in that it allows for, and quantifies, interactions between density-dependence and stochastic variation. In addition to the interaction, the partitioning can be interpreted in terms of three well-known terms: (i) the deterministic dynamics that can include population cycles, (ii) the stochastic dynamics that account for the complex sources of variation that appear unstructured, and (iii) the differences between the observed average population density and the average predicted by a deterministic model arising from Jensen's inequality.

Biologists have long relied on variance partitioning schemes to identify components of variation that have conceptual value, even if their underlying mechanistic causes are obscured. For example, analysis of variance, one of the most commonly applied statistical tools in biology, is used to distinguish differences among and between groups by comparing the average within group variation to the between group variation. Similarly, quantitative genetics has successfully partitioned the additive heritable phenotypic variation into genetic and environmental components without

knowing the specific loci responsible for this variation [64]. The study of competition and species coexistence now routinely uses a partitioning scheme that envisions counterfactual scenarios (e.g., a system in which no stabilizing mechanisms are present) to measure the impact of species differences in maintaining or eroding species diversity [65, 66]. Variance partitioning schemes for population models can play a similar role, helping us to categorize and quantify the qualitative components of a dynamical system of very different systems and improve our understanding of the mechanisms that drive fluctuations in population size over time. As a descriptor, this partition is useful for identifying, measuring, and categorizing drivers of population variation in nature and in models. We expect that a wide-scale and data-driven analysis of variance components applied to population time-series should reveal the prevalence and evidence for variance damping in nature.

Understanding the many impacts of global change on species demands an approach that embraces variability and uncertainty. Simple population models with environmental stochasticity are mathematical microscopes (sensu Cohen [67]) of natural systems through which we infer how population processes work under such uncertainty and variability. Recent analyses suggest that the prevalence of nonlinear dynamics may be much more common than previously appreciated [68, 69], so understanding the stochastic behavior of the models with nonlinear dynamics, especially relatively simple ones such as the Ricker model, is an important step toward making sound forecasts about the future of these populations. Our results suggest that our baseline understanding of environmental variability in complex ecological systems may need to be revised. Even elementary modifications to simple models may reveal paradoxical, yet plausible, phenomena such as the variance damping explored here. Our investigation into variance damping in flour beetle and Lynx-Hare dynamics shows that more complex, biologically grounded models may have similar paradoxical behavior. Given the inferential and predictive roles that stochastic population models

play in applied population and conservation biology [70, 71], understanding the basic dynamical properties of commonly used models remains imperative. As such, damping interactions between environmental stochasticity and deterministic dynamics may play a previously unrecognized role in the dynamics of ecological systems.

Acknowledgements

Many thanks to Bob Holt and the Holt lab group for comments on previous versions of this manuscript. Funding was provided by the National Science Foundation grant NSF-DMS-2052372 to JMF and JMP.

Code Statement

Code to reproduce the work found in this manuscript can be found at https://github.com/kortessis/Nonlinear-Dynamics-and-Environmental-Variability.

Competing Interests Statement

The authors declare no competing interests related to this work.

Literature Cited

[1] Céline Bellard, Cleo Bertelsmeier, Paul Leadley, Wilfried Thuiller, and Franck Courchamp. Impacts of climate change on the future of biodiversity. *Ecology Letters*, 15(4):365–377, 2012.

[2] Walter Jetz, David S Wilcove, and Andrew P Dobson. Projected impacts of climate and land-use change on the global diversity of birds. *PLoS Biology*, 5(6):e157, 2007.

[3] Henrique M Pereira, Paul W Leadley, Vânia Proença, Rob Alkemade, Jörn PW Scharlemann, Juan F Fernandez-Manjarrés, Miguel B Araújo, Patricia Balvanera, Reinette Biggs, William WL Cheung, et al. Scenarios for global biodiversity in the 21st century. *Science*, 330 (6010):1496–1501, 2010.

[4] Chris D Thomas, Alison Cameron, Rhys E Green, Michel Bakkenes, Linda J Beaumont, Yvonne C Collingham, Barend FN Erasmus, Marinez Ferreira De Siqueira, Alan Grainger, Lee Hannah, et al. Extinction risk from climate change. *Nature*, 427(6970):145–148, 2004.

[5] Mark C. Urban. Accelerating extinction risk from climate change. *Science*, 348(6234):571–573, 2015.

[6] Raymond B Huey, Curtis A Deutsch, Joshua J Tewksbury, Laurie J Vitt, Paul E Hertz, Héctor J Álvarez Pérez, and Theodore Garland Jr. Why tropical forest lizards are vulnerable to climate warming. *Proceedings of the Royal Society B: Biological Sciences*, 276(1664):1939–1948, 2009.

[7] Curtis A Deutsch, Joshua J Tewksbury, Raymond B Huey, Kimberly S Sheldon, Cameron K Ghalambor, David C Haak, and Paul R Martin. Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*, 105(18):6668–6672, 2008.

[8] S. Rahmstorf and D. Coumou. Increase of extreme events in a warming world. *Proceedings*

of the National Academy of Sciences of the United States of America, 108(44):17905–17909,

2011.

[9] N. S. Diffenbaugh, D. Singh, J. S. Mankin, D. E. Horton, D. L. Swain, D. Touma, A. Charland,

Y. J. Liu, M. Haugen, M. Tsiang, and B. Rajaratnam. Quantifying the influence of global

warming on unprecedented extreme climate events. *Proceedings of the National Academy of*

Sciences of the United States of America, 114(19):4881–4886, 2017.

[10] D. A. Vasseur, J. P. DeLong, B. Gilbert, H. S. Greig, C. D. G. Harley, K. S. McCann, V. Savage,

T. D. Tunney, and M. I. O'Connor. Increased temperature variation poses a greater risk to

species than climate warming. *Proceedings of the Royal Society B-Biological Sciences*, 281

(1779), 2014.

[11] SD Tuljapurkar and Steven Hecht Orzack. Population dynamics in variable environments I.

Long-run growth rates and extinction. *Theoretical Population Biology*, 18(3):314–342, 1980.

[12] L. A. Real and S. Ellner. Life-history evolution in stochastic environments - a graphical mean-

variance approach. *Ecology*, 73(4):1227–1236, 1992.

[13] J. R. Bernhardt, J. M. Sunday, and M. I. O'Connor. Metabolic theory and the temperature-

size rule explain the temperature dependence of population carrying capacity. *The American*

Naturalist, 192(6):687–697, 2018.

[14] Ian S Pearse, Ryan Paul, and Paul J Ode. Variation in plant defense suppresses herbivore

performance. *Current Biology*, 28(12):1981–1986, 2018.

[15] Kyle E Coblentz, Stephanie Merhoff, and Mark Novak. Quantifying the effects of intraspecific

variation on predator feeding rates through nonlinear averaging. *Functional Ecology*, 35(7): 1560–1571, 2021.

[16] Joey R. Bernhardt, M. I. O'Connor, Jennifer M. Sunday, and Andrew Gonzalez. Life in fluctuating environments. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1814):20190454, 2020. doi: 10.1098/rstb.2019.0454.

[17] Nicholas Kortessis, Margaret W. Simon, Michael Barfield, Gregory E. Glass, Burton H. Singer, and Robert D. Holt. The interplay of movement and spatiotemporal variation in transmission degrades pandemic control. *Proceedings of the National Academy of Sciences*, 117(48):30104–30106, 2020. doi: 10.1073/pnas.2018286117.

[18] Nicholas Kortessis, Gregory Glass, Andrew Gonzalez, Nick W. Ruktanonchai, Margaret W. Simon, Burton Singer, and Robert D. Holt. Metapopulations, the inflationary effect, and consequences for public health. *bioRxiv*, page 2023.10.30.564450; doi: https://doi.org/10.1101/2023.10.30.564450, 2023.

[19] R. Lande. Risks of population extinction from demographic and environmental stochasticity and random catastrophes. *The American Naturalist*, 142(6):911–927, 1993.

[20] Donald Ludwig. The distribution of population survival times. *The American Naturalist*, 147(4):506–526, 1996.

[21] Steinar Engen, Øyvind Bakke, and Aminul Islam. Demographic and environmental stochasticity-concepts and definitions. *Biometrics*, pages 840–846, 1998.

[22] J. M. Ferguson and J. M. Ponciano. Evidence and implications of higher-order scaling in the

environmental variation of animal population growth. *Proceedings of the National Academy of Sciences of the United States of America*, 112(9):2782–2787, 2015.

[23] R Levins. The effect of random variations of different types on population growth. *Proceedings of the National Academy of Sciences*, 62(4):1061–1065, 1969.

[24] John M Drake and Blaine D Griffen. Speed of expansion and extinction in experimental populations. *Ecology Letters*, 12(8):772–8, 2009.

[25] Jake M. Ferguson and José M. Ponciano. Predicting the process of extinction in experimental microcosms and accounting for interspecific interactions in single-species time series. *Ecology Letters*, 17(2):251–259, 2014.

[26] Masahiro Anazawa. The mechanistic basis of discrete-time population models: the role of resource partitioning and spatial aggregation. *Theoretical Population Biology*, 77(3):213–218, 2010.

[27] Peter Turchin. *Complex Population Dynamics*. Princeton University Press, Princeton, 2003.

[28] Franz W Simon and David A Vasseur. Variation cascades: resource pulses and top-down effects across time scales. *Ecology*, 102(4):e03277, 2021.

[29] A. Hastings, K. C. Abbott, K. Cuddington, T. B. Francis, Y. C. Lai, A. Morozov, S. Petrovskii, and M. L. Zeeman. Effects of stochasticity on the length and behaviour of ecological transients. *Journal of the Royal Society Interface*, 18(180), 2021.

[30] Timothy S. Gardner, Charles R. Cantor, and James J. Collins. Construction of a genetic toggle switch in escherichia coli. *Nature*, 403(6767):339–342, 2000.

[31] Jonathan David Touboul, Ann Carla Staver, and Simon Asher Levin. On the complex dynamics of savanna landscapes. *Proceedings of the National Academy of Sciences*, 115(7):E1336–E1345, 2018.

[32] Luca Gammaitoni, Peter Hänggi, Peter Jung, and Fabio Marchesoni. Stochastic resonance. *Reviews of Modern Physics*, 70(1):223, 1998.

[33] Jonathan Dushoff, Joshua B. Plotkin, Simon A. Levin, and David J. D. Earn. Dynamical resonance can account for seasonality of influenza epidemics. *Proceedings of the National Academy of Sciences*, 101(48):16915–16916, 2004.

[34] Roberto Benzi, Giorgio Parisi, Alfonso Sutera, and Angelo Vulpiani. A theory of stochastic resonance in climatic change. *SIAM Journal on Applied Mathematics*, 43(3):565–578, 1983.

[35] Alan A Berryman. *Population cycles: causes and analysis*, pages 3–28. Oxford University Press, 1st edition, 2002.

[36] Jacob Bock Axelsen, Rami Yaari, Bryan T. Grenfell, and Lewi Stone. Multiannual forecasting of seasonal influenza dynamics reveals climatic and evolutionary drivers. *Proceedings of the National Academy of Sciences*, 111(26):9538–9542, 2014.

[37] Z. Zeng, R. M. Nowierski, M. L. Taper, B. Dennis, and W. P. Kemp. Complex population dynamics in the real world: Modeling the influence of time-varying parameters and time lags. *Ecology*, 79(6):2193–2209, 1998.

[38] N. C. Stenseth. Population cycles in voles and lemmings: density dependence and phase dependence in a stochastic world. *Oikos*, 87(3):427–461, 1999.

[39] Alfred James Lotka. *Elements of Physical Biology.* Williams & Wilkins, 1925.

[40] Roy Malcolm Anderson and Robert Mccredie May. The population dynamics of microparasites and their invertebrate hosts. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 291(1054):451–524, 1981.

[41] William Edwin Ricker. Stock and recruitment. *Journal of the Fisheries Board of Canada*, 11 (5):559–623, 1954.

[42] Tomo Royama. *Analytical Population Dynamics*. Springer Science & Business Media, 1992.

[43] Hanna TM Eskola and Stefan AH Geritz. On the mechanistic derivation of various discrete-time population models. *Bulletin of Mathematical Biology*, 69:329–346, 2007.

[44] Masahiro Anazawa. The mechanistic basis of discrete-time population models: the role of resource partitioning and spatial aggregation. *Theoretical Population Biology*, 77(3):213–218, 2010.

[45] Perry De Valpine and Alan Hastings. Fitting population models incorporating process noise and observation error. *Ecological Monographs*, 72(1):57–76, 2002.

[46] Frédéric Barraquand, Adrien Pinot, Nigel G Yoccoz, and Vincent Bretagnolle. Overcompensation and phase effects in a cyclic common vole population: between first and second-order cycles. *Journal of Animal Ecology*, 83(6):1367–1378, 2014.

[47] A. R. Ives, B. Dennis, K. L. Cottingham, and S. R. Carpenter. Estimating community stability and ecological interactions from time-series data. *Ecological Monographs*, 73(2):301–330, 2003.

[48] B. A. Melbourne and A. Hastings. Extinction risk depends strongly on factors contributing to stochasticity. *Nature*, 454(7200):100–103, 2008.

[49] Robert M May. Stability in randomly fluctuating versus deterministic environments. *The American Naturalist*, 107(957):621–650, 1973.

[50] Steven H. Strogatz. *Nonlinear Dynamics and Chaos: with Applications to Physics, Biology, Chemistry, and Engineering*. Westview Press, Boulder, CO, second edition, 2015.

[51] A. Hastings, K. C. Abbott, K. Cuddington, T. Francis, G. Gellner, Y. C. Lai, A. Morozov, S. Petrovskii, K. Scranton, and M. L. Zeeman. Transient phenomena in ecology. *Science*, 361 (6406):eaat6412, 2018.

[52] Jim Michael Cushing, Robert F Costantino, Brian Dennis, Robert Desharnais, and Shandelle Marie Henson. *Chaos in Ecology: Experimental Nonlinear Dynamics*, volume 1. Elsevier, 2003.

[53] Brian Dennis, Robert A. Desharnais, J. M. Cushing, and R. F. Costantino. Nonlinear Demographic Dynamics: Mathematical Models, Statistical Methods, and Biological Experiments. *Ecological Monographs*, 65(3):261–282, 1995.

[54] Eugene Pleasants Odum. *Fundamentals of Ecology*. Saunders Philadelphia, 1959.

[55] Kasper Kristensen, Anders Nielsen, Casper W. Berg, Hans Skaug, and Bradley M. Bell. TMB: Automatic differentiation and Laplace approximation. *Journal of Statistical Software*, 70(5): 1–21, 2016.

[56] J. F. Arnoldi, A. Bideault, M. Loreau, and B. Haegeman. How ecosystems recover from pulse perturbations: A theory of short- to long-term responses. *Journal of Theoretical Biology*, 436: 79–92, 2018.

[57] J. F. Arnoldi, M. Loreau, and B. Haegeman. The inherent multidimensionality of temporal variability: how common and rare species shape stability patterns. *Ecology Letters*, 22(10): 1557–1567, 2019.

[58] Kim M. Cuddington and Peter Yodzis. Black noise and population persistence. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266(1422):969–973, 1999.

[59] Tim Coulson, Pejman Rohani, and Mercedes Pascual. Skeletons, noise and population growth: the end of an old debate? *Trends in Ecology & Evolution*, 19(7):359–364, 2004.

[60] David Alonso, Alan J McKane, and Mercedes Pascual. Stochastic amplification in epidemics. *Journal of The Royal Society Interface*, 4(14):575–582, 2007.

[61] Muhammet Uzuntarla. Inverse stochastic resonance induced by synaptic background activity with unreliable synapses. *Physics Letters A*, 377(38):2585–2589, 2013.

[62] Pejman Rohani, Matthew J Keeling, and Bryan T Grenfell. The interplay between determinism and stochasticity in childhood diseases. *The American Naturalist*, 159(5):469–481, 2002.

[63] Crispin M Mutshinda, Aditya Mishra, Zoe V Finkel, and Andrew J Irwin. Density regulation amplifies environmentally induced population fluctuations. *PeerJ*, 11:e14701, 2023.

[64] Michael Lynch and Bruce Walsh. *Genetics and Analysis of Quantitative Traits*, volume 1. Sinauer Sunderland, MA, 1998.

[65] Peter Chesson. Updates on mechanisms of maintenance of species diversity. *Journal of Ecology*, 106(5):1773–1794, 2018. doi: 10.1111/1365-2745.13035.

[66] Stephen P Ellner, Robin E Snyder, Peter B Adler, and Giles Hooker. Toward a "modern coexistence theory" for the discrete and spatial. *Ecological Monographs*, 92(4):e1548, 2022.

[67] Joel E Cohen. Mathematics is biology's next microscope, only better; biology is mathematics' next physics, only better. *PLoS Biology*, 2(12):e439, 2004.

[68] Tanya L Rogers, Bethany J Johnson, and Stephan B Munch. Chaos is not rare in natural ecosystems. *Nature Ecology & Evolution*, 6(8):1105–1111, 2022.

[69] TJ Clark and Angela D Luis. Nonlinear population dynamics are ubiquitous in animals. *Nature Ecology & Evolution*, 4(1):75–81, 2020.

[70] William J. Sutherland, Peter N.M. Brotherton, Zoe G. Davies, Nancy Ockendon, Nathalie Pettorelli, and Juliet A. Vickery, editors. *Conservation Research, Policy and Practice*. Ecological Reviews. Cambridge University Press, 2020.

[71] Dennis L Murray and Brett K Sandercock. *Population Ecology in Practice*. John Wiley & Sons, 2020.