Universal method for the gentle isolation of intact microvessels from frozen tissue: a multiomic investigation into the neurovasculature

Marina Wakid[a,b], Daniel Almeida[a,b], Zahia Aouabed[a], Reza Rahimian[a], Maria Antonietta Davoli[a], Volodymyr Yerko[a], Elena Leonova-Erko[a], Vincent Richard[d], René Zahedi[d], Gustavo Turecki[a,c], Naguib Mechawar[a,c]

[a]McGill Group for Suicide Studies, Douglas Research Centre, Montreal, Quebec, Canada

[b]Integrated Program in Neuroscience, McGill University, Montreal, Quebec, Canada

[c]Department of Psychiatry, McGill University, Montreal, Quebec, Canada

[d]Segal Cancer Proteomics Centre, Lady Davis Institute, Montreal, Quebec, Canada

Abstract

The neurovascular unit (NVU), comprised of endothelial cells, pericytes, smooth muscle cells, astrocytic endfeet and microglia together with neurons, is paramount for the proper function of the central nervous system. The NVU gatekeeps blood-brain barrier (BBB) properties which, as a system, experiences impairment in several neurological and psychiatric diseases, and contributes to pathogenesis. To better understand function and dysfunction at the NVU, isolation and characterization of the NVU is needed. Here, we describe a singular, standardized protocol to enrich and isolate microvessels from archived snap-frozen human and frozen mouse cerebral cortex using mechanical homogenization and centrifugation-separation that preserves the structural integrity and multicellular composition of microvessel fragments. For the first time, microvessels are isolated from postmortem vmPFC tissue and are comprehensively investigated using both RNA sequencing and Liquid Chromatography with tandem mass spectrometry (LC-MS-MS). Both the transcriptome and proteome are elucidated and compared, demonstrating that the isolated brain microvessel is a robust model for the NVU and can be used to generate highly informative datasets in both physiological and disease contexts.

1. <u>Introduction</u>

To meet the metabolic needs of the 86 billion neurons in the human brain, an elaborate 400 mile-long microvascular network[1,2] supplies blood flow to the deep structures of the cerebral hemispheres and gatekeeps blood-brain barrier (BBB) properties. Extensive research efforts have underscored the BBB as a highly selective cellular system. Ultrastructurally, the BBB consists of continuous non-fenestrated brain microvascular endothelial cells (BMECs) that precisely regulate movement between the blood and brain interface through the expression of specialized solute carriers and efflux transporters[3,4,5,6,7,8,9,10]. Attached pericytes and astrocytic end-feet, which are embedded within the vascular basement membrane, together with smooth muscle cells, are the cell types that comprise the neurovascular unit (NVU)[11] and work in concert to implement coordinated vascular responses to central and peripheral signals. Such responses include the continuous delivery of oxygen and glucose to neurons[12,13,14,15,16,17], homeostatic maintenance of the brain[18,19,20,21,22], the regulation of cerebral blood flow[23,24,25,26,27] and clearance of interstitial fluid[28,29,30,31]. An underappreciated lens through which to investigate disease, NVU dysfunction contributes to cognitive decline and neuronal loss in Alzheimer's disease-related Aβ and tau pathology[32,33,34,35,36,37], traumatic brain injury[38,39,40,41,42,43,44,45,46,47], perivenous myelin lesions observed in multiple sclerosis[48,49,50,51,52,53], as well as multiphasic changes to BBB permeability exhibited after stroke[54,55,56,57,58].

Our understanding of neurovascular development and function has been advanced largely by mouse models. Functional characteristics of the BBB are regulated at the

transcriptomic level and, in recent years, different methodologies have been employed to

investigate the neurovascular transcriptome. Single-cell sequencing studies have

leveraged transgenic-reporter claudin-5-GFP[59,60], Tie2–eGFP[61], and Pdgfrb-eGFP[62]

mouse lines in conjunction with fluorescence-activated cell sorting (FACS) to generate

highly informative transcriptomic datasets of mouse BMECs and other vascular cell types,

whereas other studies have carried out RNA sequencing of BMECs isolated from Rosa-

tdTomato;VE-Cadherin-Cre$_{ERT2}$ mouse models of stroke, MS, TBI and seizure[63]. While

past mouse data have provided precious insight in defining core NVU gene expression

and underscores the relevance of transcriptomic profiling for better understanding

neurovascular function (and dysfunction), recent breakthroughs demonstrate that there

are numerous species-specific differences between mouse and human neurovasculature,

including solute carrier and efflux transporter expression[63,64,65]. Such findings reveal the

partial utility of animal models for the study of human central nervous system (CNS)

disease. Due to scarcity of well-preserved human brain tissue available for research,

transcriptomic profiling in human brain samples has been considerably more limited,

leaving the investigation of vascular cells neglected in favour of non-vascular cell types,

such as neurons and oligodendrocytes. In addition, single-cell or single-nucleus

sequencing used to profile expression in *all* cell populations have yielded very low

populations of endothelial cells and pericytes from human adult and embryonic cortex

samples. Although vessel density, as determined in different brain regions, varies

between 361 and 811 vessels/mm$^{2;66,67,68}$ at an overall endothelial cell density of 4,504 ±

2,666 cells/mm$^{2;69}$, such techniques seem to deplete vascular cells/nuclei for reasons that

are not understood and have impeded analysis of human neurovascular

transcriptomes[70,71,72,73,74]. Recently, detailed transcriptome-wide atlases of human and

mouse brain vascular nuclei in health and disease were generated by two independent

groups[64,65], both addressing the underrepresentation of vascular cell types and

elaborating on species-specific differences in NVU gene expression. Such progress and

tools deepen our understanding of human NVU function, yet certain limitations that persist

challenge further progress in the field: single-cell and single-nucleus sequencing remains

inaccessible to many due to high costs and lack of bioinformatic expertise. Moreover,

there is heavy reliance on brain banks for well-characterized frozen human brain tissue,

which also requires considerable adaptation of techniques initially optimized for fresh

tissue and creates even further disparity in how mouse and human brain tissue are

utilized, even with the same experimental question in mind. The biological and

bioinformatic biases these experimental decisions create, and their extent, are unknown.

Understanding the molecular mechanisms of NVU dysfunction can be achieved by

examining gene expression changes in brain microvessels in different disease contexts;

and the limited number of existing human neurovascular datasets motivates

transcriptomic characterization of more human samples. While it is understood that

BMECs perform the BBB function and that other vascular-associated cell types critically

regulate that function, the study of microvessels as a preserved unit provides greater

insight into the neurovasculature in a manner that dissociated cell types cannot. To this

aim, we use a singular, standardized protocol to enrich and isolate microvessels from

archived snap-frozen human and frozen mouse cerebral cortex using mechanical

homogenization and centrifugation-separation that is gentle enough to dissociate brain

tissue while preserving the structural integrity and multicellular composition of

microvessel fragments.

The common issue of multiple or contaminating cell types in samples used for tissue-

derived RNA sequencing has been largely eliminated by single-cell workflows. However,

single-cell and single-nucleus workflows introduce other significant challenges:

measurements typically suffer from large fractions of observed zeros possibly due to

technical limitations or randomness[75,76]. Moreover, tissue dissociation and storage biases

can induce unwanted transcriptomic alterations and cell type composition differences[77].

Because of this, bulk and single-cell sequencing are complementary strategies in which

the former approach warrants a versatile and effective method for isolating the NVU from

human brain. Several approaches attempting to investigate the isolated NVU have been

developed in the past, albeit with major caveats: FACS-sorting with antibodies against

PECAM1/CD31 and CD13 (targeting the endothelial cell membrane and pericyte cell

membrane, respectively) require fresh brain tissue[78], as do other iterations of microvessel

isolation for the purpose of cell culture expansion[79,80]. Past attempts at selective capture

of endothelial and vascular-associated cells from frozen human brain have relied

exclusively on laser capture microdissection (LCM)[81,82,83,84], which demands considerable

optimization if microvessels are to be used for high-throughput applications downstream

to laser-capture[85]. To date, microvessels obtained from archived postmortem brain tissue

have only been suitable for qPCR and western blot[86], and more comprehensive

knowledge obtained from high-throughput techniques is currently lacking. Critically,

microvessels isolated using the described method are in high yield, posses all major

vascular-associated cell types, and maintain their *in situ* cellular structure, making them

suitable for characterization using high-throughput techniques. The advantages of this

simple protocol are manifold: it does not require the experimental setup needed by single-

nucleus sorting, nor does it require transgenic mice[87], enzymatic dissociation[78,87,88], or

fresh brain tissue[78,87,88]. Importantly, this is the first standardized microvessel isolation

method that is demonstrated to work with snap-frozen brain tissue that is compatible

across high-throughput downstream applications, removing unknown biases introduced

by the use of varied isolation methods.

We have successfully applied the described protocol to postmortem vmPFC tissue from

individuals having died suddenly with no neurological nor psychiatric disorder, as well as

mouse forebrain tissue as proof of concept that the same procedure can be used in both

species. To demonstrate the utility of microvessels isolated from postmortem vmPFC

tissue, we processed 5 samples of extracted total RNA and 3 samples of extracted total

protein with RNA sequencing and liquid chromatography with tandem mass spectrometry

(LC-MS/MS), respectively. Moreover, we sorted BMECs from isolated microvessels using

fluorescence-activated nuclei sorting (FANS) as proof of concept that specific

neurovascular cell types may be further purified if needed. Bioinformatic processing and

analysis of human transcriptomic and proteomic data indicated that isolated samples

showed major enrichment for BMEC, pericyte, SMC, and astrocytic endfeet components

at both the mRNA and protein level, generating the first multiomic datasets from human

brain microvessels.

2. <u>Materials and methods</u>

2.1. Biological materials

Human cortex: This study was approved by the Douglas Hospital Research Ethics Board and written informed consent from next of kin was obtained for each individual included in this study. For each individual, the cause of death was determined by the Quebec Coroner's office and medical records were obtained. Samples were obtained from Caucasian individuals having died suddenly with no neurological nor psychiatric condition (Table 1). Postmortem brain tissues were provided by the Douglas–Bell Canada Brain Bank (www.douglasbrainbank.ca). Frozen gray matter samples were dissected from the ventromedial prefrontal cortex (vmPFC, Brodmann area 11) by expert brain bank staff stored at –80 °C. The postmortem interval (PMI) is metric for the delay between an individual's death and collection and processing of the brain. To assess RNA quality, RNA integrity number (RIN) was measured for brain samples using tissue homogenates, with an average value of 5.34. A total of 5 subjects were subjected to RNA-sequencing and 3 subjects were subjects to Liquid Chromatography with tandem mass spectrometry (LC-MS/MS).

Mouse cortex: Male C57BL/6J mice (n = 2) aged between 120-126 days of age were bred, housed and cared for in accordance to the Canadian Council on Animal Care guidelines (CCAC; http://ccac.ca/en_/standards/guidelines), and all methods were approved by the Animal Care Committee from the Douglas Institute Research Center under protocol number 5570. Mice were housed in standard conditions at 22 ± 1 °C with 60% relative humidity, as well as a 12-h light-dark cycle with food and water available *ad*

libitum[89]. In accordance with pertinent guidelines and regulations, the mice were

anesthetized via intraperitoneal injection of ketamine (10 mg/ml)/xylazine (1 mg/ml) and

transcardially perfused with cold PBS 1X. The frontal cortices were removed and

immediately frozen in liquid nitrogen and then stored in -80°C freezer.

2.2.   Tissue homogenization and cellular fractionation

By virtue of a preserved basement membrane, the structure isolated using the described

method contains all neurovascular-associated cell types and, therefore, are referred to as

"microvessels". The following protocol describes the specific steps used to isolate and

enrich microvessels with retained *in situ* morphology, which is achieved using semi-

automated dissociation of microdissected brain tissue into a thorough homogenate

followed by low-speed centrifugation (schematic overview in Fig. 1). These steps have

been optimized to isolate microvessels from 100 mg of frozen brain tissue, processing a

maximum of 4 samples at a time.

Frozen brain tissue was microdissected on top of dry ice using a razor blade, weighed

using an analytical scale, and transferred to a chilled 1.5ml Eppendorf tube. The

microdissected sample was then transferred to a chilled gentleMACS™ C Tube (Miltenyi

Biotec, Bergisch Gladbach, Germany) containing 2 ml of fresh, cold Homogenization

Buffer Buffer (1M sucrose + 1% Bovine serum albumin (BSA) dissolved in DEPC-treated

water; see supplementary Table 1 for commentary on DEPC-treated water), where the

tissue was fully submerged, and the tube was gently agitated for 30 seconds to encourage

thawing and osmotic equilibrium (Fig. 2a). The rotating paddle of the GentleMACS™

Dissociator (Miltenyi Biotec, Bergisch Gladbach, Germany) was set to program Lung

02.01 (Fig. 1) and after tissue homogenization was complete, the gentleMACS™ C Tube

was returned to wet ice where an additional 8 ml of cold Homogenization Buffer was

transferred into the tube, topping up the homogenate to 10 ml, and gently inverted to mix

and collect homogenate (Fig. 2b-c). The homogenate was transferred to a chilled 15 ml

falcon tube, including any foam produced during paddle rotation (Fig. 2d) and gently

inverted a second time in order to prevent the formation of a foamy seal atop of the

homogenate when left sitting. The homogenate was centrifuged (Beckman Coulter, model

Allegra X-14R) at 3200 g for 30 mins at 4°C. Once centrifugation is complete, a

microvessel-enriched pellet forms at the bottom of the falcon tube (Fig. 2e). The

supernatant was carefully vacuum-aspirated (which may include an upper layer of

clumped, dissociated myelin, similar to milk skin on top of boiled milk) without disturbing

the microvessel-enriched pellet (Fig. 2e-f). Anticipated methodological issues are posed

in supplementary Table 3, along with potential solutions.

2.3. Detection of microvessels from the enriched pellet using histology and immunohistochemistry

To first assess yield and stability of microvessel isolation, BCIP/NBT (5-bromo-4-chloro-

3'-indolyphosphate and nitro-blue tetrazolium) was used as a chromogenic substrate for

endothelial enzyme alkaline phosphatase. Within the brain, alkaline phosphatase activity

is localized to cerebral blood vessels[90,91,92,93,94,95,96]. BCIP is hydrolyzed by the alkaline

phosphatase expressed exclusively in endothelial cells to form a blue intermediate that is

then oxidized by NBT to produce a dimer, leaving an intense insoluble purple dye. To this

effect, SIGMA*FAST*™ BCIP®/NBT tablet (Sigma-Aldrich, Missouri, United States) were

crushed and then dissolved in the fresh Homogenization Buffer according to

manufacturer's instructions (1 tablet per 10 mls of solution). Crushing the tablet first

encourages faster dissolution in what is a highly viscous buffer. Greater

immunophenotypic characterization of isolated microvessels was carried out following

resuspension of additional microvessel-enriched pellets. Primary antibodies raised

against canonical expression markers for BMECs (Vimentin, Laminin and PECAM1), tight

junctions (CLDN5), astrocytic end-feet (AQP4), pericytes (PDGFRβ) were utilized, along

with appropriate fluorophore-conjugated secondary antibodies to thoroughly characterize

collected microvessels. The microvessel-enriched pellet was gently resuspended in 500

μl of cold PBS, with 50 ul transferred into each well of an 8-well chamber slide (Nunc™

Lab-Tek™ II Chamber Slide™ System, Thermo Scientific™, Massachusetts, United

States). The chamber slide was then left open faced in a 37°C oven (Fisher Scientific,

model Fisherbrand™ Isotemp™) overnight to evaporate the PBS, after which

microvessels dry flush to the surface of the slide. It is important to note that

immunofluorescent visualization must omit any steps in which the microvessel-enriched

pellet is filtered through a cellular strainer because microvessels cannot be released from

the strainer and, therefore, cannot be mounted onto a microscope slide. Microvessels

were fixed by submerging them to a depth of 2–3 cm with cold 100% methanol for 15 min

on ice, after which wells were washed with 1X PBS three times for 5 min. Subsequently,

microvessels were incubated in blocking buffer (1% BSA + 0.5% Triton X dissolved in

PBS) under agitation for 60 min at 4°C, followed by incubation of each well in 500 ul of

primary antibody dilution (1:500 in 1% BSA + 2% normal donkey serum + 0.5% Triton X

dissolved in PBS) under agitation overnight at 4°C. Wells were washed with 1X PBS three

times for 5 min and finally incubated in fluorophore-conjugated secondary antibody

dilution (1:500-1000 in 1% BSA + 2% normal donkey serum + 0.5% Triton X dissolved in

PBS) under agitation for 2 hours at room temperature. A final round of washes was

followed by removal of the media chamber according to manufacturer's instructions.

Microvessels were coverslipped using VECTASHIELD Antifade Mounting Medium with

DAPI (Vector Laboratories, California, United States) and place coverslip.

To expand upon experimental applications possible with isolated human brain

microvessels, we adapted several downstream techniques including total RNA extraction,

RNA library construction for downstream RNA-sequencing, FANS of BMECs, and total

protein extraction for downstream Liquid Chromatography with tandem mass

spectrometry (LC-MS/MS).

2.4.   RNA extraction from isolated microvessels

Total RNA extraction from isolated microvessels is carried out starting with resuspension

of the microvessel-enriched pellet. As previously described, the microvessel-enriched

pellet was gently resuspended in 500 µl of cold PBS and gradually pipetted through a 35

µm Strainer Cap for FlowTubes™ (Canada Peptide, Quebec, Canada) using vacuum-

aspiration underneath to encourage filtration. The result is intact microvessels trapped

within the mesh of the strainer, where smaller cellular debris and free-floating nuclei have

passed through. After microvessels were trapped within the 35 µm strainer, RNA was

extracted using the Single Cell RNA Purification Kit (Norgen Biotek Corp., Ontario,

Canada). Using a flat-ended spatula and point-tip forceps, the mesh of the cellular sieve

was removed from its plastic frame (Fig. 2i) and immediately submerged into 100 μl of RL

buffer in a 1.5 ml Eppendorf tube, according to step 1A of the manufacturer's protocol.

After the addition of 100 μl of fresh 70% ETOH, the solution was pipetted 10 times to

wash through the mesh. The mesh was then discarded and the remaining steps according

to manufacturer's instructions were carried out, including the on-column DNase digestion

step. Total RNA was quantified using the Agilent TapeStation 2200 (Agilent Technologies,

model 2200 TapeStation; for total RNA concentration and RIN, see supplementary Table

2).

2.5.　Protein　extraction　from　microvessel-enriched　pellet　or　strained

microvessels

Total protein has been successfully extracted from either the microvessel-enriched pellet

or microvessels stained through a cellular sieve, with one modification to Homogenization

Buffer preparation required in which cOmplete™ EDTA-free Protease Inhibitor Cocktail

(Roche, Basel, Switzerland) and Phosphatase Inhibitor Cocktail 2 (Sigma-Aldrich,

Missouri, United States) are added according to the manufacturer's instructions and

based on final volume prepared. With the procedure identical when using strained

microvessels (on a sieve), we elaborate protein extraction from frozen microvessel-

enriched pellets: Microvessel-enriched pellets were resuspended in 5% sodium dodecyl

sulfate (SDS), 100 mM TRIS (pH 7.8) and protein extracted by heating to 99°C for 10

minutes on a hotplate with tube rack. The resuspended pellet was subjected to probe-

based sonication (Fisher Scientific, model Thermo Sonic Dismembrator) at 25%

amplitude for 3 cycles of 5 seconds and lysates were clarified by centrifugation at 20,000

g for 5 minutes. For estimation of protein concentration, approximately 10% of the sample

was aliquoted and diluted to <1% SDS and used for estimation of protein concentration

by Pierce™ bicinchoninic acid assay (BCA) Protein Assay Kit (Pierce Biotechnology Inc.,

Massachusetts, United States). It is important to note that Total protein and total RNA

extraction from the same microvessel-enriched pellet has not been validated. Two

microvessel-enriched pellets per sample should be prepared, one for RNA and one for

protein extraction.

2.6.   FANS-sorting of BMECs from microvessel-enriched pellet

If further isolation of BMECs from the microvessel enriched pellet is desired, FANS-

sorting of ETS-related gene (ERG)[+] BMECs from the obtained microvessel-enriched

pellet can be performed. ERG is a transcription factor whose expression in normal

physiological conditions is found exclusively in endothelial nuclei, making it a highly

specific pan-endothelial nuclear marker[60,64,65,97,98]. FANS of BMECs can be carried out

following the resuspension of the microvessel-enriched pellet. The pellet is resuspended

and centrifuged (Eppendorf, model 5430) at 500 g for 3 min at 4°C, resulting in a pellet

once again. The supernatant was aspirated to remove any remaining sucrose, and the

washed pellet was resuspended in 250 ul of the following antibody solution: recombinant

Alexa Fluor® 647 anti-ERG (1:100 dilution; ab196149, Abcam, Massachusetts, United

States) in 0.5% BSA + 0.1% Triton in PBS, incubated under gentle agitation for 2 hours

at 4 °C. In the last 10 minutes of the 2-hour incubation, 1 ul of Hoechst 33342 dye

(Invitrogen™, Massachusetts, United States) was added to stain nuclear DNA. The

suspension was filtered through a 35 μm Strainer Cap for FlowTubes™ and transferred

the flow-through into a FACS tube. The BD FACSAria™ Fusion Flow Cytometer (BD

Biosciences, California, United States) was used for the sorting of the ERG⁺ population.

The gating strategy used for sorting was as follows: doublet discrimination was achieved

by the gating of Hoechst 33342 stained singlets in a FSC-A versus Hoechst-A plot using

a 350 nm UV laser and a 450/50 filter. The subsequent ERG⁺ population was gated in an

Alexa Fluor 647-A vs FSC-A plot using a 640 nm laser in combination with a 730/45 filter.

Gating was applied to filter singlets using physical parameters and violet fluorescence

(405-nm laser, 525/50 filter). Nonoverlapping gates were adjusted to collect endothelial

nuclei based on Alexa Fluor® 647 anti-ERG immunoreactivity (640-nm laser, 730/45

filter). This approach was chosen due to the well-known observation that forward scatter

is proportional to size.

2.7.   Library construction and RNA-sequencing

Microvessel-enriched pellets yielded an average of 8.54 ug/ul of total RNA per sample

(supplementary Table 2). Libraries were then constructed using the SMARTer Stranded

Total RNA-Seq Kit v3 - Pico Input Mammalian (Takara Bio Inc., Shiga, Japan), which

features integration of unique molecular identifiers (UMIs). Libraries were constructed

using 10 ng of RNA as input, 2 minutes of fragmentation at 94°C (Applied Biosystems

Corporation, model ProFlex PCR), 5 cycles of amplification at PCR1 (addition of Illumina

adapters and indexes), 12 cycles of amplification at PCR2 (final RNA-seq library

amplification) and clean-up of final library using NucleoMag NGS Clean-up and Size

Select beads (Takara Bio Inc., Shiga, Japan). Libraries were then quantified at the

Genome Quebec Innovation Centre (Montreal, Quebec) using a KAPA Library

Quantification kit (Kapa Biosystems, USA), and average fragment size was determined

using a LabChip GX (PerkinElmer, USA) instrument. Libraries were sequenced on the

NovaSeq 6000 system (Illumina, Inc., California, United States) using S4 flow cells with

100bp PE sequencing kits (schematic overview in Fig. 3).

2.8.   Bioinformatic pipeline and analyses of RNA sequencing data

UMI extraction, alignment, de-duplication, metrics and gene counting

RNA sequencing of microvessel libraries yielded an average ~72 million reads per library,

which were then processed following our in-house bioinformatic pipeline. Briefly, UMI

extraction based on fastq files was performed using the module extract of umi_tools

(v.1.1.2)[99]. Reads were then aligned to the Human Reference Genome (GRCh38) using

the STAR software v2.5.4b[100] with Ensembl v90 as the annotation file and using the

parameters: **--twopassMode   Basic   --outSAMprimaryFlag   AllBestScore   --**

outFilterIntronMotifs   RemoveNoncanonical   --outSAMtype   BAM

SortedByCoordinate --quantMode TranscriptomeSAM GeneCounts. Resultant bam

files were then sorted and indexed using SAMtools (v.1.3.1)[101] and duplicate reads with

the same UMI were removed using dedup module of umi_tools (v.1.1.2)[99]. Different

metrics including the fraction of the exonic, intronic and intergenic reads were calculated

using CollectRnaSeqMetrics module of Picard (version 1.129; Picard2019toolkit, Broad

Institute, GitHub repository). The expected counts and the transcripts per million (TPMs)

were generated using RSEM (v1.3.3; reverse strand mode)[102].

Computational deconvolution of RNA sequencing data

Two approaches were used for computational deconvolution of RNA sequencing data.

The first approach was performed using the web tool BrainDeconvShiny

(https://voineagulab.shinyapps.io/BrainDeconvShiny/), which implements the best

performing algorithms and all cell-type signatures for brain, as well as goodness-of-fit

calculations based on benchmark work conducted by Sutton et al. (2022)[103]. UMI counts

for each gene were converted to transcripts per million (TPMs) to account for the varying

length of gene and sequencing depth of each sample, facilitating comparisons across

samples. Genes with zero TPMs were then removed, 34370 genes from the original

58303 passed this QC criteria and were used as input into the BrainDeconvShiny tool.

Deconvolution was performed twice: the first approach used average expression in

control samples from the Velmeshev et al. (2019)[71] single nuclei dataset (raw data

available through the Sequence Read Archive, accession number PRJNA434002;

analyzed data available at https://autism.cells.ucsc.edu) as the reference signature for

annotated cell types and CIBERSORT v1.04 algorithm to deconvolute sample profiles

and estimate cell-type composition. The second approach used the MultiBrain (MB)

composite signature[103] generated by averaging the expression signatures of five datasets

for five cell types (neurons, astrocytes, oligodendrocytes, microglia, and endothelia). MB

was used as cell type signatures for deconvolution of our dataset using CIBERSORT

v1.04.

The second approach to deconvolution was performed in-house. Postmortem NVU

single-nucleus data generated on the 10X Genomics Chromium system was accessed

from Yang et al. (2022; raw sequencing data are accessible on GEO using the accession

code GSE163577)[65] and used as the reference signature. Seurat[104] was used to pre-

process raw count expression data, removing genes with less than 3 cells or cells with

less than 200 expressed genes. 23054 genes from a total of 23537 and 141468 nuclei

from a total of 143793 passed these QC criteria. To generate the Yang signature input

for the CIBERSORTX tool[105], counts per million (CPM) values were averaged across

nuclei of each cell type.

2.9. Gas Chromatography−Mass Spectrometry (LC-MS) with tandem mass tag
403
fractionation

Protein digestion and TMT labelling

Extracted proteins were reduced with 20 mM tris(2-carboxyethyl)phosphine (TCEP) at

60°C prior to alkylation with 25 mM iodoacetamide at room temperature for 30 minutes in

the dark. An equivalent of 10 µg of total protein was used for proteolytic digestion using

suspension trapping (S-TRAP). Briefly, samples were acidified with phosphoric (1.3%

final concentration), and then diluted 6-fold in STRAP loading buffer (9:1 methanol:water

in 100 mM TEAB, pH 8.5). Samples were loaded onto S-TRAP Micro cartridges (Protifi

LLC, Huntington, NY) prior to centrifugation at 2000 g for 2 minutes and washed three

times with 50 µl of STRAP loading buffer. Proteins were digested with trypsin (Sigma

Corporation, Kanagawa, Japan) at a 1:10 enzyme to substrate ratio for 2 hours at 47°C.

Peptides were sequentially eluted in 100 mM TEAB, 0.1% formic acid in water, and 50%

acetonitrile, and lyophilized to dryness prior to labelling with TMT 10plex reagents

according to the vendor's specifications (Thermo Fisher Scientific, Massachusetts, United

States).

Offline high-pH reversed-phase fractionation

Labelled peptides were pooled and again lyophilized to dryness, and then reconstituted in 5 mM ammonium formate and fractionated offline by high pH reversed-phase separation using a Waters Xbridge Peptide BEH C18 column (2.1 x 150mm, 2.5 μm) (Waters Corp., Massachusetts, United States) and an Agilent 1290 LC system (Agilent Technologies, California, United States). Binary gradient elution was performed at a flow rate of 400 μL/minute using mobile phase A) 5 mM ammonium formate adjusted with ammonium hydroxide to pH 10, and B) 100% acetonitrile using the following program: 0 min, 0% B ; 2min , 0% B ; 2.1min , 5% B ; 25min, 30% B ; 30min , 80% B ; 32min , 80% B ; 2 min post-run , 0% B. Fractions were collected every 30 seconds and the first and last 7 fractions were concatenated such that even and odd samples were pooled separately, resulting in 20 fractions in total.

LC-MS/MS

Samples were analyzed by data dependent acquisition (DDA) using an Easy-nLC 1200 online coupled to a Q Exactive Plus (both Thermo Fisher Scientific, Massachusetts, United States). Samples were first loaded onto a pre-column (Acclaim PepMap 100 C18, 3 μm particle size, 75 μm inner diameter x 2 cm length) in 0.1% formic acid (buffer A). Peptides were then separated using a 60-min binary gradient ranging from 3-40% B (84% acetonitrile, 0.1% formic acid) on the analytical column (Acclaim PepMap 100 C18, 2 μm particle size, 75 μm inner diameter x 25 cm length) at 300 nL/min. MS spectra were acquired from m/z 350-1,500 at a resolution of 70,000, with an automatic gain control

(AGC) target of 1 x 10^6 ions and a maximum injection time of 50 ms. The 15 most intense

ions (charge states +2 to +4) were isolated with a window of m/z 1.2, an AGC target of 2

x 104 and a maximum injection time of 64 ms and fragmented using a normalized higher-

energy collisional dissociation (HCD) energy of 28. MS/MS spectra were acquired at a

resolution of 17,500 and the dynamic exclusion was set to 30 s. DDA MS raw data was

processed with Proteome Discoverer 2.5 (Thermo Scientific, Massachusetts, United

States) and searched using Sequest HT against a FASTA file containing all reviewed

protein sequences of the canonical human proteome without isoforms downloaded from

Uniprot (https://www.uniprot.org). The enzyme specificity was set to trypsin with a

maximum of 2 missed cleavages. Carbamidomethylation of cysteine was set as static

modification and methionine oxidation as variable modification. The precursor ion mass

tolerance was set to 10 ppm, and the product ion mass tolerance was set to 0.02 Da. The

percolator node was used and the data was filtered using a false discovery rate (FDR)

cut-off of 1% at both the peptide and protein level. The Minora feature detector node of

Proteome Discoverer was used for precursor-based label free quantitation.

3. <u>Results</u>

3.1. Immunophenotypic characterization reveals isolated brain microvessels have

preserved morphology and expression integrity

Human brain microvessels were isolated from 5 frozen vmPFC grey matter samples

microdissected from healthy individuals who died from peripheral diseases or natural

events (Table 1). Because the same basement membrane that maintains the integrity of

the endothelium also ensheaths astrocytic endfeet as well as pericytes or smooth muscle

cells, it is impossible to isolate solely BMECs and, therefore, isolated microvessels also

contained microvessel-associated cell types. Importantly, the described method allows

for isolation of microvessels from other brain regions (data not shown) and has

additionally been performed in the dorsolateral prefrontal cortex (dlPFC, Brodmann area

8/9), the primary visual cortex (Brodmann area 17), as well as the hippocampus

(Brodmann area 28).

Following isolation of human brain microvessels, chromogenic staining of resuspended

pellets using BCIP/NBT (5-bromo-4-chloro-3'-indolyphosphate and nitro-blue

tetrazolium), a substrate for endothelial enzyme alkaline phosphatase, demonstrated

isolation and enrichment of predominantly microvessels from vmPFC tissue samples (Fig.

4a-b), with similar success when isolating microvessels from mouse cortex (Fig. 4c). The

cytoarchitecture of brain microvessels is both complex and comprised of several cell

types, thus, following isolation of brain microvessels, we aimed to characterize the

structure and morphological integrity of isolated microvessels. Immunophenotypic

characterization of several NVU markers revealed expression of vimentin (VIM; anti-

Vimentin antibody RV202, Abcam), laminins (LAM; anti-Laminin antibody L9393,

Sigma-Aldrich), claudin 5 (CLDN5, anti-Claudin 5 antibody ab15106, Abcam), platelet

derived growth factor receptor beta (PDGFRβ, anti-PDGFRβ monoclonal antibody

G.290.3, Thermo Fisher Scientific) and aquaporin 4 (AQP4, anti-Aquaporin 4 antibody

[4/18], Abcam). More precisely, vimentin (Fig. 5a-b), a regulator of actin cytoskeleton in

smooth muscle[106], and laminins (Fig. 5c), the major basement membrane component

responsible for signal transduction via interaction with cell surface receptors[107], are

expressed continuously and homogenously across the entire length of the endothelial

surface. CLDN5, a major functional constituent of tight junctions[108], was also stained with

no apparent discontinuity, suggesting that endothelial tight junctions were preserved (Fig.

5d). Pericyte coverage of BMECs, immunolabeled by regulator of angiogenesis and

vascular stability PDGFRβ[109] was detected adhering to the surface of microvessels (Fig.

5e). These results suggest that the overall *in situ* brain microvessel structure is conserved

after isolation.

Astrocytes serve multiple essential functions in supporting normal brain physiology[110].

This is, in part, due to the extension of astrocytic endfeet that surround approximately

99% of the cerebrovascular surface[111] which, in conjunction with pericytes[112], regulate

expression of molecules that form the BBB including tight junction, enzymatic, and

transporter proteins[113,114,115]. Although astrocytes were not co-isolated with microvessels,

their perivascular endfeet are ensheathed within the same vascular basement membrane

as endothelial cells and pericytes, making it possible that astrocytic endfeet remained

attached after tissue homogenization. Because of this, we further observed astrocyte

vascular coverage of isolated microvessels. AQP4, a water channel protein essential for

the maintenance of osmotic composition and volume within the interstitial, glial and

neuronal compartments[116,117], is expressed at the vessel-facing astrocytic membrane and

superimposes the walls of isolated microvessels (Fig. 5f).

Finally, we examined possible contamination of our microvessel preparations by other

cell types found within the brain. Immunolabelling for myelin basic protein (MBP; anti-

Myelin Basic Protein antibody, BioLegend) was not detected (Fig. 5g); nor did

immunolabelling for neuronal nuclear protein (NeuN; anti-NeuN antibody clone A60,

Sigma-Aldrich) reveal the presence of neuronal elements (Fig. 5h). Thus, neurons and

oligodendrocytes were consistently not observed to be co-isolated with microvessels.

Together, these results indicate that the described protocol allows for the isolation and

enrichment of structurally preserved brain microvessel fragments that are comprised of

BMECs, astrocytic endfeet, pericytes, and tight junction proteins.

3.2. Computational deconvolution and characterization of transcriptomic data

indicates high microvessel yield after isolation

To estimate the enrichment of our microvessel preparations, we performed computational

deconvolution using calculated TPMs (Fig. 6a-b) using the BrainDeconvShiny tool

(https://voineagulab.shinyapps.io/BrainDeconvShiny/). To demonstrate stability of

outcome regardless of resources used, different iterations of deconvolution were

performed utilizing single-nucleus datasets from Velmeshev et al. (2019; VL)[71] and the

MultiBrain (MB) composite signature from Sutton et al. (2022)[103] as the reference

signature, using the CIBERSORT v1.04 algorithm to deconvolute sample profiles and

estimate cell type composition. Regardless of the approach used, majorly endothelial

gene expression was returned, with an average of 94.92% (Fig. 6a) and 86.91% (Fig. 6b).

Some contamination from neurons (1.29% and 4.66%, respectively) as well as negligible

contamination from oligodendrocytes and microglia was observed. When using either

dataset as the reference signature, limited presence of astrocytic genes was observed

(1.16% and 6.11%, respectively), which may represent the contribution of astrocytic

endfeet that cover the length of the neurovasculature. To estimate the multicellular

composition of our microvessels at a finer resolution, computational deconvolution was

performed a third time using the single-nucleus data generated by Yang et al. (2022)[65],

in which the different neurovascular cell type signatures were determined, as the

reference signature (Fig. 6c). Averaged CPMs across nuclei of each cell type were used

as input for the CIBERSORTX tool[105], which estimated an average composition of 44.02%

capillary and 37.62% SMC, along with much lower estimations for pericyte, arterial,

venous, astrocyte and perivascular fibroblast genes (Fig. 6c). Although differentially

distributed along the arteriovenous axis, both SMCs and pericytes are embedded within

the vascular basement membrane[11] and, therefore, it is unlikely that our isolated method

preferentially selects one cell type over the other. Because of this, and the known

similarity in molecular signature between SMCs and pericytes[118,119], as well as the high

percentage of captured capillary segments (in which pericytes are predominantly

observed)[120,121,122,123], the surprisingly low estimation of pericyte genes may represent a

limitation in comparing single-nuclei and bulk tissue datasets to one another. Critically,

an average of 90.4% of the total TPMs across samples were assigned to NVU-constituent

cell types.

To explore our transcriptomic data, the top 10% of most highly expressed genes were

designated (a total of 3968 genes) and used as input for over-representation analysis

(ORA) using the enrichR package in R, selecting the "Descartes Cell Types and Tissue

2021" database to identify gene sets that are statistically over-represented (Fig. 7a-b and

Supplementary Tables 4). The threshold value of enrichment was selected by a p-value

<0.05 and, as shown, over-represented genes were dramatically enriched for vascular-

related terms, such as "Vascular endothelial cells in Cerebellum" and "Vascular

endothelial cells in Cerebrum". Genes behind enriched terms were extracted and the

expression of a subset of known brain endothelial, pericyte, astrocytic and smooth muscle

genes in isolated microvessels were examined (Fig. 7c-i). As expected, microvessels had

increased expression of canonical endothelial genes such as CLDN5, CDH5, SLC2A1,

ABCB1, VWF, and MFSD2A; with highest expression in endothelial genes ACTG1, B2M,

BSG, EEF1A1, HLA-B, HLA-E, SPARCL1, TMSB10, and VIM (Fig. 7a).

Additionally, there was enrichment for other neurovascular cell types, as suggested by

canonical pericyte genes PDGFRβ, MCAM, RGS5, AGRN, and NOTCH3 (Fig. 7b); as

well as canonical smooth muscle genes ACTA2, MYL6, MYL9, TAGLN, and LGALS1

(Fig. 7c). Highest expression was detected in pericyte genes CALD1, FN1, IFGBP7,

RGS5, SPARC and SPARCL1 (Fig. 7b); as well as smooth muscle genes ACTG1,

ACTN4, CALD1, MYL6, and PTMA (Fig. 7c). Complimentary to immunophenotypic

characterization of isolated microvessels, several genes whose products are involved in

junctional complex maintenance and organization (tight junctions, Fig. 7f; adherens

junctions, Fig. 7g) are found in the top 10% of most highly expressed genes, for e.g.,

CLDN5, CTNNB1, CTNND1, ITGB1, JAM1, OCLN, TJP1, and TJP2; with additional

vascular makers expressed at lower transcripts per million are present throughout the

entire dataset. Expression of astrocytic genes were observed to a lesser extent, a number

of which have been previously validated as markers of astrocytic processes or

endfeet[124,125,126], namely EZR, GJA1, RDX, SLC1A2, and SLC1A3 (Fig. 7h). To better

visualize the proportion of expression contributed by these genes, TPMs were

summarized by cell type over the total number of TPMs (Fig. 7i), demonstrating

overrepresentation of endothelial, smooth muscle cell, and pericyte genes in samples.

Intriguingly, results from ORA reveal genes that are of interest in numerous disease

contexts. Enriched in the described dataset is gene CLDN5, an indispensable junctional

protein for the correct organization of tight junctions and maintenance of BMEC

integrity[108], which was previously reported to be downregulated in the nucleus accumbens

of depressed suicides[127] and targeted altered epigenetic regulation via histone

deacetylase 1 (HDAC1)[128]. Enriched pericyte genes TXNIP, RUNX1T1, ITGA1 and

DOCK9 are also reported to be differentially expressed in Schizophrenia[129]. Similarly

enriched in this dataset are angiogenic growth factors EGFL7, FLT1, VWF and antigen-

presentation machinery B2M and HLA-E, all of which are upregulated in a subpopulation

of angiogenic BMECs that are induced in subjects with Alzheimer's disease[130],

suggesting a compensatory angiogenic and immune response in AD pathogenesis.

Likewise, endothelial genes PICALM, INPP5D, ADAMTS1, PLCG2 that are found in the

top 10% of the microvessel dataset are differentially expressed in Alzheimer's disease[65].

Recent breakthroughs in deciphering the underlying etiology of Huntington's disease

reveals aberrant downregulation of endothelial ABCB1, ABCG2, SLC2A1 and

MFSD2A as well as mural (which constitutes pericytes and vascular smooth muscle cells)

PDGFRB, SLC20A2 and FTH1[64] – mutations in which are known to cause HD-like

syndromes with primary pathology localized in the basal ganglia[131,132]. Indeed, brain

microvessel datasets such as these integrated with others, like as those from

experimental models, will expedite our understanding of neurovascular contributions to

mood disorders and neurodegenerative disease, and even further propagate hypothesis

generation for established vascular diseases, such as white matter vascular dementia in

which each cell type of the NVU exhibits a specific disease-associated expression

signature[133].

3.3. High correspondence between generated transcriptomic data and published

neurovascular dataset

As a means to assess our isolation method, our top 10% most highly expressed genes

were juxtaposed to validated neurovascular cell type-defining markers, as designated by

Garcia et al. (2022)[64] based on sequencing of 4,992 and 11,689 vascular nuclei from *ex*

vivo and postmortem brain tissue, respectively. Results indicated substantial overlap

between the three datasets for all NVU-constituent cell types (Fig. 8), including arteriole-

defining genes (Fig. 8a), capillary-defining genes (Fig. 8b), venule-defining genes (Fig.

8c), arteriolar SMC-defining genes (Fig. 8d), pericyte-defining genes (Fig. 8e), venular

SMC-defining genes (Fig. 8f), and perivascular fibroblast-defining genes (Fig. 8g).

Interestingly, overlap was consistently greater between the two postmortem datasets.

3.4. Proteomic characterization of isolated brain microvessels

While transcript level may show positive correlations with protein level, protein abundance

should not necessarily be inferred from RNA sequencing counts[134,135], hence,

interrogation of the highly dynamic proteome is needed to speculate the functional

consequences of changes in protein expression. We sought to interrogate NVU-specific

proteomic signatures using total protein extracted from isolated microvessels.

Microvessel-enriched pellets were prepared from 3 frozen vmPFC grey matter samples

microdissected from healthy individuals who died of peripheral diseases or natural events

(Table 1). Although resuspended and strained pellets yield sufficient protein needed to

perform LC-MS/MS (validated, data not shown), straining was omitted in favour of protein

extraction directly from pellets to maximize microvessel material input for proteomic

interrogation. Using Tandem Mass Tag (TMT) isobaric labeling and sample fractionation

of peptides, global, relative quantitation of a total of 1638 individual proteins were

detected from microvessel-enriched pellets (quality parametres shown in supplementary

Fig. 2). Importantly, there was significant overlap between transcriptomic and proteomic

output, with 1635/1638 (99.8%) of detected proteins likewise identified in the

transcriptomics data (albeit no corresponding transcripts were detected for proteins F9,

DCD, and SERPINB12); and with 1034/1638 (62.5%) proteins found in the top 10% of

most highly expressed transcripts, resulting in an overall 23% overlap between high

expressors in both datasets (Fig. 9a).

Proteins known to be expressed by vascular-associated cell types and perivascular

extracellular matrix were positively identified in all 3 samples. Several canonical

endothelial markers were detected with high peptide abundance, including vimentin,

several protein subunits of laminin (LAMA2, LAMC1, LAMB2, LAMA5, LAMA4, LAMA1,

LAMB1, LAMC3, LAMA3), OCLN, TJP1, TJP2, VWF, VWA1, BSG, PECAM1,

Monocarboxylate transporter 1 (SLC16A1), broad substrate specificity ATP-binding

cassette transporter (ABCG2), ESAM, CLDN5, CDH5, ATP-binding cassette sub-family

D member 3 (ABCD3), and Protocadherin-1 (PCDH1) (normalized average abundance

for these proteins are listed in Fig. 9b). As expected, enrichment terms returned by ORA

(using corresponding gene names as input) were predominantly vascular-associated

(Fig. 9c and supplementary Table 5) and normalized abundance of known brain

endothelial, pericyte, astrocytic and smooth muscle cell proteins were high (Fig. 9d-i).

The BBB poses as a major pharmacological barrier as BMECs express a vast array of

enzymes and transport systems that facilitate brain uptake processes of essential

nutrients and neuroactive agents across the BBB[136], and therefore control the rate and

extent to which drugs are able to reach the brain parenchyma via the transcellular

pathway[137]. There is a pressing need for greater knowledge surrounding the expression

and functionality of these systems at the human BBB as the majority of data comes from

either *in vitro* cell culture or animal studies, making *in vitro* to *in vivo* or interspecies scaling

less reliable. Analyses revealed high abundance of SLC2A1/GLUT1, a transmembrane

protein responsible for the facilitated diffusion of glucose[138], and the two glutamate

transporters SLC1A2/EAAT2 and SLC1A3/EAAT1 in brain microvessels. Transporters

SLC7A5/LAT1 and SLC3A2/4F2hc, which supply the brain with large neutral amino

acids[139,140,141], and SLC16A1/MCT1 and SLC16A2/MCT8, which are involved in the

transport of monocarboxylates[142] and T3 thyroid hormone[143] at the BBB, respectively, are

also found in the described proteomic dataset. Also found are ABCB1/P-glycoprotein and

breast cancer-related protein ABCG2/BCRP, the principal ABC efflux

transporters[144,145,146] that limit entry of drug candidates, toxic compounds, as well as

xenobiotics from the central nervous system[147,148,149,150,151,152]. Due to major species-

specific differences in transporter expression profiles, there is utility in obtaining absolute

protein amounts as they may elucidate the contribution of each transporter in facilitating

the entry of endogenous substances and nutrients like glucose, glutamate, and amino

and fatty acids into the brain, in addition to drugs and other xenobiotics that exploit these

mechanisms[153].

3.5. Further deconstruction of isolated brain microvessels using FANS

The use of frozen brain tissue demands sorting of target nuclei as opposed to intact cells

and, therefore, requires the use of nuclear fluorescent tags to facilitate isolation of

endothelial nuclei. Expression of transcription factor ETS-related gene (ERG) has been

previously validated as an endothelial-specific nuclear marker[98,154,155,156]. As an additional

testament to the versatility in downstream usability of the described method, resuspended

microvessels subjected to a 2-hour incubation with anti-ERG antibody under rotation is

sufficient for vascular disassembly and to expose the ERG epitope for endothelial nuclei

immunolabelling (Fig. 10a-b). Importantly, enzymatic digestion is not needed as

previously described[78,157,158] and, in combination with the appropriate gating strategy

adjusted to collect endothelial nuclei, is sufficient for the purification of endothelial nuclei

from frozen postmortem brain tissue (data not shown). The potential to isolate ERG[+]

endothelial nuclei from postmortem microvessels under different physiological

conditions[159,160,161,162,163,164,165] is an advantage of our approach and addresses a growing

interest in the use of primary BMECs, circumventing phenotypical or gene expression

changes induced in primary BMEC cultures by prolonged adherence steps used in other

isolation protocols[166,167,168,169].

4. Discussion

We describe a singular, standardized protocol to enrich and isolate microvessels from

archived snap-frozen human with the ability to apply the same protocol to frozen mouse

cerebral cortex. Integral to this method are three factors that confer gentility and simplicity:

1) the correct molarity of the sucrose-based buffer to separate and cushion microvessels

during centrifugation-separation, 2) centrifugation-separation of microvessels from the

rest of tissue homogenate occurs in a 10 ml volume, which facilitates the formation of a

microvessel-enriched pellet as heavier structures reach the bottom of the 15 ml falcon

tube, and 3) the limited number of wash steps that minimize eventual damage done to

the integrity of microvessel fragments. Microvessel enrichment and stability was

assessed by chromogenic staining of microvessels using BCIP/NBT substrate and,

through immunophenotypic characterization, we show that isolated microvessel

fragments are comprised of NVU cellular components including BMECs, astrocytic

endfeet, pericytes, as well as tight junction protein complexes. The demonstrated gentility

and simplicity of the approach, in turn, confer versatility in the high-throughput techniques

that can be successfully utilized downstream to microvessel isolation, as demonstrated

here with RNA sequencing and LC-MS/MS.

This protocol should have excellent reproducibility in isolating intact microvessels from

any region of the adult human brain. It should be noted, however, that the current version

of the protocol may need some adjustments if myelin content of brain samples used is

high. As reported, microvessels isolated using the described method are in high yield in

the absence of other contaminating cell types. However, limited expression of neuronal

markers is observed during computational deconvolution of RNA sequencing data.

Whether a minority of neurons are co-enriched with microvessels remains unclear, as co-

enrichment of neurons was not observed during immunophenotypic characterization of

isolated microvessels. While some co-enrichment is consistent with the technically

unavoidable capture of some parenchymal cells when utilizing a method suitable for

frozen postmortem brain tissue (which necessitates as few and as gentle steps as

possible), the human BBB atlases generated by Yang et al. (2022) and Garcia et al.,

(2022) reveal that "canonically neuronal" genes are also expressed in vascular-

associated cell types. Therefore, it is possible that counts for such genes indeed originate

from vascular-associated cells and not neurons. Another limitation is that the protocol

enriches for but does not purify microvessels, so a limited proportion of contamination

from non-neurovascular cell types might be found. Additionally, representation of all

vascular-associated cell types might not be uniform, as different cell types may vary

across brain regions and even brain subregions, and different cell types may be

differentially susceptible to steps taken during RNA or protein extraction. However, it is

reasonable to assume that the proportions of vascular-associated cell types found within

isolated microvessels represent the natural multi-cellular composition of vasculature

within the brain. Inevitably, studies that make use of isolated human microvessels will

encounter high inter-subject variation in both RNA and protein yield as well as

experimental read-out. Large cohorts with matched subjects (if studying disease and/or

sex differences) would be required to power a study investigating how brain vascular gene

or proteomic expression varies with factors such as brain region, age, or disease.

Although sensitivity of mass spectrometers has been greatly improved over the

years[170,171,172,173,174], as well as peptide separation for untargeted proteome

analysis[172,173,175], the literature consistently shows that high output from mass

spectrometry relies on a fine balance between sample quantity as well as sample

complexity. Because protein extracted from a structural unit, such as microvessels, as

opposed to bulk tissue could be considered a sample lacking complexity, it is reasonable

to assume that this relays to lower output, which may be further impacted by several low

signal-to-noise events that result in unidentified peptides[176] and a more limited dynamic

range of peptide detection[174]. An example of this are low-molecular weight cytokines

which are released in diluted concentrations by virtue of their high biological activity[177].

Our successfully generated datasets promise that future datasets generated from

microvessels isolated by the described method, when compared to those isolated from

mice, have the potential to improve our understanding of species-specific differences in

brain vascular gene expression, which is key to assess potential limitations of mouse

models for brain vascular and BBB development and disease. Species differences in

disease-affected neuronal and myeloid gene expression have been characterized by

single-cell or single-nucleus sequencing[178,179,180], but such differences in neurovascular

gene expression have only just begun to be comprehensively analyzed, and solely in

Alzheimer's disease and Huntington's disease[64,65]. Despite limited data, it has become

evident there are striking differences, with one study reporting 142 mouse-enriched

genes, including Vtn, Slco1c1, Slc6a20a, Atp13a5, Slc22a8, and 211 human-enriched

genes, including SLCO2A1, GIMAP7, and A2M[181]. Yang et al. (2022)[65] further reported

hundreds of species-enriched genes in BMECs and pericytes[65], finding that BECs and

pericytes exhibit the greatest transcriptional divergence in several vascular solute transporters (for e.g., GABA transporter SLC6A12) and genes of disease and pharmacological importance[65]. This observation was corroborated by Garcia et al. (2022) [64], who further detailed that species-specific DEGs were strongly enriched for marker genes of vascular-associated cell types, indicating that cell type identity markers were among those that vary the most between species[64].

In sum, the isolated brain microvessel is a robust model for the NVU and can be used to generate a variety of highly dimensional datasets. The availability of characterized human neurovascular transcriptomes and proteomes can aid investigators in identifying potential roles for BMECs and pericytes in the pathogenesis of neurological and psychiatric conditions, and should allow others to quickly assess the expression of molecules with potential relevance to drug delivery (e.g., SLCs, ABCs, and large molecule receptors) and disease in the human brain vasculature.

Acknowledgements

The authors have no competing interests to declare. This study was funded by an ERA NET Neuron grant and a CIHR Project grant to N.M. M.W. and R.R. respectively received scholarship and fellowship support from the FRQ-S. The Douglas-Bell Canada Brain Bank is supported in part by platform support grants from the Réseau Québécois sur le Suicide, les Troubles de l'Humeur et les Troubles Associés (FRQ-S), Healthy Brains for Healthy Lives (CFREF), and Brain Canada. The present study used the services of the Molecular and Cellular Microscopy Platform (MCMP) at the Douglas Research Centre.

The authors would like to thank the expert help of Douglas-Bell Canada Brain Bank staff

members (J. Prud'homme, M. Bouchouka, D. Mirault, V. Lariviere, A. Baccichet), the

technology development team at the McGill University and Genome Quebec Innovation

Centre, and the Segal Cancer Proteomics Centre at the Lady Davis Institute. The authors

would also like to thank Dr. Ghazal Fakhfouri for rearing and providing the mice used in

this study.

References

1. Iadecola, C. The Neurovascular Unit Coming of Age: A Journey through
Neurovascular Coupling in Health and Disease. *Neuron* **96**, 17-42 (2017).
https://doi.org:10.1016/j.neuron.2017.07.030
2. Kisler, K., Nelson, A. R., Montagne, A. & Zlokovic, B. V. Cerebral blood flow
regulation and neurovascular dysfunction in Alzheimer disease. *Nat Rev Neurosci*
18, 419-434 (2017). https://doi.org:10.1038/nrn.2017.48
3. Betz, A. L. & Goldstein, G. W. Polarity of the blood-brain barrier: neutral amino acid
transport into isolated brain capillaries. *Science* **202**, 225-227 (1978).
https://doi.org:10.1126/science.211586
4. Betz, A. L., Firth, J. A. & Goldstein, G. W. Polarity of the blood-brain barrier:
distribution of enzymes between the luminal and antiluminal membranes of brain
capillary endothelial cells. *Brain Res* **192**, 17-28 (1980).
https://doi.org:10.1016/0006-8993(80)91004-5
5. Cordon-Cardo, C. *et al.* Multidrug-resistance gene (P-glycoprotein) is expressed by
endothelial cells at blood-brain barrier sites. *Proc Natl Acad Sci U S A* **86**, 695-698
(1989). https://doi.org:10.1073/pnas.86.2.695
6. Thiebaut, F. *et al.* Immunohistochemical localization in normal tissues of different
epitopes in the multidrug transport protein P170: evidence for localization in brain
capillaries and crossreactivity of one antibody with a muscle protein. *J Histochem*
Cytochem **37**, 159-164 (1989). https://doi.org:10.1177/37.2.2463300
7. Loscher, W. & Potschka, H. Blood-brain barrier active efflux transporters: ATP-
binding cassette gene family. *NeuroRx* **2**, 86-98 (2005).
https://doi.org:10.1602/neurorx.2.1.86
8. Mittapalli, R. K., Manda, V. K., Adkins, C. E., Geldenhuys, W. J. & Lockman, P. R.
Exploiting nutrient transporters at the blood-brain barrier to improve brain distribution
of small molecules. *Ther Deliv* **1**, 775-784 (2010). https://doi.org:10.4155/tde.10.76
9. Zlokovic, B. V. The blood-brain barrier in health and chronic neurodegenerative
disorders. *Neuron* **57**, 178-201 (2008). https://doi.org:10.1016/j.neuron.2008.01.003

10. Daneman, R. The blood-brain barrier in health and disease. *Ann Neurol* **72**, 648-672
(2012). https://doi.org:10.1002/ana.23648
11. McConnell, H. L., Kersch, C. N., Woltjer, R. L. & Neuwelt, E. A. The Translational
Significance of the Neurovascular Unit. *J Biol Chem* **292**, 762-770 (2017).
https://doi.org:10.1074/jbc.R116.760215
12. Mintun, M. A. *et al.* Blood flow and oxygen delivery to human brain during functional
activity: theoretical modeling and experimental data. *Proc Natl Acad Sci U S A* **98**,
6859-6864 (2001). https://doi.org:10.1073/pnas.111164398
13. Hoge, R. D. *et al.* Linear coupling between cerebral blood flow and oxygen
consumption in activated human cortex. *Proc Natl Acad Sci U S A* **96**, 9403-9408
(1999). https://doi.org:10.1073/pnas.96.16.9403
14. Fox, P. T., Raichle, M. E., Mintun, M. A. & Dence, C. Nonoxidative glucose
consumption during focal physiologic neural activity. *Science* **241**, 462-464 (1988).
https://doi.org:10.1126/science.3260686
15. De Vivo, D. C. *et al.* Defective glucose transport across the blood-brain barrier as a
cause of persistent hypoglycorrhachia, seizures, and developmental delay. *N Engl J*
Med **325**, 703-709 (1991). https://doi.org:10.1056/NEJM199109053251006
16. Farrell, C. L. & Pardridge, W. M. Blood-brain barrier glucose transporter is
asymmetrically distributed on brain capillary endothelial lumenal and ablumenal
membranes: an electron microscopic immunogold study. *Proc Natl Acad Sci U S A*
88, 5779-5783 (1991). https://doi.org:10.1073/pnas.88.13.5779
17. Gerhart, D. Z., LeVasseur, R. J., Broderius, M. A. & Drewes, L. R. Glucose
transporter localization in brain using light and electron immunocytochemistry. *J*
Neurosci Res **22**, 464-472 (1989). https://doi.org:10.1002/jnr.490220413
18. Jeong, S. M. *et al.* Changes in magnesium concentration in the serum and
cerebrospinal fluid of neuropathic rats. *Acta Anaesthesiol Scand* **50**, 211-216 (2006).
https://doi.org:10.1111/j.1399-6576.2006.00925.x
19. Gendelman, H. E. *et al.* Monocyte chemotactic protein-1 regulates voltage-gated K+
channels and macrophage transmigration. *J Neuroimmune Pharmacol* **4**, 47-59
(2009). https://doi.org:10.1007/s11481-008-9135-1
20. Wilhelm, I. *et al.* Regulation of cerebral endothelial cell morphology by extracellular
calcium. *Phys Med Biol* **52**, 6261-6274 (2007). https://doi.org:10.1088/0031-
9155/52/20/012
21. Mokgokong, R., Wang, S., Taylor, C. J., Barrand, M. A. & Hladky, S. B. Ion
transporters in brain endothelial cells that contribute to formation of brain interstitial
fluid. *Pflugers Arch* **466**, 887-901 (2014). https://doi.org:10.1007/s00424-013-1342-9
22. Smith, Q. R. & Rapoport, S. I. Cerebrovascular permeability coefficients to sodium,
potassium, and chloride. *J Neurochem* **46**, 1732-1742 (1986).
https://doi.org:10.1111/j.1471-4159.1986.tb08491.x
23. Mathiesen, C., Caesar, K., Akgoren, N. & Lauritzen, M. Modification of activity-
dependent increases of cerebral blood flow by excitatory synaptic activity and spikes
in rat cerebellar cortex. *J Physiol* **512 (Pt 2)**, 555-566 (1998).
https://doi.org:10.1111/j.1469-7793.1998.555be.x
24. Caesar, K., Thomsen, K. & Lauritzen, M. Dissociation of spikes, synaptic activity,
and activity-dependent increments in rat cerebellar blood flow by tonic synaptic

inhibition. *Proc Natl Acad Sci U S A* **100**, 16000-16005 (2003).
https://doi.org:10.1073/pnas.2635195100

25. Iadecola, C. Regulation of the cerebral microcirculation during neural activity: is nitric oxide the missing link? *Trends Neurosci* **16**, 206-214 (1993). https://doi.org:10.1016/0166-2236(93)90156-g

26. Roy, C. S. & Sherrington, C. S. On the Regulation of the Blood-supply of the Brain. *J Physiol* **11**, 85-158 117 (1890). https://doi.org:10.1113/jphysiol.1890.sp000321

27. Fergus, A. & Lee, K. S. GABAergic regulation of cerebral microvascular tone in the rat. *J Cereb Blood Flow Metab* **17**, 992-1003 (1997). https://doi.org:10.1097/00004647-199709000-00009

28. Verheggen, I. C. M., Van Boxtel, M. P. J., Verhey, F. R. J., Jansen, J. F. A. & Backes, W. H. Interaction between blood-brain barrier and glymphatic system in solute clearance. *Neurosci Biobehav Rev* **90**, 26-33 (2018). https://doi.org:10.1016/j.neubiorev.2018.03.028

29. Deane, R., Bell, R. D., Sagare, A. & Zlokovic, B. V. Clearance of amyloid-beta peptide across the blood-brain barrier: implication for therapies in Alzheimer's disease. *CNS Neurol Disord Drug Targets* **8**, 16-30 (2009). https://doi.org:10.2174/187152709787601867

30. Iliff, J. J. *et al.* A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med* **4**, 147ra111 (2012). https://doi.org:10.1126/scitranslmed.3003748

31. Iliff, J. J. *et al.* Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J Neurosci* **33**, 18190-18199 (2013). https://doi.org:10.1523/JNEUROSCI.1592-13.2013

32. Iturria-Medina, Y. *et al.* Early role of vascular dysregulation on late-onset Alzheimer's disease based on multifactorial data-driven analysis. *Nat Commun* **7**, 11934 (2016). https://doi.org:10.1038/ncomms11934

33. Montagne, A. *et al.* Blood-brain barrier breakdown in the aging human hippocampus. *Neuron* **85**, 296-302 (2015). https://doi.org:10.1016/j.neuron.2014.12.032

34. Sweeney, M. D., Sagare, A. P. & Zlokovic, B. V. Cerebrospinal fluid biomarkers of neurovascular dysfunction in mild dementia and Alzheimer's disease. *J Cereb Blood Flow Metab* **35**, 1055-1068 (2015). https://doi.org:10.1038/jcbfm.2015.76

35. Arvanitakis, Z., Capuano, A. W., Leurgans, S. E., Bennett, D. A. & Schneider, J. A. Relation of cerebral vessel disease to Alzheimer's disease dementia and cognitive function in elderly people: a cross-sectional study. *Lancet Neurol* **15**, 934-943 (2016). https://doi.org:10.1016/S1474-4422(16)30029-1

36. Toledo, J. B. *et al.* Contribution of cerebrovascular disease in autopsy confirmed neurodegenerative disease cases in the National Alzheimer's Coordinating Centre. *Brain* **136**, 2697-2706 (2013). https://doi.org:10.1093/brain/awt188

37. Rosenberg, G. A. Blood-Brain Barrier Permeability in Aging and Alzheimer's Disease. *J Prev Alzheimers Dis* **1**, 138-139 (2014). https://doi.org:10.14283/jpad.2014.25

38. Stein, S. C., Chen, X. H., Sinson, G. P. & Smith, D. H. Intravascular coagulation: a major secondary insult in nonfatal traumatic brain injury. *J Neurosurg* **97**, 1373-1377 (2002). https://doi.org:10.3171/jns.2002.97.6.1373

39. Schwarzmaier, S. M., Kim, S. W., Trabold, R. & Plesnila, N. Temporal profile of
thrombogenesis in the cerebral microcirculation after traumatic brain injury in mice. *J*
Neurotrauma **27**, 121-130 (2010). https://doi.org:10.1089/neu.2009.1114
40. Dietrich, W. D., Alonso, O. & Halley, M. Early microvascular and neuronal
consequences of traumatic brain injury: a light and electron microscopic study in
rats. *J Neurotrauma* **11**, 289-301 (1994). https://doi.org:10.1089/neu.1994.11.289
41. del Zoppo, G. J. & Mabuchi, T. Cerebral microvessel responses to focal ischemia. *J*
Cereb Blood Flow Metab **23**, 879-894 (2003).
https://doi.org:10.1097/01.WCB.0000078322.96027.78
42. Schroder, M. L., Muizelaar, J. P., Fatouros, P. P., Kuta, A. J. & Choi, S. C. Regional
cerebral blood volume after severe head injury in patients with regional cerebral
ischemia. *Neurosurgery* **42**, 1276-1280; discussion 1280-1271 (1998).
https://doi.org:10.1097/00006123-199806000-00042
43. von Oettingen, G., Bergholt, B., Gyldensted, C. & Astrup, J. Blood flow and ischemia
within traumatic cerebral contusions. *Neurosurgery* **50**, 781-788; discussion 788-790
(2002). https://doi.org:10.1097/00006123-200204000-00019
44. Shapira, Y., Setton, D., Artru, A. A. & Shohami, E. Blood-brain barrier permeability,
cerebral edema, and neurologic function after closed head injury in rats. *Anesth*
Analg **77**, 141-148 (1993). https://doi.org:10.1213/00000539-199307000-00028
45. Baldwin, S. A., Fugaccia, I., Brown, D. R., Brown, L. V. & Scheff, S. W. Blood-brain
barrier breach following cortical contusion in the rat. *J Neurosurg* **85**, 476-481
(1996). https://doi.org:10.3171/jns.1996.85.3.0476
46. Hicks, R. R., Baldwin, S. A. & Scheff, S. W. Serum extravasation and cytoskeletal
alterations following traumatic brain injury in rats. Comparison of lateral fluid
percussion and cortical impact models. *Mol Chem Neuropathol* **32**, 1-16 (1997).
https://doi.org:10.1007/BF02815164
47. Baskaya, M. K., Rao, A. M., Dogan, A., Donaldson, D. & Dempsey, R. J. The
biphasic opening of the blood-brain barrier in the cortex and hippocampus after
traumatic brain injury in rats. *Neurosci Lett* **226**, 33-36 (1997).
https://doi.org:10.1016/s0304-3940(97)00239-5
48. Gaitan, M. I., de Alwis, M. P., Sati, P., Nair, G. & Reich, D. S. Multiple sclerosis
shrinks intralesional, and enlarges extralesional, brain parenchymal veins.
Neurology **80**, 145-151 (2013). https://doi.org:10.1212/WNL.0b013e31827b916f
49. Buch, S. *et al.* Revealing vascular abnormalities and measuring small vessel density
in multiple sclerosis lesions using USPIO. *Neuroimage Clin* **29**, 102525 (2021).
https://doi.org:10.1016/j.nicl.2020.102525
50. Geraldes, R. *et al.* Vascular disease and multiple sclerosis: a post-mortem study
exploring their relationships. *Brain* **143**, 2998-3012 (2020).
https://doi.org:10.1093/brain/awaa255
51. Al-Louzi, O. *et al.* Central Vein Sign Profile of Newly Developing Lesions in Multiple
Sclerosis: A 3-Year Longitudinal Study. *Neurol Neuroimmunol Neuroinflamm* **9**
(2022). https://doi.org:10.1212/NXI.0000000000001120
52. Khan, O. *et al.* Chronic cerebrospinal venous insufficiency and multiple sclerosis.
Ann Neurol **67**, 286-290 (2010). https://doi.org:10.1002/ana.22001

53. Doepp, F. *et al.* Venous drainage in multiple sclerosis: a combined MRI and
ultrasound study. *Neurology* **77**, 1745-1751 (2011).
https://doi.org:10.1212/WNL.0b013e318236f0ea
54. Liu, P. *et al.* Time-course investigation of blood-brain barrier permeability and tight
junction protein changes in a rat model of permanent focal ischemia. *J Physiol Sci*
68, 121-127 (2018). https://doi.org:10.1007/s12576-016-0516-6
55. Lin, C. Y. *et al.* Dynamic changes in vascular permeability, cerebral blood volume,
vascular density, and size after transient focal cerebral ischemia in rats: evaluation
with contrast-enhanced magnetic resonance imaging. *J Cereb Blood Flow Metab* **28**,
1491-1501 (2008). https://doi.org:10.1038/jcbfm.2008.42
56. Strbian, D. *et al.* The blood-brain barrier is continuously open for several weeks
following transient focal cerebral ischemia. *Neuroscience* **153**, 175-181 (2008).
https://doi.org:10.1016/j.neuroscience.2008.02.012
57. Durukan, A. *et al.* Post-ischemic blood-brain barrier leakage in rats: one-week
follow-up by MRI. *Brain Res* **1280**, 158-165 (2009).
https://doi.org:10.1016/j.brainres.2009.05.025
58. Pillai, D. R. *et al.* Cerebral ischemia-reperfusion injury in rats--a 3 T MRI study on
biphasic blood-brain barrier opening and the dynamics of edema formation. *J Cereb*
Blood Flow Metab **29**, 1846-1855 (2009). https://doi.org:10.1038/jcbfm.2009.106
59. Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain
vasculature. *Nature* **554**, 475-480 (2018). https://doi.org:10.1038/s41586-018-0232-x
60. He, L. *et al.* Single-cell RNA sequencing of mouse brain and lung vascular and
vessel-associated cell types. *Sci Data* **5**, 180160 (2018).
https://doi.org:10.1038/sdata.2018.160
61. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia,
neurons, and vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929-11947
(2014). https://doi.org:10.1523/JNEUROSCI.1860-14.2014
62. He, L. *et al.* Analysis of the brain mural cell transcriptome. *Sci Rep* **6**, 35108 (2016).
https://doi.org:10.1038/srep35108
63. Munji, R. N. *et al.* Profiling the mouse brain endothelial transcriptome in health and
disease models reveals a core blood-brain barrier dysfunction module. *Nat Neurosci*
22, 1892-1902 (2019). https://doi.org:10.1038/s41593-019-0497-x
64. [dataset] Garcia, F. J. *et al.* Single-cell dissection of the human brain vasculature.
Nature **603**, 893-899 (2022). https://doi.org:10.1038/s41586-022-04521-7
65. [dataset] Yang, A. C. *et al.* A human brain vascular atlas reveals diverse mediators
of Alzheimer's risk. *Nature* **603**, 885-892 (2022). https://doi.org:10.1038/s41586-021-
04369-3
66. Klein, B., Kuschinsky, W., Schrock, H. & Vetterlein, F. Interdependency of local
capillary density, blood flow, and metabolism in rat brains. *Am J Physiol* **251**, H1333-
1340 (1986). https://doi.org:10.1152/ajpheart.1986.251.6.H1333
67. Wu, E. X., Tang, H. & Jensen, J. H. High-resolution MR imaging of mouse brain
microvasculature using the relaxation rate shift index Q. *NMR Biomed* **17**, 507-512
(2004). https://doi.org:10.1002/nbm.921
68. Weber, B., Keller, A. L., Reichold, J. & Logothetis, N. K. The microvascular system
of the striate and extrastriate visual cortex of the macaque. *Cereb Cortex* **18**, 2318-
2330 (2008). https://doi.org:10.1093/cercor/bhm259

69. Ventura-Antunes, L., Dasgupta, O. M. & Herculano-Houzel, S. Resting Rates of
Blood Flow and Glucose Use per Neuron Are Proportional to Number of Endothelial
Cells Available per Neuron Across Sites in the Rat Brain. *Front Integr Neurosci* **16**,
821850 (2022). https://doi.org:10.3389/fnint.2022.821850
70. Nagy, C. *et al.* Single-nucleus transcriptomics of the prefrontal cortex in major
depressive disorder implicates oligodendrocyte precursor cells and excitatory
neurons. *Nat Neurosci* **23**, 771-781 (2020). https://doi.org:10.1038/s41593-020-
0621-y
71. [dataset] Velmeshev, D. *et al.* Single-cell genomics identifies cell type-specific
molecular changes in autism. *Science* **364**, 685-689 (2019).
https://doi.org:10.1126/science.aav8130
72. Mathys, H. *et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature*
570, 332-337 (2019). https://doi.org:10.1038/s41586-019-1195-2
73. Grubman, A. *et al.* A single-cell atlas of entorhinal cortex from individuals with
Alzheimer's disease reveals cell-type-specific gene expression regulation. *Nat*
Neurosci **22**, 2087-2097 (2019). https://doi.org:10.1038/s41593-019-0539-4
74. Jakel, S. *et al.* Altered human oligodendrocyte heterogeneity in multiple sclerosis.
Nature **566**, 543-547 (2019). https://doi.org:10.1038/s41586-019-0903-2
75. Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and technical
variability in single-cell RNA-sequencing experiments. *Biostatistics* **19**, 562-578
(2018). https://doi.org:10.1093/biostatistics/kxx053
76. Bacher, R. & Kendziorski, C. Design and computational analysis of single-cell RNA-
sequencing experiments. *Genome Biol* **17**, 63 (2016).
https://doi.org:10.1186/s13059-016-0927-y
77. Denisenko, E. *et al.* Systematic assessment of tissue dissociation and storage
biases in single-cell and single-nucleus RNA-seq workflows. *Genome Biol* **21**, 130
(2020). https://doi.org:10.1186/s13059-020-02048-6
78. Crouch, E. E. & Doetsch, F. FACS isolation of endothelial cells and pericytes from
mouse brain microregions. *Nat Protoc* **13**, 738-751 (2018).
https://doi.org:10.1038/nprot.2017.158
79. Navone, S. E. *et al.* Isolation and expansion of human and mouse brain
microvascular endothelial cells. *Nat Protoc* **8**, 1680-1693 (2013).
https://doi.org:10.1038/nprot.2013.107
80. van Beijnum, J. R., Rousch, M., Castermans, K., van der Linden, E. & Griffioen, A.
W. Isolation of endothelial cells from fresh tissues. *Nat Protoc* **3**, 1085-1091 (2008).
https://doi.org:10.1038/nprot.2008.71
81. Kinnecom, K. & Pachter, J. S. Selective capture of endothelial and perivascular cells
from brain microvessels using laser capture microdissection. *Brain Res Brain Res*
Protoc **16**, 1-9 (2005). https://doi.org:10.1016/j.brainresprot.2005.08.002
82. Mojsilovic-Petrovic, J., Nesic, M., Pen, A., Zhang, W. & Stanimirovic, D.
Development of rapid staining protocols for laser-capture microdissection of brain
vessels from human and rat coupled to gene expression analyses. *J Neurosci*
Methods **133**, 39-48 (2004). https://doi.org:10.1016/j.jneumeth.2003.09.026
83. Harris, L. W. *et al.* The cerebral microvasculature in schizophrenia: a laser capture
microdissection study. *PLoS One* **3**, e3964 (2008).
https://doi.org:10.1371/journal.pone.0003964

84. Song, H. W. *et al.* Transcriptomic comparison of human and mouse brain microvessels. *Sci Rep* **10**, 12358 (2020). https://doi.org:10.1038/s41598-020-69096-7

85. Almeida, D. & Turecki, G. Profiling cell-type specific gene expression in post-mortem human brain samples through laser capture microdissection. *Methods* **207**, 3-10 (2022). https://doi.org:10.1016/j.ymeth.2022.08.013

86. Bourassa, P., Tremblay, C., Schneider, J. A., Bennett, D. A. & Calon, F. Beta-amyloid pathology in human brain microvessel extracts from the parietal cortex: relation with cerebral amyloid angiopathy and Alzheimer's disease. *Acta Neuropathol* **137**, 801-823 (2019). https://doi.org:10.1007/s00401-019-01967-4

87. Lee, Y. K., Uchida, H., Smith, H., Ito, A. & Sanchez, T. The isolation and molecular characterization of cerebral microvessels. *Nat Protoc* **14**, 3059-3081 (2019). https://doi.org:10.1038/s41596-019-0212-0

88. Spitzer, D. *et al.* A flow cytometry-based protocol for syngenic isolation of neurovascular unit cells from mouse and human tissues. *Nat Protoc* (2023). https://doi.org:10.1038/s41596-023-00805-y

89. Isingrini, E. *et al.* Genetic elimination of dopamine vesicular stocks in the nigrostriatal pathway replicates Parkinson's disease motor symptoms without neuronal degeneration in adult mice. *Sci Rep* **7**, 12432 (2017). https://doi.org:10.1038/s41598-017-12810-9

90. Shimizu, N. Histochemical studies on the phosphatase of the nervous system. *J Comp Neurol* **93**, 201-217 (1950). https://doi.org:10.1002/cne.900930203

91. Leduc, E. H. & Wislocki, G. B. The histochemical localization of acid and alkaline phosphatases, non-specific esterase and succinic dehydrogenase in the structures comprising the hemato-encephalic barrier of the rat. *J Comp Neurol* **97**, 241-279 (1952). https://doi.org:10.1002/cne.900970203

92. Bourne, G. H. Histochemical demonstration of phosphatases in the central nervous system of the rat. *Exp Cell Res* **14**, 101-117 (1958).

93. Becker, N. H., Goldfischer, S., Shin, W. Y. & Novikoff, A. B. The localization of enzyme activities in the rat brain. *J Biophys Biochem Cytol* **8**, 649-663 (1960). https://doi.org:10.1083/jcb.8.3.649

94. Bannister, R. G. & Romanul, F. C. The Localization of Alkaline Phosphatase Activity in Cerebral Blood Vessels. *J Neurol Neurosurg Psychiatry* **26**, 333-340 (1963). https://doi.org:10.1136/jnnp.26.4.333

95. Romanul, F. C. & Bannister, R. G. Localized areas of high alkaline phosphatase activity in the terminal arterial tree. *J Cell Biol* **15**, 73-84 (1962). https://doi.org:10.1083/jcb.15.1.73

96. Ball, H. J., McParland, B., Driussi, C. & Hunt, N. H. Isolating vessels from the mouse brain for gene expression analysis using laser capture microdissection. *Brain Res Brain Res Protoc* **9**, 206-213 (2002). https://doi.org:10.1016/s1385-299x(02)00147-2

97. Nikolova-Krstevski, V. *et al.* ERG is required for the differentiation of embryonic stem cells along the endothelial lineage. *BMC Dev Biol* **9**, 72 (2009). https://doi.org:10.1186/1471-213X-9-72

98. Miettinen, M. *et al.* ERG transcription factor as an immunohistochemical marker for vascular endothelial tumors and prostatic carcinoma. *Am J Surg Pathol* **35**, 432-441 (2011). https://doi.org:10.1097/PAS.0b013e318206b67b

99. Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**, 491-499 (2017). https://doi.org:10.1101/gr.209601.116

100. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013). https://doi.org:10.1093/bioinformatics/bts635

101. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009). https://doi.org:10.1093/bioinformatics/btp352

102. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011). https://doi.org:10.1186/1471-2105-12-323

103. Sutton, G. J. *et al.* Comprehensive evaluation of deconvolution methods for human brain gene expression. *Nat Commun* **13**, 1358 (2022). https://doi.org:10.1038/s41467-022-28655-4

104. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019). https://doi.org:10.1016/j.cell.2019.05.031

105. Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues with digital cytometry. *Nat Biotechnol* **37**, 773-782 (2019). https://doi.org:10.1038/s41587-019-0114-2

106. Chang, L. & Goldman, R. D. Intermediate filaments mediate cytoskeletal crosstalk. *Nat Rev Mol Cell Biol* **5**, 601-613 (2004). https://doi.org:10.1038/nrm1438

107. Aumailley, M. & Smyth, N. The role of laminins in basement membrane function. *J Anat* **193 (Pt 1)**, 1-21 (1998). https://doi.org:10.1046/j.1469-7580.1998.19310001.x

108. Greene, C., Hanley, N. & Campbell, M. Claudin-5: gatekeeper of neurological function. *Fluids Barriers CNS* **16**, 3 (2019). https://doi.org:10.1186/s12987-019-0123-z

109. Winkler, E. A., Bell, R. D. & Zlokovic, B. V. Pericyte-specific expression of PDGF beta receptor in mouse models with normal and deficient PDGF beta receptor signaling. *Mol Neurodegener* **5**, 32 (2010). https://doi.org:10.1186/1750-1326-5-32

110. Kimelberg, H. K. & Nedergaard, M. Functions of astrocytes and their potential as therapeutic targets. *Neurotherapeutics* **7**, 338-353 (2010). https://doi.org:10.1016/j.nurt.2010.07.006

111. Mathiisen, T. M., Lehre, K. P., Danbolt, N. C. & Ottersen, O. P. The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction. *Glia* **58**, 1094-1103 (2010). https://doi.org:10.1002/glia.20990

112. Winkler, E. A., Bell, R. D. & Zlokovic, B. V. Central nervous system pericytes in health and disease. *Nat Neurosci* **14**, 1398-1405 (2011). https://doi.org:10.1038/nn.2946

113. Abbott, N. J., Patabendige, A. A., Dolman, D. E., Yusof, S. R. & Begley, D. J. Structure and function of the blood-brain barrier. *Neurobiol Dis* **37**, 13-25 (2010). https://doi.org:10.1016/j.nbd.2009.07.030

114. Abbott, N. J., Ronnback, L. & Hansson, E. Astrocyte-endothelial interactions at the blood-brain barrier. *Nat Rev Neurosci* **7**, 41-53 (2006). https://doi.org:10.1038/nrn1824

115. Wolburg, H., Noell, S., Mack, A., Wolburg-Buchholz, K. & Fallier-Becker, P. Brain
endothelial cells and the glio-vascular complex. *Cell Tissue Res* **335**, 75-96 (2009).
https://doi.org:10.1007/s00441-008-0658-9
116. Nagelhus, E. A. & Ottersen, O. P. Physiological roles of aquaporin-4 in brain.
Physiol Rev **93**, 1543-1562 (2013). https://doi.org:10.1152/physrev.00011.2013
117. Papadopoulos, M. C. & Verkman, A. S. Aquaporin water channels in the nervous
system. *Nat Rev Neurosci* **14**, 265-277 (2013). https://doi.org:10.1038/nrn3468
118. Chasseigneaux, S. *et al.* Isolation and differential transcriptome of vascular
smooth muscle cells and mid-capillary pericytes from the rat brain. *Sci Rep* **8**, 12272
(2018). https://doi.org:10.1038/s41598-018-30739-5
119. Muhl, L. *et al.* Single-cell analysis uncovers fibroblast heterogeneity and criteria for
fibroblast and mural cell identification and discrimination. *Nat Commun* **11**, 3953
(2020). https://doi.org:10.1038/s41467-020-17740-1
120. Sweeney, M. D., Ayyadurai, S. & Zlokovic, B. V. Pericytes of the neurovascular
unit: key functions and signaling pathways. *Nat Neurosci* **19**, 771-783 (2016).
https://doi.org:10.1038/nn.4288
121. Gonzales, A. L. *et al.* Contractile pericytes determine the direction of blood flow at
capillary junctions. *Proc Natl Acad Sci U S A* **117**, 27022-27033 (2020).
https://doi.org:10.1073/pnas.1922755117
122. Hartmann, D. A. *et al.* Brain capillary pericytes exert a substantial but slow
influence on blood flow. *Nat Neurosci* **24**, 633-645 (2021).
https://doi.org:10.1038/s41593-020-00793-2
123. Alarcon-Martinez, L. *et al.* Capillary pericytes express alpha-smooth muscle actin,
which requires prevention of filamentous-actin depolymerization for detection. *Elife* **7**
(2018). https://doi.org:10.7554/eLife.34861
124. Boulay, A. C. *et al.* Translation in astrocyte distal processes sets molecular
heterogeneity at the gliovascular interface. *Cell Discov* **3**, 17005 (2017).
https://doi.org:10.1038/celldisc.2017.5
125. Derouiche, A. & Geiger, K. D. Perspectives for Ezrin and Radixin in Astrocytes:
Kinases, Functions and Pathology. *Int J Mol Sci* **20** (2019).
https://doi.org:10.3390/ijms20153776
126. Sakers, K. *et al.* Astrocytes locally translate transcripts in their peripheral
processes. *Proc Natl Acad Sci U S A* **114**, E3830-E3838 (2017).
https://doi.org:10.1073/pnas.1617782114
127. Menard, C. *et al.* Social stress induces neurovascular pathology promoting
depression. *Nat Neurosci* **20**, 1752-1760 (2017). https://doi.org:10.1038/s41593-
017-0010-3
128. Dudek, K. A. *et al.* Molecular adaptations of the blood-brain barrier promote stress
resilience vs. depression. *Proc Natl Acad Sci U S A* **117**, 3326-3336 (2020).
https://doi.org:10.1073/pnas.1914655117
129. Puvogel, S. *et al.* Single-nucleus RNA sequencing of midbrain blood-brain barrier
cells in schizophrenia reveals subtle transcriptional changes with overall
preservation of cellular proportions and phenotypes. *Mol Psychiatry* **27**, 4731-4740
(2022). https://doi.org:10.1038/s41380-022-01796-0
130. Lau, S. F., Cao, H., Fu, A. K. Y. & Ip, N. Y. Single-nucleus transcriptome analysis
reveals dysregulation of angiogenic endothelial cells and neuroprotective glia in

Alzheimer's disease. *Proc Natl Acad Sci U S A* **117**, 25800-25809 (2020).
https://doi.org:10.1073/pnas.2008762117
131. Chinnery, P. F. *et al.* Clinical features and natural history of neuroferritinopathy
caused by the FTL1 460InsA mutation. *Brain* **130**, 110-119 (2007).
https://doi.org:10.1093/brain/awl319
132. Tadic, V. *et al.* Primary familial brain calcification with known gene mutations: a
systematic review and challenges of phenotypic characterization. *JAMA Neurol* **72**,
460-467 (2015). https://doi.org:10.1001/jamaneurol.2014.3889
133. Mitroi, D. N., Tian, M., Kawaguchi, R., Lowry, W. E. & Carmichael, S. T. Single-
nucleus transcriptome analysis reveals disease- and regeneration-associated
endothelial cells in white matter vascular dementia. *J Cell Mol Med* **26**, 3183-3195
(2022). https://doi.org:10.1111/jcmm.17315
134. Nie, L., Wu, G. & Zhang, W. Correlation of mRNA expression and protein
abundance affected by multiple sequence features related to translational efficiency
in Desulfovibrio vulgaris: a quantitative analysis. *Genetics* **174**, 2229-2243 (2006).
https://doi.org:10.1534/genetics.106.065862
135. Vogel, C. & Marcotte, E. M. Insights into the regulation of protein abundance from
proteomic and transcriptomic analyses. *Nat Rev Genet* **13**, 227-232 (2012).
https://doi.org:10.1038/nrg3185
136. Hediger, M. A. *et al.* The ABCs of solute carriers: physiological, pathological and
therapeutic implications of human membrane transport proteins. *Pflugers Arch* **447**,
465-468 (2004). https://doi.org:10.1007/s00424-003-1192-y
137. Ballabh, P., Braun, A. & Nedergaard, M. The blood-brain barrier: an overview:
structure, regulation, and clinical implications. *Neurobiol Dis* **16**, 1-13 (2004).
https://doi.org:10.1016/j.nbd.2003.12.016
138. Mueckler, M. & Thorens, B. The SLC2 (GLUT) family of membrane transporters.
Mol Aspects Med **34**, 121-138 (2013). https://doi.org:10.1016/j.mam.2012.07.001
139. Yanagida, O. *et al.* Human L-type amino acid transporter 1 (LAT1):
characterization of function and expression in tumor cell lines. *Biochim Biophys Acta*
1514, 291-302 (2001). https://doi.org:10.1016/s0005-2736(01)00384-4
140. Nakamura, E. *et al.* 4F2 (CD98) heavy chain is associated covalently with an
amino acid transporter and controls intracellular trafficking and membrane topology
of 4F2 heterodimer. *J Biol Chem* **274**, 3009-3016 (1999).
https://doi.org:10.1074/jbc.274.5.3009
141. Nicklin, P. *et al.* Bidirectional transport of amino acids regulates mTOR and
autophagy. *Cell* **136**, 521-534 (2009). https://doi.org:10.1016/j.cell.2008.11.044
142. Vijay, N. & Morris, M. E. Role of monocarboxylate transporters in drug delivery to
the brain. *Curr Pharm Des* **20**, 1487-1498 (2014).
https://doi.org:10.2174/13816128113199990462
143. Trajkovic, M. *et al.* Abnormal thyroid hormone metabolism in mice lacking the
monocarboxylate transporter 8. *J Clin Invest* **117**, 627-635 (2007).
https://doi.org:10.1172/JCI28253
144. Begley, D. J. ABC transporters and the blood-brain barrier. *Curr Pharm Des* **10**,
1295-1312 (2004). https://doi.org:10.2174/1381612043384844

145. Loscher, W. & Potschka, H. Blood-brain barrier active efflux transporters: ATP-
binding cassette gene family. *NeuroRx* **2**, 86-98 (2005)
https://doi.org:10.1602/neurorx.2.1.86
146. Sun, H., Dai, H., Shaik, N. & Elmquist, W. F. Drug efflux transporters in the CNS.
Adv Drug Deliv Rev **55**, 83-105 (2003). https://doi.org:10.1016/s0169-
409x(02)00172-2
147. Agarwal, S., Hartz, A. M., Elmquist, W. F. & Bauer, B. Breast cancer resistance
protein and P-glycoprotein in brain cancer: two gatekeepers team up. *Curr Pharm
Des* **17**, 2793-2802 (2011). https://doi.org:10.2174/138161211797440186
148. Agarwal, S., Sane, R., Gallardo, J. L., Ohlfest, J. R. & Elmquist, W. F. Distribution
of gefitinib to the brain is limited by P-glycoprotein (ABCB1) and breast cancer
resistance protein (ABCG2)-mediated active efflux. *J Pharmacol Exp Ther* **334**, 147-
155 (2010). https://doi.org:10.1124/jpet.110.167601
149. Agarwal, S., Sane, R., Ohlfest, J. R. & Elmquist, W. F. The role of the breast
cancer resistance protein (ABCG2) in the distribution of sorafenib to the brain. *J
Pharmacol Exp Ther* **336**, 223-233 (2011). https://doi.org:10.1124/jpet.110.175034
150. Chen, Y. *et al.* P-glycoprotein and breast cancer resistance protein influence brain
distribution of dasatinib. *J Pharmacol Exp Ther* **330**, 956-963 (2009).
https://doi.org:10.1124/jpet.109.154781
151. de Vries, N. A. *et al.* P-glycoprotein and breast cancer resistance protein: two
dominant transporters working together in limiting the brain penetration of topotecan.
Clin Cancer Res **13**, 6440-6449 (2007). https://doi.org:10.1158/1078-0432.CCR-07-
1335
152. Polli, J. W. *et al.* An unexpected synergist role of P-glycoprotein and breast cancer
resistance protein on the central nervous system penetration of the tyrosine kinase
inhibitor lapatinib (N-3-chloro-4-[(3-fluorobenzyl)oxy]phenyl-6-[5-([2-
(methylsulfonyl)ethyl]aminomethyl)-2-furyl]-4-quinazolinamine; GW572016). *Drug
Metab Dispos* **37**, 439-442 (2009). https://doi.org:10.1124/dmd.108.024646
153. Hindle, S. J. *et al.* Evolutionarily Conserved Roles for Blood-Brain Barrier
Xenobiotic Transporters in Endogenous Steroid Partitioning and Behavior. *Cell Rep*
21, 1304-1316 (2017). https://doi.org:10.1016/j.celrep.2017.10.026
154. Haber, M. A. *et al.* ERG is a novel and reliable marker for endothelial cells in
central nervous system tumors. *Clin Neuropathol* **34**, 117-127 (2015).
https://doi.org:10.5414/NP300817
155. Kim, S. *et al.* ERG Immunohistochemistry as an Endothelial Marker for Assessing
Lymphovascular Invasion. *Korean J Pathol* **47**, 355-364 (2013).
https://doi.org:10.4132/KoreanJPathol.2013.47.4.355
156. Birdsey, G. M. *et al.* The endothelial transcription factor ERG promotes vascular
stability and growth through Wnt/beta-catenin signaling. *Dev Cell* **32**, 82-96 (2015).
https://doi.org:10.1016/j.devcel.2014.11.016
157. Pastrana, E., Cheng, L. C. & Doetsch, F. Simultaneous prospective purification of
adult subventricular zone neural stem cells and their progeny. *Proc Natl Acad Sci U
S A* **106**, 6387-6392 (2009). https://doi.org:10.1073/pnas.0810407106
158. Codega, P. *et al.* Prospective identification and purification of quiescent adult
neural stem cells from their in vivo niche. *Neuron* **82**, 545-559 (2014).
https://doi.org:10.1016/j.neuron.2014.02.039

159. Plane, J. M., Andjelkovic, A. V., Keep, R. F. & Parent, J. M. Intact and injured
endothelial cells differentially modulate postnatal murine forebrain neural stem cells.
Neurobiol Dis **37**, 218-227 (2010). https://doi.org:10.1016/j.nbd.2009.10.008

160. Ohab, J. J., Fleming, S., Blesch, A. & Carmichael, S. T. A neurovascular niche for
neurogenesis after stroke. *J Neurosci* **26**, 13007-13016 (2006).
https://doi.org:10.1523/JNEUROSCI.4323-06.2006

161. Yamashita, T. *et al.* Subventricular zone-derived neuroblasts migrate and
differentiate into mature neurons in the post-stroke adult striatum. *J Neurosci* **26**,
6627-6636 (2006). https://doi.org:10.1523/JNEUROSCI.0149-06.2006

162. Kojima, T. *et al.* Subventricular zone-derived neural progenitor cells migrate along
a blood vessel scaffold toward the post-stroke striatum. *Stem Cells* **28**, 545-554
(2010). https://doi.org:10.1002/stem.306

163. Bardehle, S. *et al.* Live imaging of astrocyte responses to acute injury reveals
selective juxtavascular proliferation. *Nat Neurosci* **16**, 580-586 (2013).
https://doi.org:10.1038/nn.3371

164. Greenberg, D. A. Cerebral angiogenesis: a realistic therapy for ischemic disease?
Methods Mol Biol **1135**, 21-24 (2014). https://doi.org:10.1007/978-1-4939-0320-7_2

165. Paul, G. *et al.* The adult human brain harbors multipotent perivascular
mesenchymal stem cells. *PLoS One* **7**, e35577 (2012).
https://doi.org:10.1371/journal.pone.0035577

166. Durafourt, B. A., Moore, C. S., Blain, M. & Antel, J. P. Isolating, culturing, and
polarizing primary human adult and fetal microglia. *Methods Mol Biol* **1041**, 199-211
(2013). https://doi.org:10.1007/978-1-62703-520-0_19

167. De Groot, C. J. *et al.* Isolation and characterization of adult microglial cells and
oligodendrocytes derived from postmortem human brain tissue. *Brain Res Brain Res*
Protoc **5**, 85-94 (2000). https://doi.org:10.1016/s1385-299x(99)00059-8

168. Goldeman, C., Ozgur, B. & Brodin, B. Culture-induced changes in mRNA
expression levels of efflux and SLC-transporters in brain endothelial cells. *Fluids*
Barriers CNS **17**, 32 (2020). https://doi.org:10.1186/s12987-020-00193-5

169. Lyck, R. *et al.* Culture-induced changes in blood-brain barrier transcriptome:
implications for amino-acid transporters in vivo. *J Cereb Blood Flow Metab* **29**, 1491-
1502 (2009). https://doi.org:10.1038/jcbfm.2009.72

170. Hahne, H. *et al.* DMSO enhances electrospray response, boosting sensitivity of
proteomic experiments. *Nat Methods* **10**, 989-991 (2013).
https://doi.org:10.1038/nmeth.2610

171. Hebert, A. S. *et al.* Improved Precursor Characterization for Data-Dependent Mass
Spectrometry. *Anal Chem* **90**, 2333-2340 (2018).
https://doi.org:10.1021/acs.analchem.7b04808

172. Shishkova, E., Hebert, A. S. & Coon, J. J. Now, More Than Ever, Proteomics
Needs Better Chromatography. *Cell Syst* **3**, 321-324 (2016).
https://doi.org:10.1016/j.cels.2016.10.007

173. Shishkova, E., Hebert, A. S., Westphall, M. S. & Coon, J. J. Ultra-High Pressure
(>30,000 psi) Packing of Capillary Columns Enhancing Depth of Shotgun Proteomic
Analyses. *Anal Chem* **90**, 11503-11508 (2018).
https://doi.org:10.1021/acs.analchem.8b02766

174. Timp, W. & Timp, G. Beyond mass spectrometry, the next step in proteomics. *Sci
Adv* **6**, eaax8978 (2020). https://doi.org:10.1126/sciadv.aax8978
175. Toth, G., Panic-Jankovic, T. & Mitulovic, G. Pillar array columns for peptide
separations in nanoscale reversed-phase chromatography. *J Chromatogr A* **1603**,
426-432 (2019). https://doi.org:10.1016/j.chroma.2019.06.067
176. Griss, J. *et al.* Recognizing millions of consistently unidentified spectra across
hundreds of shotgun proteomics datasets. *Nat Methods* **13**, 651-656 (2016).
https://doi.org:10.1038/nmeth.3902
177. Angel, T. E. *et al.* Mass spectrometry-based proteomics: existing capabilities and
future directions. *Chem Soc Rev* **41**, 3912-3928 (2012).
https://doi.org:10.1039/c2cs15331a
178. Cosacak, M. I. *et al.* Single Cell/Nucleus Transcriptomics Comparison in Zebrafish
and Humans Reveals Common and Distinct Molecular Responses to Alzheimer's
Disease. *Cells* **11** (2022). https://doi.org:10.3390/cells11111807
179. Friedman, B. A. *et al.* Diverse Brain Myeloid Expression Profiles Reveal Distinct
Microglial Activation States and Aspects of Alzheimer's Disease Not Evident in
Mouse Models. *Cell Rep* **22**, 832-847 (2018).
https://doi.org:10.1016/j.celrep.2017.12.066
180. Kamath, T. *et al.* Single-cell genomic profiling of human dopamine neurons
identifies a population that selectively degenerates in Parkinson's disease. *Nat
Neurosci* **25**, 588-595 (2022). https://doi.org:10.1038/s41593-022-01061-1
181. Song, H. W. *et al.* Transcriptomic comparison of human and mouse brain
microvessels. *Sci Rep* **10**, 12358 (2020). https://doi.org:10.1038/s41598-020-69096-
7

Tables

Table 1: Subject demographics and covariates

Experiment type	Brain ID	Sex	Age (years)	Cause of death	PMI (hours)	Tissue pH
RNA seq	1	Female	76	Polytrauma (accident)	26.50	6.50
RNA seq	2	Female	51	Pulmonary embolism	111.32	6.5
RNA seq	3	Male	26	Polytrauma (car accident)	12	6.75
RNA seq	4	Female	28	Undetermined	80	6.45
RNA seq	5	Female	45	Pulmonary embolism	39.5	5.66
Mass spec	6	Male	45	Gun wound	20.50	6.57
Mass spec	7	Male	54	Cardiac arrest	25.25	6.61
Mass spec	8	Male	63	Fall from several metres	13.00	6.84

Background information on human subjects whose vmPFC tissue was studied. Information includes sex, age, cause of death, postmortem interval (PMI), and tissue pH.

Figures

Fig. 1: Schematic overview of microvessel isolation from brain tissue samples

Figure 1. Schematic overview of microvessel isolation. After the vmPFC is dissected from

coronal slice containing the frontal lobe, 100 mg of vmPFC tissue is more precisely

microdissected using a sterile blade or scalpel. Afterwards, the vmPFC tissue sample is dissociated in Homogenization Buffer using the GentleMACS™ Dissociator. Following dissociation, the sample is centrifuged at 3200 g for 30 mins in order to pellet dissociated microvessels. Next, the microvessel-enriched pellet is resuspended in PBS and filtered through a cellular strainer to trap microvessels within the mesh and to allow debris to flow through. The microvessel fragments on top of the mesh can then be subjected to total RNA extraction for downstream, NVU-specific transcriptomic characterization. Additional high-throughput applications can be applied using microvessel-enriched pellets, as discussed in this article. Image generated using BioRender.

Fig. 2: Overview of experimental steps taken to isolate, enrich and collect microvessels

Figure 2. a-g) Overview of experimental steps taken to collect and resuspend pelleted microvessels. a) 100 mg of vmPFC tissue submerged in 2 ml of Homogenization Buffer. b) 100 mg of vmPFC tissue after homogenization. c) Lysate topped-up to 10 ml with Homogenization Buffer. d) Lysate transferred to 15 ml falcon tube. e-f) After centrifugation at 3200 g for 30 min, a microvessel-enriched pellet forms at the bottom of the tube. g) microvessel—enriched pellet is resuspended in 500 ul of PBS. h) Example of BCIP/NBT-stained microvessel pellet, which lends a purple colour to the pellet. i) BCIP/NBT-stained microvessels trapped within the meshwork of a cellular strainer.

Fig. 3: Overview of library preparation and construction for subsequent RNA-sequencing using total RNA extracted from isolated microvessels

1. First-strand cDNA synthesis
2. Addition of Illumina adapters and barcodes
3. Library purification with NucleoMag beads
4. Depletion of ribosomal cDNA
5. Final PCR amplification
6. Library purification with NucleoMag beads
7. NGS sequencing on NovaSeq 6000 platform
8. Mapping of sequencing reads to the transcriptome

Figure 3. Overview of library preparation and construction for subsequent RNA-sequencing using total RNA extracted from isolated microvessels. Image generated using BioRender.

Fig. 4: Isolated microvesels are enriched from postmortem vmPFC tissue following the described protocol

Figure 4. a-c) Isolated microvessels are enriched from brain tissue following the described protocol. a-c) Brightfield images of chromogenically stained microvessels using BCIP/NBT substrate demonstrated isolation and enrichment of predominantly microvessels from postmortem vmPFC tissue samples. b) example of preserved microvessel morphology and integrity isolated from postmortem vmPFC tissue. c) Similar microvessel enrichment was observed when the same protocol was carried out using mouse cortex.

Fig. 5: Isolated microvessels retain their *in situ* structure and express canonical markers

Figure 5. a-b) Endothelial and smooth muscle cells immunolabelled for vimentin (green). c) Extracellular matrix laminins expressed by vascular-related cell types are immunolabelled (red). d) Endothelial cell tight junctions immunolabelled for CLDN5 (red). e) Pericytes immunolabelled for PDGFRβ (green). f) AQP4 expressed by the vessel-facing astrocyte membrane is immunolabelled (magenta) and laminins (green). g) Oligodendrocyte marker MBP (red) shows an absence of immunoreactivity in microvessel preparations, similarly, h) neuronal marker NeuN (red) shows an absence of immunoreactivity in microvessel preparations. Nuclei are stained with DAPI (blue) in all micrographs.

Fig. 6: **Computational deconvolution indicates high microvessel yield after isolation**

a

Deconvolution against VL dataset using CIBERSORT v1.04

	Cell Type				
Subject	Neurons	Astrocytes	Oligodendrocytes	Microglia	Endothelia
1	4.00	4.92	2.15	0.00	88.93
2	0.00	0.62	1.51	0.00	97.86
3	0.00	0.51	2.37	0.78	96.34
4	5.81	0.25	0.97	0.00	92.96
5	0.00	0.99	0.50	0.00	98.52
Average	1.29	1.16	1.85	0.16	94.92

Dataset provided by Velmeshev et al. 36 and performed using BrainDeconvShiny tool

b

Deconvolution against MB dataset using CIBERSORT v1.04

	Cell Type				
Subject	Neurons	Astrocytes	Oligodendrocytes	Microglia	Endothelia
1	12.35	11.82	0	0.00	75.83
2	0.00	5.78	0	0.00	94.22
3	0.00	3.30	0	3.97	92.73
4	16.52	5.53	0	0.00	77.95
5	0.00	6.19	0	0.00	93.81
Average	4.66	6.11	0	1.06	86.91

Multibrain composite signature generated by Sutton et al. 202213 and performed using BrainDeconvShiny

c

Deconvolution against Yang dataset using CIBERSORTX

	Cell Type													
Subject	Neuron	Oligo	OPC	Ependymal	M. Fibro	Microglia	T cell	P. Fibro	Astrocyte	Venous	Arterial	Capillary	SMC	Pericyte	Total Vascular
1	6.40	0.00	0	0	3.63	0.00	5.31	2.84	1.87	0	0.00	33.17	46.77	0.00	84.66
2	0.00	0.00	0	0	1.36	0.00	1.49	0.00	1.76	0	0.00	66.82	28.58	0.00	97.15
3	0.60	0.00	0	0	0.81	0.69	5.79	3.18	0.00	0	8.44	36.11	44.39	0.00	92.11
4	8.41	0.00	0	0	2.58	0.00	6.49	4.35	0.00	0	6.74	26.55	44.87	0.00	82.52
5	0.00	0.00	0	0	1.11	0.00	3.25	1.03	0.37	0	0.00	65.74	25.73	2.77	95.64
Average	4.01	0.14	0	0	2.01	0.24	5.64	2.33	0.98	0	2.53	44.02	37.62	0.46	90.42

Figure 6. a-b) Using the BrainDeconvShiny tool, different iterations of computational deconvolution were performed to demonstrate stability of outcome when using our microvessel isolation preparations. a) The average expression from every control sample from the VL dataset was calculated and used as cell type signatures for deconvolution of

our dataset using CIBERSORT v1.04. b) The MB (MultiBrain) is a composite signature generated by Sutton et al. (2022) by averaging the expression signatures of five datasets for five cell types (neurons, astrocytes, oligodendrocytes, microglia, and endothelia). MB was used as cell type signatures for deconvolution of our dataset using CIBERSORT v1.04. c) In-house analysis where the average expression from every control sample from the Yang dataset was calculated and used as cell type signatures for deconvolution of our dataset using CIBERSORTX. Heat tables generated using the gt package in R.

Fig. 7: Analysis of microvessel gene expression indicates overrepresentation of neurovascular-related gene expression

Figure 7. a-b) Enrichment analysis of top 10% most highly expressed genes (3968 genes) returns predominantly vascular-related terms. Over-representation analysis

(ORA) using the enrichR package in R and the "Descartes Cell Types and Tissue 2021" database was used to identify gene sets that are statistically over-represented. The threshold value of enrichment was selected by a p-value <0.05, indicating over-represented genes were significantly enriched for vascular-related terms. a) Count for genes in our dataset that are present in returned gene sets. b) Ratio for genes in our dataset that are present in returned gene sets, determined by the total number of genes in each set. c-i) Isolated microvessels have increased expression of canonical neurovascular-related genes. c) Bar plot showing TPMs for endothelial-defining genes. Highest expression found in endothelial genes B2M, BSG, FLT1, IFITM3, MT2A, SLC2A1, VIM, VWF. d) Bar plot showing TPMs for pericyte-defining genes. Highest expression was detected in pericyte genes CALD1, FN1, IGFBP7, RGS5, SPARCL1. e) Bar plot showing TPMs for smooth muscle cell-defining genes. Highest expression was detected in smooth muscle genes ACTG1, ACTN4, MYL6, PTMA and TAGLN. f) Bar plot showing TPMs for tight junction-defining genes and g) Bar plot showing TPMs for adherens junction-defining genes. Several genes encoding for junctional proteins are found in the top 10% of most highly expressed genes, including CLDN5, CTNNB1, CTNND1, OCLN, JAM1, TJP1, and TJP2. h) Bar plot showing TPMs for astrocyte and astrocytic endfeet-defining genes. Expression of astrocytic genes were predominantly limited to markers of astrocytic processes or endfeet, namely CLU, GFAP, and GLUL. i) TPMs from neurovascular-related genes were summarized according to cell type expression, demonstrating overrepresentation of endothelial, smooth muscle cell, and pericyte genes. Bar plots generated using the ggplot package in R.

Fig. 8: Comparison of RNA sequencing data to published neurovascular dataset

Figure 8. a-g) Overlap between top 10% of highly expressed genes from our RNA sequencing data and published NVU dataset. Validated neurovascular cell type markers were obtained from Garcia et al. (2022) postmortem and human *ex vivo* single-nuclei sequencing datasets (found under supplementary table 2) and compared for potential overlap with the top 10% of most highly expressed genes from our data. Results indicated significant overlap between the three datasets for all NVU-related cell types. a) Arteriole-defining genes. b) Capillary-defining genes. c) Venule-defining genes. d) Arteriolar SMC-defining genes. e) Pericyte-defining genes. f) Venular SMC-defining genes. g) Fibroblast-defining genes. Venn diagrams generated using the ggVennDiagram package in R.

Fig. 9: Overlap between transcriptomic and proteomic data output

Figure 9. a-i) There is substantial overlap between transcriptomic and proteomic data

output. a) 1635/1637 (99.9%) of proteins likewise identified in the transcriptomics data

and with 1024/1637 (62.6%) proteins found in the top 10% of most highly expressed genes. b) Several canonical BMEC markers were detected with high normalized average abundance. c) Over-representation analysis using the enrichR package in R and the "Descartes Cell Types and Tissue 2021" database was used to identify sets that are statistically over-represented. Genes corresponding to proteins detected during LC-MS/MS were used as input and the threshold value of enrichment was selected by a p-value <0.05. Output indicated over-represented proteins were significantly enriched for vascular-related terms. a) Count for peptides in our dataset that are present in returned sets. b) Ratio for peptides in our dataset that are present in returned sets, determined by the total number in each set. d) Bar plot showing normalized abundance for endothelial cell-defining proteins. e) Bar plot showing normalized abundance for pericyte-defining proteins. f) Bar plot showing normalized abundance for smooth muscle cell defining-proteins. g) Bar plot showing normalized abundance for tight junction defining-proteins. h) Bar plot showing normalized abundance for adherens junction-defining proteins. i) Bar plot showing normalized abundance for astrocyte and astrocytic endfeet-defining proteins. Bar plots generated using the ggplot package and Venn diagram using the ggVennDiagram package in R.

Fig. 10: Isolated microvessels can be used to further sort endothelial nuclei

Figure 10. a-b) Isolated microvessels can be used to further sort endothelial nuclei. a-b) Dissociated endothelial nuclei immunolabelled with ERG conjugated to Alexa-647 antibody prior to FANS protocol. Resuspended microvessels subjected to a 2-hour incubation with anti-ERG antibody under rotation is sufficient for immunolabelling of endothelial nuclei prior to FANS sorting.