Leveraging inter-individual transcriptional correlation structure to infer discrete signaling mechanisms across metabolic tissues

Mingqi Zhou[1,2*], Ian J. Tamburini[1,2*], Cassandra Van[1,2*], Jeffrey Molendijk[3], Christy M Nguyen[1,2], Ivan Yao-Yi Chang[1], Casey Johnson[1,2], Leandro M. Velez[1,2], Youngseo Cheon[1,2], Reichelle X. Yeo[4], Hosung Bae[1,2], Johnny Le[1,2], Natalie Larson[1,2], Ron Pulido[1,2], Carlos Filho[1,2], Cholsoon Jang[1,2], Ivan Marazzi, [1,2], Jamie N. Justice[7], Nicholas Pannunzio[1,2], Andrea Hevener[5,6], Lauren M. Sparks[4], Erin E. Kershaw[8], Dequina Nicholas[9], Benjamin Parker[3], Selma Masri[1,2] and Marcus Seldin[1,12]

[1]Department of Biological Chemistry and [2]Center for Epigenetics and Metabolism, UC Irvine. Irvine, CA, USA
[3]Department of Anatomy and Physiology, University of Melbourne, Melbourne, VIC, Australia
[4]Translational Research Institute, AdventHealth, Orlando, FL, USA
[5]Department of Medicine, Division of Endocrinology, Diabetes, and Hypertension, and [6]Iris Cantor-UCLA Women's Health Research Center, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA
[7]Veterans Administration Greater Los Angeles Healthcare System, Geriatric Research Education and Clinical Center (GRECC), Los Angeles, CA, USA
[8]Department of Internal Medicine, Section On Gerontology and Geriatric Medicine, Wake Forest School of Medicine, Winston-Salem, NC, USA
[9]Division of Endocrinology, Department of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, USA
[10]Divison of Hematology/Oncology, Department of Medicine, University of California Irvine, Irvine, CA USA
[11]Department of Molecular Biology and Biochemistry, School of Biological Sciences, University of California Irvine, Irvine CA, USA

*Authors contributed equally
[12]Corresponding author

To whom correspondence should be addressed:
Marcus Seldin
UC Irvine Department of Biological Chemistry
314 Sprague Hall
Irvine, CA 92697
Phone: 949-824-6765
Email: mseldin@uci.edu

Running title: A targeted exploration of gene-centric correlations across metabolic organs

Keywords: Endocrinology, Inter-tissue communication, Organ cross-talk, Systems genetics, Inter-individual variation, Web resource

Abstract/Introduction

Inter-organ communication is a vital process to maintain physiologic homeostasis, and its dysregulation contributes to many human diseases. Beginning with the discovery of insulin over a century ago, characterization of molecules responsible for signal between tissues has required careful and elegant experimentation where these observations have been integral to deciphering physiology and disease. Given that circulating bioactive factors are stable in serum, occur naturally, and are easily assayed from blood, they present obvious focal molecules for therapeutic intervention and biomarker development. For example, physiologic dissection of the actions of soluble proteins such as proprotein convertase subtilisin/kexin type 9 (*PCSK9*) and glucagon-like peptide 1 (*GLP1*) have yielded among the most promising therapeutics to treat cardiovascular disease and obesity, respectively[1-4]. A major obstacle in the characterization of such soluble factors is that defining their tissues and pathways of action requires extensive experimental testing in cells and animal models. Recently, studies have shown that secreted proteins mediating inter-tissue signaling could be identified by "brute-force" surveys of all genes within RNA-sequencing measures across tissues within a population[5-9]. Expanding on this intuition, we reasoned that parallel strategies could be used to understand how individual genes mediate signaling across metabolic tissues through correlative analyses of gene variation between individuals. Thus, comparison of quantitative levels of gene expression relationships between organs in a population could aid in understanding cross-organ signaling. Here, we surveyed gene-gene correlation structure across 18 metabolic tissues in 310 human individuals and 7 tissues in 103 diverse strains of mice fed a normal chow or HFHS diet. Variation of genes such as *FGF21, ADIPOQ, GCG* and *IL6* showed enrichments which recapitulate experimental observations. Further, similar analyses were applied to explore both within-tissue signaling mechanisms (liver *PCSK9*) as well as genes encoding enzymes producing metabolites (adipose *PNPLA2*), where inter-individual correlation structure aligned with known roles for these critical metabolic pathways. Examination of sex hormone receptor correlations in mice highlighted the difference of tissue-specific variation in relationships with metabolic traits. We refer to this resource as **G**ene-**D**erived **C**orrelations **A**cross **T**issues (GD-CAT) where all tools and data are built into a web portal enabling users to perform these analyses without a single line of code (gdcat.org). This resource enables querying of any gene in any tissue to find correlated patterns of genes, cell types, pathways and network architectures across metabolic organs.

Results

Construction of a web tool to survey transcript correlations across tissues and individuals (GD-CAT) – Previous studies have established that "brute force" analyses of correlation structure across tissues from population expression data can identify new several known and mechanisms of organ

cross-talk. These were accomplished by surveying the global correlation structure using all genes,
whereby skewed upper-limits of significance distributions were sufficient to prioritize proteins
which elicit signaling[5–9]. Following this intuition, we hypothesized that a paralleled but alternative
approach to inter-individual correlation structure could be exploited to understand the functional
consequences of specific genes. Our initial goal was to establish a user-friendly interface where
all of these analyses and gene-centric queries could be performed without running any code. To
accomplish this, we assembled a complete analysis pipeline (Fig 1A) as a shiny-app and docker
image hosted in a freely-available web address (gdcat.org). Here, users can readily-search gene
correlation structure between individuals from filtered human (gene-by-tissue expression project
- GTEx) and mouse (hybrid mouse diversity panel - HMDP) across tissues. GTEx is presently the
most comprehensive pan-tissue dataset in humans[10], which was filtered for individuals where most
metabolic tissues were sequenced[9]. Collectively, this dataset contains 310 individuals, consisting
of 210 male and 100 female (self-reported) subjects between the ages of 20-79. Data from the
HMDP consisted of 96 diverse mouse strains fed a normal chow (5 tissues) or high-fat/high-
sucrose diet (7 tissues) as well as carefully characterized clinical traits [11–16]. Initially, users select
a given species, followed by reported sex or diet (mouse) which loads the specified environment.
Subsequent downstream analyses are then implemented accordingly from a specific gene in a
given tissue.. This selection prompts individual gene correlations across all other gene-tissue
combinations using biweight midcorrelation[17]. From these charts, users are able to select a given
tissue, where gene set enrichment analysis testing using clusterprofiler[18] and enrichR [19] are applied
to the correlated set of genes to determine the positively (activated) and negatively (suppressed)
pathways which occur in each tissue. In addition to general queries of gene ~ gene correlation
structure, comparison of expression changes are also visualized between age groups as well as
reported sexes. In addition, we included the top cell-type abundance correlations with each gene.
To compute cell abundance estimates from the same individuals, we used single-nucleus RNA-
seq available from GTEx[20] and applied cellular deconvolution methods to the bulk RNA-seq[21]
(methods). Comparison of deconvolution methods[21] showed that DeconRNASeq[22] captured the
most cell types within several tissues (Supplemental Figures 1-3) and therefore was applied to all
tissues where sn-RNA-seq was available. We note that visceral adipose, subcutaneous adipose,
aortic artery, coronary artery, transverse colon, sigmoid colon, the heart left ventricle, the kidney
cortex, liver, lung, skeletal muscle, spleen, and small intestine are the only tissues where sn-seq is
available and not other tissues, such as brain, stomach and thyroid.
We initially examined pan-tissue transcript correlation structures for several well-established
mechanisms of tissue crosstalk via secreted proteins which contribute to metabolic homeostasis.
Here, binning of the significant tissues and pathways related to each of these established secreted
proteins resembled their known mechanisms of action (Fig 1B-E). For example, variation with
subcutaneous adipose expression of *ADIPOQ* was enriched with genes in several metabolic tissues
where it has been known to act (Fig 1B, left). In particular, subcutaneous adipose *ADIPOQ*
expression correlated with fatty acid oxidative process within adipose (Fig 1B, middle) and was
enriched with ECM, chemotaxis and ribosomal biogenesis in skeletal muscle (Fig 1B, right).
These correlated pathways align with the established physiologic roles of the protein in that fat
secreted adiponectin when oxidation is stimulated[23,24] and muscle is a major site of action[25].
Beyond adiponectin, inter-individual correlation structure additionally recapitulated broad
signaling mechanisms for other relevant endocrine proteins. For example, intestinal *GCG*
(encoding GLP1, Fig 1C), liver *FGF21* (Fig 1D) and skeletal muscle *IL6* (Fig 1E) showed binning
patterns and pathway enrichments related to their known functions in pancreas[1,26], adipose

tissue[27,28] and other metabolic organs[29], respectively. These analyses and web tool show some
examples of exploring transcriptional correlation structure to confirm and identify mechanisms of
signaling, where we note that additional limitations should be considered.
136
Pathway-based examination of gene correlation structure and significance thresholds across
tissues - While the select observations shown in Fig1 provide examples of support in exploring
correlation structure of genes across interindividual differences to investigate endocrinology,
several limitations in these analyses should be considered. First, an additional explanation for a
given gene showing strong correlation between the tissues could arise from a general pattern of
correlation between the two tissues and not necessarily due to the discrete signaling mechanisms.
In previous studies surveying correlation structure and network model architectures in the HMDP
and STARNET populations, genes appeared generally stronger correlated between liver and
adipose tissue compared to all other organ combinations explored[5–7]. To investigate this global
pattern of gene correlation structure between metabolic organs, we selected key GO terms, KEGG
pathways and randomly sampled equal numbers of genes and evaluated relative significance of
inter-tissue correlations across multiple statistical thresholds. These analyses suggested that usage
of empirical student correlation pvalues recapitulated a clear pattern of inter-tissue correlations
between pathways (Fig 2). For example, comparison of the number of genes achieving
significance of correlation between tissues among select GO terms revealed that tissues such as
adipose and muscle appeared more correlated than spleen and other tissues at pvalues less than 1e-
3 (Fig 2A, left column). These global patterns of gene correlation between tissues among select
pathways were reduced when the pvalue threshold was lowered to 1e-6 (Fig 2A, middle column)
or qvalue adjustments (methods) were performed (Fig 2A, right two columns). For these reasons,
only qvalue adjusted value were used and implemented into pie charts providing the tissue-specific
occurrences of correlated genes at 3 thresholds (q<0.1, q<0.01, q<0.001) within the web tool.
Next, in order to further evaluate these global patterns of innate transcript correlation structure and
determine whether they reflected concordance between known metabolic pathways or innate to
the dataset used, tissues were rank-ordered by the number of genes which meet pvalue thresholds
and compared to randomly sampled genes of similar pathway sized (Fig 2B). Among KEGG
Pathways selects (hsa04062 − Chemokine signaling pathway, hsa04640 − Hematopoietic cell
lineage and hsa00190 − Oxidative phosphorylation), the top-ranked organs by correlated gene
numbers differed (Skeletal muscle, Colon and Thyroid, respectively); however, a general trend of
specific tissues ranking higher than others were observed (Fig 2B). For example, skeletal muscle
and heart appeared among the strongest correlated across pathways and organs, compared to
kidney cortex and spleen which were observed to rank among the lowest (Fig 2B, pathways). We
note that when the same analysis was performed on randomly sampled genes from each organ
consisting of the same number as genes within each KEGG pathway, these rankings and number
of significant correlating genes were no longer observed (Fig 2B, random genes), suggesting that
in certain instances differences between organs in general connectivity to others might reflect
concordance between known pathways. It is important to consider here that for the organs ranking
lower, the lack of relative correlating numbers is likely due to sparsity of available data and not
necessarily general patterns of gene correlation. This point is supported by the fact that among the
lowest-ranked 33% of tissues across pathways, we observed a significant negative overall
correlation (bicor = -0.45, pvalue = 2.3e-5) between number of NA values per individual and the
gene count for significance shown in Fig 2B. This negative correlation between missing data and
number of significant correlations for pathways across tissues was not observed when binning the

top 33% (bicor = 0.09, pvalue = 0.42) or middle 33% (bicor = -0.12, pvalue = 0.27) of organs.
Collectively, these analyses show that innate correlations structures exist between organs which
differ depending on pathways investigated and that tissues which don't show broad correlation
structure could potentially be attributed to areas of missing data among GTEx.

PSCK9 signaling and lipid exchange between adipose and muscle apparent in simple network
models of correlation structure - Next, we wanted to ask whether our approach of analyzing inter-
individual correlation structure across tissue for endocrine proteins was also sufficient to define
within-tissue signaling mechanisms or actions of enzymes producing metabolites that signal across
organs. Dissimilar to the cross-tissue distributions of significance in Fig 1, the same analysis of
liver *PCSK9* highlighted exclusively liver genes which were varied together (Fig 2A), in particular
those involved in cholesterol metabolism/homeostasis (Fig 2B). Consistent with the established
role for PCSK9 as a primary degradation mechanism of LDLR[4,30], network model construction of
correlated genes highlighted the gene as a central node linking cholesterol biosynthetic pathways
with those involved in other metabolic pathways such as insulin signaling (Fig 2C). Given that
organ signaling via metabolites comprises many critical processes among multicellular organisms,
our next goal was to apply this gene-centric analyses to established mechanisms of metabolite
signaling.   The gene *PNPLA2* encodes adipose triglyceride lipase (ATGL) which localizes to
lipid droplets and breaks down triglycerides for oxidation or mobilization as free fatty acids for
peripheral tissues[31]. Variation in expression of *PNPLA2* showed highly significant enrichments
with beta oxidation pathways in adipose tissue (Fig 2D). Muscle pathways enriched for the gene
were represented by sarcomere organization and muscle contraction (Fig 2F). Construction of an
undirected network from these expression data placed the gene as a central node between the two
tissues, linking regulators of adipose oxidation (Fig 2F, red) to muscle contractile process (Fig 2F,
purple) where additional strongly co-correlated genes were implicated as additional candidates
(Fig 2F). In sum, these analyses provide two examples of within-liver signaling via *PCSK9* and
adipose-muscle communication through *PNPLA2* where the top-correlated genes and network
models recapitulate known mechanisms. Given the utility of these undirected network models, a
function in GD-CAT was added to enable users to generate network models for any gene-tissue
combination and select parameters such as number of within-tissue and peripheral correlated genes
to include.

Inter-individual correlation analysis of hybrid mouse diversity panel highlights tissue- and diet-
specific phenotype relationships with sex hormones - Genetic reference panels in model organisms,
such as mice, present appeal in studying complex traits in that environmental conditions can be
tightly controlled, tissues and invasive traits readily accessible and the same, often renewable,
genetic background can be studied and compared among multiple exposures such as diets or drug
treatments [15,32–34]. For this resource, we utilized data from the HMDP fed a normal chow[15,16] or
HFHS diet for 8 weeks[11–14]. While the number of tissues available was less than in GTEx, these
panels allow for comparison of how gene correlations shift depending on diet. Therefore, queries
of gene correlation queries in mice were segregated into either chow or HFHS diet and an
additional panel to download a table or visualize the relationship between genes and clinical
measures was added. The inferred abundances of cell types from each individual are correlated
across user-defined genes, with the bicor coefficient plotted for each cell type.
One advantage of hybrid mouse diversity panel data compared to GTEx is the abundance of
phenotypic measures available within each cohort. To show the utility of examining correlations

within this reference panel, we selected sex hormone receptors androgen receptor (*Ar*), estrogen
receptor alpha (*Esr1*) or estrogen receptor beta (*Esr2*) and binned the top 10 phenotypes which
were correlated. These analyses were segregated based on where sex hormones were expressed
(either liver or adipose tissue) or dietary regiment of the ~100 strains (normal chow or HFHS diet).
This analysis demonstrated the difference in relationships between tissue location of sex hormone
receptor and dietary context with metabolic traits. For example, expression of *Ar* in adipose tissue
among HMDP mice fed a HFHS diet was negatively correlated with fat mass and body weight
traits, whereas expression in liver oppositely correlated with the same traits in a positive direction
(Fig 4A). The top traits which correlated also differed by tissue or expression for *Ar,* such as
plasma lipid parameters in adipose tissue compared to blood cell traits in chow-fed mice (Fig 4A).
We note that among the three hormone receptors investigated, *Esr2* appeared the most consistently
correlated between tissues and diets with metabolic traits (Fig 4B). Expression of *Esr1* also
showed a clear tissue and diet difference in the traits which were the most strongly co-regulated.
Under HFHS dietary conditions, a negative correlation with insulin and fat pad weights were
observed exclusively with adipose expression, while positive correlations with liver lipids were
observed with expression in liver (Fig 4C). These analyses highlight how phenotype correlations
in mouse populations can help to determine contexts relevant for gene regulation and point to the
diversity of potential contexts relevant for sex hormone receptors in metabolic tissues.
243
Discussion
Limitations and Conclusions – Here, we provide a new resource to explore correlations across
organ gene expression in the context of interindividual differences. We highlight areas where
these align with established and relevant mechanisms of physiology and suggest that similar
explorations could be used as a discovery tool. Several key limitations should be considered when
exploring GD-CAT for mechanisms of inter-tissue signaling though. Primarily, the fact that
correlation-based analyses could reflect both causal or reactive patterns of variation. While several
statistical methods such as mediation[35,36] and mendelian randomization[37,38] exist to further refine
causal inferences, likely the only definitive method to distinguish is in carefully-designed
experimentation. Further, analyses of genetic correlation (ex. correlations considering genetic loci
to infer causality) also present appeal in refining some causal mechanisms. Correlation between
molecular and phenotypic variables can occur for a variety of reasons, not just between their
individual relationships, but often more broadly, from a variety of complex genetic and
environmental factors. Further, many correlations tend to be dominated by genes expressed within
the same organ. This could be due to the fact that, within-tissue correlations could capture both
the pathways regulating expression of a gene, as well as potential consequences of changes in
expression/function, and distinguishing between the two presents a significant challenge. For
example, a GD-CAT query of insulin (*INS*) expression in pancreas shows exclusive enrichments
in pancreas and corresponding pathway terms reflect regulatory mechanisms such as secretion and
ion transport (Supplemental Fig 4). Representation of given genes may also differ significantly
depending on the dataset used. For example, while queries of other tissues for the critical X
Inactive Specific Transcript (*XIST*), in liver no significant correlations appear. This is due to the
fact that the gene operates in a sex-dependent manner, where females are significantly less
represented in GTEx and liver exists as a sparser tissue compared to others (Fig 2). In addition,
the analyses presented are derived from differences in gene expression across individuals which
arise from complex interaction of genetic and environmental variables. Expression of a gene and
its corresponding protein can show substantial discordances depending on the dataset used. These

have been discussed in detail[39–41], but ranges of co-correlation can vary widely depending on the
datasets used and approaches taken. We note that for genes encoding proteins where actions from
acute secretion grossly outweigh patterns of gene expression, such as insulin, caution should be
taken when interpreting results. As the depth and availability of tissue-specific proteomic levels
across diverse individuals continues to increase, an exciting opportunity is presented to explore
the applicability of these analyses and identify areas when gene expression is not a sufficient
measure. For example, mass-spec proteomics was recently performed on GTEx[42]; however, given
that these data represent 6 individuals, analyses utilizing well-powered inter-individual
correlations such as ours which contain 310 individuals remain limited in applications.
The queries provided in GD-CAT use fairly simple linear models to infer organ-organ signaling;
however, more sophisticated methods can also be applied in an informative fashion. For example,
Koplev et al generated co-expression modules from 9 tissues in the STARNET dataset, where
construction of a massive Bayesian network uncovered interactions between correlated modules[6].
These approaches expanded on analysis of STAGE data to construct network models using
WGCNA across tissues and relating these resulting eigenvectors to outcomes[43]. The generalized
approach of constructing cross-tissue gene regulatory modules presents appeal in that genes are
able to be viewed in the context of a network with respect to all other gene-tissue combinations.
In searching through these types of expanded networks, individuals can identify where the most
compelling global relationships occur. One challenge with this type of approach; however, is that
coregulated pathways and module members are highly subjective to parameters used to construct
GRNs (for example reassignment threshold in WGCNA) and can be difficult in arriving at a
"ground truth" for parameter selection. We note that the WGCNA package is also implemented
in these analyses, but solely to perform gene-focused correlations using biweight midcorrelation
to limit outlier inflation. While the midweight bicorrelation approach to calculate correlations
could also be replaced with more sophisticated models, one consideration would be a concern of
overfitting models and thus, biasing outcomes.
In another notable example MultiCens was developed as a tool to uncover communication
between genes and tissues and applied to suggest central processes which exist in multi-layered
data relevant for Alzheimer's disease[44]. In addition, Jadhav and colleagues adopted a machine
learning approach to mine published literature for relationships between hormones and genes[45].
Further, association mapping of plasma proteomics data has been extensively applied and
intersection with genome-wide association disease loci has offered intriguing potential disease
mechanisms[46,47]. Another common application to single-cell sequencing data is to search for
overrepresentation of known ligand-receptor pairs between cell types[48]. These and additional
applications to explore tissue communication/coordination present unique strengths and caveats,
depending on the specific usage desired. Regardless of methods used to decipher, one important
limitation to consider in all these analyses is the nature of underlying data. For example, our
evaluation of GTEx data structure suggested that important organs such as spleen and kidney were
insufficient due to availability in matching expression data between individuals. Further, GTEx
sample vary as to the collection times, sample processing times and other important parameters
such as cause of death. Mouse population data such as the HMDP or BxD cohorts offer appeal in
these regards, as environmental conditions and collection times are easily fixed. Regardless,
careful consideration of how data was generated and normalized are fundamental to interpreting
results.
In sum we demonstrate that adopting a gene-centric approach to surveying correlation structure
of transcripts across organs and individuals can inform mechanism of coordination between

metabolic tissues. Initially, we queried several well-established and key mediators of physiologic homeostasis, such as *FGF21, GCG* and *PCSK9*. These approaches are further suggested to be applicable to mechanisms of metabolite signaling, as evident by pan-tissue investigation of adipose *PNPLA2*. Exploration of hybrid mouse diversity panel data highlighted the diverse phenotype correlations depending on tissue and diet for sex hormone receptors. To facilitate widespread access and use of this transcript isoform-centric analysis of inter-individual correlations, a full suite of analyses such as those performed here can be performed from a lab-hosted server (gdcat.org) or in isolation from a shiny app or docker image.

Material and methods

Availability of web tool and analyses: All analyses, datasets and scripts used to generate the associated web tool (GD-CAT) can be accessed via: https://github.com/mingqizh/GD-CAT or within the associated docker image. In addition, access to the GD-CAT web tool is also available through the web portal gdcat.org. This portal was created to provide a user-friendly interface for accessing and using the GD-CAT tool without the need to download or install any software or packages. Users can simply visit the website, process data and start using the tool. Corresponding tutorial and the other resources were made available to facilitate the utilization of the web tool on GitHub. The interface and server of the web were built and linked based on the shiny package using R (v. 4.2.0). Shiny package provides a powerful tool for building interactive web applications using R, allowing for fast and flexible development of custom applications with minimal coding required.

Pathway-specific gene correlations across tissues: Detailed scripts and analyses for pathway-specific investigations across tissues in Fig2 are provided in: https://github.com/itamburi/gtex-app-kegg-pathways. Briefly, to interrogate broad tissue correlation structure, the number of genes which passed each biweight midcorrelation pvalue cutoff are shown normalized to the total number of genes corresponding to that pathway term. Pathways were selected by accessing all available GO annotations for all genes using the Universal Protein Resource[49] and subletting genes where a given term is listed.   To determine which tissues show the most co-correlation across genes and organs, KEGG terms shown were selected and each corresponding gene-tissue combinations were correlated. Tissues were then binned by the number of significant correlations which occurring both within and across organs among each selected KEGG pathway at indication correlation pvalues. Rank-ordering on the figure was shown by chemokine signaling at P<0.01 and each term was compared to a randomly sampled set of genes corresponding to the same number contained in each pathway.

Data sources and availability: All human data used in this study can be immediately accessed via web tool or docker to facilitate analysis. Metabolic tissue data was accessed through GTEx V8 downloads portal on August 18, 2021 and previously described[9,10]. These raw data can also be readily accessed from the associated R-based walkthrough: https://github.com/Leandromvelez/myokine-signaling. Briefly, these data were filtered to retain genes which were detected across tissues where individuals were required to show counts > 0 across all data. Given that our goal was to look across tissues at enrichments, this was done to limit spurious influence of genes only expressed in specific tissues in specific individuals. Hybrid mouse diversity panel data was collected from previously described studies[11,15,16,34] and

362 inter-individual differences were compared at the strain-level to maximize possible comparisons
363 between historical data.
364
Correlation analyses across tissues – biweight midcorrelation coefficients and corresponding p-
values within and across tissues were generated using WGCNA bicorandpvalue() function[17]. We
note that while the WGCNA package was used to calculate coefficients and corresponding
students pvalues, this generalized framework does not utilize any module generation. Associated
qvalue adjustments were applied using the Benjamini-Hochberg FDR from the R package
"stats". These BH adjustments, as opposed to standard qvalue adjustments, were selected given
their efficiency in CPU usage on the hosted server.
372
Pathway enrichment analyses – Pathway enrichments were generated using gene set enrichment
analyses available from the r package clusterprofiler. Specifically, the bicor coefficients were used
as the rank-weight of each gene and enrichment tests performed by permuting against the human
or mouse reference transcriptome. Terms used for the enrichment analyses were derived from
Gene Ontology (Biological Process, Cellular Component and Molecular Function) which were
accessed using the R package enrichR. For this analysis and on the available app, input genes
were determined at indicated qvalue threshold.
Deconvolution of bulk tissue seq data on web tool. All scripts and deconvolution data produced is
available at: https://github.com/cvan859/deconvolution. Briefly, sn-RNA-seq data was accessed
from the Human cell atlas[20] for matching organ datasets with metabolic tissues. From these data,
4 deconvolution methods were applied using ADAPTS[21] where DeconRNA-Seq[22] was selected
for its ability to capture the abundances of the most cell types across tissues such as liver heart and
skeletal muscle (Supplemental Fig 1-3). The full combined matrix was assembled for DeconRNA-
Seq results across individuals in GTEx where correlations between cell types and genes was
performed also using the bicorandpvalue() in WGCNA[17].

Acknowledgements
We acknowledge the following funding sources for supporting these studies: MZ, CF, IJT, CMN,
LMV, CV, CJ and MMS were supported by NIH grants HL138193, DK130640 and DK097771.
ALH is supported by NIH grants U54 DK120342, R01 DK109724, and P30 DK063491. NRP is
supported by NIH grant R37 CA266042. HB was supported by National Research Foundation of
Korea (2021R1A6A3A14039132). JL was supported by an NIH grant F31DK134173-01A1. CJ
was supported by AASLD Foundation Pinnacle Research Award in Liver Disease, Edward
Mallinckrodt, Jr. Foundation Award, and an NIH grant AA029124.

396
Author contributions
MZ, IJT, CV and MMS accessed raw data, performed analyses and drafted the manuscript. MZ
assembled the shiny application, where IV and RP designed the UCI infrastructure to enable
access. CJ, CMN, LMV, JM, CF, IC, RY, HB, JL, NL, IM, AH, LMS, JNJ, EEK, IM, NP, DN
and BM provided critical insight into data use and interpretation, as well as guided the study. CJ,
NP and SM guided tool design involving metabolite signaling and circadian rhythms, respectively,
as well as provided app infrastructure. All authors have approved the current manuscript. All
authors read and approved this manuscript.
405
Conflict of interest

The authors have no conflicts of interest to declare

References

1. Drucker, D. J. GLP-1 physiology informs the pharmacotherapy of obesity. *Molecular Metabolism* **57**, 101351 (2022).

2. Trapp, S. & Stanford, S. C. New developments in the prospects for GLP-1 therapy. *British J Pharmacology* **179**, 489–491 (2022).

3. Dadu, R. T. & Ballantyne, C. M. Lipid lowering with PCSK9 inhibitors. *Nat Rev Cardiol* **11**, 563–575 (2014).

4. Lambert, G., Sjouke, B., Choque, B., Kastelein, J. J. P. & Hovingh, G. K. The PCSK9 decade. *Journal of Lipid Research* **53**, 2515–2524 (2012).

5. Seldin, M. M. *et al.* A Strategy for Discovery of Endocrine Interactions with Application to Whole-Body Metabolism. *Cell Metab.* **27**, 1138-1155.e6 (2018).

6. Koplev, S. *et al.* A mechanistic framework for cardiometabolic and coronary artery diseases. *Nat Cardiovasc Res* **1**, 85–100 (2022).

7. Seldin, M. M. & Lusis, A. J. Systems-based approaches for investigation of inter-tissue communication. *J Lipid Res* **60**, 450–455 (2019).

8. Cao, Y. *et al.* Liver-heart cross-talk mediated by coagulation factor XI protects against heart failure. *Science* **377**, 1399–1406 (2022).

9. Velez, L. M. *et al.* Genetic variation of putative myokine signaling is dominated by biological sex and sex hormones. *eLife* **11**, e76887 (2022).

10. GTEx Consortium *et al.* Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).

11. Parks, B. W. *et al.* Genetic architecture of insulin resistance in the mouse. *Cell Metab* **21**, 334–347 (2015).

12. Hui, S. T. *et al.* The genetic architecture of NAFLD among inbred strains of mice. *Elife* **4**,

e05607 (2015).

13. Norheim, F. *et al.* Genetic regulation of liver lipids in a mouse model of insulin resistance

and hepatic steatosis. *Mol Syst Biol* **17**, e9684 (2021).

14. Org, E. *et al.* Genetic and environmental control of host-gut microbiota interactions. *Genome*

Res **25**, 1558–1569 (2015).

15. Lusis, A. J. *et al.* The Hybrid Mouse Diversity Panel: a resource for systems genetics

analyses of metabolic and cardiovascular traits. *J Lipid Res* **57**, 925–942 (2016).

16. Bennett, B. J. *et al.* A high-resolution association mapping panel for the dissection of

complex traits in mice. *Genome Res* **20**, 281–290 (2010).

17. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

18. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative Biology* **16**,

284–287 (2012).

19. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server

2016 update. *Nucleic Acids Res* **44**, W90-97 (2016).

20. The Tabula Sapiens Consortium* *et al.* The Tabula Sapiens: A multiple-organ, single-cell

transcriptomic atlas of humans. *Science* **376**, eabl4896 (2022).

21. Danziger, S. A. *et al.* ADAPTS: Automated deconvolution augmentation of profiles for

tissue specific cells. *PLoS ONE* **14**, e0224693 (2019).

22. Gong, T. & Szustakowski, J. D. DeconRNASeq: a statistical framework for deconvolution of

heterogeneous tissue samples based on mRNA-Seq data. *Bioinformatics* **29**, 1083–1085

(2013).

23. da Silva Rosa, S. C., Liu, M. & Sweeney, G. Adiponectin Synthesis, Secretion and

Extravasation from Circulation to Interstitial Space. *Physiology* **36**, 134–149 (2021).

24. Straub, L. G. & Scherer, P. E. Metabolic Messengers: adiponectin. *Nat Metab* **1**, 334–339

(2019).

25. Ruan, H. & Dong, L. Q. Adiponectin signaling and function in insulin target tissues. *J Mol*

Cell Biol **8**, 101–109 (2016).

26. McLean, B. A. *et al.* Revisiting the Complexity of GLP-1 Action from Sites of Synthesis to

Receptor Activation. *Endocrine Reviews* **42**, 101–132 (2021).

27. Fisher, F. M. & Maratos-Flier, E. Understanding the Physiology of FGF21. *Annu. Rev.*

Physiol. **78**, 223–241 (2016).

28. Flippo, K. H. & Potthoff, M. J. Metabolic Messengers: FGF21. *Nat Metab* **3**, 309–317

(2021).

29. Pedersen, B. K. & Febbraio, M. A. Muscle as an Endocrine Organ: Focus on Muscle-

Derived Interleukin-6. *Physiological Reviews* **88**, 1379–1406 (2008).

30. Peterson, A. S., Fong, L. G. & Young, S. G. PCSK9 function and physiology. *Journal of*

Lipid Research **49**, 1152–1156 (2008).

31. Zechner, R., Kienesberger, P. C., Haemmerle, G., Zimmermann, R. & Lass, A. Adipose

triglyceride lipase and the lipolytic catabolism of cellular fat stores. *Journal of Lipid*

Research **50**, 3–21 (2009).

32. Li, H. & Auwerx, J. Mouse Systems Genetics as a Prelude to Precision Medicine. *Trends in*

Genetics **36**, 259–272 (2020).

33. Andreux, P. A. *et al.* Systems genetics of metabolism: the use of the BXD murine reference

panel for multiscalar integration of traits. *Cell* **150**, 1287–1299 (2012).

34. Seldin, M., Yang, X. & Lusis, A. J. Systems genetics applications in metabolism research.

Nat Metab **1**, 1038–1050 (2019).

35. Richiardi, L., Bellocco, R. & Zugna, D. Mediation analysis in epidemiology: methods,

interpretation and bias. *International Journal of Epidemiology* **42**, 1511–1519 (2013).

36. Zeng, P., Shao, Z. & Zhou, X. Statistical methods for mediation analysis in the era of high-

throughput genomics: Current successes and future challenges. *Computational and*

Structural Biotechnology Journal **19**, 3209–3224 (2021).

37. Emdin, C. A., Khera, A. V. & Kathiresan, S. Mendelian Randomization. *JAMA* **318**, 1925

(2017).

38. Sanderson, E. *et al.* Mendelian randomization. *Nat Rev Methods Primers* **2**, 6 (2022).

39. Liu, Y., Beyer, A. & Aebersold, R. On the Dependency of Cellular Protein Levels on mRNA

Abundance. *Cell* **165**, 535–550 (2016).

40. Maier, T., Güell, M. & Serrano, L. Correlation of mRNA and protein in complex biological

samples. *FEBS Letters* **583**, 3966–3973 (2009).

41. Buccitelli, C. & Selbach, M. mRNAs, proteins and the emerging principles of gene

expression control. *Nat Rev Genet* **21**, 630–644 (2020).

42. Jiang, L. *et al.* A Quantitative Proteome Map of the Human Body. *Cell* **183**, 269-283.e19

(2020).

43. Talukdar, H. A. *et al.* Cross-Tissue Regulatory Gene Networks in Coronary Artery Disease.

Cell Systems **2**, 196–208 (2016).

44. Kumar, T., Sethuraman, R., Mitra, S., Ravindran, B. & Narayanan, M. *MultiCens:*

Multilayer network centrality measures to uncover molecular mediators of tissue-tissue

communication. http://biorxiv.org/lookup/doi/10.1101/2022.05.15.492007 (2022)

doi:10.1101/2022.05.15.492007.

45. Jadhav, A., Kumar, T., Raghavendra, M., Loganathan, T. & Narayanan, M. Predicting cross-

tissue hormone–gene relations using balanced word embeddings. *Bioinformatics* **38**, 4771–

4781 (2022).

46. Ferkingstad, E. *et al.* Large-scale integration of the plasma proteome with genetics and

disease. *Nat Genet* **53**, 1712–1721 (2021).

47. Suhre, K., McCarthy, M. I. & Schwenk, J. M. Genetics meets proteomics: perspectives for

large population-based studies. *Nat Rev Genet* **22**, 19–37 (2021).

48. Armingol, E., Officer, A., Harismendy, O. & Lewis, N. E. Deciphering cell–cell interactions

and communication from gene expression. *Nat Rev Genet* **22**, 71–88 (2021).

49. The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids Res*

45, D158–D169 (2017).

514
515
516
517
518   **Figure Legends**
519
Figure 1. Web tool overview and inter-individual correlation structure of established
endocrine proteins. A, Web server structure for user-defined interactions, as well as server and
shiny app implementation scheme for GD-CAT. B, All genes across the 18 metabolic tissues in
310 individuals were correlated with expression of *ADIPOQ* in subcutaneous adipose tissue,
where a qvalue cutoff of q<0.1 showed the strongest enrichments with subcutaneous and muscle
gene expression (pie chart, left). Gene set enrichment analysis (GSEA) was performed using the

bicor coefficient of all genes to *ADIPOQ* using gene ontology biological process annotations and
network construction of top pathways using clusterprofiler, where pathways related to fatty acid
oxidation were observed in adipose (left) and chemotaxis/ECM remodeling in skeletal muscle
(right). B-D, The same qvalue binning, top within-tissue and top peripheral enrichments were
applied to intestinal *GCG* (B), liver *FGF21* (C) and muscle *IL6* (D). For these analyses all 310
individuals (across both sexes) were used and qvalue adjustments calculated using a Benjamini-
Hochberg FDR adjustment.

Figure 2. Tissue-specific contributions to pan-organ gene-gene correlation structure. A
Heatmap showing all the number of gene-gene correlations across tissues which achieve
significance relative to total number of genes in each pathway at biweigth midcorrelation student
pvalue < 1e-3 (left column), pvalue < 1e-6 (left middle column) of BH-corrected qvalue <0.1
(right middle column) or BH-corrected qvalue<0.01 (right column). Within-tissue correlations
are omitted from this analysis. B-D, Genes corresponding to each KEGG pathway shown were
correlated both within and across all other organs where the number of genes which meet each
students pvalue threshold are shown (y-axis). Tissues (x-axis) are rank-ordered by the number of
genes which correlate for hsa04062 − Chemokine signaling pathway at pvalue<0.01 and shown
for other KEGG terms, hsa04640 − Hematopoietic cell lineage (C) and hsa00190 − Oxidative
phosphorylation (D) and additionally pvalue<1e-4 (right side).

Figure 3. Inter-individual transcript correlation structure and network architecture of
liver *PCSK9* and adipose *PNPLA2*. A, distribution of pan-tissue genes correlated with liver
PCSK9 expression (q<0.1), where 93% of genes were within liver (purple). B, Gene ontology
(BP) overrepresentation test for the top 500 hepatic genes correlated with *PCSK9* expression in
liver. C, Undirected network constructed from liver genes (aqua) correlated with *PCSK9*, where
those annotated for "cholesterol biosynthetic process" are colored in red. D-E, over-
representation tests corresponding to the top-correlated genes with adipose (subcutaneous)
PNPLA2 expression residing in adipose (D) or peripherally in skeletal muscle (E). F, Undirected
network constructed from from the strongest correlated subcutaneous adipose tissue (light aqua)
and muscle genes (dark blue) with PNPLA2 (black), where genes corresponding to GO terms
annotated as "fatty acid beta oxidation" or "Muscle contraction" are colored purple or red,
respectively. For these analyses all 310 individuals (across both sexes) were used and qvalue
adjustments calculated using a Benjamini-Hochberg FDR adjustment. Network graphs
generated based in Biweight midcorrelation coefficients, where edges are colored blue for
positive correlations or red for negative correlations. Network edges represent positive (blue)
and negative (red) correlations and the thicknesses are determined by coefficients. They are set
for a range of bicor=0.6 (minimum to include) to bicor=0.99

Figure 4. HMDP tissue- and diet-specific correlations of sex hormone receptors. The top
10 phenotypic traits which correlated to expression of androgen receptor (A), estrogen receptor 1
(B) or estrogen receptor 2 (C) colored by direction in the hybrid mouse diversity panel. Positive
correlations are shown in light blue and negative correlations as sunset orange, where phenotypes
(y-axis) are ordered by significance (x-axis, -log(pvalue) of correlation). Correlations are
segregated by whether sex hormone receptors are expressed by gonadal adipose tissue (left two
columns) in ~100 HMDP strains fed a HFHS diet (left), normal chow diet (left middle) or liver-
expressed receptors fed a HFHS diet (right middle) or normal chow diet (right).

Supplemental Figure 1: Performance across 4 methods of cell-type deconvolution where relative proportions of cells (y-axis) are shown for all cell types annotated in single-cell reference (x-axis) in Liver.

Supplemental Figure 2: Performance across 4 methods of cell-type deconvolution where relative proportions of cells (y-axis) are shown for all cell types annotated in single-cell reference (x-axis) in Heart.

Supplemental Figure 3: Performance across 4 methods of cell-type deconvolution where relative proportions of cells (y-axis) are shown for all cell types annotated in single-cell reference (x-axis) in Skeletal Muscle.

Supplemental Figure 4: Pancreatic *INS* expression correlations across tissues in GTEx were binned according to q<0.1 (top) and corresponding pancreatic GSEA network graph is shown (bottom)

A

Web-based app to query Gene-Derived Correlations Across Tissues (GD-CAT)

B

Pan-tissue correlations (q<0.1)

ADIPOQ (Adipose - Subcutaneous)

Top within organ pathways — Pvalue

Adipose - Subcutaneous

Top peripheral organ pathways — Pvalue

Muscle - Skeletal

C

GCG (Small Intesine - Terminal Illeum)

Small Intestine

Pancreas

D

FGF21 (Liver)

Liver

Adipose (Visceral)

E

IL6 (Muscle - Skeletal)

Muscle - Skeletal

Heart

A
GO term pathway correlation structure across tissues

B
Numer of significant genes correlated pan-tissue per KEGG term

Liver PCSK9

A Pan-tissue correlations

B Top liver pathways

C Local liver network

Subcutaneous Adipose PNPLA2

D Top adipose pathways

E Top muscle pathways

F Adipose-muscle network

Phenotype correlations for hormone receptors in mouse diversity panel

negative correlation · positive correlation