The influence of the larval microbiome on susceptibility to Zika virus is mosquito genotype dependent

Anastasia Accoti[1], Laura C. Multini[1], Babakar Diouf[2], Margaret Becker[1,3], Julia Vulcan[1], Massamba Sylla[4], Dianne aster Y Yap[5], Kamil Khanipov[5], Scott C. Weaver[1,3,6], Mawlouth Diallo[2,3], Alioune Gaye[2,3], Laura B. Dickson[1,3,6,7]

1: Department of Microbiology and Immunology, University of Texas Medical Branch, Galveston, Texas

2: Medical Zoology Unit, Institute Pasteur Dakar, Dakar, Senegal

3: West African Center for Emerging Infectious Diseases, Centers for Research in Emerging Infectious Diseases

4**:** Laboratory Vectors & Parasites, Department of Livestock Sciences and Techniques Sine Saloum University El Hadji Ibrahima NIASS (USSEIN), Kaffrine Campus, Senegal

5: Department of Pharmacology and Toxicology, University of Texas Medical Branch, Galveston, Texas

6: Institute for Human Infections and Immunity, University of Texas Medical Branch, Galveston, Texas

7: Center for Vector-borne and Zoonotic Diseases, University of Texas Medical Branch, Galveston, Texas

Abstract

The microbiome of the mosquito *Aedes aegypti* is largely determined by the environment

and influences mosquito susceptibility for arthropod-borne viruses (arboviruses). Larval

interactions with different bacteria can influence adult *Ae. aegypti* replication of

arboviruses, but little is known about the role that mosquito host genetics play in

determining how larval-bacterial interactions shape *Ae aegypti* susceptibility to

arboviruses. To address this question, we isolated single bacterial isolates and complex

microbiomes from *Ae. aegypti* larvae from various field sites in Senegal. Either single

bacterial isolates or complex microbiomes were added to two different genetic

backgrounds of *Ae. aegypti* in a gnotobiotic larval system. Using 16S amplicon

sequencing we show that similarities in bacterial community structures when given

identical microbiomes between different genetic backgrounds of *Ae. aegypti* was

dependent on the source microbiome, and the abundance of single bacterial taxa

differed between *Ae. aegypti* genotypes. Using single bacterial isolates or the entire

preserved complex microbiome, we tested the ability of specific microbiomes to drive

differences in infection rates for Zika virus in different genetic backgrounds of *Ae.*

aegypti. We observed that the proportion of Zika virus-infected adults was dependent on

the interaction between the larval microbiome and *Ae. aegypti* host genetics. By using

the larval microbiome as a component of the environment, these results demonstrate

that interactions between the *Ae. aegypti* genotype and its environment can influence

Zika virus infection. As *Ae. aegypti* expands and adapts to new environments under

climate change, an understanding of how different genotypes interact with the same

environment will be crucial for implementing arbovirus transmission control strategies.

Introduction

Arthropod-borne viruses (arboviruses) transmitted by mosquitoes represent a major

cause of morbidity and mortality worldwide [1, 2]. The mosquito, *Aedes aegypti* is the

main vector for arboviruses worldwide including dengue (DENV), Zika (ZIKV), yellow

fever (YFV), and chikungunya (CHIKV) viruses. Climate change and a warming world

exacerbate this risk from vector borne diseases [3, 4] and arbovirus epidemics are

poised to be a major threat in sub-Saharan Africa [5]. Given the ongoing and increasing

risk of mosquito-borne viruses, especially in Africa, it is crucial to understand factors that

contribute to their emergence and transmission.

Aedes aegypti demonstrates large phenotypic variability in its interactions with

arboviruses largely driven by genetic and environmental variation. *Aedes aegypti* is

genetically diverse worldwide [6-8] and the ability of *Ae. aegypti* to be an efficient vector

of arboviruses is a partially genetically controlled trait [9]. Variation in vector competence

has been widely documented among different genotypes of *Ae. aegypti* [10-12]. In some

cases, vector competence is dependent on the specific mosquito-virus interactions [13-

15].

Additionally, abiotic (non-living) environmental factors [16-29] and biotic (living)

environmental factors [30-36] are known to contribute to the vector competence of *Ae.*

aegypti. An important biotic ecological parameter influenced by the environment is the

larval microbiome. Globally, *Ae. aegypti* occupies a variety of environments and diverse

larval habitats. Outside Africa, domesticated *Ae. aegypti* oviposits in human artificial

containers such as discarded buckets or cans and tires around human habitats. In

Africa, *Ae. aegypti* will oviposit and develop in a variety of container types ranging from

human artificial containers such as discarded buckets, cans, or tires around human

habitats, to tree holes and rockpools in forested habitats. Larval development sites

represent different microbiomes [37]. The larval microbiome is largely determined by the

environment and is critical for establishing the nutritional status of the mosquito [38].

Importantly, interactions between *Ae. aegypti* larvae and different bacterial isolates drive

variation in DENV susceptibility in adults [37, 38]. However, whether different genotypes

of *Ae. aegypti* interact differently with the same larval microbiome to drive variation in

arbovirus susceptibility is remains unknown.

Here we expand on previous work [37] to determine if the impact of the larval

microbiome on arboviruses susceptibility is dependent on mosquito genotype. Using

single bacterial isolates, we demonstrate that the influence of specific bacterial isolates

on adults ZIKV susceptibility is dependent on mosquito genotype. Using complex

microbiomes isolated from larvae in Senegal, we demonstrate that different genotypes of

Ae. aegypti retain different members of the larval microbiome and that ZIKV

susceptibility is dependent on the specific pairing between *Ae. aegypti* genotype and

complex microbiome. Our results provide empirical evidence that *Ae. aegypti* genotype

by environment interactions drive variation in ZIKV susceptibility. As the earth becomes

warmer and drier and de-forestation and urbanization increase, *Ae. aegypti* will expand

into new environments [39, 40] and exploit different oviposition container types. This is

especially true in Africa where the larval development site is tied to the environment.

Aedes aegypti genotypes accustomed to ovipositing in forest or natural breeding sites

will be forced to adapt and oviposit in urban artificial container types. Understanding how

climate driven changes to the environment will increasingly shape the ability of *Ae.*

aegypti to transmit important human pathogens is required to guide the development

and deployment of control strategies.

Results

Single bacterial Isolates

Previously we observed that larval development in the presence of different single

isolates alters DENV replication in *Ae. aegypti* [37], but it remains unknown if the

influence of larval development in the presence of different single bacterial isolates on

arbovirus infection is consistent across *Ae. aegypti* genotypes. To address this, we

isolated bacteria colonies from larvae collected in Thiés, Senegal. A total of 64 bacteria

were isolated from larvae homogenized in 1X PBS originating from a collection site in

Thies. Three bacteria isolates were chosen (*Serratia spp.* (Bacterial Isolate A),

Chyrseobacterim spp. (Bacterial Isolate B), and *Serratia spp.* (Bacterial Isolate C)) for

further characterization (see Material and Methods for information on bacteria selection).

To understand if larval development in the presence of specific bacterial isolates alters

pupation rates in an *Ae. aegypti* genotype dependent manner, two lines of *Ae. aegypti*

from Senegal with known genetic differences [7], Thiés (THI) and Kedougou (KED),

were exposed to equal amounts of bacteria isolates in gnotobiotic system and the

proportion of pupae was recorded each day. Pupation rates were measured in triplicate

gnotobiotic flask daily from the onset of pupation (day five post hatching) until day 10.

Overall, the THI line of *Ae. aegypti* pupated slower than the KED line (p. value < 0.0001).

In both lines of *Ae. aegypti*, the time it took to reach 50% pupae (PD_{50}) was different

between all three isolates (Figure 1A) (Supplemental Table 1). In both lines the bacterial

isolate that had the slowest pupation rate was Bacterial Isolate A with a PD_{50} of 8.5 days

in the THI line and 7.4 days in KED line. Conversely, the bacterial isolated that resulted

in quickest pupation rate was Bacterial Isolate B where it took 6.9 days in the THI line

and 6.1 days in the KED line. Bacteria C resulted in intermediate pupation rate of 7.4

days in the THI line and 6.7 days in the KED line. Interestingly, Bacteria A and C both

belong to the genus *Serratia* but exert different effects on pupation rates. Overall, we

observed that the PD_{50} was dependent on the mosquito genotype, bacterial isolate, or

the interaction between the two variables (ANOVA on a general linearized model:

mosquito genotype: $p < 0.0001$, bacterial isolate: $p < 0.0001$, mosquito genotype x

bacterial isolate: $p < 0.0001$).

To determine if larval development in the presence of different single isolates influences

ZIKV infection rates in a mosquito genotype dependent manner, the THI and KED line

were exposed to ZIKV following larval development in Bacteria A, Bacteria B, or Bacteria

C. Seven days post exposure to ZIKV, infection rates were determined by detection of

viral RNA by RT-PCR. Infection rates ranged from 20-50% and was not dependent on

the bacteria isolate (p-value = 0.614) or the mosquito genotype (p-value = 0.372), but

was dependent on the interaction between the bacteria isolate and the mosquito

genotype (p-value = 0.010) (Figure 1B).

Complex microbiomes

To establish if the effect of the interaction between mosquito genotype and bacteria

isolate on Zika virus infection can be extended to complex microbiomes collected from

larvae in the field, whole microbiomes from the larvae pools collected in Thiés, Senegal

were introduced to both genotypes of *Ae. aegypti.* We first characterized the bacterial

communities by 16S amplicon sequencing. Out of 80 individual larvae sequenced, a total

of 87 OTUs were found which represent 30 genera after filtering for low abundance

OTUs and OTUs present in the negative controls. Rarefaction curves (Supplemental

Figure 1) show that sufficient sequencing depth was achieved. Principal component

analysis (PCA) was performed on a Bray-Curtis dissimilarity matrix on the larvae from

the same sites to test if the bacterial communities from each collection site that were

introduced to the larvae were in fact different. The community structure differed between

all five of the bacterial communities (Figure 2) (PERMANOVA, p-value = 0.001). The

bacterial community from Site 1 was the most different from the other sites, and Sites 2-

5 showed greater similarity.

To assess if the two genotypes of *Ae. aegypti* acquire and maintain different bacterial

taxa after rearing in an identical bacterial community in a gnotobiotic system, the percent

abundance of different genera was compared between genotypes and collection sites.

The abundance of the top 20 most prevalent genera differed in both genotypes between

the collection sites (Figure 3A). Gnotobiotic larvae harboring the bacterial community

from Sites four or five are most similar between the genotypes. Gnotobiotic larvae

harboring the bacterial community from Sites one, two, and three are most different

between the genotypes. Abundance of specific genera was consistent between

individuals from each treatment (Supplemental Figure 2). To determine if there are

specific genera that are differentially abundant between genotypes across all the

collection sites in the gnotobiotic system, pairwise differential abundances of each genus

were compared between mosquito genotypes. Of the 30 genera identified in this study,

nine genera were more abundant in the genotype from THI, and seven genera were

more abundant in the genotype from KED (Figure 3B). Additionally, the overall bacterial

community structure differed between mosquito genotypes when fed identical

microbiomes. When all sites were analyzed together the overall bacterial community

structure differed between sites and genotypes (PERMANOVA p=0.001) (Supplemental

Figure 3). When each microbiome source was analyzed independently, the community

structure differed between microbiomes from Sites 1, 2, and 3, but not Sites 4, or 5

(Supplemental Figure 4).

To evaluate if the mosquito genotype by larval microbiome interactions on adult

susceptibility to arboviruses extends to complex microbiomes, the two different

genotypes of *Ae. aegypti* were reared in preserved complex microbiomes harvested

from larvae in natural breeding sites in Senegal and challenged with ZIKV. The

proportion of infected individuals was determined seven days post exposure by detecting

viral RNA by RT-PCR. The proportion of infected bodies varied based on the larval

microbiome in which it developed and the mosquito genotype. Specifically, the

proportion of the infected bodies was not dependent on the bacterial community (p-value

= 0.265) or the mosquito genotype (p-value = 0.392), but was dependent on the specific

pairing of larval microbiome and genotype (p-value = 0.017) (Figure 4A). To determine if

this phenotype extends to dissemination titers, the number of infectious particles was

enumerated in the heads seven days post infection by focus forming assay. The titer of

ZIKV in the heads was dependent on the bacteria community (p-value = 0.0415), but

was not dependent on the mosquito genotype (p-value = 0.057) or on the interaction

between mosquito genotype by larval microbiome interaction (p-value = 0.538) (Figure

4B).

Discussion

In this study, we explored the contribution of mosquito genotype and larval microbiome in

driving variation in ZIKV susceptibility. Having previously observed that adult replication of

DENV was dependent on the specific bacteria the mosquito was exposed to during larval

development [37], we sought to determine if the effect of larval gnotobiotic treatment on

arbovirus susceptibility was mosquito genotype dependent. We found that proportion of ZIKV

infected adults is dependent on the interaction between a single bacteria isolate during larval

development and mosquito genotype. When reared in identical complex microbiomes in a

gnotobiotic system, the mosquito genotypes differed in the abundance specific genera. Finally,

we observed that the proportion of infected bodies, but not the head titers, was dependent on

the specific pairing between larval microbiome and mosquito genotype. Instead, the head titers

were dependent on the larval microbiome. Together these data demonstrate that different

genotypes of *Ae. aegypti* interact differently with their larval microbiome and these genotype

dependent interactions are important for ZIKV susceptibility.

In accordance with previously published work [37], we did not see an effect of larval

development in single bacterial isolates on the proportion of infected bodies. In Dickson et al.

2017 [37], the influence of larval development in different bacteria was only observed in the

amount of infectious DENV particles in the head. In the current study, we did not assay

dissemination titers. If done, we might also observe differences in the amount of infectious virus

outside the midgut dependent on which bacterial isolate the larvae developed in. None-the-less,

we still detect an interaction between mosquito genotype and larval microbiome on the

proportion of ZIKV infected bodies. Interestingly, we did observe that the amount of infectious

ZIKV in the head was dependent on complex bacterial community that the larvae were exposed

to in accordance with Dickson et al. 2017 [37], but no interaction between mosquito genotype

and larval microbiome was detected. This might suggest that mosquito genotype by larval

microbiome interactions are only important for infection rates, while the influence of the larval

microbiome on the amount of disseminated virus is largely driven by the larval microbiome

across mosquito genotypes.

The nutrition status and larval microbiome have previously been shown to influence mosquito

fitness [41, 42] and susceptibility to arboviruses [37, 38, 43, 44]. The microbiome is composed

of diverse microorganisms that colonize the mosquito's gut, interacting with the host's metabolic

processes [38, 45, 46] and modulating its immune response [36, 47-49]. Additionally, recent

research has shed light on the critical role of nutrition in determining the outcome of infection in

mosquitoes. In particular, lipid metabolism plays a crucial role in the replication and

dissemination of arboviruses within the mosquito's body [50-55], while amino acids are involved

in the mosquito's interaction with the microbiome [45, 56]. Given that the mosquito's nutritional

status is strongly influences by its microbiome, and that the nutritional status of the mosquito

can influence arbovirus infection, it is not surprising that we observe significant variation in the

rates of ZIKV infection when mosquitoes are reared in different bacterial communities.

Additionally, there have been several mosquito genes identified that impact the microbiome

composition and gut equilibrium [45, 57-60]. These genes can control the overall abundance of

the microbiome or specific taxa, through their involvement in bloodmeal digestion and immune

factors [57] [60]. Moreover, the mosquito microbiome has the potential to influence the

expression of particular genes, which can shape the mosquito's immune response and facilitate

efficient colonization of specific microbes [59]. Although it remains uncertain which mosquito

genes are responsible for the observed phenotypes, it is possible that genetic variation in genes

regulating specific bacterial taxa exist between the two genotypes of *Ae. aegypti* utilized in this

study.

An important finding of this study is that the different genotypes of *Ae. aegypti* larvae harbor a

different abundance of specific taxa when fed identical microbiomes. Multiple studies have

sought to determine if different genotypes or lines of mosquitoes have the same microbiome

when maintained in the same environment. While some studies observed no differences

between the microbiome between genotypes of *Ae. aegypt*i [61], other studies observe different

microbiomes between lines in the same environment [62] and these changes hold up across

microbially diverse environments [63]. While it is assumed that the bacteria the mosquitoes are

exposed to in the same insectary is the same across lines, this is not absolute and it is very

plausible that, during rearing, some larval trays could contain different microbes. By using a

gnotobiotic system, we are ensuring that the different genotypes are receiving the same

microbiome in a highly controlled environment. We observed differences in the abundance of

specific genera between different genetic backgrounds when fed identical microbiomes. This

demonstrates that the mosquito genetic background plays a role in microbiome composition.

Other factors that contribute to microbiome composition are the environmentally available

bacteria [46], and competition between bacteria [64]. Given that the microbiome is remodeled

from larvae to adult development it is likely that that adult microbiomes are different than what is

reported here and we cannot conclude that the adult microbiome is different between the two

genotypes. Nevertheless, this work further highlights the importance of mosquito genotype or

mosquito line in shaping the microbiome.

The results observed in this study could be dependent on the ZIKV isolate used. The isolate of

ZIKV originated from Senegal and has higher infection and dissemination rates than epidemic

isolates of ZIKV [65]. Perhaps we could detect an interaction between mosquito genotype and

larval microbiome on dissemination titers if we had used a different isolate of ZIKV.

Even though these microbiomes originated from larvae collected in Senegal, we cannot make

any conclusions of their field relevance given that we likely partially altered the composition of

the microbiomes compared to larvae in the original habitat through preservation and transport.

How well the microbiomes recapitulate microbiomes in nature is not relevant for this study. We

simply aim to show that different complex bacterial communities can impact ZIKV infection in a

mosquito genotype dependent manner and using microbiomes harvested from larvae in the field

is more relevant than ad hoc mixing single bacterial isolates, even if our complex bacterial

communities are not identical to those in the field. Perhaps if we seeded our gnotobiotic flasks

with the microbiomes from more larvae in accordance with recent studies showing you can

transplant and preserve the microbiome [66], we could make conclusions based on the origin of

microbiomes.

Two of the single bacterial isolates used in this study (Bacteria A and Bacteria C) belong to the

same genus, yet we observed differences in the pupation rate and infection rates between these

two bacteria. The influence of these isolates on pupation rate is consistent across mosquito

genotypes, but the influence of these isolates on ZIKV infection is the opposite direction across

mosquito genotypes. In fact, the influence of these two isolates is likely driving the observed

interaction between bacterial isolate and mosquito genotype. Perhaps genetic differences

between these closely related bacteria species have variable interactions with different

mosquito genotypes which are important for ZIKV infection. This system could provide a highly

tractable system to investigate the mechanism of larval microbiome by mosquito interactions on

arboviruses susceptibility.

Taken together these results show that different genotypes of *Ae. aegypti* interact with their

environment differently to influence ZIKV infection. Future studies should expand on this work to

identify how different microbiomes influence the nutrition status of the mosquito in a mosquito

genotype dependent manner.

Materials and Methods

Bacterial Isolation

Mosquito larvae were collected in two sites (Dixième and Keur Dabo Ndione) in Thiès, Senegal

from plastic drums. A pool of 3 larvae from each site was rinsed in sterile 1X PBS (Phosphate-

buffered saline), incubated in 70% Ethanol for 5 minute (min), and then rinsed in sterile 1X PBS

three times. Next, they were homogenized in 500 µl sterile 1X PBS and had 30% glycerol

added.

A portion of the glycerol stock containing homogenized larvae was plated on Trypticase Soy

Agar (rich media) and incubated for 3 days at 30 °C. Individual colonies were picked from the

plates and used to inoculate 3 ml of LB media, which were shaken at 30°C until bacterial growth

occurred (1 OD) and used to create new glycerol stocks of the individual isolates. DNA was

extracted from each colony with the QIAGEN Dnaesy blood and tissue kit following the

manufacture's protocol. The bacterial DNA was used to amplify the entire 16S region by PCR

[5′-AGAGTTTGATCCTGGCTCAG-3′ (forward) and 5′-AAGGAGGTGATCCAGCCGCA-3′

(reverse)] using Expand High-Fidelity Polymerase (Sigma-Aldrich). The PCR products were

purified using the QIAquick PCR Purification kit (Qiagen), quantified by NanoDrop (NanoDrop

Technologies Inc.), and sequenced by Sanger sequencing (Molecular Genetics Facility at

University of Texas Medical Branch). The sequences were aligned and classified at the genus

level using the SILVA database (www.arb-silva.de/). To standardize the amount of bacteria that

would be introduced into the gnotobiotic system, aliquots of equal amounts of bacteria were

made. The amount of aliquoted bacteria was then quantified by enumerating colony forming unit

(CFU) for each bacterial isolate. To make the aliquots, 200 μl of each bacterial glycerol was

disposed in 200 ml of LB and shaken at 30°C until bacterial growth occurred, then 50 ml was

pelleted by 3000 rpm centrifugation for 15 min. The pellet was washed two times with 50 ml

fresh LB broth. After the second wash, the pellet was resuspended in 50 ml of LB and aliquots

were made by mixing 500 μl of resuspended bacteria an 500 μl of 50% glycerol to make 1ml

aliquots. To quantify the amount of bacteria in each stock, 10 μl was taken from an aliquot and

serially diluted and plated on LB plates. The number of colonies were counted and the number

of CFU/ml were calculated.

Gnotobiotic larvae

To create axenic larvae, *Aedes aegypti* eggs were collected from seventh-generation and

eighth-generation laboratory colonies of Thiés (THI) and Kédougou (KED), respectively, derived

from natural populations from Thiés, Senegal, and Kédougou, Senegal [7]. Genomic data for

these lines exist and they represent different genotypes of *Ae. aegypti* [7]. Eggs were gently

scrapped off the paper into a 50 ml falcon tube. The eggs were sterilized by incubation in 70%

ethanol for 5 min, 3% bleach for 3 min, and 70% ethanol for 5 min. The eggs were then rinsed in

distillated (d) sterile water three times and then they were allowed to hatch in 30 ml of d-water in

a 50ml falcon tube with a 0.2 μM filter lid. Upon hatching, as a control, 10-15 axenic larvae

were transferred to a sterile 25 cm^2 tissue-culture flask containing 15ml of d-water and 50 μl of

sterile fish food (1 g ground fish food flakes per 10 ml d water autoclaved for 20 minutes at 121

°C). These axenic larvae were used as an egg-sterilization control and did not develop past the

1st instar larval stage in accordance with previously published work [67]. Gnotobiotic larvae were

made by distributing 50 $\pm$ 5 (T-75 cm^2 tissue-culture flasks) or 80 $\pm$ 20 (T-150 cm^2 tissue-

culture flasks) axenic larvae to sterile 75 or 150 cm^2 tissue-culture flasks in duplicate or

triplicate containing either 45 or 120 ml of d-sterile water and 1 ml of sterile fish food. For the

single isolate, 5 x10^5 CFUs/ml of washed-bacteria (for details go to bacteria growth section)

was added to each flask. For each bacterial isolate tested, three replicate flasks were used. A

total of three independent experiments was performed. To make gnotobiotic larvae with

complex microbiomes, equal amounts of the glycerol stocks from each collection site (Sites 1-5)

were added to each of two duplicate T-150 flasks. Data represents one experimental replicate

due to availability of field material. Control and gnotobiotic larvae were maintained on 50 (T-75

flask) and 500 (T-150 flask) µl sterile fish food every other day, respectively. Bacterial isolates

were chosen based on the presence of the corresponding genera in the 16S amplicon

sequencing dataset.

16S and metagenomic analysis

To characterize the microbiome in the gnotobiotic larvae seeded with the complex microbiomes,

eight individual larvae were collected from each treatment flask and transferred to a 96-well cell

culture plate. The larvae were surface sterilized in 70% ethanol for 5 min and rinsed 3 times in

sterile water. Next, individual larvae were transferred to 2 ml tubes containing a 5mm grinding

bead and placed in the -80°C freezer until DNA was extracted. Individual mosquitoes were

homogenized for 3 minutes at a 30Hz/s frequency in a TissueLyser II grinder (Qiagen). DNA

extraction of individual larvae was carried out using the QIAamp DNA Kit (Qiagen, Germany)

following the manufacturer's protocol. No-mosquito controls were used for each extraction batch

and included in the sequencing run.

Sequencing libraries for each isolate were generated using universal 16S rRNA V3-V4 region

primers [68] in accordance with Illumina 16S rRNA metagenomic sequencing library protocols.

DNA concentrations of each library were determined by Qubit and equal amounts of DNA from

each barcoded library were pooled prior to sequencing. The samples were barcoded for

multiplexing using Nextera XT Index Kit v2. The pooled libraries were diluted to 4 pM and run on

the Illumina Miseq using a MiSeq Reagent Kit v2 (500-cycles).

To identify known bacteria, sequences were analyzed using the CLC Genomics Workbench

21.0.5 Microbial Genomics Module (CLC MGM). Reads containing nucleotides below the quality

threshold of 0.05 (using the modified Richard Mott algorithm) and those with two or more

unknown nucleotides or sequencing adapters were trimmed out. Reference-based Operational

Taxonomic Unit (OTU) picking was performed using the SILVA SSU v132 97% database [69].

Sequences present in more than one copy but not clustered to the database were placed into

de novo OTUs (97% similarity) and aligned against the reference database with an 80%

similarity threshold to assign the "closest" taxonomical name where possible. Chimeras were

removed from the dataset if the absolute crossover cost was three using a k-mer size of six.

OTUs with a combined abundance of less than two were removed from the analysis. Low

abundance OTUs were removed from the analysis if their combined abundance was below 10

or 0.1% of reads. Differential abundance analysis was performed using CLC MGM at the genus

level to compare the differences between the groups using trimmed mean of M-values. Each

OTU was modeled as a separate generalized linear model, where it is assumed that

abundances follow a negative binomial distribution. The Wald test was used to determine

significance between groups. Multiple comparisons were corrected for with Bonferroni

correction. Tables of differentially abundant taxa are on the complete unfiltered data set

(Supplemental Tables 2).

Abundance profiling was performed using MicrobiomeAnalyst [70, 71]. The analysis parameters

were set so that OTUs had to have a count of at least 10 in 20% of the samples and above 10%

inter-quantile range. Analysis was performed using actual and total sum scale abundances.

Alpha diversity was measured using the observed features to identify the community richness

using Chao1. Statistical significance was calculated using T-test/ANOVA. Beta diversity was

calculated using the Bray-Curtis dissimilarity measure (genus level). Permutational Multivariate

Analysis of Variance (PERMANOVA) analysis was used to measure effect size and significance

on beta diversity for grouping variables [72]. Relative abundance analysis was done in

MicrobiomeAnalyst at the level of genera.

Out of 80 individual larvae sequenced, a total of 1768 OTUs were identified. After filtering, 92

OTUs remain which represent 30 genera. Sequences from three individuals were removed from

the analysis because they did not achieve enough reads, one from Site 1, 4, and 6. After

removing OTUs belonging to the negative control, 87 OTUs remained. These final 87 OTUs

were used for the analysis in the MicrobiomeAnalyst.

Pupal development

To determine the rate of pupation, pupae were counted from the onset of pupation (Day 5) until pupation finished (Day 10) in the same triplicate flasks used for the adult viral challenge assays. The leftover larvae were counted and considered as total amount of individuals for the statistical analysis. For the pupal development, data from two independent experiments was used, each with three internal replicates (three replicate flasks). Graphpad was used to generate a simple logistic regression that computed the day that 50% of larvae pupated (PD_{50}). An ANOVA was run on the summary statistics of the PD_{50} generated from the logistic regression to determine if the PD_{50} was dependent on the bacterial isolate, the mosquito genotype, or an interaction between the two. Multiple comparison by-two-way-ANOVAs were performed to compare the mean PD_{50} between populations (Sidak's test), and between each of three different bacterial treatments within population (Tukey's test) and between each bacterium in the two populations (Sidak's test). Data are a summary of two biological experiments done each time in triplicates. The number of larvae used per experiment along with statistical information associated with each comparison are listed in Supplementary Table 1.

Mosquito infections

Mosquito infection assays was conducted using the ZIKV DAKAR 41524 isolate received from the World Reference Center for Emerging Viruses and Arboviruses at UTMB. After pupation, pupae were transferred to a 1-pint carton box with netting and 10% sucrose solution until adult emergence. After adult emergence, 4 to 5-day-old females were sorted and transferred into a new cup with netting and deprived of sucrose solution for 24 hours and transferred to an Arthropod Containment level 2 facility (ACL-2). Females were offered an artificial blood meal for 15 minutes using the Hemotek system with de-salted pig intestine as the membrane. The infectious blood meal consisted of a 2:1 mixture of defibrinated sheep blood (Colorado Serum

Company) and virus at a final concentration of 1.49 x 10^7 focus-forming units (FFU)/ml. The

blood meal was supplemented with 10 mM adenosine triphosphate (ATP). Prior to addition to

the blood, sodium bicarbonate was mixed with the virus stock at 1% final concentration.

Following exposure to an infectious bloodmeal, fully engorged females were sorted into 1-pint

carton boxes with *ad libidum* access to 10% sucrose solution and kept in an incubator under

controlled conditions (28°C, 12h:12h light: dark cycle). After 7 days of incubation, the head and

body of ZIKV-exposed mosquitoes were separated to determine infection rate (the proportion of

blood-fed mosquitoes with ZIKV-positive body) and dissemination titer (the amount of virus in

the head tissues of ZIKV-infected mosquitoes). To determine the infection rate, female bodies

were homogenized in 200 µl of a crude RNA extraction buffer (10 mM Tris HCl, 50 mM NaCl,

1.25 mM EDTA, fresh 0.35 g/L proteinase K) during two rounds of 3 minutes at a 30Hz/s

frequency in a TissueLyser II grinder (Qiagen). Total RNA was converted into complementary

DNA (cDNA) using M-MLV reverse transcriptase (Invitrogen) and random hexamers, the

reaction was carried out as follows: 10 min at 25°C, 50 min at 37°C, and 15 min 70°C. The

cDNA was amplified by PCR carried out in a 25µl reaction containing 12.5µl of 1x DreamTaq

DNA polymerase (Thermo Fisher Scientific) and 10 µM of each ZIKV primer (forward: 5'-

GTATGGAATGGAGATAAGGCCCA-3', and reverse: 5'-ACCAGCACTGCCATTGATGTGC-3').

Cycling conditions were as follow, 2 min at 95°C, followed by 35 cycles of 30s at 95°C, 30s at

60°C, and 30s at 72°C with a final extension step of 7 min at 72°C. Amplicons were visualized

on a 2% agarose gel. The proportion of ZIKV-infected females was analyzed by binomial logistic

regression as a function of treatment, colony, and their interaction in R.

To determine the dissemination titer, the heads of females with positive ZIKV-infected bodies

were titrated by focus-forming assay in Vero cells. Heads were homogenized individually in 200

µl of Vero cell media (DMEM 1X) supplemented with 2% heat inactivated fetal bovine serum

(FBS) and 1X Antibiotic-Antimycotic (Life Technologies) for 3 minutes at a 30Hz/s frequency in

a TissueLyser II grinder (Qiagen). Vero cells were seeded in 24-well plates and incubated for 24

hours to reach confluency. Each well was inoculated with 200µl of head homogenate in 10-fold

dilutions (from 10^1 to 10^6) and incubated at 37°C (5% CO^2) for 1 hour, rocking every 15 minutes.

Infected cells were overlaid with α-MEM media supplemented with 1.25% carboxymethyl

cellulose, 5% FBS, and 1% Pen-Strep. After three days of incubation at 37°C, infected cells

were fixed with 10 % formaldehyde for at least 1 hour and cells were washed three times in 1X

PBS. Approximately 500 µl of blocking solution (5% w/v non-fat powdered milk in 1X PBS) was

added to each well and the plates were placed on the plate rocker for 30 minutes. The blocking

solution was discarded and 200 µl of primary antibody solution (ZIKV antibody diluted 1:1000 in

blocking solution) was added to each well and plates were placed on plate rocker overnight. The

primary antibody solution was discarded, and plates were washed three times with 1X PBS

again, and 200 µl of secondary antibody (peroxidase-labeled goat anti-mouse IgG human

serum KPL-474-1806) solution (secondary antibody diluted 1:2000 in blocking solution) was

added to each well. Plates were placed on plate rocker for 1 hour. The secondary antibody

solution was discarded, and plates were washed three times with 1X PBS. To develop visible

foci, 100µl of TrueBlue peroxidase substrate (KPL 5510-0050) was added to each well, and

plates were placed on the plate rocker until foci could be seen, around 10 min. Plates were

washed with deionized water and FFU was counted with the help of a light. Focus-forming units

were Log_{10} transformed to represent the concentration of infectious ZIKV particles detected in

Ae. aegypti heads. Head titer data were analyzed by two-way ANOVA as a function of bacterial

treatment, mosquito genotype, and their interaction in R. Infection rate data was analyzed by a

two-way ANOVA on a binomial logistic regress in R.

Acknowledgements:

We would like to thank Jiehua Zhou and Ruimei Yun of the UTMB insectary core. We would

also like to thank Assyatou Gueye and Marieme Gueye for their assistance in field sampling.

We would like to thank Dr. Noah Rose and Dr. Lindy McBride for sharing the mosquito colonies

with us. LBD was supported by UTMB start-up funds and U01AI151801 West African Center for

Emerging Infectious Diseases. AG was supported by U01AI151801 West African Center for

Emerging Infectious Diseases.

Figure 1: Larval exposure to different bacteria isolates alters pupation rate and

ZIKV infection rates. Variation in pupation rate (A) and ZIKV infection rates (B) are

shown for *Ae. aegypti* reared in the presence of single bacteria isolates in a gnotobiotic

system. (A) The PD_{50} (day where approximately 50% of the larvae have pupated) is

shown for two distinct genotypes of *Ae. aegypti* [7] KED and THI reared in the presence

of each bacterial isolate (Bacteria A, Bacteria B, and Bacteria C). Statistical significance

of differences between PD_{50} was determined by multiple comparisons with two-way-

ANOVA and Tukey's test (between bacterial treatments in each mosquito genotype) and

Sidak's test (between bacterial treatments within the same mosquito genotype) and is

designated by the letters. Error bars are shown within each point. (B) The proportion of

KED and THI lines of *Ae. aegypti* with a ZIKV positive body 7 days post-infection

following larval rearing in single bacteria isolates, Bacteria A, Bacteria B, and Bacteria C.

Error bars represent the 95% confidence intervals of the proportions. Data were

analyzed by a two-way ANOVA on a binomial logistic regression (bacteria isolate: p-

value = 0.614, mosquito genotype: p-value = 0.372, bacterial isolate x mosquito

genotype: p-value = 0.010).

● Site 1 ● Site 2 ● Site 3 ● Site 4 ● Site 5

Figure 2: The bacterial community structure differs between the gnotobiotic

larvae. Structure of bacterial communities was determined by deep sequencing the V3-

V4 region of the 16S gene in individual larvae reared in a gnotobiotic system inoculated

with complex microbiomes generated from larvae homogenates collected in Senegal

(Site1-Site5). Bacterial structure is represented by PCoA of a Bray-Curtis dissimilarity

matrix based on mean genera abundance (PERMANOVA p = 0.001).

A)

B)

498   **Figure 3: Abundance of specific taxa differs between lines of gnotobiotic larvae**.

499   (A) The percent abundance of the top 20 most abundant genera is plotted by larval

 treatment (Site1-5) and mosquito line (KED or THI). (B) The log$_2$fold change of bacteria

genera with significant pairwise differences between

KED and THI is plotted. Pairwise differential analysis was performed between larvae

from the KED or THI after receiving identical complex bacterial communities. The

bacteria associated with negative value are the one more likely to be associated to KED

and the one with positive value are the one associated with THI.

Figure 4: Larval development in different complex microbiomes alters ZIKV

infection rates in a mosquito genotype dependent manner. (A) The proportion of

blood-fed *Aedes aegypti* females from the KED and THI lines with a ZIKV-positive body

7 days post-infectious blood meal by RT-PCR following larval rearing in complex

microbiomes, Site1-5. The y-axis indicates the proportion of ZIKV-infected female bodies

and error bars represent the 95% confidence intervals of the proportions. Data were

analyzed by binomial logistic regression as a function of bacterial treatment, mosquito

genotype, and their interaction (bacterial treatment: p-value = 0.265, mosquito genotype:

p-value = 0.392, but was dependent on the specific pairing of bacterial treatment x

517 mosquito genotype: p-value = 0.017. (B) Boxplot showing the dissemination titers of

518 infectious ZIKV particles expressed as the Log10-transformed number of focus-forming

519 units (FFU) per ml detected in the *Ae. aegypti* head seven days post-infectious blood

520 meal. The points represent individuals and the mean is represented by a horizontal line.

521 The error bars represent the 95% confidence interval. Data were analyzed by Two-way

522 ANOVA as a function of bacterial treatment, mosquito genotype, and their interaction

523 (bacterial treatment: p-value = 0.0415, mosquito genotype: p-value = 0.057, bacterial

524 treatment x mosquito genotype: p-value = 0.538).

526 **Supplemental Figures**

528 **Supplemental Figure 1:** Rarefaction curves showing the sequencing depth of each library. The

529 number of species is shown on the Y axis, and the number of sequencing reads is shown on the

530 X axis.

Supplemental Figure 2: The percent abundance of the top 20 most abundant genera is

plotted by larval treatment (Site1-5) and mosquito line (KED or THI) and separated out

by individual.

Supplemental Figure 3: Beta diversity metrics for the *Ae. aegypti* lines in each bacterial

treatment. The dissimilarities between the two different lines of *Ae. aegypti (*KED and THI) in

each of the five different larval microbiomes was analyzed by principal component analysis of

Bray-Curtis dissimilarity index (PERMANOVA, p = 0.001).

Supplemental Figure 4: Beta diversity metrics for the *Ae. aegypti* lines in each bacterial

treatment analyzed separately. The dissimilarities between the two different lines of *Ae. aegypti*

(KED and THI) in each of the five different larval microbiomes was analyzed by principal

component analysis of Bray-Curtis dissimilarity matrixes. P-values from individual

PERMANOVA analysis are shown on the figure.

<u>**References**</u>

1.   Bhatt, S., P.W. Gething, O.J. Brady, J.P. Messina, A.W. Farlow, C.L. Moyes, J.M. Drake,
J.S. Brownstein, A.G. Hoen, O. Sankoh, M.F. Myers, D.B. George, T. Jaenisch, G.R.
Wint, C.P. Simmons, T.W. Scott, J.J. Farrar, and S.I. Hay, *The global distribution and*
burden of dengue. Nature, 2013. **496**(7446): p. 504-7 DOI: 10.1038/nature12060.
2.   Wilder-Smith, A., D.J. Gubler, S.C. Weaver, T.P. Monath, D.L. Heymann, and T.W.
Scott, *Epidemic arboviral diseases: priorities for research and public health.* Lancet
Infect Dis, 2017. **17**(3): p. e101-e106 DOI: 10.1016/S1473-3099(16)30518-7.

3.   Rocklov, J. and R. Dubrow, *Climate change: an enduring challenge for vector-borne*
562   *disease prevention and control.* Nat Immunol, 2020. **21**(5): p. 479-483 DOI:
563   10.1038/s41590-020-0648-y.
564   4.   Robert, M.A., R.C. Christofferson, P.D. Weber, and H.J. Wearing, *Temperature impacts*
565   *on dengue emergence in the United States: Investigating the role of seasonality and*
566   *climate change.* Epidemics, 2019. **28**: p. 100344 DOI: 10.1016/j.epidem.2019.05.003.
567   5.   Mordecai, E.A., S.J. Ryan, J.M. Caldwell, M.M. Shah, and A.D. LaBeaud, *Climate*
568   *change could shift disease burden from malaria to arboviruses in Africa.* Lancet Planet
569   Health, 2020. **4**(9): p. e416-e423 DOI: 10.1016/S2542-5196(20)30178-9.
570   6.   Dickson, L.B., C.L. Campbell, P. Juneja, F.M. Jiggins, M. Sylla, and W.C.t. Black, *Exon-*
571   *Enriched Libraries Reveal Large Genic Differences Between Aedes aegypti from*
572   *Senegal, West Africa, and Populations Outside Africa.* G3 (Bethesda), 2017. **7**(2): p. 571-
573   582 DOI: 10.1534/g3.116.036053.
574   7.   Rose, N.H., M. Sylla, A. Badolo, J. Lutomiah, D. Ayala, O.B. Aribodor, N. Ibe, J. Akorli,
575   S. Otoo, J.P. Mutebi, A.L. Kriete, E.G. Ewing, R. Sang, A. Gloria-Soria, J.R. Powell,
576   R.E. Baker, B.J. White, J.E. Crawford, and C.S. McBride, *Climate and Urbanization*
577   *Drive Mosquito Preference for Humans.* Curr Biol, 2020. **30**(18): p. 3570-3579 e6 DOI:
578   10.1016/j.cub.2020.06.092.
579   8.   Brown, J.E., C.S. McBride, P. Johnson, S. Ritchie, C. Paupy, H. Bossin, J. Lutomiah, I.
580   Fernandez-Salas, A. Ponlawat, A.J. Cornel, W.C.t. Black, N. Gorrochotegui-Escalante, L.
581   Urdaneta-Marquez, M. Sylla, M. Slotman, K.O. Murray, C. Walker, and J.R. Powell,
582   *Worldwide patterns of genetic differentiation imply multiple 'domestications' of Aedes*
583   *aegypti, a major vector of human diseases.* Proc Biol Sci, 2011. **278**(1717): p. 2446-54
584   DOI: 10.1098/rspb.2010.2469.
585   9.   Bosio, C.F., B.J. Beaty, and W.C.t. Black, *Quantitative genetics of vector competence for*
586   *dengue-2 virus in Aedes aegypti.* Am J Trop Med Hyg, 1998. **59**(6): p. 965-70 DOI:
587   10.4269/ajtmh.1998.59.965.
588   10.   Bennett, K.E., K.E. Olson, L. Munoz Mde, I. Fernandez-Salas, J.A. Farfan-Ale, S. Higgs,
589   W.C.t. Black, and B.J. Beaty, *Variation in vector competence for dengue 2 virus among*
590   *24 collections of Aedes aegypti from Mexico and the United States.* Am J Trop Med Hyg,
591   2002. **67**(1): p. 85-92 DOI: 10.4269/ajtmh.2002.67.85.
592   11.   Gubler, D.J., S. Nalim, R. Tan, H. Saipan, and J. Sulianti Saroso, *Variation in*
593   *susceptibility to oral infection with dengue viruses among geographic strains of Aedes*
594   *aegypti.* Am J Trop Med Hyg, 1979. **28**(6): p. 1045-52 DOI:
595   10.4269/ajtmh.1979.28.1045.
596   12.   Vazeille-Falcoz, M., L. Mousson, F. Rodhain, E. Chungue, and A.B. Failloux, *Variation*
597   *in oral susceptibility to dengue type 2 virus of populations of Aedes aegypti from the*
598   *islands of Tahiti and Moorea, French Polynesia.* Am J Trop Med Hyg, 1999. **60**(2): p.
599   292-9 DOI: 10.4269/ajtmh.1999.60.292.
600   13.   Dickson, L.B., I. Sanchez-Vargas, M. Sylla, K. Fleming, and W.C.t. Black, *Vector*
601   *competence in West African Aedes aegypti Is Flavivirus species and genotype dependent.*
602   PLoS Negl Trop Dis, 2014. **8**(10): p. e3153 DOI: 10.1371/journal.pntd.0003153.
603   14.   Lambrechts, L., *Quantitative genetics of Aedes aegypti vector competence for dengue*
604   *viruses: towards a new paradigm?* Trends Parasitol, 2011. **27**(3): p. 111-4 DOI:
605   10.1016/j.pt.2010.12.001.

606 15. Lambrechts, L., C. Chevillon, R.G. Albright, B. Thaisomboonsuk, J.H. Richardson, R.G.
607   Jarman, and T.W. Scott, *Genetic specificity and potential for local adaptation between*
608   *dengue viruses and mosquito vectors.* BMC Evol Biol, 2009. **9**: p. 160 DOI:
609   10.1186/1471-2148-9-160.
610 16. Westbrook, C.J., M.H. Reiskind, K.N. Pesko, K.E. Greene, and L.P. Lounibos, *Larval*
611   *environmental temperature and the susceptibility of Aedes albopictus Skuse (Diptera:*
612   *Culicidae) to Chikungunya virus.* Vector Borne Zoonotic Dis, 2010. **10**(3): p. 241-7 DOI:
613   10.1089/vbz.2009.0035.
614 17. Alto, B.W. and D. Bettinardi, *Temperature and dengue virus infection in mosquitoes:*
615   *independent effects on the immature and adult stages.* Am J Trop Med Hyg, 2013. **88**(3):
616   p. 497-505 DOI: 10.4269/ajtmh.12-0421.
617 18. Alto, B.W., L.P. Lounibos, S. Higgs, and S.A. Juliano, *Larval Competition Differentially*
618   *Affects Arbovirus Infection in Aedes Mosquitoes.* Ecology, 2005. **86**(12): p. 3279-3288
619   DOI: 10.1890/05-0209.
620 19. Alto, B.W., L.P. Lounibos, C.N. Mores, and M.H. Reiskind, *Larval competition alters*
621   *susceptibility of adult Aedes mosquitoes to dengue infection.* Proc Biol Sci, 2008.
622   **275**(1633): p. 463-71 DOI: 10.1098/rspb.2007.1497.
623 20. Muturi, E.J., M. Blackshear, Jr., and A. Montgomery, *Temperature and density-*
624   *dependent effects of larval environment on Aedes aegypti competence for an alphavirus.* J
625   Vector Ecol, 2012. **37**(1): p. 154-61 DOI: 10.1111/j.1948-7134.2012.00212.x.
626 21. Telang, A., A.A. Qayum, A. Parker, B.R. Sacchetta, and G.R. Byrnes, *Larval nutritional*
627   *stress affects vector immune traits in adult yellow fever mosquito Aedes aegypti*
628   *(Stegomyia aegypti).* Med Vet Entomol, 2012. **26**(3): p. 271-81 DOI: 10.1111/j.1365-
629   2915.2011.00993.x.
630 22. Joy, T.K., A.J. Arik, V. Corby-Harris, A.A. Johnson, and M.A. Riehle, *The impact of*
631   *larval and adult dietary restriction on lifespan, reproduction and growth in the mosquito*
632   *Aedes aegypti.* Exp Gerontol, 2010. **45**(9): p. 685-90 DOI: 10.1016/j.exger.2010.04.009.
633 23. Takken, W., R.C. Smallegange, A.J. Vigneau, V. Johnston, M. Brown, A.J. Mordue-
634   Luntz, and P.F. Billingsley, *Larval nutrition differentially affects adult fitness and*
635   *Plasmodium development in the malaria vectors Anopheles gambiae and Anopheles*
636   *stephensi.* Parasit Vectors, 2013. **6**(1): p. 345 DOI: 10.1186/1756-3305-6-345.
637 24. Moller-Jacobs, L.L., C.C. Murdock, and M.B. Thomas, *Capacity of mosquitoes to*
638   *transmit malaria depends on larval environment.* Parasit Vectors, 2014. **7**: p. 593 DOI:
639   10.1186/s13071-014-0593-4.
640 25. Paaijmans, K.P., S. Huijben, A.K. Githeko, and W. Takken, *Competitive interactions*
641   *between larvae of the malaria mosquitoes Anopheles arabiensis and Anopheles gambiae*
642   *under semi-field conditions in western Kenya.* Acta Trop, 2009. **109**(2): p. 124-30 DOI:
643   10.1016/j.actatropica.2008.07.010.
644 26. Ng'habi, K.R., B. John, G. Nkwengulila, B.G. Knols, G.F. Killeen, and H.M. Ferguson,
645   *Effect of larval crowding on mating competitiveness of Anopheles gambiae mosquitoes.*
646   Malar J, 2005. **4**: p. 49 DOI: 10.1186/1475-2875-4-49.
647 27. Pfaehler, O., D.O. Oulo, L.C. Gouagna, J. Githure, and P.M. Guerin, *Influence of soil*
648   *quality in the larval habitat on development of Anopheles gambiae Giles.* J Vector Ecol,
649   2006. **31**(2): p. 400-5 DOI: 10.3376/1081-1710(2006)31[400:iosqit]2.0.co;2.

28.   Okech, B.A., L.C. Gouagna, G. Yan, J.I. Githure, and J.C. Beier, *Larval habitats of*
Anopheles gambiae s.s. (Diptera: Culicidae) influences vector competence to
Plasmodium falciparum parasites. Malar J, 2007. **6**: p. 50 DOI: 10.1186/1475-2875-6-50.
29.   Roux, O., A. Vantaux, B. Roche, K.B. Yameogo, K.R. Dabire, A. Diabate, F. Simard,
and T. Lefevre, *Evidence for carry-over effects of predator exposure on pathogen*
transmission potential. Proc Biol Sci, 2015. **282**(1821): p. 20152430 DOI:
10.1098/rspb.2015.2430.
30.   Minard, G., P. Mavingui, and C.V. Moro, *Diversity and function of bacterial microbiota*
in the mosquito holobiont. Parasit Vectors, 2013. **6**: p. 146 DOI: 10.1186/1756-3305-6-
146.
31.   Douglas, A.E., *Multiorganismal insects: diversity and function of resident*
microorganisms. Annu Rev Entomol, 2015. **60**: p. 17-34 DOI: 10.1146/annurev-ento-
010814-020822.
32.   Hegde, S., J.L. Rasgon, and G.L. Hughes, *The microbiome modulates arbovirus*
transmission in mosquitoes. Curr Opin Virol, 2015. **15**: p. 97-102 DOI:
10.1016/j.coviro.2015.08.011.
33.   Bian, G., Y. Xu, P. Lu, Y. Xie, and Z. Xi, *The endosymbiotic bacterium Wolbachia*
induces resistance to dengue virus in Aedes aegypti. PLoS Pathog, 2010. **6**(4): p.
e1000833 DOI: 10.1371/journal.ppat.1000833.
34.   Cirimotich, C.M., J.L. Ramirez, and G. Dimopoulos, *Native microbiota shape insect*
vector competence for human pathogens. Cell Host Microbe, 2011. **10**(4): p. 307-10 DOI:
10.1016/j.chom.2011.09.006.
35.   Ramirez, J.L., S.M. Short, A.C. Bahia, R.G. Saraiva, Y. Dong, S. Kang, A. Tripathi, G.
Mlambo, and G. Dimopoulos, *Chromobacterium Csp_P reduces malaria and dengue*
infection in vector mosquitoes and has entomopathogenic and in vitro anti-pathogen
activities. PLoS Pathog, 2014. **10**(10): p. e1004398 DOI: 10.1371/journal.ppat.1004398.
36.   Ramirez, J.L., J. Souza-Neto, R. Torres Cosme, J. Rovira, A. Ortiz, J.M. Pascale, and G.
Dimopoulos, *Reciprocal tripartite interactions between the Aedes aegypti midgut*
microbiota, innate immune system and dengue virus influences vector competence. PLoS
Negl Trop Dis, 2012. **6**(3): p. e1561 DOI: 10.1371/journal.pntd.0001561.
37.   Dickson, L.B., D. Jiolle, G. Minard, I. Moltini-Conclois, S. Volant, A. Ghozlane, C.
Bouchier, D. Ayala, C. Paupy, C.V. Moro, and L. Lambrechts, *Carryover effects of larval*
exposure to different environmental bacteria drive adult trait variation in a mosquito
vector. Sci Adv, 2017. **3**(8): p. e1700585 DOI: 10.1126/sciadv.1700585.
38.   Giraud, E.V., H.; Legendre, R.; Sismeiro, O.; Aubry, F.; Dabo ,S.; Dickson, L.B.;
Valiente Moro C.; Louis Lambrechts, L., *Mosquito-bacteria interactions during larval*
development trigger metabolic changes with carry-over effects on adult fitness. bioRxiv,
2021 DOI: doi: https://doi.org/10.1101/2021.05.20.444942.
39.   Ryan, S.J., C.J. Carlson, E.A. Mordecai, and L.R. Johnson, *Global expansion and*
redistribution of Aedes-borne virus transmission risk with climate change. PLoS Negl
Trop Dis, 2019. **13**(3): p. e0007213 DOI: 10.1371/journal.pntd.0007213.
40.   Iwamura, T., A. Guzman-Holst, and K.A. Murray, *Accelerating invasion potential of*
disease vector Aedes aegypti under climate change. Nat Commun, 2020. **11**(1): p. 2130
DOI: 10.1038/s41467-020-16010-4.

41. MacLeod, H.J., G. Dimopoulos, and S.M. Short, *Larval Diet Abundance Influences Size*
and Composition of the Midgut Microbiota of Aedes aegypti Mosquitoes. Front
Microbiol, 2021. **12**: p. 645362 DOI: 10.3389/fmicb.2021.645362.
42. Martinson, V.G. and M.R. Strand, *Diet-Microbiota Interactions Alter Mosquito*
Development. Front Microbiol, 2021. **12**: p. 650743 DOI: 10.3389/fmicb.2021.650743.
43. Louie, W. and L.L. Coffey, *Microbial Composition in Larval Water Enhances Aedes*
aegypti Development but Reduces Transmissibility of Zika Virus. mSphere, 2021. **6**(6): p.
e0068721 DOI: 10.1128/msphere.00687-21.
44. Herd, C.S., D.G. Grant, J. Lin, and A.W.E. Franz, *Starvation at the larval stage increases*
the vector competence of Aedes aegypti females for Zika virus. PLoS Negl Trop Dis,
2021. **15**(11): p. e0010003 DOI: 10.1371/journal.pntd.0010003.
45. Short, S.M., E.F. Mongodin, H.J. MacLeod, O.A.C. Talyuli, and G. Dimopoulos, *Amino*
acid metabolic signaling influences Aedes aegypti midgut microbiome variability. PLoS
Negl Trop Dis, 2017. **11**(7): p. e0005677 DOI: 10.1371/journal.pntd.0005677.
46. Coon, K.L., L. Valzania, D.A. McKinney, K.J. Vogel, M.R. Brown, and M.R. Strand,
Bacteria-mediated hypoxia functions as a signal for mosquito development. Proc Natl
Acad Sci U S A, 2017. **114**(27): p. E5362-E5369 DOI: 10.1073/pnas.1702983114.
47. Hyde, J., M.A. Correa, G.L. Hughes, B. Steven, and D.E. Brackney, *Limited influence of*
the microbiome on the transcriptional profile of female Aedes aegypti mosquitoes. Sci
Rep, 2020. **10**(1): p. 10880 DOI: 10.1038/s41598-020-67811-y.
48. Barletta, A.B., M.C. Nascimento-Silva, O.A. Talyuli, J.H. Oliveira, L.O. Pereira, P.L.
Oliveira, and M.H. Sorgine, *Microbiota activates IMD pathway and limits Sindbis*
infection in Aedes aegypti. Parasit Vectors, 2017. **10**(1): p. 103 DOI: 10.1186/s13071-
017-2040-9.
49. Gabrieli, P., S. Caccia, I. Varotto-Boccazzi, I. Arnoldi, G. Barbieri, F. Comandatore, and
S. Epis, *Mosquito Trilogy: Microbiota, Immunity and Pathogens, and Their Implications*
for the Control of Disease Transmission. Front Microbiol, 2021. **12**: p. 630438 DOI:
10.3389/fmicb.2021.630438.
50. Koh, C., M.N. Islam, Y.H. Ye, N. Chotiwan, B. Graham, J.T. Belisle, K.A. Kouremenos,
S. Dayalan, D.L. Tull, S. Klatt, R. Perera, and E.A. McGraw, *Dengue virus dominates*
lipid metabolism modulations in Wolbachia-coinfected Aedes aegypti. Commun Biol,
2020. **3**(1): p. 518 DOI: 10.1038/s42003-020-01254-z.
51. Barletta, A.B., L.R. Alves, M.C. Silva, S. Sim, G. Dimopoulos, S. Liechocki, C.M.
Maya-Monteiro, and M.H. Sorgine, *Emerging role of lipid droplets in Aedes aegypti*
immune response against bacteria and Dengue virus. Sci Rep, 2016. **6**: p. 19928 DOI:
10.1038/srep19928.
52. Vial, T., G. Marti, D. Misse, and J. Pompon, *Lipid Interactions Between Flaviviruses and*
Mosquito Vectors. Front Physiol, 2021. **12**: p. 763195 DOI: 10.3389/fphys.2021.763195.
53. Marten, A.D., C.T. Tift, M.O. Tree, J. Bakke, and M.J. Conway, *Chronic depletion of*
vertebrate lipids in Aedes aegypti cells dysregulates lipid metabolism and inhibits innate
immunity without altering dengue infectivity. PLoS Negl Trop Dis, 2022. **16**(10): p.
e0010890 DOI: 10.1371/journal.pntd.0010890.
54. Vial, T., W.L. Tan, E. Deharo, D. Misse, G. Marti, and J. Pompon, *Mosquito*
metabolomics reveal that dengue virus replication requires phospholipid reconfiguration
via the remodeling cycle. Proc Natl Acad Sci U S A, 2020. **117**(44): p. 27627-27636
DOI: 10.1073/pnas.2015095117.

55. Chotiwan, N., B.G. Andre, I. Sanchez-Vargas, M.N. Islam, J.M. Grabowski, A. Hopf-
Jannasch, E. Gough, E. Nakayasu, C.D. Blair, J.T. Belisle, C.A. Hill, R.J. Kuhn, and R.
Perera, *Dynamic remodeling of lipids coincides with dengue virus replication in the*
midgut of Aedes aegypti mosquitoes. PLoS Pathog, 2018. **14**(2): p. e1006853 DOI:
10.1371/journal.ppat.1006853.
56. Caragata, E.P., E. Rances, S.L. O'Neill, and E.A. McGraw, *Competition for amino acids*
between Wolbachia and the mosquito host, Aedes aegypti. Microb Ecol, 2014. **67**(1): p.
205-18 DOI: 10.1007/s00248-013-0339-4.
57. Mitri, C., E. Bischoff, E. Belda Cuesta, S. Volant, A. Ghozlane, K. Eiglmeier, I. Holm, C.
Dieme, E. Brito-Fravallo, W.M. Guelbeogo, N. Sagnon, M.M. Riehle, and K.D. Vernick,
Leucine-Rich Immune Factor APL1 Is Associated With Specific Modulation of Enteric
Microbiome Taxa in the Asian Malaria Mosquito Anopheles stephensi. Front Microbiol,
2020. **11**: p. 306 DOI: 10.3389/fmicb.2020.00306.
58. Xiao, X., L. Yang, X. Pang, R. Zhang, Y. Zhu, P. Wang, G. Gao, and G. Cheng, *A Mesh-*
Duox pathway regulates homeostasis in the insect gut. Nat Microbiol, 2017. **2**: p. 17020
DOI: 10.1038/nmicrobiol.2017.20.
59. Pang, X., X. Xiao, Y. Liu, R. Zhang, J. Liu, Q. Liu, P. Wang, and G. Cheng, *Mosquito C-*
type lectins maintain gut microbiome homeostasis. Nat Microbiol, 2016. **1**: p. 16023
DOI: 10.1038/nmicrobiol.2016.23.
60. Zhao, B., K.J. Lucas, T.T. Saha, J. Ha, L. Ling, V.A. Kokoza, S. Roy, and A.S. Raikhel,
MicroRNA-275 targets sarco/endoplasmic reticulum Ca2+ adenosine triphosphatase
(SERCA) to control key functions in the mosquito gut. PLoS Genet, 2017. **13**(8): p.
e1006943 DOI: 10.1371/journal.pgen.1006943.
61. Dickson, L.B., A. Ghozlane, S. Volant, C. Bouchier, L. Ma, A. Vega-Rua, I. Dusfour, D.
Jiolle, C. Paupy, M.N. Mayanja, A. Kohl, J.J. Lutwama, V. Duong, and L. Lambrechts,
Diverse laboratory colonies of Aedes aegypti harbor the same adult midgut bacterial
microbiome. Parasit Vectors, 2018. **11**(1): p. 207 DOI: 10.1186/s13071-018-2780-1.
62. Kozlova, E.V., S. Hegde, C.M. Roundy, G. Golovko, M.A. Saldana, C.E. Hart, E.R.
Anderson, E.A. Hornett, K. Khanipov, V.L. Popov, M. Pimenova, Y. Zhou, Y. Fovanov,
S.C. Weaver, A.L. Routh, E. Heinz, and G.L. Hughes, *Microbial interactions in the*
mosquito gut determine Serratia colonization and blood-feeding propensity. ISME J,
2021. **15**(1): p. 93-108 DOI: 10.1038/s41396-020-00763-3.
63. Accoti, A.Q., Shannon; Vulcan, Julia; Cansado-Utrilla, Cintia; Anderson, Enyia R.;
Alsing, Jessica; Narra, Hema P.; Khanipov, Kamil; Hughes, Grant L.; Dickson, Laura B.
, *Microbiome variability of mosquito lines is consistent over time and across*
environments. bioRxiv, 2023 DOI: https://doi.org/10.1101/2023.04.17.537119.
64. Hegde, S., K. Khanipov, L. Albayrak, G. Golovko, M. Pimenova, M.A. Saldana, M.M.
Rojas, E.A. Hornett, G.C. Motl, C.L. Fredregill, J.A. Dennett, M. Debboun, Y. Fofanov,
and G.L. Hughes, *Microbiome Interaction Networks and Community Structure From*
Laboratory-Reared and Field-Collected Aedes aegypti, Aedes albopictus, and Culex
quinquefasciatus Mosquito Vectors. Front Microbiol, 2018. **9**: p. 2160 DOI:
10.3389/fmicb.2018.02160.
65. Roundy, C.M., S.R. Azar, S.L. Rossi, J.H. Huang, G. Leal, R. Yun, I. Fernandez-Salas,
C.J. Vitek, I.A. Paploski, U. Kitron, G.S. Ribeiro, K.A. Hanley, S.C. Weaver, and N.
Vasilakis, *Variation in Aedes aegypti Mosquito Competence for Zika Virus Transmission.*
Emerg Infect Dis, 2017. **23**(4): p. 625-632 DOI: 10.3201/eid2304.161484.

66. Coon, K.L., S. Hegde, and G.L. Hughes, *Interspecies microbiome transplantation*
recapitulates microbial acquisition in mosquitoes. Microbiome, 2022. **10**(1): p. 58 DOI:
10.1186/s40168-022-01256-5.
67. Coon, K.L., K.J. Vogel, M.R. Brown, and M.R. Strand, *Mosquitoes rely on their gut*
microbiota for development. Mol Ecol, 2014. **23**(11): p. 2727-39 DOI:
10.1111/mec.12771.
68. Klindworth, A., E. Pruesse, T. Schweer, J. Peplies, C. Quast, M. Horn, and F.O.
Glockner, *Evaluation of general 16S ribosomal RNA gene PCR primers for classical and*
next-generation sequencing-based diversity studies. Nucleic Acids Res, 2013. **41**(1): p.
e1 DOI: 10.1093/nar/gks808.
69. Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F.O.
Glockner, *The SILVA ribosomal RNA gene database project: improved data processing*
and web-based tools. Nucleic Acids Res, 2013. **41**(Database issue): p. D590-6 DOI:
10.1093/nar/gks1219.
70. Dhariwal, A., J. Chong, S. Habib, I.L. King, L.B. Agellon, and J. Xia,
MicrobiomeAnalyst: a web-based tool for comprehensive statistical, visual and meta-
analysis of microbiome data. Nucleic Acids Res, 2017. **45**(W1): p. W180-W188 DOI:
10.1093/nar/gkx295.
71. Chong, J., P. Liu, G. Zhou, and J. Xia, *Using MicrobiomeAnalyst for comprehensive*
statistical, functional, and meta-analysis of microbiome data. Nat Protoc, 2020. **15**(3): p.
799-821 DOI: 10.1038/s41596-019-0264-1.
72. Anderson, M.J. and J. Santana-Garcon, *Measures of precision for dissimilarity-based*
multivariate analysis of ecological communities. Ecol Lett, 2015. **18**(1): p. 66-73 DOI:
10.1111/ele.12385.
810