Carbonic anhydrase is a novel diagnostic biomarker for the detection of the parasitic

trematode *Schistosoma mansoni* in the intermediate snail host, *Biomphalaria glabrata*

Simone Parn[1], Gabriela Lewis[1], and Matty Knight[1,2*]

1. Division of Science & Mathematics, University of the District of Columbia, 4200 Connecticut Ave.

NW Washington, D.C. 20008, USA

2.Department of Microbiology, Immunology & Tropical Medicine, Research Center for

Neglected Diseases of Poverty, School of Medicine & Health Sciences, The George Washington

University Ross Hall, 2300 I Street, NW, Washington, DC 20037, USA

*Corresponding author

Name: Professor Matty Knight

Email: mathilde.knight@udc.edu; matty_knight@gwu.edu

ABSTRACT

Schistosomiasis is a major public health concern worldwide. Although praziquantel is currently available as the only treatment option for schistosomiasis, the absence of reliable diagnostic and prognostic tools highlights the need for the identification and characterization of new drug targets. Recently, we identified the *B. glabrata* homolog (accession number XP_013075832.1) of human CAXIV, showing 37% amino acid sequence identity, from a BLAST search in NCBI. CA's are metalloenzymes that catalyze the reversible hydration/dehydration of CO_2/HCO_3. These enzymes are associated with many physiological processes, and their role in tumorigenesis has been widely implicated. CAs create an acidic extracellular environment that facilitates the survival, metastasis, and growth of cancer cells. In this study, we investigated CA as a potential biomarker for infection in *B. glabrata* snails. We analysed the expression of the *B. glabrata* CA encoding transcript in juvenile susceptible and resistant snails, with and without exposure to *S. mansoni*. Our results showed that the expression of the CA encoding transcript was upregulated during early and prolonged infection susceptible snails (BBO2 and NMRI), but not in the resistant BS-90. Notably, sodium salicylate, a form of aspirin, inhibited the expression of CA, post-exposure, to the parasite. These data show that the expression of CA in the snail-schistosome interaction, represents a novel biomarker for the detection of schistosomes in the snail host. Moreover, this model system may aid in understanding the role of CA in the metastasis and proliferation of cancer cells. Further studies are needed to explore the potential of CA as a biomarker for infection in other schistosomiasis-causing parasites, including *S. japonicum* and *S. haematobium.*

Keywords: carbonic anhydrase, infection, biomarker*,* salicylic acid (aspirin), *Schistosoma mansoni*, *Biomphalaria glabrata.*

INTRODUCTION

Schistosomiasis is a neglected tropical disease which is estimated to impact more than 600 million people in the world each year [1]. According to the Centers for Disease Control and Prevention, it is the second most persistent parasitic disease after malaria [2]. While schistosomiasis is mostly transmitted in sub-Saharan Africa, cases of its spread to Europe have been extensively reported [3]. The parasites that cause schistosomiasis utilize the freshwater gastropod snails as obligate hosts for the development of its asexual stages. Contact with contaminated water inadvertently leads to infection by skin penetration with infectious cercaria that are released into the water from infected snails. After penetrating the skin, cercariae lose their tail, transforming into schistosomula that develop into mature adult male and female worms that inhabit the vasculature where they mate in perpetuity to produce eggs that cause widespread chronic morbidity and infertility among human populations [4]. Specifically, infections caused by the species *Schistosoma haematobium* are known to lead to bladder cancer and female genital schistosomiasis [5]. As part of a prevention program, the World Health Organization (WHO) has proposed that school-aged children aged 5-15 years, who face the highest infection rates, receive initial parasitological assessment [6]. Additionally, the WHO estimates that by 2030, schistosomiasis will be eliminated as a public health concern [7]. Although several attempts have been made toward effective preventative and therapeutic measures to control the transmission of schistosomiasis, long-term reduction remains elusive. To combat the widespread disease, better diagnostic and prognostic tools, especially biomarkers to track the presence of larval parasites in the snail are of immediate necessity.

The high impact of this disease on global health burden has promoted scientific efforts to tackle the complex mechanisms underlaying successful schistosome infection. The freshwater snail *Biomphalaria glabrata,* an obligate intermediate host for the parasitic trematode *Schistosoma mansoni*, has been studied for decades at the molecular level to understand its interaction with the

parasite. The genome characterization of the snail *B. glabrata* has allowed for investigation of the

molecular determinants of the snail relationship with the schistosome [8]. Studies of comparative

immunology have enabled scientists to identify genes and pathways involved in the parasite

development [9,10]. Additionally, differential gene expression studies have led to the identification of

several cancer-and stress-related transcripts that are elevated early and significantly in the

susceptible *B. glabrata* snail stocks compared to their resistant counterparts [11].

Recently, Carbonic Anhydrases (CAs) have been identified as potential targets in the development

of novel drugs for parasitic diseases [12,13]. CAs are a family of zinc-containing enzymes,

responsible for the reversible conversion of CO_2 into bicarbonate and protons [14]. In humans, CAs

exist in at least 15 isoforms and differ based on their enzymatic activity and cellular location: hCA I,

II, III, VII, and XIII reside in the cytosol; hCA IV, IX, XII, and XIV are associated with the cell

membrane; hCA VA, and VB are mitochondrial; and hCAVI is found in saliva. These enzymes are

important for pH and CO_2 homeostasis regulation, and for other biosynthesis processes, such as

lipogenesis, gluconeogenesis, ureagenesis, and calcification [15]. Aberrant expression of some CA

isoforms can lead to pathogenic outcomes, like carcinogenesis, obesity, and epilepsy [16]. Notably,

the expression of carbonic anhydrase IX is widely described in several hypoxic solid tumors

[17,18,19]. In the acidic microenvironment, CAIX allows tumor cells to adapt by seizing the high

CO_2 in the form of hydrogen and bicarbonate ions.

In this study, we investigated the expression of carbonic anhydrase transcript in susceptible and

resistant *B. glabrata* snail stocks in response to early and prolonged exposure to the parasite *S.*

mansoni. Given CA's involvement in tumor progression and its similarity to the human enzyme, we

hypothesized that the snail/schistosome relationship could mimic the activity of human CA in cancer

cells. Here, we show that CA serves as a potentially useful biomarker for detecting schistosomiasis

infection in infected snails, with ability to differentiate disease from normal unaffected snails.

Furthermore, we found that sodium salicylate, a water-soluble form of aspirin, inhibits the CA

transcript in susceptible snail stocks. These findings provide insight into the potential role of CA in

the development and progression of schistosomiasis and its impending as a therapeutic target.

MATERIALS AND METHODS

Biomphalaria glabrata **Stocks**

The *Biomphalaria glabrata* juvenile (3-5 mm in diameter) susceptible (BBO2) and parasite resistant

(BS90) snails were utilized in the study. Snails were maintained in aquaria in de-aerated water and

fed with romaine lettuce twice a week.

Snail exposure to *S. mansoni* miracidia

Miracidia hatched eggs isolated from 7 weeks infected mice livers were collected from the

Biomedical Research Institute (Rockville, MD). Both BBO2 and BS90 snails were exposed

individually to 10-12 *S. mansoni* miracidia in 1.0 ml de-aerated water in a 6-well microtiter plate at

room temperature. The snails were exposed for different time points at 0 min, 30 min, 1hr, 2hr and

4hr. Following exposure, snails were either processed in 500 µl of RNAzol immediately or kept at -

80°C until needed.

BLAST analysis and primer design

Several bioinformatics approaches were used to test the working hypothesis. First, we searched for

B. glabrata homologs of human related Carbonic Anhydrase encoding transcript using the protein

database Uniprot (www.uniprt.org). The amino acid or FASTA sequences were individually

deposited into the Basic Local Alignment Search Tool (BLAST) [20] to identify the *B. glabrata*

homolog. The analysis was followed by a SMART BLAST analysis to validate the identity of the *B.*

glabrata homolog and evolutionary relatedness to other vertebrate and invertebrate organisms'

amino acid sequence, including human. We then used the mRNA transcript of corresponding *B.*

glabrata protein to design gene specific primers [21]. Oligonucleotide Primers for qPCR were

obtained from Eurofins Genomics (Louisville 13 KY, 40204).

RNA extraction

Total RNA was extracted from whole juvenile snails (unexposed and exposed) using the RNAzol

method, according to the manufacturer's instructions (Sigma- Aldrich, USA). Briefly, 500 µl of

RNAzol reagent was added to the snail samples and homogenized using a pestle and a motor.

Homogenate was mixed with 200 µl of sterile dH_2O, vortexed and incubated at room temperature

for 10 min before centrifugation at 13, 000 x g for 10 min at 26°C. Ethanol (75%) was added to the

recovered supernatant after centrifugation and samples were incubated on ice for 10 min before

another centrifugation at 13, 000 x g for 10 min at 4°C. Supernatant was removed while RNA pellet

was left undisturbed. The pellet was washed with 70% ethanol and centrifuged at 13,000x g for 10

min at 4°C. Alcohol wash was removed, and pellet was air-dried. RNA pellets were dissolved in 20

µl of RNase-free dH_2O. RNA yield was determined by quantitation under UV absorbance at

wavelength 260 nm using the NanoDrop 1000 spectrophotometer (Thermo Scientific). RNA

samples were used either immediately or kept frozen at -80°C until needed to synthesize cDNA.

Trace amounts of contaminating genomic DNA were removed by treating all RNA samples with

RNAse-free DNase I prior to use for cDNA synthesis according to the manufacturer's instructions

(Promega, WI).

cDNA synthesis and qualitative RT-PCR

Complementary DNA was synthesized from total RNA using a cDNA synthesis kit (TermoFisher

Scientific). Briefly, a master mix was prepared using 5X Reverse Transcriptase Buffer, 100 mM

DTT, RNase OUT, Oligo DT, 100 mM dNTPs and Reverse transcriptase, which was aliquoted to

labelled 1.5ml Eppendorf tubes. 1 ul of RNA (500ng) was used per reaction. The tubes were

incubated at 42°C water bath for 2 hours and DNA concentrations were measured using NanoDrop

at the wavelength of 260 nm. DNA samples were then diluted to 200ng each for qualitative PCR

reaction. Experimental and reference (myoglobin) primers were diluted to 15 uM. In addition to

diluted primers, the reaction was synthesized using nuclease free water, Taq Polymerase, dNTPs

and Mg^{+2} 10X Buffer in a preheated thermocycler for 2 hours. Amplified DNA transcripts were

analyzed by 1.2% TBE- agarose electrophoresis.

Real-time quantitative RT-PCR

The real-time PCR measurement of cDNAs was performed using PowerUp SYBR Green Master

Mix (Applied Biosystems) and normalized to the expression of myoglobin as standard for the

housekeeping gene. The primers were: Carbonic Anhydrase forward: *caggagcagtttaggaagggc*;

Carbonic Anhydrase reverse: *tcggctcaaaactcacctcc*; Myoglobin forward: *gatgttcgccaatgttccc*;

Myoglobin reverse: *agcgatcaagtttccccag*. Each sample was run in technical triplicate. qRT-PCR was

performed using Applied Biosystems 7300 RT PCR system, as described previously [21].

Treatment of susceptible snails with sodium salicylate

To determine the effect of sodium salicylate on juvenile snail infections, snails were treated with

100ng/ml of sodium salicylate (Sigma Aldrich, St. Louis, MO) overnight (18 hours) at room

temperature in 6-well microtiter plates [21]. Following overnight treatment, snails were washed in

nuclease free water and placed in 24-well microtiter plates. These snails were either unexposed (0

min) or individually exposed to *S. mansoni* 10-12 miracidia for 120 minutes at 25 ☐C. Exposed and

unexposed snails were processed for RNA as described above or kept in beakers to evaluate

cercarial shedding at 4, 6 -weeks post-exposure to infection. The inhibitory concentration was

investigated by means of 2-fold dilution of inhibitor.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 8 software. All data are presented as

mean ±SD. Differences between the groups were assessed using Student's t test, Welch's t test

and 2-way analysis of variance (ANOVA) and Tukey's test wherever relevant by comparing the

differential expression (delta-Ct value) of the transcripts among treatment and control groups. Fold

change was determined by utilizing uniform expression of the housekeeping myoglobin transcript as

standard. Differences were considered statistically significant if $p<0.05$, with level of significance

denoted as follows, ****, $p \le 0.0001$, ***, $p \le 0.001$, **, $p \le 0.01$, *, $p \le 0.05$, and ns, $p > 0.05$.

RESULTS

Carbonic anhydrase is highly conserved in *B. glabrata*

Based on previous RNAseq analysis [21], we identified and validated the differential expression of

carbonic anhydrase protein (Accession number XP_013085564) through a BLAST analysis.

Utilizing SMART BLAST analysis, we revealed the evolutionary relatedness of the Carbonic

Anhydrase in *B. glabrata* to that of other organisms (**Fig 1**). Our phylogenetics results indicate that

B. glabrata is closely related to an invertebrate fruit fly (*D. melanogaster)* and to other vertebrate

animal orthologs (house mouse, zebrafish and human), indicating co-evolution of parasitism in

these organisms. Likely, the CA gene has been selected over millions of years to accommodate the

parasite in these invertebrate and vertebrate hosts (**Table 1, Fig 1**). Further analysis showed that

the CA encoding transcript denotes a single copy gene in the *B. glabrata* genome.

	Query Cover	E Value	Percent Identity	Accession
carbonic anhydrase 14 isoform X1 [Homo sapiens]	85%	6,00E-19	37.82%	XP_005245116.1
carbonic anhydrase 14 precursor [Danio rerio]	85%	1,00E-17	34.71%	NP_001315073.1

carbonic anhydrase 1 [Drosophila melanogaster]	66%	1,00E-17	45.45%	NP_523561.1
carbonic anhydrase 12 isoform X1 [Mus musculus]	71%	3,00E-17	41.18%	XP_006511611.1
carbonic anhydrase 14 isoform X6 [Homo sapiens]	71%	4,00E-17	43.00%	XP_016856382.1

Table 1. Amino acid sequence alignment analysis of *B. glabrata* Carbonic Anhydrase 14-like protein indicates high evolutionary conservancy in relation to several organisms.

Figure 1. Phylogenetic tree analysis highlighting *B. glabrata* carbonic anhydrase homolog's evolutionary relatedness to the human and other organisms. The phylogenic tree is based on amino-acid sequence alignment (ClustalW). Results indicate that *B. glabrata* is closely related to fruit fly (*D. melanogaster)* and other vertebrate organisms (house mouse, zebrafish and human).

Carbonic anhydrase is a surrogate marker of infection

To determine whether changes in the expression of CA occur in juvenile *B. glabrata* snails after parasite exposure, qPCR was utilized to measure temporal changes at different time-points after exposure to *S. mansoni* miracidia. Results showed that CA was differentially expressed between susceptible (BBO2) and resistant (BS-90) snail stocks upon early exposure to the parasite. Thus, a significant upregulation (1.42-fold) was observed following 2h post-exposure, increasing to 1.77-fold after 4hr infection in the susceptible snail (**Fig 2**). Moreover, sustained expression of CA mRNA was observed throughout the prolonged 6- week infection period of the susceptible snail (**Fig 3**). The CA mRNA transcript was significantly upregulated between 1 to 6 weeks (1.9- to 8-fold) post- exposure in the susceptible BBO2 snails but not in the resistant BS90 snails.

Figure 2. qPCR analysis of RNA from susceptible BBO2 (black histograms) and resistant BS-90 (gray histograms) juvenile snails exposed for 30 min, 1 hour, 2 hours and 4 hours to the parasite *S. mansoni* miracidia. Histograms show the expression of CA encoding transcript in *B. glabrata* snails at specific time points from four biological replicates (the total snails used for four biological replicates N=40). Note the increase in fold change in susceptible BBO2 snails compared to the resistant BS90 snails after parasite exposure. Fold change was determined as described previously by utilizing uniform expression of the reference transcript, ****, $p \leq 0.0001$, ***, $p \leq 0.001$, **, $p \leq 0.01$, *, $p \leq 0.05$, and ns, $p > 0.05$.

Figure 3. qPCR analysis of RNA from susceptible BBO2 (black histograms) and resistant BS-90 (gray histograms) juvenile snails exposed for 2 hours and kept up to 6 weeks to the parasite *S. mansoni*. Histograms show expression of CA transcripts in juvenile snails from three biological replicates (N=36). Unexposed (0min) juvenile snails served as controls. A significant increase in fold change can be observed in the susceptible BBO2 snails compared to the resistant BS90 snails. ****, $p \leq 0.0001$, ***, $p \leq 0.001$, **, $p \leq 0.01$, *, $p \leq 0.05$, and ns, $p > 0.05$.

Sodium salicylate is a potent inhibitor of carbonic anhydrase

We also investigated the potential of aspirin to inhibit CA expression in susceptible BBO2 snails. To achieve this, we utilized sodium salicylate, a major metabolite of aspirin, which is a water soluble non-steroidal anti-inflammatory agent known to induce apoptosis of cancer cells and reduce tumor growth [22]. Treatment of BBO2 snails with 100 ng/ml sodium salicylate overnight (18 hours) prior to schistosome exposure was shown to suppress CA expression (**Fig 4**). Through a 2-fold serial dilution assay of the drug, we determined the required inhibitory concentration of sodium salicylate for suppressing CA expression. Our findings, as seen from **Figure 5**, consistently showed downregulation of CA encoding RNA transcript. Notably, a concentration as low as 12.5 ng/mL of sodium salicylate was sufficient to inhibit CA. Importantly, CA expression remained downregulated in sodium salicylate treated snails with exposure to the parasite *S. mansoni*.

Figure 4. Schistosome recovery from infected *B. glabrata* following treatment with 100 ng/ml sodium salicylate. Susceptible BBO2 snails were treated overnight with 100 ng/mL sodium salicylate and exposed to *S. mansoni* miracidia (10-12 miracidia per snail) for two hours. The histograms show cercarial shedding in non-drug treated/infected snails (red); one-time drug treated (16 hours)/infected snails (green); and daily drug-treated/infected snails (yellow) from four biological replicates (N=12). The fold changes of the histograms were noted as non-significant (ns) through 2-way ANOVA analysis.

266
Figure 5. qPCR analysis of RNA from susceptible juvenile BBO2 snails unexposed (gray histograms) and exposed (blue histograms) for 2 to the parasite *S. mansoni* miracidia. Histograms show expression of carbonic anhydrase encoding transcript in normal non-drug treated juvenile snails (0 ng/ml) and those treated with sodium salicylate (12.5 ng/ml, 25 ng/ml, 50 ng/ml or 100 ng/ml) from three biological replicates (N=30). Note the upregulation of CA transcript in non-drug treated exposed snails and the downregulation in unexposed and exposed drug-treated snail groups. ****, $p \leq 0.0001$, ***, $p \leq 0.001$, **, $p \leq 0.01$, *, $p \leq 0.05$, and ns, $p > 0.05$.

Sodium salicylate may reduce infection burden in susceptible *B. glabrata* snails

To test the effects of suppression of CA gene *in vivo*, sodium salicylate treated BBO2 snails were maintained after exposure to parasite to examine for cercariae shedding after 4-6 weeks. Our results indicated that a one-time overnight treatment with 100 ng/mL sodium salicylate prior to schistosome infection (2-hour exposure) was insufficient to block infection in the snails. While neither the non-treated infected snails nor the drug- treated infected snails shed at 4 weeks, following 6 weeks' time, all snails shed cercaria (**Fig 4**). Non drug-treated infected control snails shed the highest number of cercariae. Similarly, one-time drug treated snails shed cercariae, although at lower levels. Following the experiment, we repeated the sodium salicylate treatment in susceptible BBO2 snails, this time maintaining the infected snails continuously in 100 ng/ml sodium

salicylate both before (overnight) and after 2-hour exposure to the parasite. Following 4 weeks of

exposure, we observed a decrease (1 cercariae shed) or complete absence of cercarial shedding in

the treated/exposed snails compared to the non-treated/exposed control group which shed an

average of 9 cercaria. Surprisingly, cercarial shedding was observed in the daily drug- treated

infected snails at 6 weeks, however, at lower levels compared to the non-drug treated infected

control group and one-time drug-treated infected snails (**Fig 4**). However, these results also indicate

that there was no statistically significant difference between the number of cercarial shedding

between BBO2 exposed to either sodium salicylate once or daily or normal miracidia.

DISCUSSION

Schistosomiasis remains a serious public health concern, especially in many low-income sub-

Saharan African areas. To date, however, the only known effective treatment against the causative

agent of schistosomiasis- the parasitic trematode, is praziquantel [4]. Schistosomiasis is endemic in

78 countries and at least 600 million of people are at risk of becoming infected every year [4]. The

WHO has announced that by 2030 schistosomiasis will be eradicated [7]. However, without a

preventative vaccine, and only a single therapeutic drug currently available, this might prove

elusive. Efforts to control schistosomiasis have been attempted by a mass drug chemotherapy

program that provided treatment for school aged children [23]. The detection of infection in patients

relies on the detection of parasite eggs in stool (*S. mansoni*, and *S. japonicum*) and urine (*S.*

hematobium) [4]. For the past 18 years this method of controlling schistosomiasis, although

successful, has not eliminated the disease. This is because the parasite is difficult to completely

eliminate without attention to the transmission in the intermediate snail host.

The control of the freshwater snail has been made possible largely by the use of molluscicides that

can be environmentally toxic as well as disturbing fragile ecosystems. There are currently no

reliable convenient biomarkers available for the detection of the parasite in infected snails. Results

from this study show that the presence of carbonic anhydrase is a good and accurate biomarker for

the early detection of infection by the parasite *Schistosoma mansoni* in the intermediate snail host*,*

Biomphalaria glabrata.

From qPCR analysis temporal regulation during the initial 30 minutes to 4 hours post exposure to

the parasite revealed that CA is upregulated in susceptible BBO2 snails during early infection but

downregulated in their resistant counterparts. Given that CAs are predominantly involved in hypoxia

and pH regulation in cancer cells, we observed a significant upregulation of CA expression in

susceptible snail lines, with a 1.42-fold increase after two hours of exposure and 1.77-fold increase

post four hours (**Fig 2**). Moreover, we found that CA mRNA levels were further increased in

susceptible snails during prolonged infection up to 6 weeks (**Fig 3**), indicating the critical role of CA

in upholding the proliferative function. To further validate the results, we used the *B. glabrata*

embryonic cell line (BGE) co-cultured with *S. mansoni* miracidia and observed a significant 1.4-fold

upregulation of CA transcripts at the two-hour infection period (data not shown). Our previous

research has demonstrated that transcripts encoding the heat shock proteins Hsp70 and Hsp90,

including the reverse transcriptase (RT) domain, the non-LTR-transposon nimbus, are similarly

upregulated in the susceptible snails but not in resistant [24]. Further studies from spatial

repositioning of gene loci have confirmed that schistosomes orchestrate transcription of Hsp70

transcript in susceptible *B. glabrata* snails quickly after infection [25].

CAs also serve as important therapeutic targets in various biological processes, including acid-base

balance, inflammation and angiogenesis. In humans, there are 15 CA isoforms (alpha-class CAs),

with CA IX and CA XII having been linked to cancer [14]. These enzymes are transmembrane

isoforms with an extracellular catalytic domain, showing high expression in solid tumors and low

expression in normal tissues. The overexpression of CA IX and CA XII in tumors is associated with

the survival and proliferation of cancer cells, making them attractive targets for cancer therapy [26].

In this study, we specifically examined CA isoform XIV, which has a medium-low catalytic activity

similar to that of hCA XII [27]. This isoform is highly abundant in the brain, kidneys, colon, small

intestine, urinary bladder, liver, and spinal cord [28]. A Clustal Omega sequence alignment [29]

showed high amino acid conservation among the human α-CAs, specifically in the catalytic sites.

Results indicated that the active site of these enzymes are highly superimposable, suggesting

similar evolutionary conserved enzymatic activity among the CA encoding transcripts between the

snail homolog and the other organisms. Of particular interest, structural comparison of the hCA IX

catalytic domain with the CA 14-like *B. glabrata* homolog yielded a substantial sequence homology

(**Fig 6**). While the two enzyme orthologs share 35.97% sequence identity in the catalytic sites,

further interrogations of the sequences revealed that the two proteins consist of identical zinc

binding sites (ion binding), specifically, the three conserved histidine residues (His94, His96, and

His119). This homology suggests that the isoforms may share similar functions, and one isoform

may be able to compensate for the loss of the other [30].

Figure 6. Structure-based sequence alignment of the catalytic regions of the human

Carbonic Anhydrase IX and *B. glabrata* Carbonic Anhydrase 14-like protein homolog

obtained from Clustal Omega. The two enzyme orthologs share 35.97% amino acid sequence

identity. 100% match is indicated by an asterisk.

We have shown there is a link between metastatic cancers and parasitic diseases, highlighting the

snail host/parasite relationship as a valuable animal model for studying the regulation of cancer-

associated transcripts. Similar to cancer, the schistosome parasite survives in the blood stream

while evading the immune system and migrating throughout the host system for extended periods.

Deposition of *S. mansoni* eggs also causes acute and chronic inflammation of the colorectal

mucosa [31]. Our results indicate that CA plays a key role in the progression of schistosomiasis

infection in the snail, with prolonged infection leading to increased expression of CA transcripts.

CA's role in facilitating the transport of CO_2 across the cell membrane creating acidic environment,

is one of the hallmarks of cancer. In accord with studies from CA in cancer, several human CA

isoforms have been shown to be increased in tumor tissue, correlating with cancer growth and

survival [32-35]. This supports the notion that the snail-schistosome model can provide useful

insights into the mechanisms underlaying both parasitic infections and cancer (manuscript in

preparation).

Targeting CA has emerged as a promising therapeutic strategy for cancer treatment with numerous

CA inhibitors currently undergoing preclinical and clinical trials [36, 37]. It has been well established

that a positive correlation exists between risk of cancer and aspirin intake. As part of this, several

clinical trials have investigated the use of aspirin as an adjuvant therapy for cancer patients,

particularly those with colorectal cancer [38, 39]. We show in this study that sodium salicylate, a

major subunit of aspirin, can inhibit CA in susceptible BBO2 snails. Although our results from qPCR

show that sodium salicylate down regulates the transcript encoding carbonic anhydrase, we found

that a single dose of 100 ng/ml sodium salicylate is not sufficient to block infection, likely due to its

metabolism and short half-life. However, daily administration of 100 ng/ml sodium salicylate can

further reduce the infection burden compared to a single dose and normal infected snails.

Combination therapy involving aspirin may offer greater therapeutic benefits, as evidenced by

Feitosa, et al. demonstrating a reduction in parasite load in *S. mansoni* mice [40]. Furthermore, the

combination of paraziquantel and aspirin has been shown to decrease liver pathology, highlighting

the potential of this treatment approach [41].

In summary, we have shown in this study that CA overexpression is correlated with infection and

parasite progression in the snail host. We provide the first evidence of CA in mollusk species upon

infection with *S. mansoni* and show that CA is a promising biomarker to track schistosomiasis in

early and prolonged infection stages. To gain a better understanding of the mechanism of action of

CA, further studies to determine its enzymatic activity are needed. To this end, we are currently

conducting an enzymatic assay to evaluate CA activity. In addition, antibody detection assays are

being developed to make the use of CA for rapid and practical use in the field to detect occurrence

of infected snails in endemic high prevalence area before using molluscicides which can be

expensive to use on a wide scale. To validate the presented findings, we will examine the

expression of CA infected with other schistosomiasis-causing trematodes, *S. japonicum* and *S.*

haematobium.

Acknowledgements

We thank Dr. Margaret Mentink-Kane at the NIAID Schistosomiasis Resource Center of the

Biomedical Research Institute for the snail and parasite material used in this study. We also thank

Dr. Carolyn Cousin for resources and support. This work was funded by the Clement BT Knight

Cancer Foundation.

CRediT authorship contribution statement

Matty Knight: Conceptualization, Methodology, Investigation, Resources, Project Administration,

Supervision, Funding Acquisition, Writing – Original Draft, Writing – Review & Editing, Funding

acquisition. **Simone Parn:** Investigation, Methodology, Data Curation, Formal Analysis, Validation,

Visualization, Writing – Original Draft, Writing – Review & Editing; **Gabriela Lewis:** Investigation

Methodology, Data Curation, Validation, Writing – Review & Editing.

Conflict of Interest

The authors declare no conflict of interest.

Declaration of Competing Interests

The authors declare that they have no known competing financial interests or personal relationships

that could have appeared to influence the work reported in this paper.

Data Availability

Data will be made available on request.

REFERENCES

1. Verjee MA. Schistosomiasis: Still a Cause of Significant Morbidity and Mortality. Res Rep Trop Med. 2019 Dec 31;10:153-163. doi: 10.2147/RRTM.S204345. Accessed March 8, 2023.

2.Parasites- schistosomiasis. Centers for Disease Control and Prevention. 2018. https://www.cdc.gov/parasites/schistosomiasis/index.html. Accessed April 8, 2023.

3. Boissier J, Grech-Angelini S, Webster BL, Allienne JF, Huyse T, Mas-Coma S, Toulza E, Barré-Cardi H, Rollinson D, Kincaid-Smith J, Oleaga A, Galinier R, Foata J, Rognon A, Berry A, Mouahid G, Henneron R, Moné H, Noel H, Mitta G. Outbreak of urogenital schistosomiasis in Corsica (France): an epidemiological case study. Lancet Infect Dis. 2016 Aug;16(8):971-9. doi: 10.1016/S1473-3099(16)00175-4. Accessed April 8, 2023.

4. Schistosomiasis. The World Health Organization. 2023. https://www.who.int/news-room/fact-sheets/detail/schistosomiasis. Accessed April 5, 2023.

5. Hotez PJ, Engels D, Gyapong M, Ducker C, Malecela MN. Female Genital Schistosomiasis. N Engl J Med. 2019 Dec 26;381(26):2493-2495. doi: 10.1056/NEJMp1914709. Accessed March 30, 2023.

6. WHO. (2018). Schistosomiasis: Progress report 2001–2011, strategic plan 2012–2020. Geneva, Switzerland: World Health Organization. Retrieved from https://www.who.int/schistosomiasis/resources/9789241503174/en/. Accessed March 15, 2023.

7. Ending the neglect to attain the Sustainable Development Goals: A road map for neglected

tropical diseases 2021–2030. Geneva: World Health Organization; 2020. Licence:

CC BY-NC-SA 3.0 IGO. Accessed April 10, 2023.

8. Adema, C., Hillier, L., Jones, C. *et al.* Whole genome analysis of a schistosomiasis-transmitting

freshwater snail. *Nat Commun* 8, 15451 (2017). https://doi.org/10.1038/ncomms15451. Accessed

February 25, 2023.

9. Bridger JM, Brindley PJ, Knight M (2018) The snail *Biomphalaria glabrata* as a model to

interrogate the molecular basis of complex human diseases. PLoS Negl Trop Dis 12(8): e0006552.

https://doi.org/10.1371/journal.pntd.0006552. Accessed April 12, 2023.

10. Raghavan N, Miller AN, Gardner M, FitzGerald PC, Kerlavage AR, Johnston DA, Lewis FA,

Knight M. Comparative gene analysis of Biomphalaria glabrata hemocytes pre- and post-exposure

to miracidia of Schistosoma mansoni. Mol Biochem Parasitol. 2003 Feb;126(2):181-91. doi:

10.1016/s0166-6851(02)00272-4. Accessed April 8, 2023.

11. Knight M, Arican-Goktas HD, Ittiprasert W, Odoemelam EC, Miller AN, Bridger JM.

Schistosomes and snails: a molecular encounter. Front Genet. 2014 Jul 21;5:230. doi:

10.3389/fgene.2014.00230. Accessed April 10, 2023.

12. Zolfaghari Emameh R, Barker HR, Turpeinen H, Parkkila S, Hytönen VP. A reverse vaccinology

approach on transmembrane carbonic anhydrases from Plasmodium species as vaccine candidates

for malaria prevention. Malar J. 2022 Jun 15;21(1):189. doi: 10.1186/s12936-022-04186-7.

Accessed April 11, 2023.

13. Angeli A, Pinteala M, Maier SS, Simionescu BC, Da'dara AA, Skelly PJ, Supuran CT.

Sulfonamide Inhibition Studies of an α-Carbonic Anhydrase from *Schistosoma mansoni*, a

Platyhelminth Parasite Responsible for Schistosomiasis. Int J Mol Sci. 2020 Mar 7;21(5):1842. doi:

10.3390/ijms21051842. Accessed April 11, 2023.

14. Supuran CT. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators.

Nature Reviews Drug Discovery 7, no. 2 (2008): 168-181. doi: 10.1038/nrd2467. Accessed April 13,

2023.

15. Pinard MA, Mahon B, McKenna R. Probing the surface of human carbonic anhydrase for clues

towards the design of isoform specific inhibitors. Biomed Res Int. 2015;2015:453543. doi:

10.1155/2015/453543. Accessed April 13, 2023.

16. Poggetti V, Salerno S, Baglini E, Barresi E, Da Settimo F, Taliani S. Carbonic Anhydrase

Activators for Neurodegeneration: An Overview. Molecules. 2022 Apr 14;27(8):2544. doi:

10.3390/molecules27082544. Accessed March 27, 2023.

17. Svastová E, Hulíková A, Rafajová M, Zaťovicová M, Gibadulinová A, Casini A, Cecchi A,

Scozzafava A, Supuran CT, Pastorek J, Pastoreková S. Hypoxia activates the capacity of tumor-

associated carbonic anhydrase IX to acidify extracellular pH. FEBS Lett. 2004 Nov 19;577(3):439-

45. doi: 10.1016/j.febslet.2004.10.043. Accessed April 11, 2023.

18. Pastorekova S, Gillies RJ. The role of carbonic anhydrase IX in cancer development: links to

hypoxia, acidosis, and beyond. Cancer Metastasis Rev. 2019 Jun;38(1-2):65-77. doi:

10.1007/s10555-019-09799-0. Accessed March 27, 2023.

19. Becker HM. Carbonic anhydrase IX and acid transport in cancer. *Br J Cancer* 122, 157–167

(2020). https://doi.org/10.1038/s41416-019-0642-z. Accessed April 13, 2023.

20. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol

Biol. 1990 Oct 5;215(3):403-10. doi: 10.1016/S0022-2836(05)80360-2. Accessed April 13, 2023.

21. Smith M, Yadav S, Fagunloye OG, Pels NA, Horton DA, Alsultan N, Borns A, Cousin C, Dixon

F, Mann VH, Lee C, Brindley PJ, El-Sayed NM, Bridger JM, Knight M. PIWI silencing mechanism

involving the retrotransposon nimbus orchestrates resistance to infection with Schistosoma mansoni

in the snail vector, Biomphalaria glabrata. PLoS Negl Trop Dis. 2021 Sep 8;15(9):e0009094. doi:

10.1371/journal.pntd.0009094. Accessed February 27, 2023.

22. National Center for Biotechnology Information (2023). PubChem Compound Summary for CID

16760658, Sodium Salicylate. Accessed April 13, 2023.

23. Kokaliaris C, Garba A, Matuska M, Bronzan RN, Colley DG, et al. Effect of preventive

chemotherapy with praziquantel on schistosomiasis among school-aged children in sub-Saharan

Africa: a spatiotemporal modelling study. Lancet Infect Dis. 2022 Jan;22(1):136-149. doi:

10.1016/S1473-3099(21)00090-6. Epub 2021 Dec 2. Erratum in: Lancet Infect Dis. 2022

Jan;22(1):e1. Accessed April 13, 2023.

24. Ittiprasert W, Nene R, Miller A, Raghavan N, Lewis F, Hodgson J, Knight M. Schistosoma

mansoni infection of juvenile Biomphalaria glabrata induces a differential stress response between

resistant and susceptible snails. Exp Parasitol. 2009 Nov;123(3):203-11. doi:

10.1016/j.exppara.2009.07.015. Accessed March 30, 2023.

25. Arican-Goktas HD, Ittiprasert W, Bridger JM, Knight M. Differential spatial repositioning of

activated genes in Biomphalaria glabrata snails infected with Schistosoma mansoni. PLoS Negl

Trop Dis. 2014 Sep 11;8(9):e3013. doi: 10.1371/journal.pntd.0003013. Accessed March 30, 2023.

26. Mahon BP, Pinard MA, McKenna R. Targeting carbonic anhydrase IX activity and expression.

Molecules. 2015 Jan 30;20(2):2323-48. doi: 10.3390/molecules20022323. Accessed April 13, 2023.

27. Nishimori I, Vullo D, Innocenti A, Scozzafava A, Mastrolorenzo A, Supuran CT. Carbonic

anhydrase inhibitors: inhibition of the transmembrane isozyme XIV with sulfonamides. Bioorg Med

Chem Lett. 2005 Sep 1;15(17):3828-33. doi: 10.1016/j.bmcl.2005.06.055. Accessed April 10, 2023.

28. Alterio V, Pan P, Parkkila S, Buonanno M, Supuran CT, Monti SM, De Simone G. The structural

comparison between membrane-associated human carbonic anhydrases provides insights into drug

design of selective inhibitors. Biopolymers. 2014 Jul;101(7):769-78. doi:

10.1002/bip.22456. Accessed April 4, 2023.

29. Sievers F, Wilm A, Dineen D, et al. Fast, scalable generation of high-quality protein multiple

sequence alignments using Clustal Omega. Molecular Systems Biology. 2011 Oct;7:539. DOI:

10.1038/msb.2011.75. Accessed April 13, 2023.

30. Aggarwal M, Boone CD, Kondeti B, McKenna R. (2013) Structural annotation of human carbonic

anhydrases, Journal of Enzyme Inhibition and Medicinal Chemistry,28:2, 267-

277, DOI: 10.3109/14756366.2012.737323. Accessed April 10, 2023.

31. McManus, D.P., Dunne, D.W., Sacko, M. *et al.* Schistosomiasis. *Nat Rev Dis Primers* 4, 13

(2018). https://doi.org/10.1038/s41572-018-0013-8. Accessed April 13, 2023.

32.Ning WR, Jiang D, Liu XC, Huang YF, Peng ZP, Jiang ZZ, Kang T, Zhuang SM, Wu Y, Zheng L.

Carbonic anhydrase XII mediates the survival and prometastatic functions of macrophages in

human hepatocellular carcinoma. J Clin Invest. 2022 Apr 1;132(7):e153110. doi:

10.1172/JCI153110. Accessed April 3, 2023.

33.McDonald PC, Chafe SC, Brown WS, Saberi S, Swayampakula M, Venkateswaran G,

Nemirovsky O, Gillespie JA, Karasinska JM, Kalloger SE, Supuran CT, Schaeffer DF, Bashashati A,

Shah SP, Topham JT, Yapp DT, Li J, Renouf DJ, Stanger BZ, Dedhar S. Regulation of pH by

Carbonic Anhydrase 9 Mediates Survival of Pancreatic Cancer Cells With Activated KRAS in Response to Hypoxia. Gastroenterology. 2019 Sep;157(3):823-837. doi: 10.1053/j.gastro.2019.05.004. Accessed April 3, 2023.

34.Hsieh M, Huang PJ, Chou PY, Wang SW, Lu HC, Su WW, Chung YC, Wu MH. Carbonic Anhydrase VIII (CAVIII) Gene Mediated Colorectal Cancer Growth and Angiogenesis through Mediated miRNA 16-5p. Biomedicines. 2022 Apr 29;10(5):1030. doi: 10.3390/biomedicines10051030. Accessed April 3, 2023.

35.Schmidt J, Oppermann E, Blaheta RA, Schreckenbach T, Lunger I, Rieger MA, Bechstein WO, Holzer K, Malkomes P. Carbonic-anhydrase IX expression is increased in thyroid cancer tissue and represents a potential therapeutic target to eradicate thyroid tumor-initiating cells. Mol Cell Endocrinol. 2021 Sep 15;535:111382. doi: 10.1016/j.mce.2021.111382. Accessed April 3, 2023.

36.Mussi S, Rezzola S, Chiodelli P, Nocentini A, Supuran CT, Ronca R. Antiproliferative effects of sulphonamide carbonic anhydrase inhibitors C18, SLC-0111 and acetazolamide on bladder, glioblastoma and pancreatic cancer cell lines. J Enzyme Inhib Med Chem. 2022 Dec;37(1):280-286. doi: 10.1080/14756366.2021.2004592. Accessed April 3, 2023.

37.Dvořanová J, Kugler M, Holub J, Šícha V, Das V, Nekvinda J, El Anwar S, Havránek M, Pospíšilová K, Fábry M, Král V, Medvedíková M, Matějková S, Lišková B, Gurská S, Džubák P, Brynda J, Hajdúch M, Grüner B, Řezáčová P. Sulfonamido carboranes as highly selective inhibitors of cancer-specific carbonic anhydrase IX. Eur J Med Chem. 2020 Aug 15;200:112460. doi: 10.1016/j.ejmech.2020.112460. Accessed April 3, 2023.

38.Sostres C, Gargallo CJ, Lanas A. Aspirin, cyclooxygenase inhibition and colorectal cancer. World J Gastrointest Pharmacol Ther. 2014 Feb 6;5(1):40-9. doi: 10.4292/wjgpt.v5.i1.40. Accessed April 5, 2023.

39.Rothwell PM, Wilson M, Elwin CE, Norrving B, Algra A, Warlow CP, Meade TW. Long-term effect

of aspirin on colorectal cancer incidence and mortality: 20-year follow-up of five randomised trials.

Lancet. 2010 Nov 20;376(9754):1741-50. doi: 10.1016/S0140-6736(10)61543-7. Accessed April 5,

2023.

40.Feitosa KA, Zaia MG, Rodrigues V, Castro CA, Correia RO, Pinto FG, Rossi KNZP, Avó LRS,

Afonso A, Anibal FF. *Menthol and Menthone* Associated with Acetylsalicylic Acid and Their Relation

to the Hepatic Fibrosis in *Schistosoma mansoni* Infected Mice. Front Pharmacol. 2018 Jan

18;8:1000. doi: 10.3389/fphar.2017.01000. Accessed April 14, 2023.

41.Sudsarn P, Boonmars T, Ruangjirachuporn W, Namwat N, Loilome W, Sriraj P, Aukkanimart R,

Nadchanan W, Jiraporn S. Combination of Praziquantel and Aspirin Minimizes Liver Pathology of

Hamster Opisthorchis viverrini Infection Associated Cholangiocarcinoma. Pathol Oncol Res. 2016

Jan;22(1):57-65. doi: 10.1007/s12253-015-9967-y. Accessed April 14, 2023.

CA in Sodium Salicylate-treated BBO2