SMYD5 is a novel epigenetic gatekeeper of the mild hypothermia response

Salvor Rafnsdottir[1], Kijin Jang[1], Sara Tholl Halldorsdottir[1], Arnhildur Tomasdottir[1], Meghna Vinod[1],

Katrin Möller[1], Tinna Reynisdottir[1], Laufey Halla Atladottir[1], Kristin Elisabet Allison[2], Jin He[3], Li

Zhang[4], Frances J. Northington[5], Raul Chavez-Valdez[5], Kimberley Jade Anderson[1], Hans Tomas

Bjornsson[1,4,6,7*]

[1]Louma G. Laboratory of Epigenetic Research, Faculty of Medicine, University of Iceland, Reykjavik,

Iceland; [2]Faculty of Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland;

[3]Department of Biochemistry and Molecular Biology, College of Natural Science, Michigan State

University, MI, USA; [4]McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University

School of Medicine, MD, USA; [5]Division of Neonatology, Department of Pediatrics, Johns Hopkins

University School of Medicine, Baltimore, MD, USA; [6]Department of Pediatrics, Johns Hopkins

University, MD, USA; [7]Department of Genetics and Molecular Medicine, Landspitali University

Hospital, Reykjavik, Iceland.

*Corresponding author: Hans Tomas Bjornsson, htb@hi.is

Summary

Organisms have homeostatic mechanisms to respond to cold temperature to ensure survival

including the activation of the mammalian neuroprotective mild hypothermia response (MHR) at

32°C. We show activation of the MHR at euthermia by an FDA-approved medication Entacapone,

proof-of-principle that the MHR can be medically manipulated. Utilizing a forward CRISPR-Cas9

mutagenesis screen, we identify the histone lysine methyltransferase SMYD5 as an epigenetic

gatekeeper of the MHR. SMYD5 represses the key MHR gene *SP1* at euthermia but not at 32°C. This

repression is mirrored by temperature-dependent levels of H3K36me3 at the *SP1*-locus and globally

indicating that the mammalian MHR is regulated at the level of histone modifications. We identified

45 additional SMYD5-temperature dependent genes suggesting a broader MHR-related role for

SMYD5. Our study provides an example of how the epigenetic machinery integrates environmental

cues into the genetic circuitry of mammalian cells and suggests novel therapeutic avenues for

neuroprotection after catastrophic events.

Keywords: Targeted Temperature Management, SP1, Hypothermia, Neonatal Hypoxic-Ischemic Brain

Injury, Cardiac Arrest

Introduction

The defining feature of mammals is their ability to maintain a euthermic body temperature (37°C) and most mammals maintain temperature at a relatively tight range to ensure normal cellular homeostasis[1]. The main exception is the hibernation of some mammals, which is a type of controlled temperature decrease during times of prolonged inactivity[2]. In clinical practice, physicians imitate hibernation with the use of Targeted Temperature Management (TTM) as a therapeutic strategy to minimize neurological damage following neonatal hypoxic-ischemic brain injury and cardiac arrest[3,4]. The reasons why lowered internal core temperature (32-36°C) yields benefits are currently not fully understood[5,6]. Although some argue that the positive neurological benefit from mild hypothermia relates to a lower metabolic rate, others point out that the specific upregulation of several genes during mild hypothermia does not fit that idea[7]. An alternative hypothesis is that hypothermia is beneficial because of the activation of the mild hypothermia response (MHR) itself, and if so, it would be important for researchers and clinicians to understand both the mechanisms and the extent of the response. In support of this, researchers have found that several genes (e.g., *CIRBP/CIRP*, *RBM3*) consistently show increased expression during mild hypothermia.[8] Two of these, *SP1* and *RBM3,* appear to have roles in decreasing neuronal death during disease states[9–13].

SP1, a transcription factor for many genes[14], binds to the promoter region of *CIRBP*[15] upon cold exposure at a sequence called the Mild Cold Response Element (MCRE)[15], and its binding leads to increased expression of *CIRBP* at 32°C, indicating that this regulation occurs at the transcriptional level[15]. Similarly, RBM3 is known to regulate at least one downstream gene, *RTN3*[11], a potential neuroprotective factor in a neurodegeneration model[11]. Although this suggests that mammalian cells have a pathway culminating in SP1/CIRBP and RBM3/RTN3, little has been done to further elucidate the upstream regulation of this pathway and relatively little knowledge is available regarding mammalian cellular responses to cooling compared to heat shock responses, which have been previously characterized[16].

Here, we use complementary unbiased strategies to gain clues into the MHR upstream of *SP1*. Specifically, we use a novel *SP1* mild hypothermia indicator (MHI) containing the *SP1* promoter and the MCRE enhancer (SP1+MCRE-MHI) to perform compound screening of 1953 FDA-approved agents. We have successfully identified and validated that the drug Entacapone activates the endogenous MHR at the transcriptional level, which indicates that it is possible to manipulate the response without exposure to hypothermia. To identify additional regulators, we have performed a genome-wide CRISPR-Cas9 forward mutagenesis screen[17–19], reading out *SP1*-promoter activity using the SP1-MHI. Our study has yielded multiple candidate regulators and compounds that may regulate

the MHR. We focus on one of these regulators, *SMYD5*, a histone methyltransferase, and

demonstrate that the presence of SMYD5 is temperature dependent and that several other SMYD5-

bound genes show significant differential expression upon mild hypothermia exposure. Our insights

elucidate how an external cue (mild hypothermia) specifically mediates its effects on the mammalian

genome through the actions of the epigenetic machinery and this paves the way towards the

development of a novel therapeutic means to treat neonatal hypoxic-ischemic brain injury, a major

cause of mortality and morbidity in otherwise healthy newborns[20], without exposure to TTM.

Results

Mild hypothermia indicators show increased fluorescence at mild hypothermia (32°C).

Core temperature (euthermia) is kept at a very tight range in humans (36.5-37.8°C)[1,21] and by

definition, mild hypothermia involves temperature ranges from 32-36°C (**Figure S1**, bold). To directly

visualize the promoter activity of three genes previously shown[8,15] to respond to mild hypothermia

(*CIRBP*, *RBM3*, *SP1*), we have made eight distinct MHIs (**Supplementary Table 1** and **Supplementary**

File 1) all of which have promoter and/or MCRE sequences, driving the expression of either GFP or

mCherry (**Figure 1A**). When transfected into HEK293 cells, the MHIs demonstrate increased

fluorescence at 32°C compared to 37°C (**Figure 1B–1C**). These findings are mirrored by increased

endogenous SP1 levels in a Western blot from cells collected after 16 hours (h) at 32°C (**Figure 1D**).

We assessed the temporal dynamics of promoter activity using flow cytometry analysis of the three

indicators and observed a maximal increase in fluorescence of SP1-MHI between 6-8 h after a

temperature shift to 32°C (P<0.05, **Figure 1E**). This finding was supported by continuous monitoring

of fluorescence from an Incucyte S3 (Sartorius) where again maximal intensity was reached within 6-

8 h, with SP1-MHI, rising fastest, followed by RBM3-MHI, and then CIRBP-MHI (**Figure S2A-S2C**).

Indicators were also transiently transfected into HEK293 cells and then exposed to five distinct

temperatures (26°C, 29°C, 32°C, 37°C, and 40°C), for 16 h followed by flow cytometry (**Figure 1F-1H;**

Figure S2D-S2E). Consistent with previous literature[8,15] and our own validation in HEK293 cells

(**Figure 1B-D**), indicators representing all three genes (*SP1*, *CIRBP*, *RBM3)* had significantly (P<0.0001)

increased fluorescence at 32°C compared to 37°C/40°C (**Figure 1F-1H**) and the response was

uniformly strongest at 32°C (**Figure 1F-1H**). At the same time, the SP1- and CIRBP-MHIs do not show

increased activity at lower (26°C, 29°C, **Figure S2D-S2E**) or higher (40°C) temperatures (**Figure 1F-1G**),

supporting the notion that our MHIs specifically reflect the MHR and are not activated by general cell

stress (i.e., moderate hypo- or hyperthermia). These data suggest that all three genes are regulated

at the transcriptional level and the time-lag in the response upon hypothermia is measured in hours

rather than minutes, indicating that there must be additional regulators operating upstream of these

genes. Some would argue that storing cells on ice (i.e., during fluorescent activating cell sorting

(FACS)) could possibly lower the temperature of cells to the hypothermic range and activate the

MHR, but we found that even prolonged storage on ice (8 h) in a room at room temperature was

unable to activate the MHI (**Figure S3**), ensuring that storage on ice during the study could not act as

a potential confounder.

Figure 1. MHIs based on all three genes (*SP1*, *CIRBP*, *RBM3*) show the strongest effect at 32°C. (A) A

schematic of MHI structures. (B-C) Representative images of SP1-Lenti-MHIs and CIRBP-Lenti-MHI,

respectively, at 37°C and 32°C as visualized by fluorescent microscope. (D) Western blot

quantification and a representative figure demonstrating increased SP1 after 16 h at 32°C compared

to 37°C. Shown are data points, each data point is a biological replicate and the mean is depicted.

Significance levels were calculated with an unpaired two-tailed t-test in GraphPad Prism. (E)

Geometric mean fluorescence (gMFI, flow cytometry) after varying lengths of hypothermia exposure

for SP1-MHI. Shown are data points, each data point is a technical replicate, mean and SD where

applicable. Significance levels were calculated with an unpaired one-tailed t-test in GraphPad Prism.

(F-G) Mean fluorescence (MFI, flow cytometry) of SP1-MHI and CIRBP-MHI, respectively, (16 h at

32°C, 37°C, 40°C). Shown are data points, each point is a technical replicate, mean and SD where

applicable. Significance levels were calculated in GraphPad Prism with Šidák's multiple comparison

test. (H) gMFI of RBM3-MHI (16 h at 32°C and 37°C). Significance levels were calculated with an

unpaired one-tailed t-test in GraphPad Prism otherwise data shown as in (G-H). * = P<0.05, ** =

P<0.01, *** = P<0.001, **** = P<0.0001.

Activation of the MHR at 32°C is uniformly seen in cells of diverse origins.

In addition to HEK293 cells, which have been widely used in prior MHR studies[11,15,22], we

sought to verify that our novel MHIs also work in diverse cellular systems. We transiently transfected

three individual MHIs (SP1-, RBM3-, SP1+MCRE-MHI) into seven distinct cell lines: (HEK293WT,

HEK293T, HCT-116, HeLa, Jurkat, K562, and SK-N-SH), representing diverse cellular origins including embryonic kidney, colon, cervix, T lymphocytes, lymphoblasts, and neurons, respectively. After exposing these cells to 26°C, 29°C, 32°C, 37°C, and 40°C for 16 h and quantifying the fluorescence by flow cytometry, we observed highest fluorescence at 32°C independent of cellular origin (**Figure S4**) and no effect for moderate hypothermia (26-29°C, **Figure S5**, and **Figure S2D-S2E**) with the exception of SK-N-SH and K562 cell lines (**Figure S5D-S5E**). The strongest activation of MHI at 32°C was detected in the cell-types previously described to have neurological characteristics (HEK293 and SK-N-SH)[23–25] and these were also the cell lines that showed significant activation of all three MHIs at 32°C (**Figure 1F-1H**, and **Figure S4**). This is consistent with prior literature showing that the RBM3-MHR response is strongest in young neurons[26]. Since many of the key studies in the field have used HEK293 [11,15,22] and they have neuronal characteristics[23,25] we used this cell line for our screening approaches.

Compound screening reveals MHR activators at euthermia.

In our search for therapeutic targets for TTM, we sought to identify compounds that could activate the MHR at 37°C as this could yield insights into the involved biochemical pathways. To explore this in an unbiased manner, we performed a systematic library drug screen (1953 FDA-approved inhibitors, **Figure 2A**) using HEK293T cells transfected with the SP1+MCRE-MHI. Cells transfected with SP1+MCRE-MHI were exposed to the drugs (20 µM concentration) and fluorescence was measured after 16, 28, and 40 h of drug exposure (**Figure 2A**). We focused on drugs that increase the MHI fluorescence at least ≥3.75 times at one of the time points (**Supplementary Table 2**). Among these drugs we saw an apparent albeit non-significant enrichment of Tyrosine Kinase Inhibitors (TKI, 7/80, p<0.075) including Orantinib, Dasatinib (hydrochloride, monohydrate), Poziotinib, Rociletinib, Afatinib, and Lapatinib[27–31]. We validated three from this group (Dasatinib, Orantinib, and Poziotinib) as well as selecting another 5 drugs that showed one of the greatest responses in the screen (Cinacalcet, Entacapone, Erythromycin, Naloxone, and Tamoxifen, **Figure 2B**). For validation of the drug screen HEK293 cells expressing SP1-MHI were individually exposed to each of the 8 compounds (20 µM) for 16 h at either 32°C or 37°C followed by flow cytometry. Two out of the eight (25%), showed significantly (P<0.001, P<0.0001) increased fluorescence at 37°C (Entacapone, Poziotinib, **Figure 2C-2D**), indicating these two drugs can potentially induce the MHR. Entacapone also had significantly (P<0.01) enhanced fluorescence levels at 32°C (**Figure 2C**), suggesting that the effect is additive and thus synergistic to the regular response and thus likely independent of the SP1 target itself. Other compounds did not increase the fluorescence of the SP1-MHI for the dose tested (**Figure S6A-S6F**). To exclude the possibility of cell cycle arrest or increased apoptosis[31,32], we evaluated the cell cycle phase in HEK293 cells exposed to Poziotinib and Entacapone, and no apparent signs of cell

cycle arrest were observed for either compound, other than a slight increase in the percentage of

total cells in the S phase of the cell cycle after Entacapone exposure (**Figure S6G**). After exposing

HEK293 cells to the pro-oxidant agent H_2O_2 in order to induce apoptosis, we could not see any signs

of activation of the SP1-MHI (**Figure S6H**), but we observed a significant (P<0.01) increase in

fluorescence after only 4 h at 32°C (**Figure S6H**) suggesting that cell stress may potentiate the

response. To our surprise, three drugs (Cinacalcet, Dasatinib, and Tamoxifen) showed significantly

decreased fluorescence of the SP1-MHI at 32°C during this validation (**Figure S6A**, **S6B**, and **S6F**). To

evaluate the endogenous effects on SP1 we carried out Western blotting after 24 h exposure to the

compounds at either 32°C or 37°C. For Entacapone (20 μM concentration) we observed a significant

increase of SP1 at 37°C (P<0.05, **Figure 2E**). We were unable to test Poziotinib at 20 μM due to low

protein yields, therefore, we performed the Western blot at a lower dose (4 μM) and for this dose we

only saw a trend towards increased SP1 at 37°C (**Figure 2F**). When we looked at mRNA levels for *SP1*,

CIRBP, and *RBM3* at 32°C and 37°C after 16 h of Entacapone and Poziotinib exposure (20 μM), we

only observed a significant (P<0.01) elevated *RBM3* mRNA levels upon Poziotinib exposure at 37°C

(**Figure S7**). This could suggest that the drugs may impact both transcriptional and post-

transcriptional levels of key factors of the MHR.

Figure 2. Poziotinib and Entacapone activate the MHR. (A) A schematic overview of our drug screen in HEK293T cells transfected with SP1+MCRE-MHI. (B) Representative drugs that were found to cause more than ≥3.75 times increase of SP1+MCRE-MHI fluorescence. The entire list is provided in **Supplementary Table 2**. (C-D) gMFI of SP1-MHI as measured by flow cytometry after 16 h of either 32°C or 37°C with or without Entacapone (ENT, C, 20 μM) and Poziotinib (POZ, D, 20 μM) exposure. Shown are data points, each point is a technical replicate, mean and SD where applicable. Significance levels were calculated in GraphPad Prism with Šidák's multiple comparison test. (E-F) Western blot quantification and a representative figure of SP1 after exposure to Entacapone (ENT, E, 20 μM) and Poziotinib (POZ, F, 4 μM) or vehicle (DMSO) after 24 h. Shown are data points, each data point is a biological replicate, mean and SD where applicable. Significance levels were calculated with an unpaired one-tailed t-test in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

CRISPR-Cas9 Knockout Screen with SP1-MHIs reveals candidate regulators.

To map regulators of the cooling response (*SP1*) in an unbiased manner, we used an available genome-coverage lentiGuide-Puro pooled sgRNA library[17–19] in combination with our SP1-MHI to uncover transcriptional regulators of *SP1*. We chose to use the SP1-MHI, as our prior data indicate that this gene is regulated at the transcriptional level (**Figure 1**) and is the quickest key MHR gene to achieve maximal response at 32°C (**Figure S2A-S2C**). However, given that it does not reach its maximal response until 6-8h of hypothermia exposure, we hypothesized that there may be unknown regulators of the *SP1* locus that function earlier in the MHR process. We performed the screen on HEK293WT+Cas9 cells expressing SPI-MHI (HEK293WT+Cas9+SP1), transducing the sgRNA library at the rate of 0.3 sgRNA per cell. After puromycin selection and exposure to hypothermia (32°C) for 16 h (to activate the MHR), we sorted out the 5% most and least fluorescent cells (**Figure 3A**) to capture repressors and activators of *SP1*, respectively. We saw a consistent shift towards lower fluorescence for the transduced HEK293WT+Cas9+SP1-MHI (green) compared to the positive control (non-transduced HEK293WT+Cas9+SP1-MHI, grey, **Figure 3B**). The observation of such a large fluorescent shift of cells (less SP1-MHI fluorescence) at hypothermia suggests the involvement of many sgRNAs, and thus there are likely many genes that are upstream of *SP1*. Furthermore, this suggests that the majority are activators rather than repressors of the SP1 arm of the MHR. We amplified guides for each sorted group as well as the negative control (transduced HEK293WT+Cas9) and performed deep next-generation sequencing of their distribution from sorted and un-sorted fractions (negative control). To identify guides enriched in the sorted populations, we used the MaGeCK pipeline[33] to computationally predict genes with enrichment (**Figure 3C-3D**). We decided to cast a wide net and look at all genes that had a -Log10(RRA) score above 2.5 to not miss possible regulators (**Supplementary Table 3**). We excluded 21 genes that were seen in lists for both the high (repressor list) and low (activator list) sorts of transduced HEK293+Cas9+SP1-MHI (**Supplementary Table 4**), as these may be related to proliferation rather than the MHR itself, and a validation of two of these

(*RGS21* and *MALT1*) supported that idea (**Figure S8**). We were left with a list of 495 genes for the

SP1-repressors and 61 for SP1-activators – a clear overrepresentation of repressors to activators

(**Supplementary Table 3**). Of note, there were two microRNAs on the top list of SP1-repressors and

activators (*hsa-mir-4314* and *hsa-mir-6739*) and both are predicted to target SP1 (mirdb.org).

Since the ERBB2/HER2 signaling is known to regulate SP1 binding through phosphorylation[34],

and is the target of tyrosine kinases inhibitors such as Poziotinib (**Figure S9A**), we looked through

both lists and noted that ERK was found on the SP1-repressor list (**Figure S9A**). On the Ingenuity

Qiagen ERBB2 Pathway there are 22 genes upstream of *SP1*, and among those, 3 were found on the

SP1-repressor list (*RAS*, *ERK*, *STAT5*, P<0.02). We also used the Molecular Signature Database

(MSigDB) to establish likely pathways from genes that had a -Log10(RRA) score above 0.5 from each

screen (**Figure S9B-S9C**) and observed one shared pathway (Pentose Phosphate Pathway (3P)). Of

interest there were two categories in the SP1-activator analysis that implicate cytosolic calcium

regulation; calcium influx has been previously shown to be an early event in cells exposed to

hypothermic stimuli[35]. As we were particularly interested in a possible transcriptional regulator of

SP1, we annotated our lists and identified 50 (10%) on the SP1-repressor list and 4 (6%) on the SP1-

activator list (**Figure 3E**) as transcriptional regulators. Of these, ten (*GATAD2*, *POU4F3*, *ZNF71*, *KAT5*,

ZNF260, *ANKZF1*, *SMYD5*, *DMRT1*, *TAF13*, *NKX6-2*) had a -Log10(RRA) score above 3,5 in either the

repressor or activator screen (**Figure 3F**). For a gene from our list to act as a direct repressor or

activator for *SP1* we reasoned that regulators from the repressor list should have a known function

as repressors and genes from the activator list should have known activating functions. Four genes

from the list appeared to have opposing functions (*KAT5*, *ZNF260*, *NKX6-2, ANKZF1*, **Figure 3F**, not

bold)[36–39] leaving six genes (**Figure 3F**, bold)[40–46] on our list. Histone modification has previously been

known to dictate temperature-dependent sex determination in reptiles[47] and vernalization in

plants[48], and one of these six targets (*SMYD5*) is a histone methyltransferase, known to catalyze

H4K20, H3K9, and H3K36 trimethylation[44,45]. Interestingly, SMYD5 is a likely interactor with KDM6B in

fish[49] and KDM6B mediates the previously described epigenetic temperature dependent sex

determination switch in reptiles[47]. Also, like CIRBP and RBM3, SMYD5 is a known resident of stress

granules[50]. Since SMYD5 histone tail modifications are known and their levels can be measured at

SP1 and other targets we argued that this might be useful for validation purposes. Thus, we decided

to further focus our efforts on *SMYD5*.

Figure 3. Genome wide CRISPR-Cas9 knockout screen on SP1-MHI reveals multiple potential

inhibitors and activators of the MHR. (A) Overview of the genome-wide CRISPR-Cas9 Knockout

approach for the HEK293WT+Cas9+SP1 cell line. (B) Fluorescence and sort gates of 4 replicates of

transduced HEK293WT+Cas9+SP1 cells (green), negative control (HEK293WT, black) and positive

control (HEK293WT+Cas9+SP1, grey). (C-D) Genes marked with their name in red are transcription

regulators that have a -Log10(RRA) score >3.5 and have a repressive functions and genes marked

with their name in green are transcription regulators that have an activating functions and a

-Log10(RRA) score >3.5. Colored dots represent genes that have a -Log10(RRA) score >2.5, where the

orange dots indicate genes that have a positive log fold change (LFC) and the blue dots indicate genes

that have a negative LFC, from either the SP1 activator (C) or repressor (D) screen. (E) A summary of

transcriptional regulators from our screens, activators (green) on left, repressors (red) on right. (F) A

summary of top 10 transcriptional regulators ranked by their -Log10(RRA score). Genes in bold are

254 known in the literature to act in the same way as the screen results indicated, that is either as a
255 direct repressor (red) or activator (green).

SMYD5 is a direct repressor of *SP1*.

To further explore the role of SMYD5, we took advantage of a newly published dataset[44]
from a CUT&TAG assay in mouse Embryonic Stem Cells (mES) with an overexpressed SMYD5-Flag
plasmid. Upon re-analysis of these data, we noticed that SMYD5 has strong peaks at the promoters
of both *Sp1* (130K) and *Cirbp (70K),* but not *Rbm3* (**Figure 4A**). These peaks occur at evolutionarily
conserved sites and overlap with CpG islands (**Figure 4A**). We took advantage of an RNASeq dataset
on *Smyd5* knockout (KO) mESC from the same study[44] and reanalyzed it which yielded approximately
equal amounts of upregulated and downregulated genes among SMYD5 bound loci (**Figure 4B**)
supporting prior reports that histone marks deposited by SMYD5 can either repress or activate[44,45].
When looking at the three loci (*Sp1*, *Cirbp*, *Rbm3*), we found that *Smyd5*-KO yielded increased
expression of *Sp1* but not *Cirbp* or *Rbm3* (**Figure 4C**) suggesting a direct role for SMYD5 as an *Sp1*
repressor in a murine system. In this dataset, *Cirbp* showed lower expression compared to *Sp1* and
Rbm3 (**Figure S10**) which may impede interpretation. In order to validate these findings in our own
system we decided to knockdown *SMYD5* with siRNA in HEK293WT+Cas9+SP1 cell lines, we observed
an increased fluorescence of SP1-MHI at 32°C and 37°C (**Figure 4D**). *SMYD5*-KO with a guide RNA
(gRNA) and Cas9 significantly decreased amounts of *SMYD5* mRNA expression by qRT-PCR (**Figure 4E**)
and SMYD5 protein levels by Western blot (**Figure 4G**), which are rescued upon overexpression of a
gRNA-resistant Flag-SMYD5 plasmid (Flag-SMYD5 sgRNAres plasmid [labelled SMYD5sgRNA#6res],
Figure 4E and 4G). The same experiment revealed de-repression of the *SP1* transcript in KO cells by
RT-qPCR, with repression restored in rescued cells (**Figure 4F**). Furthermore, we observed a
significant increase of SP1 and CIRBP, but not RBM3 following *SMYD5*-KO at 37°C using Western blot
of the endogenous loci (**Figure 4H-4J**).

Figure 4. SMYD5 is a repressor of SP1 at 37°C. (A) Overexpressed FLAG-tagged SMYD5 in mESc binds at promoters of *SP1* and *CIRBP* but not of *RBM3*. (B) Venn graph showing both up- and down-regulated SMYD5-bound genes in an RNASeq of *Smyd5*-KO cells. Statistical analysis was done with GeneOverlap package in R using Fisher's Exact test. (C) *Smyd5*-KO leads to increased expression of *SP1* but not *RBM3* or *CIRBP* in mESCs. (D) *SMYD5* knockdown by siRNA yields higher levels of fluorescence of SP1-MHI at 32°C and 37°C compared to empty vector control. Shown are data points, each data point is a biological replicate that has been normalized against the same non-transfected HEK293WT+Cas9+SP1-MHI cell line and a mean of data points. Significance levels were calculated with Šidák's multiple comparison test in GraphPad Prism. (E) Relative expression of *SMYD5* compared to GAPDH from *SMYD5*-KO cells, measured by RT-qPCR at 37°C, with or without rescue with Flag-SMYD5 sgRNAres plasmid (labelled SMYD5 sgRNA#6res) which demonstrates that SMYD5-KO was

successful. Shown are data points, each data point is a biological replicate, mean and SD where applicable. Significance levels were calculated with an unpaired one-tailed t-test in GraphPad Prism. (F) Western blot quantification and representative figure using antibodies against SMYD5 and Lamin B in a *SMYD5*-KO cell line with and without rescue with Flag-SMYD5 sgRNAres plasmid (labelled SMYD5 sgRNA#6res) shows that *SMYD5*-KO was successful on protein level at 37°C. Data shown as in (E). (G) Relative expression of *SP1* compared to GAPDH from *SMYD5*-KO cells, measured by RT-qPCR at 37°C, with or without rescue with Flag-SMYD5 sgRNAres plasmid (labelled SMYD5 sgRNA#6res). *SMYD5*-KO leads to increased expression of *SP1* showing that effect is secondary to targeting of *SMYD5* but not something else. Data shown as in (E). (H-J) Western blot quantification and a representative figure using antibodies against SP1, CIRBP and RBM3, respectively, in *SMYD5*-KO cells. Shown are data points, each data point is a biological replicate, mean and SD where applicable. Significance levels were calculated with an unpaired one-tailed t-test in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

SMYD5 shows temperature dependent expression *in vitro* and *in vivo.*

To test whether SMYD5 shows differential expression levels at 37°C and 32°C, we performed a Western blot on HEK293 for SMYD5 using a custom antibody against SMYD5[51] that showed significantly decreased SMYD5 at 32°C (P<0.01, **Figure 5A**). This difference was not observed at the transcriptional level (**Figure 5B**) indicating that SMYD5, is primarily regulated post-transcriptionally. To validate this further we performed immunofluorescence staining where we observe nuclear and cytoplasmic staining of SMYD5 at 37°C (SMYD5; white or green, DAPI; purple, **Figure 5Ca**) and at 32°C (**Figure 5Cb**). The expression of SMYD5 was significantly (p<0.05) decreased at 32°C compared to 37°C (**Figure 5C-5D** and **Figure S11A**). We validated this by using an anti-FLAG antibody in cells with overexpression of SMYD5-FLAG[44], which yielded a similar result (**Figure S11B**). We observed a non-significant difference in nuclear to whole cell SMYD5 mean intensity ratio at the two temperatures (**Figure 5E**), suggesting that the levels are equally affected in both compartments at 32°C. Finally, we show that the brains of mice cooled in a standard regimen used to treat neonatal hypoxic-ischemia[52] show a similar temperature-dependent nuclear decrease in SMYD5 after 6 h of cooling, both in neurons of the dentate gyrus (DG) region of the hippocampus (**Figure 5Fa-d**) and cortical neurons (**Figure 5Fe-h**), indicating a SMYD5-dependent impact on the MHR during standard TTM regimen in mice.

Figure 5. SMYD5 is depleted at 32°C *in vitro* and *in vivo*. (A) Western blot quantification and a representative figure with a SMYD5 antibody at 37°C and 32°C after 6 h incubation. Significance levels calculated with an unpaired two-tailed t-test in GraphPad Prism. Shown are data points, each data point is a biological replicate, SD and mean are depicted where applicable. (B) qRT-PCR results for SMYD5 expression at 37°C and 32°C after 6 h incubation. Significance levels calculated with an unpaired one-tailed t-test in GraphPad Prism, otherwise data shown as in (A). (C) Intensity of endogenous SMYD5 at 37°C (top, a) and 32°C (bottom, b) after 6 h incubation. Nuclei are marked with a dotted line. (D) Cellular SMYD5 mean intensity levels. Data shown as in (B). (E) SMYD5 nuclear to whole cell mean intensity ratio. Data shown as in (B). (F) SMYD5 staining in brain slices of mice kept at euthermia (37°C) compared to mice that had been cooled down to 32°C for 6 h, mouse model of TTM. Brain slices are from the hippocampal DG (a-d) and cortical (e-h) region. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Epigenetic and transcriptional consequences of temperature-dependent changes in SMYD5.

As SMYD5 is decreased at 32°C and SMYD5-binding peaks overlap a CpG-island (**Figure 4A**),

we performed DNA methylation (DNAm) analysis using the 850 K EPIC Illumina array and observed

limited DNAm at 16/18 sites and no differential DNAm at *SP1*, *CIRBP* or *RBM3* promoters at 32°C for

6 h compared to 37°C at CpGs spanning each of the gene regions (**Figure S12** and **Supplementary**

Table 5). However, there was differential DNAm at other CpGs loci after 6 h exposure at 32°C

(**Supplementary Table 6**) supporting the idea that the MHR is regulated by the epigenetic machinery.

To further explore the epigenetic consequences of differential SMYD5 availability at 37°C versus

32°C, we performed CUT&RUN using antibodies against H3K36me3, H4K20me3, H3K9me3,

H3K4me3, H3K27me3, SMYD5 and H3 (control) in HEK293 cells and human neural progenitor cells

(NPCs). We chose H3K36me3, H4K20me3, and H3K9me3 since these modifications have been

previously shown to be deposited by SMYD5[44,45]. We also chose to look at H3K4me3 and H3K27me3

as these are general open and closed chromatin modifications that are also known to change during

vernalization in plants[53]. We saw less H3K36me3 over *SP1* at 32°C compared to 37°C in NPCs (**Figure**

6A) and HEK293 cells (**Figure S13A**) as well as decreased global SMYD5 binding (**Figure 6B**), all

consistent with lower SMYD5 amounts at 32°C (**Figure 5**). We further observed a global deficiency of

H3K36me3 at 32°C compared to 37°C seen in both HEK293 cells (**Figure 6C**) and NPCs (**Figure S13B**)

albeit more prominent in NPCs indicating that this modification is used to regulate additional genes

perhaps through SMYD5 but levels of SMYD5 mirror this effect (**Figure 6B** and **Figure 5**). We also

observed a global H3K4me3 deficiency at 32°C compared to 37°C (**Figure 6D**), although there were

also 1288 genes that had a unique H3K4me3 modification at 32°C which was not seen at 37°C

(**Supplementary Table 8**). For H3K9me3 we saw a deficiency of histone modification marks upstream

of gene bodies in NPCs (**Figure 6E**). We did not observe a difference for other histone modifications

when comparing 32°C and 37°C in NPCs (**Figure S13C-S13D**). CUT&RUN for H3K36me3 also revealed

several significantly differentially modified genes at 32°C when compared to 37°C (**Figure 6F**, **Figure**

S13E, **Figure S14**, and **Supplementary Table 7**) including a number of epigenetic regulators, such as

BAZ2A and *SETD1B* (**Figure 6F**, **Figure S14A**, and **Figure S14E**). To identify additional genes regulated

by SMYD5 and temperature, we performed an RNASeq in mouse NPCs (mNPCs) at the two

temperatures. This yielded 2890 differentially expressed genes (DEGS) that were upregulated and

3133 DEGS that were downregulated at 32°C compared to 37°C. Interestingly, when we overlap

upregulated DEGS in mNPCs at 32°C with SMYD5-bound and repressed genes (*Smyd5*-KO

upregulated) at 37°C in mESCs we observe a statistically significant overlap of 26 genes (P=5.4e-04,

Figure 6G). Similarly, when we overlap down-regulated DEGS in mNPCs at 32°C with SMYD5-bound

and upregulated genes (*Smyd5*-KO downregulated) at 37°C in mESC we also observe an overlap of 19

genes (P=0.029, **Figure 6H**). This suggests that SMYD5 regulates additional genes in a similar manner

to *SP1* (as a repressor at euthermia) as well as regulating some in an inverse manner (normally as an

activator at euthermia) in murine models. This includes several genes upregulated at 32°C with a

theoretically appealing role for hypothermia responses such as *Klf15* (role in brown adipose tissue

metabolism)[54] and *Thrap3* (role in thyroid metabolism and adipocyte differentiation)[55,56]. Among the

genes normally upregulated at 37°C but downregulated at 32°C, *Bnip3L*, a pro-apoptotic gene[57] is of

particular interest given the previously observed cytoprotective role of mild hypotherma[9–13]. We

provide the entire list of 45 SMYD5-temperature dependent genes in **Table 1.** When we looked more

closely at the list of genes that had unique H3K4me3 modification in NPCs at 32°C and overlapped

those genes with human orthologs of upregulated DEGS in mNPCs at 32°C we saw a significant

overlap of 169 genes (P=2.4e-03, **Figure 6J,** and **Supplementary Table 9**). Only 6 genes are bound by

SMYD5 and upregulated with *Smyd5*-KO at 37°C in mESCs as well as upregulated at 32°C in mNPCs

(**Table 1**) and have a unique H3K4me3 peak at 32°C in NPCs (**Supplementary Table 8**). Interestingly

Asxl1, a member of the Polycomb family[58] a family which is known to take part in vernalization in

plants[48], is among these 6 genes. This indicates that there may be additional epigenetic MHR-

responses that are independent of SMYD5. Finally, we demonstrate that on average these 45 genes,

regulated by SMYD5, have lower H3K36me3 modifications at 32°C versus 37°C at their gene bodies as

well as fewer modifications over the 3ʹ-UTR region at 32°C in NPCs and HEK293 cells (**Figure S13F-**

S13G). In addition, these genes also show a different H3K4me3 and H3K9me3 modification pattern at

32°C and 37°C in NPCs when looking at their 3ʹ- and 5ʹ-UTR regions (**Figure S13H-S13J**).

Figure 6. SMYD5 is a novel epigenetic gatekeeper of the MHR. (A) Decreased H3K36me3 is observed
at the *SP1* promoter and gene body at 32°C compared to 37°C. (B) SMYD5 chromatin binding is
globally decreased (all genes) at 32°C compared to 37°C. (C) There is a global deficiency of H3K36me3
at 32°C compared to 37°C. (D) There is a global H3K4me3 deficiency at 32°C compared to 37°C. (E)
Upstream of gene bodies there is a global H3K9me3 deficiency at 32°C compared to 37°C. (F) At the
gene-specific levels there are a number of genes, that have significantly altered H3K36me3 at 32°C
compared to 37°C in NPCs. Red circles indicate genes that have Log_2 fold change over 1 and p-value
less than 10^{-6}, gray circles indicate Log_2 fold change over 1 but non-significant p-value ($>10^{-6}$) and
black circles indicate non-significant Log_2 fold change and p-values for those genes. (G-H) DEGS from

398   RNASeq in mNPCs at both temperatures show overlap with SMYD5-bound genes that are

399   differentially expressed upon *Smyd5*-KO in mESCs at 37°C. Statistical analysis was done with

400   GeneOverlap package in R using Fisher's Exact test. (J) There is a significant overlap of DEGS and

401   genes that have a unique H3K4me3 modification at 32°C. Statistical analysis as in (G-H).

SMYD5-repressed genes ↑ at 32°C (SMYD5 represses at 37°C)

Gene	Function	Role
Mybl2	Transcription factor	Cell-cycle control
Klf15	Transcription factor	Brown-tissue metabolism
Thrap3	Transcriptional co-activator	Thyroid hormone receptor binding
Tex15	DNA repair protein	Role in testicular meiosis
Ewsr1	RNA binding	Role in Ewing sarcoma
Tfcp2l1	Transcription factor	Pluripotency in stem cells
Amotl2	Angiostatin binding factor	Role in angiogenesis
Phf20	Methyl-lysine-binding protein	Histone modification
Cd24a	Sialoglycoprotein	Cell differentiation
Nt5e	Nucleotidase	Nucleotides metabolism
Neo1	Cell surface receptor	Immunoglobulin superfamily
Gabbr1	GABA receptor	Inhibitory neurotransmitter
Asxl1	Chromatin binding protein	Polycomb family member
Tulp4	Tubby Superfamily Protein	Ubiquitination function
Pkn1	Serine/threonine-protein kinase	Protein kinase C superfamily
Hnrnpul1	RNA-binding protein	hnRNP family
Mier1	Transcriptional regulator	Mesoderm induction early response
Fam21	WASH complex member	Endosomal protein sorting
Trip12	E3 ubiquitin-protein ligase	Ubiquitin regulation and DNA repair
Lamc1	Extracellular matrix glycoprotein	A laminin
Rell1	RELT-like protein	Regulates the p38MAPK cascade
Xrn2	5'-3' exonuclease	Promotes transcription termination
Hspg2	Multidomain proteoglycan	Encodes Perlecan protein
Ctc1	Telomere maintenance protein	Limits telomere degradation
Stag1	Component of cohesin complex	Role in cohesion of sister chromatids
Ass1	Enzyme	Role in urea cycle

SMYD5-upregulated genes ↓ at 32°C (SMYD5 activates at 37°C)

Gene	Function	Role
Usp28	Ubiquitin Thioesterase	DNA damage checkpoint
Ccnd1	Cyclin D	Cell cycle regulation
Dzip3	Ubiquitin-protein transferase	Spermatogenesis
Rad50	DNA double-strand break repair	Cell cycle regulation
Arglu1	Enables cadherin binding	Mediator complex
Sltm	RNA binding activity	Regulation of transcription
Slc16a3	Solute transporter	Lactic acid and pyruvate transport
Ier5l	Unknown function	Parolog of IER5
Acly	Enzyme	Synthesis of cytosolic acetyl-CoA
Agps	Enzyme	Lipid biosynthesis
Apobec3	Cytidine deaminase	RNA editing

Ivns1abp	Viral binding protein	Response to viral infection
Dus3l	Synthesis of dihydrouridine	RNA editing
Upp1	Uridine phosphorylase	Ribonucleosides dynamics
Ldha	Enzyme	Lactate to Pyruvate conversion
Scd2	Enzyme	Fatty acid biosynthesis
Smarcad1	Chromatin remodeler	Role in heterochromatin formation
Bnip3l	Pro-apoptotic protein	Induces apoptosis
Esrra	Nuclear receptor	Mitochondrial biogenesis regulator

Table 1: Summary of temperature responsive SMYD5-regulated genes.

Discussion

Histone modification has previously been found to help integrate the effects of cold

temperature exposure in diverse organisms. For example, in plants, vernalization, the basis of how

cold exposure influences the rate of flowering, is mediated through a Trithorax-Polycomb switch[48]. In

some reptiles, sex determination is achieved through a temperature dependent mechanism and for

one such, the red eared slider turtle (*Trachemys Scripta Elegans*), this regulation acts through a

histone methylation switch[47]. To uncover regulators of the mammalian MHR we used our novel MHIs

to identify an FDA-approved drug, Entacapone, that can induce the SP1-mild hypothermia response

as well as conducting a forward CRISPR-Cas9 mutagenesis screen resulting in the identification of

SMYD5 as a repressor of the MHR at euthermia. Such insights can be useful to expand the

mechanistic understanding of the MHR, but our data also provide both a feasible current strategy

(Entacapone) as well as a future therapeutic target (SMYD5) for efforts to activate the MHR without

cooling in patients at risk of neurological sequela after catastrophic events. Finally, a detailed

understanding of the mammalian MHR is important to understand how temperature can

systematically bias results in biology and medicine. One such example involves the process of

harvesting tissue from human cadavers but uniformly these are kept at a low temperature which

may activate the mild hypothermia response in the cells being harvested. Similarly, countless

researchers store samples at room temperature between experimental steps which in some

scenarios may activate the MHR. This contrasts with storage on ice but in our own hands that does

not appear to activate the MHR. Such data may be of use to a large number of scientists.

Our current MHIs, a novel tool for the MHR community, allow for some temporal dissection,

yet they all currently harbor stable fluorescent protein, which may capture average responses over

time rather than dynamic difference, but the latter can be detected by real time qPCR. Thus, our

current MHIs may be the most useful to map the start of the response (as we have done in **Figure 1**).

However, in future work, we hope to create unstable fluorescent molecules that break down quickly

to help us map the kinetics of different parts of the response, for example, how quickly the response

is resolved (i.e., termination).

Although our drug screen and validation may have limitations (only three timepoints, single

dose), we observe an overrepresentation of TKIs. The reason why some TKIs as well as other drugs

did not validate could be that the drug screen was performed using an MHI containing the MCRE (to

increase signal) but validation was performed on an MHI lacking the MCRE (the endogenous situation

in mammals[15]). Of the TKIs we have further shown that Poziotinib, a known pan-HER inhibitor[29], was

able to increase *RBM3* mRNA but not *SP1* or *CIRBP* mRNA (**Figure S7**) or SP1 (**Figure 2F**) at 37°C with

the doses tested. The lack of increased SP1 quantified by Western blot could be caused by the

relatively low (4 µM) Poziotinib dose compared to what was used for RT-qPCR and flow cytometry

(20 µM). Here, we identify and validate Entacapone as a compound that can activate the SP1-MHR at

euthermia (**Figure 2E**). Entacapone is a COMT-inhibitor, which is frequently used in the clinic due to

its ability to increase dopamine availability by blocking its degradation in synapses[59]. It is interesting

that synaptic plasticity is known to be influenced by temperature changes[60]. This suggests that

Entacapone may act at the cell membrane since Entacapone is a neurotransmitter breakdown

inhibitor (COMT), which acts at the neural synapse. In fact, several candidates (**Figure 3C-3D**) from

the CRISPR-Cas9 forward mutagenesis screen appear to operate at the neural synapse or bind to

Syntaxin (**Supplementary table 3**) such as *STXBP1* (SP1 repressor) and *TXLNB* (SP1 activator). Both

bind to Syntaxin (**www.genecards.org**) and play a role at the neuronal synapse as well as having a

role in dopaminergic synaptic transmission[61], and such distal regulation may be worth exploring in

future work.

Our data support the notion that SP1 and RBM3 may be parts of distinct arms of the

mammalian hypothermia response as SMYD5 only regulates *SP1* but not *RBM3*. We interpret the lag

in the response (at least 6 hours to signal) and the obvious shift in fluorescence of mutagenized cells

to mean that there are likely many genes upstream of the currently tested point of the response

(*SP1*). Despite a global shift towards activators, the yield from the enrichment screen yielded more

putative *SP1* repressors compared to *SP1* activators. This could be caused by the screen setup as it

may be easier in general to find genes that increase fluorescence (repressors) than those that

decrease fluorescence (inhibitors), as the latter will overlap with cells that have lost the SP1-MHI for

various reasons. Another limitation of our forward screen is that it can only identify cell-autonomous

genes and cannot find essential genes. However, it is a starting point, and here we provide 556

candidate regulators that can be subsequently validated using our MHIs or other strategies. Of

interest, SMYD5 and ERBB2/HER2 (the receptor Poziotinib blocks) have been previously found to

functionally interact in a forward mutagenesis screen of genes that play a role in metastatic

reactivation and tumor dormancy[62] and additionally, *CIRBP* transcript levels correlate with prognosis

in luminal A/B breast cancer, which is influenced by HER2 status[63]. In fact, our SP1-repressor screen

revealed several downstream factors of ERBB2 signaling (*RAS*, *ERK*, *STAT5*, **Figure S9A**), indicating

that this pathway may be upstream of SMYD5 and worth exploring. Alternatively, SP1 itself may be a

target of phosphorylation, and as SP1 is predicted to bind at its own promoter[64] and perhaps both

signals are required for full effect.

Here, by uncovering *SMYD5* as a negative epigenetic regulator of SP1, a key factor of the

mammalian MHR, we provide an example of how histone methylation integrates temperature cues

into a fundamental mammalian response. Our present model is that SMYD5 exerts its effects at 37°C

to ensure that *SP1*/*CIRBP* are not upregulated without a cold stimulus, however, upon cold stimulus

(32°C), SMYD5 is degraded and SMYD5 repression is released, thus activating the MHR. The nuclear

SMYD5 repression release upon mild hypothermia could happen because of one of the following:

nuclear exclusion of SMYD5, less *SMYD5* mRNA being translated, less protein synthesis, or SMYD5

could undergo proteasomal degradation. Per bioGRID, SMYD5 is known to bind to four ubiquitinators

(PARK2, UBA52, TRIM25, USP34) so this may be an avenue for further study as such factors often

target proteins to the proteasome. Previously, a generalized decrease of protein production upon

cold stimulus has been shown to act through AMPK mediated phosphorylation of eEF2K[22], and thus

decreased translational efficiency could play a part. However, we are unaware of biological examples

of targeted protein degradation in response to mild hypothermia.

SMYD5 itself appears to regulate at least 45 other factors that show temperature

dependence, 26 in which SMYD5 acts as a repressor (like *SP1*) and another 19 where SMYD5 acts as

an activator at euthermia. Some of these genes look like very plausible MHR participants, especially

BNIP3L,[57] a pro-apoptotic factor that is downregulated with hypothermia and may thus contribute to

the neuroprotective/cytoprotective effect of mild hypothermia. There are also genes that play a role

in thyroid and adipocyte function (*THRAP3*)[55,56], brown adipose tissue (*KLF15*)[54] and testes

(*TEX15*)[65,66], all target tissues of physiological mammalian temperature responses. Coinciding with

decreased global H3K4me3 at 32°C, we also observed decreased H3K36me3 at 32°C over the gene

SETD1B (**Figure 6F** and **S14E**) which encodes an H3K4me3-methyltransferase. In oocytes, *Setd1b*-KO

can cause gain or loss of H3K4me3[67]. Accordingly, this could be one potential mechanism underlying

H3K4me3 changes at 32°C and warrants further investigation. It is interesting that H3K36me3 has

previously been shown to regulate temperature-regulated alternative splicing in plants[68] and in the

fungus *Saccharomyces cerevisiae,* H3K4 and H3K36 methyltransferases (Set1 and Set2) regulate

alternative polyadenylation[69]. Furthermore, in mice, H3K36me3 has a direct impact on alternative

splicing of neurobiological factors in the brain[70]. In mammals' temperature dependent alternative

splicing[71] and alternative splicing of *CIRBP* and *RBM3* have been observed[72–74]. Perhaps alternative

splicing or polyadenylation play a role in controlling levels of SP1 or one of the other 45 SMYD5-

bound factors elucidated here.

In summary, we have developed and validated novel tools to interrogate the MHR and used

them to show, for the first time, that a compound exists that can activate the MHR without cooling.

We provide a list of 556 candidate of the MHR pathway and validate *SMYD5* as a temperature-

dependent epigenetic gatekeeper of the MHR and, for the first time, show how histone methylation

(H3K36me3, H3K9me3, and H3K4me3) changes in mammals with MHR exposure, providing a specific

example of how mammalian cells use epigenetics to respond to temperature cues.

Acknowledgements: We thank SCRU lab (UI) for gifting us pMD2.G (Addgene, #12259) and psPAX2

(Addgene, #12260) and H.C. Dietz and D. Valle labs (JHU), for gifting us the HCT-116, SK-N-SH, HeLa,

Jurkat and K562 cell lines. We thank Alan Long at the ChemCore facility at JHU and the Pharmacy of

Landspitali for support regarding the drug screen and validation. We thank the Immunology

department of Landspitali University Hospital for the access and assistance with the FACS sorting. We

thank FUNlab at deCODE genetics for the usage of their flow cytometry machines as well as for

gifting us the HEK293T cell line. We thank Kári Stefánsson and Ólafur Þ. Magnússon at deCODE

Genetics for providing some of the next generation sequencing for this project. We thank Erna

Magnúsdóttir for providing pAG-MNase for the CUT&RUN assays as well as guidance with the

CUT&RUN experiment and data analysis. This study was supported with funding from the Rannis

Technology Development Fund (H.T.B., #2010588), Eimskip University Fund (S.R.), Fulbright Iceland

(S.R.), Icelandic Cancer Society Research Fund (H.T.B.), Göngum Saman Research Fund (H.T.B.),

Landspitali Research Fund (H.T.B.), Félag Háskólakvenna Research Fund (S.R.). H.T.B. is also

supported by funding from Rannís (#217988, #195835, #206806) and the Louma G. Foundation. R.C-

V. is supported by Thomas Wilson Foundation and the Johns Hopkins University Pediatric Innovation

Grant. R.C-V. and F.J.N. are supported by the NIH R01NS126549, 1R21NS123814, and 1R01

HD110091 and F.J.N. additionally by R01HD074593, U01NS114144. Finally, some images in this

manuscript were created with BioRender.com.

Author contributions: H.T.B. and S.R. conceptualized the project. S.R., L.Z., and H.T.B. created and

designed the MHIs. S.R., A.T., L.H.A., M.V., K.E.A., and K.J. performed cell culture. S.R., A.T. L.H.A.,

and M.V. performed flow cytometry experiments. L.Z. and H.T.B. designed the approach to the drug

screen. S.R., and H.T.B. designed the approach to the forward mutagenesis assay. S.R. performed the

forward mutagenesis assay and drug screen. S.R. and K.J.A. performed the PCR for the NGS. S.R., A.T.,

and K.J. validated results from drug and mutagenesis screens (Western blots (A.T., K.J.), RT-qPCR

(S.R., K.J.) and flow cytometry (S.R., A.T.)). K.J. and K.M. performed IF on cells from culture. K.J. and

K.M. performed the confocal imaging and K.M., K.J. S.Th.H., and S.R. did the data analysis. S.R. and

M.V. performed siRNA knockdown. R.C-V. and F.J.N. produced the euthermic and hypothermic

mouse pups, performed microdissections and perfusions. K.J.A. isolated and cultured mNPCs. M.V.

performed cryosectioning and IF on mouse brain slices and M.V. and K.M. performed confocal

imaging as well as the analysis of the mouse brain slices. S.R., A.T., and S.Th.H. performed CUT&RUN

experiments. K.J.A. did the mNPCs isolation and RNASeq and data analysis. S.R., K.J.A., K.J., A.T.

S.Th.H., and T.R. analyzed data. J.H. provided SMYD5 antibody and K.E.A. provided human NPCs. S.R.
and H.T.B. wrote the manuscript and all authors critically reviewed the manuscript. Project
administration was carried out by H.T.B, S.R., and K.J.A.

Declaration of interest: H.T.B. is a consultant for Mahzi therapeutics. S.R. and H.T.B. have a European
patent application (number 23167505.9).

STAR Methods

Creation of novel indicators: We used the PGL4.10 plasmid (Promega, #9PIE665) as the backbone for
our CIRBP- and SP1-MHI, where the *luc2* sequence was cut from the plasmid with NcoI (NEB) and
XbaI (NEB). Promoter sequence for *CIRBP*[75] was cloned from mouse DNA using *CIRBP* forward primer
5'-tcgataggtaccTGGCTTCACAAATGCGCCTCAGT-3' and *CIRBP* reverse primer 3'-
cctaaggcagatctGCGAGGGGGAGCGCAAGAGT-5'[75]. Restriction enzymes KpnI (NEB) and BglII (NEB)
were used to insert the promoter into the plasmid. Promoter sequence for *SP1*[76] was cloned from
human DNA using the forward primer 5'-tcaagtcaggctagcGCAACTTAGTCTCACACGCCTTGG-3' and
reverse primer 3'-cagtgctgcctcgagGCTCAAGGGGGTCCTGTCCGG-5'[76]. Restriction enzymes NheI (NEB)
and XhoI (NEB) were used to insert the promoter into the plasmid. AcGFP1 was cloned from
PT7XbG2-AcGFP1 (Novopro, #V002843) vector using the reverse primer 3'-
cggcggagTCTAGAATTACTTGTACAGCTCGTCC-5' and the forward primer 5'-
taagccaccATGGTGAGCAAGGGCGAGGAGC-3'. Restriction enzymes NheI (NEB) and XhoI (NEB) were
used to insert the acGFP1 into the plasmid. The neomycin selection cassette was cloned from
pROSA26-dest plasmid (Addgene, #21189) using the forward primer 5'-
CATTATCGTCGACTCTACCGGGTAGGGGAGGCGCTT-3' and reverse primer 3'-
CGCCGCCGACGATAGTCAAGCTTCTGATGGAATTAGAACTTGGC-5'. Restriction enzymes SalI (NEB) and
PshAI (NEB) were used to insert the neomycin cassette into the plasmid. CIRBP- and SP1-MHI were
made with and without MCRE[15]. The MCRE sequence was inserted in 5 repeats in front of the
promoter sequences with a linker sequence between. Acc65I (NEB) restriction enzyme was used to
insert the MCRE sequence into the CIRBP-MHI and Acc65I (NEB) and SacI (NEB) restriction enzymes
were used to insert the MCRE into the SP1-MHI. Sanger Sequencing was used to validate insertion
sites, size, and sequence of insertion. The *RBM3* promoter with MCRE enhancer (6 repeats) in front
of the promoter was cloned into the multiple cloning site pCMV6-AC-GFP (BlueHeron, #PS100010).
We also made lentiviral vectors for *CIRP*, *SP1* and *RBM3* on a pLV-Ex (VectorBuilder) lentiviral vector.
The *CIRBP* promoter for the CIRBP-lenti-MHI was designed from the human promoter and the *SP1*
promoter was moved upstream of the TSS for the SP1-lenti-MHI but *RBM3* promoter sequence
remained unchanged. Included were MCRE (8 repeats for CIRBP-lenti-MHI, five repeats for other),
promoter, neomycin selection cassette and a mCherry fluorescent protein behind each promoter
sequence. The promoter region, size and backbone plasmid for each indicator is detailed in
Supplementary Table 1 and sequence of each MHI in **Supplementary File 1**. We also observed some
variation in absolute levels among individual indicators/experiments, but the pattern was always
consistent; given this, we only show data from individual experiments in any figure.

Culture of cell lines: HEK293 (gifted and DSMZ, CRL-1573, referred to as 293/293WT) cells were
cultured in DMEM/F12 with GlutaMAX (Gibco, 10565018) with 10% filtered FBS (Gibco, 10270106)
under 5% CO_2. Cells were cultured at 37°C unless otherwise stated, then 37°C served as a control. In
addition to HEK293, we used the following cell types HEK293T (gifted, CRL-3216), HCT-116 (gifted,
CCL-247), HeLa (gifted, CCL-2), Jurkat (gifted, TIB-152), K562 (gifted, CCL-243), SK-N-SH (gifted, HTB-
11). Cells were sub-cultured when they had reached 70-90% confluence and media was changed
every other day. Killing curves were used for all cell lines to determine lowest selection concentration
for all selection agents. We used culture recommendations from ATCC when possible. We also used
human iPS cells (ATCC, ACS-1019) that were used to generate NPCs by neural induction of iPSCs using
dual SMAD inhibition and embryoid body generation (StemCell Technologies; 08581, 05832,
05838)[77], prior to experiments described here the NPCs were cultured in STEMdiff Neural Progenitor

Medium (StemCell Technologies, 05833) on Matrigel (Corning, 354234). As well as murine primary
NPCs (see section mNPC isolation for further information).

Transfection of MHI and SMYD5-Flag overexpression plasmid: HCT116, 293WT, HEK293T, HeLa and
SK-N-SH cell lines were transfected with MHI when they had reached 70-90% confluency. We used
either Lipofectamine™ 2000 or 3000 (Thermo Fisher Scientific, 11668027, L3000015) according to
the manufacturer's protocol. For transfection of K5622 and Jurkat cell lines we used electroporation
for transfecting the MHIs into the cell lines. The fluorescence of the indicators was measured via flow
cytometry up to 48 h after transfection. We analyzed flow cytometry data with the FlowJo Software,
where we gated living cells, single cells, GFP positive cells and measured the fluorescence for that
population. For further analysis, we used Excel from Microsoft, GraphPad Prism (v.9) and FlowJo
(v.10.8.1). For transfection of SMYD5-Flag overexpression vector[44] we used the same lipofectamine
method as described above.

Construction of 293WT+Cas9 and 293WT+Cas9+SP1 cell lines: We made two cell lines from 293
cells, one that expressed Cas9 (lentiCas9-blast, Addgene #52962), referred to as 293WT+Cas9, and
another that expressed both Cas9 and SP1-MHI, referred to as 293WT+Cas9+SP1. These cell lines
were made by transfecting the plasmids with lipofectamine as described above and selecting stably
transfected cells using Blasticidin selection for Cas9 and Neomycin selection for SP1-MHI.

Drug screen: Briefly, we used Lipofectamine 3000 (Thermo Fisher Scientific, L3000015) to transfect
the SP1+MCRE-MHI into the 293T cell line. 24 h after transfection, 30000 cells were plated into each
well of a black 96 well plate. 48 h after transfection the cells were exposed to FDA Approved Drug
Screening Library (L1300-Z298012) via the High Throughput Screening Services, ChemCore Facilities
(Johns Hopkins University). Each drug was exposed in 20 µM final concentration. Total well GFP was
read with the CLARIOStar Plus Plate Reader (BMG LABTECH, settings: gain 1500, 470-15 excitation
and emission 515-20 at focal height 2.6) 16, 28, and 40 h after exposure of the library. Ratio of
increased fluorescence was calculated by dividing GFP reads from positive control (SP1+MCRE-MHI
transfected but untreated) at each time point with GFP reads of each whole well exposed to drug at
the same timepoint for wells in the same lane. For further analysis, we used Excel from Microsoft and
GraphPad Prism (v.9).

Validation of drug screen: We individually validated 8 compounds in same concentration (20 µM).
Here, we exposed the drug for 16 h either at 32 or 37°C, and evaluated by flow cytometry as
previously described. In a 96-well plate 20,000 of either 293 or 293+SP1-MHI cells were seeded per
well in a total volume of 200 µL of media. After cells had attached to the wells (the morning after),
cells were exposed to 20 µM of each drug, controls were exposed to the same quantity of DMSO as
drug, for 16 h and harvested for flow cytometry analysis. Flow cytometry analysis was conducted as
previously described. Drugs that were tested for further validation after initial screen were: Orantinib
(Selleckchem, S1470), Dasatinib (Selleckchem, S7782), Poziotinib (Selleckchem, S7358), Tamoxifen
Mylan (Mylan AB*), Erythromycin (Stragen Pharma GmbH*), Naloxone (B. Braun*), Entacapone
(Orion Corporation*) and Cinacalcet (Ratiopharm GmbH*). Drugs labeled with "*" were
pharmaceutical grade medications gifted from the Pharmacy at Landspítali University Hospital.
Analysis was performed as described above.

Apoptosis assay: We based our apoptosis assay on a protocol published by Xiang et al.[78]. We exposed 293WT+Cas9+SP1 cell line either to 1-5 µM H_2O_2 or vehicle (H_2O) for 4 h at 32°C or 37°C. 4 h after exposure gMFI of the cells were analyzed with flow cytometry as described above.

Cell cycle assay: The cell cycle assay was based on a protocol from Rutgers Flow Cytometry/Cell Sorting & Confocal Microscopy Core Facility (available online)[79]. HEK293 cells were seeded in four 12-well culture plates. The day after, the media was replaced with fresh media containing Poziotinib, Entacapone, or DMSO (vehicle) either at 32 or 37°C. 16 h later, the cells were harvested. To include dead cells in the analysis, the media and the PBS used to wash the cells were collected with the cells. Cells were fixed with 1% paraformaldehyde for 1 hour at 4°C and then permeabilized with 66% EtOH at 4°C for 2 hours. Next, cells were stained with 1 x propidium iodide + RNase staining solution (Abcam, ab139418) and incubated at 37°C in the dark for 30 minutes. Cells were placed on ice and underwent flow cytometry.

Genome Wide CRISPR-Cas9 screen on fluorescent HEK293WT cell lines that stably express SP1-MHI: Using the 293WT+Cas9+SP1 and 293WT+Cas9 (negative control) cell lines we performed a genome wide CRISPR-Cas9 knockout screen (GeCKO) screen, where we transduced the sgRNA pool made from library A and B (Addgene, #1000000049) in MOI 0.3[17–19]. We mostly followed the Joung et al. protocol[19], the exceptions are as follows. For making the lentivirus, a protocol described by Kutner et al.[80] was used. Lentiviral sgRNA library concentration was done with Amicon ultracentrifugal filters (Millipore, UFC9003). 20 h after transduction with the sgRNA library selection was started with Puromycin and continued for 6 days and 8 h before the cells were moved to 32°C, where they were incubated for 16 h before fluorescent activating cell sorting (FACS) with Cell sorter SH800Z (Sony). The 5% highest fluorescent cells for 293WT+Cas9+SP1 and 5% lowest fluorescent cells for 293WT+Cas9+SP1 were sorted. Next, we isolated gDNA and performed a two step PCR for all samples (sorted as well as negative control). According to cell number after puromycin selection, the sgRNA library coverage was >700 times coverage for HEK293WT+Cas9+SP1. Next generation sequencing (NGS) was performed with single read, 80 cycles and 8 indexing cycles with PhiX spike in of 20% on NovaSeq 6000 S4 at deCODE genetics for HEK293WT+Cas9+SP1 and HEK293WT+Cas9, 32-280 million aligned reads per sample. To analyze the screen data, single read fastq files for each replicate and condition were merged using cat command. Then, MAGeCK (v.0.5.9) was used to identify enriched sgRNA's in sorted samples[33,81] and sgRNA counts were normalized to internal control for sgRNA's (control sgRNAs). We used RStudio (R, v.4.1.1)[82] to visualize the results and to filter genes. We exported a list of genes that had both FDR value under 0.25 and LFC value over 2.5 for each condition in both screens. Then, we excluded genes, overrepresented in both the repressor and activators screen when compared to control as they most likely promote cell growth and therefore were present in both lists (**Supplementary Table 4**). For the pathway enrichment analysis we used EnrichAnalyzer() from MAGeCKFlute (v.1.12.0)[81].

Making of *SMYD5* knockout (*SMYD5*-KO) cells: Two ready-made sgRNA vectors (Genscript, *SMYD5* plasmid vector #2 and #6) were packed into lentivirus according to protocol described by Kutner et al.[80], for concentration we used the 4x Lentivirus concentrator solution protocol from the MD Anderson Cancer center at University of Texas (available online)[83]. We selected for stable *SMYD5*-KO cells with Puromycin selection for at least 7 days.

Making of *MALT1* and *RGS21* knockout cells: Ready-made sgRNA vectors (Genscript, plasmid vector *MALT1* #1 and #2, *RGS21* plasmid vector #2 and #3) were packed into lentivirus and selected as described above in the making of *SMYD5-KO* cells.

Making of gRNA resistant SMYD5-Flag plasmid (Flag-SMYD5 sgRNAres plasmid)**:** In an effort to make SMYD5-Flag resistant to CRISPR-Cas9 cutting from the two guides (Genscript, *SMYD5* plasmid vector #2 and #6), we used site-directed mutagenesis (Q5® Site-Directed Mutagenesis Kit from NEB, E0554S) to modify predicted PAM sites of the guides. We used the following primers to induce the mutations: SMYD5-2mut-F: 5'-gctctttacgAGGAAGCAGTCAGCCAGT-'3; SMYD5-2-mut-R: 5'-ttccgtaaagAGTCTCCGCAGAAGTTCC-'3; SMYD5-6-mut-F: 5'-accgatatcgAGCCTGTGACCACTGCCT-'3; SMYD5-6-mut-R: 5'-atagagcgtt-CCAGAGAAACTGTGCAGCC-'3. After site-directed mutagenesis, we isolated plasmid with Monarch Midiprep kit (NEB, T1010) and Sanger sequenced to verify changes in PAM-sites. This plasmid was then used for rescue experiments (**Figure 4E-4G**).

siRNA: We used the TriFECTa RNAi kit (IDT; TYE 563 Transfection Control DsiRNA, 51-01-20-19 (transfection control); Negative Control DsiRNA, 51-01-14-03 (negative control)) and used predesigned DsiRNA (IDT, hs.Ri.SMYD5.13.8) to knockdown *SMYD5* mRNA in two biological cell lines of HEK293+Cas9+SP1-MHI. For the reverse transfection we used Lipofectamine 2000 (Thermo Fisher Scientific, 11668027) where we seeded $1*10^6$ cells per well in a 6 well plate and exposed to siRNA. Around 24 h after the reverse transfection one batch of cells was exposed to 32°C and the other kept at 37°C for 6 hours before the cells underwent flow cytometry. Further statistical analysis of the flow cytometry data was similar to that described above.

Immunofluorescence of cell culture: HEK293 cells were seeded into two 8-well chamber slides (Falcon, 354118) and cultured for 24-48 h before one of the chamber slides was moved to 32°C. After 6 h cells from both slides at 32°C and 37°C were fixed and permeabilized with 4% PFA and 0.1% Triton-X 100 diluted in PBS for 10 minutes at room temperature. Then cells were washed twice with a blocking solution (PBS with 2.5% BSA and 10% normal goat serum) and followed by blocking for 30 min at room temperature in the blocking solution. Cells were incubated overnight with primary antibodies (rabbit anti-SMYD5, 1:1000[51]; anti-FLAG, 1:500, (Sigma-Aldrich, F1804-200UG)) at 4°C. The following day, cells were incubated at room temperature for 30 mins and then washed three times every 10 min with a washing solution composed of PBS with 0.25% BSA. Cells were incubated with secondary antibodies (goat anti-mouse IgG (H+L) Alexa Fluor™ 488, 1:1000 (Invitrogen™, A-11001); goat anti-rabbit IgG (H+L) Alexa Fluor™ 647, 1:1000 (Invitrogen™, A21244)) for 1 h at room temperature in the dark. Cells were washed three times every 10 mins with the washing solution and mounted with Fluoromount-G Mounting Medium with DAPI (Invitrogen, 00-4959-52). Cells were imaged using confocal microscopy (FV 1200, Olympus Fluoroview) using a 30x silicon objective (NA1.05). Images were processed by ImageJ[84,85] and quantified using CellProfiler (v.4.2.5.)[86]. Statistics were calculated in GraphPad Prism.

Real Time Quantitative Polymerase Chain Reaction (RT-qPCR): Total RNA from cells incubated at 32 or 37°C was isolated, using Direct-zol™ RNA Microprep (Zymo Research, R2062) according to the manufacturer's instructions. The concentration of RNA was measured by NanoDrop (Thermo Fisher Scientific) followed by cDNA synthesis with High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, 4368814) on MiniAmp™ Thermal Cycler (Applied Biosystems™). RT-qPCR was performed using Luna® Universal qPCR Master Mix (NEB, M3003) on CFX384™ Real-Time PCR Detection System

(Bio-Rad). Each biological replicate of the RT-qPCR assay in this study was carried out in technical
triplicates. Any technical replicate that deviated from other replicates by ≥ 0.4 cycle threshold (Ct)
was removed from calculations of average Ct values. The primers used are as follows: SP1 (fwd): 5'-
CACCCAATTCAAGGCCTGCCGT-3'; SP1 (rev): 5'-GGGTTGGGCATCTGGGCTGTTT-3'; RBM3 (fwd): 5'-
GAGACTCAGCGGTCCAGGGGTT-3'; RBM3 (rev): 5'-CCTCTGGTTCCCCGAGCAGACT-3'; CIRBP (fwd): 5'-
CCGAGTTGACCAGGCTGGCAAG-3'; CIRBP (rev): 5'-TCCATAGCCCCGGTCTCCTCCT-3'; GAPDH (fwd): 5'-
TCAAGGCTGAGAACGGGAAG-3'; GAPDH (rev): 5'-CGCCCCACTTGATTTTGGAG-3'. SMYD5 1 (fwd): 5'-
GCACTGTGCGCAAAGACCTCCA-3', SMYD5 2 (fwd): 5'-GGAAACCAGGCCAGGTTCTGCC-3', SMYD5 3
(fwd) 5'-CGTGGAAGTCCGTTTCGTGA-3', SMYD5 1 (rev:) 5'-CTGGGCACAGGACCTGGTGGTA-3', SMYD5
2 (rev): 5'-GGCTGCCAACCGACATTCTGCA-3', SMYD5 3 (rev) 5'-CCAGAGAAACTGTGCAGCCA-3'.

Western blot assay: Cells were washed with PBS and then lysed (whole cell lysate) for 30 min on ice
in RIPA buffer (50mM Tris HCl ph8, 150mM NaCl, 1% NP-40, 1% Sodium deoxycholate, 0.1% SDS,
2mM EDTA, phosphatase inhibitor (either Cell signaling, 5870S or Thermo Fisher Scientific, 78437)).
The lysate was centrifuged at 16,000 xg for 20 min at 4°C and the supernatant was collected. The
supernatant was diluted with 4X loading buffer (LI-COR, 928-40004) and heated at 95°C for 5 min.
The protein concentration was measured using Pierce™ BCA Assay Kit (Thermo Fisher Scientific,
23225). 20-30 μg of protein was loaded into each well and separated by SDS-PAGE. They were
transferred to polyvinylidene fluoride (PVDF) membranes and blocked in 5% bovine serum albumin in
Tris-buffered saline with Tween 20 for 1 h. Primary antibodies: rabbit anti-CIRBP, 1:2000
(Proteintech, 10209-2-AP); rabbit anti-RBM3, 1:1000 (Proteintech, 14363-1-AP); rabbit anti-SP1
1:1000 (Proteintech, 21962-1-AP); mouse anti-Lamin B1, 1:5000 (Proteintech 66095-1-Ig); rabbit anti-
SMYD5, 1:1000[51]; mouse anti-GAPDH, 1:5000 (Abcam, ab8245) were applied to respective
membranes after washing, and incubated at 4°C overnight. Membranes were incubated with IRDye®
secondary antibodies; anti-rabbit (LI-COR,926-32213); anti-mouse (LI-COR, 926-68072), at room
temperature for 90 mins. After washing, the protein bands were visualized on the Odyssey® CLx
Infrared Imaging System and protein band intensity quantified by Image J[84,85] software.

DNAm studies and analysis: HEK293WT (biological triplicates) were split into two batches and
exposed to 37°C and 32°C for 6 h, next cells were harvested, and DNA isolated with the Zymo Quick-
DNA™ Midiprep Plus Kit (Zymo Research, D4075). Samples were submitted to the Genetic Resources
Core Facility (Johns Hopkins University), bisulfite treated, and run on the Infinium MethylationEPIC
BeadChip Kit (Illumina) with two technical controls (50% and 100% DNAm, 8 samples total). Technical
controls were created using mixtures of 100% and 0% methylated HCT116 DKO (Zymo Research
D50414-1/2). DNA methylation data (IDAT files) containing unmethylated and methylated intensity
values were imported to R (v.4.2.1) for analysis[82]. Raw data was processed using single sample
approach with preprocessNoob() function and methylation values (beta values) were obtained using
the getBeta() function from minfi package (v.1.42.0)[87–89]. The model matrix was created for the two
temperatures using the model.matrix() function from stats R package and sva() function from sva
package (v.3.44.0) was applied with two estimated surrogate variables[82,90]. The model was fit with
contrast matrix using lmFit(), with the estimated coefficients and standard errors generated using
contrasts.fit() from limma package (v.3.52.4)[91]. The p-values were moderated using eBayes() function
from limma package (v.3.52.4) and CpG cites ranked according to their p-value[91].

CUT&RUN: 293WT and human NPCs (ATCC, ACS-1019) were harvested after 6 h at 32°C or kept at
37°C. CUT&RUN performed according to Epicypher CUTANA protocol (v.1.7)[92] on 300.000 cells per

sample. We permeabilized the cells with 0.01% digitonin (Sigma, D141). For normalization we spiked
in E. Coli DNA (Epicypher, 23618-1401) at the final concentration of 0.2 ng per sample. The library
preparation was performed with TrueSeq-Chip Sample preparation Kit (Illumina, IP-202-9001). The
following antibodies were used for the CUT&RUN: CTCF antibody (Cell Signaling, 2899S), H3K36me3
(Thermo Fisher Scientific, MA5-24687), H4K20me3 (EpiCypher, 13-0054), H3K9me3 (Abcam,
ab176916), H3K4me3 (EpiCypher, 13-0041), H3K27me3 (Thermo Fisher Scientific, MA5-11198),
SMYD5 [51], and Rabbit IgG Antibody (Epicypher, 23613-0042). The sequencing was performed on a
NovaSeq 6000 S4 at deCODE genetics, paired-end 8 cycles of index1, 8 cycles of index 2, 150 cycles
read 1 and, 150 cycles read 2, where we aimed for 10 million reads per sample. For the CUT&RUN
analysis and the CUT&TAG re-analysis we followed published STAR protocols[93]. Briefly, first we
trimmed the raw Fastq reads in paired mode with TrimGalore (v.0.6.7)[94] then we used Bowtie2
(v.2.4.4.)[95] to align the reads to Hg38 or mm10 and E. Coli reference genome, all downloaded from
UCSC. Overall alignment to Hg38 were from 83.6-98.63% and to E. Coli spike in were from 0.1-4.18%
for our data. The aligned Sam files were then converted to Bam files, indexed, and sorted via
SAMtools (v.1.15)[96] and BedGraphtoBigWig (v.4)[97] from BEDTools (v.2.3)[98]. We used SEACR (v.1.3)[99]
for peak calling after normalization with the normalization factor calculated as 1/(percent alignment
to E. coli). We used DiffBind (v.3.9.6)[100] for the differential peak analysis, ChIPseeker (v.1.35.3)[101],
TxDb.Hsapiens.UCSC.hg38.knownGene (v.3.4.6)[102] and org.Hs.eg.db (v.3.8.2)[103] for annotation. For
further analysis and image creation we used deepTools (v.3.5.1.)[104] and R [82].

Mouse NPC isolation: Mouse NPCs were isolated from the cortex of E17.5 C57BL6/NTac embryos.
The protocol was adapted from Bernas et al, 2017[105]. Embryo cortices were manually dissected from
brains and the tissue was dissociated in 1X TrypLE™ Select Enzyme (Thermo Fisher Scientific,
A1217701) for 10 min, with manual dissociation using a 1000 µL pipette. The cell suspension was
washed in Neurobasal medium (Thermo Fisher Scientific, 21103049) and filtered through a 70 µm
cell strainer (Miltenyi Biotech, 130-110-916). Cells were washed twice in 5 mL Neurobasal medium
(Gibco, 21103049) by centrifugation at 200 xg for 10 min. Cells were resuspended in 1 mL Neurobasal
growth medium containing 1X B27 supplement (Thermo Fisher Scientific, 17504044), 1X
Penicillin/Streptomycin (Thermo Fisher Scientific, 15140122), 1X Glutamax (Thermo Fisher Scientific,
35050038), 20ng/mL FGF-2 (Peprotech, 100-18B), 20 ng/mL EFG (Peprotech, AF-100-15), and 2
µg/mL Heparin (MP Biomedicals, 210193125). Cells were seeded onto 12-well plates, previously
coated for 2 h at 37°C in 1:100 dilution of Matrigel (Corning, 354234). After passaging, NPCs were
cultured in media described above on plates coated overnight in poly-D-lysine (Sigma, P7280), and
then coated for 2 h in laminin (Sigma, L2020). Cells were either sub-cultured or media was changed
every 2 days.

RNAseq: Primary mouse NPC lines from 4 separate embryos were seeded 0.3×10^6 cells per well, in
two batches, in a poly-D-lysine/laminin coated 6-well plate after 9 days *in vitro*. Following 36 h of
incubation at 37°C, one plate with 4 lines was incubated at 32°C for 6 h. Cells from both 37°C and
32°C conditions were harvested in tri-reagent and RNA was isolated using the Direct-zol RNA
Microprep kit (Zymo Research, R2063). RNAseq library construction and Illumina next generation
paired-end sequencing was performed by Novogene, generating 30 million paired-end reads per
sample. For the RNAseq analysis, Fastq files were pseudo-aligned to the GRCm39 mouse
transcriptome, downloaded from NCBI, using Kallisto (v.0.48.0) with paired-end mode on and 100
bootstraps. Kallisto output files were imported into R using the tximport package[106]. Transcripts were
assigned to genes using the TxDb.Mmusculus.UCSC.mm10.ensGene package (v.3.4.0) [107] in R .

Differential expression analysis was performed using the DESeq2 package (v.3.16)[108] in R, after
discarding transcripts with less than 10 counts. Differentially expressed genes were defined as those
with an adjusted p-value less than 0.1. Overlaps of gene lists were calculated using the GeneOverlap
package (v.1.34.0)[109] in R, with overlaps tested using Fisher's exact test. Gene set enrichment analysis
was performed using the GSEA-preranked program[110,111] with the Mouse Ortholog Hallmark Gene
Sets[112]. For the GSEA-pre-ranked run, we generated a ranked gene list comprising all detected genes
assigned to a rank value, which was calculated as -log10(adjusted p-value) × (sign of fold change).
The genes were then ranked from the highest rank value to the lowest before inputting into the
program.

Mouse cooling followed by immunostaining on brain slices: C57BL6 mice (Jackson) were bred to
create a litter (n=2). At P10, we split littermates into two groups. One group (n=6) was cooled to a
core temperature of 32°C and the other (n=6) maintained at 37°C, for 6 hours respectively. These
animals were then euthanized, and their brains were isolated immediately after the cooling period
and flashed with PBS. One of the hemispheres was immersed in 4% PFA for 72-96 h for fixation and
the other hemisphere was dissected for the cerebellum, thalamus/basal ganglia, hippocampus,
anterior and posterior cortex. After fixation, samples were cryopreserved in a sucrose gradient and
then flash-frozen. Frozen hemispheres were cryosectioned with the help of a cryostat (Keldur, UI).
The sections were washed in TBS+ (TBS (LiCor, 927-60001) with 0.05% triton X-100 (Sigma-Aldrich,
T8787-100ML)) twice for 10 minutes. Next, the slides were immersed in DAKO (target retrieval
solution, sodium citrate buffer, pH 6) at 95°C for 20 minutes. Slides were then washed again in TBS+
twice for 10 minutes. Next, the slides were blocked in TBS++ (TBS (LiCor, 927-60001) with 3% goat
serum (Abcam, ab7481)) for 1 h at room temperature. The slides were then stained with primary
antibodies and anti-rabbit-SMYD5[51], 1:200, for 24 h at room temperature and carried out in the dark.
On the next day, slides were allowed to reach room temperature before they are washed 3x for 10
minutes in TBS+. The slides were incubated with secondary antibodies, goat anti-mouse IgG (H+L)
Alexa Fluor™ 488, 1:1000 (Invitrogen™, A-11001); goat anti-rabbit IgG (H+L) Alexa Fluor™ 647, 1:1000
(Invitrogen™, A21244), for 24 h in the dark and washed 3x for 10 minutes in TBS+. All the liquid on
the slides was briefly dried off. Fluoromount-G Mounting Medium with DAPI (Invitrogen, 00-4959-52)
was placed on each section and covered with a cover slip. The slides were incubated for 24 h at room
temperature before being imaged using confocal microscopy (FV 1200, Olympus Fluoroview) using a
30x silicon objective (NA1.05). Images were processed by ImageJ[84,85] and quantified using CellProfiler
(v.4.2.5.)[86]. Statistics were calculated in GraphPad Prism.

Supplementary Tables, Files and Figure titles and legends

List of Supplementary Tables. All Word documents.

Supplementary Table 1: Specifics of our MHI, including sequence material used, vector, selection
markers, plasmid size and individual fluorophores.

Supplementary Table 2: Top 91 drug hits from MHI screen.

Supplementary Table 3: Top candidate genes found enriched in mutation screening.

Supplementary Table 4: Genes found in both repressor and activator SP1 lists and thus excluded for
further evaluation.

Supplementary Table 5: List of CpGs represented at three genes.

Supplementary Table 6: List of differentially methylated locations from EPIC DNAm array.

Supplementary Table 7: Top differentially modified locations for H3K36me3 in NPCs from CUT&RUN
experiment.

Supplementary Table 8: List of genes that have a unique H3K4me3 histone modification peak at
32°C.

Supplementary Table 9: List of genes that show DEGS at 32°C and have a unique H3K4me3 histone
modification peak at 32°C.

List of Supplementary Files. Word document.

Supplementary File 1: Sequences of individual MHIs.

Figure S1. Different temperature ranges. Temperatures known to induce distinct responses

(moderate and mild hypothermia, euthermia and hyperthermia). Our range of interest (mild

hypothermia) is bolded.

Figure S2. MHI activation peaks during mild hypothermia (32°C) and is limited at other temperatures. (A-C) We measured fluorescence hourly using an IncucyteS3 at 32°C using SP1-Lenti-MHI, CIRBP-Lenti-MHI, RBM3-Lenti-MHI respectively. (D-E) No increased fluorescence was observed at moderate hypothermia for either SP1-MHI or CIRBP-MHI in HEK293 cells. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S3. Storage of samples on ice does not lead to activation of MHI. (A-D) gMFI or percentage of fluorescent cells of HEK293WT+Cas9+SP1 left on ice for different periods of time before flow cytometry analysis. Shown are data points, each data point is a technical replicate, mean and SD where applicable. Dunnett's multiple comparison test calculated in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S4. MHI demonstrate consistent activation at 32°C in diverse cellular systems. (A-E) Left
column shows mean fluorescence (MF, flow cytometry) after transient transfection of SP1-MHI into
HCT116, HeLa, Jurkat, K562 and SK-N-SH (16 h at 32°C, 37°C and 40°C). Shown are data points, each
data point is a technical replicate, mean and SD where applicable, Šidák's multiple comparison test
calculated in GraphPad Prism. (F-K) Middle column shows gMFI (flow cytometry) after transient
transfection of RBM3-MHI into HCT116, HeLa, Jurkat, K562, SK-N-SH, and HEK293T, 16 h at 32°C.
Shown are data points, each data point is a technical replicate, mean and SD where applicable,
Tukey's multiple comparison test calculated in GraphPad Prism. (L-Q) Right column shows gMFI (flow
cytometry) after transient transfection of SP1+MCRE-MHI into HCT116, HeLa, Jurkat, K562, SK-N-SH,
and HEK293T, 16 h at 32°C. Data as in (F-K). * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S5. Moderate hyperthermia does not activate SP1 MHI in diverse systems. (A-E) Mean fluorescence of SP1-MHI as measured by flow cytometry after transient transfection of MHI into HCT116, HeLa, Jurkat, K562, and SK-N-SH, 16 h at 26°C or 29°C. Shown are data points, each data point is a technical replicate, mean and SD where applicable. Significance levels calculated with Šidák's multiple comparison test in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S6: Validation of potential compound hits from drug screen and cell cycle analysis. (A-F)
gMFI measured with flow cytometry of HEK293WT cell line that stably expresses SP1-MHI after 16 h
of Cinacalcet (A), Dasatinib (B), Erythromycin (C), Naloxone (D), Orantinib (E), and Tamoxifen (F) at
32°C or 37°C. Shown are data points, mean and SD where applicable. Significance levels calculated
with Šidák's multiple comparison calculated with GraphPad Prism. (G) Cell cycle analysis done with PI
staining. Mean fluorescence measured for G1, S and G2 phases of the cell cycle with flow cytometry
of HEK293WT cell line 16 h after Entacapone or Poziotinib exposure at 32°C or 37°C. Data shown as in
(A-F). (H) Apoptosis induction analysis done with H_2O_2 exposure. Shown are data points, each point a
technical replicate, mean and SD where applicable. Significance levels calculated with an unpaired
two-tailed t-test calculated in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** =
P<0.0001.

Figure S7: Expression of endogenous loci after exposure to Entacapone and Poziotinib. Relative expression of *SP1* (top), *CIRBP* (middle), *RBM3* (bottom) when compared to GAPDH after 16 h at 32°C or 37°C with and without Entacapone (right) or Poziotinib (left). Shown are data points, each data point is a biological replicate (mean of technical replicates), mean and SD where applicable. Significance levels were calculated with an unpaired one-tailed t-test in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S8: *RGS21* and *MALT1* knockout did not show any significant fluorescent change. sgRNA KO

for *MALT1* and *RGS21* did not show any changes in fluorescence of HEK293 cells with SP1-MHI,

suggesting that *MALT1* and *RGS21* do not have an impact on SP1-MHI. Shown are data points, each

data point is a technical replicate, mean and SD where applicable. Significance levels were calculated

with Šidák's multiple comparison test in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001,

**** = P<0.0001.

Figure S9. Pathway analysis using genes from forward mutagenesis. (A) Schematic depiction of
repressors identified which land on the Neuregulin pathway. (B-C) Unbiased MSigDB pathway
enrichment of highly ranked (-Log10(RRA) >0.5) genes from the SP1 activator (B) and SP1 repressor
(C) lists.

Figure S10. *Cirbp* has fewer normalized counts compared to *Sp1* and *Rbm3.* Normalized counts from
RNAseq in mESc (referring to data from **Figure 4C**).

Figure S11. Validation of temperature dependence of SMYD5. (A) An independent biological replicate with anti-SMYD5 antibody showing striking differences between two temperatures. Green or white (SMYD5), purple (DAPI). (B) A validation with anti-FLAG antibody, showing the same difference at the two temperatures. Green (SMYD5), purple (DAPI).

Figure S12. DNAm status at *SP1*, *CIRBP* and *RBM3* promoters. (A) DNAm status of CpG dinucleotides
from Illumina EPIC array 5 kb upstream and 2 kb downstream of *SP1* gene. (B) DNAm status of CpG
dinucleotides from Illumina EPIC array 5 kb upstream and 2 kb downstream of *CIRBP* gene. (C) DNAm
status of CpG dinucleotides from Illumina EPIC array 5 kb upstream and 2 kb downstream of *RBM3*
gene.

Figure S13. Methylation modification levels in NPCs and HEK293 cells. (A) H3K36me3 levels at 32°C and 37°C at *SP1* promoter in HEK293 cells. (B) Global H3K36me3 levels (all genes) in HEK293 cells. (C-D) Global H3K27me3, H4K20me3 levels, respectively (all genes) in NPCs (E) At the gene-specific levels there is a number of genes, which have more or less H3K36me3 at 32°C compared to 37°C in HEK293 cells. Red circles indicate genes that have Log2 fold change over 1 and p-value less than 10-6, gray circles indicate Log2 fold change over 1 but non-significant p-value (>10-6) and black circles indicate non-significant Log2 fold change and p-values for those genes. (F-G) H3K36me3 levels at SMYD5-bound genes identified to be DEGS at the two temperatures in NPC and HEK293 cells, respectively. (H-J) H3K9me3 and H3K4me3, respectively, levels at SMYD5-bound genes identified to be DEGS at the two temperatures in NPCs.

Figure S14: H3K36me3 levels on representative peaks showing differential binding at 32°C when compared to 37°C. (A-C) Increased H3K36me3 is observed for *BAZ2A, RIMOC1 and USP1,* respectively, at 32°C compared to 37°C in NPCs. (D-E) Decreased H3K36me3 is observed for *CENPE, SETD1B* and *YEATS2I,* respectively, at 32°C compared to 37°C in NPCs.

References

1.　Osilla, E. V., Marsidi, J.L., and Sharma, S. (2022). Physiology, Temperature Regulation
(StatPearls Publishing).

2.　Turbill, C., Bieber, C., and Ruf, T. (2011). Hibernation is associated with increased survival and
the evolution of slow life histories among mammals. Proceedings of the Royal Society B:
Biological Sciences *278*, 3355–3363. 10.1098/RSPB.2011.0190.

3.　Mathew, J.L., Kaur, N., and Dsouza, J.M. (2022). Therapeutic hypothermia in neonatal hypoxic
encephalopathy: A systematic review and meta-analysis. J Glob Health *12*.
10.7189/JOGH.12.04030.

4.　Luedke, M.W., Graffagnino, C., McKinney, B.G., Piper, J., Iversen, E., and Kolls, B. (2022).
Association of time-temperature curves with outcomes in temperature management for
cardiac arrest. BMJ Neurol Open *4*. 10.1136/BMJNO-2022-000273.

5.　Dietrichs, E.S., and Dietrichs, E. (2015). Neuroprotective effects of hypothermia. Tidsskr Nor
Laegeforen *135*, 1646–1651. 10.4045/tidsskr.14.1250.

6.　Sun, Y.J., Zhang, Z.Y., Fan, B., and Li, G.Y. (2019). Neuroprotection by Therapeutic
Hypothermia. Front Neurosci *13*. 10.3389/FNINS.2019.00586.

7.　Fujita, J. (1999). Cold shock response in mammalian cells. J Mol Microbiol Biotechnol *1*, 243–
255.

8.　Zhu, X., Bührer, C., and Wellmann, S. (2016). Cold-inducible proteins CIRP and RBM3, a unique
couple with activities far beyond the cold. Cellular and Molecular Life Sciences, 1–21.
10.1007/s00018-016-2253-7.

9.　Ryu, H., Lee, J., Olofsson, B.A., Mwidau, A., Deodoglu, A., Escudero, M., Flemington, E.,
Azizkhan-Clifford, J., Ferrante, R.J., and Ratan, R.R. (2003). Histone deacetylase inhibitors
prevent oxidative neuronal death independent of expanded polyglutamine repeats via an Sp1-
dependent pathway. Proc Natl Acad Sci U S A *100*, 4281–4286. 10.1073/PNAS.0737363100.

10.　Yang, H.J., Zhuang, R.J., Li, Y.B., Li, T., Yuan, X., Lei, B.B., Xie, Y.F., and Wang, M. (2019). Cold-
inducible protein RBM3 mediates hypothermic neuroprotection against neurotoxin rotenone
via inhibition on MAPK signalling. J Cell Mol Med *23*, 7010–7020. 10.1111/JCMM.14588.

11.　Bastide, A., Peretti, D., Knight, J.R.P., Grosso, S., Spriggs, R. V., Pichon, X., Sbarrato, T., Roobol,
A., Roobol, J., Vito, D., et al. (2017). RTN3 Is a Novel Cold-Induced Protein and Mediates
Neuroprotective Effects of RBM3. Current Biology *27*, 638. 10.1016/J.CUB.2017.01.047.

12.　Chip, S., Zelmer, A., Ogunshola, O.O., Felderhoff-Mueser, U., Nitsch, C., Bührer, C., and
Wellmann, S. (2011). The RNA-binding protein RBM3 is involved in hypothermia induced
neuroprotection. Neurobiol Dis *43*, 388–396. 10.1016/j.nbd.2011.04.010.

13.　Peretti, D., Bastide, A., Radford, H., Verity, N., Molloy, C., Martin, M.G., Moreno, J.A., Steinert,
J.R., Smith, T., Dinsdale, D., et al. (2015). RBM3 mediates structural plasticity and protective

effects of cooling in neurodegeneration. Nature 2015 518:7538 *518*, 236–239.
10.1038/nature14142.

14. Wierstra, I. (2008). Sp1: Emerging roles—Beyond constitutive activation of TATA-less
housekeeping genes. Biochem Biophys Res Commun *372*, 1–13. 10.1016/J.BBRC.2008.03.074.

15. Sumitomo, Y., Higashitsuji, H., Higashitsuji, H., Liu, Y., Fujita, T., Sakurai, T., Candeias, M.M.,
Itoh, K., Chiba, T., and Fujita, J. (2012). Identification of a novel enhancer that binds Sp1 and
contributes to induction of cold-inducible RNA-binding protein (cirp) expression in
mammalian cells. BMC Biotechnol *12*, 72. 10.1186/1472-6750-12-72.

16. Gomez-Pastor, R., Burchfiel, E.T., and Thiele, D.J. (2018). Regulation of heat shock
transcription factors and their roles in physiology and disease. Nat Rev Mol Cell Biol *19*, 4–19.
10.1038/NRM.2017.73.

17. Sanjana, N.E., Shalem, O., and Zhang, F. Improved vectors and genome - wide libraries for
CRISPR screening.

18. Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelson, T., Heckl, D., Ebert, B.L.,
Root, D.E., Doench, J.G., et al. (2014). Genome-scale CRISPR-Cas9 knockout screening in
human cells. Science *343*, 84–87. 10.1126/science.1247005.

19. Joung, J., Konermann, S., Gootenberg, J.S., Abudayyeh, O.O., Platt, R.J., Brigham, M.D.,
Sanjana, N.E., and Zhang, F. (2017). Genome-scale CRISPR-Cas9 knockout and transcriptional
activation screening. Nat Protoc *12*, 828–863. 10.1038/nprot.2017.016.

20. McIntyre, S., Nelson, K.B., Mulkey, S.B., Lechpammer, M., Molloy, E., and Badawi, N. (2021).
Neonatal encephalopathy: Focus on epidemiology and underexplored aspects of etiology.
Semin Fetal Neonatal Med *26*. 10.1016/J.SINY.2021.101265.

21. Geneva, I.I., Cuzzo, B., Fazili, T., and Javaid, W. (2019). Normal Body Temperature: A
Systematic Review. Open Forum Infect Dis *6*. 10.1093/OFID/OFZ032.

22. Knight, J.R.P., Bastide, A., Roobol, A., Roobol, J., Jackson, T.J., Utami, W., Barrett, D.A., Smales,
C.M., and Willis, A.E. (2015). Eukaryotic elongation factor 2 kinase regulates the cold stress
response by slowing translation elongation. Biochem J *465*, 227–238. 10.1042/BJ20141014.

23. Shaw, G., Morse, S., Ararat, M., and Graham, F.L. (2002). Preferential transformation of
human neuronal cells by human adenoviruses and the origin of HEK 293 cells. The FASEB
Journal. 10.1096/fj.01-0995fje.

24. Barnes, E.N., Biedler, J.L., Spengler, B.A., and Lyser, K.M. (1981). The fine structure of
continuous human neuroblastoma lines SK-N-SH, SK-N-BE(2), and SK-N-MC. In Vitro *17*, 619–
631. 10.1007/BF02618461.

25. Stepanenko, A.A., and Dmitrenko, V. V (2015). HEK293 in cell biology and cancer research:
phenotype, karyotype, tumorigenicity, and stress-induced genome-phenotype evolution.
Gene *569*, 182–190. 10.1016/j.gene.2015.05.065.

26. Jackson, T.C., Manole, M.D., Kotermanski, S.E., Jackson, E.K., Clark, R.S.B., and Kochanek, P.M.
(2015). Cold stress protein RBM3 responds to temperature change in an ultra-sensitive
manner in young neurons. Neuroscience *305*, 268–278. 10.1016/j.neuroscience.2015.08.012.

27. Hartmann, J., Haap, M., Kopp, H.-G., and Lipp, H.-P. (2009). Tyrosine kinase inhibitors - a
review on pharmacology, metabolism and side effects. Curr Drug Metab *10*, 470–481.
10.2174/138920009788897975.

28. Nelson, V., Ziehr, J., Agulnik, M., and Johnson, M. (2013). Afatinib: emerging next-generation
tyrosine kinase inhibitor for NSCLC. Onco Targets Ther *6*, 135–143. 10.2147/OTT.S23165.

29. Kim, T.M., Lee, K.W., Oh, D.Y., Lee, J.S., Im, S.A., Kim, D.W., Han, S.W., Kim, Y.J., Kim, T.Y., Kim,
J.H., et al. (2018). Phase 1 Studies of Poziotinib, an Irreversible Pan-HER Tyrosine Kinase
Inhibitor in Patients with Advanced Solid Tumors. Cancer Res Treat *50*, 835–842.
10.4143/CRT.2017.303.

30. Erber, R., Thurnher, A., Katsen, A.D., Groth, G., Kerger, H., Hammes, H.P., Menger, M.D.,
Ullrich, A., and Vajkoczy, P. (2004). Combined inhibition of VEGF and PDGF signaling enforces
tumor vessel regression by interfering with pericyte-mediated endothelial cell survival
mechanisms. FASEB J *18*, 338–340. 10.1096/FJ.03-0271FJE.

31. Lee, H., Kim, J.W., Choi, D.K., Yu, J.H., Kim, J.H., Lee, D.S., and Min, S.H. (2020). Poziotinib
suppresses ovarian cancer stem cell growth via inhibition of HER4-mediated STAT5 pathway.
Biochem Biophys Res Commun *526*, 158–164. 10.1016/J.BBRC.2020.03.046.

32. Grinstein, E., Jundt, F., Weinert, I., Wernet, P., and Royer, H.D. (2002). Sp1 as G1 cell cycle
phase specific transcription factor in epithelial cells. Oncogene 2002 21:10 *21*, 1485–1492.
10.1038/sj.onc.1205211.

33. Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S., Brown, M., and Liu,
X.S. (2014). MAGeCK enables robust identification of essential genes from genome-scale
CRISPR/Cas9 knockout screens. Genome Biol *15*, 554. 10.1186/s13059-014-0554-4.

34. Tan, N.Y., and Khachigian, L.M. (2009). Sp1 phosphorylation and its regulation of gene
transcription. Mol Cell Biol *29*, 2483–2488. 10.1128/mcb.01828-08.

35. Nagai, G., and Nakaoka, Y. (1998). Cooling sensitive [Ca2+](i) response associated with
signaling of G protein-coupled receptors. Biochem Biophys Res Commun *248*, 733–737.
10.1006/BBRC.1998.8873.

36. Chelban, V., Patel, N., Vandrovcova, J., Zanetti, M.N., Lynch, D.S., Ryten, M., Botía, J.A., Bello,
O., Tribollet, E., Efthymiou, S., et al. (2017). Mutations in NKX6-2 Cause Progressive Spastic
Ataxia and Hypomyelination. Am J Hum Genet *100*, 969–977. 10.1016/J.AJHG.2017.05.009.

37. ZNF260 zinc finger protein 260 [Homo sapiens (human)] - Gene - NCBI
https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=339324#gene-
expression.

38. Burke, T.L., and Grant, P.A. (2009). Histone Acetylation Complexes. Handbook of Cell
Signaling, Second Edition *3*, 2369–2378. 10.1016/B978-0-12-374145-5.00285-0.

39. Zhou, X., Shang, Y.N., Lu, R., Fan, C.W., and Mo, X.M. (2019). High ANKZF1 expression is
associated with poor overall survival and recurrence-free survival in colon cancer.
https://doi.org/10.2217/fon-2018-0920 *15*, 2093–2106. 10.2217/FON-2018-0920.

40. Tawamie, H., Martianov, I., Wohlfahrt, N., Buchert, R., Mengus, G., Uebe, S., Janiri, L., Hirsch,
F.W., Schumacher, J., Ferrazzi, F., et al. (2017). Hypomorphic Pathogenic Variants in TAF13 Are
Associated with Autosomal-Recessive Intellectual Disability and Microcephaly. Am J Hum
Genet *100*, 555. 10.1016/J.AJHG.2017.01.032.

41. ZNF71 zinc finger protein 71 [Homo sapiens (human)] - Gene - NCBI
https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=58491.

42. Yu, H. V., Tao, L., Llamas, J., Wang, X., Nguyen, J.D., Trecek, T., and Segil, N. (2021). POU4F3
pioneer activity enables ATOH1 to drive diverse mechanoreceptor differentiation through a
feed-forward epigenetic mechanism. Proc Natl Acad Sci U S A *118*.
10.1073/PNAS.2105137118/-/DCSUPPLEMENTAL.

43. GATAD2A GATA zinc finger domain containing 2A [Homo sapiens (human)] - Gene - NCBI
https://www.ncbi.nlm.nih.gov/gene/54815.

44. Zhang, Y., Fang, Y., Tang, Y., Han, S., Jia, J., Wan, X., Chen, J., Yuan, Y., Zhao, B., and Fang, D.
(2022). SMYD5 catalyzes histone H3 lysine 36 trimethylation at promoters. Nature
Communications 2022 13:1 *13*, 1–19. 10.1038/s41467-022-30940-1.

45. Kidder, B.L., He, R., Wangsa, D., Padilla-Nash, H.M., Bernardo, M.M., Sheng, S., Ried, T., and
Zhao, K. (2017). SMYD5 Controls Heterochromatin and Chromosome Integrity during
Embryonic Stem Cell Differentiation. Cancer Res *77*, 6729–6745. 10.1158/0008-5472.CAN-17-
0828.

46. Murphy, M.W., Sarver, A.L., Rice, D., Hatzi, K., Ye, K., Melnick, A., Heckert, L.L., Zarkower, D.,
and Bardwell, V.J. (2010). Genome-wide analysis of DNA binding and transcriptional
regulation by the mammalian Doublesex homolog DMRT1 in the juvenile testis. Proc Natl
Acad Sci U S A *107*, 13360–13365. 10.1073/PNAS.1006243107/SUPPL_FILE/SD01.XLS.

47. Weber, C., Zhou, Y., Lee, J.G., Looger, L.L., Qian, G., Ge, C., and Capel, B. (2020). Temperature-
dependent sex determination is mediated by pSTAT3 repression of Kdm6b. Science (1979)
368, 303–306. 10.1126/SCIENCE.AAZ4165/SUPPL_FILE/AAZ4165_S1.MP4.

48. Kim, D.H., Doyle, M.R., Sung, S., and Amasino, R.M. (2009). Vernalization: winter and the
timing of flowering in plants. Annu Rev Cell Dev Biol *25*, 277–299.
10.1146/annurev.cellbio.042308.113411.

49. Fellous, A., Earley, R.L., and Silvestre, F. (2019). The Kdm/Kmt gene families in the self-
fertilizing mangrove rivulus fish, Kryptolebias marmoratus, suggest involvement of histone
methylation machinery in development and reproduction. Gene *687*, 173–187.
10.1016/J.GENE.2018.11.046.

50.   Youn, J.Y., Dyakov, B.J.A., Zhang, J., Knight, J.D.R., Vernon, R.M., Forman-Kay, J.D., and
Gingras, A.C. (2019). Properties of Stress Granule and P-Body Proteomes. Mol Cell *76*, 286–
294. 10.1016/J.MOLCEL.2019.09.014.

51.   Aljazi, M.B., Gao, Y., Wu, Y., and He, J. (2022). SMYD5 is a histone H3-specific
methyltransferase mediating mono-methylation of histone H3 lysine 36 and 37. Biochem
Biophys Res Commun *599*, 142–147. 10.1016/J.BBRC.2022.02.043.

52.   Diaz, J., Abiola, S., Kim, N., Avaritt, O., Flock, D., Yu, J., Northington, F.J., and Chavez-Valdez, R.
(2017). Therapeutic hypothermia provides variable protection against behavioral deficits after
neonatal HI: A potential role for BDNF. Dev Neurosci *39*, 257. 10.1159/000454949.

53.   Xi, Y., Park, S.R., Kim, D.H., Kim, E.D., and Sung, S. (2020). Transcriptome and epigenome
analyses of vernalization in Arabidopsis thaliana. Plant J *103*, 1490. 10.1111/TPJ.14817.

54.   Fan, L., Lesser, A.F., Sweet, D.R., Keerthy, K.S., Lu, Y., Chan, E.R., Vinayachandran, V., Ilkayeva,
O., Das, T., Newgard, C.B., et al. (2022). KLF15 controls brown adipose tissue transcriptional
flexibility and metabolism in response to various energetic demands. iScience *25*, 105292.
10.1016/J.ISCI.2022.105292.

55.   Katano-Toki, A., Satoh, T., Tomaru, T., Yoshino, S., Ishizuka, T., Ishii, S., Ozawa, A., Shibusawa,
N., Tsuchiya, T., Saito, T., et al. (2013). THRAP3 Interacts with HELZ2 and Plays a Novel Role in
Adipocyte Differentiation. Molecular Endocrinology *27*, 769. 10.1210/ME.2012-1332.

56.   THRAP3 thyroid hormone receptor associated protein 3 [Homo sapiens (human)] - Gene -
NCBI https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=9967.

57.   Guo, K., Searfoss, G., Krolikowski, D., Pagnoni, M., Franks, C., Clark, K., Yu, K.T., Jaye, M., and
Ivashchenko, Y. (2001). Hypoxia induces the expression of the pro-apoptotic gene BNIP3. Cell
Death Differ *8*, 367–376. 10.1038/SJ.CDD.4400810.

58.   Abdel-Wahab, O., Adli, M., LaFave, L.M., Gao, J., Hricik, T., Shih, A.H., Pandey, S., Patel, J.P.,
Chung, Y.R., Koche, R., et al. (2012). ASXL1 mutations promote myeloid transformation
through loss of PRC2-mediated gene repression. Cancer Cell *22*, 180–193.
10.1016/J.CCR.2012.06.032.

59.   Kaakkola, S. (2000). Clinical pharmacology, therapeutic use and potential of COMT inhibitors
in Parkinson's disease. Drugs *59*, 1233–1250. 10.2165/00003495-200059060-00004.

60.   Feng, Z., Saha, L., Dritsa, C., Wan, Q., and Glebov, O.O. (2022). Temperature-dependent
structural plasticity of hippocampal synapses. Front Cell Neurosci *16*, 520.
10.3389/FNCEL.2022.1009970/BIBTEX.

61.   Carvelli, L., Blakely, R.D., and DeFelice, L.J. (2008). Dopamine transporter/syntaxin 1A
interactions regulate transporter channel activity and dopaminergic synaptic transmission.
Proc Natl Acad Sci U S A *105*, 14192–14197. 10.1073/PNAS.0802214105.

62.   Gao, H., Chakraborty, G., Lee-Lim, A.P., Mavrakis, K.J., Wendel, H.G., and Giancotti, F.G.
(2014). Forward genetic screens in mice uncover mediators and suppressors of metastatic

reactivation. Proc Natl Acad Sci U S A *111*, 16532–16537.
10.1073/PNAS.1403234111/SUPPL_FILE/PNAS.201403234SI.PDF.

63.   INDACOCHEA, A., GUERRERO, S., UREÑA, M., ARAUJO, F., COLL, O., LLEONART, M.E., and
1141   GEBAUER, F. (2021). Cold-inducible RNA binding protein promotes breast cancer cell
1142   malignancy by regulating Cystatin C levels. RNA *27*, 190–201. 10.1261/RNA.076422.120.

1143   64.   Nicolas, M., Noe, V., Jensen, K., and Ciudad CJ (2001). Cloning and characterization of the 5'-
1144   flanking region of the human transcription factor Sp1 gene. . J Biol Chem, 22126-22132.

1145   65.   Yang, F., Eckardt, S., Leu, N.A., McLaughlin, K.J., and Wang, P.J. (2008). Mouse TEX15 is
1146   essential for DNA double-strand break repair and chromosomal synapsis during male meiosis.
1147   J Cell Biol *180*, 673. 10.1083/JCB.200709057.

1148   66.   TEX15 testis expressed 15, meiosis and synapsis associated [Homo sapiens (human)] - Gene -
1149   NCBI https://www.ncbi.nlm.nih.gov/gene/56154.

1150   67.   Hanna, C.W., Huang, J., Belton, C., Reinhardt, S., Dahl, A., Andrews, S., Francis Stewart, A.,
1151   Kranz, A., and Kelsey, G. (2022). Loss of histone methyltransferase SETD1B in oogenesis
1152   results in the redistribution of genomic histone 3 lysine 4 trimethylation. Nucleic Acids Res *50*,
1153   1993. 10.1093/NAR/GKAC051.

1154   68.   Pajoro, A., Severing, E., Angenent, G.C., and Immink, R.G.H. (2017). Histone H3 lysine 36
1155   methylation affects temperature-induced alternative splicing and flowering in plants. Genome
1156   Biol *18*, 1–12. 10.1186/S13059-017-1235-X/FIGURES/4.

1157   69.   Michaels, K.K., Mostafa, S.M., Capella, J.R., and Moore, C.L. (2020). Regulation of alternative
1158   polyadenylation in the yeast Saccharomyces cerevisiae by histone H3K4 and H3K36
1159   methyltransferases. Nucleic Acids Res *48*, 5407–5425. 10.1093/NAR/GKAA292.

1160   70.   Xu, S.J., Lombroso, S.I., Fischer, D.K., Carpenter, M.D., Marchione, D.M., Hamilton, P.J., Lim,
1161   C.J., Neve, R.L., Garcia, B.A., Wimmer, M.E., et al. (2021). Chromatin-mediated alternative
1162   splicing regulates cocaine-reward behavior. Neuron *109*, 2943-2966.e8.
1163   10.1016/J.NEURON.2021.08.008.

1164   71.   Preußner, M., Goldammer, G., Neumann, A., Haltenhof, T., Rautenstrauch, P., Müller-
1165   McNicoll, M., and Heyd, F. (2017). Body Temperature Cycles Control Rhythmic Alternative
1166   Splicing in Mammals. Mol Cell *67*, 433-446.e4. 10.1016/J.MOLCEL.2017.06.006.

1167   72.   Horii, Y., Shiina, T., Uehara, S., Nomura, K., Shimaoka, H., Horii, K., and Shimizu, Y. (2019).
1168   Hypothermia induces changes in the alternative splicing pattern of cold-inducible RNA-binding
1169   protein transcripts in a non-hibernator, the mouse. Biomed Res *40*, 153–161.
1170   10.2220/BIOMEDRES.40.153.

1171   73.   Sano, Y., Shiina, T., Naitou, K., Nakamori, H., and Shimizu, Y. (2015). Hibernation-specific
1172   alternative splicing of the mRNA encoding cold-inducible RNA-binding protein in the hearts of
1173   hamsters. Biochem Biophys Res Commun *462*, 322–325. 10.1016/j.bbrc.2015.04.135.

74. Smart, F., Aschrafi, A., Atkins, A., Owens, G.C., Pilotte, J., Cunningham, B.A., and Vanderklish,
P.W. (2007). Two isoforms of the cold-inducible mRNA-binding protein RBM3 localize to
dendrites and promote translation. J Neurochem *101*, 1367–1379. 10.1111/J.1471-
4159.2007.04521.X.

75. Al-Fageeh, M.B., and Smales, C.M. (2013). Alternative promoters regulate cold inducible RNA-
binding (CIRP) gene expression and enhance transgene expression in mammalian cells. Mol
Biotechnol *54*, 238–249. 10.1007/s12033-013-9649-5.

76. Nicolas, M., Noe, V., Jensen, K.B., and Ciudad, C.J. (2001). Cloning and characterization of the
5'-flanking region of the human transcription factor Sp1 gene. J Biol Chem *276*, 22126–22132.
10.1074/jbc.M010740200.

77. Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., Studer, L., and
Author, N.B. (2009). Highly efficient neural conversion of human ES and iPS cells by dual
inhibition of SMAD signaling HHS Public Access Author manuscript. Nat Biotechnol *27*, 275–
280. 10.1038/nbt.1529.

78. Xiang, J., Wan, C., Guo, R., and Guo, D. (2016). Is Hydrogen Peroxide a Suitable Apoptosis
Inducer for All Cell Types? Biomed Res Int *2016*. 10.1155/2016/7343965.

79. Cell cycle staining with Propidium Iodide for the cytoplasmic GFP transfected cells using
Paraformaldehyde and Ethanol (2010). 1–2. https://flowcyt.rutgers.edu/wp-
content/uploads/2017/10/Cell-cycle-staining-with-PI-for-GFP-transfected-cells-using-
PFA_EtOH_2.pdf.

80. Kutner, R.H., Zhang, X.Y., and Reiser, J. (2009). Production, concentration and titration of
pseudotyped HIV-1-based lentiviral vectors. Nat Protoc *4*, 495–505. 10.1038/nprot.2009.22.

81. Wang, B., Wang, M., Zhang, W., Xiao, T., Chen, C.H., Wu, A., Wu, F., Traugh, N., Wang, X., Li,
Z., et al. (2019). Integrative analysis of pooled CRISPR genetic screens using MAGeCKFlute. Nat
Protoc *14*, 756–780. 10.1038/s41596-018-0113-7.

82. Statistical R Core Team (2017). R Foundation for Statistical Computing Vienna Austria. R: A
language and environment for computing.

83. 4×Lentivirus Concentrator Solution https://www.mdanderson.org/documents/core-
facilities/Functional%20Genomics%20Core/Homemade%204fold%20lentivirus%20concentrat
or.pdf.

84. Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and Eliceiri,
K.W. (2017). ImageJ2: ImageJ for the next generation of scientific image data. BMC
Bioinformatics *18*, 1–26. 10.1186/S12859-017-1934-Z/FIGURES/7.

85. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nature Methods 2012 9:7 *9*, 676–682. 10.1038/nmeth.2019.

86.   Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and Goodman, A.
1211   (2021). CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics *22*.
1212   10.1186/S12859-021-04344-9.

1213   87.   Fortin, J.P., Triche, T.J., and Hansen, K.D. (2017). Preprocessing, normalization and integration
1214   of the Illumina HumanMethylationEPIC array with minfi. Bioinformatics *33*, 558.
1215   10.1093/BIOINFORMATICS/BTW691.

1216   88.   Triche, T.J., Weisenberger, D.J., Van Den Berg, D., Laird, P.W., and Siegmund, K.D. (2013). Low-
1217   level processing of Illumina Infinium DNA Methylation BeadArrays. Nucleic Acids Res *41*, e90.
1218   10.1093/NAR/GKT090.

1219   89.   Aryee, M.J., Jaffe, A.E., Corrada-Bravo, H., Ladd-Acosta, C., Feinberg, A.P., Hansen, K.D., and
1220   Irizarry, R.A. (2014). Minfi: a flexible and comprehensive Bioconductor package for the
1221   analysis of Infinium DNA methylation microarrays. Bioinformatics *30*, 1363.
1222   10.1093/BIOINFORMATICS/BTU049.

1223   90.   Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E., and Storey, J.D. (2012). The sva package for
1224   removing batch effects and other unwanted variation in high-throughput experiments.
1225   Bioinformatics *28*, 882. 10.1093/BIOINFORMATICS/BTS034.

1226   91.   Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
1227   powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
1228   Acids Res *43*, e47. 10.1093/NAR/GKV007.

1229   92.   CUT&RUN Protocol v1.7 (2021).
1230   https://www.epicypher.com/content/documents/protocols/cutana-cut&run-
1231   protocol_1.7.pdf.

1232   93.   Kong, N.R., Chai, L., Tenen, D.G., and Bassal, M.A. (2021). A modified CUT&RUN protocol and
1233   analysis pipeline to identify transcription factor binding sites in human cell lines. STAR Protoc
1234   *2*. 10.1016/J.XPRO.2021.100750.

1235   94.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
1236   reads. EMBnet J *17*, 10–12. 10.14806/EJ.17.1.200.

1237   95.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
1238   Methods *9*, 357–359. 10.1038/nmeth.1923.

1239   96.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and
1240   Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*,
1241   2078–2079. 10.1093/BIOINFORMATICS/BTP352.

1242   97.   Luo, Y., Hitz, B.C., Gabdank, I., Hilton, J.A., Kagda, M.S., Lam, B., Myers, Z., Sud, P., Jou, J., Lin,
1243   K., et al. (2020). New developments on the Encyclopedia of DNA Elements (ENCODE) data
1244   portal. Nucleic Acids Res *48*, D882–D889. 10.1093/NAR/GKZ1062.

1245   98.   Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
1246   genomic features. Bioinformatics *26*, 841–842. 10.1093/BIOINFORMATICS/BTQ033.

99.   Meers, M.P., Tenenbaum, D., and Henikoff, S. (2019). Peak calling by Sparse Enrichment
Analysis for CUT&RUN chromatin profiling. Epigenetics Chromatin *12*, 1–11. 10.1186/S13072-
019-0287-4/FIGURES/6.

100. Stark, R., and Brown, G. (2012). DiffBind : differential binding analysis of ChIP-Seq peak data.

101. Yu, G., Wang, L.G., and He, Q.Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. Bioinformatics *31*, 2382–2383.
10.1093/BIOINFORMATICS/BTV145.

102. Team BC, and Maintainer BP (2019). TxDb.Hsapiens.UCSC.hg38.knownGene: Annotation
package for TxDb object(s).

103. Carlson M (2019). org.Hs.eg.db: Genome wide annotation for Human. .

104. Ramírez, F., Dündar, F., Diehl, S., Grüning, B.A., and Manke, T. (2014). deepTools: a flexible
platform for exploring deep-sequencing data. Nucleic Acids Res *42*, W187.
10.1093/NAR/GKU365.

105. Bernas, S.N., Leiter, O., Walker, T.L., and Kempermann, G. (2017). Isolation, Culture and
Differentiation of Adult Hippocampal Precursor Cells. Bio Protoc *7*.
10.21769/BIOPROTOC.2603.

106. Soneson, C., Love, M.I., and Robinson, M.D. (2016). Differential analyses for RNA-seq:
Transcript-level estimates improve gene-level inferences. F1000Res *4*.
10.12688/F1000RESEARCH.7563.2/DOI.

107. Team BC, and Maintainer BP (2016). TxDb.Mmusculus.UCSC.mm10.ensGene: Annotation
package for TxDb object(s)..

108. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 1–21. 10.1186/S13059-014-0550-
8/FIGURES/9.

109. Shen L, and Sinai ISoMaM (2022). GeneOverlap: Test and visualize gene overlaps. .

110. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment
analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc
Natl Acad Sci U S A *102*, 15545–15550. 10.1073/PNAS.0506580102.

111. Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver,
P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1alpha-responsive genes
involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat
Genet *34*, 267–273. 10.1038/NG1180.

112. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015).
The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst *1*, 417–
425. 10.1016/J.CELS.2015.12.004.