SMYD5 is a novel epigenetic gatekeeper of the mild hypothermia response

Salvor Rafnsdottir[1], Kijin Jang[1], Sara Tholl Halldorsdottir[1], Arnhildur Tomasdottir[1], Meghna Vinod[1],

Katrin Möller[1], Tinna Reynisdottir[1], Laufey Halla Atladottir[1], Kristin Elisabet Allison[2], Jin He[3], Li

Zhang[4], Frances J. Northington[5], Raul Chavez-Valdez[5], Kimberley Jade Anderson[1], Hans Tomas

Bjornsson[1,4,6,7*]

[1]Louma G. Laboratory of Epigenetic Research, Faculty of Medicine, University of Iceland, Reykjavik,

Iceland; [2]Faculty of Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland;

[3]Department of Biochemistry and Molecular Biology, College of Natural Science, Michigan State

University, MI, USA; [4]McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University

School of Medicine, MD, USA; [5]Division of Neonatology, Department of Pediatrics, Johns Hopkins

University School of Medicine, Baltimore, MD, USA; [6]Department of Pediatrics, Johns Hopkins

University, MD, USA; [7]Department of Genetics and Molecular Medicine, Landspitali University

Hospital, Reykjavik, Iceland.

*Corresponding author: Hans Tomas Bjornsson, htb@hi.is

Summary

Organisms have homeostatic mechanisms to respond to cold to ensure survival including the

activation of the neuroprotective mild hypothermia response (MHR) in mammals at 32°C. We show

activation of the MHR by one FDA-approved medication, Entacapone, a proof-of-principle that the

MHR can be medically manipulated. Utilizing a forward CRISPR-Cas9 mutagenesis screen, we identify

the histone lysine methyltransferase SMYD5 as an epigenetic gatekeeper of the MHR. SMYD5

represses the key MHR gene *SP1* at euthermia but not at 32°C. This repression is mirrored by

temperature-dependent levels of H3K36me3 at the *SP1*-locus and globally indicating the mammalian

MHR is regulated at the level of histone modifications. We identified 45 additional SMYD5-

temperature dependent genes suggesting a broader MHR-related role for SMYD5. Our study

provides an example of how the epigenetic machinery integrates environmental cues into the

genetic circuitry of mammalian cells and suggests novel therapeutic avenues for neuroprotection

after catastrophic events.

Keywords: Targeted Temperature Management, SP1, Hypothermia, Neonatal asphyxia, cardiac arrest

Introduction

The defining feature of mammals is their ability to maintain an elevated body temperature and most maintain a relatively tight range to ensure normal cellular homeostasis[1]. The main exception is hibernation of some mammals, which is a type of controlled decrease of temperature during times of prolonged inactivity[2]. In clinical practice, physicians imitate hibernation with the use of Targeted Temperature Management (TTM) as a therapeutic strategy to minimize neurological damage following neonatal asphyxia and cardiac arrest[3,4]. The reasons why lowered internal core temperature (32°C-36°C) yields benefits are currently not fully understood[5,6]. Although some argue that the positive neurological benefit from mild hypothermia relates to lower metabolic rate, others point out that the specific upregulation of several genes during mild hypothermia does not fit that idea[7]. An obvious alternative hypothesis is that hypothermia is beneficial, because of the activation of the mild hypothermia response (MHR) itself and if so, it would be important for researchers and clinicians to understand the extent of the response. In support of this, researchers have found that several genes (e.g., *CIRBP/CIRP*, *RBM3*) consistently show increased expression during mild hypothermia[8]; two of these, *SP1* and *RBM3,* appear to have roles in decreasing neuronal death during disease states[9–13].

Scientists have recently found that SP1, a transcription factor known to bind at the promoter of many genes[14], binds to the promoter region of *CIRBP*[15] upon cold exposure at a sequence called the Mild Cold Response Element (MCRE)[15], and its binding leads to increased expression of *CIRBP* at 32°C, indicating that this regulation occurs at the transcriptional level[15]. Similarly, RBM3 is known to regulate at least one down-stream gene, *RTN3*[11], a potential neuroprotective factor in a neurodegeneration model[11]. Although this suggests that mammalian cells have a pathway culminating in SP1/CIRBP and RBM3/RTN3, little has been done to further elucidate the upstream regulation of this pathway and relatively little knowledge is available regarding mammalian cellular responses to cooling compared to heat shock responses, which have been previously extensively characterized[16].

Here, we use complementary and unbiased strategies to gain clues into the MHR upstream of *SP1*. Specifically, we use a novel *SP1* mild hypothermia indicator (MHI) containing the MCRE enhancer (SP1+MCRE-MHI) to perform compound screening of 1953 FDA-approved agents. We successfully validated the drug Entacapone that activates the endogenous MHR at the transcriptional level; this indicates that it is possible to manipulate the response without exposure to hypothermia. To identify additional gene regulators, we performed a genome-wide CRISPR-Cas9 forward mutagenesis screen[17–19], reading out *SP1*-promoter activity using the SP1-MHI. Our study yielded

multiple candidate regulators and compounds that may regulate the MHR. We focus on one of these

regulators, *SMYD5*, a histone methyltransferase, and demonstrate that the presence of SMYD5 is

temperature dependent and that a number of other SMYD5-bound genes show significant

differential expression upon mild hypothermia exposure. Our insights elucidate how an external cue

(mild hypothermia) specifically mediates its effects on the human genome through the actions of the

epigenetic machinery. These results paves the way towards the development of a novel therapeutic

means to treat asphyxia, a major cause of mortality and morbidity in otherwise healthy newborns[20].

Results

Mild hypothermia indicators show increased fluorescence at mild hypothermia (32°C).

Core temperature (euthermia) is kept at a very tight range in humans (36,5-37,8°C)[1,21] and by

definition mild hypothermia involves temperature ranging from 32-36°C (**Figure S1**, **bold**). In order to

directly visualize the promoter activity of three genes previously shown[8] to respond to mild

hypothermia (*CIRBP*, *RBM3*, *SP1*), we made eight distinct MHIs (**Supplementary table 1;**

Supplementary file 1) all of which have promoter and/or MCRE sequences, driving the expression of

either GFP or mCherry (**Figure 1A**). When transfected into HEK293 cells, the MHIs demonstrate

increased fluorescence at 32°C compared to 37°C (**Figure 1B–1C**). These findings are mirrored by

endogenous SP1 levels in a Western blot from cells collected after 16 hours at 32°C (h, **Figure 1D**).

We assessed the temporal dynamics of promoter activity using flow cytometry analysis of the three

indicators and observed a maximal increase in fluorescence of SP1-MHI between 6-8 h after a

temperature shift to 32°C (P<0.05, **Figure 1E**). This finding was supported by a separate method

(continuous monitoring of fluorescence from a Incucyte S3), again maximal intensity was reached

within 6-8 h, with SP1-MHI, rising fastest, followed by RBM3-MHI, and then CIRBP-MHI (**Figure S2A-**

S2C). Indicators were also transiently transfected into HEK293 cells and then exposed to five distinct

temperatures (26°C, 29°C, 32°C, 37°C and 40°C), for 16 h followed by flow cytometry (**Figure 1F-1H;**

Figure S2D-S2E). Consistent with previous literature[8,15], and our own validation in HEK293 cells

(**Figure 1C**), indicators representing all three genes (*SP1*, *CIRBP*, *RBM3)* had significantly (P<0.0001)

increased fluorescence at 32°C compared to 37°C/40°C (**Figure 1F-1H**) and the response was

uniformly strongest at 32°C (**Figure 1F-1H**). At the same time, the SP1- and CIRBP-MHIs do not show

increased activity at lower (26°C, 29°C, **Figure S2D-S2E**) or higher (40°C) temperatures (**Figure 1F-1G**),

supporting the notion that our MHIs specifically reflect the MHR and are not activated by general cell

stress (i.e., heat stress or moderate hypothermia). These data suggest that all three genes are

regulated at the transcriptional level and the time-lag in the response upon hypothermia, measured

in hours rather than minutes, indicates that there must be additional regulators operating upstream

of these genes. Some would argue that storing cells on ice (i.e., during fluorescent activating cell

sorting (FACS)) could possibly lower the temperature of cells to the hypothermic range and activate

the MHR, but we found that even prolonged storage on ice (8 h) was unable to activate the MHI

(**Figure S3A-S3D**), ensuring that storage on ice during the study could not act as a potential

confounder.

Figure 1. MHIs based on all three genes (*SP1*, *CIRBP*, *RBM3*) show strongest effect at 32°C. (A) A
schematic of MHI structures. (B-C) Representative images of SP1-Lenti-MHIs and CIRBP-Lenti-MHI,
respectively, at 37°C and 32°C as visualized by fluorescent microscope. (D) Western blot
quantification demonstrating increased SP1 after 16 h at 32°C compared to 37°C. Shown are data
points, each data point is a biological replicate, mean and SD where applicable. Significance levels
were calculated with an unpaired one-tailed t-test in GraphPad Prism. (E) Geometric mean
fluorescence (gMFI, flow cytometry) at different timing of hypothermia exposure for SP1-MHI. Each
data point is a technical replicate. (F-G) Mean fluorescence (MFI, flow cytometry) of SP1-MHI and
CIRBP-MHI (16 h at 32°C, 37°C, 40°C), respectively. Shown are data points, each point is a technical
replicate, mean and SD where applicable. Significance levels were calculated in GraphPad Prism with
Šidák's multiple comparison test. (H) gMFI of RBM3-MHI (16 h at 32°C and 37°C). Significance level
were calculated with unpaired one-tailed t-test in GraphPad Prism otherwise data shown as in (G-H).
* = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Activation of the MHR at 32°C is uniformly seen in cells of diverse origins.

In addition to HEK293 cells, which has been widely used in prior studies[11,15,22], we sought to

verify that our novel MHIs also work in diverse cellular systems. We transiently transfected three

individual MHIs (SP1-, RBM3-, SP1+MCRE-MHI) into six distinct cell lines: (HEK293, HCT-116, HeLa,

Jurkat, K562 and SK-N-SH), representing diverse cellular origins including embryonic kidney, colon,

cervix, T lymphocytes, lymphoblasts and neurons, respectively. After exposing these cells to 26°C,

29°C, 32°C, 37°C and 40°C for 16 h and quantifying the fluorescence by flow cytometry, we observed

highest fluorescence at 32°C independent of cellular origin (**Figure S4**) and no effect for moderate

hypothermia (26-29°C, **Figure S5, Figure S2E-S2F**) with the exception of SK-N-SH and K562 cell lines

(**Figure S5D-S5E**). The strongest activation of MHI at 32°C was detected in the cell-types previously

described to have neurological characteristics (HEK293 and SK-N-SH)[23–25] and these were also the cell

lines that showed significant activation of all three MHIs at 32°C (**Figure 1F-1H; Figure S4**). This is

consistent with prior literature showing that the RBM3-MHR response is strongest in young

neurons[26]. Since many of the key studies in the field have used HEK293 [11,15,22] and they have neuronal

characteristics[23,25] we used this cell line for experiments going forward.

Compound screening reveals MHR activators at euthermia.

In our search for therapeutic targets for TTM, we sought to identify compounds that could

activate the MHR at 37°C as this could yield insights into the involved biochemical pathways. To

explore this in an unbiased manner, we performed a systematic library drug screen (1953 FDA-

approved inhibitors, **Figure 2A**) using HEK293T cells transfected with the SP1+MCRE-MHI. Cells

transfected with SP1+MCRE-MHI were exposed to the drugs (20 µM concentration) and fluorescence

was measured after 16, 28 and 40 h of drug exposure (**Figure 2A**). We focused on drugs that increase

the MHI fluorescence at least ≥3.75 times at one of the time points (**Supplementary Table 2**). Among

these drugs we saw an apparent albeit non-significant enrichment of Tyrosine Kinase Inhibitors (TKI,

7/80, p<0.075) including Orantinib, Dasatinib (hydrochloride, monohydrate), Poziotinib, Rociletinib,

Afatinib, and Lapatinib[27–31]. We validated three from this group (Dasatinib, Orantinib, and Pozitinib)

as well as selecting another 5 drugs that showed one of the greatest responses in the screen

(Cinacalcet, Entacapone, Erythromycin, Naloxone, and Tamoxifen, **Figure 2B**). For validation of the

drug screen HEK293 cells expressing SP1-MHI were individually exposed to each of the 8 compounds

(20 µM) for 16 h at either 32°C or 37°C followed by flow cytometry. Two out of the eight (25%),

showed significantly (P<0.001, P<0.0001) increased fluorescence at 37°C (Entacapone, Poziotinib,

Figure 2C-2D), indicating these two drugs can potentially induce the MHR. Entacapone also had

significantly (P<0.01) enhanced fluorescence levels at 32°C (**Figure 2C**), suggesting that the effect is

additive and thus synergistic to the regular response and thus likely independent of the SP1 target

itself. Other compounds did not increase the fluorescence of the SP1-MHI for the dose tested (**Figure**

S6A-S6F). To exclude the possibility of cell cycle arrest or increased apoptosis[31,32], we evaluated the

cell cycle phase in HEK293 cells exposed to Poziotinib and Entacapone and no apparent signs of cell

cycle arrest were observed for either compound, other than a slight increase in the percentage of

total cells in the S phase of the cell cycle after Entacapone exposure (**Figure S6G**). After exposing

HEK293 cells to the apoptotic agent H_2O_2 we could not see any signs of activation of the SP1-MHI

(**Figure S6H**), but we observed a significant (P<0.01) increase of fluorescence after only 4 h at 32°C

(**Figure S6H**) suggesting that cell stress may potentiate the response. To our surprise, three drugs

(Cinacalcet, Dasatinib, and Tamoxifen) showed significantly decreased fluorescence of the SP1-MHI

at 32°C during this validation (**Figure S6A, S6B and S6F**). To evaluate the endogenous effects on SP1

we carried out Western blotting after 24 h exposure to the compounds at either 32°C or 37°C. For

Entacapone (20 µM concentration) we observed a significant increase of SP1 at 37°C (P<0.05, **Figure**

2F). We were unable to test Poziotinib at 20 µM due to low protein yields therefore we performed

the western blot at a lower dose (4 µM) and for this dose we only saw a trend towards increased SP1

at 37°C (**Figure 2G**). When we looked at mRNA levels for *SP1*, *CIRBP* and *RBM3* at 32°C and 37°C after

16 h of Entacapone and Poziotinib exposure (20 µM), we only observed a significant (P<0.01)

elevated *RBM3* mRNA levels upon Poziotinib exposure at 37°C (**Figure S7**). This could suggest that the

drugs may impact both transcriptional and post-transcriptional levels of key factors of the MHR.

Figure 2. Poziotinib and Entacapone activate the MHR. (A) A schematic overview of our drug screen

in HEK293T cells transfected with SP1+MCRE-MHI. (B) Representative drugs were found to cause

more than ≥3.75 times increase of SP1+MCRE-MHI fluorescence. The entire list is provided as

Supplementary Table 2. (C-D) gMFI of SP1-MHI as measured by flow cytometry after 16 h of either

32°C or 37°C with or without Entacapone (ENT, C, 20 µM) and Poziotinib (POZ, D, 20 µM) exposure.

Shown are data points, each point is a technical replicate, mean and SD where applicable.

Significance levels were calculated in GraphPad Prism with Šidák's multiple comparison test.). (E-F)

Western blot of SP1 after exposure to Entacapone (ENT, E, 20 µM) and Poziotinib (POZ, F, 4 µM) or

vehicle (DMSO) after 24 h. Shown are data points, each data point is a biological replicate, mean and

SD where applicable. Significance levels were calculated with unpaired one-tailed t-test in GraphPad

Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

CRISPR-Cas9 Knockout Screen with SP1-MHIs reveals candidate regulators.

In an effort to map regulators of the cooling response (*SP1*) in an unbiased manner, we used

an available genome-coverage lentiGuide-Puro pooled sgRNA library[17–19] in combination with our

SP1-MHI to uncover transcriptional regulators of *SP1* during MHR. We chose to use the SP1-MHI, as

our prior data indicate that this gene is regulated at the transcriptional level (**Figure 1**) and is the

earliest key MHR gene to achieve maximal response at 32°C (**Figure S2A-S2C**). However, given that it

does not reach its maximal response until 6-8h of hypothermia exposure, we hypothesized that there

may be unknown regulators of the *SP1* locus that function earlier in the MHR process. We performed

the screen on HEK293WT+Cas9 cells expressing SPI-MHI (HEK293WT+Cas9+SP1), transducing the

sgRNA library at the rate of 0.3 sgRNA per cell. After puromycin selection and exposure to

hypothermia (32°C) for 16 h (to activate the MHR), we sorted out the 5% most and least fluorescent

cells (**Figure 3A**) to capture repressors and activators of *SP1*, respectively. We saw a consistent shift

towards lower fluorescence for the transduced HEK293WT+Cas9+SP1-MHI (green) compared to the

positive control (non-transduced HEK293WT+Cas9+SP1-MHI, grey) (**Figure 3B**). The observation of

such a large fluorescent shift of cells (less SP1-MHI fluorescence) at hypothermia suggests the

involvement of many sgRNAs and therefore likely many genes are upstream of *SP1*. Furthermore, this

suggests that the majority are activators rather than repressors of the SP1 arm of the MHR. We

amplified guides for each sorted group as well as the negative control (transduced HEK293WT+Cas9)

and performed deep next-generation sequencing of their distribution from sorted and un-sorted

fractions (negative control). To identify guides enriched in the sorted populations, we used the

MaGeCK pipeline[33] to computationally predict genes with enrichment (**Figure 3C-3D**). We decided to

cast a wide net and look at all genes that had a -Log10 (RRA score) > 2.5 to not miss possible

regulators (**Supplementary Table 3**). We excluded 21 genes that were seen in lists for both the high

(repressor list) and low (activator list) sorts of transduced HEK293+Cas9+SP1-MHI (**Supplementary**

Table 4), as these may be related to proliferation rather than the MHR itself, and a validation of two

of these (*RGS21* and *MALT1*) supported that idea (**Figure S8**). We were left with a list of 495 genes for

the SP1-repressors and 61 for SP1-activators – a clear overrepresentation of repressors to activators

(**Supplementary Table 3**). Of note, there were two microRNAs on the top list of SP1-repressors and

activators (*hsa-mir-4314* and *hsa-mir-6739*) and both are predicted to target SP1 (mirdb.org).

Since the ERBB2/HER2 signaling is known to regulate SP1 binding through phosphorylation[34],

and is the target of tyrosine kinases inhibitors such as Poziotinib (**Figure S9A**), we looked through

both lists and noted that ERK was found on the SP1-repressor list (**Figure S9A**). On the Qiagen ERBB2 Pathway (from Qiagen Inguinity Pathways) there are 22 genes upstream of *SP1*, and among those 3 were found on the SP1-repressor list (*RAS*, *ERK*, *STAT5*, P<0.02). We also used the Molecular Signature Database (MSigDB) to establish likely pathways from genes that had a -Log10(RRA) score over 0.5 from each screen (**Figure S9B-S9C**) and observe one shared pathway (Pentose Phosphate Pathway (3P)). Of interest there were two categories in the SP1-activator analysis that implicate cytosolic calcium regulation; calcium influx has been shown to be an immediate event in cells exposed to hypothermic stimuli[35]. As we were particularly interested in a possible transcriptional regulator of *SP1*, we annotated our lists and identified 50 (10%) on the SP1-repressor list and 4 (6%) on the SP1-activator list (**Figure 3E**) as transcriptional regulators. Of these, ten (*GATAD2*, *POU4F3*, *ZNF71*, *KAT5*, *ZNF260*, *ANKZF1*, *SMYD5*, *DMRT1*, *TAF13*, *NKX6-2*) had a -Log10(RRA) score above 3,5 in either the repressor or activator screen (**Figure 3F**). For a gene from our list to act as a direct repressor or activator for *SP1* we reasoned that direct regulators from the repressor list should have known functions as repressors and genes from the activator list should have known activating functions. Four genes from the list appeared to have opposing functions (*KAT5*, *ZNF260*, *NKX6-2*, *ANKZF1*, **Figure 3F**, not bold)[36–39] leaving six genes (**Figure 3F,** bold)[40–46] on our list. Histone modification has previously been known to dictate temperature-dependent sex determination in reptiles[47] and vernalization in plants[48], and one of these six targets (*SMYD5*), is a histone methyltransferase, known to catalyze H4K20, H3K9 and H3K36 trimethylation[44,45]. Interestingly, *SMYD5* is a known interactor with *KDM6B* in fish[49] and *KDM6B* mediates the previously described epigenetic temperature dependent sex determination switch in reptiles[47]. Also, like CIRBP and RBM3, SMYD5 is a known resident of stress granules[50]. Since SMYD5 histone tail modifications are known and their levels can be measured at its targets we argued that this might be useful for validation purposes. Thus, we decided to further focus our efforts on *SMYD5*.

Figure 3. Genome wide CRISPR-Cas9 knockout screen on SP1-MHI reveals multiple potential inhibitors/activators of the MHR. (A) Overview over the genome-wide CRISPR-Cas9 Knockout approach for the HEK293WT+Cas9+SP1 cell line. (B) Fluorescence and sort gates of 4 replicates of transduced HEK293WT+Cas9+SP1 cells (green), negative control (HEK293WT, black) and positive control (HEK293WT+Cas9+SP1, grey). (C-D) Genes marked with their name in red are transcription regulators that have -Log10(RRA) score >3.5 and have repressive functions and genes marked with their name in green are transcription regulators that have activating functions and -Log10(RRA) score >3.5. Colored dots represent genes that have -Log10(RRA) score >2.5, where orange dots indicate genes that have positive log fold change (LFC) and blue dots indicate genes that have a negative LFC, from either the SP1 activator (C) or repressor (D) screen. (E) A summary of transcriptional regulators

from our screens, activators (green) on left, repressors (red) on right. (F) A summary of top 10

transcriptional regulators ranked by their -Log10 (RRA score). Bold genes can act as a direct repressor

(red) or direct activator (green).

SMYD5 is a direct repressor of *SP1*.

To further explore the role of SMYD5, we took advantage of a newly published dataset[44]

from a CUT&TAG assay in mouse Embryonic Stem Cells (mES) with an overexpressed SMYD5-Flag

plasmid. Upon re-analysis of their data, we noticed that SMYD5, has strong peaks at the promoters of

both *Sp1* (130K) and *Cirbp (70K),* but not *Rbm3* (**Figure 4A**). These peaks occur at evolutionarily

conserved sites and overlap with CpG islands (**Figure 4A**). We took advantage of an RNASeq dataset

on *Smyd5* knockout (KO) mESC from the same study[44] and reanalyzed it which yielded approximately

equal amounts of upregulated and down-regulated genes among SMYD5 bound loci (**Figure 4B**)

supporting prior reports that histone marks deposited by SMYD5 can either repress or activate[44,45].

When looking at the three loci (*Sp1*, *Cirbp*, *Rbm3*), we found that *Smyd5*-KO yielded increased

expression of *Sp1* but not *Cirbp* or *Rbm3* (**Figure 4C**) suggesting a direct role for SMYD5 as a *Sp1*

repressor in a murine system. In this dataset, *Cirbp* showed lower expression compared to *Sp1* and

Rbm3 (**Figure S10**) which may impede interpretation. In order to validate this findings in our own

system we decided to knockdown *SMYD5* with siRNA in HEK293WT+Cas9+SP1 cell lines (**Figure 4D**),

we observed an increased fluorescence of SP1-MHI at 32°C and 37°C. *SMYD5*-KO with a guide RNA

(gRNA) and Cas9 significantly decreased amounts of *SMYD5* mRNA expression by qRT-PCR (**Figure 4E**)

and SMYD5 protein levels by Western blot (**Figure 4G**), which are rescued upon overexpression of a

gRNA-resistant Flag-SMYD5 plasmid (Flag-SMYD5 sgRNAres plasmid (labelled SMYD5sgRNA#6res),

Figure 4E and 4G). The same experiment revealed de-repression of the *SP1* transcript in KO cells by

RT-qPCR, with repression restored in rescued cells (**Figure 4F**). Furthermore, we observed a

significant increase of SP1 and CIRBP, but not RBM3 following *SMYD5*-KO at 37°C using Western blot

of the endogenous loci (**Figure 4H-4J**).

Figure 4. SMYD5 is a repressor of SP1 at 37°C. (A) Overexpressed FLAG-tagged SMYD5 in mESc, binds at promoters of *SP1* and *CIRBP* but not of *RBM3*. (B) Venn graph showing both up- and down-regulated SMYD5-target genes in an RNASeq of *Smyd5*-KO cells. Statistical analysis done with GeneOverlap package in R using Fisher's Exact test. (C) *Smyd5*-KO leads to increased expression of *SP1* but not *RBM3* or *CIRBP* in mESCs. (D) *SMYD5* knockdown by siRNA yields higher levels of fluorescence of SP1-MHI at 32°C and 37°C compared to empty vector control. Shown are data points, each data point is a biological replicate that has been normalized against same non-transfected HEK293WT+Cas9+SP1-MHI cell line and a mean of data points. Significance levels were calculated with Šidák's multiple comparison test in GraphPad Prism. (E) Relative expression of *SMYD5* compared to GAPDH from *SMYD5*-KO cells, measured by RT-qPCR at 37°C, with or without rescue with Flag-SMYD5 sgRNAres plasmid (labelled SMYD5 sgRNA#6res) which demonstrates that SMYD5-KO was

successful. Shown are data points, each data point is a biological replicate, mean and SD where
applicable. Significance levels were calculated with unpaired one-tailed t-test in GraphPad Prism. (F)
Relative expression of *SP1* compared to GAPDH from *SMYD5*-KO cells, measured by RT-qPCR at 37°C,
with or without rescue with Flag-SMYD5 sgRNAres plasmid (labelled SMYD5 sgRNA#6res). *SMYD5*-KO
leads to increased expression of *SP1* showing that effect is secondary to targeting of *SMYD5* but not
something else. Data shown as in (E). (G) Western blot quantification and representative figure using
antibodies against SMYD5 and Lamin B in a *SMYD5*-KO cell line with and without rescue with Flag-
SMYD5 sgRNAres plasmid (labelled SMYD5 sgRNA#6res) shows that *SMYD5*-KO was successful on
protein level at 37°C. Data shown as in (E). (H-J) Western blot quantification and representative
figure using antibodies against SP1, CIRBP and RBM3, respectively, in *SMYD5*-KO cells. Shown are
data points, each data point is a biological replicate, mean and SD where applicable. Significance
levels were calculated with unpaired one-tailed t-test in GraphPad Prism. * = P<0.05, ** = P<0.01,
*** = P<0.001, **** = P<0.0001.

SMYD5 shows temperature dependent expression *in vitro* and *in vivo.*

To test whether SMYD5 shows differential expression levels at 37°C and 32°C, we performed
immunofluorescence staining using a custom antibody against SMYD5[51]. We observe nuclear and
cytoplasmic staining of SMYD5 at 37°C (SMYD5; white or green, DAPI; purple, **Figure 5A, top**). The
expression of SMYD5 was significantly decreased at 32°C compared to 37°C (**Figure 5A and S11A**).
We observed a statistically significant (P<0.05) difference of SMYD5 intensity comparing the two
temperatures at the whole cell level (**Figure 5B**). Upon limiting to nuclear SMYD5-levels at the two
temperatures we see a trend towards less SMYD5 in the nucleus at 32°C when compared to 37°C
(**Figure 5C**). We also validated using an anti-FLAG antibody in cells with overexpression of SMYD5-
FLAG[44], which yielded a similar result (**Figure S11B**). We validated further by performing a Western
blot from HEK293 at 32°C and 37°C, showing significantly less SMYD5 at 32°C when compared to 37°C
(**Figure 5D**). This difference was not noted at the transcriptional level (**Figure 5E**) indicating that
SMYD5, is primarily regulated post-transcriptionally. Finally, we show that the brains of mice cooled
in a standard regimen used to treat neonatal asphyxia[52] show a similar temperature-dependent
decrease in SMYD5 after 6 h of cooling (**Figure 5F**), indicating that SMYD5-dependent impact on the
MHR during a standard TTM regimen in mice.

Figure 5. SMYD5 is degraded at 32°C *in vitro* and *in vivo*. (A) Intensity of endogenous SMYD5 at 37°C (left) and 32°C (right), 6 h incubation. (B) Quantification of cellular SMYD5 intensity. Shown are data points, each data point is a biological replicate, mean where applicable. Significance levels were calculated with unpaired two-tailed t-test in GraphPad Prism. (C) Quantification of nuclear SMYD5 intensity. Data shown as in (B). (D) Western blot and quantification using SMYD5 antibodies at 37°C

and 32°C, 6 h incubation. Significance levels calculated with one-tailed t-test in GraphPad Prism
otherwise data shown as in (B). (E) qRT-PCR results for SMYD5 expression at 37°C and 32°C, 6 h
incubation. Data shown as in (D). (G) Less SMYD5 staining in mice that had been cooled down to 32°C
compared to mice kept at euthermia 37°C in brain slices from the hippocampal region of a mouse
model of TTM. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Epigenetic and transcriptional consequences of temperature-dependent changes in
SMYD5.

As SMYD5 is decreased at 32°C and SMYD5-binding peaks overlap a CpG-island (**Figure 4A**),
we also performed DNA methylation (DNAm) analysis using the 850 K EPIC Illumina array and
observed limited DNAm at 16/18 sites and no differential DNAm at *SP1*, *CIRBP* or *RBM3* promoters at
32°C for 6 h compared to 37°C at CpG´s spanning the gene regions (**Figure S12, Supplementary Table**
6). However, there was differential DNAm at other CpGs loci after 6 h exposure at 32°C
(**Supplementary Table 7**) supporting the idea that the MHR may be regulated by the epigenetic
machinery. To further explore the epigenetic consequences of differential SMYD5 availability at 37°C
versus 32°C, we performed CUT&RUN using antibodies against H3K36me3, H4K20me3, H3K9me3,
H3K4me3, H3K27me3, SMYD5 and H3 (control) in HEK293 cells and human neural progenitor cells
(NPCs). We choose H3K36me3, H4K20me3 and H3K9me3 since these modifications have been
previously shown to be deposited by SMYD5[44,45]. We also chose to look at H3K4me3 and H3K27me3
as these are general open and closed chromatin modifications and are also known to change during
vernalization in plants[53]. We saw less H3K36me3 over *SP1* at 32°C compared to 37°C in NPCs (**Figure**
6A) and HEK293 cells (**Figure S13A**) as well as decreased global SMYD5 binding (**Figure 6B**), all
consistent with lower SMYD5 amounts at 32°C (**Figure 5**). We further observed a global deficiency of
H3K36me3 at 32°C compared to 37°C seen in both HEK293 cells and NPCs but more prominent in
NPCs (**Figure 6C, Figure S13B**) indicating that this modification is used to regulate additional genes
perhaps through SMYD5 but levels of SMYD5 mirror this effect (**Figure 6B**, **Figure 5B-5E**). We also
observed a global H3K4me3 deficiency at 32°C compared to 37°C (**Figure 6D**), although there were
also 1288 genes that had a unique H3K4me3 modification at 32°C which was not seen at 37°C
(**Supplementary Table 8**). For H3K9me3 we saw a deficiency of histone modification marks upstream
of gene bodies in NPCs (**Figure 6E**). We did not see a difference for other histone modification when
comparing 32°C and 37°C in NPCs (**Figure S13C-S13D**). CUT&RUN for H3K36me3 also revealed a
number of significantly differentially modified genes at 32°C when compared to 37°C (**Figure 6F,**
Figure S13E, Figure S14, Supplementary Table 5) including a number of epigenetic regulators such as
SETD1B and *BAZ2A* (**Figure 6F, Figure S14A, Figure S14E**). To identify additional genes regulated by
SMYD5 and temperature, we performed RNASeq in mouse NPCs (mNPCs) at the two temperatures.
This yielded 2890 differentially expressed genes (DEGS) that were upregulated and 3133 that were

downregulated at 32°C compared to 37°C. Interestingly, when we overlap upregulated DEGS in

mNPCs at 32°C with SMYD5-bound and repressed genes (*Smyd5*-KO upregulated) at 37°C in mESCs

we observe a statistically significant overlap of 26 genes (P=5.4e-04, **Figure 6G**). Similarly, when we

overlap down-regulated DEGS in mNPCs at 32°C with SMYD5-bound and upregulated genes (*Smyd5*-

KO downregulated) at 37°C in mESC we also observe an overlap of 19 genes (P=0.029, **Figure 6H**).

This suggests that SMYD5 regulates additional genes in a similar manner to *SP1* (as a repressor at

euthermia) as well as regulating some in an inverse manner (normally as an activator at euthermia)

in murine models. This includes several genes upregulated at 32°C with a theoretically appealing role

for hypothermia responses such as *Klf15* (role in brown adipose tissue metabolism)[54] and *Thrap3*

(role in thyroid metabolism and adipocyte differentiation)[55,56]. Among the genes normally

upregulated at 37°C but down-regulated at 32°C, *Bnip3L*, a pro-apoptotic gene[57] is of particular

interest given the previously observed cytoprotective role of mild hypotherma[10–12]. We provide the

entire list of 45 SMYD5-temperature dependent genes as **Table 1.** When we looked more closely at

the list of genes that had unique H3K4me3 modification in NPCs at 32°C and overlapped those genes

with human orthologs of upregulated DEGS in mNPCs at 32°C we saw a significant overlap of 169

genes (P=2.4e-03, **Figure 6J, Supplementary table 9**). Only 6 genes are bound by SMYD5 and

upregulated with *Smyd5*-KO at 37°C in mESCs as well as upregulated at 32°C in mNPCs (**Table 1**) and

have a unique H3K4me3 peak at 32°C in NPCs (**Supplementary table 8**). Interestingly *Asxl1,* a

member of the Polycomb family[58] a family which is known to take part in vernalization in plants[48], is

among these 6 genes. This indicates that there may be additional epigenetic MHR-responses that are

independent of SMYD5. Finally, we demonstrate that on average these 45 genes, regulated by

SMYD5, have lower H3K36me3 modifications at 32°C versus 37°C at their gene bodies as well as

fewer modifications over the 3ʹ-UTR region at 32°C in NPCs and HEK293 cells (**Figure S13F-S13G**). As

well as, these genes show a different H3K4me3 and H3K9me3 modification pattern at 32°C and 37°C

when looking at their 3ʹ- and 5ʹ-UTR regions (**Figure S13H-S13J**).

Figure 6. SMYD5 is a novel epigenetic gatekeeper of the MHR. (A) Increased H3K36me3 is observed at the *SP1* promoter and genebody at 37°C compared to 32°C. (B) SMYD5 chromatin binding is globally decreased (all genes) at 32°C compared to 37°C. (C) There is a global deficiency of H3K36me3 at 32°C compared to 37°C. (D) There is a global H3K4me3 deficiency at 32°C compared to 37°C. (E) Upstream of gene bodies there is a global H3K9me3 deficiency at 32°C compared to 37°C. (F) At the gene-specific levels there are a number of genes, that have more or less H3K36me3 at 32°C compared to 37°C in NPCs. Red circles indicated genes that have Log$_2$ fold change over 1 and p-value

less than 10^{-6}, gray circles indicate Log_2 fold change over 1 but non-significant p-value and black
circles indicate non-significant genes. (G-H) DEGS from RNASeq in mNPCs at both temperatures,
show overlap with SMYD5-bound genes that are differentially expressed upon *Smyd5*-KO in mESCs at
37°C. Statistical analysis done with GeneOverlap package in R using Fisher's Exact test. (J) There is an
significant overlap of DEGS and genes that have a unique H3K4me3 modification at 32°C. Statistical
analysis as in (G-H).

SMYD5-repressed genes ↑ at 32°C (SMYD5 represses at 37°C)

Gene	Function	Role
Mybl2	Transcription factor	Cell-cycle control
Klf15	Transcription factor	Brown-tissue metabolism
Thrap3	Transcriptional co-activator	Thyroid hormone receptor binding
Tex15	DNA repair protein	Role in testicular meiosis
Ewsr1	RNA binding	Role in Ewing sarcoma
Tfcp2l1	Transcription factor	Pluripotency in stem cells
Amotl2	Angiostatin binding factor	Role in angiogenesis
Phf20	Methyllysine-binding protein	Histone modification
Cd24a	Sialoglycoprotein	Cell differentiation
Nt5e	Nucleotidase	Nucleotides metabolism
Neo1	Cell surface receptor	Immunoglobulin superfamily
Gabbr1	GABA receptor	Inhibitory neurotransmitter
Asxl1	Chromatin binding protein	Polycomb family member
Tulp4	Tubby Superfamily Protein	Ubiquitination function
Pkn1	Serine/threonine-protein kinase	Protein kinase C superfamily
Hnrnpul1	RNA-binding protein	hnRNP family
Mier1	Transcriptional regulator	Mesoderm induction early response
Fam21	WASH complex member	Endosomal protein sorting
Trip12	E3 ubiquitin-protein ligase	Ubiquitin regulation and DNA repair
Lamc1	Extracellular matrix glycoprotein	A laminin
Rell1	RELT-like protein	Regulates the p38MAPK cascade
Xrn2	5'-3' exonuclease	Promotes transcription termination
Hspg2	Multidomain proteoglycan	Encodes Perlecan protein
Ctc1	Telomere maintenance protein	Limits telomere degradation
Stag1	Component of cohesin complex	Role in cohesion of sister chromatids
Ass1	Enzyme	Role in urea cycle

SMYD5-upregulated genes ↓ at 32°C (SMYD5 activates at 37°C)

Gene	Function	Role
Usp28	Ubiquitin Thioesterase	DNA damage checkpoint
Ccnd1	Cyclin D	Cell cycle regulation
Dzip3	Ubiquitin-protein transferase	Spermatogenesis
Rad50	DNA double-strand break repair	Cell cycle regulation
Arglu1	Enables cadherin binding	Mediator complex
Sltm	RNA binding activity	Regulation of transcription
Slc16a3	Solute transporter	Lactic acid and pyruvate transport
Ier5l	Unknown function	Parolog of IER5
Acly	Enzyme	Synthesis of cytosolic acetyl-CoA
Agps	Enzyme	Lipid biosynthesis

Apobec3	Cytidine deaminase	RNA editing
Ivns1abp	Viral binding protein	Response to viral infection
Dus3l	Synthesis of dihydrouridine	RNA editing
Upp1	Uridine phosphorylase	Ribonucleosides dynamics
Ldha	Enzyme	Lactate to Pyruvate conversion
Scd2	Enzyme	Fatty acid biosynthesis
Smarcad1	Chromatin remodeler	Role in heterochromatin formation
Bnip3l	Pro-apoptotic protein	Induces apoptosis
Esrra	Nuclear receptor	Mitochondrial biogenesis regulator

Table 1: Summary of temperature responsive SMYD5-regulated genes.

Discussion

Histone modification has previously been found to help integrate the effects of cold
temperature exposure in diverse organisms. For example, in plants, vernalization is the basis of how
cold exposure influences the rate of flowering is mediated through a Trithorax-Polycomb switch[48]. In
some reptiles, sex determination is achieved through a temperature dependent mechanism and for
one such, the red eared slider turtle (*Trachemys Scripta Elegans*), the mechanistic basis of this
regulation acts through a histone methylation switch[47]. To uncover additional regulators of the MHR
we used the MHI to identify an FDA-approved drug, Entacapone, that can induce the SP1-mild
hypothermia response as well as conducting a forward CRISPR-Cas9 mutagenesis screen resulting in
recognition of *SMYD5* as a repressor of the MHR at euthermia. Such agents can be useful to
understand the mechanistic underpinnings of the MHR but also for therapeutic development for
patients at risk of neurological sequela after catastrophic events.

Our current MHIs allow for some temporal dissection yet they all currently harbor stable
fluorescent protein, which may be less time sensitive to detecting dynamic differences compared to
real time qPCR. Thus, our current MHIs may be the most useful to map the start of the response (as
we have done in **Figure 1**). However, in future work, we hope to create unstable fluorescent
molecules that break down quickly to help us map the kinetics of different parts of the response, for
example how quickly the response is resolved (i.e. termination).

Although, our drug screen and validation may have limitations (only three timepoints, single
dose), we observe an overrepresentation of TKIs. The reason why some TKIs as well as other drugs
did not validate could be as the drug screen was performed using an MHI containing the MCRE (to
increase signal) but validation was performed on a MHI lacking the MCRE (endogenous situation in
mammals[15]). Of the TKIs we have further shown that Poziotinib, a known pan-HER inhibitor[29], was
able to increase *RBM3* mRNA but not *SP1* or *CIRBP* mRNA (**Figure S7**) or SP1 (**Figure 2F**) at 37°C with

the single dose tested. The lack of increased SP1 quantified by Western blot could be caused by low

(4 µM) Poziotinib dosage compared to RT-qPCR and flow cytometry (20 µM). Here, we did identify

and validated Entacapone as a compound that can activate the SP1-MHR at euthermia (**Figure 2E**).

Interestingly, Entacapone is COMT-inhibitor which increases dopamine availability with blocking its

degradation in synapses[59] and it is known that synaptic plasticity can be influenced with temperature

changes.[60] Furthermore, Entacapone is a neurotransmitter breakdown inhibitor (COMT), supporting

the fact that several candidates (**Figure 3C-3D**) from the CRISPR-Cas9 forward mutagenesis screen

appear to bind the neural synapse and/or Syntaxin (**Supplementary table 3**) such as *STXBP1* (SP1

repressor) and *TXLNB* (SP1 activator) which bind to Syntaxin (**www.genecards.org**), and play a role at

the neuronal synapse as well as having a role in dopaminergic synaptic transmission[61]. which may be

worth exploring in future work.

Our data support the notion that SP1 and RBM3 may be parts of individual arms of the

mammalian hypothermia response as SMYD5 only regulates *SP1* but not *RBM3*. We interpret the lag

in the response (at least 6 hours to signal) and the obvious shift in fluorescence of mutagenized cells

to mean that there are likely many genes upstream of the currently tested point of the response

(*SP1*). Despite a global shift towards activators, the yield from the enrichment screen yielded more

putative *SP1* repressors compared to *SP1* activators. This could be caused by the setup of the screen

as it may be easier to find genes that increase fluorescence (repressors) than those with decreased

fluorescence (inhibitors) which may overlap with cells that have lost the SP1-MHI for various reasons.

Another limitation to our forward screen is it can only identify cell-autonomous genes and cannot

find essential genes. However, it is a starting point, and it has already provided several candidates

that can be subsequently tested for effect using our MHIs. Of interest, SMYD5 and ERBB2/HER2 (the

receptor Poziotinib blocks) have been previously found to functionally interact in a forward

mutagenesis screen of genes that play a role in metastatic reactivation and tumor dormancy[62] and

additionally *CIRBP* transcript levels, correlates with prognosis in for luminal A/B breast cancer which

is influenced by HER2 status[63]. In fact, our SP1-repressor screen revealed several downstream factors

of ERBB2 signaling (*RAS*, *ERK*, *STAT5,* **Figure S7A**), indicating that this pathway may be upstream of

SMYD5 and worth exploring. Alternatively, SP1 itself may be a target of phosphorylation, and as SP1

is predicted to bind at its own promoter[64], perhaps both signals are required for full effect.

Here, by uncovering *SMYD5* as a negative epigenetic regulator of SP1, a key factor of the

mammalian MHR, we provide an example of how histone methylation integrates temperature cues

into a fundamental pathway in mammals. Our present model is that SMYD5 exerts its effects at 37°C

to ensure that *SP1/CIRBP* are not upregulated without the cold stimulus, however, upon cold

stimulus (32°C), SMYD5 is degraded or its protein synthesis is degraded enough to lead to loss of

SMYD5 levels and thus allows the pathway to proceed. Global cellular SMYD5 levels are decreased at

32°C *in vivo* and *in vitro* (**Figure 5**) as well as there is a trend towards less SMYD5 in the nucleus at

32°C (**Figure 5C and 5F**) this could happen as a result of nuclear exclusion of SMYD5 or there could be

less *SMYD5* translation or SMYD5 could undergo proteasomal degradation. Per bioGRID, SMYD5 is

bound to four ubiquitinators (PARK2, UBA52, TRIM25, USP34) so this may be an avenue for further

study as such factors often target proteins to the proteasome. Previously, decreased production of

protein upon cold stimulus has shown to act through AMPK mediated phosphorylation of eEF2K[22],

and decreased translational efficiency could play a part. But we are unaware of biological examples

of directed protein degradation in response to mild hypothermia. This requires further investigation.

SMYD5 itself appears to regulate at least 45 other factors that show temperature

dependence, 26 in which SMYD5 acts as a repressor (like *SP1*) and another 19 where SMYD5 acts as

an activator at euthermia. Some of these genes look very plausible MHR participants, especially

BNIP3L,[57] a pro-apoptotic factor which is down-regulated with hypothermia and may thus contribute

to the neuroprotective effect of mild hypothermia. There are also genes that play a role in thyroid

and adipocyte function (*THRAP3*)[55,56], brown adipose tissue (*KLF15*)[54] and testes (*TEX15*)[65,66], all

target tissues for mammalian temperature responses. Coinciding with decreased global H3K4me3 at

32°C, we also observed decreased H3K36me3 at 32°C over the gene *SETD1B* which encodes a

H3K4me3-methyltransferase (**Figure 6 and S14E**). In oocytes *Setd1b*-KO can cause gain or loss of

H3K4me3[67]. Accordingly, this could be one potential mechanism underlying H3K4me3 changes at

32°C and warrants further investigation. It is interesting that H3K36me3 has previously been shown

to regulate temperature-regulated alternative splicing in plants[68] and in the fungus *Saccharomyces*

cerevisiae, H3K4 and H3K36 methyltransferases (Set1 and Set2) regulate alternative

polyadenylation[69] as well as in mice H3K36me3 has a direct impact on alternative splicing of

rewarding neurobiological factors in the brain[70]. In mammals temperature dependent alternative

splicing[71] and alternative splicing of *CIRBP* and *RBM3* has been observed[72–74]. Perhaps alternative

splicing or polyadenylation plays a role for *SP1* or one of the other 45 SMYD5-bound factors

elucidated here.

In summary, we have developed and validated tools to interrogate the MHR, and used them

to show for the first time that a compound exists that can activate the MHR as well as shone light on

potential regulatory genes in the MHR pathway. We show that *SMYD5* is a temperature-dependent

epigenetic gatekeeper of the MHR and for the first time we show how histone methylation signature

(H3K36me3, H3K9me3 and H3K4me3) change in mammals with MHR exposure, providing a specific

example of how mammalian cells use epigenetics to respond to temperature cues.

Acknowledgements: We thank SCRU lab (UI) for gifting us pMD2.G (Addgene, #12259) and psPAX2
(Addgene, #12260) and H.C. Dietz and D. Valle labs (JHU), for gifting us the HCT-116, SK-N-SH, HeLa,
Jurkat and K562 cell lines. We want to thank Alan Long at the ChemCore facility at JHU and Pharmacy
of Landspitali for support regarding the drug screen and validation. We thank the Immunology
department of Landspitali University Hospital for the access and assistance with the FACS sorting. We
thank Funlab at deCODE genetics for the usage of their flow cytometry machines as well as for gifting
us the HEK293T cell line. We thank Kári Stefánsson and Ólafur Þ. Magnússon at deCODE Genetics for
providing some of the next generation sequencing for this project. We thank Erna Magnúsdóttir for
providing pAG-MNase for the CUT&RUN assays as well as guidance with the CUT&RUN experiment
and data analysis. This study was supported with funding from the Rannis Technology Development
Fund (H.T.B., #2010588), Eimskip University Fund (S.R.), Fulbright Iceland (S.R.), Icelandic Cancer
Society Research Fund (H.T.B.), Göngum Saman Research Fund (H.T.B.), Landspitali Research Fund
(H.T.B.), Félag Háskólakvenna Research Fund (S.R.). HTB is also supported by funding from Rannís
(#217988, #195835, #206806) and the Louma G. Foundation. Finally, some images in this manuscript
were created with BioRender.com.

Author contributions: H.T.B. and S.R. conceptualized project. S.R., L.Z. and H.T.B. created and
designed the MHIs. S.R., A.T., L.H.A., M.V., K.E.A. and K.J. performed cell culture. S.R., A.T. L.H.A. and
M.V. performed flow cytometry experiments. L.Z. and H.T.B. designed the approach to the drug
screen. S.R. and H.T.B. designed the approach to the forward mutagenesis assay. S.R. performed the
forward mutagenesis assay and drug screen. S.R. and K.J.A. performed the PCR for the NGS. S.R., A.T.
and K.J. validated results from drug and mutagenesis screens (Western blots (A.T., K.J.), RT-qPCR
(S.R., K.J.) and flow cytometry (S.R., A.T.)). K.J and K.M. performed IF on cells from culture and K.J.
and K.M performed the confocal imaging and K.M., K.J. S.Th.H. and S.R. did the analysis. S.R. and
M.V. performed siRNA knockdown. R.C-V. and F.J.N. provided mouse brains treated with TTM. K.J.A.
isolated and cultured mNPCs. M.V. performed cryosectioning and IF on mouse brain slices and M.V.
and K.M. performed confocal imaging as well as the analysis of the mouse brain slices. S.R., A.T. and
S.Th.H. performed CUT&RUN experiments. K.J.A. did the mNPCs isolation and RNASeq and data
analysis. S.R., K.J.A., K.J., A.T. S.Th.H. and T.R. analyzed data. J.H. provided SMYD5 antibody and K.E.A.
provided human NPCs. S.R. and H.T.B. wrote the manuscript. Project administration was carried out
by H.T.B, S.R and K.J.A.

Declaration of interest: Dr. Bjornsson is a consultant for Mahzi therapeutics. Dr. Rafnsdottir and Dr.
Bjornsson have a European patent application pending with the number 23167505.9.

  STAR Methods

Creation of novel indicators: We used the PGL4.10 plasmid (Promega, #9PIE665) as the backbone for
our CIRBP- and SP1-MHI, where the *luc2* sequence was cut from the plasmid with NcoI (NEB,) and
XbaI (NEB). Promoter sequence for *CIRBP*[75] was cloned from mouse sequence using *CIRBP* forward
primer 5'tcgataggtaccTGGCTTCACAAATGCGCCTCAGT3' and *CIRBP* reverse primer
3'cctaaggcagatctGCGAGGGGGAGCGCAAGAGT5'[75]. Restriction enzymes KpnI (NEB) and BglII (NEB)
were used to insert the promoter into the plasmid. Promoter sequence for *SP1*[76] was cloned from
human DNA using the forward primer 5'tcaagtcaggctagcGCAACTTAGTCTCACACGCCTTGG3' and
reverse primer 3'cagtgctgcctcgagGCTCAAGGGGGTCCTGTCCGG5'[76]. Restriction enzymes NheI (NEB)
and XhoI (NEB) were used to insert the promoter into the plasmid. AcGFP1 was cloned from
PT7XbG2-AcGFP1 (Novopro, #V002843) vector using the reverse primer
3'cggcggagTCTAGAATTACTTGTACAGCTCGTCC5' and the forward primer
5'taagccaccATGGTGAGCAAGGGCGAGGAGC3'. Restriction enzymes NheI (NEB) and XhoI (NEB) were
used to insert the acGFP1 into the plasmid. The neomycin selection cassette was cloned from
pROSA26-dest plasmid (Addgene, #21189) using the forward primer
5'CATTATCGTCGACTCTACCGGGTAGGGGAGGCGCTT3' and reverse primer
3'CGCCGCCGACGATAGTCAAGCTTCTGATGGAATTAGAACTTGGC5'. Restriction enzymes SalI (NEB) and
PshAI (NEB) were used to insert the neomycin cassette into the plasmid. CIRBP- and SP1-MHI were
made with and without MCRE[15]. The MCRE sequence was inserted in 5 repeats in front of the
promoter sequences with a linker sequence between. Acc65I (NEB) restriction enzyme was used to
insert the MCRE sequence into the CIRBP-MHI and Acc65I (NEB) and SacI (NEB) restriction enzymes
were used to insert the MCRE into the SP1-MHI. Sanger Sequencing was used to validate insertion
sites, size, and sequence of insertion. The *RBM3* promoter with MCRE enhancer (6 repeats) in front
of the promoter was cloned into the multiple cloning site pCMV6-AC-GFP (BlueHeron, #PS100010).
We also made lentiviral vectors for *CIRP*, *SP1* and *RBM3* on a pLV-Ex (VectorBuilder) lentiviral vector.
The *CIRBP* promoter for the CIRBP-lenti-MHI was designed from the human promoter and the *SP1*
promoter was moved upstream of the TSS for the SP1-lenti-MHI but *RBM3* promoter sequence
remained unchanged. Included were MCRE (8 repeats for CIRBP-lenti-MHI, 5 repeats for other),
promoter, neomycin selection cassette and mCherry fluorescent protein behind each promoter
sequence. The promoter region, size and backbone plasmid for each indicator is detailed in
Supplementary Table 1 and sequence of each MHI in **Supplementary File 1**. We also observed some
variation in absolute levels among individual indicators/experiments, but the pattern was always
consistent; given this, we only show data from individual experiments in any figure.

Culture of cell lines: HEK293 (gifted and DSMZ, CRL-1573, referred to as 293/293WT) cells were
cultured in DMEM/F12 with GlutaMAX (Gibco, 10565018) with 10% filtered FBS (Gibco, 10270106)
under 5% CO_2. Cells were cultured at 37°C unless otherwise stated, then 37°C served as a control. In
addition to HEK293, we used the following cell types HEK293T (gifted, CRL-3216), HCT-116 (gifted,
CCL-247), HeLa (gifted, CCL-2), Jurkat (gifted, TIB-152), K562 (gifted, CCL-243), SK-N-SH (gifted, HTB-
11), human iPS cells (ATCC, 1019) that were used to generate NPCs by neural induction of iPSCs using
dual SMAD inhibition and embryoid body generation (StemCell Technologies; 08581, 05832,
05838)[77], prior to experiments described here the NPCs were cultured in STEMdiff Neural Progenitor
Medium (StemCell Technologies, 05833) on Matrigel (Corning). As well as murine primary NPCs (see
section mNPC isolation for further information). We used culture recommendations from ATCC when
possible. Cells were sub-cultured when they had reached 70-90% confluence and media was changed

every other day. Killing curves were used for all cell lines to determine lowest selection concentration
for all selection agents.

Transfection of MHI and SMYD5-Flag overexpression plasmid: HCT116, 293WT, HEK293T, HeLa and
SK-N-SH cell lines were transfected with MHI when they had reached 70-90% confluency. We used
either Lipofectamine™ 2000 or 3000 (Thermo Fisher Scientific, 11668027, L3000015) according to
the manufacturer's protocol. The fluorescence of the indicators was measured via flow cytometry up
to 48 h after transfection. We analyzed flow cytometry data with the FlowJo Software, where we
gated on living cells, single cells, GFP positive cells and measured the fluorescence for that
population. For further analysis, we used Excel from Microsoft, GraphPad Prism (v.9) and FlowJo
(v.10.8.1). All statistical testing was done with GraphPad Prism (v.9). For transfection of SMYD5-Flag
overexpression vector[44] we used the same method as described above.

Construction of 293WT+Cas9 and 293WT+Cas9+SP1 cell lines: We made two cell lines from 293
cells, one that expressed Cas9 (lentiCas9-blast, Addgene #52962), referred to as 293WT+Cas9, and
another that expressed both Cas9 and our SP1-MHI, referred to as 293WT+Cas9+SP1. These cell lines
were made by transfecting the plasmids with lipofectamine as described above and selecting stably
transfected cells using Blasticidin selection for Cas9 and Neomycin selection for our SP1-MHI.

Drug screen: Briefly, we used Lipofectamine 3000 (Thermo Fisher Scientific, L3000015) to transfect
the SP1+MCRE-MHI into the 293T cell line. 24 h after transfection, 30000 cells were plated into each
well of a black 96 well plate. 48 h after transfection the cells were exposed to FDA Approved Drug
Screening Library (L1300-Z298012) via the High Throughput Screening Services, ChemCore Facilities
(Johns Hopkins University). Each drug was exposed in 20 μM final concentration. Total well GFP was
read with the CLARIOStar Plus Plate Reader (BMG LABTECH, settings: gain 1500, 470-15 excitation
and emission 515-20 at focal height 2.6) 16, 28, and 40 h after exposure of the library. Ratio of
increased fluorescence was calculated by dividing GFP reads from positive control (SP1+MCRE-MHI
transfected but untreated) at each time point with GFP reads of each whole well exposed to drug at
the same timepoint for wells in the same lane. For further analysis, we used Excel from Microsoft and
GraphPad Prism (v.9).

Validation of drug screen: We individually validated 8 compounds in same concentration (20 μM).
Here, we exposed the drug for 16 h, and evaluated by flow cytometry as previously described. In a
96-well plate 20,000 of either 293 or 293+SP1-MHI cells were seeded per well in a total volume of
200 μL of media. After cells had attached to the wells (the morning after), cells were exposed to 20
μM of each drug, controls were exposed to the same quantity of DMSO as drug, for 16 h and
harvested for flow cytometry analysis. Flow cytometry analysis was conducted as previously
described. Drugs that were tested for further validation after initial screen were: Orantinib
(Selleckchem, S1470), Dasatinib (Selleckchem, S7782), Poziotinib (Selleckchem, S7358), Tamoxifen
Mylan (Mylan AB*), Erythromycin (Stragen Pharma GmbH*), Naloxone (B. Braun*), Entacapone
(Orion Corporation*) and Cinacalcet (Ratiopharm GmbH*). Drugs labeled with "*" were
pharmaceutical grade medications gifted from the Pharmacy at Landspítali University Hospital.
Analysis was performed as described above.

Apoptosis assay: We based our apoptosis assay on that published by Xiang et al[78]. We exposed
293WT+Cas9+SP1 cell line either to 1-5 μM H_2O_2 or vehicle (H_2O) for 4 h at 32°C or 37°C. 4 h after
exposure gMFI of the cells were analyzed with flow cytometry as described above.

Genome Wide CRISPR-Cas9 screen on fluorescent HEK293WT cell lines that stably express SP1-MHI: Using the 293WT+Cas9+SP1 and 293WT+Cas9 (negative control) cell lines we performed a GeCKO screen, where we transduced the sgRNA pool made from library A and B (Addgene, #1000000049) in MOI 0.3[17–19]. We mostly followed the Joung et. al. protocol[19], the exceptions are as follows. For making the lentivirus, a protocol described by Kutner et. al[79] was used. Lentiviral sgRNA library concentration was done with Amicon ultracentrifugal filters (Millipore, UFC9003). 20 h after transduction with the sgRNA library selection was started with puromycin and continued for 6 days and 8 h before the cells were moved to 32°C, where they were incubated 16 h before fluorescent activating cell sorting (FACS) sorting with Cell sorter SH800Z (Sony). The 5% highest fluorescent cells for 293WT+Cas9+SP1 and and 5% lowest fluorescent cells for 293WT+Cas9+SP1 were sorted. Next, we isolated gDNA and performed a two step PCR for all samples (sorted as well as negative control). According to cell number after puromycin selection, the sgRNA library coverage was >700 times coverage for HEK293WT+Cas9+SP1. Next generation sequencing (NGS) with single read with 80 cycles and 8 indexing cycles with PhiX spike in of 20% on NovaSeq S4 at deCODE genetics for HEK293WT+Cas9+SP1 and HEK293WT+Cas9, 32-280 million aligned reads per sample. To analyze the screen data, single read fastq files for each replicate and condition were merged using cat command. Then, MAGeCK (v.0.5.9) was used to identify enriched sgRNA's in sorted samples[33,80] and sgRNA counts were normalized to internal control for sgRNA's (control sgRNAs). We used RStudio (R, v.4.1.1)[81] to visualize the results and to filter genes. We exported a list of genes that had both FDR value under 0.25 and LFC value over 2.5 for each condition in both screens. Then, we excluded genes, overrepresented in both the repressor and activators screen when compared to control as they most likely promote cell growth and therefore were present in both lists. For the pathway enrichment analysis we used EnrichAnalyzer() from MAGeCKFlute (v.1.12.0)[80].

Making of *SMYD5* knockout (*SMYD5*-KO) cells: Two ready-made sgRNA vectors (Genscript, *SMYD5* plasmid vector #2 and #6) were packed into lentivirus according to protocol described by Kutner et. al[79], for concentration we used the 4x Lentivirus concentrator solution protocol from the MD Anderson Cancer center at University of Texas (available online). We selected for stable *SMYD5*-KO cells with puromycin selection for at least 7 days.

Making of *MALT1* and *RGS21* knockout cells: Ready made sgRNA vectors (Genscript, plasmid vector *MALT1* #1 and #2, *RGS21* plasmid vector #2 and #3) were packed into lentivirus and selected as described above in the making of *SMYD5-KO* cells.

Making of gRNA resistant SMYD5-Flag plasmid (Flag-SMYD5 sgRNAres plasmid)**:** In an effort to make SMYD5-Flag resistant to CRISPR-Cas9 cutting from our two guides (guide 2&6), we used site-directed mutagenesis (Q5® Site-Directed Mutagenesis Kit, New England Biolabs) to modify predicted PAM sites of the guides (Genscript, SC1678, plasmid vector #2 and #6). We used the following primers to induce the mutations: SMYD5-2mut-F: 5'-gctctttacgAGGAAGCAGTCAGCCAGT-'3; SMYD5-2-mut-R: 5'-ttccgtaaagAGTCTCCGCAGAAGTTCC-'3; SMYD5-6-mut-F: 5'-accgatatcgAGCCTGTGACCACTGCCT-'3; SMYD5-6-mut-R: 5'-atagagcgtt-CCAGAGAAACTGTGCAGCC-'3. After site-directed mutagenesis, we isolated plasmid with Monarch Midiprep kit (New England Biolabs) and Sanger sequenced to verify changes in PAM-sites. This plasmid was then used for rescue experiments (**Figure 4E-4F**).

siRNA: We used the TriFECTa RNAi kit (IDT; TYE 563 Transfection Control DsiRNA, 51-01-20-19 (transfection control); Negative Control DsiRNA, 51-01-14-03 (negative control)) and used

predesigned DsiRNA (IDT, hs.Ri.SMYD5.13.8) to knockdown *SMYD5* mRNA in two biological clonal cell
lines of HEK293+Cas9+SP1-MHI with Lipofectamine 2000 (Thermo Fisher Scientific, 11668027) with
reverse transfection, where we seeded $1*10^6$ cells per well in a 6 well plate. One batch of cells was
exposed to 32°C and the other to 37°C for 6 hours. Data were analyzed with flow cytometry 36 h
after transfection. Further statistical analysis of the flow cytometry data was similar to that described
above.

Immunofluorescence of cell culture: HEK293 cells were seeded into an 8-well chamber slide and
then fixed and permeabilized with 4% PFA and 0.1% Triton-X 100 diluted in PBS for 10 mins at room
temperature. Then cells were washed twice with a blocking solution comprising PBS with 2.5% BSA
and 10% normal goat serum and followed by blocking for 30 mins in the blocking solution. Cells were
incubated overnight with primary antibodies (rabbit anti-SMYD5, 1:1000[51]; anti-FLAG, 1:500, (Sigma-
Aldrich, F1804-200UG)) at 4°C. The following day, cells were incubated at room temperature for 30
mins and then washed three times every 10 mins with a washing solution composed of PBS with
0.25% BSA. Cells were incubated with secondary antibodies (goat anti-mouse IgG (H+L) Alexa Fluor™
488, 1:1000 (Invitrogen™, A-11001); goat anti-rabbit IgG (H+L) Alexa Fluor™ 647, 1:1000
(Invitrogen™, A21244)) for 1 h at room temperature in the dark. Cells were washed three times every
10 mins with the washing solution and added with Fluoromount-G Mounting Medium with DAPI
(Invitrogen, 00-4959-52). Cells were imaged using confocal microscopy (FV 1200, Olympus
Fluoroview) using a 30x silicon objective (NA1.05). Images were processed by ImageJ[82,83] and
quantified using CellProfiler (v.4.2.5.)[84] . Statistics were calculated in GraphPad Prism.

Real Time Quantitative Polymerase Chain Reaction (RT-qPCR): Total RNA from cells incubated at
32°C or 37°C was isolated, using Direct-zol™ RNA Microprep (Zymo Research, R2062), according to
the manufacturer's instructions. The concentration of RNA was measured by NanoDrop (Thermo
Fisher Scientific) followed by cDNA synthesis with High-Capacity cDNA Reverse Transcription Kit
(Applied Biosystems, 4368814) on MiniAmp™ Thermal Cycler (Applied Biosystems™). RT-qPCR was
performed using Luna® Universal qPCR Master Mix (NEB, M3003) on CFX384™ Real-Time PCR
Detection System (Bio-Rad). Each biological replicate of the RT-qPCR assay in this study was carried
out in technical triplicates. Any technical replicate that deviated from other replicates by ≥ 0.4 cycle
threshold (Ct) was removed from calculations of average Ct values. The primers used are as follows:
SP1 (fwd): 5'-CACCCAATTCAAGGCCTGCCGT-3'; SP1 (rev): 5'-GGGTTGGGCATCTGGGCTGTTT-3'; RBM3
(fwd): 5'-GAGACTCAGCGGTCCAGGGGTT-3'; RBM3 (rev): 5'-CCTCTGGTTCCCCGAGCAGACT-3'; CIRBP
(fwd): 5'-CCGAGTTGACCAGGCTGGCAAG-3'; CIRBP (rev): 5'-TCCATAGCCCCGGTCTCCTCCT-3'; GAPDH
(fwd): 5'-TCAAGGCTGAGAACGGGAAG-3'; GAPDH (rev): 5'-CGCCCCACTTGATTTTGGAG-3'. SMYD5 1
(fwd): 5'-GCACTGTGCGCAAAGACCTCCA-3', SMYD5 2 (fwd): 5'-GGAAACCAGGCCAGGTTCTGCC-3',
SMYD5 3 (fwd) 5'-CGTGGAAGTCCGTTTCGTGA-3', SMYD5 1 (rev:) 5'-CTGGGCACAGGACCTGGTGGTA-
3', SMYD5 2 (rev): 5'-GGCTGCCAACCGACATTCTGCA-3', SMYD5 3 (rev) 5'-CCAGAGAAACTGTGCAGCCA-
3'.

Western blot assay: Cells were washed with PBS and then lysed for 30 min on ice in RIPA buffer
(50mM Tris HCl ph8, 150mM NaCl, 1% NP-40, 1% Sodium deoxycholate, 0.1% SDS, 2mM EDTA,
phosphatase inhibitor (either Cell signaling, 5870S or Thermo Fisher Scientific, 78437)). The lysate
was centrifuged at 16,000 xg for 20 min at 4°C and the supernatant was collected. The supernatant
was diluted with 4X loading buffer (LI-COR, 928-40004) and heated at 95°C for 5 min. The protein
concentration was measured using Pierce™ BCA Assay Kit (Thermo Fisher Scientific, 23225). 20-30 µg

of protein was loaded into each well and separated by SDS-PAGE. They were transferred to
polyvinylidene fluoride (PVDF) membranes and blocked in 5% bovine serum albumin in Tris-buffered
saline with Tween 20 for 1 hour. Primary antibodies: rabbit anti-CIRBP, 1:2000 (Proteintech, 10209-2-
AP); rabbit anti-RBM3, 1:1000 (Proteintech, 14363-1-AP); rabbit anti-SP1 1:1000 (Proteintech, 21962-
1-AP); mouse anti-Lamin B1, 1:5000 (Proteintech 66095-1-Ig); rabbit anti-SMYD5, 1:1000[51]; mouse
anti-GAPDH, 1:5000 (Abcam, ab8245) were applied to respective membranes after washing, and
incubated at 4°C overnight. Membranes were incubated with IRDye® secondary antibodies; anti-
rabbit (LI-COR,926-32213); anti-mouse (LI-COR, 926-68072), at room temperature for 90 mins. After
washing, the protein bands were visualized on the Odyssey® CLx Infrared Imaging System and protein
band intensity quantified by Image J software.

DNAm studies and analysis: 293WT (biological triplicates) were split into two batches and exposed
to 37°C and 32°C for 6 h, next cells were harvested and DNA isolated with the Zymo Quick-DNA™
Midiprep Plus Kit (Zymo Research, D4075). Samples were submitted to the Genetic Resources Core
Facility (Johns Hopkins University), bisulfite treated, and run on the Infinium MethylationEPIC
BeadChip Kit (Illumina) with two technical controls (50% and 100% DNAm, 8 samples total). Technical
controls were created using mixtures of 100% and 0% methylated HCT116 DKO (Zymo, Research
D50414-1/2). DNA methylation data (IDAT files) containing unmethylated and methylated intensity
values were imported to R (v.4.2.1) for analysis[81]. Raw data was processed using single sample
approach with preprocessNoob() function and methylation values (beta values) were obtained using
the getBeta() function from minfi package (v.1.42.0)[85–87]. The model matrix was created for the two
temperatures using the model.matrix() function from stats R package and sva() function from sva
package (v.3.44.0) was applied with two estimated surrogate variables[81,88]. The model was fit with
contrast matrix using lmFit(), with the estimated coefficients and standard errors generated using
contrasts.fit() from limma package (v.3.52.4)[85,89]. The p-values were moderated using eBayes()
function from limma package (v.3.52.4) and CpG cites ranked according to their p-value[89].

CUT&RUN: 293WT and human NPCs (ATCC, cat. ACS-1019) were harvested and CUT&RUN performed
according to Epicypher CUTANA protocol (v.1.7) on 300.000 cells per sample. We permeabilized the
cells with 0.01% digitonin (Sigma, D141). For normalization we spiked in E. Coli DNA (Epicypher,
23618-1401) at the final concentration of 0.2 ng per sample. The library preparation was performed
with TrueSeq-Chip Sample preparation Kit (Illumina, IP-202-9001). The following antibodies were
used for the CUT&RUN: CTCF antibody (Cell Signaling, 2899S), H3K36me3 (Thermo Fisher Scientific,
MA5-24687), H4K20me3 (EpiCypher, 13-0054), H3K9me3 (Abcam, ab176916), H3K4me3 (EpiCypher,
13-0041), H3K27me3 (Thermo Fisher Scientific, MA5-11198), SMYD5 [51], and Rabbit IgG Antibody
(Epicypher, 23613-0042). For the CUT&RUN analysis and the CUT&TAG re-analysis we followed
published STAR protocols[90]. Briefly, first we trimmed the raw Fastq reads in paired mode with
TrimGalore (v.0.6.7)[91] then we used Bowtie2 (v.2.4.4.)[92] to align the reads to Hg38 or mm10 and E.
Coli reference genome, all downloaded from UCSC. Overall alignment to Hg38 were from 83.6-
98.63% and to E. Coli spike in were from 0.1-4.18% for our data. The aligned Sam files were then
converted to Bam files, indexed, and sorted via SAMtools (v.1.15)[93] and BedGraphtoBigWig (v.4)[94]
from BEDTools (v.2.3)[95]. We used SEACR (v.1.3)[96] for peak calling after normalization with the
normalization factor calculated as 1/(percent alignment to E. coli). We used DiffBind (v.3.9.6)[97] for
the differential peak analysis, ChIPseeker (v.1.35.3)[98], TxDb.Hsapiens.UCSC.hg38.knownGene
(v.3.4.6)[99] and org.Hs.eg.db (v.3.8.2)[100] for annotation. For further analysis and image creation we
used deepTools (v.3.5.1.)[101] and R [81].

Mouse NPC isolation: Mouse NPCs were isolated from the cortex of E17.5 C57BL6/NTac embryos. The protocol was adapted from Bernas et al, 2017[102]. Embryo cortices were manually dissected from brains and the tissue was dissociated in 1X TrypLE™ Select Enzyme (Thermo Fisher Scientific, A1217701) for 10 min, with manual dissociation using a 1000 µL pipette. The cell suspension was washed in Neurobasal medium (Thermo Fisher Scientific, 21103049) and filtered through a 70 µm cell strainer (Miltenyi Biotech, 130-110-916). Cells were washed 2X in 5 mL Neurobasal medium (Gibco, 21103049) by centrifugation at 200 xg for 10 min. Cells were resuspended in 1 mL Neurobasal growth medium containing 1X B27 supplement (Thermo Fisher Scientific, 17504044), 1X Penicillin/Streptomycin (Thermo Fisher Scientific, 15140122), 1X Glutamax (Thermo Fisher Scientific, 35050038), 20ng/mL FGF-2 (Peprotech, 100-18B), 20 ng/mL EFG (Peprotech, AF-100-15), and 2 µg/mL Heparin (MP Biomedicals, 210193125). Cells were seeded onto 12-well plates, previously coated for 2 h at 37°C in 1:100 dilution of Matrigel (Corning, 354234). After passaging, NPCs were cultured in media described above on plates coated overnight in poly-D-lysine (Sigma, P7280), and then coated for 2 h in laminin (Sigma, L2020). Cells were either sub-cultured or media was changed every 2 days.

RNAseq: Primary mouse NPC lines from 4 separate embryos were seeded 0.3 x 10^6 cells per well, in two batches, in a poly-D-lysine/laminin coated 6-well plate after 9 days *in vitro*. Following 36 h of incubation at 37°C, one plate with 4 lines was incubated at 32°C for 6 h. Cells from both 37°C and 32°C conditions were harvested in tri-reagent and RNA was isolated using the Direct-zol RNA Microprep kit (Zymo Research, R2063). RNAseq library construction and Illumina next generation sequencing was performed by Novogene, generating 30 million paired-end reads per sample. For the RNAseq analysis, Fastq files were pseudo-aligned to the GRCm39 mouse transcriptome, downloaded from NCBI, using Kallisto (v.0.48.0) with paired-end mode on and 100 bootstraps. Kallisto output files were imported into R using the tximport package[103]. Transcripts were assigned to genes using the TxDb.Mmusculus.UCSC.mm10.ensGene package (v.3.4.0)[104] in R . Differential expression analysis was performed using the DESeq2 package (v.3.16)[105] in R after discarding transcripts with less than 10 counts. Differentially expressed genes were defined as those with an adjusted p-value less than 0.1. Overlaps of gene lists were calculated using the GeneOverlap package (v.1.34.0)[106] in R, with overlaps tested using Fisher's exact test. Gene set enrichment analysis was performed using the GSEA-preranked program[107,108] with the Mouse Ortholog Hallmark Gene Sets[109]. For the GSEA-pre-ranked run, we generated a ranked gene list comprising all detected genes assigned to a rank value, which was calculated as -log10(adjusted p-value) × (sign of fold change). The genes were then ranked from the highest rank value to the lowest before inputting into the program.

Mouse cooling followed by immunostaining on brain slices: C57BL6 mice (Jackson) were bred to create a litter (n=2). At P10, we split littermates into two groups. One group (n=6) was cooled to a core temperature of 32°C and the other (n=6) maintained at 37°C, for 6 hours respectively. These animals were then euthanized, and their brains were isolated immediately after the cooling period and flashed with PBS. One of the hemispheres was immersed in 4% PFA for 72-96 h for fixation and the other hemisphere was dissected for the cerebellum, thalamus/basal ganglia, hippocampus, anterior and posterior cortex. After fixation, samples were cryopreserved in a sucrose gradient and then flash-frozen. Frozen hemispheres were cryosectioned with the help of a cryostat (Keldur, UI). The sections were washed in TBS+ (TBS (LiCor, 927-60001) with 0.05% triton X-100(Sigma-Aldrich, T8787-100ML)) twice for 10 minutes. Next, the slides were immersed in DAKO (target retrieval solution, sodium citrate buffer, pH 6) at 95°C for 20 minutes. Slides were then washed again in TBS+

twice for 10 minutes. Next, the slides were blocked in TBS++ (TBS (LiCor, 927-60001) with 3% goat

serum (Abcam, ab7481)) for 1 h at room temperature. The slides were then stained with primary

antibodies and anti-rabbit-SMYD5, 1:200, for 24 h at room temperature and carried out in the dark.

On the next day, slides were allowed to reach room temperature before they are washed 3x for 10

minutes in TBS+. The slides were incubated with secondary antibodies, goat anti-mouse IgG (H+L)

Alexa Fluor™ 488, 1:1000 (Invitrogen™, A-11001); goat anti-rabbit IgG (H+L) Alexa Fluor™ 647, 1:1000

(Invitrogen™, A21244), for 24 h in the dark and washed 3x for 10 minutes in TBS+. All the liquid on

the slides was briefly dried off. Fluoromount-G Mounting Medium with DAPI (Invitrogen, 00-4959-52)

was placed on each section and covered with a cover slip. The slides were incubated for 24 h at room

temperature before being imaged using confocal microscopy (FV 1200, Olympus Fluoroview) using a

30x silicon objective (NA1.05). Images were processed by ImageJ[82,83] and quantified using CellProfiler

(v.4.2.5.)[84] . Statistics were calculated in GraphPad Prism.

Supplementary Tables and Files

List of Supplementary Tables

Supplementary Table 1: Specifics of MHIs (word file).

Supplementary Table 2: Top 91 drug hits from MHI screen (excel file).

Supplementary Table 3: Top candidate genes found enriched in mutation screening (excel file).

Supplementary Table 4: Genes found in both repressor and activator SP1 lists and thus excluded for
further evaluation (excel file).

Supplementary Table 5: Top differentially modified locations for H3K36me3 in NPCs from CUT&RUN
experiment (excel file).

Supplementary Table 6: List of CpGs represented at three genes (word document).

Supplementary Table 7: List of differentially methylated locations from EPIC DNAm array (excel file).
Supplementary Table 8: List of genes that have a unique H3K4me3 histone modification peak at 32°C
(excel file).

Supplementary Table 9: List of genes that show DEGS at 32°C and have a unique H3K4me3 histone
modification peak at 32°C (excel file).

Supplementary files

Supplementary File 1: Sequences of individual MHIs.

Supplementary figures

Figure S1. Different temperature ranges. Temperatures known to induce distinct responses
(moderate and mild hypothermia, euthermia and hyperthermia). Our range of interest (mild
hypothermia) is bolded.

Figure S2. MHI activation peaks during mild hypothermia (32°C) and is limited at other temperatures. (A-C) We measured fluorescence hourly using an IncucyteS3 at 32°C using SP1-Lenti-MHI, CIRBP-Lenti-MHI, RBM3-Lenti-MHI respectively. (D-E) No increased fluorescence was observed at moderate hypothermia for either SP1-MHI or CIRBP-MHI in HEK293 cells. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S3. Storage of samples on ice does not lead to activation of MHI. (A-B) gMFI or percentage of

fluorescent cells of HEK293WT+Cas9+SP1 (293WT cell line expressing SP1-MHI) left on ice for

different periods of time before flow cytometry analysis. Shown are data points, each data point is a

technical replicate, mean and SD where applicable. Dunnett's multiple comparison test calculated in

GraphPad Prism. (C-D) Mean fluorescence or percentage of fluorescent cells of HEK293WT+Cas9+SP1

left on ice for different periods of time before flow cytometry analysis. Data shown as in (A),

independent replicate. * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S4. MHI demonstrate consistent activation at 32°C in diverse cellular systems. (A-E) Left
column shows mean fluorescence (MF, flow cytometry) after transient transfection of SP1-MHI
into HCT116, HeLa, Jurkat, K562 and SK-N-SH (16 h at 32°C, 37°C and 40°C). Shown are data
points, each data point is a technical replicate, mean and SD where applicable, Šidák's multiple
comparison test calculated in GraphPad Prism. (F-J) Middle column shows gMFI (flow cytometry)
after transient transfection of RBM3-MHI into HCT116, HeLa, Jurkat, K562 and SK-N-SH (16 h at
32°C and 37°C). Data shown as in (A-E). (K-O) Right column shows gMFI (flow cytometry) after
transient transfection of SP1+MCRE-MHI into HCT116, HeLa, Jurkat, K562 and SK-N-SH (last 16 h
at 32°C and 37°C). Data as in (A-E). * = P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S5. Moderate hyperthermia does not activate SP1 MHI in diverse systems. (A-E) Mean
fluorescence of SP1-MHI as measured by flow cytometry after transient transfection of MHI into
HCT116, HeLa, Jurkat, K562, SK-N-SH (16 h at 26°C, 29°C and 37°C). Shown are data points, each
data point is a technical replicate, mean and SD where applicable. Šidák's multiple comparison. *
= P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

853
854

Figure S6: Individual validation of potential hits from drug screen. (A-F) gMFI measured with flow cytometry of HEK293WT cell line that stably expresses MHI after 16 h of Cinacalcet (A), Dasatinib (B), Erythromycin (C), Naloxone (D), Orantinib (E), and Tamoxifen (F) at 32°C or 37°C.

Shown are data points, mean and SD where applicable. Šidák's multiple comparison calculated with GraphPad Prism. (G) Cell cycle analysis done with PI staining. Mean fluorescence measured for G1, S and G2 phases of the cell cycle with flow cytometry of HEK293WT cell line 16 h after Entacapone or Poziotinib exposure at 32°C or 37°C. Data shown as in (A-F). (H) Apoptosis analysis done with H_2O_2 exposure. Shown are data points, mean and SD where applicable. Unpaired t test calculated with Graphpad Prism. * = $P<0.05$, ** = $P<0.01$, *** = $P<0.001$, **** = $P<0.0001$.

Figure S7: Expression of endogenous loci after exposure to Entacapone and Poziotinib. Relative
expression of *SP1* (top), *CIRBP* (middle), *RBM3* (bottom) when compared to GAPDH after 16 h at 32°C
or 37°C with and without Entacapone (left) or Poziotinib (right). Shown are data points, each data
point is a biological replicate (mean of technical replicates), mean and SD where applicable.
Significance levels were calculated with unpaired one tailed Student t test in GraphPad Prism. * =
P<0.05, ** = P<0.01, *** = P<0.001, **** = P<0.0001.

Figure S8: *RGS21* and *MALT1* knockout did not show any significant fluorescent change. sgRNA KO
for *MALT1* and *RGS21* did not show any changes in fluorescence of HEK293 cells with SP1-MHI,
suggesting that *MALT1* and *RGS21* do not have an impact on SP1-MHI. Shown are data points, each
data point is a technical replicate, mean and SD where applicable. Significance levels were calculated
with Šidák's multiple comparison in GraphPad Prism. * = P<0.05, ** = P<0.01, *** = P<0.001, **** =
P<0.0001.

Figure S9. Pathway analysis using genes from forward mutagenesis. (A) Schematic depiction of

repressors identified from both screens which land on the Neuregulin pathway. (B-C) Unbiased

MSigDB pathway enrichment of highly ranked (FDR<0.25 and -Log10(RRA) >0.5) genes from the SP1

activator and SP1 repressor lists, respectively.

Figure S10. *Cirbp* has fewer normalized counts compared to *Sp1* and *Rbm3*. Normalized counts from
RNAseq in mESc (referring to data from **Figure 4C**).

Figure S11. Validation of temperature dependence of SMYD5. (A) An independent biological
replicate with anti-SMYD5 antibody showing striking differences between two temperatures. Green
or white (SMYD5), purple (DAPI). (B) A validation with anti-FLAG antibody, showing same difference.
Green (SMYD5), purple (DAPI).

Figure S12. DNAm status at SP1, CIRBP and RBM3 promoters. (A) DNAm status of CpG dinucleotides
from Illumina EPIC array 5 kb upstream and 2 kb downstream of *SP1* gene. (B) DNAm status of CpG
dinucleotides from Illumina EPIC array 5 kb upstream and 2 kb downstream of *CIRBP* gene. (C) DNAm
status of CpG dinucleotides from Illumina EPIC array 5 kb upstream and 2 kb downstream of *RBM3*
gene.

Figure S13. Methylation modification levels in NPCs and HEK293 cells. (A) H3K36me3 levels at 32°C and 37°C at *SP1* promoter in HEK293 cells. (B) Global H3K36me3 levels (all genes) in HEK293 cells. (C-D) Global H3K27me3, H4K20me3 levels, respectively (all genes) in NPCs (E) Similarly, at the gene-specific levels there is a number of genes, that have more or less H3K36me3 at 32°C compared to 37°C in HEK293 cells. Red circles indicate genes that have Log$_2$ fold change over 1 and p-value less than 10^{-6}, gray circles indicate Log$_2$ fold change over 1 but non-significant p-value and black circles indicate non-significant genes. (F-G) H3K36me3 levels at SMYD5-bound genes identified to be DEGS at the two temperatures in NPC and HEK293 cells, respectively. (H-J) H3K9me3 and H3K4me3, respectively, levels at SMYD5-bound genes identified to be DEGS at the two temperatures in NPCs.

Figure S14: H3K36me3 levels on representative peaks showing differential binding at 32°C when compared to 37°C. (A-C) Increased H3K36me3 is observed for *BAZ2A, RIMOC1 and USP1I,* respectively, at 32°C compared to 37°C. (D-E) Decreased H3K36me3 is observed for *CENPE*, *SETD1B* and *YEATS2I,* respectively, at 37°C compared to 32°C.

References

1.   Osilla, E. V., Marsidi, J.L., and Sharma, S. (2022). Physiology, Temperature Regulation
(StatPearls Publishing).

2.   Turbill, C., Bieber, C., and Ruf, T. (2011). Hibernation is associated with increased survival and
the evolution of slow life histories among mammals. Proceedings of the Royal Society B:
Biological Sciences *278*, 3355–3363. 10.1098/RSPB.2011.0190.

3.   Mathew, J.L., Kaur, N., and Dsouza, J.M. (2022). Therapeutic hypothermia in neonatal hypoxic
encephalopathy: A systematic review and meta-analysis. J Glob Health *12*.
10.7189/JOGH.12.04030.

4.   Luedke, M.W., Graffagnino, C., McKinney, B.G., Piper, J., Iversen, E., and Kolls, B. (2022).
Association of time-temperature curves with outcomes in temperature management for
cardiac arrest. BMJ Neurol Open *4*. 10.1136/BMJNO-2022-000273.

5.   Dietrichs, E.S., and Dietrichs, E. (2015). Neuroprotective effects of hypothermia. Tidsskr Nor
Laegeforen *135*, 1646–1651. 10.4045/tidsskr.14.1250.

6.   Sun, Y.J., Zhang, Z.Y., Fan, B., and Li, G.Y. (2019). Neuroprotection by Therapeutic
Hypothermia. Front Neurosci *13*. 10.3389/FNINS.2019.00586.

7.   Fujita, J. (1999). Cold shock response in mammalian cells. J Mol Microbiol Biotechnol *1*, 243–
255.

8.   Zhu, X., Bührer, C., and Wellmann, S. (2016). Cold-inducible proteins CIRP and RBM3, a unique
couple with activities far beyond the cold. Cellular and Molecular Life Sciences, 1–21.
10.1007/s00018-016-2253-7.

9.   Ryu, H., Lee, J., Olofsson, B.A., Mwidau, A., Deodoglu, A., Escudero, M., Flemington, E.,
Azizkhan-Clifford, J., Ferrante, R.J., and Ratan, R.R. (2003). Histone deacetylase inhibitors
prevent oxidative neuronal death independent of expanded polyglutamine repeats via an Sp1-
dependent pathway. Proc Natl Acad Sci U S A *100*, 4281–4286. 10.1073/PNAS.0737363100.

10. Yang, H.J., Zhuang, R.J., Li, Y.B., Li, T., Yuan, X., Lei, B.B., Xie, Y.F., and Wang, M. (2019). Cold-
inducible protein RBM3 mediates hypothermic neuroprotection against neurotoxin rotenone
via inhibition on MAPK signalling. J Cell Mol Med *23*, 7010–7020. 10.1111/JCMM.14588.

11. Bastide, A., Peretti, D., Knight, J.R.P., Grosso, S., Spriggs, R. V., Pichon, X., Sbarrato, T., Roobol,
A., Roobol, J., Vito, D., et al. (2017). RTN3 Is a Novel Cold-Induced Protein and Mediates
Neuroprotective Effects of RBM3. Current Biology *27*, 638. 10.1016/J.CUB.2017.01.047.

12. Chip, S., Zelmer, A., Ogunshola, O.O., Felderhoff-Mueser, U., Nitsch, C., Bührer, C., and
Wellmann, S. (2011). The RNA-binding protein RBM3 is involved in hypothermia induced
neuroprotection. Neurobiol Dis *43*, 388–396. 10.1016/j.nbd.2011.04.010.

13. Peretti, D., Bastide, A., Radford, H., Verity, N., Molloy, C., Martin, M.G., Moreno, J.A., Steinert,
J.R., Smith, T., Dinsdale, D., et al. (2015). RBM3 mediates structural plasticity and protective

effects of cooling in neurodegeneration. Nature 2015 518:7538 *518*, 236–239.
10.1038/nature14142.

14. Wierstra, I. (2008). Sp1: Emerging roles—Beyond constitutive activation of TATA-less
housekeeping genes. Biochem Biophys Res Commun *372*, 1–13. 10.1016/J.BBRC.2008.03.074.

15. Sumitomo, Y., Higashitsuji, H., Higashitsuji, H., Liu, Y., Fujita, T., Sakurai, T., Candeias, M.M.,
Itoh, K., Chiba, T., and Fujita, J. (2012). Identification of a novel enhancer that binds Sp1 and
contributes to induction of cold-inducible RNA-binding protein (cirp) expression in
mammalian cells. BMC Biotechnol *12*, 72. 10.1186/1472-6750-12-72.

16. Gomez-Pastor, R., Burchfiel, E.T., and Thiele, D.J. (2018). Regulation of heat shock
transcription factors and their roles in physiology and disease. Nat Rev Mol Cell Biol *19*, 4–19.
10.1038/NRM.2017.73.

17. Sanjana, N.E., Shalem, O., and Zhang, F. Improved vectors and genome - wide libraries for
CRISPR screening.

18. Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelson, T., Heckl, D., Ebert, B.L.,
Root, D.E., Doench, J.G., et al. (2014). Genome-scale CRISPR-Cas9 knockout screening in
human cells. Science *343*, 84–87. 10.1126/science.1247005.

19. Joung, J., Konermann, S., Gootenberg, J.S., Abudayyeh, O.O., Platt, R.J., Brigham, M.D.,
Sanjana, N.E., and Zhang, F. (2017). Genome-scale CRISPR-Cas9 knockout and transcriptional
activation screening. Nat Protoc *12*, 828–863. 10.1038/nprot.2017.016.

20. McIntyre, S., Nelson, K.B., Mulkey, S.B., Lechpammer, M., Molloy, E., and Badawi, N. (2021).
Neonatal encephalopathy: Focus on epidemiology and underexplored aspects of etiology.
Semin Fetal Neonatal Med *26*. 10.1016/J.SINY.2021.101265.

21. Geneva, I.I., Cuzzo, B., Fazili, T., and Javaid, W. (2019). Normal Body Temperature: A
Systematic Review. Open Forum Infect Dis *6*. 10.1093/OFID/OFZ032.

22. Knight, J.R.P., Bastide, A., Roobol, A., Roobol, J., Jackson, T.J., Utami, W., Barrett, D.A., Smales,
C.M., and Willis, A.E. (2015). Eukaryotic elongation factor 2 kinase regulates the cold stress
response by slowing translation elongation. Biochem J *465*, 227–238. 10.1042/BJ20141014.

23. Shaw, G., Morse, S., Ararat, M., and Graham, F.L. (2002). Preferential transformation of
human neuronal cells by human adenoviruses and the origin of HEK 293 cells. The FASEB
Journal. 10.1096/fj.01-0995fje.

24. Barnes, E.N., Biedler, J.L., Spengler, B.A., and Lyser, K.M. (1981). The fine structure of
continuous human neuroblastoma lines SK-N-SH, SK-N-BE(2), and SK-N-MC. In Vitro *17*, 619–
631. 10.1007/BF02618461.

25. Stepanenko, A.A., and Dmitrenko, V. V (2015). HEK293 in cell biology and cancer research:
phenotype, karyotype, tumorigenicity, and stress-induced genome-phenotype evolution.
Gene *569*, 182–190. 10.1016/j.gene.2015.05.065.

26.   Jackson, T.C., Manole, M.D., Kotermanski, S.E., Jackson, E.K., Clark, R.S.B., and Kochanek, P.M.
1001   (2015). Cold stress protein RBM3 responds to temperature change in an ultra-sensitive
1002   manner in young neurons. Neuroscience *305*, 268–278. 10.1016/j.neuroscience.2015.08.012.

1003   27.   Hartmann, J., Haap, M., Kopp, H.-G., and Lipp, H.-P. (2009). Tyrosine kinase inhibitors - a
1004   review on pharmacology, metabolism and side effects. Curr Drug Metab *10*, 470–481.
1005   10.2174/138920009788897975.

1006   28.   Nelson, V., Ziehr, J., Agulnik, M., and Johnson, M. (2013). Afatinib: emerging next-generation
1007   tyrosine kinase inhibitor for NSCLC. Onco Targets Ther *6*, 135–143. 10.2147/OTT.S23165.

1008   29.   Kim, T.M., Lee, K.W., Oh, D.Y., Lee, J.S., Im, S.A., Kim, D.W., Han, S.W., Kim, Y.J., Kim, T.Y., Kim,
1009   J.H., et al. (2018). Phase 1 Studies of Poziotinib, an Irreversible Pan-HER Tyrosine Kinase
1010   Inhibitor in Patients with Advanced Solid Tumors. Cancer Res Treat *50*, 835–842.
1011   10.4143/CRT.2017.303.

1012   30.   Erber, R., Thurnher, A., Katsen, A.D., Groth, G., Kerger, H., Hammes, H.P., Menger, M.D.,
1013   Ullrich, A., and Vajkoczy, P. (2004). Combined inhibition of VEGF and PDGF signaling enforces
1014   tumor vessel regression by interfering with pericyte-mediated endothelial cell survival
1015   mechanisms. FASEB J *18*, 338–340. 10.1096/FJ.03-0271FJE.

1016   31.   Lee, H., Kim, J.W., Choi, D.K., Yu, J.H., Kim, J.H., Lee, D.S., and Min, S.H. (2020). Poziotinib
1017   suppresses ovarian cancer stem cell growth via inhibition of HER4-mediated STAT5 pathway.
1018   Biochem Biophys Res Commun *526*, 158–164. 10.1016/J.BBRC.2020.03.046.

1019   32.   Grinstein, E., Jundt, F., Weinert, I., Wernet, P., and Royer, H.D. (2002). Sp1 as G1 cell cycle
1020   phase specific transcription factor in epithelial cells. Oncogene 2002 21:10 *21*, 1485–1492.
1021   10.1038/sj.onc.1205211.

1022   33.   Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S., Brown, M., and Liu,
1023   X.S. (2014). MAGeCK enables robust identification of essential genes from genome-scale
1024   CRISPR/Cas9 knockout screens. Genome Biol *15*, 554. 10.1186/s13059-014-0554-4.

1025   34.   Hsu, P.D., Lander, E.S., and Zhang, F. (2014). Development and Applications of CRISPR-Cas9 for
1026   Genome Engineering. Cell *157*, 1262–1278. 10.1016/j.cell.2014.05.010.

1027   35.   Nagai, G., and Nakaoka, Y. (1998). Cooling sensitive [Ca2+](i) response associated with
1028   signaling of G protein-coupled receptors. Biochem Biophys Res Commun *248*, 733–737.
1029   10.1006/BBRC.1998.8873.

1030   36.   Chelban, V., Patel, N., Vandrovcova, J., Zanetti, M.N., Lynch, D.S., Ryten, M., Botía, J.A., Bello,
1031   O., Tribollet, E., Efthymiou, S., et al. (2017). Mutations in NKX6-2 Cause Progressive Spastic
1032   Ataxia and Hypomyelination. Am J Hum Genet *100*, 969–977. 10.1016/J.AJHG.2017.05.009.

1033   37.   ZNF260 zinc finger protein 260 [Homo sapiens (human)] - Gene - NCBI
1034   https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=339324#gene-
1035   expression.

38. Burke, T.L., and Grant, P.A. (2009). Histone Acetylation Complexes. Handbook of Cell
Signaling, Second Edition *3*, 2369–2378. 10.1016/B978-0-12-374145-5.00285-0.

39. Zhou, X., Shang, Y.N., Lu, R., Fan, C.W., and Mo, X.M. (2019). High ANKZF1 expression is
associated with poor overall survival and recurrence-free survival in colon cancer.
https://doi.org/10.2217/fon-2018-0920 *15*, 2093–2106. 10.2217/FON-2018-0920.

40. Tawamie, H., Martianov, I., Wohlfahrt, N., Buchert, R., Mengus, G., Uebe, S., Janiri, L., Hirsch,
F.W., Schumacher, J., Ferrazzi, F., et al. (2017). Hypomorphic Pathogenic Variants in TAF13 Are
Associated with Autosomal-Recessive Intellectual Disability and Microcephaly. Am J Hum
Genet *100*, 555. 10.1016/J.AJHG.2017.01.032.

41. ZNF71 zinc finger protein 71 [Homo sapiens (human)] - Gene - NCBI
https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=58491.

42. Yu, H. V., Tao, L., Llamas, J., Wang, X., Nguyen, J.D., Trecek, T., and Segil, N. (2021). POU4F3
pioneer activity enables ATOH1 to drive diverse mechanoreceptor differentiation through a
feed-forward epigenetic mechanism. Proc Natl Acad Sci U S A *118*.
10.1073/PNAS.2105137118/-/DCSUPPLEMENTAL.

43. GATAD2A GATA zinc finger domain containing 2A [Homo sapiens (human)] - Gene - NCBI
https://www.ncbi.nlm.nih.gov/gene/54815.

44. Zhang, Y., Fang, Y., Tang, Y., Han, S., Jia, J., Wan, X., Chen, J., Yuan, Y., Zhao, B., and Fang, D.
(2022). SMYD5 catalyzes histone H3 lysine 36 trimethylation at promoters. Nature
Communications 2022 13:1 *13*, 1–19. 10.1038/s41467-022-30940-1.

45. Kidder, B.L., He, R., Wangsa, D., Padilla-Nash, H.M., Bernardo, M.M., Sheng, S., Ried, T., and
Zhao, K. (2017). SMYD5 Controls Heterochromatin and Chromosome Integrity during
Embryonic Stem Cell Differentiation. Cancer Res *77*, 6729–6745. 10.1158/0008-5472.CAN-17-
0828.

46. Murphy, M.W., Sarver, A.L., Rice, D., Hatzi, K., Ye, K., Melnick, A., Heckert, L.L., Zarkower, D.,
and Bardwell, V.J. (2010). Genome-wide analysis of DNA binding and transcriptional
regulation by the mammalian Doublesex homolog DMRT1 in the juvenile testis. Proc Natl
Acad Sci U S A *107*, 13360–13365. 10.1073/PNAS.1006243107/SUPPL_FILE/SD01.XLS.

47. Weber, C., Zhou, Y., Lee, J.G., Looger, L.L., Qian, G., Ge, C., and Capel, B. (2020). Temperature-
dependent sex determination is mediated by pSTAT3 repression of Kdm6b. Science (1979)
368, 303–306. 10.1126/SCIENCE.AAZ4165/SUPPL_FILE/AAZ4165_S1.MP4.

48. Kim, D.H., Doyle, M.R., Sung, S., and Amasino, R.M. (2009). Vernalization: winter and the
timing of flowering in plants. Annu Rev Cell Dev Biol *25*, 277–299.
10.1146/annurev.cellbio.042308.113411.

49. Fellous, A., Earley, R.L., and Silvestre, F. (2019). The Kdm/Kmt gene families in the self-
fertilizing mangrove rivulus fish, Kryptolebias marmoratus, suggest involvement of histone
methylation machinery in development and reproduction. Gene *687*, 173–187.
10.1016/J.GENE.2018.11.046.

50.   Youn, J.Y., Dyakov, B.J.A., Zhang, J., Knight, J.D.R., Vernon, R.M., Forman-Kay, J.D., and
Gingras, A.C. (2019). Properties of Stress Granule and P-Body Proteomes. Mol Cell *76*, 286–
294. 10.1016/J.MOLCEL.2019.09.014.

51.   Aljazi, M.B., Gao, Y., Wu, Y., and He, J. (2022). SMYD5 is a histone H3-specific
methyltransferase mediating mono-methylation of histone H3 lysine 36 and 37. Biochem
Biophys Res Commun *599*, 142–147. 10.1016/J.BBRC.2022.02.043.

52.   Diaz, J., Abiola, S., Kim, N., Avaritt, O., Flock, D., Yu, J., Northington, F.J., and Chavez-Valdez, R.
(2017). Therapeutic hypothermia provides variable protection against behavioral deficits after
neonatal HI: A potential role for BDNF. Dev Neurosci *39*, 257. 10.1159/000454949.

53.   Xi, Y., Park, S.R., Kim, D.H., Kim, E.D., and Sung, S. (2020). Transcriptome and epigenome
analyses of vernalization in Arabidopsis thaliana. Plant J *103*, 1490. 10.1111/TPJ.14817.

54.   Fan, L., Lesser, A.F., Sweet, D.R., Keerthy, K.S., Lu, Y., Chan, E.R., Vinayachandran, V., Ilkayeva,
O., Das, T., Newgard, C.B., et al. (2022). KLF15 controls brown adipose tissue transcriptional
flexibility and metabolism in response to various energetic demands. iScience *25*, 105292.
10.1016/J.ISCI.2022.105292.

55.   Katano-Toki, A., Satoh, T., Tomaru, T., Yoshino, S., Ishizuka, T., Ishii, S., Ozawa, A., Shibusawa,
N., Tsuchiya, T., Saito, T., et al. (2013). THRAP3 Interacts with HELZ2 and Plays a Novel Role in
Adipocyte Differentiation. Molecular Endocrinology *27*, 769. 10.1210/ME.2012-1332.

56.   THRAP3 thyroid hormone receptor associated protein 3 [Homo sapiens (human)] - Gene -
NCBI https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=9967.

57.   Guo, K., Searfoss, G., Krolikowski, D., Pagnoni, M., Franks, C., Clark, K., Yu, K.T., Jaye, M., and
Ivashchenko, Y. (2001). Hypoxia induces the expression of the pro-apoptotic gene BNIP3. Cell
Death Differ *8*, 367–376. 10.1038/SJ.CDD.4400810.

58.   Abdel-Wahab, O., Adli, M., LaFave, L.M., Gao, J., Hricik, T., Shih, A.H., Pandey, S., Patel, J.P.,
Chung, Y.R., Koche, R., et al. (2012). ASXL1 mutations promote myeloid transformation
through loss of PRC2-mediated gene repression. Cancer Cell *22*, 180–193.
10.1016/J.CCR.2012.06.032.

59.   Kaakkola, S. (2000). Clinical pharmacology, therapeutic use and potential of COMT inhibitors
in Parkinson's disease. Drugs *59*, 1233–1250. 10.2165/00003495-200059060-00004.

60.   Feng, Z., Saha, L., Dritsa, C., Wan, Q., and Glebov, O.O. (2022). Temperature-dependent
structural plasticity of hippocampal synapses. Front Cell Neurosci *16*, 520.
10.3389/FNCEL.2022.1009970/BIBTEX.

61.   Carvelli, L., Blakely, R.D., and DeFelice, L.J. (2008). Dopamine transporter/syntaxin 1A
interactions regulate transporter channel activity and dopaminergic synaptic transmission.
Proc Natl Acad Sci U S A *105*, 14192–14197. 10.1073/PNAS.0802214105.

62.   Gao, H., Chakraborty, G., Lee-Lim, A.P., Mavrakis, K.J., Wendel, H.G., and Giancotti, F.G.
(2014). Forward genetic screens in mice uncover mediators and suppressors of metastatic

1111  reactivation. Proc Natl Acad Sci U S A *111*, 16532–16537.
1112  10.1073/PNAS.1403234111/SUPPL_FILE/PNAS.201403234SI.PDF.

1113 63. INDACOCHEA, A., GUERRERO, S., UREÑA, M., ARAUJO, F., COLL, O., LLEONART, M.E., and
1114  GEBAUER, F. (2021). Cold-inducible RNA binding protein promotes breast cancer cell
1115  malignancy by regulating Cystatin C levels. RNA *27*, 190–201. 10.1261/RNA.076422.120.

1116 64. Nicolas, M., Noe, V., Jensen, K., and Ciudad CJ (2001). Cloning and characterization of the 5'-
1117  flanking region of the human transcription factor Sp1 gene. . J Biol Chem, 22126-22132.

1118 65. Yang, F., Eckardt, S., Leu, N.A., McLaughlin, K.J., and Wang, P.J. (2008). Mouse TEX15 is
1119  essential for DNA double-strand break repair and chromosomal synapsis during male meiosis.
1120  J Cell Biol *180*, 673. 10.1083/JCB.200709057.

1121 66. TEX15 testis expressed 15, meiosis and synapsis associated [Homo sapiens (human)] - Gene -
1122  NCBI https://www.ncbi.nlm.nih.gov/gene/56154.

1123 67. Hanna, C.W., Huang, J., Belton, C., Reinhardt, S., Dahl, A., Andrews, S., Francis Stewart, A.,
1124  Kranz, A., and Kelsey, G. (2022). Loss of histone methyltransferase SETD1B in oogenesis
1125  results in the redistribution of genomic histone 3 lysine 4 trimethylation. Nucleic Acids Res *50*,
1126  1993. 10.1093/NAR/GKAC051.

1127 68. Pajoro, A., Severing, E., Angenent, G.C., and Immink, R.G.H. (2017). Histone H3 lysine 36
1128  methylation affects temperature-induced alternative splicing and flowering in plants. Genome
1129  Biol *18*, 1–12. 10.1186/S13059-017-1235-X/FIGURES/4.

1130 69. Michaels, K.K., Mostafa, S.M., Capella, J.R., and Moore, C.L. (2020). Regulation of alternative
1131  polyadenylation in the yeast Saccharomyces cerevisiae by histone H3K4 and H3K36
1132  methyltransferases. Nucleic Acids Res *48*, 5407–5425. 10.1093/NAR/GKAA292.

1133 70. Xu, S.J., Lombroso, S.I., Fischer, D.K., Carpenter, M.D., Marchione, D.M., Hamilton, P.J., Lim,
1134  C.J., Neve, R.L., Garcia, B.A., Wimmer, M.E., et al. (2021). Chromatin-mediated alternative
1135  splicing regulates cocaine-reward behavior. Neuron *109*, 2943-2966.e8.
1136  10.1016/J.NEURON.2021.08.008.

1137 71. Preußner, M., Goldammer, G., Neumann, A., Haltenhof, T., Rautenstrauch, P., Müller-
1138  McNicoll, M., and Heyd, F. (2017). Body Temperature Cycles Control Rhythmic Alternative
1139  Splicing in Mammals. Mol Cell *67*, 433-446.e4. 10.1016/J.MOLCEL.2017.06.006.

1140 72. Horii, Y., Shiina, T., Uehara, S., Nomura, K., Shimaoka, H., Horii, K., and Shimizu, Y. (2019).
1141  Hypothermia induces changes in the alternative splicing pattern of cold-inducible RNA-binding
1142  protein transcripts in a non-hibernator, the mouse. Biomed Res *40*, 153–161.
1143  10.2220/BIOMEDRES.40.153.

1144 73. Sano, Y., Shiina, T., Naitou, K., Nakamori, H., and Shimizu, Y. (2015). Hibernation-specific
1145  alternative splicing of the mRNA encoding cold-inducible RNA-binding protein in the hearts of
1146  hamsters. Biochem Biophys Res Commun *462*, 322–325. 10.1016/j.bbrc.2015.04.135.

74. Smart, F., Aschrafi, A., Atkins, A., Owens, G.C., Pilotte, J., Cunningham, B.A., and Vanderklish,
P.W. (2007). Two isoforms of the cold-inducible mRNA-binding protein RBM3 localize to
dendrites and promote translation. J Neurochem *101*, 1367–1379. 10.1111/J.1471-
4159.2007.04521.X.

75. Al-Fageeh, M.B., and Smales, C.M. (2013). Alternative promoters regulate cold inducible RNA-
binding (CIRP) gene expression and enhance transgene expression in mammalian cells. Mol
Biotechnol *54*, 238–249. 10.1007/s12033-013-9649-5.

76. Nicolas, M., Noe, V., Jensen, K.B., and Ciudad, C.J. (2001). Cloning and characterization of the
5'-flanking region of the human transcription factor Sp1 gene. J Biol Chem *276*, 22126–22132.
10.1074/jbc.M010740200.

77. Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., Studer, L., and
Author, N.B. (2009). Highly efficient neural conversion of human ES and iPS cells by dual
inhibition of SMAD signaling HHS Public Access Author manuscript. Nat Biotechnol *27*, 275–
280. 10.1038/nbt.1529.

78. Xiang, J., Wan, C., Guo, R., and Guo, D. (2016). Is Hydrogen Peroxide a Suitable Apoptosis
Inducer for All Cell Types? Biomed Res Int *2016*. 10.1155/2016/7343965.

79. Kutner, R.H., Zhang, X.Y., and Reiser, J. (2009). Production, concentration and titration of
pseudotyped HIV-1-based lentiviral vectors. Nat Protoc *4*, 495–505. 10.1038/nprot.2009.22.

80. Wang, B., Wang, M., Zhang, W., Xiao, T., Chen, C.H., Wu, A., Wu, F., Traugh, N., Wang, X., Li,
Z., et al. (2019). Integrative analysis of pooled CRISPR genetic screens using MAGeCKFlute. Nat
Protoc *14*, 756–780. 10.1038/s41596-018-0113-7.

81. Statistical R Core Team (2017). R Foundation for Statistical Computing Vienna Austria. R: A
language and environment for computing.

82. Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and Eliceiri,
K.W. (2017). ImageJ2: ImageJ for the next generation of scientific image data. BMC
Bioinformatics *18*, 1–26. 10.1186/S12859-017-1934-Z/FIGURES/7.

83. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nature Methods 2012 9:7 *9*, 676–682. 10.1038/nmeth.2019.

84. Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and Goodman, A.
(2021). CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics *22*.
10.1186/S12859-021-04344-9.

85. Fortin, J.P., Triche, T.J., and Hansen, K.D. (2017). Preprocessing, normalization and integration
of the Illumina HumanMethylationEPIC array with minfi. Bioinformatics *33*, 558.
10.1093/BIOINFORMATICS/BTW691.

86.   Triche, T.J., Weisenberger, D.J., Van Den Berg, D., Laird, P.W., and Siegmund, K.D. (2013). Low-
level processing of Illumina Infinium DNA Methylation BeadArrays. Nucleic Acids Res *41*, e90.
10.1093/NAR/GKT090.

87.   Aryee, M.J., Jaffe, A.E., Corrada-Bravo, H., Ladd-Acosta, C., Feinberg, A.P., Hansen, K.D., and
Irizarry, R.A. (2014). Minfi: a flexible and comprehensive Bioconductor package for the
analysis of Infinium DNA methylation microarrays. Bioinformatics *30*, 1363.
10.1093/BIOINFORMATICS/BTU049.

88.   Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E., and Storey, J.D. (2012). The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics *28*, 882. 10.1093/BIOINFORMATICS/BTS034.

89.   Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res *43*, e47. 10.1093/NAR/GKV007.

90.   Kong, N.R., Chai, L., Tenen, D.G., and Bassal, M.A. (2021). A modified CUT&RUN protocol and
analysis pipeline to identify transcription factor binding sites in human cell lines. STAR Protoc
2. 10.1016/J.XPRO.2021.100750.

91.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet J *17*, 10–12. 10.14806/EJ.17.1.200.

92.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359. 10.1038/nmeth.1923.

93.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and
Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*,
2078–2079. 10.1093/BIOINFORMATICS/BTP352.

94.   Luo, Y., Hitz, B.C., Gabdank, I., Hilton, J.A., Kagda, M.S., Lam, B., Myers, Z., Sud, P., Jou, J., Lin,
K., et al. (2020). New developments on the Encyclopedia of DNA Elements (ENCODE) data
portal. Nucleic Acids Res *48*, D882–D889. 10.1093/NAR/GKZ1062.

95.   Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842. 10.1093/BIOINFORMATICS/BTQ033.

96.   Meers, M.P., Tenenbaum, D., and Henikoff, S. (2019). Peak calling by Sparse Enrichment
Analysis for CUT&RUN chromatin profiling. Epigenetics Chromatin *12*, 1–11. 10.1186/S13072-
019-0287-4/FIGURES/6.

97.   Stark, R., and Brown, G. (2012). DiffBind : differential binding analysis of ChIP-Seq peak data.

98.   Yu, G., Wang, L.G., and He, Q.Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. Bioinformatics *31*, 2382–2383.
10.1093/BIOINFORMATICS/BTV145.

99.   Team BC, and Maintainer BP (2019). TxDb.Hsapiens.UCSC.hg38.knownGene: Annotation
package for TxDb object(s).

100.   Carlson M (2019). org.Hs.eg.db: Genome wide annotation for Human. .

101.   Ramírez, F., Dündar, F., Diehl, S., Grüning, B.A., and Manke, T. (2014). deepTools: a flexible
1221   platform for exploring deep-sequencing data. Nucleic Acids Res *42*, W187.
1222   10.1093/NAR/GKU365.

1223   102.   Bernas, S.N., Leiter, O., Walker, T.L., and Kempermann, G. (2017). Isolation, Culture and
1224   Differentiation of Adult Hippocampal Precursor Cells. Bio Protoc *7*.
1225   10.21769/BIOPROTOC.2603.

1226   103.   Soneson, C., Love, M.I., and Robinson, M.D. (2016). Differential analyses for RNA-seq:
1227   Transcript-level estimates improve gene-level inferences. F1000Res *4*.
1228   10.12688/F1000RESEARCH.7563.2/DOI.

1229   104.   Team BC, and Maintainer BP (2016). TxDb.Mmusculus.UCSC.mm10.ensGene: Annotation
1230   package for TxDb object(s)..

1231   105.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
1232   dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 1–21. 10.1186/S13059-014-0550-
1233   8/FIGURES/9.

1234   106.   Shen L, and Sinai ISoMaM (2022). GeneOverlap: Test and visualize gene overlaps. .

1235   107.   Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
1236   Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment
1237   analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc
1238   Natl Acad Sci U S A *102*, 15545–15550. 10.1073/PNAS.0506580102.

1239   108.   Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver,
1240   P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1alpha-responsive genes
1241   involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat
1242   Genet *34*, 267–273. 10.1038/NG1180.

1243   109.   Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015).
1244   The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst *1*, 417–
1245   425. 10.1016/J.CELS.2015.12.004.

1246

1247