Circadian PERIOD proteins regulate TC-DSB repair through anchoring to the nuclear envelope.

Benjamin Le Bozec[1], Laure Guitton-Sert[1], Sarah Collins[1], Anne-Laure Finoux[1], Charlotte Payrault[1], Jessica Frison[1], Romane Carette[1], Vincent Rocher[1], Emmanuelle Guillou[1], Coline Arnould[1], Aude Guénolé[1], Marion Aguirrebengoa[2], Ikrame Lazar[1], Aline Marnef[1], Philippe Frit[3,4], Patrick Calsou[3,4], Thomas Mangeat[5], Nadine Puget[1*] and Gaëlle Legube[1,6*]

1. *MCD, Centre de Biologie Intégrative (CBI), CNRS, Université de Toulouse, UT3, 31062 Toulouse, France*

2. *BigA, NGS data analysis facility, Centre de Biologie Intégrative (CBI), CNRS, Université de Toulouse, UT3, 31062 Toulouse, France*

3. *Institut de Pharmacologie et de Biologie Structurale (IPBS), Université de Toulouse, CNRS, Université Toulouse III - Paul Sabatier (UT3), Toulouse, France*

4. *Equipe Labellisée Ligue Contre le Cancer 2018, Toulouse, France*

5. *LITC Core Facility, Centre de Biologie Intégrative (CBI), CNRS, Université de Toulouse, UT3, 31062 Toulouse, France*

6. *Lead author*

* Corresponding authors: gaelle.legube@univ-tlse3.fr, nadine.puget1@univ-tlse3.fr

Running title: Circadian regulation of TC-DSB repair

Abstract

Repair of DNA Double-Strand Breaks (DSBs) produced in transcriptionally active chromatin occurs through a poorly characterized pathway called Transcription-Coupled DSB repair (TC-DSBR). Here, using a screening approach scoring multiple outputs in human cells, we identified proteins from the PERIOD complex, a key module ensuring circadian oscillations, as novel TC-DSBR players. We show that the core PER complex protein PER2 is recruited at TC-DSBs and that it contributes to the targeting of TC-DSBs at the nuclear envelope (NE). At the NE, SUN1 and the Nuclear Pore Complex (NPC) act as docking sites for TC-DSBs and TC-DSB anchoring fosters RAD51 assembly. Impaired DSB localization at the NE results in elevated DSB clustering and translocation rate. In agreement, the circadian clock regulates TC-DSB anchoring to the NE, RAD51 assembly, and DSB clustering. Our study shows that DSB localization to the NPC is a conserved molecular pathway that also occurs in human cells and provides a direct link between the circadian rhythm and the response to DSBs occurring in active genes. This opens new therapeutic strategies for chemotherapies based on drugs that are inducing DSBs in active loci such as topoisomerase poisons.

Main

DNA Double-Strand breaks (DSBs) are among the most toxic lesions that occur on the genome, given their potential to elicit large chromosomal rearrangements. While global genome DSB repair (GG-DSBR) mechanisms have been well characterized, a transcription-coupled arm of DSB repair (TC-DSBR) has only recently been identified[1]. Yet, physiological DSBs mostly appear in transcriptionally active loci (TC-DSBs)[2–5] and topoisomerase II inhibitors, classically used for cancer treatment, mainly induce TC-DSBs[2,4] emphasizing the need to better understand TC-DSBR mechanisms. TC-DSBR entails the ATM- and DNA-PK-dependent transcriptional repression of damaged genes which is necessary for the initiation of resection[6,7]. TC-DSBs are preferentially repaired by Homologous Recombination (HR) in G2[8,9], while in G1, they are refractory to repair and cluster in large subnuclear structures[10] that we recently identified as being a novel, DSB-induced, chromatin compartment (the D-compartment)[11] whose formation enhances translocation frequency and checkpoint activation[11,12]. Additionally, when occurring on human ribosomal DNA, TC-DSBs are physically relocated at the nucleolar periphery where they contact invaginations of the Nuclear Envelope (NE)[13]. Whether localization to the NE also occurs when DSBs arise within RNAPII-transcribed loci is yet unknown, although NE targeting of other types of persistent DSBs and collapsed replication forks have been reported in *Drosophila* and *S. Cerevisiae*[14–19]. Altogether the TC-DSBR pathway is still poorly understood.

Multi-output screen to identify new TC-DSBR factors

In search of novel TC-DSBR players, we performed a siRNA library screen scoring multiple outputs in the well-characterized human DIvA cell line (Fig. 1a). In this cell line, annotated DSBs are induced throughout the genome in a temporally controlled manner, thanks to a restriction enzyme fused to the ligand binding domain of the estrogen receptor (AsiSI-

ER)[20]. BLESS experiments (enabling mapping and quantification of DSBs throughout the

genome using streptavidin-capture and sequencing of fragments adjacent to DSBs ligated to

biotinylated linkers[21]) allowed to identify 174 sites robustly cleaved *in vivo* out of the 1211

AsiSI sites annotated on the human genome (mainly due to the fact that AsiSI activity is

inhibited by DNA methylation)[20,22]. A large fraction of these DSBs occur in RNAPII-enriched

(transcribed) loci hereafter called TC-DSBs[20,22]. The siRNA library comprised 130 candidate

proteins previously identified to be associated with the SMC1 subunit of the cohesin complex,

which was previously shown to be recruited at DSBs post-DSB induction[13]. The screen was

designed to quantify the effect of candidate protein depletion on 1) γH2AX foci intensity using

high-content imaging, 2) cell survival using a cell proliferation assay, 3) chromosome

rearrangement frequency using quantitative PCR and 4) DSB clustering using high content

microscopy (Fig. 1a, Extended Data Fig. 1a). A positive control siRNA against SETX, an R-

loop resolving factor previously reported to be involved in TC-DSBR[23,24], triggered an expected

increase in γH2AX foci intensity, translocation frequency and DSB clustering as well as

decreased cell proliferation (Fig. 1b red label, Extended Data Fig. 1b-e). We further validated

our screening dataset with the candidate protein Periphilin (PPHLN1, Fig. 1b green label,

Extended Data Fig. 1b-e), an H3K9me3 reader of the HUSH complex that we previously

reported to be involved in the repair of DSB occurring on ribosomal DNA[13]. In agreement with

the screen results (Fig. 1b), depletion of PPHLN1 triggered an increase in γH2AX foci intensity

(Extended Data Fig. 1f-g). We further used a degradable version of AsiSI-ER (AID-tagged

DIvA cell line[8]) to measure clonogenic potential and repair post-DSB induction. PPHLN1

depletion increased two distinct translocation events and cell death, in agreement with our

screen results (Extended Data Fig. 1h-i).

Importantly, two-parameter correlations established that γH2AX intensity was inversely

correlated with cell survival but positively correlated with translocation frequency (Fig. 1d-e),

consistent with a decreased repair capacity respectively impairing cell proliferation and

promoting genome rearrangements. Moreover, increased DSB clustering positively correlated

with translocation frequency (Fig. 1f), and increased translocation frequency inversely

correlated with cell survival (Fig. 1g) in agreement with previous work[11,12]. Altogether, results

from our siRNA screen establish that TC-DSBR deficiency severely affects cell survival and

further confirm the relationship between DSB clustering and genome rearrangements as

recently reported[12,25].

The circadian clock PERIOD proteins contribute to TC-DSB repair.

Hierarchical clustering identified two groups displaying statistically significant

differences in clustering, translocation, γH2AX and cell survival (Fig. 1b-c). Interestingly,

among the candidate proteins whose depletion gave similar phenotypes to SETX (*i.e.*, increased

γH2AX staining, decreased cell survival, increased chromosomal rearrangement and increased

DSB clustering, Group 1, Fig. 1b-c, Extended Data Fig. 1b-e), we found NONO, SFPQ and

DDX5 which all belong to the same complex, the mammalian PERIOD (PER) complex

involved in the regulation of the circadian rhythm[26–29]. Similar outputs were also seen for

another candidate, the DDX5-paralog DDX17, both DDX5 and DDX17 being human orthologs

of the *Neurospora* PRD1 protein involved in circadian rythmicity[30] and existing as a

heterodimer[31]. The circadian clock is molecularly controlled by the BMAL1/CLOCK

transcriptional activator complex, which regulates the transcription of thousands of genes.

Among those, the PER-CRY transcriptional repressor can downregulate BMAL1/CLOCK

activity, creating a negative transcriptional feedback loop that ensures circadian oscillations[32].

In order to directly investigate whether the core clock components of the PER complex play a

role in TC-DSBR, we depleted PER1 and PER2 proteins (Extended Data Fig. 2a). As found for

DDX17, NONO, DDX5 and SETX (Fig. 1b; Extended Data Fig. 1b), PER1 and PER2

knockdown increased γH2AX foci intensity following DSB induction in DIvA cells (Fig. 2a)

with no major changes in cell cycle distribution (Extended Data Fig. 2b). In agreement, PER2-

depleted cells also displayed elevated γH2AX levels around DSBs as detected by chromatin

immunoprecipitation (ChIP) (Fig. 2b). In order to validate this finding in another, non-

cancerous, cell line, we further depleted PER1 and PER2, alone or in combination, in hTERT

RPE-1 cells (RPE1) stably infected with a mAID-AsiSI-ER lentiviral construct (RPE-DIvA).

PER1 and/or PER2 depleted cells displayed elevated γH2AX levels post DSB induction by

AsiSI (Extended Data Fig. 2c-d). Moreover, γH2AX increase following PER1/2 depletion also

hold true when inducing DSB using etoposide, a Topoisomerase 2 (TOP2) inhibitor known to

mainly trigger DSBs in transcribed loci and regulatory elements[2,4] (Extended Data Fig. 2e). Of

note in RPE1 cells, PER1/2 depletion also significantly increased γH2AX levels even before

DSB induction using AsiSI or etoposide (Extended Data Fig. 2c-e), suggesting that PER1/2

contribute to endogenous DSB repair.

As found for DDX17, NONO, SETX and DDX5 (Fig. 1b; Fig. 2c-d; Extended Data Fig. 1c-d;

Extended Data Fig. 2g-i), PER1 and PER2 depletion increased translocation frequency (Fig.

2c) and impaired cell survival following induction of TC-DSBs in DIvA cells (Fig. 2d;

Extended Data Fig. 2f). Concomitant depletion of PER1 and PER2 triggered exacerbated

translocation frequency and cell death following DSB induction, as compared to PER1 or PER2

depletion alone (Extended Data Fig. 2h-i). PER1/2 depletion also impaired cell survival upon

treatment with the TOP2 inhibitors etoposide and doxorubicin in U2OS and RPE1 (Extended

Data Fig. 2j-k).

In contrast and surprisingly, depletion of CRY2, known to function in the PER complex, did

not result in an elevated translocation rate nor increased cell death (Extended Data Fig. 2h-i).

Similarly, depletion of neither BMAL1 nor CLOCK transcriptional activators, alone or in

combination, altered the translocation frequency or cell survival post DSB induction (Fig. 2c-

d, Extended Data Fig. 2h-i). Altogether, these data show that the PERIOD proteins PER1 and

PER2, as well as some other members of the PER complex (NONO, DDX17, SETX and

DDX5), contribute to the response to DSBs induced in transcribed loci.

In order to determine whether the function of the PER complex is direct, we assessed the

recruitment of PER complex proteins at DSBs using ChIP. PER2, DDX17 and DDX5 showed

significant recruitment at a TC-DSB induced in DIvA cells (already observed for DDX5[33]),

while this was not the case for BMAL1 (Fig. 2e). We further performed ChIP-seq against PER2

and BMAL1 in damaged and undamaged DIvA cells. Both proteins displayed the expected

pattern on the genome (see examples Extended Data Fig. 3a). Indeed, in absence of DSB

induction, BMAL1 was enriched on genomic loci previously identified by BMAL1 ChIP-seq[34]

(Extended Data Fig. 3b) and PER2 accumulated at promoters (Extended Data Fig. 3c) as

expected for a transcriptional repressor and from previous studies[35]. Moreover, BMAL1-bound

genes displayed significant enrichment for the "rhythmic process" GO term (Extended Data

Fig. 3d). Post-DSB induction, we interestingly observed that PER2 accumulated at TC-DSBs

(see an example Fig. 2f top panel, purple track) but not at a silent locus (Fig. 2f bottom panel),

despite equivalent cleavage (BLESS tracks). In contrast, BMAL1 showed no accumulation at

DSBs (Fig. 2f, blue tracks). On average, PER2 was significantly targeted on ~2kb around DSBs,

while BMAL1 was not (Fig. 2g, Extended Data Fig. 3e). In agreement with our above data

showing that PER1/2 depletion triggers increased γH2AX even before DSB induction

(Extended Data Fig. 2c-e), we also observed a significant enrichment of PER2 at previously

determined endogenous DSB hotspots[11] (Extended Data Fig. 3f). We previously identified a

subset of equivalently cleaved DSBs in DIvA cells with two distinct repair behaviors: the Non-

Homologous End-Joining (NHEJ)-prone DSBs and the HR-prone DSBs with the latter having

a tendency to locate within transcriptionally active loci (*i.e.* TC-DSBs)[8,22]. Importantly, PER2

preferentially accumulates at HR-prone TC-DSBs compared to NHEJ-prone DSBs (*i.e.*

occurring in non-transcribed regions) (Fig. 2h). Altogether these data show that the PER

complex is recruited at HR-repaired, TC-DSBs, suggesting a direct role of the PER complex in

TC-DSB repair.

168 TC-DSBs are targeted to the nuclear envelope via a PER2 -dependent mechanism.

Of interest, the nuclear envelope (NE) contributes to the regulation of the circadian clock[36] and

PER proteins have been involved in the physical targeting of clock-regulated genes to the NE

in *Drosophila*[37]. On another hand, persistent DSBs[14,17,38–40], sub-telomeric DSBs[41,42],

dysfunctional telomeres[43,44] and arrested replication forks[18,19] have been previously found to be

relocated at the NE in yeast as were stressed replication forks in mammals[45] and

heterochromatic (HC) DSBs in *Drosophila* in order to complete HR[15,46]. While in mammalian

cells, physical relocation of DSBs to the NE has not yet been documented, we previously

observed that ribosomal DNA DSBs can contact NE invaginations inside the nucleoplasm[13].

Hence, we set out to determine whether TC-DSBs could be targeted to the NE. As a first

approach, we performed fixed and live super-resolution imaging using Random Illumination

Microscopy (RIM)[47]. We observed that γH2AX foci can establish close contacts with LaminB1

filaments (Extended data Fig. 4a). Live imaging after 1 hour of DSB induction in DIvA cells

expressing 53BP1-GFP and mCherry-LaminB1, showed that such relocalization of DSBs to the

nuclear lamina were fast (<60s) and transient (Fig. 3a and Supplementary Videos 1-4). Contacts

between repair foci and the nuclear lamina were further confirmed by Proximity Ligation Assay

(PLA) performed between LaminB1 and 53BP1 (Fig. 3b) or between γH2AX and LaminB1

(Extended Data Fig. 4b). DSBs induced by etoposide in U2OS cells also displayed interactions

with LaminB1 (Extended data Fig. 4c). Moreover, we could recapitulate DSB/LaminB1 interactions in RPE1 primary cells subjected to AsiSI or etoposide induced DSBs (Extended data Fig. 4d-e). In agreement, ChIP against LaminB1 (Extended Data Fig. 4f) also revealed increased LaminB1 occupancy at TC-DSBs in contrast to a DSB induced in a silent locus and repaired by NHEJ, in DIvA cells post-DSB induction (Fig. 3c). Taken together, these data suggest that TC-DSBs can be physically relocated at the NE in human cells.

In order to evaluate the function of PER2 in targeting TC-DSB at the NE, we performed LaminB1 ChIP in DIvA cells upon PER2 siRNA. Of interest PER2 depletion abolished LaminB1 recruitment at TC-DSBs (Fig. 3d). Additionally, PER1 or PER2-depleted cells also displayed decreased 53BP1/LMNB1 PLA signal post-DSB induction when compared to PER1/2-proficient cells (siRNA CTRL) (Fig. 3e, Extended Data Fig. 4g). Similarly, PER1/2 depletion also decreased γH2AX/LMNB1 PLA signal in response to etoposide (Extended Data Fig. 4h). In order to investigate a potential cell cycle dependency for PER2-mediated DSB anchoring to the NE, we then monitored γH2AX/LaminB1 PLA signal combined with EdU incorporation by quantitative high throughput microscopy in DIvA cells (Extended Data Fig. 4i). PER2 depletion decreased PLA foci in G1, S and G2 cells (Extended Data Fig. 4j). Collectively, these data suggest that PER2 contributes to TC-DSB targeting at the NE.

Tethering of TC-DSBs to the NE depends on SUN1 and NUP153

Interestingly, we and others previously reported a function for SUN domain containing-proteins, which are components of the Inner Nuclear Membrane (INM), in DSB mobility in mammals[10,48]. Moreover, SUN protein orthologs in yeast and *Drosophila* (respectively Mps3, and Koi/Spag4) were also found to act as DSB anchoring points in the NE[15–17]. We thus investigated the potential involvement of the two main SUN proteins in mammals, SUN1 and

SUN2, in anchoring TC-DSBs to the NE. Interestingly, ChIP against SUN proteins (Extended Data Fig. 5a) revealed that, upon damage, SUN1 displayed enrichment at TC-DSBs compared to NHEJ-DSBs (Extended Data Fig. 5b, top panel). Surprisingly, SUN2 rather showed decreased occupancy post-DSB induction (Extended Data Fig. 5b, bottom panel), suggesting that TC-DSBs are specifically interacting with SUN1 and not SUN2. Of note SUN1 recruitment at TC-DSB was independent of the activity of the three main DNA Damage Response (DDR) kinases, namely ATM, ATR and DNAPK (Extended Data Fig. 5c). To further investigate SUN protein recruitment at DSBs, we performed ChIP-seq against SUN1 and SUN2 before and after DSB induction in DIvA cells. Visual inspection and peak calling on SUN1 and SUN2 ChIP-seq datasets revealed that in undamaged conditions, both proteins are enriched on genes (Extended Data Fig. 5d-e), accumulated at promoters (Extended Data Fig. 5f) and largely overlapped on the genome (Extended Data Fig. 5g). Of interest, post-DSB induction, SUN1 (purple track), was recruited on approximately 1kb around the break, while SUN2 rather displayed eviction (blue track) (see an example Fig. 3f top panel, and average profiles at all DSBs Fig. 3g). Both SUN1 recruitment and SUN2 eviction at DSBs were statistically significant when compared to random, undamaged genomic positions (Extended Data Fig. 5h). Notably, the profile of SUN1 at DSBs strongly resembled to that of PER2 (Extended Data Fig. 5i), and as PER2, SUN1 also showed enrichment at endogenous DSB hotspots (Extended Data Fig. 5j). Moreover, as observed for PER2, SUN1 was specifically recruited at a TC-DSB (Fig. 3f, top panel) but not at a DSB induced in a silent locus (Fig. 3f, bottom panel). On average, SUN1 displayed enhanced accumulation at DSBs with high RNAPII levels compared to those with low RNAPII levels (Extended data Fig. 5k). In agreement, as for PER2, HR-prone DSBs showed higher level of SUN1 as compared to NHEJ-prone DSBs (Extended Data Fig. 5l left panel) while SUN2 depletion was exacerbated at HR-prone DSBs compared to NHEJ-prone DSBs (Extended Data Fig. 5l right panel).

We then performed 3D-super resolution imaging by RIM to detect SUN1 and SUN2. Both

proteins localized to the NE and displayed punctuated patterns (Extended Data Fig. 6a). Co-

staining with γH2AX enabled the detection of a significant colocalization (ICQ) with SUN1,

but not with SUN2 (Extended Data Fig. 6b). Van Steensel Cross-correlation function (CCF)

analysis showed a non-random overlap between γH2AX foci and SUN1 while non-random

exclusion was observed between γH2AX foci and SUN2 (see examples Fig. 3h and

quantification Fig. 3i). In agreement with a function of SUN1 in tethering TC-DSB to the NE,

depletion of SUN1 (Extended Data Fig. 6c) abolished Lamin B1 enrichment at TC-DSBs (Fig.

3j), and also decreased PLA signal between LaminB1 and γH2AX following etoposide

treatment (Extended Data Fig. 6d). Altogether, these data suggest that TC-DSBs are physically

targeted to the NE through an interaction with SUN1 and not SUN2.

Interestingly, SUN1 has been reported to interact with the Nuclear Pore Complex (NPC), as

opposed to SUN2[49,50]. In agreement, in DIvA cells, we found that SUN1 exhibits significant

colocalization with NUP153, a protein part of the NPC basket, as compared to SUN2 (Extended

Data Fig. 6e). Given that the NPC has also been identified as a docking site for persistent DSBs

and collapsed replication forks in yeast[14,18,19,38,51] and heterochromatic DSBs in drosophila[15] we

thus postulated that the specificity of SUN1 as opposed to SUN2 to anchor DSBs at the NE

may be related to its association to the NPC. Indeed, NUP153 accumulates at TC-DSBs but not

at a DSB induced in an inactive locus (Extended Data Fig. 6f). 3D-super resolution RIM and

CCF analysis further indicated that, as SUN1, NUP153 displays significant colocalization with

γH2AX foci (Fig. 3k, examples on the left panel and quantification on the right panel). Of

importance, depletion of NUP153 (Extended Data Fig. 6c) triggered a decreased association of

TC-DSB with the NE, as measured by PLA between γH2AX and LaminB1 (Extended Data Fig.

6g), further confirming a function of NUP153 in TC-DSB anchoring. Overall, our data suggest

that TC-DSBs are tethered to the NE through a SUN1 and NPC-dependent mechanism.

PER2 and SUN1-dependent DSB anchoring at the NPC promotes HR

Given that PER2, SUN1 and NUP153 are recruited at HR-prone DSBs (Fig. 2h, Extended Data Fig. 5l, Extended Data Fig. 6f) and that previous work showed that targeting DSBs and collapsed replication forks to the NPC promotes HR in yeast and drosophila[14,15,17–19,38,40,46], we further investigated whether this pathway contributes to HR-repair at TC-DSBs in human cells. One of the first steps of TC-DSB repair is the transcriptional repression of nearby genes (including the damaged gene itself) which is essential for the proper execution of resection and HR repair[1,52]. Given that the PER complex is a transcriptional repressor, we envisaged that it could promote HR repair at TC-DSBs via a role in transcriptional repression. However, RT-qPCR analyses revealed that both PER2 and PER1 are dispensable for transcriptional repression following DSB induction (Fig. 4a). Similarly, SUN1 depletion did not modify the ability of TC-DSB to trigger transcriptional repression (Fig. 4b) showing that the PER2/SUN1-dependent TC-DSB targeting pathway to the NE is not required for the transcriptional inactivation of genes *in cis* to DSBs. We further investigated the effect of SUN1 and PER1 or PER2 depletion on end-resection, by quantification of single-stranded DNA at DSBs[53] (Extended Data Fig. 6h). siRNA depletion of SUN1 did not alter resection (Extended Data Fig. 6i) while both PER1 and PER2 triggered a mild decrease of resection (Extended Data Fig. 6j), although a lot less pronounced than upon CtIP depletion. Altogether, these data suggest that the contribution of NE anchoring in HR repair takes place downstream of transcriptional repression and resection.

In drosophila, the targeting of HC breaks to the NE favors RAD51 loading[15,46]. We thus further assayed in DIvA cells, whether PER2-dependent TC-DSB targeting at the NE regulates RAD51 recruitment. Indeed, RAD51 loading at TC-DSBs was significantly reduced in PER2-depleted cells (Fig. 4c). Of interest, in yeast, the eviction of the single strand DNA binding protein RPA,

required to ensure Rad51 nucleofilament formation, is promoted by its ubiquitination by the

Sumo-Targeted Ubiquitin Ligase (STUbL) Slx5/8[19], which is also required to relocate collapsed

forks and DSBs to the NE[15,18,19,38,40,46]. We thus further tested RPA and RAD51 recruitment at

TC-DSBs upon depletion of SUN1 and RNF4, the Slx5/8 ortholog in human cells which has

also been previously reported to be involved in HR repair[54,55]. SUN1 and RNF4 depletion

(Extended Data Fig. 6c, Extended Data Fig. 6k) altered the balance between RAD51 and RPA

at TC-DSB (Fig. 4d, Extended Data Fig. 6l).

In agreement with our above findings, depletion of SUN1, NUP153 and RNF4 all impaired cell

survival post TC-DSB induction, while SUN2 did not, suggesting deficient repair upon

impairment of the TC-DSB NE-anchoring pathway (Fig. 4e-g). SUN1 and NUP153 depletion

also impaired cell survival following etoposide in RPE1 cells (Extended Data. Fig. 6m).

Importantly co-depletion of both SUN1 and PER1 or PER2 did not exacerbate the impaired

survival observed upon depletion of PER1 or PER2 proteins independently (Fig. 4h). Similarly,

co-depletion of NUP153 and PER2 or SUN1 did not trigger a significant decrease of cell

survival as compared to NUP153 depletion alone (Fig. 4i), nor did co-depletion of RNF4 and

PER2 (Fig. 4j), indicating they belong to the same pathway.

Taken together, our data suggest that TC-DSB anchoring to the NE mediated by a PER2, SUN1,

NUP153 and RNF4-dependent pathway, fosters RAD51 assembly, thereby ensuring efficient

repair and cell survival post DSB induction.

PER2-mediated DSB anchoring to the NE counteracts DSB clustering and D-

compartment formation.

In mammalian nuclei, principal component analysis (PCA) of Hi-C experiments revealed that

chromatin compartmentalizes to form the so-called "A" and "B" compartments, with the B-

compartment corresponding to microscopically visible heterochromatin foci, while the A-

compartment correspond to euchromatin[56]. Of interest, both experimental data and

computational modeling of chromosome behaviour showed that targeting at the nuclear lamina

prevents heterochromatin to coalesce into internal larger nuclear bodies[57]. The current model

postulates that upon detachment of chromatin from the NE, phase separation allows multiple

HC foci to cluster thus increasing B-compartmentalization. Importantly, we recently reported

that upon DSB induction, another chromatin compartment forms, so called the D-compartment

(for DSB-induced), through polymer-polymer phase separation[11]. This D-compartment that

arises via DSB clustering gathers γH2AX-modified topologically associated domains (TADs)

as well as additional undamaged loci, including a subclass of DSB-activated genes[11].

Of interest, our siRNA screen indicated that the depletion of NONO, SETX and DDX17

increased clustering (Fig. 1b, Extended data Fig. 1f) and we found that PER2 contributes in

DSBs targeting to the NE (Fig. 3). Hence, we set out to investigate whether PER2 depletion

could also increase DSB clustering and D-compartment formation as a result of a lack of

damaged TADs anchoring at the NE. To address this point, we performed Hi-C experiments

before and after DSB induction in PER2-proficient and deficient cells. PER2 depletion did not

drastically affect chromosome organization in undamaged cells (Extended Data Fig. 7a), nor

the overall distribution of A and B compartments (Extended Data Fig. 7b). However, PER2

depletion triggered increased compartmentalization as visualized by increased A-A and B-B

interaction, as well as decreased A-B interaction (Extended Data Fig. 7c). This agrees with the

previously reported role of the PER complex in chromatin anchoring at the NE which would

predict an increased compartmentalization upon NE detachment in PER-depleted cells. Upon

DSB induction, we could recapitulate our previous finding showing DSB clustering (Extended

Data Fig. 7d, arrows). Importantly, PER2 depletion increased DSB clustering as shown on

individual events (Fig. 5a, Extended Data Fig. 7d), average aggregate peak analysis (APA) plots

of inter-chromosomal or intra-chromosomal DSB-DSB contacts centered on all 80 best-induced

DSBs (Fig. 5b, Extended Data Fig. 7e) and box plots quantifying the interactions between DSBs

(Fig. 5c). PER2 depletion also triggered enhanced clustering of endogenous DSB hotspots

(Extended Data Fig. 7f) and of etoposide-induced DSBs as measured using quantitative high

throughput microscopy (Extended data Fig. 7g).

We further computed Chromosomal Eigen Vectors (CEV) using the first PC of Principal

Component Analysis (PCA) on differential Hi-C matrices to identify the D-compartment as

previously described[11]. As expected, we could recapitulate D-compartment formation upon

damage in DIvA cells transfected with a control siRNA (Extended Data Fig. 7h). PER2

deficiency did not alter D-compartment detection on chromosomes previously found to display

D-compartment (Extended Data Fig. 7h). However, we could observe D-compartment

formation on additional chromosomes upon PER2 depletion (Fig. 5d). Collectively our data

suggest that PER proteins prevent DSB clustering and D-compartment formation by fostering

the targeting of TC-DSBs to the nuclear lamina.

D-compartment formation fosters the DNA damage response, but it also comes at the expense

of increased translocation rate[12,25,58]. In agreement with the above data involving PER2 in

counteracting D-compartment formation, increased translocation frequency was observed upon

depletion of the PER complex proteins (Fig. 2c). Similarly, and in line with their involvement

in TC-DSB tethering at NE (Extended data Fig. 6d, Fig. 3g, Extended data Fig. 6g), SUN1 and

NUP153 depletion increased translocation (Fig. 5e) suggesting enhanced DSB clustering in

SUN1 and NUP153 deficient cells. Of note, this was not the case when depleting SUN2, in

agreement with the absence of SUN2 recruitment at DSBs. Moreover, in agreement with the

previously reported role of Slx5/8 in the relocation of persistent DSBs and arrested forks in

yeast[14,19,40,51], as well as of HC breaks in drosophila[15], RNF4 depletion also increased

translocation frequency (Fig. 5e), suggesting enhanced DSB clustering in absence of the STUbL

RNF4. Importantly co-depletion of both SUN1 and PER1 or PER2 did not further increased

translocation compared to SUN1 depletion alone (Fig. 5f). Similarly, co-depletion of NUP153

and PER2 or SUN1 did not trigger a significant increase in translocation frequency as compared

to NUP153 depletion alone (Fig. 5g), nor did co-depletion of RNF4 and PER2 (Fig. 5h). Taken

together, our data suggest that TC-DSB anchoring to the NE mediated by a PER2, SUN1,

NUP153 and RNF4-dependent pathway, counteracts DSB clustering and translocations in

human cells.

The circadian clock regulates the response to TC-DSBs and their anchoring to the NE.

Given that the PER complex is a core component of the circadian clock, we set out to investigate

whether the circadian clock could also regulate the response to TC-DSBs. As expected,

dexamethasone treatment synchronized the circadian clock in both U2OS and DIvA cells, as

assessed by the cycling expression of PER2, BMAL1 and CRY2 (Extended Data Fig. 8a-b).

We further monitored γH2AX foci formation in different cell cycle phases, at different time

points after dexamethasone-mediated circadian clock synchronization by combining γH2AX

staining and EdU labeling, using quantitative high throughput microscopy in DIvA cells. In line

with the results obtained with the depletion of PER2 by siRNA, our data indicate that γH2AX

foci intensity was increased post-DSB induction during the low-PER2 circadian phase (Fig. 6a,

white bars; Extended Data Fig. 8c) as compared to the high-PER2 phase (Fig. 6a blue bars;

Extended Data Fig. 8c). This was independent of the cell cycle phase (Fig. 6a). Moreover,

increased accumulation of γH2AX at TC-DSBs during the low-PER2 phase, was further

confirmed by ChIP (Fig. 6b). Of importance the same holds true in S and G2 phases when DSBs

were induced using etoposide (Extended Data Fig. 8d). To further reinforce the notion that the

low-PER2 phase of the circadian clock impacts TC-DSB repair, we performed circadian clock

synchronization and analyzed several outputs that were affected by the depletion of PER2. First,

the circadian clock did not alter transcriptional repression in *cis* to DSBs (Extended Data Fig.

8e) as expected from PER2 depletion data (Fig. 4a). Second, as found using PER2 depletion

(Fig. 3d), the low-PER2 circadian phase reduced the recruitment of LaminB1 (Fig. 6c) and

SUN1 (Fig. 6d) to TC-DSBs when monitored by ChIP-qPCR, suggesting that DSB anchoring

at the NE is defective when the PER1/2 proteins are expressed at low levels. Indeed,

γH2AX/LaminB1 PLA signal decreased during the low PER2 phase (Fig. 6e). Third, given that

our data indicated a role of PER2 in RAD51 loading at TC-DSBs (Fig.4c), we assessed RAD51

recruitment using ChIP. As expected from PER2 depletion data, RAD51 recruitment at TC-

DSBs decreased during the low PER phase (Fig. 6f). In agreement, RAD51 foci formation

monitored using high throughput microscopy at different time points after dexamethasone

treatment revealed an impaired RAD51 foci formation during the low-PER2 phase (Extended

Data Fig. 8f). Fourth, quantitative high throughput microscopy further showed an increase in

γH2AX foci area during the low-PER2 phase, in agreement with enhanced DSB clustering (Fig.

6g).

We previously reported that, in addition to γH2AX domains, the D-compartment also physically

attracts additional loci including some, but not all, DNA damage responsive (DDR) genes, in

order to potentiate their activation post-DSB induction[25]. Notably, we found that *RNF19B* and

PLK3, two upregulated DDR genes found within the D-compartment, were significantly more

activated post-DSB during the low-PER2 phase when compared to the high-PER2 phase (Fig.

6h). This agrees with an increased D-compartment formation in absence of PER2 (Fig. 5a-d,

Extended Data Fig. 7h). Of importance, this was not the case for other DDR genes which are

not targeted to the D-compartment (*PPM1D*, *SLC9A1*)[25], or for other non-induced genes

(*LPHN2*, *UTP18*). Altogether these data indicate that the circadian rhythm regulates the repair

and signaling of TC-DSBs by controlling their PER2-dependent anchoring to the NE and,

therefore, the formation of the D-compartment, further fine-tuning the response to DSBs.

Discussion

In this study, we uncovered that the repair of DSBs occurring in transcriptionally active

chromatin in human requires their anchoring to the NE, a process that is under the direct control

of PERIOD proteins and is therefore sensitive to circadian oscillations (Fig. 6i). Notably, in a

recent preprint, the Mekhail lab reports that etoposide induces nuclear envelope tubules

(dsbNET) inside the nucleus which establish contacts with DSBs to foster repair[59]. Since

persistent DSBs and arrested forks in yeast as well as HC breaks in *Drosophila*, also relocate

to the NE and NPC, altogether our findings suggest that targeting repair refractory DSBs to this

specific nuclear microenvironment is evolutionary conserved.

Of interest, previous work uncovered that the maintenance of genomic integrity and cancer

progression are tightly linked to the circadian clock[60–62]. In general, genes involved in DNA

repair pathways are transcriptionally regulated by the circadian clock[61,63–68]. Here we found that

in human cells, PER2 is directly targeted at DSBs. In contrast to a very recent report[69], we

observed neither BMAL1 binding at TC-DSBs, although BMAL1 distribution on the genome

behaved as expected, nor a function of BMAL1/CLOCK heterodimer in promoting cell survival

and counteracting translocation upon TC-DSBs induction in DIvA cells. This difference may

arise from analyzing overexpressed[69] versus endogenous (our study) BMAL1, from the DSB-

inducible systems used in both studies (AsiSI versus Zeocin and I-PpoI) and/or from difference

in the time points analyzed post-DSB induction.

Upon PER1 or PER2 depletion, as well as during the low-PER phase, γH2AX level was

increased while RAD51 assembly was reduced suggesting a decreased repair capacity. Of

interest, in the accompanying paper, the Huertas lab reports that CRY1 is also recruited at radiation-induced DSBs, where it directly represses resection and HR. Altogether our studies suggest a two-arm control of HR repair during the 24h circadian cycle, with minimum HR occurring at night in diurnal species. Indeed, large transcriptomic surveys, performed in baboon[63] and very recently in human[70], allowed to carefully establish the transcriptional pattern of core clock regulators during the day. PERIOD genes *Per1*, *Per2* and *Cry2* display a maximum expression level around noon, while *Cry1* expression peaks at sunset, in agreement with a PERIOD complex-independent function of CRY1. This allows us to propose the following model (Extended Data Fig. 8g): in the morning, the presence of the PERIOD PER1/PER2 proteins combined with the absence of CRY1, establishes an environment that favors HR repair of TC-DSBs by allowing resection to occur, DSB anchoring to the NE and RAD51 loading. Subsequently, the gradual increase of CRY1 during the afternoon causes minimum resection at nightfall. The absence of PERIOD complex expression early at night establishes a second layer of HR repression during the night. The gradual decrease of CRY1, combined with the re-expression of PER proteins, then allows HR to resume the next morning.

Of interest, sleep was found to regulate the repair of physiological DSBs in neurons[71], so whether such a circadian regulation of TC-DSB repair also takes place in post-mitotic cells will deserve further investigation.

In summary, our findings have implications for disease etiology and treatment. Indeed, evidence suggests that a dysregulation of the circadian clock contributes to the progression of neurodegenerative disorders[72] and cancer initiation and progression[60,62], both types of diseases being tightly coupled to DSB repair mechanisms and the maintenance of genome integrity. Moreover, and importantly, topoisomerase II poison-based chemotherapies are a first-line treatment against a number of cancers and given that these poisons mainly trigger DSBs in

active chromatin, our findings that circadian rhythm regulates the repair of TC-DSBs may be an important feature to take into account for chronochemotherapy.

Data and Code availability

Code is available at https://github.com/LegubeDNAREPAIR/CircadianClock. High throughput data are available at Array Express (E-MTAB-12712) using the link https://www.ebi.ac.uk/biostudies/arrayexpress/studies/E-MTAB-12712?key=2f8af589-d077-466a-a563-725b33761cda

Acknowledgments

We thank the genomics core facility of EMBL for high-throughput sequencing. We thank Vanessa Dougados for her help during the RIM acquisitions and the LITC TRI-GENOTOUL photonic imaging platform. We thank L. Ligat from the CRCT Cell Imaging Platform (INSERM-UMR1037) for technical assistance in high-content microscopy. We thank T. Clouaire and N. Firmin for technical help. We thank T. Clouaire, and E. Cau for their advice throughout the project and critical reading of the manuscript. Development of 3D RIM on the LITC was supported by ANR-20-CE45-0024. Funding in GL laboratory was provided by grants from the European Research Council (ERC-2014-CoG 647344; and ERC-AdG-101019963), the Agence Nationale pour la Recherche (ANR-18-CE12-0015), the Fondation ARC pour la recherche sur le cancer, the ITMO Cancer, the Ligue Nationale Contre le Cancer, comité de l'Aude and the Fondation Bettencourt-Schueller. C.A was a recipient of a FRM fellowship (FRM FDT201904007941), S.C. was a recipient of a PhD fellowship from Joint Training and Research Programme on Chromatin Dynamics & the DNA Damage Response (H2020 ITN aDDRess, grant N° 812829), J.F is a recipient of a PhD fellowship from the Horizon Europe

478 Marie Skłodowska-Curie Actions Doctoral Networks CohesiNet (Grant Agreement No.

479 859853). I.L. is supported by a funding from the Fondation ARC pour la recherche sur le cancer.

480 N.P and P.C are INSERM researchers.

Authors contributions

483 B.L.B, L.G.S, J.F, R.C, E.G, C.P, C.A, I.L, A.G, A.L.F, A.M. and N.P, performed and analyzed

484 experiments. P.F and P.C optimized the mAID-AsiSIER lentiviral construct. S.C, M.A, and

485 V.R performed bioinformatic analyses of all high-throughput sequencing datasets. T.M

486 performed RIM acquisition and analysis. G.L. and N.P. supervised the work. G.L. wrote the

487 manuscript. All authors commented, edited and approved the manuscript.

Conflict of interest

491 The authors declare no competing interest

References

1. Marnef, A., Cohen, S., and Legube, G. (2017). Transcription-Coupled DNA Double-
Strand Break Repair: Active Genes Need Special Care. Journal of Molecular Biology *429*,
1277–1288. 10.1016/j.jmb.2017.03.024.

2. Canela, A., Maman, Y., Huang, S.N., Wutz, G., Tang, W., Zagnoli-Vieira, G., Callen, E.,
Wong, N., Day, A., Peters, J.-M., et al. (2019). Topoisomerase II-Induced Chromosome
Breakage and Translocation Is Determined by Chromosome Architecture and
Transcriptional Activity. Molecular Cell *75*, 252-266.e8. 10.1016/j.molcel.2019.04.030.

3. Ballarino, R., Bouwman, B.A.M., Agostini, F., Harbers, L., Diekmann, C., Wernersson,
E., Bienko, M., and Crosetto, N. (2022). An atlas of endogenous DNA double-strand
breaks arising during human neural cell fate determination. Sci Data *9*, 400.
10.1038/s41597-022-01508-x.

4. Gothe, H.J., Bouwman, B.A.M., Gusmao, E.G., Piccinno, R., Petrosino, G., Sayols, S.,
Drechsel, O., Minneker, V., Josipovic, N., Mizi, A., et al. (2019). Spatial Chromosome
Folding and Active Transcription Drive DNA Fragility and Formation of Oncogenic MLL
Translocations. Mol Cell *75*, 267-283.e12. 10.1016/j.molcel.2019.05.015.

5. Madabhushi, R., Gao, F., Pfenning, A.R., Pan, L., Yamakawa, S., Seo, J., Rueda, R.,
Phan, T.X., Yamakawa, H., Pao, P.-C., et al. (2015). Activity-Induced DNA Breaks
Govern the Expression of Neuronal Early-Response Genes. Cell *161*, 1592–1605.
10.1016/j.cell.2015.05.032.

6. Pankotai, T., Bonhomme, C., Chen, D., and Soutoglou, E. (2012). DNAPKcs-dependent
arrest of RNA polymerase II transcription in the presence of DNA breaks. Nat Struct Mol
Biol *19*, 276–282. 10.1038/nsmb.2224.

7. Shanbhag, N.M., Rafalska-Metcalf, I.U., Balane-Bolivar, C., Janicki, S.M., and
Greenberg, R.A. (2010). ATM-dependent chromatin changes silence transcription in cis
to DNA double-strand breaks. Cell *141*, 970–981. 10.1016/j.cell.2010.04.038.

8. Aymard, F., Bugler, B., Schmidt, C.K., Guillou, E., Caron, P., Briois, S., Iacovoni, J.S.,
Daburon, V., Miller, K.M., Jackson, S.P., et al. (2014). Transcriptionally active chromatin
recruits homologous recombination at DNA double-strand breaks. Nat Struct Mol Biol *21*,
366–374. 10.1038/nsmb.2796.

9. Ouyang, J., Yadav, T., Zhang, J.-M., Yang, H., Rheinbay, E., Guo, H., Haber, D.A., Lan,
L., and Zou, L. (2021). RNA transcripts stimulate homologous recombination by forming
DR-loops. Nature *594*, 283–288. 10.1038/s41586-021-03538-8.

10. Aymard, F., Aguirrebengoa, M., Guillou, E., Javierre, B.M., Bugler, B., Arnould, C.,
Rocher, V., Iacovoni, J.S., Biernacka, A., Skrzypczak, M., et al. (2017). Genome-wide
mapping of long-range contacts unveils clustering of DNA double-strand breaks at
damaged active genes. Nat Struct Mol Biol *24*, 353–361. 10.1038/nsmb.3387.

11. Arnould, C., Rocher, V., Saur, F., Bader, A.S., Muzzopappa, F., Collins, S., Lesage, E.,
Le Bozec, B., Puget, N., Clouaire, T., et al. (2023). Chromatin compartmentalization
regulates the response to DNA damage. Nature *623*, 183–192. 10.1038/s41586-023-
06635-y.

12. Zagelbaum, J., Schooley, A., Zhao, J., Schrank, B.R., Callen, E., Zha, S., Gottesman,
M.E., Nussenzweig, A., Rabadan, R., Dekker, J., et al. (2022). Multiscale reorganization
of the genome following DNA damage facilitates chromosome translocations via nuclear
actin polymerization. Nat Struct Mol Biol. 10.1038/s41594-022-00893-6.

13. Marnef, A., Finoux, A.-L., Arnould, C., Guillou, E., Daburon, V., Rocher, V., Mangeat,
T., Mangeot, P.E., Ricci, E.P., and Legube, G. (2019). A cohesin/HUSH- and LINC-
dependent pathway controls ribosomal DNA double-strand break repair. Genes Dev. *33*,
1175–1190. 10.1101/gad.324012.119.

14. Nagai, S., Dubrana, K., Tsai-Pflugfelder, M., Davidson, M.B., Roberts, T.M., Brown,
G.W., Varela, E., Hediger, F., Gasser, S.M., and Krogan, N.J. (2008). Functional
Targeting of DNA Damage to a Nuclear Pore-Associated SUMO-Dependent Ubiquitin
Ligase. Science *322*, 597–602. 10.1126/science.1162790.

15. Ryu, T., Spatola, B., Delabaere, L., Bowlin, K., Hopp, H., Kunitake, R., Karpen, G.H.,
and Chiolo, I. (2015). Heterochromatic breaks move to the nuclear periphery to continue
recombinational repair. Nat Cell Biol *17*, 1401–1411. 10.1038/ncb3258.

16. Oza, P., Jaspersen, S.L., Miele, A., Dekker, J., and Peterson, C.L. (2009). Mechanisms
that regulate localization of a DNA double-strand break to the nuclear periphery. Genes
Dev. *23*, 912–927. 10.1101/gad.1782209.

17. Horigome, C., Oma, Y., Konishi, T., Schmid, R., Marcomini, I., Hauer, M.H., Dion, V.,
Harata, M., and Gasser, S.M. (2014). SWR1 and INO80 Chromatin Remodelers
Contribute to DNA Double-Strand Break Perinuclear Anchorage Site Choice. Molecular
Cell *55*, 626–639. 10.1016/j.molcel.2014.06.027.

18. Kramarz, K., Schirmeisen, K., Boucherit, V., Ait Saada, A., Lovo, C., Palancade, B.,
Freudenreich, C., and Lambert, S.A.E. (2020). The nuclear pore primes recombination-
dependent DNA synthesis at arrested forks by promoting SUMO removal. Nat Commun
11, 5643. 10.1038/s41467-020-19516-z.

19. Whalen, J.M., Dhingra, N., Wei, L., Zhao, X., and Freudenreich, C.H. (2020). Relocation
of Collapsed Forks to the Nuclear Pore Complex Depends on Sumoylation of DNA
Repair Proteins and Permits Rad51 Association. Cell Rep *31*, 107635.
10.1016/j.celrep.2020.107635.

20. Iacovoni, J.S., Caron, P., Lassadi, I., Nicolas, E., Massip, L., Trouche, D., and Legube, G.
(2010). High-resolution profiling of γH2AX around DNA double strand breaks in the
mammalian genome. EMBO J *29*, 1446–1457. 10.1038/emboj.2010.38.

21. Crosetto, N., Mitra, A., Silva, M.J., Bienko, M., Dojer, N., Wang, Q., Karaca, E., Chiarle,
R., Skrzypczak, M., Ginalski, K., et al. (2013). Nucleotide-resolution DNA double-strand
break mapping by next-generation sequencing. Nat Methods *10*, 361–365.
10.1038/nmeth.2408.

22. Clouaire, T., Rocher, V., Lashgari, A., Arnould, C., Aguirrebengoa, M., Biernacka, A.,
Skrzypczak, M., Aymard, F., Fongang, B., Dojer, N., et al. (2018). Comprehensive
Mapping of Histone Modifications at DNA Double-Strand Breaks Deciphers Repair
Pathway Chromatin Signatures. Molecular Cell *72*, 250-262.e6.
10.1016/j.molcel.2018.08.020.

23. Cohen, S., Puget, N., Lin, Y.-L., Clouaire, T., Aguirrebengoa, M., Rocher, V., Pasero, P.,
Canitrot, Y., and Legube, G. (2018). Senataxin resolves RNA:DNA hybrids forming at
DNA double-strand breaks to prevent translocations. Nat Commun *9*, 533.
10.1038/s41467-018-02894-w.

24. Cohen, S., Guenolé, A., Lazar, I., Marnef, A., Clouaire, T., Vernekar, D.V., Puget, N.,
Rocher, V., Arnould, C., Aguirrebengoa, M., et al. (2022). A POLD3/BLM dependent
pathway handles DSBs in transcribed chromatin upon excessive RNA:DNA hybrid
accumulation. Nat Commun *13*, 2012. 10.1038/s41467-022-29629-2.

25. Arnould, C., Rocher, V., Bader, A.S., Lesage, E., Puget, N., Clouaire, T., Mourad, R.,
Noordermeer, D., Bushell, M., and Legube, G. (2021). ATM-dependent formation of a
novel chromatin compartment regulates the Response to DNA Double Strand Breaks and
the biogenesis of translocations (Molecular Biology) 10.1101/2021.11.07.467654.

26. Brown, S.A., Ripperger, J., Kadener, S., Fleury-Olela, F., Vilbois, F., Rosbash, M., and
Schibler, U. (2005). PERIOD1-Associated Proteins Modulate the Negative Limb of the
Mammalian Circadian Oscillator. Science *308*, 693–696. 10.1126/science.1107373.

27. Duong, H.A., Robles, M.S., Knutti, D., and Weitz, C.J. (2011). A Molecular Mechanism
for Circadian Clock Negative Feedback. Science *332*, 1436–1439.
10.1126/science.1196766.

28. Padmanabhan, K., Robles, M.S., Westerling, T., and Weitz, C.J. (2012). Feedback
Regulation of Transcriptional Termination by the Mammalian Circadian Clock PERIOD
Complex. Science *337*, 599–602. 10.1126/science.1221592.

29. Kim, J.Y., Kwak, P.B., and Weitz, C.J. (2014). Specificity in Circadian Clock Feedback
from Targeted Reconstitution of the NuRD Corepressor. Molecular Cell *56*, 738–748.
10.1016/j.molcel.2014.10.017.

30. Emerson, J.M., Bartholomai, B.M., Ringelberg, C.S., Baker, S.E., Loros, J.J., and Dunlap,
J.C. (2015). *period* -1 encodes an ATP-dependent RNA helicase that influences
nutritional compensation of the *Neurospora* circadian clock. Proc. Natl. Acad. Sci. U.S.A.
112, 15707–15712. 10.1073/pnas.1521918112.

31. Ogilvie, V.C. (2003). The highly related DEAD box RNA helicases p68 and p72 exist as
heterodimers in cells. Nucleic Acids Research *31*, 1470–1480. 10.1093/nar/gkg236.

32. Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock.
Nat Rev Genet *18*, 164–179. 10.1038/nrg.2016.150.

33. Sessa, G., Gómez-González, B., Silva, S., Pérez-Calero, C., Beaurepere, R., Barroso, S.,
Martineau, S., Martin, C., Ehlén, Å., Martínez, J.S., et al. (2021). BRCA2 promotes
DNA-RNA hybrid resolution by DDX5 helicase at DNA breaks to facilitate their repair‡.
EMBO J *40*, e106018. 10.15252/embj.2020106018.

34. Qu, M., Qu, H., Jia, Z., and Kay, S.A. (2021). HNF4A defines tissue-specific circadian
rhythms by beaconing BMAL1::CLOCK chromatin binding and shaping the rhythmic
chromatin landscape. Nat Commun *12*, 6350. 10.1038/s41467-021-26567-3.

35. Koike, N., Yoo, S.-H., Huang, H.-C., Kumar, V., Lee, C., Kim, T.-K., and Takahashi, J.S.
(2012). Transcriptional Architecture and Chromatin Landscape of the Core Circadian
Clock in Mammals. Science *338*, 349–354. 10.1126/science.1226339.

36. Tartour, K., and Padmanabhan, K. (2022). The Clock Takes Shape—24 h Dynamics in
Genome Topology. Front. Cell Dev. Biol. *9*, 799971. 10.3389/fcell.2021.799971.

37. Xiao, Y., Yuan, Y., Jimenez, M., Soni, N., and Yadlapalli, S. (2021). Clock proteins
regulate spatiotemporal organization of clock genes to control circadian rhythms. Proc.
Natl. Acad. Sci. U.S.A. *118*, e2019756118. 10.1073/pnas.2019756118.

38. Kalocsay, M., Hiller, N.J., and Jentsch, S. (2009). Chromosome-wide Rad51 Spreading
and SUMO-H2A.Z-Dependent Chromosome Fixation in Response to a Persistent DNA
Double-Strand Break. Molecular Cell *33*, 335–343. 10.1016/j.molcel.2009.01.016.

39. Horigome, C., Unozawa, E., Ooki, T., and Kobayashi, T. (2019). Ribosomal RNA gene repeats associate with the nuclear pore complex for maintenance after DNA damage. PLoS Genet *15*, e1008103. 10.1371/journal.pgen.1008103.

40. Horigome, C., Bustard, D.E., Marcomini, I., Delgoshaie, N., Tsai-Pflugfelder, M., Cobb, J.A., and Gasser, S.M. (2016). PolySUMOylation by Siz2 and Mms21 triggers relocation of DNA breaks to nuclear pores through the Slx5/Slx8 STUbL. Genes Dev. *30*, 931–945. 10.1101/gad.277665.116.

41. Oshidari, R., Strecker, J., Chung, D.K.C., Abraham, K.J., Chan, J.N.Y., Damaren, C.J., and Mekhail, K. (2018). Nuclear microtubule filaments mediate non-linear directional motion of chromatin and promote DNA repair. Nat Commun *9*, 2567. 10.1038/s41467-018-05009-7.

42. Chung, D.K.C., Chan, J.N.Y., Strecker, J., Zhang, W., Ebrahimi-Ardebili, S., Lu, T., Abraham, K.J., Durocher, D., and Mekhail, K. (2015). Perinuclear tethers license telomeric DSBs for a broad kinesin- and NPC-dependent DNA repair process. Nat Commun *6*, 7742. 10.1038/ncomms8742.

43. Khadaroo, B., Teixeira, M.T., Luciano, P., Eckert-Boulet, N., Germann, S.M., Simon, M.N., Gallina, I., Abdallah, P., Gilson, E., Géli, V., et al. (2009). The DNA damage response at eroded telomeres and tethering to the nuclear pore complex. Nat Cell Biol *11*, 980–987. 10.1038/ncb1910.

44. Pinzaru, A.M., Kareh, M., Lamm, N., Lazzerini-Denchi, E., Cesare, A.J., and Sfeir, A. (2020). Replication stress conferred by POT1 dysfunction promotes telomere relocalization to the nuclear pore. Genes Dev *34*, 1619–1636. 10.1101/gad.337287.120.

45. Lamm, N., Read, M.N., Nobis, M., Van Ly, D., Page, S.G., Masamsetti, V.P., Timpson, P., Biro, M., and Cesare, A.J. (2020). Nuclear F-actin counteracts nuclear deformation and promotes fork repair during replication stress. Nat Cell Biol *22*, 1460–1470. 10.1038/s41556-020-00605-6.

46. Caridi, C.P., D'Agostino, C., Ryu, T., Zapotoczny, G., Delabaere, L., Li, X., Khodaverdian, V.Y., Amaral, N., Lin, E., Rau, A.R., et al. (2018). Nuclear F-actin and myosins drive relocalization of heterochromatic breaks. Nature *559*, 54–60. 10.1038/s41586-018-0242-8.

47. Mangeat, T., Labouesse, S., Allain, M., Negash, A., Martin, E., Guénolé, A., Poincloux, R., Estibal, C., Bouissou, A., Cantaloube, S., et al. (2021). Super-resolved live-cell imaging using random illumination microscopy. Cell Reports Methods *1*, 100009. 10.1016/j.crmeth.2021.100009.

48. Lottersberger, F., Karssemeijer, R.A., Dimitrova, N., and de Lange, T. (2015). 53BP1 and the LINC Complex Promote Microtubule-Dependent DSB Mobility and DNA Repair. Cell *163*, 880–893. 10.1016/j.cell.2015.09.057.

49. Liu, Q., Pante, N., Misteli, T., Elsagga, M., Crisp, M., Hodzic, D., Burke, B., and Roux, K.J. (2007). Functional association of Sun1 with nuclear pore complexes. Journal of Cell Biology *178*, 785–798. 10.1083/jcb.200704108.

50. Li, P., and Noegel, A.A. (2015). Inner nuclear envelope protein SUN1 plays a prominent
role in mammalian mRNA export. Nucleic Acids Res, gkv1058. 10.1093/nar/gkv1058.

51. Su, X.A., Dion, V., Gasser, S.M., and Freudenreich, C.H. (2015). Regulation of
recombination at yeast nuclear pores controls repair and triplet repeat stability. Genes Dev
29, 1006–1017. 10.1101/gad.256404.114.

52. Caron, P., van der Linden, J., and van Attikum, H. (2019). Bon voyage: A transcriptional
journey around DNA breaks. DNA Repair *82*, 102686. 10.1016/j.dnarep.2019.102686.

53. Zhou, Y., Caron, P., Legube, G., and Paull, T.T. (2014). Quantitation of DNA double-
strand break resection intermediates in human cells. Nucleic Acids Research *42*, e19–e19.
10.1093/nar/gkt1309.

54. Galanty, Y., Belotserkovskaya, R., Coates, J., and Jackson, S.P. (2012). RNF4, a SUMO-
targeted ubiquitin E3 ligase, promotes DNA double-strand break repair. Genes Dev *26*,
1179–1195. 10.1101/gad.188284.112.

55. Han, M.M., Hirakawa, M., Yamauchi, M., and Matsuda, N. (2022). Roles of the SUMO-
related enzymes, PIAS1, PIAS4, and RNF4, in DNA double-strand break repair by
homologous recombination. Biochem Biophys Res Commun *591*, 95–101.
10.1016/j.bbrc.2021.12.099.

56. Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T.,
Telling, A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009).
Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the
Human Genome. Science *326*, 289–293. 10.1126/science.1181369.

57. Falk, M., Feodorova, Y., Naumova, N., Imakaev, M., Lajoie, B.R., Leonhardt, H., Joffe,
B., Dekker, J., Fudenberg, G., Solovei, I., et al. (2019). Heterochromatin drives
compartmentalization of inverted and conventional nuclei. Nature *570*, 395–399.
10.1038/s41586-019-1275-3.

58. Roukos, V., Voss, T.C., Schmidt, C.K., Lee, S., Wangsa, D., and Misteli, T. (2013).
Spatial Dynamics of Chromosome Translocations in Living Cells. Science *341*, 660–664.
10.1126/science.1237150.

59. Shokrollahi, M., Stanic, M., Hundal, A., Chan, J. N.Y., Urman, D., Hakem, A., Garcia,
R.E., Hao, J., Maass, P.G., Dickson, B.C., et al. (2023). DNA double-strand break-
capturing nuclear envelope tubules drive DNA repair. bioRxiv, 2023.05.07.539750.
10.1101/2023.05.07.539750.

60. Sancar, A., and Van Gelder, R.N. (2021). Clocks, cancer, and chronochemotherapy.
Science *371*, eabb0738. 10.1126/science.abb0738.

61. Kondratov, R.V., and Antoch, M.P. (2007). Circadian proteins in the regulation of cell
cycle and genotoxic stress responses. Trends in Cell Biology *17*, 311–317.
10.1016/j.tcb.2007.07.001.

62. Masri, S., and Sassone-Corsi, P. (2018). The emerging link between cancer, metabolism,
and circadian rhythms. Nat Med *24*, 1795–1803. 10.1038/s41591-018-0271-8.

63. Mure, L.S., Le, H.D., Benegiamo, G., Chang, M.W., Rios, L., Jillani, N., Ngotho, M.,
Kariuki, T., Dkhissi-Benyahya, O., Cooper, H.M., et al. (2018). Diurnal transcriptome
atlas of a primate across major neural and peripheral tissues. Science *359*, eaao0318.
10.1126/science.aao0318.

64. Alexandrou, A.T., Duan, Y., Xu, S., Tepper, C., Fan, M., Tang, J., Berg, J., Basheer, W.,
Valicenti, T., Wilson, P.F., et al. (2022). PERIOD 2 regulates low-dose radioprotection
via PER2/pGSK3β/β-catenin/Per2 loop. iScience *25*, 105546. 10.1016/j.isci.2022.105546.

65. Papp, S.J., Huber, A.-L., Jordan, S.D., Kriebs, A., Nguyen, M., Moresco, J.J., Yates, J.R.,
and Lamia, K.A. (2015). DNA damage shifts circadian clock time via Hausp-dependent
Cry1 stabilization. eLife *4*, e04883. 10.7554/eLife.04883.

66. Shafi, A.A., McNair, C.M., McCann, J.J., Alshalalfa, M., Shostak, A., Severson, T.M.,
Zhu, Y., Bergman, A., Gordon, N., Mandigo, A.C., et al. (2021). The circadian
cryptochrome, CRY1, is a pro-tumorigenic factor that rhythmically modulates DNA
repair. Nat Commun *12*, 401. 10.1038/s41467-020-20513-5.

67. Anabtawi, N., Cvammen, W., and Kemp, M.G. (2021). Pharmacological inhibition of
cryptochrome and REV-ERB promotes DNA repair and cell cycle arrest in cisplatin-
treated human cells. Sci Rep *11*, 17997. 10.1038/s41598-021-97603-x.

68. Gaddameedhi, S., Selby, C.P., Kaufmann, W.K., Smart, R.C., and Sancar, A. (2011).
Control of skin cancer by the circadian rhythm. Proc. Natl. Acad. Sci. U.S.A. *108*, 18790–
18795. 10.1073/pnas.1115249108.

69. Zhang, C., Chen, L., Sun, L., Jin, H., Ren, K., Liu, S., Qian, Y., Li, S., Li, F., Zhu, C., et
al. (2023). BMAL1 collaborates with CLOCK to directly promote DNA double-strand
break repair and tumor chemoresistance. Oncogene. 10.1038/s41388-023-02603-y.

70. Talamanca, L., Gobet, C., and Naef, F. (2023). Sex-dimorphic and age-dependent
organization of 24-hour gene expression rhythms in humans. Science *379*, 478–483.
10.1126/science.add0846.

71. Zada, D., Sela, Y., Matosevich, N., Monsonego, A., Lerer-Goldshtein, T., Nir, Y., and
Appelbaum, L. (2021). Parp1 promotes sleep, which enhances DNA repair in neurons.
Molecular Cell *81*, 4979-4993.e7. 10.1016/j.molcel.2021.10.026.

72. Nassan, M., and Videnovic, A. (2022). Circadian rhythms in neurodegenerative disorders.
Nat Rev Neurol *18*, 7–24. 10.1038/s41582-021-00577-7.

73. Caron, P., Choudjaye, J., Clouaire, T., Bugler, B., Daburon, V., Aguirrebengoa, M.,
Mangeat, T., Iacovoni, J.S., Álvarez-Quilón, A., Cortés-Ledesma, F., et al. (2015). Non-
redundant Functions of ATM and DNA-PKcs in Response to DNA Double-Strand
Breaks. Cell Reports *13*, 1598–1609. 10.1016/j.celrep.2015.10.024.

74. Natsume, T., Kiyomitsu, T., Saga, Y., and Kanemaki, M.T. (2016). Rapid Protein
Depletion in Human Cells by Auxin-Inducible Degron Tagging with Short Homology
Donors. Cell Rep *15*, 210–218. 10.1016/j.celrep.2016.03.001.

75. Nemoz, C., Ropars, V., Frit, P., Gontier, A., Drevet, P., Yu, J., Guerois, R., Pitois, A.,
Comte, A., Delteil, C., et al. (2018). XLF and APLF bind Ku80 at two remote sites to

745   ensure DNA repair by non-homologous end joining. Nat Struct Mol Biol *25*, 971–980.
746   10.1038/s41594-018-0133-6.

747 76. Fu, C., Donovan, W.P., Shikapwashya-Hasser, O., Ye, X., and Cole, R.H. (2014). Hot
748   Fusion: an efficient method to clone multiple DNA fragments as well as inverted repeats
749   without ligase. PLoS One *9*, e115318. 10.1371/journal.pone.0115318.

750 77. Cheng, Q., Barboule, N., Frit, P., Gomez, D., Bombarde, O., Couderc, B., Ren, G.-S.,
751   Salles, B., and Calsou, P. (2011). Ku counteracts mobilization of PARP1 and MRN in
752   chromatin damaged with DNA double-strand breaks. Nucleic Acids Res *39*, 9605–9619.
753   10.1093/nar/gkr656.

754 78. Li, Q. (2004). A Syntaxin 1, G o, and N-Type Calcium Channel Complex at a Presynaptic
755   Nerve Terminal: Analysis by Quantitative Immunocolocalization. Journal of
756   Neuroscience *24*, 4070–4081. 10.1523/JNEUROSCI.0346-04.2004.

757 79. van Steensel, B., van Binnendijk, E.P., Hornsby, C.D., van der Voort, H.T., Krozowski,
758   Z.S., de Kloet, E.R., and van Driel, R. (1996). Partial colocalization of glucocorticoid and
759   mineralocorticoid receptors in discrete compartments in nuclei of rat hippocampus
760   neurons. Journal of Cell Science *109*, 787–792. 10.1242/jcs.109.4.787.

761 80. Arnould, C., Rocher, V., Finoux, A.-L., Clouaire, T., Li, K., Zhou, F., Caron, P.,
762   Mangeot, Philippe.E., Ricci, E.P., Mourad, R., et al. (2021). Loop extrusion as a
763   mechanism for formation of DNA damage repair foci. Nature *590*, 660–665.
764   10.1038/s41586-021-03193-z.

Figure Legends

Figure 1: A multi-output screen for the identification of new TC-DSBR factors.

a. Experimental scheme: DIvA cells allow to temporally induce, following treatment with 4-hydroxytamoxifen (4-OHT), DSBs at annotated positions on the human genome, among which >50% fall in active loci (TC-DSBs). TC-DSB potential interacting partners were identified through a proteomic approach[13]. 130 proteins were further subjected to a multi-output screen in AID-DIvA cells using a siRNA library, to score DSB-induced γH2AX foci intensity (using quantitative high throughput imaging), DSB clustering (γH2AX foci mean area), chromosome rearrangement (illegitimate rejoining between two TC-DSBs *t(MIS12;TRIM37)* analyzed by qPCR following DSB induction and repair) and cell survival (quantified eight days after DSB induction, using a colorimetric assay based on mitochondrial dehydrogenase activity in viable cells).

b. Heatmap of the z-score for the 4 outputs measured (γH2AX foci intensity, Clustering, Translocation and Survival) in the screen of the 130 genes of interest ordered by hierarchical clustering (Pearson correlation distance/ward.D method) after normalization of the data to a control siRNA (see Extended Data Fig. 1b-e). A positive control (SETX) was included in the experiment (red). PPHLN1 and PER complex siRNAs are highlighted in green and blue respectively.

c. Violin plots of scored outputs for the two groups of genes defined by the hierarchical clustering. *P*, student t-test.

d-g. Scatter plots showing the correlations for each siRNA between γH2AX intensity and cell survival (**d**); γH2AX intensity and translocations (**e**); translocations and DSB clustering (**f**) and

cell survival and translocations (**g**). The 95% confidence intervals and the coefficient of
correlation (r, Pearson) are indicated.

Figure 2. PER proteins are recruited at TC-DSBs and regulate γH2AX foci formation,
translocation, and cell survival upon DSB induction

a. γH2AX staining performed in control (CTRL), PER1 or PER2-siRNA transfected DIvA
cells, before (-DSB) and after (+DSB) DSB induction. Quantification (normalized against
Hoechst intensity) is shown on the bottom panel (>7000 nuclei, from a representative
experiment). Center line: median; box limits: 1st and 3rd quartiles; whiskers: maximum and
minimum without outliers. *P*, unpaired t-test.

b. γH2AX ChIP performed in DIvA cells transfected with siRNA control (CTRL) or PER2
before and after DSB induction and analyzed by qPCR at two DSBs (DSB1-2). Data are
normalized to a control location devoid of a DSB. Mean and SEM are shown for n=4 biological
replicates. *P,* paired t-test (two-sided).

c. *t(MIS12:TRIM37)* rejoining frequencies after DSB induction measured by qPCR in AID-
DIvA cells transfected with siRNA CTRL, PER1 PER2, DDX17, SETX, NONO, DDX5 and
BMAL1. Mean and SD (n≥4 biological replicates) are shown. *P*, paired t-test (two-sided).

d. Clonogenic assays in AID-DIvA cells transfected with siRNA CTRL, PER1, PER2, DDX17,
NONO, DDX5 or BMAL1. Upper and lower panels show the mean and SEM after 4-OHT
treatment (+DSB) and after IAA treatment (+DSB+Repair) respectively (n≥4 biological
replicates). *P*, paired t-test (two-sided).

e. PER2, DDX5, DDX17 and BMAL1 ChIP efficiency (expressed as % of input
immunoprecipitated) before (−DSB) and after (+DSB) DSB induction, at a control genomic

locus devoid of DSB, a promoter (*BTG2* or *PER1* as indicated), and DSB1 or DSB4. Mean and

SEM are shown for n≥10 replicates. *P*, paired t-test (two-sided).

f. Genomic tracks showing RNA Polymerase II before DSB induction (RNAPII, green), BLESS

after DSB induction (black), as well as BMAL1 (blue) and PER2 (violet) (log2(+DSB/−DSB))

at a TC-DSB (upper panel; chr21:460221789) and a DSB induced in a silent locus (lower panel;

chr12:130091880). DSBs are indicated by arrows.

g. Average PER2 (left) and BMAL1 (right) ChIP-seq profiles on a ±5 kb window centered on

the eighty best-induced AsiSI-DSBs (top) or eighty random sites (bottom, no DSB). Data are

presented in log2(+DSB/-DSB).

h. Box plots representing Log2 (+DSB/-DSB) PER2 ChIP-seq count on a ±2kb at HR-DSB,

NHEJ-DSB (n=30 in each category, among the 80 best-induced DSBs) or random sites (n=30).

Center line: median; box limits: 1st and 3rd quartiles; whiskers: maximum and minimum

without outliers. *P*, Wilcoxon test.

Figure 3: TC-DSBs are targeted to the nuclear envelope via a PER2-dependent

mechanism

a. Super-resolution live imaging performed using Random Illumination microscopy (RIM) in

DIvA cells expressing 53BP1-GFP and mCherry-LaminB1, starting >1h after DSB induction.

Arrows show the establishment of a contact between a focus and nuclear lamina. Images were

acquired every 40s.

b. Proximity Ligation Assay (PLA) performed using either 53BP1 antibody (top panels), Lamin

B1 antibody (middle panels) or both 53BP1 and LaminB1 antibodies (bottom panels) before (-

DSB) and after DSB (+DSB) induction in DIvA cells. PLA quantification is shown on the right

panel. An average of 190 cells were analyzed per condition. Center line: median; box limits: 1st and 3rd quartiles; whiskers: maximum and minimum without outliers. *P*, Wilcoxon test.

c. LaminB1 ChIP-qPCR before (−DSB) and after (+DSB) DSB induction, at TC-DSBs (HR-DSBs, DSB1-4) and one DSB induced in a silent locus and repaired by NHEJ (DSB5). Data are normalized to a control location devoid of a DSB. Mean and SEM are shown for n=7 biological replicates. *P*, paired t-test (two-sided).

d. LaminB1 ChIP-qPCR before (−DSB) and after (+DSB) DSB induction in control and PER2 siRNA-depleted DIvA cells, at TC-DSBs (DSB1-4, HR-DSBs) and a DSB in a silent locus (DSB5, NHEJ-DSBs). Data are normalized to a control location devoid of DSB. Mean and SEM are shown for n=4 biological replicates. *P*, paired t-test (two-sided).

e. Number of PLA foci (53BP1-LaminB1) in CTRL, PER1 or PER2-siRNA transfected DIvA cells as indicated, before (−DSB) and after (+DSB) DSB induction. Data were normalized to the untreated samples. Mean and SEM (n=3 biological replicates) are shown. *P*, paired t-test (two-sided).

f. Genomic tracks of SUN1(purple) and SUN2 (dark blue) ChIP-seq (log2(+DSB/-DSB)) at a TC-DSB (upper panel; chr17:57184296) and a DSB induced in a silent locus (lower panel; chr12:130091880).

g. Average profiles of SUN1 (left) and SUN2 (right) on a ±5 kb window centered on the eighty best-induced AsiSI-DSBs (top) or eighty random sites (bottom, no DSB). Data are presented as log2(+DSB/-DSB).

h. Magnifications of SUN2 (top panels) and SUN1 (bottom panels) staining in nuclear envelope (dotted line) together with γH2AX foci in DIvA cells after DSB induction. Arrows indicate sites of colocalization.

i. Cross Correlation Function (CCF) plots between γH2AX foci and NE-embedded SUN2 (top panel, N=70 γH2AX foci) or SUN1 (bottom panel, N=46 γH2AX foci). CCF of bicolor 3D RIM images was determined by plotting the value of Pearson's correlation coefficient P[0.1] against Δx for each voxel of the image (see methods). The arrows show a non-random exclusion between γH2AX and SUN2 (dip at Δx=0) or a non-random overlap between the signals of SUN1and γH2AX (peak at Δx=0).

j. LaminB1 ChIP-qPCR before (−DSB) and after (+DSB) DSB induction in control and SUN1 siRNA-depleted DIvA cells, at two TC-DSBs (DSB1 and DSB3, HR-DSBs) and two DSBs in silent loci (DSB5 and DSB6, NHEJ-DSBs). Data are normalized to a control location devoid of DSB. Mean and SEM are shown for n=3 biological replicates. *P*, paired t-test (two-sided).

k. Left panel: Magnification of NUP153 staining in nuclear envelope (dotted line) together with γH2AX foci in DIvA cells after DSB induction. Arrows indicate sites of colocalization. Right panel: CCF plot as above between γH2AX foci and NPC protein NUP153 (N=28 γH2AX foci) showing a non-random overlap between both signals (peak at Δx=0).

Figure 4. PER2 and SUN1-dependent anchoring at the Nuclear Pore Complex promotes Homologous Recombination repair

a. RT-qPCR performed before (−DSB) and after (+DSB) DSB induction in control (CTRL), PER1 or PER2 siRNA-depleted DIvA cells for six genes carrying a TC-DSB (relative to untreated sample (-DSB)). Mean and SEM are shown for n=6 biological replicates. *P*, paired t-test (two-sided).

b. RT-qPCR performed before (−DSB) and after (+DSB) DSB induction in control (CTRL) or SUN1 siRNA-depleted DIvA cells for six genes carrying a TC-DSB (relative to untreated

sample (-DSB)). Mean and SEM are shown for n=3 biological replicates. *P*, paired t-test (two-

sided).

c. RAD51 ChIP-qPCR in DIvA cells transfected with control or PER2 siRNA before and after

DSB induction, at four TC-DSBs (DSB1-4) and one DSB in a silent locus (DSB5). Data are

normalized to a control location devoid of a DSB, and further expressed relative to data obtained

at DSB4. Mean and SEM are shown for n=3 biological replicates. *P*, paired t-test (two-sided).

d. Ratios of RAD51 over RPA ChIP-qPCR results in DIvA cells transfected with control

(CTRL) or SUN1 siRNA after DSB induction, at four TC-DSBs (DSB1-4). Data are normalized

to a control location devoid of a DSB, and further expressed relative to data obtained at DSB4.

Mean and SEM are shown for n=3 biological replicates. *P*, paired t-test (two-sided).

e-j. Clonogenic assays in AID-DIvA cells transfected with the indicated siRNAs. Mean and

SEM of n biologically independent experiments after OHT and IAA treatment (DSB induction

and repair) are shown. *P*, paired t-test (two-sided). **e**. Control (CTRL), SUN1 or SUN2 siRNAs

(n=5). **f.** CTRL or NUP153 siRNAs (n=6). **g.** CTRL or RNF4 siRNAs (n=6). **h**. CTRL, PER1,

PER2, and double PER1+SUN1 and PER2+SUN1 siRNAs (n=4). **i.** CTRL, NUP153, and

double PER2+NUP153 and SUN1+NUP153 siRNAs (n=3). **j.** CTRL, PER2, RNF4 and double

PER2+RNF4 siRNAs (n=3).

Figure 5. PER proteins, SUN1, NUP153 and RNF4-mediated TC-DSB anchoring to the

nuclear envelope counteract DSB clustering and D-compartment formation

a. Differential Hi-C contact matrix [(+DSB)-(-DSB)] on a region located on chromosome 2 at

25kb resolution in DIvA cells transfected with a control siRNA or a siRNA directed against

PER2 as indicated. γH2AX ChIP-seq track (+DSB) are also shown (dark red).

b. Interchromosomal DSB interaction shown as aggregate peak analysis (APA) plotted on a

200kb window (10kb resolution) before (-DSB) and after DSB (+DSB) induction in control and

PER2 siRNA-transfected DIvA cells for the eighty best-induced AsiSI-DSBs (left) or eighty

random sites (right, no DSB). FC: Fold change calculated between the central pixel and a square

of 3x3 pixels on the bottom left corner of the matrix. APAs show an increased signal at the

center (DSB-DSB interaction) after DSBs induction only for DSB sites (FC= 3.49, compared

to 0.89 for random sites)). The signal increases after PER2 depletion (FC= 3.85).

c. Box plot showing the differential Hi-C read counts (as (log2 +DSB/-DSB)) between the 80

best induced DSBs (+/- 500kb) (left) or 80 random sites (right) in control (CTRL, grey) and

PER2 (purple) siRNA-transfected DIvA cells. Center line: median; box limits: 1st and 3rd

quartiles; whiskers: maximum and minimum without outliers. *P*, non-parametric Wilcoxon test.

d. Genomic tracks of γH2AX ChIP-seq after DSB induction (dark red) and of Chromosomal

Eigen vectors (CEV) obtained from PCA analyses performed on differential +DSB/-DSB Hi-

C matrices. CEV in siCTRL (grey) and siPER2 (blue). The D-compartment is represented as

positive values. siPER2 triggered the appearance of the D-compartment on the chromosome 2

normally devoid of D compartmentalization post-DSB induction in DIvA cells (areas

highlighted in light grey).

e-h. *t(MIS12:TRIM37)* rejoining frequency after DSB induction and repair measured by qPCR

in AID-DIvA cells transfected with the indicated siRNAs. Mean and SEM of n biological

replicates are shown. *P*, paired t-test (two-sided). **e**. Control (CTRL), SUN1, SUN2, NUP153

and RNF4 siRNAs (n≥4). **f**. CTRL, SUN1 and double PER1+SUN1 and PER2+SUN1 siRNAs

(n=3). **g**. CTRL, PER2, SUN1 and double PER2+NUP153 and SUN1+NUP153 siRNAs (n=3).

h. CTRL, PER2, RNF4 and double PER2+RNF4 siRNAs (n=3).

Figure 6. The Circadian rhythm regulates TC-DSB targeting to nuclear envelope and RAD51 loading

a. γH2AX foci intensity, measured by high content microscopy, in DIvA cells after DSB induction (+DSB 2h), in G1, S and G2 phases at different time points after dexamethasone (Dex) treatment (synchronization of the circadian rhythm). Bars at the different time points are colored according to the PER2 expression level (from highest in dark cyan, to lowest in white). Data were normalized to the average. Mean and SEM (n = 3 biological replicates, with >3000 nuclei acquired per experiment) are shown. *P*, paired t-test (two-sided) was computed using high PER time-points (28h, 30h, 32h) compared to low PER time-points (40h, 42h, 44h).

b. γH2AX ChIP-qPCR performed in DIvA cells before (-DSB) and after DSB (+DSB) induction at 28h and 40h after dexamethasone treatment (Dex) at three TC-DSBs (DSB1-3). Data were normalized to a control location devoid of DSB. Mean and SEM for n=4 biological replicates. *P,* paired t-test (two-sided).

c. LaminB1 ChIP-qPCR performed in DIvA cells before (-DSB) and after DSB induction (+DSB) at 28h and 40h after dexamethasone (Dex) treatment at four TC-DSBs (DSB1-4) and DSB5 (DSB in silent locus). Data were normalized to a control location devoid of DSB and expressed relative to the undamaged condition. Mean and SEM for n=5 biological replicates. *P*, paired t-test (two-sided).

d. SUN1 ChIP-qPCR performed in DIvA cells before (-DSB) and after DSB induction (+DSB) at 28h and 40h after dexamethasone (Dex) treatment at four TC-DSBs (DSB1-4) and DSB5. Data are normalized to a control location devoid of DSB and further expressed related to data obtained in the undamaged condition. Mean and SEM for n=3 biological replicates. *P*, paired t-test (two-sided).

e. Number of PLA foci (γH2AX-LaminB1) per nucleus in DIvA cells before (-DSB) and after DSB induction (+DSB) at 28h and 40h after dexamethasone (Dex) treatment. Data were normalized to the untreated samples. Mean and SEM (n=3 biological replicates) are shown. *P*, paired t-test (two-sided).

f. RAD51 ChIP-qPCR performed in DIvA cells before (-DSB) and after DSB (+DSB) induction at 28h and 40h after dexamethasone (Dex) treatment at four TC-DSBs (DSB1-4) and two NHEJ-DSBs (DSB5-6). Data were normalized to a control location devoid of DSB and expressed relative to DSB4-28h. Mean and SEM for n=6 biological replicates. *P*, paired t-test (two-sided).

g. γH2AX foci mean area, measured by high content microscopy, in DIvA cells after DSB induction (+DSB 2h), in G1, S and G2 phase at different time points after dexamethasone (Dex) treatment. Data were normalized to the average. Mean and SEM (n=3 biological replicates) are shown. *P*, paired t-test (two-sided) was computed using high PER time points (28h, 30h, 32h) compared to low PER time points (40h, 42h, 44h)..

h. DSB responsive genes expression was measured by RT-qPCR before (−DSB) and after (+DSB) DSB induction in DIvA cells at 28h and 40h after dexamethasone treatment. cDNA level of two D-compartment DSB responsive gene (*RNF19B* and *PLK3*), two non-D compartment DSB responsive genes (*PPM1D* and *SLC9A1*) and two control genes not regulated post DSB induction (*LPHN2*, *UTP18*) are shown. Mean and SEM for n=4 biological replicates. *P,* paired t-test (two-sided).

i. Model. During the day (PER phase), PERIOD proteins are recruited at DSBs occurring in transcribed loci (TC-DSBs). This triggers the targeting of the TC-DSBs to the NE and their tethering to SUN1 and the Nuclear Pore Complex, further ensuring RAD51 loading and HR repair. During the night (BMAL1 phase), in absence of PER1/2, impaired TC-DSB anchoring

to the NE leads to impaired RAD51 loading, and enhanced DSB clustering and D-compartment

formation.

Methods

Cell culture and treatment

U2OS, DIvA (AsiSI-ER-U20S)[20], AID-DIvA (AID-AsiSI-ER-U20S)[8], 53BP1-GFP DIvA[73],

hTERT RPE1 and RPE-DIvA (mAID-AsiSI-ER-RPE1) cells were grown in Dulbecco's

modified Eagle's medium (DMEM) supplemented with 10%SVF (Invitrogen), antibiotics and

either 1µg/mL puromycin (DIvA cells) or 800µg/mL G418 (AID-DIvA cells) or both

puromycin and G418 (53BP1-GFP DIvA) or 10µg/ml Blasticidine (RPE-DIvA) at 37°C under

a humidified atmosphere with 5%CO_2. The cell lines were regularly checked for mycoplasma

contamination. For AsiSI-dependent DSB induction, DIvA cell lines were treated with 300nM

4-hydroxytamoxifen (OHT) (Sigma, H7904) for 4h (unless otherwise indicated). For DSB

induction with etoposide (Sigma, E1383), U2OS cells were treated with 0,5µM etoposide for

4h, dexamethasone-synchronized cells were treated with 1µM etoposide for 2h and hTERT

RPE1 cells were treated with 1µM etoposide for 4h. For DSB induction with doxorubicin

(Sigma, D1515), U2OS cells were treated with 0,1µM doxorubicin for 4h. For DDR kinase

inhibition, DIvA cells were pretreated for 1h and during subsequent DSB induction with either

20µM ATMi (KU-55933 from Sigma, SML1109), 2µM ATRi (ETP-46464 from Sigma,

SML1321) or 2µM DNAPKi (NU-7441 from Selleckchem, S2638). To stop inducing AsiSI-

DSBs, AID-DIvA cells were washed twice in pre-warmed PBS after OHT treatment and further

incubated with 500µM auxin (IAA) (Sigma, I5148) for 2h (unless otherwise indicated). For

circadian rhythm synchronization, cells were incubated for 1h with 500nM dexamethasone

(Dex) Sigma, D4902), then the medium was replaced (time = 0h). Dex treatment was done

every 2h during 24h and the DSB induction was started for all synchronized cells at the same

time, 24h after the last Dex treatment, to obtain a full period circadian rhythm (analyses are

shown from 24h to 46h post-Dex). To induce DSBs, synchronized cells are treated during 2h

or 4h surrounding the synchronization timepoint (*e.g.*, from 23h to 25h after Dex for a 2h

treatment corresponding at the synchronized time = 24h).

1007

RPE-DIvA cell line generation

Generation of mAID-AsiSI expression vectors

pLV-OsTIR1-2Myc-T2A-mAID-HA-ER-NLS-AsiSI-P2A-PuroR. The lentiviral vector allowing

the coexpression of (i) the AsiSI restriction enzyme amino-terminally fused to a mini-auxin

inducible degron (mAID)[74], a human influenza hemagglutinin (HA) tag, a modified estrogen

receptor ligand binding domain (ER) and a nuclear localization signal (NLS) of SV40 large T

antigen, (ii) the Myc-tagged rice F-box protein TIR1, and (iii) an antibiotic resistance selectable

marker, was generated from the previously described pLV3 plasmid[75] by the following

sequential modifications. First, a coding sequence for the T2A self-cleaving peptide was

inserted between the Kpn2I and MluI restriction sites using pre-annealed T2A-F and T2A-R

oligonucleotides. The mAID cDNA was then PCR-amplified using mAID-F and mAID-R

primers and the pAID1.1-N plasmid (BioROIS, Japan) as a template, and inserted between

XmaI and EcoRI restriction sites of the previous plasmid. A P2A cassette encoding the self-

cleaving peptide (pre-annealed P2A-F and P2A-R oligonucleotides) was further inserted

between MluI and EcoRI sites. Next, a cDNA fragment coding for puromycin-resistance was

amplified by PCR with primers Puro-F and Puro-R, and added by Hot-Fusion[76] between EcoRI

and NdeI restriction sites. The HA-ER-NLS-AsiSI coding sequence was then PCR-amplified

from a previously described pBabe-Puro-HA-ER-NLS-AsiSI vector[20] with primers AsiSI-F and

AsiSI-R and inserted by Hot-Fusion between BamHI and MluI restriction sites. Finally, the

OsTIR1-2Myc coding sequence was amplified by PCR (primers TIR1-F and TIR1-R) from a

synthetic DNA construct with optimized codons and inserted by Hot-Fusion at the MssI site.

pLV-OsTIR1-2Myc-T2A-mAID-HA-ER-NLS-AsiSI-P2A-BSD. As hTERT RPE1 are already

resistant to puromycin, puromycin-resistance gene was switched to blasticidin-resistance by

NEBuilder HiFi DNA Assembly strategy. Amplification of pLV-OsTIR1-2Myc-T2A-mAID-

HA-ER-NLS-AsiSI-P2A-BSD was done by PCR using F1_fwd, F1_rev, F2_fwd, F2_rev,

F3_fwd, F3_rev primers. Coding sequence for blasticidin resistance was amplified from

pCRIS-PITCHv2_BSD_dTAG plasmid (Addgene, #9179) using primers BSD_fwd and

BSD_rev. PCR were performed using PrimeSTAR Max DNA Polymerase (Takara, ref R045A).

0.1pmol of each fragment were then assembled for 1h at 50°C using NEBuilder HiFi DNA

Assembly master mix from the NEBuilder HiFi DNA Assembly Cloning Kit (NEB, ref

E5520S). Reaction products were amplified in provided bacteria and purified using QIAprep

Spin Miniprep Kit.

The intermediate and final constructs were checked by sequencing (Eurofins Genomics,

Ebersberg, Germany). All restriction and modification enzymes were purchased from

Thermofisher Scientific (Illkirch, France). Oligonucleotides were from Eurofins Genomics.

Production of lentiviral particles in HEK-293T cells and transduction of cells were performed

as previously described[77].

Generation of RPE-DIvA cell line

3.10^5 cells were infected with lentiviral particules in DMEM, Hepes pH 7.4 10mM and

polybrene 8µg/ml. Media was changed 18h after and transduced cells were selected with

12µg/ml blasticidin for 10 days. Clones were isolated and selected according to γH2AX and

AsiSI staining by immunofluorescence, and survival after DSB induction by 300nM OHT

treatment for 4 hours.

1051

Multi-output screen methodology

1052

Each of the 130 SMARTpool siRNAs from the TC-DSBR-focused siRNA library (Extended

Data Table 1) was transfected in 10^6 AID-DIvA cells as described below (siRNA and plasmid

transfection section), always including a negative (CTRL) and a positive (SETX) control

siRNAs. Transfected cells were plated (i) in 96-well Cell Carrier Ultra plates (20.000 cells/well

in triplicates per condition) for γH2AX foci analysis (intensity and area) with quantitative high

throughput imaging as described below (High-Content microscopy and Immunofluorescence

sections); (ii) in 96-well plates (1000 cells/well in quadruplicates per condition for cell survival

analysis as described below (WST-1 cell survival assay section); and in 10cm diameter dish for

genomic DNA extraction and translocation frequency analysis as described below (DSB-

induced rearrangement /Translocation assay section). All data from these different outputs post-

DSB induction were expressed normalized to the negative control siRNA (CTRL) and

compared to the positive control siRNA (SETX). For the 130 genes of interest, each measured

variable ("γH2AX.foci Intensity", "Clustering", "Translocation" and "Survival") was centered

and scaled to obtain the z-score displayed in a heatmap. Genes were ordered according

hierarchical clustering computed with Pearson correlation distance and ward.D clustering

method. Then genes were divided in 2 groups based on the hierarchical clustering and means

of z-scores for each group were computed and displayed in independent boxplot for each

variable. Mean differences between groups were tested using student t-test.

siRNA and plasmid transfection

1071

In U2OS and U2OS derived cell lines, siRNA transfections were performed using the 4D-

Nucleofector and the SE cell line 4D-Nucleofector X kit**s** (Lonza) according to the

manufacturer's instructions. Briefly, 1-10μL of 100μM annealed siRNA was transfected in 1-

20x10^6 cells with 20-100μL SE solution in Nucleocuvettes using the U2OS program CM-104.

In RPE1 cell lines, 20nM of siRNA were transfected using Lipofectamine RNAiMAX reagent

(Invitrogen 13778075) according to the manufacturer's instructions. Subsequent cell treatments

were performed ~42-48h post-transfection. Most siRNAs used in this study were siGENOME

SMARTpools (Dharmacon) which are a mixture of 4 individual siRNAs, see Extended Data

Table 1. Individual siRNA sequences (Eurogentec) were used for SUN1, SUN2 or RBBP8/CtIP

(Extended Data Table 1). For all experiments, the control siRNA (CTRL) was the siGENOME

Non-Targeting Control siRNA Pool #2 (Dharmacon), except for Fig. 4h and Fig. 4i where the

control siRNA used with SUN1 and SUN2 siRNAs was Ctrl1 (Extended Data Table 1). Plasmid

transfections were performed using the 4D-Nucleofector and the SE cell line 4D-Nucleofector

X S kit (Lonza) according to the manufacturer's instructions. Briefly, 0.5µg DNA was

transfected in 10^6 cells with 20µL SE solution in Nucleocuvette strips using the U2OS program

CM-104, and subsequent OHT treatments were performed 48h later. mCherry-LaminB1-10

plasmid was a gift from Michael Davidson (Addgene plasmid #55069;

http://n2t.net/addgene:55069 ; RRID:Addgene_55069).

Immunofluorescence

Transfected 53BP1-GFP DIvA cells were grown on glass coverslips and fixed with 4%

Paraformaldehyde during 15min at room temperature. After, 2 washes with PBS, the

permeabilization step was performed by treating cells with 0,5%Triton X-100 in PBS for

10min, then cells were blocked with PBS-BSA3% for 30min. Primary antibody against γH2AX

(Extended Data Table 2) was diluted in PBS-BSA3% and incubated with cells overnight at 4°C.

After 3 washes for 5min in PBS-BSA3%, cells were incubated with anti-mouse secondary

antibody (conjugated to Alexa594 or Alexa488, Invitrogen), diluted 1:1000 in PBS-BSA3%,

for 1h at room temperature. After a Hoechst 33342 (Invitrogen) staining (5µg/ml for 5min at

room temperature), Citifluor (AF1-25, Clinisciences) was used for coverslip mounting.

High content microscopy (QIBC)

U2OS, DIvA (AsiSI-ER-U20S)[20], AID-DIvA (AID-AsiSI-ER-Transfected or Dex-synchronized DIvA and U20S cells were plated in 96-well Cell Carrier Ultra plates (Perkin Elmer). Prior to the end of OHT, etoposide or bleomycin treatments, cells are treated with 10µM EdU (Invitrogen, C10340) at 37 °C for 15min. For γH2AX staining, cells are fixed and permeabilized as described above. For RAD51 staining, buffer II pre-extraction (20mM NaCl, 0.5% NP-40, 20mM HEPES at pH 7.5, 5mM $MgCl_2$, 1 mM DTT) was added to cells for 20min on ice followed by fixation in 4% Paraformaldehyde during 15min at room temperature and washed twice with PBS. Then EdU was labeled by Click-iT AlexaFluor 647 reaction (Invitrogen, C10340) according to the manufacturer's instructions (cells protected from light from now). Cells were washed twice for 5min in PBS-BSA3% and then blocked with PBS-BSA3% for 30min at room temperature. Primary antibodies targeting γH2AX or RAD51 (Extended Data Table 2) were diluted in PBS-BSA3% and incubated with cells overnight at 4°C. After 3 washes of 3min in PBS-BSA3%, cells were incubated with anti-mouse Alexa 488 secondary antibody (Invitrogen) diluted 1:1000 in PBS-BSA3%, for 1h at room temperature. After 3 washes of 3min in PBS, nuclei were labeled with 5µg/ml Hoechst 33342 (Invitrogen) for 30min at room temperature. The cells were then washed twice with PBS and stored at 4°C. γH2AX foci were further analyzed with an Operetta CLS High-Content Imaging System (Perkin Elmer) and Harmony software (version 4.9). For quantitative image analysis, ~20-80 fields per well were acquired with a 20X (for γH2AX circadian synchronization) or 40X objective lens to visualize >3000 cells per well in triplicate. Subsequent analyses were performed with Columbus software (version 2.8.2). Briefly, the Hoechst-stained nuclei were selected with B method, and the size and roundness of nuclei were used as parameters to eliminate false positive compounds. Then, the Find Spots D method was used (Detection sensitivity =0.2-0.4; Splitting coefficient =1; Background correction =0.5) to determine the

number, mean area and intensity of repair foci in each nucleus. For cell cycle analysis, the sum

of the Hoechst intensity and the mean of the EdU intensity were plotted in order to select G1,

S, and G2 cells.

Determination of the number (y axis) and size of γH2AX foci (mean area on the x axis) in each

nucleus has been used to infer DSB clustering (>8000 nuclei analyzed per sample) and the

resulting scatter plots are divided into four quadrants on the basis of the medians[10,11].

Percentages of cell populations with low number of γH2AX foci of large size (lower right-side

area of the scatter plot), called "cluster positive cells", or with high number of γH2AX foci of

small average size (upper left area of the scatter plot), called "cluster negative cells", are

determined.

DSB- induced rearrangement/translocation assay

AID-DIvA cells were treated as indicated and DNA was extracted using the DNeasy kit

(Qiagen). Illegitimate rejoining frequencies between different AsiSI sites, *t(MIS12;TRIM37)*

and *t(LINC00217;LYRM2)*, were analyzed by qPCR (in 3-4 replicates) using primers referenced

in Extended Data Table 1[23]. Results were normalized using two control regions (Norm1 and

Norm17) both far from any AsiSI sites and γH2AX domains (primers in Extended Data Table

1). Normalized translocation frequencies were calculated using the Bio-Rad CFX Manager 3.1

software.

WST-1 cell survival assay

Transfected AID-DIvA cells were plated in 96-well plates for each of the 3 conditions: -DSB,

+DSB, +DSB+repair. 48h later, cells were treated or not with OHT for 4h and further incubated

or not with IAA for 4h. Eight days after DSB induction, the medium is replaced by a mix of

100µl warmed medium + 10µl WST-1 ready-to-use solution per well and the cells were

incubated for 2h at 37°C. The tetrazolium salts from Premix WST-1 Cell Proliferation Assay

System (TAKARA) are cleaved by mitochondrial dehydrogenase in viable cells to produce

formazan dye, which is quantified by measuring its absorbance at 450nm (Multiskan GO

Microplate Spectrophotometer from Thermo Fisher Scientific). The percentage of viable cells

is normalized to the undamaged cells and compared to the control condition.

Clonogenic assays

Transfected AID-DIvA cells were seeded, in duplicate, at a clonal density in 10cm diameter

dishes and, 48h later, were treated or not with 4OHT for 4h and washed twice in pre-warmed

PBS and further incubated with auxin (IAA) for another 4h when indicated. Then, the medium

was replaced for all cell dishes. After 8 days at 37°C under a humidified atmosphere with

5%CO_2, cells were washed in PBS and stained with crystal violet (Sigma, V5265). Images of

stained colonies were obtained using the White Sample Tray for the ChemiDoc Touch Imaging

System (Bio-Rad) and colonies were counted using the Spot Detector plugin of Icy software,

Version 2.0.2.0 (icy.exe program). Only colonies containing more than 50 cells were scored.

The same protocol was applied to transfected U2OS and hTERT RPE1 cells treated with

etoposide or doxorubicin.

Western blots

DIvA cells were incubated in RIPA buffer (50mM Tris pH8.0, 150mM NaCl, 0.5% sodium

deoxycholate, 1% NP-40, 0.1% SDS) on ice for 20 min. Samples were further incubated with

250 units of Benzonase (Sigma, E1014) per million of cells for 10 min at room temperature,

then centrifuged at 13,000 rpm for 10 min at 4°C. The supernatants, containing soluble protein

extracts, were then mixed with SDS loading buffer and reducing agent, and incubated 10 min

at 70°C. Electrophoreses were performed using 4–12% NuPAGE Bis-Tris gels (Invitrogen) and

semi-dry western blotting on nitrocellulose membranes (Bio-Rad) was done with the Trans-

Blot Turbo System (Bio-Rad) according to the manufacturer's instructions. Membranes were

incubated in TBS containing 0.1% Tween 20 (Sigma, P1379) and 4% BSA during 1h at room

temperature for blocking, followed by overnight incubation at 4 °C using primary antibody

diluted in TBS-Tween-4% BSA. The corresponding mouse or rabbit horseradish peroxidase-

coupled secondary antibodies (Sigma, A2554 and A0545) were used at 1:10,000 to reveal the

proteins, using a luminol-based enhanced chemiluminescence HRP substrate (Super Signal

West Dura Extended Duration Substrate, ThermoScientific). Picture acquisition of the

membranes was done using the ChemiDoc Touch Imaging System (Bio-Rad) and pictures were

analyzed using Image Lab software (Bio-Rad). All primary antibodies used in this study are

detailed in Extended Data Table 2. Note: although different conditions and several human PER1

antibodies were tested, none of them resulted in a ~140kDa signal corresponding to the PER1

protein on the western-blots.

RT-qPCR

Total RNA was extracted from U2OS and DIvA cell lines before and after DSB induction using

homogenization with QIAshredder and the RNeasy kit (Qiagen) following the manufacturer's

instructions. RNA was then reverse transcribed to cDNA using the AMV reverse transcriptase

(Promega, M510F). qPCR experiments were performed to assess the levels of cDNA using

primers referenced in Extended Data Table 1. cDNA levels were then normalized with GAPDH

(Gene ID: 2597), TBP (Gene ID: 6908) and 18S (RNA18SN5, Gene ID: 100008588) cDNA

levels using the CFX Manager 3.1 software (Bio-Rad).

Resection assay

Measure of resection was performed as described in[53] with the following modifications.

Genomic DNA was extracted from transfected DIvA cells just after DSB induction using the

DNeasy kit (Qiagen) without any vortexing step. 1 µg of DNA was incubated 20min at 37°C

with 15 units of RNAseH1 (New England Biolabs) and first digested for 2h at 37°C with 35

units of AsiSI restriction enzyme (New England Biolabs). Then, 250ng of DNA were further

digested overnight at 37°C using 10 units of BanI restriction enzyme, that cuts at 200bp from

the DSB1, or of BanII restriction enzyme, that cuts at 1122bp from the DSB7 (Extended Data

Table 1). Both enzymes were heat inactivated for 20 min at 80°C. Digested and undigested

samples were analyzed by qPCR (10ng/well) using primers in Extended Data Table 1. ssDNA%

was calculated with the following equation: Percent of ssDNA= 1/((PUISSANCE(2;(Ct

Digested − Ct Non Digested) − 1) + 0.5))*100.

Proximity Ligation Assay (PLA)

PLA experiments were performed using Duolink® In Situ Detection assay from Sigma-Aldrich

according to the manufacturer's instructions. Primary antibodies used for PLA are detailed in

Extended Data Table 2.

High throughput γH2AX-LaminB1 PLA. siRNA-transfected or Dexamethasone-synchronized

cells were plated in 96-well Cell Carrier Ultra plates (Perkin Elmer). At the end of DSB

inducing treatment, cells are treated with 10μM EdU (Invitrogen, C10340) for 15min. Cells are

fixed with 4% paraformaldehyde during 15min at room temperature. Permeabilization step was

performed by treating cells with 0,5% Triton X-100 in PBS for 5min. Then EdU was labeled

by Click-iT Alexa488 reaction (Invitrogen, C10340) according to the manufacturer's

instructions. Cells were blocked with 1 drop of blocking solution provided with the Duolink®

PLA Probes (DUO92002 and DUO92004, Sigma-Aldrich) for 1h at 37°C in a humid chamber.

Primary antibodies targeting γH2AX and LaminB1 were diluted in the antibody dilution buffer

provided with the Duolink® PLA Probes and incubated with cells overnight at 4°C. After two

5min washes in Duolink® In Situ Wash Buffer A (DUO82049, sigma-Aldrich), cells were

incubated with anti-mouse plus and anti-rabbit minus probes for 1h at 37°C, under slow

rotation. After two washes in wash buffer A for 5min, cells were incubated for 30min at 37°C

in the ligation reaction mix from the Duolink® In Situ Detection Reagents Red (DUO92008,

Sigma-Aldrich). The cells were washed again twice in wash buffer A for 5min and incubated

for 1h40 at 37°C under slow rotation for the amplification step. Finally, after two 10min washes

in wash buffer B (DUO82049, sigma-Aldrich), nuclei were stained in the staining solution from

the Duolink® In Situ Microplate Nuclear Stain (DUO82064, Sigma-Aldrich) for 30min at 37°C

and cells were kept in the Anti-Fade solution (DUO82064, Sigma-Aldrich) until image

acquisition. For quantitative image analysis, images were acquired with a 40X objective lens to

visualize >3000 cells per well in duplicate. Subsequent analyses were performed with

Columbus software (version 2.8.2) to determine the numbers of PLA foci in each nucleus. G1,

S and G2 nuclei were selected based on EdU and DAPI staining distribution in all cells.

53BP1-LaminB1 and γH2AX-LaminB1 PLA. siRNA-transfected cells were seeded on 13mm

coverslips and treated for 4h with 4OHT before fixation in 4%PFA. Slides were then processed

as before, except that there were no EdU staining, and that at the end of the PLA procedure,

after the two 10min washes in wash buffer B, cells were then washed for 1min in 0.01% wash

buffer B and mounted on slides with Duolink® In Situ Mounting Medium with DAPI

(DUO82040, Sigma-Aldrich). Image acquisition was done on Leica DM6000 at 40X objective

lens to visualize >175 cells per experiment in triplicate. Subsequent analyses were performed

with Cell Profiler software (version 4.2.4) to determine the numbers of PLA foci in each

nucleus.

3D super resolution using Random Illumination Microscopy (RIM)

3D RIM two cameras setup. The 3D RIM home-made setup is coupled to an inverted

microscope (TEi Nikon) equipped with 100x magnification, a 1.49 N.A. objective (CFI SR

APO 100XH ON 1.49 NIKON) and two SCMOS cameras (ORCA-Fusion, Hamamatsu)

mounted in an industrial apochromatic alignment module (abbelight SA). Fast diode lasers

(Oxxius) with wavelengths centered at 488nm (LBX-488-200-CSB) and 561nm (LMX-561L-

200-COL) are used for all experiments. The bandpass emission filters in front of the two

respective cameras are FF01-514/30-25 for camera 1 and FF01-609/54-25 for camera 2. The

binary phase modulator (QXGA fourth dimension) conjugated to the image plane combined

with polarization elements are used to generate dynamic speckle on the object plane as

described in[47]. The synchronization of the hardware (Z-platform, cameras, microscope, laser

and SLM) is performed by an improved version of the commercial software INSCOPER.

Fixed 3D RIM acquisition and reconstruction. Two-color 3D RIM imaging was performed

with the 3D RIM system. The acquisition of 40 planes was done sequentially with a 120nm step

in the image plane. For each image plane, 400 speckles were used to increase the desired super-

resolved 3D resolution (95nm,95nm,200nm). The image reconstructions were performed with

the software (ALgoRIMhttps://github.com/teamRIM/tutoRIM). The Wiener filter used is 0.01,

the deconvolution parameter is 0.02 and the regularization parameter is 0.02 for the first channel

images. The Wiener filter used is 0.02, the deconvolution parameter is 0.03 and the

regularization parameter is 0.02 for the second channel images. The misalignment between the

two cameras and the residual chromatic aberrations are corrected using the SVI (Scientific

Volume Imaging) software.

3D+t two colors RIM super-resolution acquisition and reconstruction. 3D+t RIM movies

were performed on AID-DIvA cells expressing 53BP1-GFP and mcherry-LaminB1. 3D movies

of 47min with 40sec intervals were made (60 times). The whole cell acquisition time was 6

seconds with 48 random patterns optimized for live cells for each plane. Image reconstructions

were performed with the software (ALgoRIMhttps://github.com/teamRIM/tutoRIM). The

Wiener filter used is 0.01, the deconvolution parameter is 0.151 and the regularization

parameter is 0.08 for the first channel images. The Wiener filter used is 0.1, the deconvolution

parameter is 0.25 and the regularization parameter is 0.08 for the second channel images. The

misalignment between the two cameras and the residual chromatic aberrations are corrected

using the SVI (Scientific Volume Imaging) software.

3D film editing. Bleaching correction is performed after RIM reconstruction with the open-

source FIJI 546 software (https://imagej.net/software/fiji/) based on an exponential FIT from

the background. The FIJI 3D drift correction plugin is performed for 3D registration

(https://github.com/fiji/Correct_3D_Drift). Rendering of 3D+t movies was performed with the

VTK library implemented in ICY software (https://icy.bioimageanalysis.org/) from 3D CROP

on 550 areas of interest. Chromatism correction was performed by Scientific Volume Imaging

(SVI) software.

Colocalization analyses. The colocalization analyzer from the Scientific Volume Imaging

(SVI) software is used on fixed 3D RIM images to quantify the interactions of γH2AX nanofoci

with those of SUN1 and SUN2.

Quantitative immunocolocalization (ICQ). The intensity correlation quotient was defined in[78]

by the following equation for each voxel "i":

$$\left(I_i^{D1} - I_{avg}^{D1}\right)\left(I_i^{D2} - I_{avg}^{D2}\right)$$

For a random or mixed interaction, this number will tend towards 0, and for a dependent

correlation, it will tend towards +0.5. This parameter does not directly use the intensity of each

pair of voxels and has the advantage of eliminating the bias towards particularly high or too

low intensities.

Cross-correlation function CCF. The Van Steensel cross-correlation function (VSCF) CCF is

used to quantify interactions and has been described in[79]. It is obtained by calculating the

Pearson coefficient after shifting the second camera image over a distance of Δx voxels.

Thresholding was carefully tailored for each image to reject the 10^{th} percentile of the lowest

intensity value for both channels. The CCF was measured with the x-shift set to 940nm without

rotation and the resulting three graphs were averaged and plotted with a 95 percentil. The

Pearson coefficient is the classic equation below for each voxel "i":

1296
$$P = \frac{\sum\left(I_i^{D1} - I_{avg}^{D1}\right)\left(I_i^{D2} - I_{avg}^{D2}\right)}{\sqrt{\sum\left(I_i^{D1} - I_{avg}^{D1}\right)\left(I_i^{D2} - I_{avg}^{D2}\right)}}$$

with I_{avg}^{D1} and I_{avg}^{D2} the averages of camera 1 and camera 2 of the microscope.

For better visualization of the shape of the CCF function for each condition, we normalized the

CCF function between 0 and 1 with the following normalization calculation:

$P_{(0,1)} = \frac{(P - P_{min})}{(P_{max} - P_{min})}$, for Figure 3i and 3k or $P_{(min)=}(P - P_{min})$ for Extended Data Figure 6e.

Negative values for Δx indicate a shift in nm of the red image to the left, positive values indicate

a shift to the right. Non-random overlap results in a peak at $\Delta x=0$ and non-random exclusion

results in a dip at $\Delta x=0$. Uncorrelated distributions will not show any clear peak in the CCF.

1304

ChIP-qPCR, ChIP-seq and data analysis

ChIP-qPCR. ChIP experiments were performed in DIvA cells according to the protocol

described in[8,20] with 200μg of chromatin per immunoprecipitation. The quantity of primary

antibodies used is detailed in Extended Data Table 2. qPCR experiments were performed on

both IP and input samples to assess the percent of input DNA immunoprecipitated, using AsiSI-

induced DSBs and primers referenced in Extended Data Table 1. When indicated in the figure

legends, data are normalized to a control location devoid of DSB (Ctrl genomic locus) and could

be further expressed related to data obtained at DSB4 or in undamaged condition.

Library preparation and sequencing. Multiple ChIP experiment samples were pooled and

sonicated for 15 cycles (30 sec on, 30 sec off, high setting) with a Bioruptor (Diagenode), then

concentrated with a vacuum (Eppendorf). 10ng of purified DNA (average size 250–300bp) was

used to prepare sequencing libraries with the Next Ultra II Library Prep Kit for Illumina (New

England Biolabs) using the application note for "Low input ChIP-seq", and subjected to 75bp

single-end sequencing using Illumina NextSeq500 at the EMBL Genomics core facility

(Heidelberg, Germany).

ChIP-seq data processing. ChIP-seq data processing was performed as described[22]. In brief,

raw sequencing reads were aligned using bwa (https://bio-bwa.sourceforge.net/) to the human

reference genome (hg19) and then, were sorted, deduplicated, and indexed using samtools

(http://www.htslib.org/). Bigwig coverage tracks were subsequently extracted from the

processed bam files using bamCoverage from deepTools (HYPERLINK

https://deeptools.readthedocs.io/en/develop/), and were normalized by total read count to be

used for downstream analyses.

DSB Average profiles and boxplots. Differential coverage tracks representing the log2 ratio

between damaged and undamaged ChIP-seq samples were created using the bamCompare

function of deepTools on default settings. These bigwigs were then used as input for the

computeMatrix function of deepTools to calculate the coverage at DSBs using 200 bins

covering either 10kb or 1kb as indicated on Figures. These matrices were then processed using

a custom script in R with ggplot2.

Peak calling. Peak calling for ChIP-seq datasets (BMAL1, SUN1 and SUN2) was performed

using *MACS2* using -q 0.1 for BMAL1 resulting in 1776 peaks, and -q 0.005 for SUN1 and

SUN2 resulting in 27326 and 14571 peaks respectively. Endogenous DSBs were identified as

described in [11]. In brief, peaks were called from untreated pATM-ChIP-seq data[80] using macs2,

identifying 1,206 regions. A random set of regions was then generated from the gkmsvm R

package.

Gene Ontology Analysis. Genes for BMAL1 GO analysis were selected if the identified

BMAL1 peak was within a genes body or within its promoter region. The subsequent gene list

was   used   to   perform   gene   ontology   analysis   using   the *enrichGO* function

from *clusterprofiler* in R.

Peak annotation. The genomic locations of SUN1 and SUN2 peaks were identified by using

the *ChIPseeker* package in R, specifically the *annotatePeak* function to annotate the regions

and *plotAnnoPie* for subsequent visualization. To calculate the coverage of SUN1 at SUN2

regions, the *computeMatrix* function of deepTools was used using 200 bins over a 10kb

window. The matrix was then processed using a custom script in R with ggplot2.

1348

Hi-C and data analyses

Hi-C libraries. Hi-C experiments were performed in DIvA cells following transfection with

CTRL or PER2 siRNAs and with or without DSB induction as described in[80]. Briefly, 10^6 cells

were used per condition. Hi-C libraries were generated using the Arima-HiC+ for HiC (Arima

Genomics) by following the manufacturer's instructions. DNA was sheared using the Covaris

S220 to obtain an average fragment size of 350-400pb. Sequencing libraries were prepared on

beads using the Next Ultra II DNA Library Prep Kit for Illumina and Next Multiplex Oligos

for Illumina (New England Biolabs) following instructions from the Arima-HiC+ for HiC

(Arima Genomics). Hi-C data were processed as described in[25,80]see below.

Hi-C heatmaps. Hi-C reads were mapped to hg19 and processed with Juicer v2.0 using default

settings (https://github.com/aidenlab/juicer). Hi-C count matrices were generated using Juicer

at multiple resolutions: 100kb, 50kb, 25kb, 10kb, 5kb and 1kb. Hi-C heatmap screenshots were

obtained using Juicebox (https://github.com/aidenlab/Juicebox/wiki/Download).

Aggregate Peak Analysis (APA). APA plots of inter- or intra-chromosomal interactions of

DSBs and control regions were created using the APA program of Juicer tools (https://github.

com/aidenlab/juicer/wiki/APA) with a 10kb resolution -/+ 100kb.

Trans contact quantification. To determine inter-chromosomal interaction changes (in *trans*),

we built the whole-genome Hi-C matrix for each experiment by merging all chromosome-

chromosome interaction matrices using Juicer and R to obtain a genome matrix with 33kx33k

bin interactions for 100kb resolution. Interactions between bins inside damaged TADs

(240X240 for the 80 best-induced AsiSI DSBs) were extracted and counted for each condition,

then log2 ratio was calculated on normalized count (cpm), and plotted as boxplots or heatmaps

(https://github.com/LegubeDNAREPAIR/ATMcompD/blob/main/scriptR/Heatmap_D_trans.

R).

A/B compartment. To identify A and B main chromosomal compartments, the extraction of the

first Eigen vector of the correlation matrix (PC1) was done on the Observed/Expected matrix

at 500kb resolution using juicer eigenvector command. The resulting values were then

correlated with ATAC-seq signal in order to attributes positive and negative values to the A

and B compartment, respectively, on each chromosome. The Observed/Expected bins were then

arranged based on the PC1 values and aggregated into 21 percentiles, to visualize A-B

interactions on the siPER2 and siCTRL experiments (saddle plots), and differences were

computed between PER2 and control conditions for each percentile.

D-compartment. For D-compartment identification, we retrieved the first component (PC1) of

a PCA made on the differential observed over expected Hi-C matrix $log2\left(\frac{damaged}{undamaged}\right)$ at 100kb

resolution. Each matrix was extracted from the .hic files using Juicer and the ratio was

computed bin per bin. Pearson Correlation matrices were then computed for each chromosome,

and PCA was applied to each matrix. The first component of each PCA was then extracted and

correlated with the position of DSBs. A PC1 showing a positive correlation with DSBs was

then called D-compartment, and PC1 showing negative correlation with DSBs were multiplied

by -1. We were able to extract the D-compartment on several chromosomes for +DSB/-DSB[25].

D-compartment (first component of the PCA) was converted into a coverage file using

rtracklayer R package. For analysis of DSB responsive gene expression by RT-qPCR, we used

the previously determined D compartment genes as described in[25].

1392

1393 **Contact for Reagent and Resource Sharing**

1394 Further requests for reagents and resources should be asked to the lab PI, Gaëlle Legube

1395 (gaelle.legube@univ-tlse3.fr). DIvA cell lines are subjected to an MTA with the CNRS.

1396

a

siRNA

CTRL   PER2

γH2AX

chr2

siRNA PER2/siRNA CTRL (+DSB)

HiC read count (+DSB)-(-DSB)

-20   +20

b

HiC read count

0 20 40 60

siRNA   CTRL   PER2

-DSB   +DSB   -DSB   +DSB
DSB +/-100kb   DSB +/-100kb   DSB +/-100kb   DSB +/-100kb

DSBs (80)   FC = 1.08   FC = 3.49   FC = 1.34   FC = 3.85

random sites (80)   FC = 0.84   FC = 0.89   FC = 0.95   FC = 0.99

c

Trans interaction of each damaged domains with the other damaged domains (80 DSBs)

8e-26   *2e-4*

HiC read count Log2(+DSB/-DSB)

siRNA   CTRL   PER2   CTRL   PER2

80 DSBs   no DSBs (80 randoms)

d

Chr2p

γH2AX

D Comp siCTRL

D Comp siPER2

CEV (+DSB-DSB) PC1   read count

10   80   MB

e

t (MIS12;TRIM37)

Translocation frequency (normalized to CTRL+DSB)

0.015 *0.015* *1e-3* *0.0092*

siRNA   CTRL SUN1 SUN2 NUP153 RNF4

f

t (MIS12;TRIM37)

Translocation frequency (normalized to CTRL+DSB)

0.88
0.72

siRNA   CTRL SUN1 PER1+SUN1 PER2+SUN1

g

t (MIS12;TRIM37)

Translocation frequency (normalized to CTRL+DSB)

0.72
0.42

siRNA   CTRL PER2 SUN1 PER2+NUP153 SUN1+NUP153

h

t (MIS12;TRIM37)

Translocation frequency (normalized to CTRL+DSB)

0.74

siRNA   CTRL PER2 RNF4 PER2+RNF4