Distinct guard cell specific remodeling of chromatin accessibility during abscisic acid and CO₂ dependent stomatal regulation

Authors
Charles A. Seller[1*] and Julian I. Schroeder[1*]

Affiliations
1. School of Biological Sciences
Cell and Developmental Biology Department
University of California San Diego,
La Jolla, CA 92093-0116
*Corresponding authors. Email: cseller@ucsd.edu; jischroeder@ucsd.edu

Classification: Plant Biology

Keywords: Guard cells, stomata, chromatin, plant hormones, abscisic acid, transcription factors, carbon dioxide

1 **Abstract**

In plants, epidermal guard cells integrate and respond to numerous environmental signals to
control stomatal pore apertures thereby regulating gas exchange. Chromatin structure controls
transcription factor access to the genome, but whether large-scale chromatin remodeling occurs
in guard cells during stomatal movements, and in response to the hormone abscisic acid (ABA)
in general, remain unknown. Here we isolate guard cell nuclei from *Arabidopsis thaliana* plants
to examine whether the physiological signals, ABA and CO_2, regulate guard cell chromatin
during stomatal movements. Our cell type specific analyses uncover patterns of chromatin
accessibility specific to guard cells and define novel cis-regulatory sequences supporting guard
cell specific gene expression. We find that ABA triggers extensive and dynamic chromatin
remodeling in guard cells, roots, and mesophyll cells with clear patterns of cell-type specificity.
DNA motif analyses uncover binding sites for distinct transcription factors enriched in ABA-
induced and ABA-repressed chromatin. We identify the ABF/AREB bZIP-type transcription
factors that are required for ABA-triggered chromatin opening in guard cells and implicate the
inhibition of a set of bHLH-type transcription factors in controlling ABA-repressed chromatin.
Moreover, we demonstrate that ABA and CO_2 induce distinct programs of chromatin
remodeling. We provide insight into the control of guard cell chromatin dynamics and propose
that ABA-induced chromatin remodeling primes the genome for abiotic stress resistance.

19
20
21 **Significance statement**

22
Specialized leaf cells called guard cells integrate environmental cues to optimally control the
size of microscopic stomatal pores. The hormone abscisic acid (ABA), a key regulator of plant
drought responses, and changes in atmospheric CO_2 concentration are signals that control
stomatal aperture size, but whether these signals also regulate genome packaging into
chromatin is unknown. Using guard cell specific chromatin profiling we uncovered regulatory
DNA sequences driving specific gene expression in this cell-type. We also discovered that ABA
triggers extensive and persistent changes to chromatin structure in guard cells. Unexpectedly,
exposure of plants to elevated atmospheric CO_2 had only minimal impact on chromatin
dynamics. Furthermore, we identified the specific transcription factors that regulate ABA-
induced chromatin dynamics in guard cells.

INTRODUCTION

Organisms evolved mechanisms that connect the activity of their genome to the conditions in their environment. Developmental and environmental signals can impact the genome by regulating transcription factor (TF) binding to cognate DNA sequences known as cis-regulatory elements (CRE) (1). In eukaryotes, this regulation occurs in the context of chromatin where nucleosomes can impede transcription factor binding to target sequences (2, 3). Consequently, the remodeling of chromatin structure to allow access of transcription factors to target DNA is thought to be a key step in gene regulation. Plant hormones are key signaling molecules controlling numerous aspects of plant life that coordinate genome activity with environmental conditions (4, 5). Recent studies have provided insight into how chromatin structure changes during plant development (6–12) and in response to environmental and hormonal stimuli (13–15).

Here we focus on the abscisic acid (ABA) signal transduction pathway to explore the connection between hormone signaling and chromatin structure in plants. ABA is a major plant stress hormone that accumulates in cells and tissues experiencing abiotic stress, especially those linked to plant water-status (16–18). ABA signals through a well understood core module consisting of PYR/PYL/RCAR ABA receptor proteins, PP2C protein phosphatases, and SnRK2 protein kinases (19–21). Upon ABA binding, PYR/PYL/RCAR receptor proteins inactivate PP2C phosphatases resulting in the derepression of SnRK2 kinases. ABA-activated SnRK2 kinases then phosphorylate and thereby directly regulate downstream proteins including numerous transcription factors (22–28). Consequently, ABA triggers the differential expression of thousands of genes in the *Arabidopsis thaliana* genome (4, 29–31). However, it remains unclear, especially at the genome scale, whether ABA signaling reshapes chromatin structure, and how ABA-regulated TFs might function in the context of chromatin.

One well known action of ABA is to trigger the closure of the stomatal aperture (17). Stomata

are small pores on the surface of the leaf that mediate gas exchange and are formed by a pair

of specialized epidermal cells known as guard cells. Guard cells perceive and respond to

multiple environmental and hormonal cues in order to optimally regulate the size of the stomatal

pore (32–34). For instance, some signals such as elevated atmospheric carbon dioxide (CO_2),

ABA, and the immune elicitor Flg22 trigger stomatal closure, while reduced CO_2, light, and heat

trigger stomatal opening. Because of their exquisite environmental reactivity, guard cells are an

excellent model cell system to investigate how different physiological signals regulate chromatin

structure in a cell-type specific manner. Although the recent application of single-cell RNA-seq

to *Arabidopsis* has expanded our understanding of gene regulation, these studies often focus on

plate-grown seedlings (35, 36) and/or resulted in datasets with limited representation of guard

cells (37, 38). To apply epigenomic techniques to the question of how guard cells respond to

different stimuli new protocols are needed that can isolate large numbers of pure guard cells

from soil grown plants, a more relevant setting for physiological studies.

In this paper, we describe the development of an approach to purify guard cell nuclei and by

deploying this method, uncover how guard cell chromatin structure changes in response to

different stimuli that drive stomatal movements. We profiled the chromatin and transcriptional

reprogramming in response to ABA in three different developmental contexts – roots, mesophyll

cells, and guard cells. We map thousands of loci that gain or lose chromatin accessibility in

response to ABA and link these regions to co-regulated transcripts, uncovering a critical role for

chromatin dynamics in controlling ABA-dependent transcription. Furthermore, we show that

genome-wide and persistent changes to chromatin structure accompany ABA-induced stomatal

closure and that four related bZIP transcription factors known as ABRE Binding Factors (ABFs)

are required for initiating the majority of ABA-induced chromatin opening. In contrast, we

implicate a family of related bHLH-type TFs known as ABA-Kinase Substrates (AKSs) in the

maintenance of open chromatin upstream of ABA-repressed genes. Finally, we show that in

guard cells, changes to atmospheric CO_2-concentration trigger distinct and more limited

chromatin remodeling and gene regulatory programs from those mediated by abscisic acid.

RESULTS

ABA induces rapid and genome-wide chromatin remodeling in roots

As an initial test of the relationship between ABA and chromatin we focused on *Arabidopsis*

thaliana seedlings. To measure changes to chromatin structure we combined fluorescence

activated nuclei sorting (FANS) with the Assay for Transposase-Accessible Chromatin with

sequencing (ATAC-seq), a quantitative measurement of chromatin accessibility across the

genome (39, 40). 50,000 nuclei per sample were isolated using FANS (Fig. S1A). We called

25,753 ATAC-seq peaks across all whole seedling and seedling root libraries (Datasets S1 and

S2). Genomic regions coinciding with these peaks are known as Accessible Chromatin Regions

(ACRs) and predominantly aligned with upstream regulatory regions in the genome (Fig. S1B-

E). In contrast, ATAC-seq reads derived from purified genomic DNA did not cluster into defined

peaks (Fig. S1B,C). We next treated whole seedlings with ABA and profiled chromatin

accessibility after 4 hours. Differential analysis revealed hundreds of regions that significantly

gain and lose accessibility in response to ABA (Fig. 1A, Dataset S3).

To capture the dynamics of ABA-induced chromatin remodeling we focused on roots which can

rapidly take up ABA from their surroundings (41, 42). Using FANS, we isolated root nuclei at

three different timepoints following ABA treatment (after 45, 120, and 240 minutes) and

performed ATAC-seq. Differential analysis revealed that ABA triggers time-dependent changes

in chromatin accessibility in roots (Dataset S3). Heatmaps centered over these ABA regulated

ACRs illustrate the progressive impact of ABA on chromatin structure, with more regions of the

genome changing in ATAC-signal over time (Fig. 1B,C). We observed significant changes to

chromatin after only 45 minutes of ABA treatment, with 163 ACRs located in upstream

regulatory regions gaining accessibility and 64 losing accessibility (Fig. 1C,E). After 4 hours,

ABA had activated 1293 ACRs and repressed 1119 ACRs (Dataset S3). Interestingly, we found

that ABA-regulated ACRs tend to be located further away from transcription start sites (> 1kb)

than static regions (Fig. 1D). ABA-induced ACRs were associated with downstream genes

enriched for gene ontology (GO) terms such as "response to water deprivation", "response to

osmotic stress", and "response to abscisic acid" (Dataset S6).

To explore the role of specific transcription factors in orchestrating these changes we performed

DNA motif analysis using a set of motifs empirically generated by the in vitro DNA binding assay

DAP-seq (43). We found that motifs recognized by ABF/AREB transcription factors such as

ABF1/2/3/4 and ABI5 were highly enriched in ABA-induced ACRs (Fig. 1F, example ABF1, p =

1e-63, Dataset S7). In contrast, we found that ABA-repressed ACRs showed enrichment for

motifs recognized by the related bHLH type transcription factors AKS1, AKS3, and bHLH80

(Fig. 1F, example AKS3, p = 1e-32, Dataset S7). Notably, both ABF1/2/3/4 and AKS1/2/3

proteins are direct targets of ABA-regulated SnRK2 kinases (22, 24). Previous research has

shown a role for ABA-dependent inhibition of bHLH-type AKS transcription factors in the

transcriptional repression of the guard cell K^{+} channel *KAT1* (24, 44). Interestingly, our findings

suggest a larger role for the AKS clade of closely related bHLH proteins in the regulation of

ABA-repressed genes. We conclude that ABA can trigger rapid and genome-wide chromatin

remodeling in root tissue.

Revealing guard cell specific patterns of chromatin accessibility

Encouraged by our findings in whole seedlings and roots we next decided to investigate the

effect of ABA on chromatin in mature guard cells. We first developed a FACS-based strategy to

purify guard cell nuclei from soil-grown plants. Nuclei were labeled by expressing the histone

H2B fused to GFP under the control of the guard cell *pGC1* promoter (45). We first confirmed

the guard cell specificity of the H2B-GFP fusion in our transgenic lines using fluorescence

microscopy (Fig. 2A) and confirmed that ABA treatment did not alter the guard cell specific

expression (Fig. S2A). We first isolated nuclei from leaf epidermal samples harvested from

GC1:H2B-GFP plants. DAPI stained nuclei preparations were then analyzed using FACS (Fig

S2B). We could define a clearly separated GFP-positive population among all detected nuclei

(Fig. 2B, Fig. S2C). In contrast to many other cell types in the *Arabidopsis* leaf, guard cells are

diploid, and GFP-positive nuclei from *pGC1:H2B-GFP* expressing plants had a 2n DNA content

while GFP-negative nuclei were distributed among higher ploidy levels (Fig. 2C). Approximately

5% of GFP-positive particles (41 out of 5037 total from a single FACS run) had DAPI content

greater than 2N, however we sorted only 2N nuclei for downstream applications. This method

allowed us to obtain ~30,000 guard cell nuclei with an estimated purity of 98% (Fig. S2D) from

the leaves of 40-50 plants. To assay guard cell chromatin structure, nuclei isolated using this

method were used to generate ATAC-seq libraries. The resulting libraries were of high quality

with clear concentration of peaks in annotated upstream regulatory regions (Fig. S2E),

enrichment of open chromatin fragments (< 100 bp) immediately upstream of transcription start

sites (Fig. S2F), strong correlation between biological replicates (Fig. S2G), and high FRiP

(fraction of reads in peaks) scores (Dataset S1).

In parallel, we generated ATAC-seq samples from root nuclei and from FANS-isolated

mesophyll nuclei labeled by expression of H2B-GFP from the promoter of the mesophyll

expressed Rubisco small subunit 2B gene (Fig. S3A-C) (6). For each tissue/cell-type we

compared ATAC-seq libraries derived from three biological replicates. We identified 30,985

ATAC-seq peaks across all samples (Dataset S2) and called cell-type enriched peaks using

pairwise differentially accessibility analysis. Although most ATAC-seq peaks did not differ

significantly among cell types, we found 3272 guard cell enriched ACRs, 1472 mesophyll

enriched ACRs, and 3364 root enriched ACRs (Dataset S2). Heatmaps centered over these

regions highlight characteristics of cell/tissue specificity in the observed patterns of chromatin

accessibility (Fig. 2D, Fig. S4A). Importantly, biological replicates derived from the same

tissue/cell-type clustered together in principal component analysis (PCA) underscoring the

robustness of this approach (Fig. 2E).

To evaluate the relationship between guard cell enriched chromatin accessibility and gene

expression we performed RNA-seq on both whole leaves and guard cell-enriched samples. We

found that regions with high chromatin accessibility specifically in guard cells were associated

with elevated transcript levels from the adjacent downstream genes, indicating that we are

identifying functional cis-regulatory elements (CREs) (Fig. 2F). Although the correlation between

overall promoter chromatin accessibility and downstream transcript level was weak in guard

cells (Pearson's r = 0.19, Fig. S4B), we found that differentially accessible ACRs showed higher

correlation with downstream gene differential expression (Pearson's r = 0.55, Fig. S4C).

Examination of individual gene loci like the guard cell expressed gene *KAT1* (46–48) illustrated

the connection between increased upstream accessibility and increased transcript levels (Fig.

2G and Fig S4D). By contrast, the mesophyll expressed gene *AOC1* had high chromatin

accessibility in mesophyll cells, but minimal upstream ATAC-seq signal in guard cells (Fig. 2G).

TF binding motif analysis on guard cell enriched ACRs returned de novo motifs with strong

similarity to motifs recognized by known stomatal lineage transcriptional regulators such as the

bHLH proteins SPCH, MUTE, and FAMA as well as EDT1 and DOF-type TFs (49–53) (Fig. 2H

and Dataset S7). In contrast, we find that WRKY-type TF binding motifs are enriched in regions

that are significantly less accessible in guard cells compared to roots and mesophyll cells (Table

S7). However, these motifs are recognized by large transcription factor families containing many

transcription factors that are not specific to guard cells, for instance SPCH and MUTE are active

only early during stomatal lineage development (49, 50). By connecting guard cell specific

ACRs with downstream guard cell enriched transcripts we were able to define a set of 827

putative regulatory elements preferentially active in guard cells (Table S4). To further confirm

that our approach identified functional cis-regulatory regions, we generated transcriptional

reporter constructs expressing nuclear localized GFP under the control of the guard cell specific

ACRs upstream of the malate-transporter encoding gene *ALMT6* (54) and the wax biosynthesis

enzyme-encoding gene *AAE9* (55). Both reporter constructs produced guard cell specific GFP

signal in transgenic *Arabidopsis* leaves (Fig. 2I)*.* In summary, we have developed a robust

protocol for isolating pure populations of guard cell nuclei allowing us to discover cis-regulatory

regions that are active in mature guard cells.

ABA induces persistent chromatin remodeling in guard cells during stomatal closure

We next asked whether epigenomic reprogramming occurs during ABA-induced stomatal

closure. Soil grown 6-week-old plants were sprayed with 50 uM ABA or control solution. After 3

hours, we confirmed a stomatal response by infrared photography (Fig. 3A). To determine if our

ABA treatment led to significant transcriptional changes we performed RNA-sequencing on both

whole leaf samples (Fig. S5A) and guard cell enriched samples (Fig. 3B). We observed that

ABA triggered the differential expression of thousands of transcripts in both whole leaves and

guard cells (2142 upregulated and 1574 downregulated in guard cells). As expected, ABA-

induced genes are enriched for the pathways "response to water deprivation" (p_value = 1.7e-

22), "response to abscisic acid" (pvalue = 4.47e-18), and "response to salt stress (Fig. S5D).

Although most of the transcriptional changes induced by ABA were shared between guard cells

and whole leaves, we found that 22.7% (488) of ABA-induced transcripts and 42.7% (673) of

ABA-repressed transcripts were specific to guard cells (Fig. S5C, Dataset S5). For instance,

ABA-regulated expression of the major facilitator protein-encoding gene *AT2G37900* and the

gene *AAE9* was only evident in guard cells (Fig. S5E).

To evaluate if ABA changed chromatin accessibility and whether these changes are sustained

over time following a single treatment, we isolated guard cell nuclei at 4 hours and 24 hours

after spraying (Fig. 3A). We generated ATAC-seq libraries from three biological replicates per

time point. PCA analysis showed that samples clustered together according to treatment (Fig.

S6A). Differential chromatin accessibility analysis found that ABA extensively reshapes

chromatin structure in guard cells with 2323 ACRs gaining accessibility and 1657 ACRs losing

accessibility across the treatment time course (Fig. 3C, Fig. S7A and B, and Dataset S3). We

next analyzed the temporal characteristics of this response by comparing the ATAC-seq

datasets from the 4-hour and 24-hour timepoints. Our results allowed us to define three

categories of ABA-regulated ACRs – early, sustained, and late (Fig. 3D). Strikingly, we found

that 1093 of the 2079 (52.6%) ACRs opened by ABA at 4 hours remained open after 24 hours.

Of the 975 ACRs closed by ABA at 4 hours 594 (61%) remained closed at 24 hours.

Additionally, we found 230 ACRs gained and 682 ACRs lost accessibility only 24 hours after

ABA treatment (Fig. 3D). Interestingly, we found differences in downstream gene GO term

enrichment between early and late ACRs. For example, the terms "response to light" and

"defense response" were strongly enriched in ABA-repressed ACRs only at 24 hours after ABA

treatment (Table S6).

Consistent with our observations in seedling roots (Fig. 1D), we found that ABA-regulated

ATAC-seq peaks in guard cells reside further away from transcription start sites (TSS) than non-

regulated peaks (> 1kb), indicating that distal regulation may be a characteristic of ABA-

controlled CREs (Fig. S6B). ABA-induced ACRs in guard cells are clearly associated with

downstream genes that are upregulated transcriptionally by ABA, while ABA-repressed ACRs

are associated with genes downregulated by ABA (Fig. 3E). Indeed, ABA-induced changes in

chromatin accessibility were correlated with ABA-induced differential expression of downstream

genes (r = 52, Fig. S6C). Examination of individual genes like the ABA-upregulated *MYB121*

and the ABA-downregulated *MYB60* (56) illustrate the connection between upstream chromatin

accessibility and transcript levels (Fig. 3F). Interestingly, we found that ABA modifies chromatin

accessibility upstream of many known regulators of stomatal conductance including *KAT1*,

NCED3, and *HT1* (57) as well as many other genes with no known stomatal functions (Fig.

S6D-K, Table S3).

We next performed TF binding motif analysis on early, sustained, and late ABA-regulated ACRs

using a set of DAP-seq derived motifs (43) restricted to those TFs with guard cell expression in

our RNA-seq datasets. Both early and sustained ABA-induced ACRs had strong enrichment for

motifs recognized by ABRE- binding TFs including ABI5, ABF1,2, and 3 (Fig. 3G). Interestingly,

we found that the motifs recognized by CBF2, CBF3, and MYB61 (58–60) were preferentially

enriched in early ABA-induced ACRs. ABA-repressed ACRs showed strong enrichment for

motifs recognized by a clade of related bHLH-type TFs (Fig. S6L) including AKS1, AKS3, and

bHLH80 in both sustained and late categories (Fig. 3H). In contrast, MYB96 motifs were

enriched only in late appearing ABA-repressed ACRs (Fig. 3H). Collectively, our results

demonstrate that ABA can induce persistent changes to chromatin in guard cells and expose

the temporal regulation of different TFs during this response.

By analyzing published genome-wide DAP-seq datasets (39, *56*) we found strong ABF2 DAP-

seq signal coincident with ABA-induced ACRs in guard cells (Fig. 3C). DAP-seq signals for

ABF1, ABF3, and ABF4 were also enriched over regions that gain accessibility in response to

ABA, although to a lesser extent than ABF2 (Fig. S6M). In contrast, ABA-repressed ACRs

contained AKS1 DAP-seq binding sites (Fig. 3C). Additionally, unbiased de novo motif discovery

(62) returned motifs similar to ABF1 and AKS3 binding sites as the most enriched in ABA-

activated and ABA-repressed ACRs, respectively (p = 1e-119 and p = 1e-57). Metaplots of

ATAC-seq signal centered over peaks containing these motifs further supported our observation

that ABA elevates chromatin accessibility at ABF binding sites and represses chromatin

accessibility at AKS binding sites (Fig. S6N). Interestingly, ABA closed chromatin upstream of

AKS3 and decreased *AKS3* transcript levels in guard cells (Fig. 3I). We also observed that

GFP-AKS3 signal decreased in ABA-treated guard cells when expressed from its native

promoter (Fig. 3J).

Furthermore, by comparing ABA-induced ACRs from guard cells, mesophyll cells (Fig. S3E-G),

and root cells (Fig. 1) we uncovered significant tissue/cell-type specificity in this response (Fig.

3K, Fig S7C-L). Interesting, we found that ABA had a much larger impact on chromatin

accessibility in guard cells than in the other cell types with more ABA regulated regions (2079

ACRs compared with 1293 in roots and 926 in mesophyll) and larger fold changes in

accessibility (Dataset S3). In conclusion, we show that ABA drives large scale changes to

chromatin structure in guard cells and implicate major roles for the activation of ABFs and

inhibition of AKSs in this response.

ABF transcription factors are required for ABA-induced chromatin opening in guard cells

The mechanisms that cause ABA-specific chromatin opening are unknown. Given the strong

enrichment of ABF binding motifs in ABA-activated ACRs, and their coincidence with ABF *in*

vitro binding sites, we wondered if these TFs might be required for reshaping chromatin during

ABA-induced stomatal closure. To test this, we used the *abf1/2/3/4* (*abfx4*) quadruple mutant

which lacks four related ABF proteins that are prominent during vegetative growth (25, 26). We

first performed RNA-seq following ABA treatment on leaves from 6-week-old *abf1/2/3/4* (*abfx4*)

mutant plants (Fig. S8A). Of the 2257 transcripts that are upregulated by ABA in wild type

leaves, 1897 (~84%) depend on ABF1/2/3/4 (Fig. S8B). Interestingly, although ABA increased

ABF1/2/3/4 transcript levels in guard cells we did not find that ABA altered chromatin

accessibility in the upstream regions near the corresponding genes (Fig. S8C). We conclude

that ABF1/2/3/4 are required for the bulk of ABA-induced gene expression in mature leaf tissue.

Plants lacking *ABF1/2/3/4* (*abfx4*) are highly sensitive to drought stress and lose water faster

than wild type plants in detached leaf assays (26). Consistent with this, we observed that *abfx4*

mutant plants exhibited higher steady-state stomatal conductances than *Col-0* control plants

(Fig. S8D).

Under non-stressed conditions, guard cells have a higher ABA concentration than other leaf cell

types (42, 63). Recent studies have provided evidence for basal ABA signaling in guard cells

and defined important roles for basal ABA in stomatal function (64, 65). By generating guard cell

FANS lines in an *abfx4* mutant background (Fig. S9A) we were able to assess the impact of the

loss of ABF1/2/3/4 on guard cell chromatin in non-stressed plants. In the absence of ABA

treatment, we observed minimal differences in genome-wide patterns of chromatin accessibility

between *WT* and *abfx4* guard cells (Fig S9B, C and Dataset S3) suggesting that basal ABA

signaling in guard cells may regulate mainly post-transcriptional stomatal closing mechanisms

but does not significantly regulate chromatin structure.

Next, we asked if the ABF1/2/3/4 are required for the ability of ABA to reshape chromatin

structure. Strikingly, loss of *abf1/2/3/4* strongly impaired ABA-induced chromatin opening in

guard cells but had less impact on ABA-triggered chromatin closing (Fig. 4A, Dataset S7). Of

the 2079 ABA-induced ACRs, 1539 (~74%) required ABF1/2/3/4 for gained accessibility (Fig.

4B). ABF-dependent ABA-induced ACRs were strongly enriched for ABREs (example ABF1, p

= 1e-86), while ABF-independent ABA-induced ACRs were most strongly enriched for CBF1/2/3

motifs (example CBF1, p = 1e-16) (Dataset S7). Examination of individual loci like the ABA-

activated genes *DTX37* and *HSFA6B* revealed a close connection between abf-dependent

increases in upstream chromatin accessibility and abf-dependent transcription (Fig. 4C).

We next performed TF motif analysis on ABA-regulated accessible chromatin regions in both

WT and *abfx4* mutant guard cells. As expected, *abfx4* mutants were depleted for ABRE motif

enrichment in ABA-induced ACRs but notably retained enrichment for AKS1, AKS3, and

bHLH80 motifs in ABA-repressed ACRs (Fig. 4D, Dataset S7). Additionally, although most TF

binding sites lost enrichment in ABA-induced ACRs in the absence of ABF1/2/3/4, we found that

enrichment for motifs bound by CBF1/2/3 was preserved. TF binding motifs are often

information poor and the presence of a motif at a genomic location does not always lead to TF

action. Analysis of published ABF1/2/3/4 binding assays conducted on naked genomic DNA

(61) uncovered 11334 binding sites, but only 1315 resided within ABA-induced ACRs in guard

cells (Fig. 4E). However, ABA-induced chromatin opening at these ACRs containing ABF-

binding sites was strongly impaired in *abfx4* mutant guard cells (Fig. 4F). We reasoned that

chromatin structure may determine which *in vitro* ABF binding sites are functional. To test this

idea, we examined if ABF DAP-seq peaks found within ABA-induced ACRs could predict ABA-

activated transcription. Using our guard cell RNA-seq data, we found that ABF binding sites that

reside within regions of ABA regulated open chromatin were more strongly associated with

downstream ABA-induced genes than were an equal number of random ABF-binding sites (Fig.

4G).

To examine the temporal characteristics of ABF-dependent chromatin remodeling we focused

on roots where ABA can reshape chromatin by 45 mins (Fig. 1C). We performed ATAC-seq on

isolated *abfx4* mutant root nuclei over an ABA treatment time course and found that ABA fails to

drive progressive chromatin opening in this background (Fig. S10). In *abfx4* mutant roots, 45

mins of ABA treatment induced only 2 ACRs compared to 163 in wild type roots. By 2 hours of

ABA treatment, only 27 ABA-induced ACRs were present in *abfx4* mutants compared with 601

in wild type (Fig. 4H , Fig. S10A, and Dataset S3). However, ABA repressed a similar number of

root ACRs in the absence of ABF1/2/3/4 (Fig. 4H and Fig. S10A). Although our data support a

general requirement for ABFs in ABA-induced chromatin opening in roots and guard cells, we

do find that some guard cell specific ABA-induced ACRs are ABF-dependent (Fig. S10D,E).

Collectively, these results demonstrate that the ABF1/2/3/4 transcription factors are required for

triggering genome-wide chromatin opening in response to ABA.

ABA and changes to atmospheric CO_2 concentration induce distinct chromatin

remodeling programs in guard cells

Because we observed that ABA has extensive effects on chromatin accessibility in guard cells,

we wondered whether chromatin structure may respond to other stimuli that drive stomatal

movements. Here we focus on changes to atmospheric CO_2 concentration. Exposure of plants

to low CO_2 stimulates stomatal opening while high CO_2 drives stomatal closure (Fig 5A). The

exact role of ABA signaling during CO_2 induced stomatal responses is unclear and is a matter of

some debate (64–66). Recent studies suggest that although high-CO_2 induced stomatal closure

requires basal ABA signaling it does not involve further SnRK2 kinase activation (64, 65).

Comparing how these signals impact genome activity could provide new insight into whether

and how the mechanisms underlying guard cell responses to ABA and CO_2 may differ.

To generate low and high CO_2 guard cell ATAC-seq libraries we exposed 6-week-old plants to

either 100 ppm CO_2 or 1000 ppm CO_2 for 4 hrs, two treatments that induce robust stomatal

responses (Fig 5B). Surprisingly, in contrast to the extensive effect of ABA on chromatin

accessibility in guard cells (Fig. 5C), we observed smaller changes in response to low CO_2 (Fig.

5D), and minimal changes following treatment with high CO_2 (Fig. 5E). Following exposure of

plants to 100 ppm CO_2, 329 ACRs gained accessibility and 487 ACRs lost accessibility while

treatment with 1000 ppm CO_2 induced only 11 ACRs and repressed 79 ACRs (Dataset S3).

Heatmaps of ATAC-seq signal at ABA and Low CO_2-regulated regions highlighted the specificity

of these responses (Fig. 5F, Fig. S7A,B). PCA results comparing our guard cell ATAC-seq

datasets from different conditions further support these findings and revealed little difference

between control and high CO_2 treatments (Fig. 5G). Examination of individual loci highlights the

specificity of these responses, for instance chromatin opening upstream of *ERF113* only

occurred in response to ABA while the upstream region of the PP2C-encoding *AT2G25070*

gains accessibility only in response to low CO_2 (Fig. S11B-E).

Analysis of the set relationships among ABA, low CO_2, and high CO_2 regulated ACRs revealed

little overlap suggesting that these stimuli control distinct cis-regulatory elements in the genome

(Fig. S11A). We did find overlap between ABA-induced and low CO_2-repressed ACRs and

between ABA-repressed and low CO_2-induced ACRs. ABA-repressed regions had elevated

average ATAC-seq signal in guard cells isolated from plants exposed to 100 ppm [CO_2] (Fig.

5F). For instance, ABA-repressed chromatin accessibility upstream of *ALMT6* while low-CO_2

treatment enhanced accessibility (Fig. S11D). Interestingly, ABA repressed chromatin

accessibility upstream of *HT1*, *BLUS1*, and *CBC2* (67)*,* all key positive regulators of low CO_2-

and blue light-induced stomatal opening, while high CO_2 had no effect (Fig. 5H). These data

indicate that chromatin remodeling represents an important cross talk point for these stimuli,

ensuring robust stomatal closing or opening, depending on the physiological conditions. DNA

binding motif analysis further emphasized the distinct regulatory impacts of changes in CO_2

concentration and ABA on the genome (Table S7). In contrast with ABA-induced ACRs, we did

not find enrichment of ABF binding motifs in low or high CO_2 regulated chromatin (Fig. 5I).

Instead, low CO_2 gained ACRs were enriched for motifs recognized by C2C2dof-type TFs which

were oppositely enriched only in ABA-repressed ACRs. Collectively, our data demonstrate how

different signals that control stomatal movement deploy distinct gene regulatory programs in

guard cells.

DISCUSSION

Abscisic acid triggers changes in gene expression, but whether this transcriptional response

depends on chromatin remodeling remained unknown (5). Here, we reveal massive ABA-

induced chromatin remodeling pointing to an orchestrated reprogramming of the *Arabidopsis*

epigenome by specific sets of transcription factors. We identify regulatory regions of the

genome specifically active in guard cells and uncover how the accessibility of these sequences

is modified by signals that control stomatal aperture size.

Guard cell regulation of stomatal aperture size is essential for plant survival under drought

stress. Such regulation requires that the guard cells forming the stomatal pore integrate

numerous physiological and environmental cues. In this study, we profiled genome-wide

chromatin accessibility in tens of thousands of guard cells isolated from soil-grown plants

exposed to different stimuli that trigger stomatal movements. Our results lead us to propose that

guard cell ABA signaling controls the balance between two opposing chromatin states. Under

unstressed conditions, AKS proteins maintain open chromatin at upstream regulatory regions of

ABA-repressed genes while ABA-activated genes are kept silent by closed chromatin. Under

abiotic stress such as drought, ABA levels rise in guard cells thereby activating SnRK2 kinases

which subsequently inhibit AKSs and activate ABFs. This leads to chromatin opening upstream

of abiotic stress-activated genes and the loss of chromatin accessibility upstream of stress-

repressed genes.

We uncovered patterns of chromatin accessibility specific to guard cells (Fig. 2D) and identified

cis-regulatory regions supporting guard cell specific gene expression (Fig. 2F-I and Dataset S4).

We found that ABA induces extensive and cell-type specific chromatin remodeling and that

guard cells possess a more extensive response than mesophyll and root cells (Fig. 3K and Fig.

S7C-L). Importantly, we find that these ABA-induced chromatin remodeling events correlate with

ABA-regulated gene expression (Fig. 3E and S6C). By performing TF-binding motif enrichment

and analyzing published DAP-seq data, we hypothesized that the bZIP transcription factors

ABF1/2/3/4 may function in ABA-induced chromatin remodeling in guard cells (Fig. 3C,G and

Fig. S6M). Interestingly, we find that among *in vitro* ABF binding sites, those occurring within

ABA-opened chromatin are most strongly associated with ABA-induced transcription (Fig. 4G),

indicating that regulated chromatin accessibility may determine which genomic ABF binding

sites are functional. Our experiments in *abf1/2/3/4* quadruple mutant guard cells demonstrate

that these four transcription factors are essential for the majority of ABA-triggered chromatin

opening (Fig. 4). Furthermore, experiments in root nuclei show that in *abf1/2/3/4* quadruple

mutants, even early ABA-induced chromatin remodeling events are strongly impaired (Fig. 4H

and Fig. S10), suggesting that ABFs initiate chromatin opening. How ABFs accomplish

chromatin opening remains to be determined, but existing literature could be interpreted to

support two possible and not necessarily mutually exclusive explanations. (1) Although

nucleosomes can restrict TF binding to DNA, some TFs, known as pioneer factors (68, 69),

possess the ability to bind their target sequences even within inaccessible chromatin regions. At

present, only a handful of pioneer factors have been documented in plants (70). While our data

are consistent with ABFs acting as pioneer factors, future biochemical experiments

reconstituting ABF binding to nucleosomal DNA are needed to investigate this model. (2) ABFs

may recruit chromatin modifying proteins to target sequences which could increase chromatin

accessibility. Interestingly, several such molecules including the SWI/SNF ATPase BRM (71,

72) and the histone deacetylase HDA9 (73) interact with known ABA signaling components.

Furthermore, ABF-binding proteins known as AFPs, which negatively regulate ABA signaling,

have been shown to physically interact with histone and chromatin modifying factors (74).

Although our data support a general requirement for ABF proteins in both roots and guard cells,

we also observe that ABA alters chromatin accessibility in a cell-type specific manner. At

present we do not understand how ABFs might contribute to cell-type specific regulation of

chromatin. Beyond ABF1/2/3/4, our analyses implicated additional TFs in ABA-dependent

genome reprogramming (Fig. 3F and 3G), which is consistent with prior research showing that

ABA regulates the binding of multiple transcription factors (75). We observed that ~26% of ABA-

gained ACRs (540 ACRs) still open in *abfx4* mutant guard cells in response to ABA (Fig. 4B).

These ABF-independent regions retained enrichment for CBF1/2/3 binding motifs, suggesting

that CBF transcription factors display ABF-independent binding (Fig. 4D). Finally, although ABA-

induced chromatin opening is severely impaired in *abf1/2/3/4* quadruple mutant guard cells, we

cannot exclude the possibility that other ABRE-binding TFs, in particular ABI5 (76, 77),

participate in this process as well.

Notably, our data suggest that the related bHLH transcription factors AKS1/2/3 play a larger role

in the regulation of ABA-repressed genes than was previously thought. Prior research had

shown that SnRK2 kinase-dependent inhibition of AKS1/2 was required for the ABA-dependent

transcriptional repression of the K+ ion channel gene *KAT1* in guard cells (24). Surprisingly, we

found that AKS1 bound many regions in the genome that lose chromatin accessibility in guard

cells following ABA treatment (Fig. 3C). Additionally, motifs recognized by AKS1/3 and the

related protein bHLH80 were enriched in ABA-repressed ACRs in guard cells (Fig. 3H), as well

as in root and mesophyll cells (Fig. 1F, Fig. S3G, Table S7). This observation was further

supported by de novo motif analysis (Fig S6N). AKS1 and AKS2 are inhibited by direct ABA-

dependent SnRK2 phosphorylation, but their expression was not controlled by ABA signaling

(44). We found that in guard cells, ABA represses *AKS3* expression implying that ABA-

dependent inhibition of AKS proteins occurs through both transcriptional and phosphorylation-

dependent mechanisms (Fig. 3I and J).

Elevation in the CO_2 concentration in leaves triggers rapid stomatal closing (34). The extent to

which elevated atmospheric CO_2 signals through the ABA signal transduction pathway is

currently under debate. Different research groups using the same mutants have come to

different conclusions regarding direct CO_2 signaling via the early ABA receptor/SnRK2 kinase

pathway during CO_2-induced stomatal closure. Our results exposed major differences in how

guard cells interpret these stimuli. In contrast to the extensive and persistent changes to

chromatin structure induced by ABA (Fig. 3), exposure to elevated CO_2 had minimal impact on

genome-wide chromatin accessibility in guard cells (Fig. 5). While ABA controlled chromatin

accessibility at 3018 ACRs in the genome, we found that only 90 ACRs were regulated by

elevated CO_2 (Fig 5C-E). Additionally, the strong enrichments of ABF binding motifs in gained

ACRs and AKS binding motifs in lost ACRs, evident following ABA treatment, are absent during

the response to elevated CO_2 (Fig. 5I). The disparity between these two signals at the level of

chromatin mirrors the disparity between characteristics of their respective physiological

responses. Unlike ABA- and drought-triggered stomatal closure, elevated CO_2-induced stomatal

closure is rapidly reversed upon return to ambient CO_2 (64). Such rapid changes in the

intercellular CO_2 concentration in leaves occur for example with alternating light intensities (e.g.

passing cloud cover) and require rapid reversals of stomatal CO_2 responses. Furthermore, we

found that exposure to low CO_2 triggers a distinct program of chromatin remodeling in guard

cells (Fig. 5F-I).

Finally, our results may open new avenues for investigation into two poorly understood features

of stomatal biology. Studies in multiple plant species have documented a phenomenon referred

to as stomatal "memory" (78, 79). Following drought stress and re-watering, plants typically

exhibit delayed and/or slow re-opening of stomata despite rapid recovery of plant water status.

Furthermore, guard cells display transcriptional memory where repetitive dehydration stress

leads to altered transcript levels in subsequent stress events (80–82). We found that ABA

triggered changes to chromatin accessibility in guard cells that can persist for up to 24 hours,

which could prime the genome for subsequent abiotic stress exposure (Fig. 3). These shifts

occurred at thousands of locations in the genome including upstream of many genes known to

regulate stomatal movements as well as many genes with potential functions in stomatal

physiology that could be the basis for future studies (Fig. S6 and Dataset S3). Moreover, our

findings support crosstalk between ABA and low CO_2 signaling at the chromatin remodeling

level. For example, we uncovered that ABA, but not high CO_2, triggered chromatin closing

upstream of genes encoding key CO_2-sensing and signaling proteins that promote low CO_2-

induced stomatal opening, including *HT1*, *BLUS1*, and *CBC2* (Fig. 5H). This response may

contribute to the long-term stomatal closing triggered by drought and ABA by inhibiting

mechanisms for low CO_2-induced stomatal opening. ABA-triggered chromatin dynamics may

provide a basis for long-term reprogramming of the guard cell genome for abiotic stress survival.

We envision that these sustained changes to chromatin structure may store the experience of

drought stress and contribute to long-term adjustments of stomatal function.

METHODS

Plant material and growth conditions

The Columbia-0 (Col-0) accession of *Arabidopsis thaliana* was used as the wild-type

background. Experiments on seedlings were conducted using plants germinated and grown on

1/2 Murashige and Skoog (MS) media (pH 5.7-8) solidified with 1% Phyto-agar. Experiments on

mature plants were conducted using plants grown in soil (Sunshine Mix #1) filled plastic pots

incubated in a growth chamber (Conviron) under a 12hr-12hr light-dark cycle, a light intensity of

100 μmol m^{-2} s^{-1}, a temperature of 22 C, ambient $[CO_2]$, and a relative humidity of 65%. To

minimize the impact of environmental variation all experiments were performed on plants grown

in two identical growth chambers equipped with $[CO_2]$ control.

ABA treatments in plate-based experiments were performed using a mesh (100 μm polyester)

transfer method. Seeds were germinated on ½ MS agar plates overlaid with sterilized polyester

mesh (100 μm). ABA treatments were initiated by transferring 8-day-old seedlings in mass to

plates containing 50 μM ABA (Sigma cis/trans-abscisic acid) or to plates containing vehicle

control (Ethanol). ABA treatments on soil grown plants were performed by spraying plants with

50 μM ABA diluted with 0.01% Silwet (in water) or with vehicle control. To assess treatment

responses, leaf temperature was measured using an infrared thermal imaging camera (T650sc;

FLIR).

Plasmid DNA and plant transformation

Constructs for cell-type specific nuclear sorting were prepared using multi-site gateway

technology (Invitrogen). The ORF encoding the histone H2B (*AT5G22880*) was cloned in a

pDONR221 entry vector. The cell type specific promotors RBCp and GC1p, from the Rubisco

small subunit 2B gene (*AT5G38420)* and the *GASA9* (*AT1G22690*) gene respectively, were

cloned from the *Arabidopsis* genome into the gateway entry vector pDONR-p4p1r. Final plant

transformation constructs were generated by performing multisite gateway recombination into

the pP7m34GW destination vector. The plasmid for expressing *GFP-AKS3* under the *AKS3*

(*AT2G42280*) endogenous promotor was generated using GreenGate cloning (83). The cloned

AKS3 upstream regulatory sequence was defined using guard cell ATAC-seq data.

Transcriptional reporter constructs expressing nuclear localized GFP under the guard cell

specific ACRs upstream of *AAE9* (*AT1G21540*) and *ALMT6* (*AT2G17470*) were generated

using GreenGate cloning. Primer sequences are available in Dataset S8. Constructs were

transformed into *A. thaliana* Col-0 plants using floral dipping with *Agrobacterium tumefaciens*

strain GV3101.

Fluorescence-activated nuclear sorting (FANS)

FANS on seedling/root nuclei and mesophyll nuclei was performed similar to (40) with small

modifications. Briefly, whole seedlings, surgically isolated roots, or leaves were chopped

vigorously with a clean razor blade in 2-3 ml of ice-cold FANS-lysis buffer (15 mM Tris-HCl

pH7.5, 20 mM NaCl, 80 mM KCl, 0.2 mM spermine, 5 mM 2-ME, 0.5 mM spermidine, 0.2%

IGEPAL CA-630, Roche mini EDTA-free Protease Inhibitor Cocktail) for 5 minutes in a cold

glass petri dish. The subsequent lysate was filtered through a 40 µm filter unit before

transferring to a 15 ml tube containing 3 ml of FANS-Sucrose buffer (20 mM Tris-HCl pH 8.0, 2

mM $MgCl_2$, 2 mM EDTA, 15 mM 2-ME, 1.7 M sucrose, 0.2% IGEPAL CA-630). Nuclei were

pelleted at 2500xG at 4ºC for 25 mins, after which the supernatant was carefully removed. The

nuclei pellet was then gently resuspended, using clipped pipette tips, with 600 µl FANS-lysis

buffer supplemented with DAPI, and then immediately used for flow cytometry.

For experiments on guard cells, the leaves of ~40 six-week-old plants were harvested and

blended in 500 ml of ice-cold water for 2x40 secs with 1 min between pulses. The blended

sample was poured over a 100 µm mesh to remove mesophyll cells and collect epidermal

enriched fragments. The mesh was rinsed with ~200 ml of cold water and gently blotted with

paper towels to remove excess liquid. Thin sections of epidermal tissue were then flash-frozen

in liquid nitrogen. Epidermal sections were ground to a fine powder with a frozen mortar and

pestle. After grinding, powder was transferred to a new mortar with a frozen metal spatula.

Immediately 12-15 ml of ice-cold FANS-lysis buffer was added, and the powder was gently

resuspended with a clean pestle. The subsequent lysate was filtered through a 40 µm filter unit

before transferring to a 15 ml tube containing 6 ml of FANS-Sucrose buffer. Nuclei were then

pelleted and prepared for FACS as above.

Flow cytometry was conducted on a BD FACS Aria II using a 70 µm nozzle with a flow rate < 2

and the following photomultiplier parameters: Forward Scatter (FSC) – 250 eV, Side Scatter

(SSC) – 220 eV, BV421 (DAPI channel) – 550 eV, FITC (GFP channel) – 340 eV. Sort gates

were defined empirically using the FSC and SSC channels to exclude debris and potential

doublets. Before sorting GFP labeled nuclei, FACS analysis on unlabeled control samples (Col-

0 nuclei) was performed to define background fluorescence signal. When isolating guard cell

nuclei, sort gates were established to purify exclusively 2N (diploid) and GFP-positive nuclei. To

evaluate the purity of sorted samples, a fraction (~10% of total volume) was re-run under the

same sorting program (Fig. S2D). Sorted nuclei were collected in 600 µl of FANS-lysis buffer in

a refrigerated tube chamber. After sorting, nuclei were pelleted by centrifugation at 1000xg for

10 mins at 4C. Nuclei pellets were gently washed with 1 ml of 10 mM Tris-HCl pH 8.0, 5 mM

MgCl2 buffer, before centrifugation again as above. After carefully removing the supernatant,

nuclei were used immediately for ATAC-seq.

Detailed methods descriptions for microscopy and time-resolved stomatal conductance

measurements can be found in the SI Appendix, Materials and Methods. Detailed protocols for

RNA-seq and ATAC-seq library preparation and description of ATAC-seq, RNA-seq, and DAP-

seq data analyses can be found in the SI Appendix, Materials and Methods.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

Funding for this study was provided by the National Institutes of Health (R01GM060396 to J.I.S

and F32GM137544 to C.A.S). CO_2 experiments were supported by National Science

Foundation grant MCB-1900567 to J.I.S. This study includes data generated at the UC San

Diego IGM Genomics Center utilizing an Illumina NovaSeq 6000 that was purchased with

funding from a National Institutes of Health SIG grant (#S10 OD026929). Microscopy was

supported by the UC San Diego Microscopy Core with funding by grant NINDS P30NS047101.

We thank Kazuko Yamaguchi-Shinozaki (University of Tokyo) for sharing *abf1/2/3/4* quadruple

mutant seeds, and members of the Schroeder lab for helpful discussions throughout the

research.

Author contributions: C.A.S. and J.I.S. conceived of the study and designed the methodology.

C.A.S. performed all experiments and analyzed the data. C.A.S. and J.I.S. wrote and edited the

paper.

Data and materials availability: All materials (plasmids, seed stocks) are available upon

request. Raw FASTQ files generated in this study and ATAC-seq peak files in narrowPeak

format are available at NCBI's Gene Expression Omnibus under accession GSE243473. All

data are available in the main text and supplementary material.

REFERENCES

1.   R. J. Schmitz, E. Grotewold, M. Stam, Cis-regulatory sequences in plants: Their importance, discovery, and future challenges. *The Plant Cell* **34**, 718–741 (2022).

2.   W. K. M. Lai, B. F. Pugh, Understanding nucleosome dynamics and their links to gene expression and DNA replication. *Nature Reviews Molecular Cell Biology* **18**, 548–562 (2017).

3.   T. C. Voss, G. L. Hager, Dynamic regulation of transcriptional states by chromatin and transcription factors. *Nature Reviews Genetics* **15**, 69–81 (2014).

4.   J. L. Nemhauser, F. Hong, J. Chory, Different Plant Hormones Regulate Similar Processes through Largely Nonoverlapping Transcriptional Responses. *Cell* **126**, 467–475 (2006).

5.   R. Waadt, *et al.*, Plant hormone regulation of abiotic stress responses. *Nat Rev Mol Cell Biol* **23**, 680–694 (2022).

6.   P. Sijacic, M. Bajic, E. C. McKinney, R. B. Meagher, R. B. Deal, Changes in chromatin accessibility between Arabidopsis stem cells and mesophyll cells illuminate cell type-specific transcription factor networks. *The Plant Journal* **94**, 215–231 (2018).

7.   K. A. Maher, *et al.*, Profiling of Accessible Chromatin Regions across Multiple Plant Species and Cell Types Reveals Common Gene Regulatory Principles and New Control Modules. *The Plant Cell* **30**, 15–36 (2018).

8.   A. M. Sullivan, *et al.*, Mapping and Dynamics of Regulatory DNA in Maturing Arabidopsis thaliana Siliques. *Frontiers in Plant Science* **10** (2019).

9.   A. Farmer, S. Thibivilliers, K. H. Ryu, J. Schiefelbein, M. Libault, Single-nucleus RNA and ATAC sequencing reveals the impact of chromatin accessibility on gene expression in Arabidopsis roots at the single-cell level. *Molecular Plant* **14**, 372–383 (2021).

10. M. Borg, *et al.*, Epigenetic reprogramming rewires transcription during the alternation of generations in Arabidopsis. *eLife* **10**, e61894 (2021).

11. A. P. Marand, Z. Chen, A. Gallavotti, R. J. Schmitz, A cis-regulatory atlas in maize at single-cell resolution. *Cell* **184**, 3041-3055.e21 (2021).

12. E.-D. Kim, *et al.*, Dynamic chromatin accessibility deploys heterotypic cis/trans-acting factors driving stomatal cell-fate commitment. *Nat. Plants* **8**, 1453–1466 (2022).

13. K. C. Potter, J. Wang, G. E. Schaller, J. J. Kieber, Cytokinin modulates context-dependent chromatin accessibility through the type-B response regulators. *Nat. Plants* **4**, 1102–1111 (2018).

14. M. A. Reynoso, *et al.*, Evolutionary flexibility in flooding response circuitry in angiosperms. *Science* **365**, 1291–1295 (2019).

15. E. S. Karaaslan, N. Faiß, C. Liu, K. W. Berendzen, Isolation of Lineage Specific Nuclei
Based on Distinct Endoreduplication Levels and Tissue-Specific Markers to Study
Chromatin Accessibility Landscapes. *Plants* **9**, 1478 (2020).

16. S. R. Cutler, P. L. Rodriguez, R. R. Finkelstein, S. R. Abrams, Abscisic Acid: Emergence of
a Core Signaling Network. *Annual Review of Plant Biology* **61**, 651–679 (2010).

17. R. Finkelstein, Abscisic Acid Synthesis and Response. *Arabidopsis Book* **11** (2013).

18. J.-K. Zhu, Abiotic Stress Signaling and Responses in Plants. *Cell* **167**, 313–324 (2016).

19. Y. Ma, *et al.*, Regulators of PP2C Phosphatase Activity Function as Abscisic Acid Sensors.
Science **324**, 1064–1068 (2009).

20. S.-Y. Park, *et al.*, Abscisic Acid Inhibits Type 2C Protein Phosphatases via the PYR/PYL
Family of START Proteins. *Science* **324**, 1068–1071 (2009).

21. T. Umezawa, *et al.*, Type 2C protein phosphatases directly regulate abscisic acid-activated
protein kinases in Arabidopsis. *Proceedings of the National Academy of Sciences* **106**,
17588–17593 (2009).

22. H. Fujii, P. E. Verslues, J.-K. Zhu, Identification of Two Protein Kinases Required for
Abscisic Acid Regulation of Seed Germination, Root Growth, and Gene Expression in
Arabidopsis. *The Plant Cell* **19**, 485–494 (2007).

23. K. E. Hubbard, N. Nishimura, K. Hitomi, E. D. Getzoff, J. I. Schroeder, Early abscisic acid
signal transduction mechanisms: newly discovered components and newly emerging
questions. *Genes Dev.* **24**, 1695–1708 (2010).

24. Y. Takahashi, *et al.*, bHLH Transcription Factors That Facilitate K+ Uptake During
Stomatal Opening Are Repressed by Abscisic Acid Through Phosphorylation. *Science
Signaling* **6**, ra48–ra48 (2013).

25. T. Yoshida, *et al.*, AREB1, AREB2, and ABF3 are master transcription factors that
cooperatively regulate ABRE-dependent ABA signaling involved in drought stress
tolerance and require ABA for full activation. *The Plant Journal* **61**, 672–685 (2010).

26. T. Yoshida, *et al.*, Four Arabidopsis AREB/ABF transcription factors function predominantly
in gene expression downstream of SnRK2 kinases in abscisic acid signalling in response
to osmotic stress. *Plant, Cell & Environment* **38**, 35–49 (2015).

27. P. Wang, *et al.*, Quantitative phosphoproteomics identifies SnRK2 protein kinase
substrates and reveals the effectors of abscisic acid action. *Proceedings of the National
Academy of Sciences* **110**, 11205–11210 (2013).

28. T. Umezawa, *et al.*, Genetics and Phosphoproteomics Reveal a Protein Phosphorylation
Network in the Abscisic Acid Signaling Pathway in Arabidopsis thaliana. *Sci. Signal.* **6**,
rs8–rs8 (2013).

29. N. Leonhardt, *et al.*, Microarray Expression Analyses of Arabidopsis Guard Cells and
Isolation of a Recessive Abscisic Acid Hypersensitive Protein Phosphatase 2C Mutant.
The Plant Cell **16**, 596–615 (2004).

30. H. Goda, *et al.*, The AtGenExpress hormone and chemical treatment data set:
experimental design, data evaluation, model data analysis and data access. *The Plant*
Journal **55**, 526–542 (2008).

31. R.-S. Wang, *et al.*, Common and unique elements of the ABA-regulated transcriptome of
Arabidopsis guard cells. *BMC Genomics* **12**, 216 (2011).

32. A. M. Hetherington, F. I. Woodward, The role of stomata in sensing and driving
environmental change. *Nature* **424**, 901–908 (2003).

33. S. M. Assmann, T. Jegla, Guard cell sensory systems: recent insights on stomatal
responses to light, abscisic acid, and CO2. *Current Opinion in Plant Biology* **33**, 157–167
(2016).

34. G. Dubeaux, *et al.*, Deep dive into CO2-dependent molecular mechanisms driving stomatal
responses in plants. *Plant Physiology* **187**, 2032–2042 (2021).

35. Z. Liu, *et al.*, Global Dynamic Molecular Profiling of Stomatal Lineage Cell Development by
Single-Cell RNA Sequencing. *Molecular Plant* **13**, 1178–1193 (2020).

36. C. B. Lopez-Anido, *et al.*, Single-cell resolution of lineage trajectories in the Arabidopsis
stomatal lineage and developing leaf. *Developmental Cell* **56**, 1043-1055.e4 (2021).

37. J.-Y. Kim, *et al.*, Distinct identities of leaf phloem cells revealed by single cell
transcriptomics. *The Plant Cell* **33**, 511–530 (2021).

38. C. Procko, *et al.*, Leaf cell-specific and single-cell transcriptional profiling reveals a role for
the palisade layer in UV light protection. *The Plant Cell* **34**, 3261–3279 (2022).

39. J. D. Buenrostro, P. G. Giresi, L. C. Zaba, H. Y. Chang, W. J. Greenleaf, Transposition of
native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-
binding proteins and nucleosome position. *Nature Methods* **10**, 1213–1218 (2013).

40. Z. Lu, B. T. Hofmeister, C. Vollmers, R. M. DuBois, R. J. Schmitz, Combining ATAC-seq
with nuclei sorting for discovery of cis-regulatory regions in plant genomes. *Nucleic Acids*
Research **45**, e41 (2017).

41. A. M. Jones, *et al.*, Abscisic acid dynamics in roots detected with genetically encoded
FRET sensors. *eLife* **3**, e01741 (2014).

42. R. Waadt, *et al.*, FRET-based reporters for the direct visualization of abscisic acid
concentration changes and distribution in Arabidopsis. *eLife* **3**, e01739 (2014).

43. R. C. O'Malley, *et al.*, Cistrome and Epicistrome Features Shape the Regulatory DNA
Landscape. *Cell* **165**, 1280–1292 (2016).

44.   Y. Takahashi, T. Kinoshita, M. Matsumoto, K. Shimazaki, Inhibition of the Arabidopsis
741   bHLH transcription factor by monomerization through abscisic acid-induced
742   phosphorylation. *The Plant Journal* **87**, 559–567 (2016).

743   45.   Y. Yang, A. Costa, N. Leonhardt, R. S. Siegel, J. I. Schroeder, Isolation of a strong
744   Arabidopsis guard cell promoter and its potential as a research tool. *Plant Methods* **4**, 6
745   (2008).

746   46.   J. A. Anderson, S. S. Huprikar, L. V. Kochian, W. J. Lucas, R. F. Gaber, Functional
747   expression of a probable Arabidopsis thaliana potassium channel in Saccharomyces
748   cerevisiae. *Proceedings of the National Academy of Sciences* **89**, 3736–3740 (1992).

749   47.   D. P. Schachtman, J. 1. Schroeder, W. J. Lucas, J. A. Anderson, R. F. Gaber, Expression
750   of an Inward-Rectifying Potassium Channel by the Arabidopsis KAT1 cDNA. *Science* **258**,
751   1654–1658 (1992).

752   48.   R. L. Nakamura, *et al.*, Expression of an Arabidopsis Potassium Channel Gene in Guard
753   Cells. *Plant Physiology* **109**, 371–374 (1995).

754   49.   O. S. Lau, *et al.*, Direct roles of SPEECHLESS in the specification of stomatal self-
755   renewing cells. *Science* **345**, 1605–1609 (2014).

756   50.   S.-K. Han, *et al.*, MUTE Directly Orchestrates Cell-State Switch and the Single Symmetric
757   Division to Create Stomata. *Developmental Cell* **45**, 303-315.e5 (2018).

758   51.   K. Ohashi-Ito, D. C. Bergmann, Arabidopsis FAMA Controls the Final
759   Proliferation/Differentiation Switch during Stomatal Development. *The Plant Cell* **18**, 2493–
760   2505 (2006).

761   52.   H. Yu, *et al.*, Activated Expression of an Arabidopsis HD-START Protein Confers Drought
762   Tolerance with Improved Root System and Reduced Stomatal Density. *The Plant Cell* **20**,
763   1134–1151 (2008).

764   53.   J. Negi, *et al.*, A Dof Transcription Factor, SCAP1, Is Essential for the Development of
765   Functional Stomata in Arabidopsis. *Current Biology* **23**, 479–484 (2013).

766   54.   S. Meyer, *et al.*, Malate transport by the vacuolar AtALMT6 channel in guard cells is
767   subject to multiple regulation. *The Plant Journal* **67**, 247–257 (2011).

768   55.   S. Li, *et al.*, Arabidopsis ACYL-ACTIVATING ENZYME 9 (AAE9) encoding an isobutyl-CoA
769   synthetase is a key factor connecting branched-chain amino acid catabolism with iso-
770   branched wax biosynthesis. *New Phytologist* **233**, 2458–2470 (2022).

771   56.   E. Cominelli, *et al.*, A Guard-Cell-Specific MYB Transcription Factor Regulates Stomatal
772   Movements and Plant Drought Tolerance. *Current Biology* **15**, 1196–1200 (2005).

773   57.   M. Hashimoto, *et al.*, Arabidopsis HT1 kinase controls stomatal movements in response to
774   CO2. *Nat Cell Biol* **8**, 391–397 (2006).

775   58.   Y.-K. Liang, *et al.*, AtMYB61, an R2R3-MYB Transcription Factor Controlling Stomatal
776   Aperture in Arabidopsis thaliana. *Current Biology* **15**, 1201–1206 (2005).

59. M. B. Prouse, M. M. Campbell, Interactions between the R2R3-MYB Transcription Factor,
AtMYB61, and Target DNA Binding Sites. *PLOS ONE* **8**, e65132 (2013).

60. S. J. Gilmour, *et al.*, Low temperature regulation of the Arabidopsis CBF family of AP2
transcriptional activators as an early step in cold-induced COR gene expression. *The Plant*
Journal **16**, 433–442 (1998).

61. Y. Sun, *et al.*, Divergence in the ABA gene regulatory network underlies differential growth
control. *Nat. Plants*, 1–12 (2022).

62. S. Heinz, *et al.*, Simple Combinations of Lineage-Determining Transcription Factors Prime
cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell* **38**,
576–589 (2010).

63. W. Lahr, K. Raschke, Abscisic-acid contents and concentrations in protoplasts from guard
cells and mesophyll cells ofVicia faba L. *Planta* **173**, 528–531 (1988).

64. P.-K. Hsu, *et al.*, Abscisic acid-independent stomatal CO_2 signal transduction pathway and
convergence of CO_2 and ABA signaling downstream of OST1 kinase. *Proceedings of the*
National Academy of Sciences **115**, E9971–E9980 (2018).

65. L. Zhang, *et al.*, FRET kinase sensor development reveals SnRK2/OST1 activation by ABA
but not by MeJA and high CO_2 during stomatal closure. *eLife* **9**, e56351 (2020).

66. M. Dittrich, *et al.*, The role of Arabidopsis ABA receptors from the PYR/PYL/RCAR family
in stomatal acclimation and closure signal integration. *Nat. Plants* **5**, 1002–1011 (2019).

67. A. Hiyama, *et al.*, Blue light and CO_2 signals converge to regulate light-induced stomatal
opening. *Nat Commun* **8**, 1284 (2017).

68. M. Iwafuchi-Doi, K. S. Zaret, Pioneer transcription factors in cell reprogramming. *Genes*
Dev. **28**, 2679–2692 (2014).

69. A. Balsalobre, J. Drouin, Pioneer factors as master regulators of the epigenome and cell
fate. *Nat Rev Mol Cell Biol*, 1–16 (2022).

70. R. Jin, *et al.*, LEAFY is a pioneer transcription factor and licenses cell reprograming to
floral fate. *Nat Commun* **12**, 626 (2021).

71. S.-K. Han, *et al.*, The SWI2/SNF2 Chromatin Remodeling ATPase BRAHMA Represses
Abscisic Acid Responses in the Absence of the Stress Stimulus in Arabidopsis. *The Plant*
Cell **24**, 4892–4906 (2012).

72. M. Peirats-Llobet, *et al.*, A Direct Link between Abscisic Acid Sensing and the Chromatin-
Remodeling ATPase BRAHMA via Core ABA Signaling Pathway Components. *Molecular*
Plant **9**, 136–147 (2016).

73. D. Baek, *et al.*, Histone Deacetylase HDA9 With ABI4 Contributes to Abscisic Acid
Homeostasis in Drought Stress Response. *Frontiers in Plant Science* **11** (2020).

74. T. J. Lynch, B. J. Erickson, D. R. Miller, R. R. Finkelstein, ABI5-binding proteins (AFPs)
813   alter transcription of ABA-induced genes via a variety of interactions with chromatin
814   modifiers. *Plant Mol Biol* **93**, 403–418 (2017).

815   75. L. Song, *et al.*, A transcription factor hierarchy defines an environmental stress response
816   network. *Science* **354**, 1550 (2016).

817   76. R. Finkelstein, S. S. L. Gampala, T. J. Lynch, T. L. Thomas, C. D. Rock, Redundant and
818   Distinct Functions of the ABA Response Loci ABA-INSENSITIVE(ABI)5 and ABRE-
819   BINDING FACTOR (ABF)3. *Plant Mol Biol* **59**, 253–267 (2005).

820   77. R. R. Finkelstein, T. J. Lynch, The Arabidopsis Abscisic Acid Response Gene ABI5
821   Encodes a Basic Leucine Zipper Transcription Factor. *The Plant Cell* **12**, 599–609 (2000).

822   78. E. Fleta-Soriano, M. Pintó-Marijuan, S. Munné-Bosch, Evidence of Drought Stress Memory
823   in the Facultative CAM, Aptenia cordifolia: Possible Role of Phytohormones. *PLOS ONE*
824   **10**, e0135391 (2015).

825   79. P. A. Auler, F. B. S. Freire, V. F. Lima, D. M. Daloso, On the role of guard cells in sensing
826   environmental signals and memorising stress periods. *Theor. Exp. Plant Physiol.* **34**, 277–
827   299 (2022).

828   80. Y. Ding, M. Fromm, Z. Avramova, Multiple exposures to drought "train" transcriptional
829   responses in *Arabidopsis*. *Nature Communications* **3**, 740 (2012).

830   81. L. Virlouvet, M. Fromm, Physiological and transcriptional memory in guard cells during
831   repetitive dehydration stress. *New Phytologist* **205**, 596–607 (2015).

832   82. A. van Weringh, A. Pasha, E. Esteban, P. J. Gamueda, N. J. Provart, Generation of guard
833   cell RNA-seq transcriptomes during progressive drought and recovery using an adapted
834   INTACT protocol for Arabidopsis thaliana shoot tissue. 2021.04.15.439991 (2021).

835   83. A. Lampropoulos, *et al.*, GreenGate - A Novel, Versatile, and Efficient Cloning System for
836   Plant Transgenesis. *PLOS ONE* **8**, e83043 (2013).

FIGURES LEGENDS
854

Figure 1. ABA induces rapid and genome-wide chromatin remodeling in seedlings. A)
Volcano plot of differential ATAC-seq analysis showing regions that significantly change in
chromatin accessibility in seedlings after 4hrs of ABA treatment. ABA increased accessibility at
666 regions and decreased accessibility at 214 regions (FDR < 0.001 and FC > 1.5, differential
peaks colored in red). **B)** Heatmap of root nuclei ATAC-seq signal (RPKM-normalized) centered
over ATAC-seq peaks (+/- 1.5 kb) with ABA-regulated chromatin accessibility during the
indicated ABA treatment time course. Across all timepoints, ABA increased accessibility at 1293
regions and decreased accessibility at 1119 regions (FDR < 0.001 and FC > 1.5). **C)** Number of
ABA-regulated ATAC-seq peaks over time in roots. ABA-induced peaks are colored in blue
while ABA-repressed peaks are colored in pink. **D)** Distribution among annotated genomic
features of either all ATAC-seq peaks (orange) or ABA-regulated ATAC-seq peaks (grey). **E)**
Genome browser snapshots at two representative genes (*MAPKKK18* and *CML25*) showing
changes in ATAC-seq signal in upstream regions following ABA treatment. Scale bars indicate
500 bp. **F)** The transcription factor binding motifs with the highest enrichment (by binomial p-
value) in ABA-induced and ABA-repressed chromatin regions in roots.
870
871

Fig 2. Cell-type specific epigenomics identifies guard cell specific regulatory elements.
A) Left panel is a fluorescence micrograph showing guard cell specific expression of H2B-GFP.
Right panel shows a 3D reconstruction from confocal imaging of a 5-week-old *GC1p:H2B-GFP*
leaf (merged image shows GFP fluorescence in green and chlorophyll fluorescence in
magenta). **B)** FACS analysis of nuclei (DAPI positive) from *GC1p:H2b-GFP* plants. A GFP
density plot defines two distinct populations separated by GFP intensity. **C)** DAPI density plot
shows that GFP positive nuclei (blue population) are primarily diploid, while GFP negative nuclei
(pink population) are distributed over multiple ploidies. **D)** Chromatin accessibility heatmaps
comparing ATAC-seq signal (RPKM-normalized) in guard cell, mesophyll, or root cell nuclei.
ATAC-seq peaks are separated into guard cell enriched (3272 ACRs), mesophyll enriched
(1472 ACRs), and root enriched (3364 ACRs) sets defined by differential accessibility (p.adj <
0.001, Fold-Change > 2). **E)** Principal component analysis (PCA) comparing ATAC-seq
biological replicates derived from guard cell, mesophyll, or root nuclei. **F)** A peak-centered
ATAC-seq metaplot and a gene-body centered RNA-seq metaplot showing average chromatin
accessibility at guard cell specific peaks and average transcript levels of nearest downstream
genes (ATAC-seq peak within -2.5 kb and +0.5 kb of transcription start site). Guard cell derived
signal is shown in black while mesophyll/leaf signal is shown in green. **G)** Genome browser
images displaying ATAC-seq and RNA-seq signal for the indicated tracks near the guard cell
expressed gene *KAT1* and the mesophyll cell expressed gene *AOC1*. Scale bars indicate 500
bp. **H)** Results of de novo motif analysis using homer are shown along with the best matching
known TFs and the enrichment p-values. **I)** Transcriptional reporter lines generated using
cloned guard cell specific ACRs upstream of the genes *AAE9* and *ALMT6,* the positions of the
cloned sequences are indicated with orange highlighting on the genome browser images (scale
bars indicate 500 bp). Confocal micrographs showing GFP fluorescence (left panel) or merged
GFP (green) and chlorophyll (magenta) fluorescence (right panel). Scale bar represents 20 μm.

Figure 3. ABA triggers extensive and persistent chromatin remodeling in guard cells.
A) Schematic showing the experimental outline. Plants were sprayed with ABA and infrared images
were taken using a thermal camera to assess stomatal closure. Guard cell nuclei were purified at
4hrs and 24 hrs after treatment and used for ATAC-seq. **B)** Volcano plot of guard cell enriched RNA-
seq data showing that ABA upregulates 2142 transcripts and downregulates 1574 transcripts (FDR <
0.001, FC > 2.0). Differentially expressed transcripts are colored in red. **C)** Heatmaps and profile
plots of ATAC-seq signal (RPKM-normalized) in guard cell nuclei centered (+/- 1.5 kb) over 2323
regions with ABA-induced chromatin accessibility and 1657 regions with ABA-repressed chromatin
accessibility (FDR < 0.001, FC > 2.0). The adjacent heatmaps show DAP-seq signal for ABF2 (in
red) and AKS1 (in black) centered over the same ABA regulated ACRs, data from (39) and (56). **D)**
Comparison of ABA-regulated peaks from 4hr and 24hr timepoints separated ACRs into three
categories – early only, sustained (both early and late), and late only. The bar graph shows the
number of ABA-increased and -decreased regions in each category. **E)** (Top) Metaplots showing
average ATAC-seq signal centered over either ABA-induced or ABA-repressed ACRs and (bottom)
RNA-seq metaplots showing average transcript level centered over gene bodies downstream of the
same ACRs. Signal from ABA-treated plants is shown in orange and signal from control treated
plants is shown in black. **F)** Genome browser images showing guard cell ATAC-seq and RNA-seq
signal at a representative ABA-induced gene (*MYB121)* and an ABA-repressed gene (*MYB60*).
Scale bars indicated 500 bp. **G,H)** Heatmaps showing TF binding motif enrichments (-log(p-value))
in ABA-increased **(G)** and ABA-decreased **(H)** ATAC-seq peaks. **I)** Genome browser images
showing guard cell ATAC-seq and RNA-seq signal at the *AKS3* gene. Scale bars indicated 500 bp.
J) Fluorescence micrograph showing expression of *AKS3pr::GFP-AKS3* in guard cells after
treatment with control or ABA (4 hr). **K)** Overlap of ABA-induced accessible regions in guard cell
nuclei, mesophyll nuclei, and root nuclei.
928
929
Fig 4. ABF transcription factors are required for ABA-induced chromatin opening in
guard cells. A) Heatmaps of ATAC-seq signal (RPKM-normalized) in either *WT* or *abfx4*
mutant guard cells centered over 2079 regions with ABA-induced accessibility and 975 regions
with ABA-repressed accessibility (FDR < 0.001 and FC > 2.0). ACRs are subdivided by
requirement for ABF1/2/3/4 **B)** Overlap of ABA-induced ATAC-seq peaks in *WT* guard cells with
those in *abfx4* mutant guard cells. **C)** Genome browser snapshots showing ATAC-seq and
RNA-seq signal near two ABA-induced genes *DTX37* and *HSFA6B*. Scale bars indicate 1 kb. **D)**
Enrichment (-log(p-value)) of the indicated transcription factor binding motifs in ABA regulated
ATAC-seq peaks in either *WT* or *abfx4* mutant guard cell nuclei. **E)** Overlap between ABF DAP-
seq peaks and ABA-induced accessible chromatin regions (ACRs). **F)** Peak-centered metaplot
showing ATAC-seq signal in the indicated samples at ABA-induced ACRs containing ABF DAP-
seq binding sites. **G)** 1000 ABF DAP-seq peaks overlapping ABA-induced ATAC-seq peaks or
1000 random ABF DAP-seq peaks were assigned to nearest downstream genes (within -2.5 kb
and +0.5 kb of TSS). The plot shows the ABA-induced fold changes within these two sets of
downstream transcripts. Asterisks (****) indicate that the distributions are significantly different
by Welch's t-test ($p < 1E-7$) and the red bars represent the means of the distributions. **H)** Bar
plots representing the number of ABA-activated or ABA-repressed ACRs over time in *Col-0* or
abfx4 mutant roots.
948
949
Fig 5. ABA and changes in CO_2 concentration induce distinct chromatin remodeling
programs in guard cells. A) Diagram showing the effects of different stimuli on the stomatal
aperture. **B)** Representative IR image showing differences in leaf temperature due to CO_2
induced stomatal movement. **C-E)** Volcano plots of differential ATAC-seq results showing **C)**
ABA-regulated chromatin accessibility, **D)** low CO_2 regulated chromatin accessibility, and **E)**
high CO2 regulated chromatin accessibility. The numbers of differentially accessible peaks

(colored in red, FDR < 0.001 and FC > 2.0) in each condition are indicated on each plot. **F)**
Heatmaps and profile plots of guard cell ATAC-seq signal centered over either ABA-regulated
ACRs (2079 induced and 975 repressed) or low CO_2-regulated ACRs (329 induced and 487
repressed). Libraries are derived from plants exposed to the indicated treatments. **G)** Principal
component analysis (PCA) shows that ABA, low CO_2, and high CO_2 introduce separate variation
to genome-wide chromatin accessibility. **H)** Genome browser images showing ABA-repression
of chromatin accessibility upstream of *HT1*, *BLUS1*, and *CBC1* – three positive regulators of low
CO_2 induced stomatal opening. Scale bars indicate 1 kb. **I)** Enrichment (-log P_value) of the
indicated transcription factor binding motifs in ABA, Low CO_2, or High CO_2 regulated regions.
Negative enrichment scores indicate motifs overrepresented in treatment-repressed regions. No
significant enrichment for any of these motifs was found in High CO_2 regulated ACRs.
967
968

A Seedling ATACseq: ABA vs control

B Roots ATACseq ABA
ctrl 45 120 240

ABA-induced

ABA-repressed

−1.5 kb peak center peak center +1.5 kb

ATAC-seq (RPKM)
2000
250

C

DOWN UP

Number of peaks

Time (mins)
45 120 240

D

Peaks
ABA All

Percentage

Distal Downstream Promoter (< 1 kb) Promoter (1 −2 kb) Transcript

E

[0-4500]
control
45 mins
120 mins
240 mins
MAPKKK18

[0-3500]
control
45 mins
120 mins
240 mins
CML25

F

CACGTGGC ABF1 (1e-63) ↑

CAACTTG AKS3 (1e-32) ↓

A GC1p:H2B-GFP

100 um

GC1:H2B-GFP

B

Num. of nuclei

GFP
Neg.
Pos.

40000

1e+01   1e+05
GFPArea

C

Scaled Density

GFP
Neg.
Pos.

1.00

0.00

3000   10000
DAPI Area

D

| Guard Cell | Root | Mesophyll |

Guard Cell specific

Meso. specific

Root specific

ATAC-seq (RPKM)

3000

1500

-1.5 kb   peak center   +1.5 kb

E

PC2: 23%

-10

-20   0   20   40
PC1: 76%

GC
Meso.
Root

F

ATAC-seq
Guard Cell
Mesophyll

-1.5 kb   Peak center   +1.5 kb

mRNA-seq

2500

TSS   gene body   TES

G

ATAC   [0-7000]   Mesophyll
Guard Cell

mRNA   [0-9000]   Leaf
Guard Cell

KAT1

ATAC   [0-5000]   Mesophyll
Guard Cell

mRNA   [0-25000]   Leaf
Guard Cell

AOC1

H

CACGTGAC   bHLH: SPCH, MUTE, FAMA (1e-144)

GCATTTAATG   EDT1 (1e-38)

CCTACTTTAA   C2C2-DOF (1e-22)

I

Mesophyll

Guard Cell   [0-14000]

AAE9

AAE9pr:GFP-nls

Mesophyll

Guard Cell   [0-9000]

ALMT6

ALMT6pr:GFP-nls

A Open ← Low CO2 / High CO2, ABA → Closed

B CO₂ (PPM) 100 1000

C ABA vs control
975 ACRs decrease
2079 ACRs increase

D 100 ppm CO2 vs control
487 ACRs decrease
329 ACRs increase

E 1000 ppm CO2 vs control
79 ACRs decrease
11 ACRs increase

−log₁₀(p. adjusted)
log₂(fold change)

F 100 ppm
control | ABA | [CO₂]
ABA-ind. | ABA-rep.
ATAC-seq (RPKM)
4000
500
ABA induced | ABA repressed
−1.5 kb | peak center | +1.5 kb

control | ABA | [CO₂]
Low CO₂-ind. | LowCO₂-rep.
Low CO₂ induced | Low CO₂ repressed
−1.5 kb | +1.5 kb

G
PC2: 6% variance
PC1: 91% variance
control | ABA | Low CO₂ | High CO₂

H
control | ABA | Low CO₂ | High CO₂
[0-11000] | [0-13000] | [0-9000]
control | ABA
[0-6000]
HT1 | *BLUS1* | *CBC2*

I Enrichment
200
100
0
−100
High CO₂
Low CO₂
ABA
WRKY28 | WRKY50 | OBP1 | bZIP53 | bZIP3 | bHLH80 | ABF3 | AKS3 | AKS1 | Adof1 | ABI5 | ABF2 | ABF1 | C3H | C2C2dof.1 | C2C2dof.2