PI(3,4,5)P3 allosteric regulation of repressor activator protein 1 controls antigenic variation in trypanosomes

Abdoulie O. Touray[1,2], Rishi Rajesh[1], Tony Isebe[1], Tamara Sternlieb[1], Mira Loock[1], Oksana Kutova[1], Igor Cestari[1,2,*]

[1]Institute of Parasitology, McGill University, Sainte-Anne-de-Bellevue, QC H9X 3V9, Canada

[2]Division of Experimental Medicine, Department of Medicine, McGill University, Montreal, QC, H4A 3J1, Canada

* Correspondence: igor.cestari@mcgill.ca

Abstract: African trypanosomes evade host immune clearance by antigenic variation, causing persistent infections in humans and animals. These parasites express a homogeneous surface coat of variant surface glycoproteins (VSGs). They transcribe one out of hundreds of VSG genes at a time from telomeric expression sites (ESs) and periodically change the VSG expressed by transcriptional switching or recombination. The mechanisms underlying the control of VSG switching and its developmental silencing remain elusive. We report that telomeric ES activation and silencing entail an on/off genetic switch controlled by a nuclear phosphoinositide signaling system. This system includes a nuclear phosphatidylinositol 5-phosphatase (PIP5Pase), its substrate PI(3,4,5)P3, and the repressor-activator protein 1 (RAP1). RAP1 binds to ES sequences flanking VSG genes via its DNA binding domains and represses VSG transcription. In contrast, PI(3,4,5)P3 binds to the N-terminus of RAP1 and controls its DNA binding activity. Transient inactivation of PIP5Pase results in the accumulation of nuclear PI(3,4,5)P3, which binds RAP1 and displaces it from ESs, activating transcription of silent ESs and VSG switching. The system is also required for the developmental silencing of VSG genes. The data provides a mechanism controlling reversible telomere silencing essential for the periodic switching in VSG expression and its developmental regulation.

Keywords: Trypanosomes, antigenic variation, repressor-activator protein 1 (RAP1), phosphoinositides, signaling.

Abbreviations: PI(3,4,5)P3, phosphatidylinositol-(3,4,5)-triphosphate; PIP5Pase, phosphatidylinositol phosphate 5-phosphatase; RAP1, repressor activator protein 1; ES, expression site; VSG, variant surface glycoproteins.

Introduction

Antigenic variation is a strategy of immune evasion used by many pathogens to maintain persistent infections and entails changes in surface antigens to escape host immune clearance. The single-celled protozoa *Trypanosoma brucei spp.* circulate in the mammalian host bloodstream and periodically change their variant surface glycoprotein (VSG) coat to evade antibody clearance by antigenic variation (1). *T. brucei* have over 2,500 VSG genes and pseudogenes, primarily on subtelomeres, and ~20 telomeric expression sites (ESs), each containing a VSG gene (Fig 1A). Only one VSG gene is transcribed at a time from a telomeric ES. Antigenic variation occurs via transcriptional switching between ESs or VSG gene recombination within ESs (1). The active VSG gene is developmentally regulated, and its expression is silenced in parasites encountered in the tsetse fly vector, e.g., procyclic and epimastigote forms. Epimastigotes develop into transmissible metacyclic trypomastigotes, which re-activate VSG gene expression before infecting mammals. The coordinated activation and silencing of telomeric ESs is essential for the periodic switch in VSG gene expression during antigenic variation and VSG developmental regulation; however, the mechanisms underlying the control of this process remain unknown.

The expressed VSG gene is transcribed by RNA polymerase I (RNAP I) from a compartment outside the nucleolus termed ES body (ESB) (2). Transcription initiates in all ESs but only elongates through one ES (hereafter called active ES), resulting in VSG monogenic expression. The silencing of ES VSG genes involves its proximity to telomeres (3, 4) and the telomere-associated factor repressor activator protein 1 (RAP1) (5). RAP1 is conserved among eukaryotes (5-8) and functions in telomere silencing (5, 8), telomere end protection (9-11), and non-telomeric gene regulation in mammals and yeast (8, 11). Histones and chromatin-modifying enzymes, such as histone methyltransferase and bromodomain proteins, also associate with ESs and contribute to their repression (12-15). In contrast, the active ES is depleted of nucleosomes (12) and enriched in proteins that facilitate its transcription, e.g., VSG exclusion (VEX) 1 and 2 and ES body 1 (ESB1), and processing of the highly abundant VSG mRNAs (16, 17).

The mechanisms underlying the initiation or control of VSG switching remain unknown. Antigenic variation was thought to occur stochastically (18) with the DNA break and repair machinery involved in VSG recombination (19-22). However, we showed that phosphoinositide signaling plays a role in the expression and switching of VSG genes (23, 24). This system entails the plasma membrane/cytosolic-localized phosphatidylinositol phosphate 5-kinase (PIP5K) and phospholipase C (PLC) (23), and the nuclear phosphatidylinositol phosphate 5-phosphatase (PIP5Pase). The PIP5Pase enzyme interacts with RAP1 (24, 25), and either protein knockdown results in the transcription of all ESs simultaneously (5, 23, 24). RAP1 and PIP5Pase interact with other nuclear proteins, including nuclear lamina, nucleic acid binding proteins, and protein kinases and phosphatases (24). The involvement of phosphoinositide signaling in VSG expression and switching implied that signaling and regulatory processes have a role in antigenic variation linking cytosolic and nuclear proteins to control transcriptional and recombination mechanisms.

We report here that PIP5Pase, its substrate PI(3,4,5)P3, and its binding partner RAP1 form a genetic regulatory circuit that controls ES activation and repression. We show that RAP1 binds to silent

telomeric ESs sequences flanking the VSG genes, and the binding is essential for VSG transcriptional repression. This association is regulated by PI(3,4,5)P3, which binds to the N-terminus of RAP1, acting as an allosteric regulator controlling RAP1-ES interactions. The system depends on PIP5Pase catalytic activity, which dephosphorylates PI(3,4,5)P3 and prevents its binding to RAP1. Inactivation of PIP5Pase results in PI(3,4,5)P3 and RAP1 binding, which displaces RAP1 from ESs, leading to transcription and switching of VSGs in the population. Hence, PIP5Pase activity is required for RAP1 association with silent ESs and VSG gene transcriptional control. The regulatory system is essential for VSG developmental silencing, and temporal disruption of this nuclear signaling system results in VSG switching. The data indicate a molecular mechanism by which ESs are periodically turned on and off during antigenic variation and parasite development.

Results

PIP5Pase activity controls VSG switching and developmental silencing

To investigate the role of PIP5Pase activity in VSG gene expression and switching, we performed gene expression analysis in *T. brucei* bloodstream forms that exclusively express a wildtype (WT) or a catalytic mutant D360A/N362A (Mut) PIP5Pase; the latter is unable to dephosphorylate PI(3,4,5)P3 (24). The exclusive expression cells have the endogenous PIP5Pase alleles replaced by drug-selectable markers, a tetracycline (tet)-regulatable PIP5Pase allele introduced in the silent rDNA spacer, and a V5-tagged WT or Mut PIP5Pase allele in the constitutively expressed tubulin loci (24). In the absence of tet, the cells exclusively express the WT or Mut PIP5Pase allele (24). We performed RNA-seq after 24h exclusive expression of the WT or Mut PIP5Pase with Oxford nanopore sequencing (~500 bp reads) to distinguish the VSG genes expressed. The expression of the Mut PIP5Pase resulted in 1,807 genes upregulated and 33 downregulated (Figure 1B, $\geq$ 2-fold change, *p*-value $\leq$ 0.01, Supplementary File 1). Notably, all silent ES VSG genes were upregulated (Figure 1B and C), consistent with their transcription. In contrast, there was a ~10-fold decrease in the active VSG (VSG2) and ESAG mRNAs (Figure 1B and C), implying decreased expression of the active ES (BES1) genes, perhaps resulting from competition among ESs for polymerase or factors required for their transcription. The remaining upregulated genes were primarily from silent subtelomeric arrays, largely VSG genes and pseudogenes (Figure 1B and D), indicating that PIP5Pase activity is also required for subtelomeric ES repression. Other upregulated genes included primarily retrotransposons, endonucleases, and RNA polymerases, but most chromosome core genes (Pol II and Pol III transcribed genes) were not significantly affected (Figure 1B). Re-establishing expression of WT PIP5Pase for 60h after its 24h withdrawal restored VSG monogenic expression. However, analysis of this population by VSG-seq revealed the expression of several VSGs other than the initially expressed VSG2 indicating a switch in VSG expression (Fig 1E). In contrast, only a few VSGs were detected in cells expressing WT PIP5Pase by VSG-seq (Fig 1E). To verify PIP5Pase role in VSG switching, we knocked down PIP5Pase for 24h (Tet -), then restored its expression (Tet +) and isolated clones by limit dilution and growth for 5-7 days (Figure S1). Analysis of isolated clones after temporary PIP5Pase knockdown (Tet -/+) confirmed VSG switching in 93 out of 94 (99%) analyzed clones (Figure 1F, Figure 1-figure supplement 1). The cells switched to express VSGs from silent ESs or subtelomeric regions, indicating switching by transcription or recombination

mechanisms. Moreover, no switching was detected in 118 isolated clones from cells continuously expressing WT PIP5Pase (Tet +, Figure 1F). The data imply that PIP5Pase activity can control the transcription and switching of VSGs.

The active VSG gene is expressed in bloodstream forms but silenced during parasite development to the insect stage procyclic forms. To determine whether PIP5Pase is required for VSG developmental silencing, we generated procyclic conditional null cells expressing a tet-regulatable V5-tagged PIP5Pase (Figure 1-figure supplement 2). Immunofluorescence analysis showed that PIP5Pase-V5 localizes in the nucleus of procyclic forms (Figure 1-figure supplement 2), as in bloodstream forms (23). Western analysis showed that PIP5Pase-V5 proteins were eliminated 72h after knockdown (Figure 1G), and growth arrest was detected after five days of knockdown (Figure 1-figure supplement 2). Gene expression analysis showed a 5-15 log2-fold upregulation of all ES VSG genes at 72h (Figure 1H) at a time when cells are viable and dividing. The data indicate that VSG switching and developmental silencing depend on PIP5Pase activity.

RAP1 bind to sequences flanking VSG genes in silent telomeric ESs

Because PIP5Pase and its substrate PI(3,4,5)P3 interact with RAP1 (24), we postulated that VSG silencing might involve the regulation of RAP1's repressive function. We showed that RAP1 binds telomeric or 70 bp repeats (24), but it is unknown if it binds to other ES sequences or genomic loci. We performed ChIP-seq with a cell line expressing an *in-situ* HA-tagged RAP1 and nanopore sequencing (~500 bp reads). We found that RAP1-HA binds primarily to silent ESs at the 70 bp and telomeric repeats, which are sequences upstream and downstream of the VSG genes in bloodstream-form ESs, respectively (Figure 2A-B, E and Figure 2-figure supplement 1; p-values $< 10^{-4}$). There was also a slight enrichment of RAP1-HA in the 50 bp repeat sequences preceding the ES promoters (Figure 2A and Figure 2-figure supplement 1; p-values $< 10^{-4}$). However, RAP1-HA did not bind significantly to genes, including VSG genes or pseudogenes in the ESs or subtelomeric regions (Figure 2A-B, F). Notably, analysis of uniquely mapped reads showed that RAP1-HA was depleted from the active ES (Figure 2C-D and Figure 2 -figure supplement 2), consistent with RAP1 having a repressive function. The RAP1-HA bound sequences are rich in TAAs and GTs, including the 70 bp $(TTA)_7(GT/A)_9$ and telomeric $(TTAGGG)_n$ repeats (Figure 2A), but also T and G/T rich sequences downstream of VSG genes. RAP1-HA was also bound to metacyclic ESs at AT/GC-rich or (TAACCC)n repeat sequences near VSG genes (Figure 2E, p-values $< 10^{-4}$, Figure 2-figure supplement 3). RAP1-HA also bound sparsely to subtelomeric regions, including centromeres (Figure 2F, p-values $< 10^{-4}$), which are AT-rich in trypanosomes (26), and some binding sites overlapped with the centromeric protein kinetoplastid kinetochore 2 (KKT2) (27). Other poorly enriched RAP1-HA peaks in the subtelomeric regions reflect residual ES sequences from recombination (Figure 2F). Hence, RAP1-HA binds primarily to telomeric and 70 bp repeats within silent ESs. The 70 bp repeats are ES-specific sites for RAP1 binding near silent VSG genes.

To confirm RAP1 binding to ES sequences, we expressed and purified from *E. coli* recombinant 6xHis-tagged *T. brucei* RAP1 (rRAP1-His) (Figure 3A-B). We performed electrophoretic mobility shift assays (EMSA) using rRAP1-His and biotinylated telomeric repeats, 70 bp repeat, or a scrambled telomeric

repeat sequence as a control. rRAP1-His bound to telomeric and 70 bp repeats but not to the scrambled
DNA sequence (Fig 3C). We obtained similar results by microscale thermophoresis (MST) kinetics
using rRAP1-His and Cy5-labelled DNA sequences (Figure 3D, Figure 3-figure supplement 1). The
MST analysis revealed a Kd of 24.1 and 10 nM of rRAP1-His for telomeric and 70 bp repeats,
respectively (Table 1). RAP1 has two DNA binding domains, a central Myb (position E426-Q500) and a
C-terminal Myb-like (MybL, position E639-R761) domain (Figure 3A), and a divergent BRCT domain
(S198-P385) (5), which is involved in RAP1 self-interaction (28) and perhaps other protein interactions
(23, 24). To identify which RAP1 domain mediates DNA interactions, we performed EMSA with
rRAP1$^{1-300}$-His (aa positions in superscript), rRAP1$^{301-560}$-His, and rRAP1$^{561-855}$-His, the last two
fragments encompass the Myb and MybL domains, respectively (Figure 3A-B). Both rRAP1$^{301-560}$-His
and rRAP1$^{561-855}$-His proteins bound to telomeric or 70 bp repeats (Figure 3E-F, Figure 3-figure
supplement 1), but no DNA binding was detected with the N-terminal rRAP1$^{1-300}$-His (Fig 3G). Because
all proteins were His-tagged and no binding was detected to scrambled DNA nor with rRAP1$^{1-300}$-His,
unspecific DNA binding by the His-tag is ruled out. Our data show that both Myb and MybL can
independently and directly bind to 70 bp and telomeric repeats.

PIP5Pase activity controls PI(3,4,5)P3 binding to the N-terminus of RAP1

We found in the N-terminus of RAP1 a villin headpiece (VHP) domain (Fig 3A, position Y59 to F94, *e*-
value $< 10^{-4}$), which is typically present in Villin proteins and binds phosphoinositides (29). *T. brucei*
RAP1 VHP domain is conserved (~27% aa identity) with other VHPs in Villin proteins forming a three
helices fold over a hydrophobic core (30, 31) (Figure 4A), which is essential for phosphoinositide
binding (29). We performed binding assays with rRAP1-His and biotinylated phosphoinositides (25) and
confirmed that rRAP1-His binds specifically to PI(3,4,5)P3 (Figure 3-figure supplement 1) (24).
Moreover, we found that rRAP1$^{1-300}$-His binds to PI(3,4,5)P3 (Figure 4B), but not other
phosphoinositides or inositol phosphates tested (Figure 4C) and the binding was competed by a molar
excess of unlabelled PI(3,4,5)P3 (Figure 4D). However, no PI(3,4,5)P3 binding was detected with
rRAP1$^{301-560}$-His or rRAP1$^{561-855}$-His proteins, indicating that RAP1 binds to PI(3,4,5)P3 via its N-
terminus containing the VHP domain. Moreover, binding kinetics by differential scanning fluorescence
with unlabelled phosphoinositides showed that rRAP1-His binds to PI(3,4,5)P3 with a Kd of 19.7 µM
(Figure 4E, Table 1). To determine if RAP1 binds to PI(3,4,5)P3 *in vivo,* we *in-situ* HA-tagged RAP1 in
cells that express the WT or Mut PIP5Pase and analyzed endogenous PI(3,4,5)P3 levels associated with
immunoprecipitated RAP1-HA. There were ~100-fold more PI(3,4,5)P3 associated with RAP1-HA in
cells expressing the Mut than the WT PIP5Pase (Figure 4F, Figure 4-figure supplement 1). In contrast,
there was a mild but not significant difference in the total cellular PI(3,4,5)P3 levels comparing cells
expressing WT versus Mut PIP5Pase (Figure 4F), implying that PIP5Pase activity controls a localized
pool of PI(3,4,5)P3 in the nucleus available for RAP1 binding. Hence, RAP1 binds specifically to
PI(3,4,5)P3 via its N-terminus in a PIP5Pase activity-dependent fashion.

PI(3,4,5)P3 is an allosteric regulator of RAP1 and controls telomeric ES repression

Because RAP1 binds to PI(3,4,5)P3 via its N-terminus and to ESs via its central and C-terminus Myb and Myb-L domains, we posited that PI(3,4,5)P3 binds to RAP1 and controls its association with ESs. We performed EMSA binding assays and found that PI(3,4,5)P3 inhibited rRAP1-His binding to telomeric repeats in a dose-dependent fashion, but no inhibition was detected with PI(4,5)P2 (Figure 5A-B). Similar PI(3,4,5)P3 binding inhibition of rRAP1-His was observed for 70 bp repeats (Figure 5B). Moreover, MST binding kinetics with 30 µM of PI(3,4,5)P3 increased rRAP1-His Kd to telomeric and 70 bp repeats in ~6.5-fold and ~18.5-fold, respectively (Figure 5C, Table 1). Notably, PI(3,4,5)P3 did not affect rRAP1$^{301-560}$-His or rRAP1$^{561-855}$-His binding to telomeric or 70 bp repeats (Figure 3-figure supplement 1), implying that PI(3,4,5)P3 does not compete with Myb or MybL domains direct binding to DNA. PI(3,4,5)P3 inhibition of RAP1-DNA binding might be due to its association with RAP1 N-terminus causing conformational changes that affect Myb and MybL domains association with DNA.

Our data suggest a model in which PI(3,4,5)P3 levels control RAP1 binding to ESs and thus silencing and activation of VSG genes. To evaluate this model, we performed ChIP-seq with RAP1-HA in cells that exclusively express WT or Mut PIP5Pase. The Mut PIP5Pase expression results in local PI(3,4,5)P3 available for RAP1 binding (Figure 4F), whereas WT PIP5Pase dephosphorylates PI(3,4,5)P3 into PI(3,4)P2, which cannot bind RAP1 (24). The expression of Mut PIP5Pase abolished RAP1-HA binding to 70 bp, 50 bp, and telomeric repeats in all bloodstream and metacyclic ESs (Figure 5D-G). Notably, the decreased RAP1-HA binding to ESs correlates with increased expression of VSGs and ESAG genes (Figure 5D-E, RNAseq), indicating that PIP5Pase controls RAP1 binding to ESs and thus VSG gene expression. The increased level of VSG and ESAG mRNAs detected in cells expressing Mut PIP5Pase (Figure 5D) may reflect increased Pol I transcription. It is possible that the low levels of RAP1-HA at the 50 bp repeats affect Pol I accessibility to the BES promoter; alternatively, RAP1 association to telomeric or 70 bp repeats may affect chromatin compaction or folding impairing VSG and ESAG genes transcription. The expression of Mut PIP5Pase also removed RAP1-HA peaks in subtelomeric regions (Figure 2F), which might impact subtelomeric chromatin organization and thus VSG expression and recombination (Figure 1D-F). We showed that RAP1 interacts with PIP5Pase within a 0.9 MDa complex, and this association is stable in cells expressing the Mut PIP5Pase (24). Hence, RAP1 dissociation from ESs is unlikely the result of PIP5Pase mutations affecting the complex integrity. Our data indicate that PIP5Pase activity regulates RAP1 binding to DNA via PI(3,4,5)P3, thus controlling the reversible silencing of telomeric VSG genes.

Discussion

We found that the reversible silencing of telomeric VSG genes in *T. brucei* is controlled by a phosphoinositide regulatory system. The system operates via PI(3,4,5)P3 regulation of RAP1-DNA binding and is controlled by PIP5Pase enzymatic activity. The system functions as an on/off genetic switch to control telomeric ES activation and silencing and indicates a mechanism to control the periodic switching of VSG genes during antigenic variation and VSG developmental silencing. The regulation requires PIP5Pase activity for continuous ES repression via dephosphorylation of PI(3,4,5)P3. PIP5Pase temporary inactivation results in the accumulation of PI(3,4,5)P3 bound to the RAP1 N-terminus. This binding displaces RAP1 Myb and MybL domains from DNA, likely due to RAP1 conformational

changes, allowing polymerase elongation through the VSG gene. PI(3,4,5)P3 acts as a typical allosteric
regulator controlling RAP1-DNA interactions and, thus, VSG transcriptional repression.

Our data indicate that regulation of PIP5Pase activity and PI(3,4,5)P3 levels play a role in VSG
switching. Other phosphoinositide enzymes such as PIP5K or PLC, which act on PI(3,4,5)P3 precursors,
also affect VSG expression and switching (23), arguing that this system is part of a cell-wide signaling
network and includes the transcriptional and recombination machinery. The finding of RAP1 binding to
subtelomeric regions other than ESs, including centromeres, requires further validation. Nevertheless, it
suggests a role for RAP1 repressing subtelomeric chromatin. Chromatin conformational capture analysis
showed that subtelomeric regions and centromeres (many subtelomeric in *T. brucei*) have a high
frequency of interactions forming the boundaries of genome compartments in *T. brucei* (13). Disrupting
PIP5Pase activity affects RAP1 association with those regions and, thus, might impact long-range
chromatin interactions. This organization may explain RAP1's role in VSG recombinational switching
(32) and the transcription of subtelomeric VSGs upon PIP5Pase mutation, RAP1 (33) or NUP1
knockdowns (34). Although we showed that C-terminal HA-tagged RAP1 protein has telomeric
localization (23) and interactions with other telomeric proteins (24); we cannot rule out potential
differences between HA-tagged and non tagged RAP1.

RAP1 and PIP5Pase interact and localize near the nuclear periphery (23, 24). RAP1 also associates with
NUP1 (24), which knockdown derepresses silent VSG genes (34). PI(3,4,5)P3 and other
phosphoinositides are synthesized in the endoplasmic reticulum (ER) and Golgi (35, 36) and distributed
to organelles, including the nuclear membrane (23, 36). The scenario suggests a model in which RAP1
associates with silent telomeric ESs and lamina proteins at the nuclear periphery (13, 23, 34), where
regulation of silencing and activation might occur. The compartmentalization of silent ESs near the
nuclear lamina may prevent the association of factors required for ES transcription, including chromatin-
associated proteins and RNA processing factors. In contrast, the active ES is depleted of RAP1, a
process that might involve ES mRNAs competition for RAP1 binding to ES DNA (37), and it is enriched
in proteins that facilitate transcription, including VEX2 and ESB1 (17, 38). Furthermore, the active ES
localizes in a compartment that favours efficient processing of VSG mRNAs (39, 40). Hence, the active
and silent ESs seem to occupy distinct subnuclear compartments and have specific associated proteins.
Within this model, VSG switching may entail the inactivation of PIP5Pase, leading to reorganization of
silent ES chromatin, e.g., RAP1 removal from ESs, ESs relocation away from the nuclear lamina, and
perhaps changes in ES three-dimensional organization (Figure 5 -figure supplement 1). Once RAP1 is
dissociated from ESs, other factors, such as ESB1 and VEX2, may associate with the silent ESs to
initiate transcription. Although the requirements for establishing VSG monogenic expression are
unknown, it may entail the successful association of factors with ESs for their transcription or
recruitment to the ESB (2, 39).

How the phosphoinositide signaling system is initiated to control VSG switching is unknown, but our
data suggest an alternative non-stochastic model regulating antigenic variation. It is unknown if this
signaling system regulates antigenic variation *in vivo*. Nevertheless, the data indicate that trypanosomes
may have evolved a sophisticated mechanism to regulate antigenic variation that entails genetic and
signaling processes, and such processes may be conserved in other pathogens that rely on antigenic
variation for infection, e.g., *Plasmodium* and *Giardia*, broadening opportunities for drug discovery. The

allosteric regulation of RAP1 by PI(3,4,5)P3 denotes a mechanism for phosphoinositide regulation of telomere silencing. Given that telomere silencing is conserved and dependent on RAP1 in most eukaryotes, this regulatory system may be present in other eukaryotes.

Materials and Methods

Cell culture, cell lines, and growth curves

T. brucei bloodstream forms (BF) single marker 427 conditional null (CN) and V5- or HA-tagged cell lines were generated and maintained in HMI-9 at 37°C with 5% CO_2 as described (23). *T. brucei* were cultured at McGill University under permit B-07840 (for I. Cestari laboratory). Cell lines that exclusively express wildtype (WT) or mutant D360A/N360A (Mut) PIP5Pase gene (Tb927.11.6270) were generated as previously described (24). The WT or Mut PIP5Pase mRNAs exclusively expressed from tubulin loci are 1.6 and 1.3-fold the WT PIP5Pase mRNA levels expressed from endogenous alleles in the single marker 427 strain. The fold-changes were calculated from RNA-seq counts per million from this work (WT and Mut PIP5Pase, Supplementary File 1) and our previous RNA-seq from single marker 427 strain (24). *T. brucei* 29.13 were maintained in SDM-79 medium supplemented with 10% FBS, G418 (2.5 µg/mL), and hygromycin (50 µg/mL) at 27°C. To generate *T. brucei* procyclic forms (PF) CN PIP5Pase, PF 29.13 was transfected with a V5-tagged PIP5Pase gene cloned into pLEW100-3V5 (23). The plasmid was integrated into the rRNA silent spacer, and transgenic cells were selected by resistance to phleomycin (5 µg/mL). The PIP5Pase endogenous alleles were then sequentially replaced by homologous recombination with PCR constructs containing about 500 bp of the PIP5Pase 5'UTR and 3'UTR flanking a puromycin or blasticidin drug resistance gene (23). The CRISPR/Cas9 ribonucleoprotein complex was used to cut the second endogenous allele and increase homologous recombination rates. Guides targeting 5' and 3' regions (see primer sequences in Supplementary File 6) of the PIP5Pase coding sequence were produced by in vitro transcription with T7 polymerase (Promega), and co-transfected with recombinant saCas9 Nuclease NLS Protein (Applied Biological Materials Inc.), and a PCR-based puromycin resistance construct containing 5'- and 3'-UTRs of the targeted gene. Note that this procedure increased the efficiency of recombination by about 20 times. Cumulative growth curves were performed by diluting *T. brucei* PFs to 2×10^6 parasites/mL in the absence or presence of tetracycline (tet, 0.5 µg/mL). The culture growth was counted every two days using a Coulter Counter (Beckman Coulter) for twelve days. Cell lines generated in this work are available upon request.

Immunofluorescence and Western blotting

Immunofluorescence: Immunofluorescence was done using *T. brucei* PF PIP5Pase CN, which expresses V5-tagged WT PIP5Pase in the presence of 0.1 µg/mL of tet. Mid-log growth phase parasites were washed three times in PBS with 6 mM glucose (PBS-G) for 10 minutes (min). Then, 2.0×10^6 cells were fixed in 2% paraformaldehyde in PBS for 10 min at room temperature (RT) onto poly L-lysine treated 12 mm glass coverslips (Fisher Scientific). The coverslips were washed three times in PBS, and cells permeabilized in 0.2% Nonidet P-40 in PBS for 10 min. Cells were washed five times in PBS, and then blocked for 1 hour (h) in 10% nonfat dry milk in PBS. After, cells were incubated in □-V5 mouse monoclonal antibodies (mAb) (Thermo Life Technologies) diluted 1:500 in 1% milk in PBS for 2 h at RT. The cells were washed five times in PBS and incubated in goat anti-mouse IgG (H+L) cross-adsorbed secondary antibody, Alexa Fluor 488 (Thermo Fisher Scientific, catalog number

A11001) 1:1,000 in 1% milk in PBS for 2 h at RT. Cells were washed five times in PBS and stained in 10 µg/mL of 4',6-diamidino-2-phenylindole (DAPI) diluted in PBS for 15 min at RT. Cells were washed four times in PBS, twice in water, and then mounted onto microscope glass slides (Fisher Scientific) using a mounting medium (Southern Biotech). Cells' images were acquired using a Nikon E800 Upright fluorescence microscope (Nikon). *Western blotting:* Western analyses were performed as previously described (25). Briefly, cleared lysates of *T. brucei* PFs were prepared in 1% Triton X-100 in PBS with 1X protease inhibitor cocktail (Bio-Vision) and mixed in 4X Laemmli buffer (Bio-Rad) with 710 mM β-mercaptoethanol and heated for 5 min at 95°C. Proteins were resolved in 10% SDS/PAGE and transferred to nitrocellulose membranes (Sigma Aldrich). Membranes were probed for 2 h at RT (or overnight at 4°C) with mAb α-V5 (BioShop Canada Inc., catalog number TAG006.100) 1:2,500 in 6% milk in PBS 0.05% Tween (PBS-T). Membranes were incubated in 1:5,000 goat anti-mouse IgG (H+L)-HRP (Bio-Rad, catalog number 1706516) in 6% milk in PBS-T, washed in PBS-T, and developed by chemiluminescence using Supersignal West Pico Chemiluminescent Substrate (Thermo scientific). Images were acquired on a ChemiDoc MP imaging system (Bio-Rad). Membranes were stripped in 125 mM glycine, pH 2.0, and 1% SDS for 30 min, washed in PBS-T, and re-probed with mAb anti-mitochondrial heat shock protein (HSP) 70 of *T. brucei* (gift from Ken Stuart, Center for Global Infectious Diseases Research, Seattle Children's) 1:25 in 3% milk in PBS-T, followed by 1:5,000 goat anti-mouse IgG (H+L)-HRP (Bio-Rad, catalog number 1706516) in 3% milk in PBS-T and developed as described above.

Protein expression and purification

The *T. brucei* RAP1 (Tb927.11.370) recombinant protein was expressed and purified as described before (24). The RAP1 fragments rRAP1$^{1-300}$-His (aa 1-300), rRAP1$^{301-560}$-His (aa 301-560), or rRAP1$^{561-855}$-His (aa 561-855) were amplified by PCR (see Supplementary File 6 for primers) from the *T. brucei* 427 genome and cloned into the pET-29a(+) vector (Novagen) using *NdeI* and *XhoI* restriction sites for the expression of proteins with a C-terminal 6xHis tag. Proteins were expressed and purified from 2 to 4 liters of *Escherichia coli* NiCo21(DE3) (New England Biolabs) as previously described (24). Briefly, lysates were sonicated in PBS supplemented with 5 mM dithiothreitol (DTT), 0.2 mg/mL lysozyme, 0.05% NP-40, 10% glycerol, and 1 mM PMSF. For rRAP1$^{1-300}$-His, this buffer was supplemented with 7 M of urea (Fisher Scientific). After sonication, His-tagged proteins in the cleared lysate were purified using Profinity™ IMAC Nickel Charged Resin (Bio-Rad) and eluted in 50 mM sodium phosphate buffer with 300 mM NaCl pH 8 and 300 mM imidazole (Fischer Scientific). Eluted proteins were dialyzed overnight at 4°C in binding buffer (25 mM HEPES, 150 mM NaCl, 10% glycerol, pH 7.5).

Phosphoinositide binding assays

Phosphoinositide binding assays were performed as previously described (20). Briefly, 1 µg of His-tagged recombinant protein (rRAP1-His, rRAP1$^{1-300}$-His, rRAP1$^{301-560}$-His, or rRAP1$^{561-855}$-His) was incubated with 1 µM of biotin-conjugated dioctanoylglycerol (diC8) PI(3,4,5)P3, diC8 PI(4,5)P2, InsP(1,4,5)P3 or InsP(1,3,4,5)P4 (Echelon Biosciences) rotating at RT for 1 h. Afterward, magnetic streptavidin beads (Sigma Aldrich) (previously blocked overnight at 4°C with 3% BSA in PBS) were added and incubated at 4°C for 1 h rotating. Using a magnetic stand, the mix was washed 5-6 times in 25 mM HEPES pH 7.5, 300 mM NaCl, 0.2% NP-40 and 0.1% Tween 20, and then eluted in 50 µL of 2x

359 Laemmli buffer supplemented with 710 mM β-mercaptoethanol. Samples were boiled at 95°C for 5 min.
360 Samples were resolved in 10% SDS-PAGE, transferred onto a nitrocellulose membrane (Sigma Aldrich),
361 and probed with □-His mAb (Thermo Scientific) diluted 1:1,000 in 6% milk in PBS-T followed by α-
362 mouse IgG HRP (Bio-Rad) diluted 1:5,000 in 6% milk in PBS-T. Membranes developed as indicated
363 above.
364
<u>Electrophoretic mobility shift assays</u>
Synthetic single-stranded 5'-biotin-conjugated DNA sequences (Supplementary File 6) of telomeric
repeats $(TTAGGG)_{10}$, 70 bp repeats (one repeat), and scrambled DNA (derived from telomeric repeats)
were annealed (1:1 v/v, 10 µM) to reverse complementary unlabeled sequences (Supplementary File 6)
(Integrated DNA Technologies) in a thermocycler by incubation at 95°C for 10 min, 94°C for 50
seconds, and a gradual decrease of 1°C every 50 seconds for 72 cycles in 200 mM Tris–HCl pH 7.4, 20
mM $MgCl_2$, 500 mM NaCl. 100 nM of annealed DNA were mixed with 1 µg of recombinant protein,
and 2 mg/mL yeast tRNA (Life Technologies) in 20 mM HEPES, 40 mM KCl, 10 mM $MgCl_2$, 10 mM
$CaCl_2$, and 0.2% N P-40, and incubated on a thermomixer (Eppendorf) at 37°C for 1 h. Then, 10 µL of
the reaction was resolved on a 6% native gel (6% acrylamide in 200 mM Tris, 12,5 mM
ethylenediaminetetraacetic acid (EDTA), pH 7.8 adjusted with acetic acid) at 100 V in 0.5 X TAE buffer
(40 mM Tris, 20 mM acetic acid, and 1 mM EDTA). DNAs were transferred onto a nylon membrane
(Life Technologies) at 100 V for 30 min in 0.5x TAE. The membrane was blocked for 1 h at RT with
nucleic acid detection blocking buffer (Life Technologies), then probed for 1 h at RT with streptavidin-
HRP 1:5,000 (Genescript) diluted in PBS 3% BSA. The membrane was washed in PBS-T and developed
using Supersignal West Pico Chemiluminescent Substrate (Thermo Fisher Scientific). Images were
acquired with a ChemiDoc MP imaging system (Bio-Rad).
382
<u>Microscale thermophoresis binding kinetics</u>
Binding assays were performed using synthetic single-stranded 5'-Cy5 conjugated DNA sequences of
telomeric repeats (TTAGGG)10, 70 bp repeats (one repeat), or scrambled DNA (derived from telomeric
repeats) (see Supplementary File 6 for sequences). Sequences were annealed (1:1 v/v, 10 µM) to reverse
complementary unlabeled sequences (Supplementary File 6, all sequences synthesized by Integrated
DNA Technologies) in a thermocycler by incubation at 95°C for 10 min, 94°C for 50 seconds, and a
gradual decrease of 1°C every 50 seconds for 72 cycles in 200 mM Tris–HCl pH 7.4, 20 mM $MgCl_2$,
500 mM NaCl. Then, 1 µM rRAP1-His was diluted in 16 two-fold serial dilutions in 250 mM HEPES
pH 7.4, 25 mM $MgCl_2$, 500 mM NaCl, and 0.25% (v/v) N P-40 and incubated with 20 nM telomeric or
70 bp repeats for 2 h at 37 °C, gently shaking. Afterward, samples were centrifuged at 2,000 x*g* for 5
min, loaded into Monolith™ capillary tubes (NanoTemper Technologies), and analyzed using the
Monolith NT.115 MiscroScale Thermophoresis instrument with MO.Control software (NanoTemper
Technologies). For binding assays in the presence of phosphatidylinositol phosphates (PIPs), the binding
reaction was prepared as above but in the presence of 30 µM of diC8 PI(3,4,5)P3 or diC8 PI(4,5)P2.
Four to six replicates were performed and presented as mean ± standard deviation. Data were analyzed
using MO.Affinity Analysis (NanoTemper Technologies).
399
<u>Differential Scanning Fluorimetry</u>

T. brucei rRAP1-His at 10 µM was incubated with 10 mM SYPRO Orange dye (ThermoFisher
Scientific), filtered before use with a 0.22 µm sterile filter (Fisher scientific), in 15 mM HEPES, 150 mM
NaCl, pH 7.0, and 10, 20, 30, or 40 µM of diC8 PI(3,4,5)P3, diC8 PI(4,5)P2, or no PIPs in 96-well
microplates (Life Technologies). The plate was sealed with MicroAmp® Optical Adhesive Film (Life
Technologies), and the reaction mixture was placed on ice for 1 h. Afterward, 20 µL of the reaction was
added to a differential scanning fluorimetry plate (Life Technologies) and run using the Applied
Biosystems StepOnePlus instrument (ThermoFisher Scientific) according to the manufacturer's
instructions. Briefly, samples were run using continuous ramp mode with two thermal profiles: Step 1 at
25°C for 2 min and step 2 at 99°C for 2 min, with a ramp rate of 100% at step 1 and 1% at step 2.
Experiments were performed in triplicate and presented as mean ± standard deviation. Data analysis was
performed using the DSFworld (https://gestwickilab.shinyapps.io/dsfworld/) platform, and melting
temperatures were calculated as the midpoint of the resulting fluorescence versus temperature curve.
413
RNA-seq and real-time PCR
Poly-A enriched RNAs were extracted from 1.0×10^8 *T. brucei* BFs CN PIP5Pase exclusively expressing
V5-tagged PIP5Pase WT or mutant D360A/N362A for 24 h using the magnetic mRNA isolation kit
(New England Biolabs) according to manufacturer's instructions. For procyclic forms, total RNAs were
extracted from 5.0×10^8 *T. brucei* CN PIP5Pase growing in Tet + (0.5 µg/mL, no knockdown) or Tet –
(knockdown) at 5h, 11h, 24h, 48h, and 72h using TRIzol (Thermo Fisher Scientific) according to
manufacturer's instructions. The isolated mRNA samples were used to synthesize cDNA using
ProtoScript II Reverse Transcriptase (New England Biolabs) according to the manufacturer's
instructions. Real-time PCRs were performed using VSG primers as previously described (23). For
RNA-seq, cDNA samples were purified using Totalpure NGS mag-bind (Omega-BioTek) at a 1.0x ratio
(beads to cDNA volume). cDNA samples were eluted in 55 uL of water, then fragmented to a size range
of 300bp – 1kb (average 700 bp) using an M220 Focused-ultrasonicator (Covaris) with 75 peak
incidence power, 10% duty factor, and 200 cycles per burst for 35 seconds at 20 °C. The fragmented
cDNAs were then used for RNA-seq library preparation for Oxford nanopore sequencing (indicated
below). Fifty fmol of barcoded libraries were sequenced in a MinION (Oxford Nanopore Technologies)
using an FLO-MIN106 flow cell (Oxford Nanopore Technologies). Three biological replicates were
performed.
431
ChIP-seq
ChIP-seq was performed with *T. brucei* BFs CN PIP5Pase expressing endogenously HA-tagged RAP1
and exclusively expressing V5-tagged PIP5Pase WT or mutant D360A/N362A. Cells were grown for
24h in the absence of tet, which results in the exclusive expression of WT or mutant PIP5Pase, in HMI-9
media supplemented with 10% FBS, 2 µg/mL of G418, 2.5 µg/mL of phleomycin, 0.1 µg/mL of
puromycin, and 25 µg/mL of nourseothricin. A total of 3.0×10^8 cells at mid-log growth were fixed in 1%
paraformaldehyde at 4°C for 10 min, then quenched for 10 min in 140 mM glycine. Fixed cells were
spun down at 2000 x*g* for 10 min, washed twice in PBS, and processed for ChIP using truChIP
Chromatin Shearing Kit with Formaldehyde (Covaris) according to the manufacturer's instructions. DNA
was sheared to a size range of 0.5-1.5 kb (average 700 bp) using an M220 Focused-ultrasonicator

(Covaris) with 75 peak incidence power, 5% duty factor, and 200 cycles per burst for 7 min at 7°C. ChIP
was performed with a total of 15 µg of sonicated chromatin and 10 µg of Mab anti-V5 (BioShop Canada
Inc.). The antibodies were cross-linked to Protein G Mag Sepharose Xtra (GE Healthcare) according to
the manufacturer's instructions before immunoprecipitations. Antibody eluted chromatin was incubated
with 20 units of Proteinase K (Thermo Fisher Scientific) and reverse cross-linked at 65°C overnight,
followed by DNA extraction by phenol:chloroform:isoamyl alcohol (25:24:1, ThermoFisher Scientific)
and isopropanol (v/v) precipitation. Samples were resuspended in water, and the DNA was selected for
fragments above 300 bp using Totalpure NGS mag-bind (Omega-BioTek) at 0.85x ratio (beads to DNA
volume). The DNA was eluted in water and used to prepare Oxford nanopore DNA sequencing libraries
(indicated below). Fifty fmol of barcoded libraries was sequenced in a MinION (Oxford Nanopore
Technologies) using an FLO-MIN106 flow cell (Oxford Nanopore Technologies). Three biological
replicates were performed.

VSG-seq
T. brucei BFs PIP5Pase CN cell lines exclusively expressing V5-tagged PIP5Pase WT or mutant
D360A/N362A were seeded at $1.0x10^3$ parasites/mL in 50 mL of HMI-9 media supplemented with 10%
FBS, 2 µg/mL of G418, 2.5 µg/mL of phleomycin, and grown for 24 h in the absence of tet, which
results in the exclusive expression of WT or PIP5Pase mutant. Then, tet (0.5 µg/mL) was added to Mut
PIP5Pase cells to rescue WT PIP5Pase expression, and cells were grown for an additional 60 h at 37°C
and 5% CO_2 until they reached a concentration of $1.0x10^6$ parasites/mL. RNA was then isolated using
Trizol (Sigma-Aldrich) according to the manufacturer's instructions. cDNA was synthesized using
random primers with 5x Protoscript II Reverse Transcriptase (New England BioLabs Ltd). cDNA was
amplified using the VSG Splice Leader and SP6-VSG14mer primers (41) (Supplementary File 6) with
denaturing at 94°C for 3 min, followed by 22 cycles of 94°C for 1 min, 42°C for 1 min, and 72°C for 2
min (41). Amplified fragments were purified using 0.55x beads/sample ratio with Mag-Bind® TotalPure
NGS (Omega Bio-Tek). Amplicons were prepared for Oxford Nanopore sequencing using the ligation
sequencing kit SQK-LSK110 (Oxford Nanopore Technologies) and PCR Barcoding Expansion kit
(EXP-PBC001) according to the manufacturer's instructions (described below). Five fmol pooled
barcoded libraries were sequenced in a MinION using a Flongle FLO-FLG001 flow cell (Oxford
Nanopore Technologies). Experiments were performed in three biological replicates.

Clonal-VSG-seq
T. brucei BFs PIP5Pase CN cell lines seeded at $1.0x10^4$ parasites/mL in 10 mL of HMI-9 media
supplemented with 10% FBS and 2 µg/mL of G418 and 2.5 µg/mL of phleomycin in the absence of tet
(tet -) for PIP5Pase knockdown, or in the presence of tet (tet +, 0.5 µg/mL) for PIP5Pase expression.
After 24 h, tet (0.5 µg/mL) was added to the tet – cells to restore PIP5Pase expression and cells were
cloned by limited dilution. A protocol for clonal VSG switching is available (42). Briefly, the cells were
diluted to a concentration of 1 parasite/300 µL and grown in 200 µL of media in 96-well culture plates
(Thermo Scientific Inc.) and grown for 5-7 days at 37°C and 5% CO_2. Clones were collected from the
plates and transferred to 24 well culture plates (Bio Basic Inc.) containing 2 mL of HMI-9 media with
10% FBS, 2 µg/mL of G418, 2.5 µg/mL of phleomycin, and 0.5 µg/mL of tet for 24 h to reach $1.0x10^6$
parasites/mL. Afterward, cells were harvested by centrifugation at 2,000 x*g*, and pellets were collected

for RNA extraction using the 96 Well Plate Bacterial Total RNA Miniprep Super Kit (Bio Basic Inc.)
according to the manufacturer's instructions. cDNA was synthesized from extracted RNA samples using
M-MuLV reverse transcriptase kit (New England Biolabs Ltd) based on the manufacturer's instructions
and using customized primers splice leader (SL_F) and VSG barcoded primers (BCA_Rd_3'AllVSGs 1
to 96) (see Supplementary File 6 for primer sequences). The forward SL F primer is complementary to
the splice leader sequence and has an Oxford nanopore barcode adapter sequence. The reverse primer
contains a sequence complementary to the 3'-region conserved among VSG genes, a unique 6 random
nucleotide barcode sequence, and an Oxford nanopore barcode adapter sequence for library preparation.
The synthesized cDNAs were pooled in a 1.5 mL Eppendorf tube and amplified using the Oxford
nanopore library barcoding primers (PCR Barcoding Expansion kit EXP-PBC001, Oxford Nanopore
Technologies) with denaturing at 95° for 3 min, followed by 22 cycles of 95°C for 30 sec, 60°C for 30
sec, and 72°C for 2.30 min. Amplified fragments were purified using 0.55x beads/sample ratio
NucleoMag® NGS magnetic beads (Takara Bio). Amplicons were prepared for Oxford Nanopore
sequencing using the ligation sequencing kit SQK-LSK110 (Oxford Nanopore Technologies) according
to the manufacturer's instructions (details described below). Five fmol of the barcoded library were
sequenced in a MinION (Oxford Nanopore Technologies) using a Flongle FLO-FLG001 flow cell
(Oxford Nanopore Technologies). A total of 118 and 94 clones of CN PIP5Pase cells tet + (PIP5Pase
expression) and tet – (PIP5Pase knockdown), respectively, were analyzed.

Preparation of DNA libraries and Oxford nanopore sequencing
DNA libraries were prepared using the ligation sequencing kit SQK-LSK110 (Oxford Nanopore
Technologies), according to the manufacturer's instructions. Briefly, the fragmented nucleic acids were
end-repaired and A-tailed using the NEBNext Ultra II End Repair and A-Tailing Module (New England
Biolabs). The samples were cleaned with Totalpure NGS mag-bind (Omega-BioTek) at 0.85x ratio
(beads to DNA volume), then ligated with nanopore barcode adapters. The product was subsequently
used for PCR amplification with barcoding primers (Oxford Nanopore Technologies). The PCR product
was cleaned with Totalpure NGS mag-bind (Omega-BioTek) at a 0.85x ratio (beads to DNA volume)
and ligated to motor proteins for DNA sequencing. The libraries were sequenced in a MinION Mk1C
sequencer (Oxford Nanopore Technologies) using FLO-MIN106 flow cells (unless otherwise stated),
and sequences were basecalled using the Guppy software (Oxford Nanopore Technologies). Sequencing
information is available in Supplementary File 5 and fastq data is available in the Sequence Read
Archive (SRA) with the BioProject identification PRJNA934938.

Computational analysis of RNA-seq, VSG-seq and ChIP-seq
RNA-seq or ChIP-seq fastq data from nanopore sequencing were mapped to *T. brucei* 427-2018
reference genome using minimap2 (https://github.com/lh3/minimap2) and the alignment data were
processed with SAMtools (https://github.com/samtools/samtools). Reads with a nanopore sequencing Q-
score > 7 were used for analysis, and the mean Q-score was 12. RNA-seq and VSG-seq (including clonal
VSG-seq) mapped reads were quantified using featureCounts from package Subread
(https://bioconductor.org/packages/release/bioc/html/Rsubread.html) and used for differential expression
analysis using EgdeR (https://bioconductor.org/packages/release/bioc/html/edgeR.html). Alignments
were filtered to remove supplementary alignments using samtools flags. Because RNA-seq reads were

detected mapping to subtelomeric regions in cells expressing Mut PIP5Pase, differential expression analysis was also performed with alignments filtered for primary reads with a stringent mapping probability of 99,9% (mapQ 30) to eliminate potential multiple mapping reads. Read counts were obtained with featureCounts counting primary alignments only and processed using EgdeR. For ChIP-seq, aligned reads mapped with minimap2 were processed with deepTools (https://deeptools.readthedocs.io/en/develop/) for coverage analysis using bamCoverage and enrichment analysis using bamCompare comparing RAP1-HA ChIP vs Input. Peak calling and statistical analysis were performed for broad peaks with Model-based Analysis of ChIP-Seq MACS3 (https://github.com/macs3-project/MACS), and data were visualized using the integrated genomics viewer tool (Broad Institute). To identify if RAP1-HA was mapping specifically to silent vs active ESs, mapped reads were filtered to remove supplementary and secondary alignments, i.e., only primary alignments were considered. Analyses were also performed with a minimum mapping probability of 90% and 99% (mapQ 10 and 20, see Fig S4). This stringent analysis removes reads aligning to multiple regions of the genome including reads mapping to multiple ESs, i.e., it retains reads aligning to specific regions of the ESs. Filtered reads were processed as described above using deepTools and MACS3 analysis. Scripts used for ChIP-seq, RNA-seq, and VSG-seq analysis are available at https://github.com/cestari-lab/lab_scripts. A specific pipeline was developed for clonal VSG-seq analysis, available at https://github.com/cestari-lab/VSG-Bar-seq.

Quantification of PI(3,4,5)P3

T. brucei BFs PIP5Pase CN cell lines exclusively expressing V5-tagged PIP5Pase WT or mutant D360A/N362A were grown in 100 mL HMI-9 media supplemented with 10% FBS, 2 µg/mL of G418, 2.5 µg/mL of phleomycin, and grown for 24 h in the absence of tet, which results in the exclusive expression of WT or Mut PIP5Pase. A total of 1.0×10^8 cells were used for PI(3,4,5)P3 quantification using Echelon's PIP3 Mass ELISA Kit (Echelon Biosciences), according to the manufacturer's instructions. Briefly, the cells were centrifuged at 2,000 x*g* at RT for 5 min, washed in PBS 6mM Glucose, and the supernatant was discarded. The cell pellets were resuspended in 10 mL of ice-cold 0.5 M Trichloroacetic Acid (TCA), incubated on ice for 5 min, then centrifuged at 1,000 x*g* for 7 min at 4°C. The supernatant was discarded, and the pellets were washed twice using 3 mL of 5% TCA/1 mM EDTA per wash at RT. Neutral lipids were extracted by adding 3 mL of MeOH: $CHCl_3$ (2:1) to the pellets and vortexed for 10 min at RT, followed by centrifugation at 1,000 x*g* for 5 min, and the supernatant discarded. This step was repeated once more to remove neutral lipids. 2.25 mL of MeOH:$CHCl_3$:HCl (80:40:1) was added to the pellets and vortexed for 25 min at RT, then centrifuged at 1,000 x*g* for 5 min to extract the acidic lipids. The supernatants were transferred to new 15 mL centrifuge tubes. 0.75 mL $CHCl_3$ and 1.35 mL 0.1 N HCl were added to the supernatants, vortexed for 30 seconds, then centrifuged at 1,000 x*g* for 5 min to separate the organic and aqueous phases. 1.5 mL of the organic (lower) phase was transferred to new 2 mL vials and vacuum dried using a SpeedVac (Thermo Scientific) at RT. The dried lipids were stored at -20°C until used. To quantify PI(3,4,5)P3 interacting with RAP1-HA, 300 mL of *T. brucei* BFs CN PIP5Pase expressing endogenously HA-tagged RAP1 and exclusively expressing V5-tagged PIP5Pase WT or mutant D360A/N362A were grown for 24h in absence of tet, which results in the exclusive expression of WT or mutant PIP5Pase, in HMI-9 media supplemented with 10% FBS, 2 µg/mL of G418, 2.5 µg/mL of phleomycin, 0.1 µg/mL of puromycin, and 25 µg/mL of nourseothricin. A

total of $3x10^8$ cells (PIP5Pase WT and Mut) were centrifuged at 2,000 x*g* at RT for 5 min, washed in 20

mL of PBS 6 mM glucose and then lysed in 5 mL of *T. brucei* cell lysis buffer (50 mM Tris, 150 mM

NaCl, 2X protein inhibitor cocktail, 0.1% NP-40, and 1% Triton-X100), rotating at 4^{o}C for 30 min.

Lysates were centrifugated at 15,000 x*g*, 4^{o}C for 30 min, and the cleared lysate was transferred to a new

15 mL falcon tube. 2% (100 µL) of the cleared lysate was collected for Western blot of HA-tagged

RAP1 (input). The remaining cleared lysate was used for RAP1-HA immunoprecipitation with anti-HA

monoclonal antibodies (ABClonal). 20 µg of anti-HA monoclonal antibodies crosslinked to protein G

MicroBeads (GE Healthcare) were added to the cleared lysate and incubated overnight at 4 ^{o}C.

Immunoprecipitated samples on beads were washed five times in wash buffer (50 mM Tris, 150 mM

NaCl, 2X protein inhibitor cocktail, 0.1% NP-40). An aliquote corresponding to 20% of the beads-bound

samples was collected and eluted in 6M Urea/100 mM Glycine pH 2.9 for Western blot analysis of

immunoprecipitated RAP1-HA protein. The remaining samples on the beads (80%) were used for acidic

lipids extraction, as indicated above. The dried lipids were resuspended in PBS-T 0.25% Protein

Stabilizer (PBS-T 0.25% PS), and PI(3,4,5)P3 content was measured using Echelon's PIP3 Mass ELISA

Kit (Echelon Biosciences), according to the manufacturer's instructions.

<u>Data presentation and statistical analysis</u>

Data are shown as means ± SEM from at least three biological replicates. Comparisons among groups

were made by a two-tailed t-test using GraphPad Prism. P-values < 0.05 with a confidence interval of

95% were considered statistically significant. Graphs were prepared using Prism (GraphPad Software,

Inc.), MATLAB (Mathworks), or Integrated Genome Viewer (Broad Institute).

References

1.   I. Cestari, K. Stuart, Transcriptional Regulation of Telomeric Expression Sites and Antigenic Variation in
Trypanosomes. *Curr Genomics* **19**, 119-132 (2018).

2.   M. Navarro, K. Gull, A pol I transcriptional body associated with VSG mono-allelic expression in
Trypanosoma brucei. *Nature* **414**, 759-763 (2001).

3.   D. Horn, G. A. Cross, A developmentally regulated position effect at a telomeric locus in Trypanosoma
brucei. *Cell* **83**, 555-561 (1995).

4.   G. Rudenko, P. A. Blundell, A. Dirks-Mulder, R. Kieft, P. Borst, A ribosomal DNA promoter replacing the
promoter of a telomeric VSG gene expression site can be efficiently switched on and off in T. brucei. *Cell*
83, 547-553 (1995).

5.   X. Yang, L. M. Figueiredo, A. Espinal, E. Okubo, B. Li, RAP1 is essential for silencing telomeric variant
surface glycoprotein genes in Trypanosoma brucei. *Cell* **137**, 99-109 (2009).

6.   J. A. Baur, Y. Zou, J. W. Shay, W. E. Wright, Telomere position effect in human cells. *Science* **292**, 2075-
2077 (2001).

7.   L. R. Myler *et al.*, The evolution of metazoan shelterin. *Genes Dev* **35**, 1625-1641 (2021).

8.   G. Kyrion, K. Liu, C. Liu, A. J. Lustig, RAP1 and telomere structure regulate telomere position effects in
Saccharomyces cerevisiae. *Genes Dev* **7**, 1146-1159 (1993).

9.   T. de Lange, Shelterin: the protein complex that shapes and safeguards human telomeres. *Genes Dev* **19**,
2100-2110 (2005).

10.   L. Lototska *et al.*, Human RAP1 specifically protects telomeres of senescent cells from DNA damage.
EMBO Rep **21**, e49076 (2020).

11.   J. M. Platt *et al.*, Rap1 relocalization contributes to the chromatin-mediated gene expression profile and
pace of cell senescence. *Genes Dev* **27**, 1406-1420 (2013).

12.   L. M. Figueiredo, G. A. Cross, Nucleosomes are depleted at the VSG expression site transcribed by RNA
polymerase I in African trypanosomes. *Eukaryot Cell* **9**, 148-154 (2010).

13.   L. S. M. Muller *et al.*, Genome organization and DNA accessibility control antigenic variation in
trypanosomes. *Nature* **563**, 121-125 (2018).

14.   L. M. Figueiredo, C. J. Janzen, G. A. Cross, A histone methyltransferase modulates antigenic variation in
African trypanosomes. *PLoS Biol* **6**, e161 (2008).

15.   D. Schulz *et al.*, Bromodomain Proteins Contribute to Maintenance of Bloodstream Form Stage Identity
in the African Trypanosome. *PLoS Biol* **13**, e1002316 (2015).

16.   L. Glover, S. Hutchinson, S. Alsford, D. Horn, VEX1 controls the allelic exclusion required for antigenic
variation in trypanosomes. *Proc Natl Acad Sci U S A* **113**, 7225-7230 (2016).

17.   J. Faria *et al.*, Monoallelic expression and epigenetic inheritance sustained by a Trypanosoma brucei
variant surface glycoprotein exclusion complex. *Nat Commun* **10**, 3023 (2019).

18.   K. W. Deitsch, S. A. Lukehart, J. R. Stringer, Common strategies for antigenic variation by bacterial, fungal
and protozoan pathogens. *Nat Rev Microbiol* **7**, 493-503 (2009).

19.   C. E. Boothroyd *et al.*, A yeast-endonuclease-generated DNA break induces antigenic switching in
Trypanosoma brucei. *Nature* **459**, 278-281 (2009).

20.   E. Briggs, K. Crouch, L. Lemgruber, C. Lapsley, R. McCulloch, Ribonuclease H1-targeted R-loops in surface
antigen gene expression sites can direct trypanosome immune evasion. *PLoS Genet* **14**, e1007729 (2018).

21.   M. S. da Silva, G. A. Hovel-Miner, E. M. Briggs, M. C. Elias, R. McCulloch, Evaluation of mechanisms that
may generate DNA lesions triggering antigenic variation in African trypanosomes. *PLoS Pathog* **14**,
e1007321 (2018).

22.   R. McCulloch, J. D. Barry, A role for RAD51 and homologous recombination in Trypanosoma brucei
antigenic variation. *Genes Dev* **13**, 2875-2888 (1999).

23.   I. Cestari, K. Stuart, Inositol phosphate pathway controls transcription of telomeric expression sites in
trypanosomes. *Proc Natl Acad Sci U S A* **112**, E2803-2812 (2015).

24. I. Cestari, H. McLeland-Wieser, K. Stuart, Nuclear Phosphatidylinositol 5-Phosphatase Is Essential for
Allelic Exclusion of Variant Surface Glycoprotein Genes in Trypanosomes. *Mol Cell Biol* **39** (2019).
25. I. Cestari, Identification of Inositol Phosphate or Phosphoinositide Interacting Proteins by Affinity
Chromatography Coupled to Western Blot or Mass Spectrometry. *J Vis Exp* 10.3791/59865 (2019).
26. S. O. Obado, C. Bot, D. Nilsson, B. Andersson, J. M. Kelly, Repetitive DNA is associated with centromeric
domains in Trypanosoma brucei but not Trypanosoma cruzi. *Genome Biol* **8**, R37 (2007).
27. B. Akiyoshi, K. Gull, Discovery of unconventional kinetochores in kinetoplastids. *Cell* **156**, 1247-1258
(2014).
28. M. Afrin, H. Kishmiri, R. Sandhu, M. A. G. Rabbani, B. Li, Trypanosoma brucei RAP1 Has Essential
Functional Domains That Are Required for Different Protein Interactions. *mSphere* **5** (2020).
29. N. Kumar, P. Zhao, A. Tomar, C. A. Galea, S. Khurana, Association of villin with phosphatidylinositol 4,5-
bisphosphate regulates the actin cytoskeleton. *J Biol Chem* **279**, 3096-3110 (2004).
30. T. K. Chiu *et al.*, High-resolution x-ray crystal structures of the villin headpiece subdomain, an ultrafast
folding protein. *Proc Natl Acad Sci U S A* **102**, 7517-7522 (2005).
31. J. Meng, C. J. McKnight, Heterogeneity and dynamics in villin headpiece crystal structures. *Acta
Crystallogr D Biol Crystallogr* **65**, 470-476 (2009).
32. V. Nanavaty, R. Sandhu, S. E. Jehi, U. M. Pandya, B. Li, Trypanosoma brucei RAP1 maintains telomere and
subtelomere integrity by suppressing TERRA and telomeric RNA:DNA hybrids. *Nucleic Acids Res* **45**, 5785-
5796 (2017).
33. M. Afrin *et al.*, TbRAP1 has an unusual duplex DNA binding activity required for its telomere localization
and VSG silencing. *Sci Adv* **6** (2020).
34. K. N. DuBois *et al.*, NUP-1 Is a large coiled-coil nucleoskeletal protein in trypanosomes with lamin-like
functions. *PLoS Biol* **10**, e1001287 (2012).
35. K. L. Martin, T. K. Smith, Phosphatidylinositol synthesis is essential in bloodstream form Trypanosoma
brucei. *Biochem J* **396**, 287-295 (2006).
36. Y. J. Kim, M. L. Guzman-Hernandez, T. Balla, A highly dynamic ER-derived phosphatidylinositol-
synthesizing organelle supplies phosphoinositides to cellular membranes. *Dev Cell* **21**, 813-824 (2011).
37. A. K. Gaurav *et al.*, The RRM-mediated RNA binding activity in T. brucei RAP1 is essential for VSG
monoallelic expression. *Nat Commun* **14**, 1576 (2023).
38. L. Lopez-Escobar *et al.*, Stage-specific transcription activator ESB1 regulates monoallelic antigen
expression in Trypanosoma brucei. *Nat Microbiol* **7**, 1280-1290 (2022).
39. J. Faria *et al.*, Spatial integration of transcription and splicing in a dedicated compartment sustains
monogenic antigen expression in African trypanosomes. *Nat Microbiol* **6**, 289-300 (2021).
40. J. Budzak, R. Jones, C. Tschudi, N. G. Kolev, G. Rudenko, An assembly of nuclear bodies associates with
the active VSG expression site in African trypanosomes. *Nat Commun* **13**, 101 (2022).
41. M. R. Mugnier, G. A. Cross, F. N. Papavasiliou, The in vivo dynamics of antigenic variation in Trypanosoma
brucei. *Science* **347**, 1470-1473 (2015).
42. A. Touray, Sternlieb, T., Isebe, T. I., Cestari, I., Identifying antigenic switching by clonal cell barcoding and
nanopore sequencing in trypanosomes. *Bio-protocol Preprint* 10.21769/p2385 (2023).

Acknowledgments

This research was enabled in part by computational resources provided by Calcul Quebec
(https://www.calculquebec.ca/en/) and the Digital Research Alliance of Canada (alliancecan.ca).

Funding:

Canadian Institutes of Health Research grant CIHR PJT-175222 (IC)

The Natural Sciences and Engineering Research Council of Canada grant RGPIN-2019-05271
(IC)

Fonds de Recherche du Québec - Nature et technologie grant 2021-NC-288072 (IC)

Canada Foundation for Innovation grant JELF 258389 (IC)

McGill University fund 130251 (IC)

Islamic Development Bank Scholarship 600042744 (AOT)

FRQNT-Ukraine postdoctoral fellowship BUKX:2022-2023 337989 (OK)

The Natural Sciences and Engineering Research Council of Canada CGS M fellowship (ML)

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: RNA-seq and ChIP-seq sequencing data is available in the
Sequence Read Archive (SRA) with the BioProject identification PRJNA934938.
695

 Figure legends

Figure 1. PIP5Pase activity is essential for VSG gene silencing and switching. A) Diagram of
bloodstream-form ESs (BES, top) and metacyclic-form ESs (MES, bottom). B) RNA-seq analysis of *T.*
brucei bloodstream forms comparing 24h exclusive expression of Mut to WT PIP5Pase. FC, fold-
change. Horizontal dotted lines indicate *p*-values at 0.05 and 0.01. Vertical dotted lines, FC at 2-fold.
Unitig genes, genes not assembled in the reference genome. C-D) RNA-seq read coverage and FC (Mut
vs WT) of silent BES7 (C, top), the active BES1 (C, middle), a silent MES (C, bottom), and
chromosome 4 subtelomere (D). Heat-map in D shows RNA-seq bins per million (BPM) reads. Gray
rectangles represent genes. A 99.9% reads mapping probability to the genome (mapQ>30) retained
alignments to subtelomeric regions. E) VSG-seq analysis of *T. brucei* bloodstream forms after temporary
(24h) exclusive expression of Mut PIP5Pase, and re-expression for 60h of WT PIP5Pase. B1-B3,
biological replicates. The color shows normalized read counts per million. A 3'-end conserved VSG
sequence was used to capture VSG mRNAs (41). See Supplementary File 3 for data and gene IDs of
VSGs. F) VSG-seq from isolated clones after PIP5Pase temporary knockdown (24h) followed by its re-
expression (Tet -/+) and cloning for 5-7 days. Clones of non-knockdown (tet +) cells were analyzed as
controls. BES1_VSG2 (Tb427_000016000), BES12_VSG (Tb427_000008000), Chr2_5A_VSG
(Tb427_000284800), Chr9_3A_VSG (Tb427_000553800). G) Western blot of V5-tagged PIP5Pase
knockdown in *T. brucei* procyclic forms. The membrane was stripped and reprobed with anti-
mitochondrial heat shock protein 70 (MtHSP70). H) Expression analysis of ES VSG genes after
knockdown of PIP5Pase in procyclic forms by real-time PCR. Data are the result of three biological
replicates.

Figure 2. ChIP-seq analysis of RAP1-HA in *T. brucei*. A) RAP1-HA binding sites on the silent
BES17. Data show fold-change comparing ChIP vs Input. Below is the sequence bias of RAP1-HA
bound regions. Black rectangles, ES genes; black triangle, ES promoter; cyan rectangle, 70 bp and
telomeric repeats. B) RAP1-HA binding to selected regions in silent BES sequences. Each dot represents
the mean of a silent BES. C) RAP1-HA enrichment to the active BES1 (fold-change comparing ChIP vs
Input). See Fig S4 for additional read mapping and filtering analysis. D) Comparison of RAP1-HA
binding to 70 bp in silent and active ESs. E) RAP1-HA enrichment over all MESs. Flanking VSGs, DNA
sequences upstream or downstream of VSG genes in MESs. F) RAP1-HA binding to subtelomere 3B of
chromosome (Chr) 9 (left) or chromosome 1 core (right). Yellow fluorescent protein (YFP)-tagged
KKT2 protein ChIP-seq from (27) is shown. RAP1-HA ChIP-seq in cells expressing Mut PIP5Pase is
shown below. *p*-values (*p)* were calculated using Model-based Analysis of ChIP-Seq (MACS) from three
biological replicates. Data show fold-change of ChIP vs Input analysis. See Supplementary File 2 and 4
for detailed statistics. ****, *p*-value < 0.0001.

Figure 3. rRAP1-His binds to telomeric and 70 bp repeats via its Myb and MybL domain. A) Top,
RAP1 diagram shows VHP, BRCT, Myb and MybL domains. Numbers indicate residue positions;
asterisk, nuclear targeting sequence. Bottom, telomeric (10 repeats), 70 bp (one repeat), or scrambled
(generated from telomeric repeats) sequences used in binding assays. B) His-tagged rRAP1, rRAP1[1-300],

rRAP$^{301-560}$, and rRAP1$^{561-855}$ resolved in 10% SDS/PAGE and Coomassie-stained. C) EMSA of His-tagged rRAP1 with biotinylated telomeric repeats (Tel rep), 70 repeats (70 bp rep), or scrambled sequences resolved in 6% native/PAGE and developed with Streptavidin-HRP. D) MST binding kinetics of rRAP1-His with Cy5-labelled telomeric repeats or 70 bp repeats. Data shown are the mean ± SDM of four biological replicates. E-G) EMSA of rRAP1$^{301-560}$-His (E), rRAP1$^{561-855}$-His (F), or rRAP1$^{1-300}$-His (G) with telomeric, 70 bp repeats, or scrambled sequences resolved in 6% native/PAGE. rRAP1$^{301-560}$-His were used as a positive control in G.

Figure 4. rRAP1-His binds to PI(3,4,5)P3 through its N-terminus. A) Modeling and alignment of RAP1 VHP domain. *Left*, RAP1 modeled structure; *middle*, VHP domain structure of human supervillin protein (PDB accession number 2K6N); and *right*, superposition of *T. brucei* RAP1 modeled VHP and human supervilin VHP domains. Alignment of *T. brucei* RAP1 VHP with human (Hs), *Arabidopsis thaliana* (At), and *Gallus gallus* (Gg) VHP domains from Villin proteins. PDB accession numbers are indicated in parenthesis. % of aa identity to *T. brucei* sequence are shown. B) Binding assays with rRAP1-His, rRAP1$^{1-300}$-His, rRAP$^{301-560}$-His, or rRAP1$^{561-855}$-His and PI(3,4,5)P3-biotin. Beads, Streptavidin-beads; FT, flow-through. Proteins were resolved in 10% SDS/PAGE and Western developed with α-His mAbs. C) Binding of His-tagged rRAP1$^{1-300}$ to biotinylated phosphoinositides or IPs. D) Binding of rRAP1$^{1-300}$-His to PI(3,4,5)P3-biotin in presence of unlabelled PI(3,4,5)P3 or PI(4,5)P2. For C and D, proteins were analyzed as in B. E) Binding kinetics of rRAP1-His with unlabelled PI(3,4,5)P3 or PI(4,5)P2. ΔTm, change in melting temperature. Data show the mean ± SDM of three biological replicates. F) Quantification of RAP1-bound PI(3,4,5)P3 (top) or total cellular PI(3,4,5)P3 (bottom) levels in *T. brucei* exclusively expressing WT or Mut PIP5Pase. Data show the mean ± SDM of four biological replicates. ****, *p*-value < 0.0001.

Figure 5. PIP5Pase controls rRAP1-His binding to telomeric ESs via PI(3,4,5)P3. A) EMSA of rRAP1-His with biotinylated telomeric repeats and increasing concentrations of PI(3,4,5)P3. B) EMSA of rRAP1-His with biotinylated telomeric repeats (left) or 70 bp repeats (right) and 30 μM of PI(3,4,5)P3 or PI(4,5)P2. For A-B, samples were resolved in 6% native/PAGE, transferred to nylon membranes, and developed with streptavidin-HRP. C) MST binding kinetics of rRAP1-His with Cy5-labelled telomeric repeats (top) or 70 bp repeats (bottom) with 30 μM of PI(3,4,5)P3 or PI(4,5)P2. Data show the mean ± SDM of four biological replicates. D-E) ChIP-seq of RAP1-HA binding to BES7 (D) or (MES_Chr11_5A) from cells that exclusively express WT or Mut PIP5Pase for 24h. RNA-seq comparing exclusive expression of Mut vs WT PIP5Pase for 24h is shown. F-G) Violin plots show RAP1-HA mean enrichment over 70 bp or telomeric repeats from all silent BESs (F) or MESs (G). Each dot represents an ES. BPM, bin per million. ChIP-seq and RNA-seq were performed in three biological replicates. ****, *p*<0.0001.

Figure 1-figure supplement 1. Clonal VSG-seq to identify VSG switching. A) Diagram of the clonal VSG-seq approach. CN PIP5Pase cells were treated with tet + (express PIP5Pase) or tet – (knockdown

PIP5Pase) for 24h. After, tet was added back to restore PIP5Pase expres,sion and cells were cloned in
96-well plates by limiting dilution. Clones were collected between 5-7 days and expanded in 24-well
plates for 24h. Then, RNAs were isolated and cDNA synthesized using splice leader forward primer and
a reverse primer hybridizing to the 3' end of VSG genes (conserved region) to capture all VSGs (41).
The 3'-end VSG primers have a 6 bp barcode sequence unique to each clone (see Methods). cDNAs
were pooled for nanopore barcoding to multiplex libraries. Libraries were sequenced in MinION. Fastq
sequences were split using clonal barcodes and mapped against *T. brucei*'s genome. B) VSG-seq read
coverage for a Tet+ control culture (clone A) that expresses VSG2, BES1 (green). Silent BES12 or
subtelomeric region of Chr2_5A is shown as a comparison. C) VSG-seq read coverage of Tet- cultures
(clone B and clone C), indicating VSG switching (red or orange); other VSGs expressed than the initial
VSG2. D) Mean of VSG-seq read counts from VSG genes expressed by the *T. brucei* clones compared
to read counts mapped to other VSG genes. The number of read counts to a specific VSG was used to
determine which VSG is expressed by each clone.

Figure 1-figure supplement 2. A) Diagram of PIP5Pase CN cell line. Constructs containing drug
resistance markers flanked by ~500 bp of the 5'-untranslated region (UTR) and 3'-UTR of the PIP5Pase
gene were used for homologous recombination and replacement of endogenous PIP5Pase alleles in *T.*
brucei 29.13 procyclic cells. SaCas9 with guides targeting the PIP5Pase gene was used to cleave the
PIP5Pase alleles and increase recombination rates. A tet-regulatable V5-tagged PIP5Pase allele (using
pLEW100-3V5 vector) was inserted in the silent ribosomal spacer, for controlled PIP5Pase gene
expression. Arrows indicated by 5'_Amp and 3'_Amp show hybridization sites of primes used to verify
the constructs' recombination. B) Real-time PCR amplification from genomic DNA of *T. brucei* 29.13
(WT) or CN PIP5Pase cell line. Primers 5'_Amp and 3'_Amp, indicated in A, were used in
amplification. The Tubulin gene was used as endogenous control. C) Real-time PCR amplification of
PIP5Pase from cDNA of CN PIP5Pase cell line growing in the presence of tet (tet +, 0.5 µg/mL for 24h)
or after removal of tet from medium (Tet -) for several time points (indicated in graph x-axis). Tubulin
was used as endogenous control. D) Subcellular localization of V5-tagged PIP5Pase in procyclic forms
using *T. brucei* CN PIP5Pase (100 ng/mL of tet). PIP5Pase-V5 was detected with α-V5 monoclonal
antibodies (mAb) followed by goat α-mouse IgG-AlexaFluor 488 (green). DAPI stains DNA (blue). DIC
(Differential image contrast). N, nuclear DNA; K, kinetoplast DNA. E) Cumulative growth curve of *T.*
brucei CN PIP5Pase (0.5 µg/mL of tet). The data shown are the mean ± SDM of three biological
replicates.

Figure 2 – figure supplement 1. RAP1-HA binding to silent telomeric ESs by ChIP-seq in cells
expressing WT or Mut PIP5Pase. BES1 and BES7 are included in the main manuscript (Figure 2 and
Figure 5). Black boxes below indicate ES genes. The asterisks indicate the location of the VSG gene.
The data shown is a fold-change comparison of RAP1-HA ChIP versus the input. For statistical analysis,
see Supplementary File 4.

Figure 2 – figure supplement 2. ChIP-seq of RAP1-HA binding to a silent (BES17) vs the active ES (BES1) after multiple alignment filtering parameters. *All alignments include primary and secondary alignments; supplementary alignments were removed. Primary MAPQ10 and MAPQ20 are primary alignments filtered for mapping quality $\geq$ 10 ($\geq$ 90% probability of correct alignment) and $\geq$ 20 ($\geq$ 99% probability of correct alignment), respectively. Note no significant enrichment of RAP1-HA on BES1. Increased stringency in alignment filtering removes reads from telomeric repeats. See Supplementary File 4 for statistics.

Figure 2 – figure supplement 3. MES Chr11_5A data are included in the main manuscript (Figure 5). The black boxes below indicate MES VSG genes, and the triangle indicates the MES promoter. The data shown is a fold-change comparison of RAP1-HA ChIP versus the input. For statistical analysis, see Supplementary File 4.

Figure 3 – figure supplement 1. Recombinant His-tagged RAP1 (rRAP1-His) binds to 70 bp, telomeric repeats, and PI(3,4,5)P3. A) MST assay of rRAP1-His (full length), rRAP1$^{301-360}$-His, and rRAP1$^{561-855}$-His binding to telomeric repeats (Tel rep), 70 bp repeats (70 bp rep), or a scrambled DNA sequence derived from telomeric repeats (see Supplementary File 6 for sequences). The data shown are the mean $\pm$ SDM of at least three biological replicates. Fnorm Δ, change in normalized fluorescence. B) Western blot of rRAP1-His binding assay with biotin-conjugated PI(3,4,5)P3, PI(4,5)P2, and Ins(1,4,5)P3. STV, streptavidin beads. rRAP1-His was detected with anti-His tag antibodies. C-D) Gel shift assays of rRAP1$^{301-560}$-His binding to biotinylated telomeric repeats (C) or 70 bp repeats (D) with and without PI(3,4,5)P3. E-F) rRAP1$^{561-855}$-His binding to biotinylated telomeric repeats (E) or 70 bp repeats (F) with and without PI(3,4,5)P3. PI(3,4,5)P3 concentrations used are indicated in the image. Samples were resolved in 6% native polyacrylamide gels, transferred to nylon membranes, and probed with streptavidin-HRP.

Figure 4 – figure supplement 1. Quantification of PI(3,4,5)P3 from *T. brucei* bloodstream form total cell extract or associated with RAP1-HA. A) *Top*, a diagram showing acidic lipids extraction and PI(3,4,5)P3 quantification from *T. brucei* total cell extracts (PIP5Pase WT and Mut); *bottom*, diagram of acidic lipids extraction from immunoprecipitated RAP1-HA eluates (PIP5Pase WT and Mut). PI(3,4,5)P3 was quantified using a competitive ELISA assay. B) *Top*, Western blot of RAP1-HA immunoprecipitation from *T. brucei* bloodstream forms (PIP5Pase WT and Mut) using anti-HA mAbs. The Western shows biological replicate 1 with input, flowthrough (FT), wash, and eluate. *Bottom*, Western blot of immunoprecipitated RAP1-HA (eluates) from biological replicates 2, 3, and 4 (PIP5Pase WT and Mut. WT, cells expressing WT PIP5Pase; Mut, cells expressing mutant PIP5Pase; IP, immunoprecipitation; Bio, biological replicates; kDa, kilodalton).

Figure 5-figure supplement 1. PI(3,4,5)P3 control of VSG expression and model of VSG switching and developmental silencing. A) *Left*, PIP5Pase dephosphorylates PI(3,4,5)P3 to PI(3,4)P2, and thus prevents its interaction with RAP1. RAP1 binds to ES DNA via Myb and MybL domains to maintain silent ESs. *Right,* Inactivation of PIP5Pase (e.g., mutation of catalytic site) results in the accumulation of PI(3,4,5)P3, which binds RAP1 N-terminus. The binding results in the displacement of RAP1 from ES DNA, thus resulting in VSG transcription. B) PIP5Pase inactivation may result in the derepression of multiple ESs, of which one is selected for monogenic expression in the ESB by unknown mechanisms, however, it may involve proteins that associate with the active ES such as VEX1, VEX2, and ESB1. Recombination of VSG genes may also occur during this process. The remaining derepressed ESs return to repression once PI(3,4,5)P3 levels are depleted, e.g., PIP5Pase reactivation. The system may similarly function to repress the active ES during developmental silencing of VSG genes and parasites differentiation from bloodstream forms to the insect stage procyclic forms. The location of RAP1, PIP5Pase, and telomeres near nuclear lamina reflects RAP1 and PIP5Pase immunofluorescence (23), telomere analysis by fluorescence in-situ hybridization (23), and PIP5Pase and RAP1 protein interaction analysis with nuclear lamina proteins, e.g., NUP1 (24).

Supplementary File 1. RNA-seq analysis of *T. brucei* bloodstream form cells expressing mutant vs wildtype PIP5Pase. Gene expression analysis by RNA-seq using Oxford nanopore sequencing of *T. brucei* bloodstream forms exclusively expressing mutant (D360A/N362A) compared to wildtype PIP5Pase. cDNA was synthesized from poly-A enriched RNAs. The data combines three biological replicates. LogFC, log2 fold-change; LogCPM, log2 counts per million; FDR, false discovery rate, calculated using the Benjamini-Hochberg method.

Supplementary File 2. Statistical analysis of RAP1-HA ChIP-seq in *T. brucei* expressing WT PIP5Pase. Location of ChIP-seq peaks and statistical analysis were obtained using Model-based Analysis of ChIP-Seq (MACS3) set for broad peaks search. The *q*-values (false discovery rate) are adjusted *p*-values by the Benjamini-Hochberg method. Fold enrichment is calculated for the region against random Poisson distribution with background lambda. Pileup indicates the mean value across all values in the entire peak region. Data for BESs, MES, and centromeres were included in Supplementary File 4, hence not included in this dataset. Three biological replicates comparing RAP1-HA ChIP versus Input were included in the analysis.

Supplementary File 3. VSGs identified in VSG-seq experiments (related to Fig 1E). Data compares VSGs expressed in cells expressing WT (wild type) or Mut (Mutant, D360A/N362A) PIP5Pase. Mutant PIP5Pase was temporarily knocked down for 24h followed by re-expression for 60h. Values are counts per million in log2.

Supplementary File 4. Analysis of RAP1-HA binding sites on BESs, MES, and centromeres by ChiP-seq comparing ChIP vs Input. Location of peaks and statistical analysis were obtained using Model-based Analysis of ChIP-Seq (MACS3). The *q*-values (false discovery rate) are adjusted *p*-values

by the Benjamini-Hochberg method. The location of centromeres was obtained by analysis of ChIP-seq

data from the kinetoplastid kinetochore protein 2 (KKT2) generated by Akiyoshi and Gull (27)

(accession numbers SRX372731 and SRX372732) mapped to the *T. brucei* 427-2018 genome reference.

FE, fold enrichment for the region against random Poisson distribution with background lambda; Pileup,

mean value across all values in the entire peak region.

Supplementary File 5. Oxford nanopore sequencing metrics for RNA-seq and ChIP-seq. RNA-seq

in cells expressing WT or Mut (D360A/N362A) PIP5Pase, and ChIP-seq of RAP1-HA in cells

expressing WT or Mut PIP5Pase.

Supplementary File 6. List of oligonucleotides used in this study. Oligonucleotides used for cloning

RAP1 recombinant DNA, synthesis of telomeric and 70 bp repeats, synthesis of CRISPR guide RNAs,

and clonal VSG-seq library preparation. BCA_Rd, nanopore barcode adaptor and random 6 nucleotide

sequence. Oligonucleotide modifications, e.g., 5'-biotin (5Biosg) or 5'-Cy5^{TM} fluorescence modification

(5Cy5) are indicated. Oligonucleotides are shown from 5' to 3'.

912 **Table 1. Binding kinetics of rRAP1 to telomeric repeats, 70 bp repeats, and phosphoinositides.**
913 Data show the mean ± standard deviation of the mean (SDM).

Interactions	Kd [M] ± SDM
rRAP1-His + Telomeric repeats	$24.1 \times 10^{-9} \pm 2.0 \times 10^{-9}$
rRAP1-His + 70 bp repeats	$10.0 \times 10^{-9} \pm 0.33 \times 10^{-9}$
rRAP1-His + PI(3,4,5)P3	$19.7 \times 10^{-6} \pm 2.8 \times 10^{-6}$
rRAP1-His + PI(4,5)P2	No binding
rRAP1-His + Telomeric repeats + PI(3,4,5)P3	$154.6 \times 10^{-9} \pm 14.2 \times 10^{-9}$
rRAP1-His + Telomeric repeats + PI(4,5)P2	$19.4 \times 10^{-9} \pm 1.5 \times 10^{-9}$
rRAP1-His + 70 bp repeats + PI(3,4,5)P3	$187.7 \times 10^{-9} \pm 29.9 \times 10^{-9}$
rRAP1-His + 70 bp repeats + PI(4,5)P2	$17.5 \times 10^{-8} \pm 0.9 \times 10^{-9}$

A

BES 17

$p < 10^{-10}$

RAP1-HA ChIP/Input (log2)

50 bp repeats   *ESAGs*   *70 bp repeats*   *Tel*

VSG

70 bp repeats $(TAA)_7(GT/A)_9$

ATAATAATAAtAtAATAATATATAtAtGGAGAGTGTTGTGAGgGTgTGTATATCGgATATTAtAaTAAGAGCAGTAa

Downstream of VSG genes

T-rich   G/T-rich

Telomeric repeats

TTAGGG

B

Silent BESs

RAP1-HA ChIP/Input (log2)

****   ****

50 bp rep | 70 bp rep | VSGs | Tel rep

C

BES 1

RAP1-HA ChIP/Input (log2)

50 bp repeats   *ESAGs*

Ψ   *VSG*

D

RAP1-HA ChIP/Input (70 bp enrichment, log2)

BES1 | BES2-17

E

All MESs

RAP1-HA ChIP/Input (log2)

****

Flanking VSGs | VSG

F

Chr9_3B subtelomere $p < 10^{-7}$ Chr_1 core

ChIP/Input (log2)

RAP1-HA (PIP5Pase-WT)   — Centromere $p < 10^{-7}$

YFP-KKT2

RAP1-HA (PIP5Pase-Mut)